HOXB7 induces STAT3-mediated transformation and lung metastasis in an immortalized

mammary gland NMuMG cells

Kazushi Azuma[a,+], Mai Sakamoto[a,b,+], Jun Nakayama[a,b,c,*], Shota Katayama[a], Atsuka Matsui[a], Naoki

Goshima[d,e,f], Shinya Watanabe[g], and Kentaro Semba[a,g,*]

[a] Department of Life Science and Medical Bioscience, School of Advanced Science and Engineering,

Waseda University, Tokyo, Japan.

[b] Computational Bio-Big Data Open Innovation Lab. (CBBD-OIL), National Institute of Advanced

Industrial Science and Technology (AIST), Tokyo, Japan.

[c] Laboratory of Integrative Oncology, National Cancer Center Research Institute, Tokyo, Japan.

[d] Division of Transcriptome Analysis, Translational Research Center, Fukushima Medical University,

Fukushima, Japan.

[e] Functional Proteomics Team, Molecular Profiling Research Center for Drug Discovery (molprof),

National Institute of Advanced Industrial Science and Technology (AIST), Tokyo, Japan.

[f] Department of Human Sciences, Musashino University, Tokyo, Japan.

[g] Translational Research Center, Fukushima Medical University, Fukushima, Japan.

[+] These authors equality contributed

[*] Corresponding authors

Correspondence to: Jun Nakayama (junakaya@ncc.go.jp or jnakayama.re@gmail.com) and

Kentaro Semba (ksemba@waseda.jp), Department of Life Science and Medical Bioscience, School

of Advanced Science and Engineering, Waseda University, TWIns, 2-2 Wakamatsu-cho, Shinjuku-ku,

Tokyo, 162-8480, Japan, Tel/Fax: +81-3-5369-7320.

Running Head: HOXB7 induces oncogenic transformation and metastasis

Abstract:

Genes from the homeobox family are often dysregulated in cancer. Particularly, homeobox B7

(*HOXB7*) amplification and overexpression correlate with poor prognosis in various cancers, such as

gastric, pancreatic, and lung cancers. Moreover, *HOXB7* is known to contribute to cancer

progression by promoting epithelial to mesenchymal transition, drug resistance, and angiogenesis. In

this study, we show that *HOXB7* overexpression correlates with a poor prognosis in patients with

breast cancer. Coamplification of *HOXB7* with *ERBB2* enhanced the expression of *ERBB2* and the

malignancy of HER2-positive breast cancer cells. Additionally, *HOXB7* overexpression in an

immortalized murine mammary gland epithelial cell line, NMuMG, induced cellular transformation

through an activation of the JAK-STAT signaling pathway, which resulted in lung metastasis upon

orthotropic transplantation in mice. These findings reveal that *HOXB7* alone is sufficient to induce

tumorigenesis and lung metastasis using *HOXB7*-expressing NMuMG cells.

Keywords: homeobox B7, oncogene, breast cancer, metastasis, ERBB2, normal murine mammary

gland

1. Introduction

Oncogenic alterations involving gene amplifications, mutations, chromosomal translocation, and epigenetic dysregulation induce tumorigenesis *in vivo* [1]. Activated oncogenes also trigger typical phenotypic changes, so-called transformation, in immortalized cell lines such as NIH3T3 and normal murine mammary gland (NMuMG) cells *in vitro*. Therefore, assay systems evaluating phenotypic changes, including focus formation, soft agar colonization, and tumorigenesis, have been used for the screening of novel oncogenes in previous studies including ours [2,3].

Most oncogenes cause a dysregulation of signal transduction cascades relating to cellular proliferation, cell cycle, and survival [4]. Oncogenic signaling pathways, such as RAS-MAPK and PI3K-AKT are typically activated in various cancer cell types and hence targeted by several anti-cancer drugs. Recent studies have shown that the JAK-STAT signaling is also important for cancer initiation and progression [5]. STAT3 is activated in cancer stem cell-like populations [6] and regulates stemness via fatty metabolism in breast cancer [7]. In addition, aberrant activation of STAT3 within IL6-enriched tumor microenvironments is associated with an immunosuppression of tumor-infiltrating immune cells [8]. JAK-STAT signaling further coactivates and supports cancer growth and progression via the activation of the RAS-MAPK and PI3K-AKT pathways [9].

Homeobox (HOX) genes encode transcriptional factors controlling development and organ homeostasis. Homeobox genes of the HOX superfamily (HOXA, HOXB, HOXC and HOXD clusters) are often dysregulated in breast, prostate, colon, and lung cancers. The differential dysregulation of HOX family genes in various solid tumors provides an opportunity to better understand cancer development [10,11]. The HOXB clusters are composed of *HOXB1*, *HOXB2*, *HOXB3*, *HOXB4*, *HOXB6*, *HOXB7*, *HOXB8*, *HOXB9*, and *HOXB13*, which are located at 17q23.32. HOXB cluster genes are co-amplified by gene amplification in several types of cancer. Notably, an overexpression of *HOXB7* contributes to cancer progression in breast, hepatocellular, oral, and gastric cancer cell lines [12-14]. *HOXB7* induces an activation of TGFß signaling, epithelial to mesenchymal transition (EMT), and migration in breast cancer cells [15,16]. In addition, the breast cancer cell line MCF7 of the luminal subtype acquires tamoxifen resistance via *HOXB7* expression [17]. Acting as a cofactor

of estrogen receptor (ER), *HOXB7* induces cancer progression through the expression of

downstream genes [18]. In this study, we reveal that HOXB7 activates the JAK-STAT pathway and

results in cellular transformation *in vitro*. We also found that HOXB7 is sufficient to induce

tumorigenesis and metastasis in the immortalized NMuMG cell line.

2. Materials and methods

2.1. Cell culture

NMuMG and NMuMG-luc cells [19] were cultured in Dulbecco's modified Eagle's medium (DMEM;

Fujifilm Wako, Osaka Japan) supplemented with 10% fetal bovine serum (FBS; NICHIREI

BIOSCIENCES INC., Tokyo, Japan), 100 U/mL penicillin G (Meiji-Seika Pharma Co., Ltd., Tokyo,

Japan), 100 μg/mL streptomycin (Meiji-Seika Pharma Co., Ltd.), 10 μg/mL insulin (Fujifilm Wako),

and 4.5% glucose (Fujifilm Wako) at 37°C with 5% CO_2. Plat-E cells for retroviral packaging were

cultured in DMEM supplemented with 10% FBS, 100 U/mL penicillin G, and 100 μg/mL streptomycin

at 37°C with 5% CO_2. The cells were treated with 0.1–1 μM Ruxolitinib (LC Laboratories, MA, USA).

2.2. Retroviral packaging and infection

Plat-E cells were seeded in 6-well plates and transfected with pMXs-Venus-IRES-Puromycin[R] and

pMXs-*HOXB7*-IRES-Puromycin[R] plasmid DNAs using 6 μg polyethylenimine (PEI; Polysciences, Inc.,

PA, USA). The NMuMG-Venus and HOXB7-expressing NMuMG (NMuMG-HOXB7) cell lines were

established by retroviral infection using the cell supernatants and 800 μg/mL polybrene (SIGMA

ALDRICH Co., MO, USA), as previously described [2]. To select the *Venus* or *HOXB7* expressing

cells, 1 μg/mL of puromycin (Fujifilm Wako) was added to the culture medium for 2 days.

2.3. Western blotting

The infected cell lines were lysed with SDS sampling buffer (3.5% SDS, 12% glycerol, 0.05M

Tris-HCl [pH 6.8], 0.003% bromo phenol blue, and 5% 2-mercaptoethanol) for Western blotting.

Western blotting was performed as previously described [2] using the antibodies listed in the

Supplementary Table S1.

2.4. Cell proliferation assay

A total of 5.0×10^5 NMuMG cells were seeded into a 12-well plate. Cells were harvested with 0.25%

trypsin (Thermo Fisher Scientific, MA, USA) and counted at each time point (24, 48, and 72 h)

examined [20].

2.5. Focus formation assay

To perform a focus formation assay, 7.0×10^4 parental NMuMG cells and 3.5×10^4 NMuMG-HOXB7

cells were mixed and seeded into 12-well plates. Mixed cells were cultured for over 10 days. Cells

were stained with 0.01% crystal violet in 70% methanol to detect the cellular foci [2].

2.6. Tumorigenesis assay

A total of 1.0×10^6 NMuMG cells were transplanted into 6–8-weeks-old *Rag2*$^{-/-}$ female mice with 100

μL 50% Matrigel (BD Bioscience, CA, USA) as previously described [19]. All the animal experiments

were approved by the Animal Committee of Waseda University (WD19-058, 2019-A068).

2.7. Orthotopic transplantation and bioluminescence imaging

A total of 1.0×106 NMuMG-HOXB7 cells were transplanted into the fourth fat pads of 6-week-old

BALB/cSlc-nu/nu female mice as previously described [21]. After injection, the transplantation sites

were palpated for every 2–3 days to calculate tumor formation time and the mice were weekly

monitored for primary tumor formation and metastasis formation using an *in vivo* imaging system

(IVIS Lumina Xr3; Perkin-elmer, Waltham, MA, USA). Mice were intraperitonially injected with 200 μL

of D-Luciferin (15 mg/mL; Gold Biotechnology, Inc., St. Luis, MO, USA), and luminescence was taken

under anesthesia using IVIS. Sixteen weeks after injection, all mice were sacrificed, and excised

lungs. Lungs were transferred to a 12-well plate with 900 µl of D-PBS (-) (Wako Pure Chemical

Industries) and 100 µL of D-Luciferin (15 mg/ml), and then luminescence was imaged.

2.7. Quantitative real-time PCR

Total RNA was extracted using an ISOGEN kit (NIPPON GENE Co., LTD, Tokyo, Japan), then cDNA

was synthesized using the Superscript III first-strand synthesis system (Thermo Fisher Scientific).

Quantitative RT-PCR was performed using a THUNDER BIRD SYBR qPCR mix (TOYOBO, Osaka,

Japan) with 18S rRNA serving as a relative control. The primers used are listed in the Supplementary

Table S2.

2.8. Clinical database analysis

The genomic profiles and RNA-seq datasets of The Cancer Genome Atlas (TCGA) breast cancer

(BRCA) cohorts were subjected to OncoPrint visualization and expression analyses. OncoPrint

visualization of *HOXB7* and *ERBB2* was conducted using the cBioPortal [22]. The Kaplan-Meier

method was then used to perform survival analyses of molecular taxonomy according to the breast

cancer international consortium (METABRIC) [23], using the GSE1456 and GSE9893 datasets, as

previously described [24].

2.9. Statistical analysis

Statistical analyses using the Welch's t-test, Fisher's exact test and log-rank test were performed

using R 3.4.1 (https://www.r-project.org/).

3. Results

3.1. *HOXB7* expression correlates to a poor prognosis in patients with HER2-positive breast

cancer

To elucidate the clinical significance of *HOXB7* in breast cancer, we first analyzed RNA-seq data of

TCGA BRCA cohorts for *HOXB7* expression. To further examine the genomic profiles of *HOXB7* in

TCGA BRCA cohorts, the mutational status of *ERBB2* and *HOXB7* was subjected to an OncoPrint

visualization using the cBioPortal (Figure 1A). *HOXB7* was significantly co-amplified with *ERBB2* in

30 out of 48 patients with *HOXB7* amplification (Figure 1B). Gene expression analysis based on

breast cancer subtype revealed a significantly higher expression of *HOXB7* in the HER2-positive

subtype compared to other subtypes and normal tissue (Figure 1C). These results suggest that

HOXB7 was preferentially amplified and expressed in patients with HER2-positive breast cancer.

Next, to determine the relationship between *HOXB7* expression and progression in breast

cancer, we performed a survival analysis using several public clinical datasets. In the METABRIC

cohorts, a high expression of *HOXB7* was correlated with a poor prognosis in all the patients with

breast cancer (Figure 1D left) including those of the HER2-high subtype (Figure 1D right). The

correlation of poor prognosis with *HOXB7* expression was further confirmed by analyzing other

public breast cancer cohorts comprising all subtypes (Figure 1E).

3.2. *HOXB7* induces oncogenic focus formation and activation of JAK-STAT3 signaling

To elucidate the oncogenic function of *HOXB7*, we first expressed *HOXB7* in an immortalized murine

breast epithelial cell line, NMuMG, which has been used for both the *in vitro* and *in vivo* evaluation of

oncogenes [2,19]. Expressing HOXB7 changed the morphology of NMuMG cells from epithelial-like

to spindle-like shapes (Figure 2A). Furthermore, the NMuMG-HOXB7 cells formed foci (Figure 2B).

Next, to identify oncogenic signaling pathways downstream of *HOXB7*, we performed

Western blotting using antibodies against major cancer-related signaling proteins. We found that

HOXB7 overexpression increased the protein levels of vimentin (a mesenchymal marker gene)

(Figure 2C) without changing those of E-cadherin and N-cadherin, suggesting that *HOXB7* induces a

partial epithelial-mesenchymal transition. In addition, *HOXB7* overexpression significantly increased

Stat3 phosphorylation and the protein levels of Jak1, Jak2, Stat3 and Erbb2 (Figure 2D). Major

proliferative signals, such as Erk and Akt, were unchanged by HOXB7 expression. However, Stat3

and Erbb2 were increased at the transcriptional level, while Jak2 was increased at the protein level

(Figure 2E). These results indicate that *HOXB7* induces focus formation and activates JAK-STAT

signal.

3.3. JAK-STAT signaling regulates *HOXB7*-mediated focus formation in NMuMG cells

The phosphorylation of Stat3 induced by *HOXB7* expression was completely inhibited in a

dose-dependent manner by a JAK1/2 inhibitor, Ruxolitinib, confirming JAK1/2 involvement in Stat3

phosphorylation (Figures 3A and 3B). Lapatinib, which inhibits the activation of Erbb2, decreased

p-Erk and partially reduced p-Stat3 in HOXB7-expressing cells (Figure 3C), suggesting its partial

involvement in the phosphorylation of Stat3. The protein levels of Stat3, Jak2, and Erbb2 were

unchanged by Ruxolitinib treatment, indicating that the transcriptional regulation of Stat3 and Erbb2

is not taking place downstream of Stat3 signaling. To elucidate the contribution of the oncogenic

activity of p-Stat3, we next performed proliferation and focus formation *in vitro* assays following

Ruxolitinib treatment. *HOXB7* decreased the proliferative activity of NMuMG cells, but Ruxolitinib

treatment did not affect the proliferative activity of NMuMG-HOXB7 cells (Figure 3D). Nevertheless,

Ruxolitinib inhibited their focus-forming activity (Figure 3E). These results suggest that activation of

JAK-STAT signaling by *HOXB7* induces cellular transformation.

3.4. Overexpression of *HOXB7* induces tumorigenesis and metastasis

To assess oncogenic activities *in vivo*, we performed a subcutaneous tumorigenesis assay and a

metastasis assay by orthotopic transplantation. We found that NMuMG-HOXB7 cells formed tumors

when subcutaneously injected into immunodeficient mice (Figure 4A). Tumor formation was similarly

observed upon orthotopic transplantation. Primary tumors of fat pads were formed in half the mice

within one week and eleven out of twelve mice within ten days (Figure 4B). They continued to grow

until sixteen weeks after transplantation (Figure 4C), confirming that *HOXB7* induces oncogenic

activities *in vivo*. We next examined the metastatic activity of NMuMG-HOXB7 cells *in vivo* and *ex*

vivo. Using *in vivo* imaging, bioluminescence from the lung was detected in one mouse out of twelve,

sixteen weeks after transplantation (Figure 4D). To detect lung metastasis with more sensitivity, we

excised the lungs and performed *ex vivo* bioluminescence imaging. As a result, lung metastasis was

detected in four out of twelve mice (Figure 4E). Our results indicate that *HOXB7* induces not only *in*

vivo tumor formation but also lung metastasis in an orthotropic transplantation model.

4. Discussion

In this study, we demonstrated that *HOXB7* induces cellular transformation by activating

the JAK-STAT pathway, which suggests its involvement in determining the poor prognosis of

HER2-positive patients. Additionally, overexpression of *HOXB7* in NMuMG cells transplanted into

mouse models induced the transition from tumorigenesis to metastasis.

HOX family genes regulate development and homeostasis as transcriptional factors

[10,11,25]. Several HOX genes contribute to cancer malignancy [10,11]. Overexpression and

knockdown experiments together support the idea that *HOXB7* is acting as an oncogene in breast,

prostate, hepatocellular and gastric cancer cell lines [12-14]. HOXB7 also induces EMT and

migration/invasion activity in MCF10A normal human mammary epithelial cells [15]. *In vivo*, *HOXB7*

was also shown to promotes malignant progression through the activation of TGFß signaling in a

MMTV-Hoxb7/Her2 transgenic mouse model [16]. In addition, *HOXB7* overexpression in MCF7 cells

(luminal A subtype) induces acquired resistance to tamoxifen via the activation of receptor tyrosine

kinases and ERα signaling [17,18]. Previous analysis of microarray data also showed that high levels

of HOXB7 are associated with a poor prognosis in HER-2–positive, but not HER-2–negative breast

cancers [26]. Consistently, our analysis with recent genomic and RNA-seq data confirmed their study

and further demonstrated that *HOXB7* is significantly co-amplified with *ERBB2*. Another study

showed that *HOXB7* expression induces upregulation of *ERBB2* in MCF7 cells [18], a finding which

was also confirmed by our experiments using different cell lines. Taken together, these results

indicate that coamplification of *HOXB7* with *ERBB2* may enhance the expression of *ERBB2* to

enhance the malignancy of HER2-positive breast cancer cells.

Using an immortalized mammary gland epithelial cell line, NMuMG, *HOXB7* was shown to

induce cellular transformation *in vitro* and tumor formation *in vivo*. In this cellular context, *HOXB7*

increased the mRNAs of *STAT3* and protein levels of JAK2 and STAT3 as well as phosphorylated

STAT3. Intriguingly, treatment with Ruxolitinib, a JAK1/2 inhibitor, abolished cellular transformation

with inhibition of Stat3 phosphorylation, suggesting involvement of the JAK-STAT pathway in this

HOXB7-mediated transformation. As observed previously [27], Jak2 phosphorylation was found to

be upregulated by Ruxolitinib treatment. This phenomenon was also observed following treatment

with Pacritinib, another JAK2 inhibitor [28], but the molecular mechanism underlying the upregulation

of phosphorylated Jak2 remains unclear. The JAK-STAT pathway is activated in several cancers and

is considered to be involved in tumorigenesis [5]. Kinase inhibitors against JAK including Ruxolitinib

were thus developed. Even though *HOXB7* cannot be directly targeted, the downstream JAK-STAT

pathway could be targeted in cancers associated with *HOXB7* alterations [8].

A previous study revealed that *HOXB7* is not sufficient to induce tumors in *MMTV-Hoxb7*

transgenic mice [26]. By contrast, we showed that *HOXB7* alone triggers tumorigenesis and lung

metastasis using *HOXB7*-expressing NMuMG cells. While this discrepancy remains unclear, the

immortalization status of NMuMG cells may be involved. In any case, our observation that *HOXB7*

enhances lung metastasis in an orthotropic mouse model underpins the importance of *HOXB7* in

lung metastasis, consistently with a previous study of M*MTV-Hoxb7/Erbb2* transgenic mice [26]. The

transplantation models using NMuMG-HOXB7 cells are expected to be useful in discovering the role

of HOXB7 in cancer initiation and metastasis. Furthermore, a combined therapy against HOXB7 or

downstream JAK-STAT together with ERBB2 may lead to efficient therapeutic outcomes in the

"*ERBB2*- and *HOXB7*-amplified" subtype of breast cancer.

Acknowledgments

We thank Dr. Yusuke Yamamoto (National Cancer Center Research Institute) and Dr. Jiro Fujimoto (Japan Biological Informatics Consortium; JBic) for their meaningful comments and discussion, Prof. K. Miyazawa (University of Yamanashi, Yamanashi, Japan) for kindly providing the NMuMG cell line, and Prof. T. Kitamura (Institute of Medical Science, The University of Tokyo, Tokyo, Japan) for kindly providing the Plat-E cell line. This study was conducted in part through the Fukushima Translational Research Project program.

Funding

This work was supported by JSPS KAKENHI [grant no. 23241064, Grant-in-Aid for Scientific Research (A) to K.S.; grant no. 18K16269, Grant-in-Aid for Early-Career Scientist to J.N.; grant no. 20J01794, Grant-in-Aid for JSPS Fellows to J.N.] and the JBic.

Authors' contributions

J.N. and K.S. conceived and designed the study. K.A, M.S., S.K., J.N., and A.M. performed the experiments. K.A., M.S. and J.N. analyzed the data. N.G., S.W. and K.S. interpreted the data. J.N. wrote the manuscript. All the authors reviewed and edited the manuscript.

Conflict of interest

The authors declare no conflict of interest.

References

[1] D. Hanahan, R.A. Weinberg, Hallmarks of cancer: the next generation, Cell 144 (2011) 646-674. 10.1016/j.cell.2011.02.013.

[2] A. Matsui, J. Fujimoto, K. Ishikawa, E. Ito, N. Goshima, S. Watanabe, K. Semba, Hepatocyte nuclear factor 1 beta induces transformation and epithelial-to-mesenchymal transition, FEBS Lett 590 (2016) 1211-1221. 10.1002/1873-3468.12147.

[3] M. Saito, Y. Kato, E. Ito, J. Fujimoto, K. Ishikawa, A. Doi, K. Kumazawa, A. Matsui, S. Takebe, T. Ishida, S. Azuma, H. Mochizuki, Y. Kawamura, Y. Yanagisawa, R. Honma, J. Imai, H. Ohbayashi, N. Goshima, K. Semba, S. Watanabe, Expression screening of 17q12-21 amplicon reveals GRB7 as an ERBB2-dependent oncogene, FEBS Lett 586 (2012) 1708-1714. 10.1016/j.febslet.2012.05.003.

[4] W. Kolch, M. Halasz, M. Granovskaya, B.N. Kholodenko, The dynamic control of signal transduction networks in cancer cells, Nat Rev Cancer 15 (2015) 515-527. 10.1038/nrc3983.

[5] X. Hu, J. li, M. Fu, X. Zhao, W. Wang, The JAK/STAT signaling pathway: from bench to clinic, Signal Transduction and Targeted Therapy 6 (2021) 402. 10.1038/s41392-021-00791-1.

[6] L.L. Marotta, V. Almendro, A. Marusyk, M. Shipitsin, J. Schemme, S.R. Walker, N. Bloushtain-Qimron, J.J. Kim, S.A. Choudhury, R. Maruyama, Z. Wu, M. Gönen, L.A. Mulvey, M.O. Bessarabova, S.J. Huh, S.J. Silver, S.Y. Kim, S.Y. Park, H.E. Lee, K.S. Anderson, A.L. Richardson, T. Nikolskaya, Y. Nikolsky, X.S. Liu, D.E. Root, W.C. Hahn, D.A. Frank, K. Polyak, The JAK2/STAT3 signaling pathway is required for growth of CD44$^+$CD24$^-$ stem cell-like breast cancer cells in human tumors, J Clin Invest 121 (2011) 2723-2735. 10.1172/jci44745.

[7] T. Wang, J.F. Fahrmann, H. Lee, Y.J. Li, S.C. Tripathi, C. Yue, C. Zhang, V. Lifshitz, J. Song, Y. Yuan, G. Somlo, R. Jandial, D. Ann, S. Hanash, R. Jove, H. Yu, JAK/STAT3-Regulated Fatty Acid β-Oxidation Is Critical for Breast Cancer Stem Cell Self-Renewal and Chemoresistance, Cell Metab 27 (2018) 136-150.e135. 10.1016/j.cmet.2017.11.001.

[8] Z. Qureshy, D.E. Johnson, J.R. Grandis, Targeting the JAK/STAT pathway in solid tumors, Journal of Cancer Metastasis and Treatment 6 (2020) 27. 10.20517/2394-4722.2020.58.

[9] H. Fazilaty, P. Mehdipour, Genetics of breast cancer bone metastasis: a sequential multistep pattern, Clinical & Experimental Metastasis 31 (2014) 595-612. 10.1007/s10585-014-9642-9.

[10] S. Bhatlekar, J.Z. Fields, B.M. Boman, HOX genes and their role in the development of human cancers, J Mol Med (Berl) 92 (2014) 811-823. 10.1007/s00109-014-1181-y.

[11] B. Li, Q. Huang, G.H. Wei, The Role of HOX Transcription Factors in Cancer Predisposition and Progression, Cancers (Basel) 11 (2019). 10.3390/cancers11040528.

[12] W.M. Wang, Y. Xu, Y.H. Wang, H.X. Sun, Y.F. Sun, Y.F. He, Q.F. Zhu, B. Hu, X. Zhang, J.L. Xia, S.J. Qiu, J. Zhou, X.R. Yang, J. Fan, HOXB7 promotes tumor progression via bFGF-induced activation of MAPK/ERK pathway and indicated poor prognosis in hepatocellular carcinoma, Oncotarget 8 (2017) 47121-47135. 10.18632/oncotarget.17004.

[13] X. He, Z. Liu, Y. Xia, J. Xu, G. Lv, L. Wang, T. Ma, L. Jiang, Y. Mou, X. Jiang, J. Ma, Z. Zhao, H. Ni, W. Xu, G. Ru, D. Huang, H. Tao, HOXB7 overexpression promotes cell proliferation and correlates with poor prognosis in gastric cancer patients by inducing expression of both AKT and MARKs, Oncotarget 8 (2017) 1247-1261. 10.18632/oncotarget.13604.

[14] M.C. Errico, K. Jin, S. Sukumar, A. Carè, The Widening Sphere of Influence of HOXB7 in Solid Tumors, Cancer Res 76 (2016) 2857-2862. 10.1158/0008-5472.Can-15-3444.

[15] X. Wu, H. Chen, B. Parker, E. Rubin, T. Zhu, J.S. Lee, P. Argani, S. Sukumar, HOXB7, a
homeodomain protein, is overexpressed in breast cancer and confers epithelial-mesenchymal
transition, Cancer Res 66 (2006) 9527-9534. 10.1158/0008-5472.Can-05-4470.

[16] S. Liu, K. Jin, Y. Hui, J. Fu, C. Jie, S. Feng, D. Reisman, Q. Wang, D. Fan, S. Sukumar, H. Chen,
HOXB7 promotes malignant progression by activating the TGFβ signaling pathway, Cancer Res 75
(2015) 709-719. 10.1158/0008-5472.Can-14-3100.

[17] K. Jin, X. Kong, T. Shah, M.F. Penet, F. Wildes, D.C. Sgroi, X.J. Ma, Y. Huang, A. Kallioniemi, G.
Landberg, I. Bieche, X. Wu, P.E. Lobie, N.E. Davidson, Z.M. Bhujwalla, T. Zhu, S. Sukumar, The
HOXB7 protein renders breast cancer cells resistant to tamoxifen through activation of the EGFR
pathway, Proc Natl Acad Sci U S A 109 (2012) 2736-2741. 10.1073/pnas.1018859108.

[18] K. Jin, S. Park, W.W. Teo, P. Korangath, S.S. Cho, T. Yoshida, B. Győrffy, C.P. Goswami, H.
Nakshatri, L.A. Cruz, W. Zhou, H. Ji, Y. Su, M. Ekram, Z. Wu, T. Zhu, K. Polyak, S. Sukumar, HOXB7
Is an ERα Cofactor in the Activation of HER2 and Multiple ER Target Genes Leading to Endocrine
Resistance, Cancer Discov 5 (2015) 944-959. 10.1158/2159-8290.Cd-15-0090.

[19] T. Ihara, Y. Hosokawa, K. Kumazawa, K. Ishikawa, J. Fujimoto, M. Yamamoto, T. Muramkami, N.
Goshima, E. Ito, S. Watanabe, K. Semba, An in vivo screening system to identify tumorigenic genes,
Oncogene 36 (2017) 2023-2029. 10.1038/onc.2016.351.

[20] Y. Kuroiwa, J. Nakayama, C. Adachi, T. Inoue, S. Watanabe, K. Semba, Proliferative
Classification of Intracranially Injected HER2-positive Breast Cancer Cell Lines, Cancers (Basel) 12
(2020) 1811. 10.3390/cancers12071811.

[21] J. Nakayama, E. Ito, J. Fujimoto, S. Watanabe, K. Semba, Comparative analysis of gene
regulatory networks of highly metastatic breast cancer cells established by orthotopic transplantation
and intra-circulation injection, Int J Oncol 50 (2017) 497-504. 10.3892/ijo.2016.3809.

[22] E. Cerami, J. Gao, U. Dogrusoz, B.E. Gross, S.O. Sumer, B.A. Aksoy, A. Jacobsen, C.J. Byrne,
M.L. Heuer, E. Larsson, Y. Antipin, B. Reva, A.P. Goldberg, C. Sander, N. Schultz, The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data, Cancer
Discov 2 (2012) 401-404. 10.1158/2159-8290.Cd-12-0095.

[23] B. Pereira, S.-F. Chin, O.M. Rueda, H.-K.M. Vollan, E. Provenzano, H.A. Bardwell, M. Pugh, L.
Jones, R. Russell, S.-J. Sammut, D.W.Y. Tsui, B. Liu, S.-J. Dawson, J. Abraham, H. Northen, J.F.
Peden, A. Mukherjee, G. Turashvili, A.R. Green, S. McKinney, A. Oloumi, S. Shah, N. Rosenfeld, L.
Murphy, D.R. Bentley, I.O. Ellis, A. Purushotham, S.E. Pinder, A.-L. Børresen-Dale, H.M. Earl, P.D.
Pharoah, M.T. Ross, S. Aparicio, C. Caldas, The somatic mutation profiles of 2,433 breast cancers
refine their genomic and transcriptomic landscapes, Nat Commun 7 (2016) 11479.
10.1038/ncomms11479.

[24] Y. Han, K. Azuma, S. Watanabe, K. Semba, J. Nakayama, Metastatic profiling of HER2-positive
breast cancer cell lines in xenograft models, Clinical & Experimental Metastasis (2022).
10.1007/s10585-022-10150-1.

[25] R. Rezsohazy, A.J. Saurin, C. Maurel-Zaffran, Y. Graba, Cellular and molecular insights into Hox
protein action, Development 142 (2015) 1212-1227. 10.1242/dev.109785.

[26] H. Chen, J.S. Lee, X. Liang, H. Zhang, T. Zhu, Z. Zhang, M.E. Taylor, C. Zahnow, L. Feigenbaum,
A. Rein, S. Sukumar, Hoxb7 Inhibits Transgenic HER-2/neu–Induced Mouse Mammary Tumor Onset
but Promotes Progression and Lung Metastasis, Cancer Research 68 (2008) 3637-3644.

10.1158/0008-5472.Can-07-2926.

[27] S. Lee, T. Shah, C. Yin, J. Hochberg, J. Ayello, E. Morris, C. van de Ven, M.S. Cairo, Ruxolitinib

significantly enhances in vitro apoptosis in Hodgkin lymphoma and primary mediastinal B-cell

lymphoma and survival in a lymphoma xenograft murine model, Oncotarget 9 (2018) 9776-9788.

10.18632/oncotarget.24267.

[28] J.R. Hart, L. Liao, J.R. Yates, 3rd, P.K. Vogt, Essential role of Stat3 in PI3K-induced oncogenic

transformation, Proc Natl Acad Sci U S A 108 (2011) 13247-13252. 10.1073/pnas.1110486108.

Figure legends

Figure 1. High expression of HOXB7 is correlated to a poor prognosis in breast cancer

patients. (A) OncoPrint visualization of *ERBB2* and *HOXB7* genomic status in TCGA. (B) Altered

genetic profiles of ERBB2 and HOXB7. Fisher's exact test ($p < 0.05$). (C) Comparative expression

analysis of *HOXB7* in breast cancer subtypes of TCGA. Luminal A (n = 499), luminal B (n = 197),

HER2-positive (n = 78), basal-like (n = 171), and normal-like (n = 36) subtypes. Welch's t-test ($p <$

0.005). (D left) Survival analysis of all METABRIC cohorts with *HOXB7* expression and the overall

survival status shown in a Kaplan-Meier plot. Shading along the curves shows 95% confidence

interval. Log-rank test ($p = 0.0056$) (D right) METABRIC cohorts with high HER2 expression (476

patients; Top 25% of all METABRIC cohorts) Log-rank test ($p = 0.0088$) (E) Survival analysis of other

breast cancer cohorts (GSE1456; disease specific survival; GSE9893; overall survival). GSE1456,

log-rank test ($p = 0.00022$); GSE9893, log-rank test ($p < 0.0001$).

Figure 2. HOXB7 expression induces transformation and activation of the JAK-STAT and

ERBB2 signaling pathways. (A) Brightfield imaging of HOXB7-expressing cells. (B) Focus

formation assay in HOXB7-expressing NMuMG cells. Left: crystal violet staining. Right: brightfield

focus. (C) Western blotting analysis of HOXB7-expressing NMuMG cells. Venus-expressing NMuMG

cells were used as a negative control. (D) Western blotting analysis of HOXB7-expressing cells. (E)

qRT-PCR of *Stat3*, *Erbb2*, and *Jak2*. Welch's t-test (* $p < 0.001$, n = 3).

Figure 3. Focus formation by HOXB7 is regulated by Stat3 activation. (A) Western blotting of

p-Stat3 analysis following Ruxolitinib treatment. (B) Signal analysis following Ruxolitinib treatment of

HOXB7-expressing NMuMG cells. (C) Signal analysis following Lapatinib treatment of

HOXB7-expressing NMuMG cells. (D) *In vitro* growth curve of HOXB7-expressing NMuMG cells

treated with Ruxolitinib. (n.s.: Not significant, n = 3) (E) Focus formation assay of HOXB7-expressing

NMuMG cells treated with Ruxolitinib. Foci were stained with crystal violet.

Figure 4. Overexpression of HOXB7 induces tumorigenesis and lung metastasis. (A)

Tumorigenesis assay of HOXB7-expressing NMuMG cells. Left: Tumor growth curve with volume.

HOXB7-expressing NMuMG cells formed tumors following subcutaneous transplantation in *Rag2*$^{-/-}$

mice (n = 10). Right: Image of HOXB7-positve tumor. (B) Primary tumor formation curve for

orthotopic transplantation. HOXB7-expressiong NMuMG cells were transplanted to BALB/cSlc-nu/nu

mice (n = 12). (C) Representative *in vivo* luminescence images of primary tumor in orthotopic

transplantation. (D) Representative *in vivo* luminescence image of lung metastasis in orthotopic

transplantation. (E) *Ex vivo* luminescence images of lungs excised from orthotopically transplanted

mice. Left: Lungs with no metastasis. Right: Lungs with metastasis.

Figure 1

Figure 2

Figure 3

Figure 4

A Venus / HOXB7 — Tumor Volume (mm³) vs Time(Weeks)

B Tumor formation at fatpad
Cells: NMuMG-HOXB7
Tumor formation ratio (%) vs Days

C Primary tumor — Week 0, Week 8, Week 16
Radiance (p/sec/cm²/sr)

D Lung metastasis
Radiance (p/sec/cm²/sr)

E Lung metastasis − / Lung metastasis +
Radiance (p/sec/cm²/sr)