1 *HOXB7* **induces STAT3-mediated transformation and lung metastasis in immortalized**
2 **mammary gland NMuMG cells**
3

4 Kazushi Azuma[a, †], Mai Sakamoto[a,b, †], Shota Katayama[a, †], Atsuka Matsui[a], Kazuya Nakamichi[a],
5 Naoki Goshima[c,d,e], Shinya Watanabe[f], Jun Nakayama[a,b,g,*], Kentaro Semba[a,f,*]
6
[a] Department of Life Science and Medical Bioscience, School of Advanced Science and Engineering,
Waseda University, Tokyo, Japan.
[b] Computational Bio-Big Data Open Innovation Lab. (CBBD-OIL), National Institute of Advanced
Industrial Science and Technology (AIST), Tokyo, Japan.
[c] Division of Transcriptome Analysis, Translational Research Center, Fukushima Medical University,
Fukushima, Japan.
[d] Functional Proteomics Team, Molecular Profiling Research Center for Drug Discovery (molprof),
National Institute of Advanced Industrial Science and Technology (AIST), Tokyo, Japan.
[e] Department of Human Sciences, Musashino University, Tokyo, Japan.
[f] Translational Research Center, Fukushima Medical University, Fukushima, Japan.
[g] Laboratory of Integrative Oncology, National Cancer Center Research Institute, Tokyo, Japan.
18
19 [†] These authors equality contributed
20 [*] Corresponding authors
21
22
23 **Correspondence to:** Jun Nakayama (junakaya@ncc.go.jp or jnakayama.re@gmail.com) and
24 Kentaro Semba (ksemba@waseda.jp), Department of Life Science and Medical Bioscience, School
25 of Advanced Science and Engineering, Waseda University, TWIns, 2-2 Wakamatsu-cho, Shinjuku-ku,
26 Tokyo, 162-8480, Japan, Tel/Fax: +81-3-5369-7320.

27
28 **Running Head:** HOXB7 induces oncogenic transformation and metastasis.
29
30 **Keywords:** HOXB7, oncogene, breast cancer, metastasis, ERBB2, NmuMG
31
Abbreviations
HOXB7, homeobox B7: NMuMG, normal murine mammary gland: HER2, human epidermal growth
factor receptor 2: TCGA, the cancer genome atlas: TGFß, transforming growth factor beta: EMT,
epithelial to mesenchymal transition: DMEM, Dulbecco's modified Eagle's medium: FBS, fetal bovine
serum: IRES, internal ribosome entry site: PEI, polyethylenimine: METABRIC, the molecular

taxonomy of breast cancer international consortium: IVIS, in vivo imaging system: SOCS,
Suppressor of cytokine signaling.
39

Abstract:

The homeobox family genes are often dysregulated in a various cancer type. Particularly *HOXB7*
amplification and overexpression correlate with poor prognosis in various cancer such as gastric,
pancreatic, and lung cancers. Moreover, *HOXB7* is known to contribute to cancer progression by
promoting epithelial to mesenchymal transition, anti-cancer drug resistance, and angiogenesis. In
this study, we show that *HOXB7* is coamplified with *ERBB2* in a subset of breast cancer patients and
HOXB7 expression correlates with poor prognosis in HER2-positive breast cancer patients. This
clinical observation is supported by the following results: *HOXB7* overexpression in an immortalized
murine mammary gland epithelial cell line NMuMG induces cellular transformation *in vitro*,
tumorigenesis and lung metastasis through the activation of JAK-STAT signaling.

50
51

1. Introduction

Oncogenic alterations, such as gene amplifications, mutations, chromosomal translocation, and epigenetic dysregulation, cause tumorigenesis *in vivo* (Hanahan & Weinberg 2011). Activated oncogenes also induce typical phenotypic changes, so-called transformation, in immortalized normal cells such as a NIH3T3 and normal murine mammary gland (NMuMG) cells *in vitro*. Therefore, assay systems evaluating such phenotypic changes, namely focus formation, soft agar colonization, and tumorigenesis have been used for the screening of novel oncogenes in previous studies including ours (Saito *et al.* 2012; Matsui *et al.* 2016).

Most oncogenes cause dysregulation of the signal transduction cascades that relate to proliferation, cell cycle, and survival (Kolch *et al.* 2015). Typically, oncogenic signaling pathways, such as RAS-MAPK and PI3K-AKT are often activated in various cancer cell types and thus molecular targeted-drugs against them have been successfully developed. Recent studies showed that the JAK-STAT signaling is also important in cancer initiation and progression (Hu *et al.* 2021). STAT3 is activated in cancer stem cell-like populations (Marotta *et al.* 2011), and regulates stemness via fatty metabolism in breast cancer (Wang *et al.* 2018). In addition, aberrant activation of STAT3 with IL6-enriched tumor microenvironment is associated with an immunosuppression of tumor-infiltrating immune cells (Qureshy *et al.* 2020). JAK-STAT signaling also coactivates and supports cancer growth and progression with the activation of RAS-MAPK and PI3K-AKT pathways (Fazilaty & Mehdipour 2014).

The homeobox (HOX) genes encode transcriptional factors, that controls development and organ homeostasis. The HOX superfamily of homeobox genes (HOXA, HOXB, HOXC and HOXD clusters) are often dysregulated in breast, prostate, colon, and lung cancer. The differential dysregulation of HOX family genes in various solid tumors provides an opportunity to understand cancer development (Bhatlekar *et al.* 2014; Li *et al.* 2019). The HOXB clusters is composed of *HOXB1*, *HOXB2*, *HOXB3*, *HOXB4*, *HOXB6*, *HOXB7*, *HOXB8*, *HOXB9*, and *HOXB13* at 17q23.32. HOXB cluster genes are co-amplified by gene amplification in several types of cancer. Notably, overexpression of *HOXB7* contributes to cancer progression in breast, hepatocellular, oral, and gastric cancer cell lines (Errico *et al.* 2016; He *et al.* 2017; Wang *et al.* 2017). *HOXB7* induces the activation of TGFß signaling, epithelial to mesenchymal transition (EMT) and migration in breast cancer (Wu *et al.* 2006; Liu *et al.* 2015). In addition, the luminal subtype breast cancer cell line MCF7 acquires tamoxifen resistance via *HOXB7* expression (Jin *et al.* 2012). *HOXB7* as a cofactor of estrogen receptor (ER) causes cancer progression through the expression of downstream genes (Jin *et al.* 2015). In this study, we revealed HOXB7-mediated enhancement of JAK-STAT pathway and its possible involvement in cellular transformation *in vitro*. We also found that HOXB7 is enough to induce tumorigenesis and metastasis *in vivo* in the context of immortalized NMuMG cells.

2. Materials and methods

2.1. Cell culture

Normal murine mammary gland (NMuMG) and NMuMG-luc cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Fujifilm Wako, Osaka Japan) supplemented with 10% fetal bovine serum (FBS; NICHIREI BIOSCIENCES INC., Tokyo, Japan), 100 U/mL penicillin G (Meiji-Seika Pharma Co., Ltd., Tokyo, Japan), 100 µg/mL streptomycin (Meiji-Seika Pharma Co., Ltd.), 10 µg/mL insulin (Fujifilm Wako), and 0.45% glucose (Fujifilm Wako) at 37°C with 5% CO_2. Plat-E cells for retroviral packaging were cultured in DMEM supplemented with 10% FBS, 100 U/mL penicillin G, and 100 µg/mL streptomycin at 37°C with 5% CO_2. Cells were treated with 0.1-1 µM ruxolitinib (LC Laboratories, MA, USA) under experimental conditions.

2.2. Retroviral packaging and infection

To establish NMuMG-Venus, *HOXB7*-expressing NMuMG (NMuMG-HOXB7), *HOXB7*-expressing NMuMG-luc (NMuMG-luc-HOXB7) cell lines by using retroviral infection, Plat-E cells were seeded into 6-well plates and transfected with pMXs-Venus-IRES-Puromycin^R and pMXs-*HOXB7*-IRES-Puromycin^R plasmid DNAs using 6 µg polyethylenimine (PEI; Polysciences, Inc., PA, USA). Then, the cell supernatants and 800 µg/mL polybrene (SIGMA ALDRICH Co., MO, USA), as previously described (Matsui *et al.* 2016). To select the expressing cells, 1 µg/mL puromycin (Fujifilm Wako) was added to the culture medium for 2 days.

To establish NMuMG-luc-HOXB7-shGFP, NMuMG-luc-HOXB7-shSTAT3#2 and NMuMG-luc-HOXB7-shSTAT3#3 cell lines by using retroviral infection, pSuper-shGFP-Hygromycin^R, pSuper-shSTAT3#2-Hygromycin^R and pSuper-shSTAT3#3-Hygromycin^R plasmid DNAs were used with the method described above. To select the expressing cells, 800 µg/mL hygromycin was added to the culture medium for 5 days. The following pairs of oligonucleotides were used to target specified genes: shGFP, ACAACAGCCACAACGTCTATAGTGTGCTGTCCTATAGACGTTGTGGCTGTTGTTTTTTT and AAAAAAACAACAGCCACAACGTCTATAGGACAGCACACTATAGACGTTGTGGCTGTTGT; shSTAT3#2, CCTGGGTTGAATTGTCAGTTTGTGTGCTGTCCAAGCTGATAATTCAACTCAGGTTTT TGGAA and TTCCAAAAACCTGAGTTGAATTATCAGCTTGGACAGCACACAAACTGACAATTCAAC CCAGG; shSTAT3#3, CGACTTTGGTTTCAATTACGAGTGTGCTGTCCTTGTAGTTGAAATCAAAGT CGTTTTTGGAA and TTCCAAAAACGACTTTGATTTCAACTACAAGGACAGCACACTCGTAATTGA AACCAAAGTCG.

2.3. Western blotting

The infected cell lines were lysed with SDS sampling buffer (3.5 % SDS, 12% glycerol, 0.05M Tris-HCl (pH 6.8), 0.003% bromo phenol blue, and 5% 2-mercaptoethanol) for western blotting.

Western blotting was performed as previously described (Matsui *et al.* 2016) with the antibodies
shown in Supplementary Table S1. The cells were treated with 1μM ruxolitinib for 24 hours before
harvesting.

2.4. Cell proliferation assay
A total of 2.5×10^4 of NMuMG-Venus cells or NMuMG-HOXB7 cells were seeded into a 12-well plate.
Cells were harvested with 0.25% trypsin (Thermo Fisher Scientific, MA, USA) and counted at each
time point (24, 48 and 72 h) (Kuroiwa *et al.* 2020). Cells were continuously treated with ruxolitinib
from 24 hours after cell seeding.

2.5. Focus formation assay
For focus formation assay, 7.0×10^4 parental NMuMG cells and 3.5×10^4 NMuMG-HOXB7 cells
were mixed and seeded into 12-well plates. In STAT3 knock down experiment, 1.5×10^5 of parental
NMuMG cells and 7.5×10^2 of NMuMG-luc-HOXB7 cell lines (NMuMG-luc-HOXB7-shGFP cells or
NMuMG-luc-HOXB7-shSTAT3#3 cells) were mixed and seeded into 12-well plates. Cells were
stained with 0.01% crystal violet/70% methanol to detect the cellular foci (Matsui *et al.* 2016). Cells
were continuously treated with ruxolitinib from 24 hours after cell seeding. Photos were taken at the
indicated day described in each figure legend.

2.6. Tumorigenesis assay
A total of 1.0×10^6 NMuMG-Venus cells and NMuMG-HOXB7 cells were transplanted into
6-8-weeks-old *Rag2*$^{-/-}$ female mice with 100 μL 50% Matrigel (BD Bioscience, CA, USA) as
previously described (Ihara *et al.* 2017). All animal experiments were approved by the Animal
Committee of Waseda University (WD19-058, 2019-A068).

2.7. Orthotopic Transplantation and bioluminescence imaging
A total of 1.0×10^6 NMuMG-luc-HOXB7 cells, NMuMG-luc-HOXB7-shGFP cells, or
NMuMG-luc-HOXB7-shSTAT3#3 cells were transplanted into fourth fat pads of 6-weeks-old
BALB/cSlc-nu/nu female mice as previously described (Nakayama *et al.* 2017; Nakayama *et al.*
2022). After the injection, the transplantation sites were palpated for every 2-3 days to measure
tumor formation time and mice were monitored primary tumor formation and metastasis formation by
in vivo imaging system (IVIS Lumina Xr3; Perkin-elmer, Waltham, MA, USA) for every week. Mice
were intraperitonially injected with 200 μL D-Luciferin (15 mg/mL; Gold Biotechnology, Inc., St. Luis,
MO, USA), and Luminescence was taken under anesthesia using IVIS. Sixteen weeks after injection,
all mice were sacrificed, and excised lungs. Lungs were transferred to a 12-well plate with 900 μL of
D-PBS (-) (Fujifilm Wako Pure Chemical Industries) and 100 μL of D-Luciferin (15 mg/mL), and then
luminescence was taken by IVIS, as previously described (Han *et al.* 2020). All animal experiments

were approved by the Animal Committee of Waseda University (WD21-082, 2021-A074)

2.8. Quantitative real-time PCR

Total RNA was extracted using ISOGEN (NIPPON GENE Co., LTD, Tokyo, Japan). cDNA synthesis

was performed using the Superscript III first-strand synthesis system (Thermo Fisher Scientific).

Quantitative RT-PCR was performed using THUNDER BIRD SYBR qPCR mix (TOYOBO, Osaka,

Japan) and 18S rRNA was used as a relative control. Primers are listed in Supplementary Table S2.

2.9. Clinical database analysis

The genomic profiles and RNA-seq datasets of The Cancer Genome Atlas (TCGA) breast cancer

(BRCA) cohorts were subjected to OncoPrint and expression analyses. OncoPrint of *HOXB7* and

ERBB2 was conducted using cBioPortal (Cerami *et al.* 2012). The Kaplan-Meier method was used

perform survival analyses of the molecular taxonomy of breast cancer international consortium

(METABRIC) (Pereira *et al.* 2016), GSE1456 and GSE9893 data sets, as previously described (Han

et al. 2022).

2.10. Statistical analysis

Welch's t-test, Fisher's exact test and log-rank test were performed using R 3.4.1

(https://www.r-project.org/).

3. Results

3.1. Expression of *HOXB7* correlated to poor prognosis in the HER2-positive breast cancer

patients.

To elucidate the clinical significance of *HOXB7* in breast cancer, we first analyzed RNA-seq data of

TCGA BRCA cohorts for *HOXB7* expression. Gene expression analysis in each breast cancer

subtype revealed that the HER2-positive subtype significantly highly expressed *HOXB7* compared

with the other subtypes and normal tissue (Figure 1a). To examine the genomic profiles of *HOXB7* in

TCGA BRCA cohorts, the mutational status of *ERBB2* (the gene that encodes HER2) and *HOXB7*

was illustrated as an OncoPrint by cBioPortal (Figure 1b). *HOXB7* was significantly co-amplified with

ERBB2 in 30 patients out of 48 patients with *HOXB7* amplification (Figure 1c). HOXB7 expression

also significantly increased in the *ERBB2/HOXB7*-amplified patients compared with the

ERBB2-amplified patients (Figure 1d). These results suggested that *HOXB7* was preferentially

amplified and expressed in HER2-positive breast cancer patients.

Next, to examine the relationship between *HOXB7* expression and progression in breast

cancer, we performed a survival analysis using several public clinical data sets. In METABRIC

cohorts, high expression of *HOXB7* was correlated with poor prognosis in all breast cancer patients (Figure 1e left) and especially in HER2-high patients (Figure 1e right). The correlation of poor prognosis and *HOXB7* expression was further confirmed by an analysis of other public breast cancer cohorts that included all subtypes (Figure 1f).

3.2. *HOXB7* induced oncogenic focus formation and activation of JAK-STAT3 signaling

To elucidate the oncogenic function of *HOXB7*, we first established NMuMG-HOXB7 cells that expressed *HOXB7* in an immortalized murine breast epithelial cell line NMuMG as a model system for both *in vitro* and *in vivo* evaluation of oncogenes (Matsui *et al.* 2016; Ihara *et al.* 2017). Expression of HOXB7 changed morphology of NMuMG cells from epithelial-like to spindle-like shape (Figure 2a). Furthermore NMuMG-HOXB7 cells formed foci (Figure 2b).

Next, to identify oncogenic signaling pathways of *HOXB7*, we performed western blotting using antibodies against major cancer-related signaling proteins. Among them, we detected the *HOXB7* overexpression increased the protein level of vimentin (a mesenchymal marker gene) (Figure 2c), but did not change those of E-cadherin and N-cadherin, suggesting *HOXB7* induces partial epithelial-mesenchymal transition. Notably, *HOXB7* overexpression drastically increased Stat3 phosphorylation and protein levels of Jak1, Jak2, and Stat3 (Figure 2d). Major proliferative signals, such as Erk and Akt, were not changed by *HOXB7* expression. Stat3 was increased at the transcriptional level, while Jak2 was increased at the protein level (Figure 2e).

Additionally, phosphorylation of Stat5 was also increased by *HOXB7* expression (Figure 3a). Phosphorylation of Stat3 and Stat5 induced by *HOXB7* expression was completely inhibited by a JAK1/2 inhibitor, ruxolitinib in a dose-dependent manner confirming JAK1/2 involvement in Stat3 and Stat5 phosphorylation (Figure 3a). Next, to elucidate the contribution of the oncogenic activity of p-Stat3, we performed *in vitro* proliferation assays and focus formation assays following ruxolitinib treatment. *HOXB7* decreased the proliferative activity of NMuMG cells *in vitro*, however, ruxolitinib treatment did not affect the proliferative activity of NMuMG-HOXB7 cells (Figure 3b). Nevertheless, ruxolitinib inhibited their focus-forming activity (Figure 3c). These results suggest that the activation of JAK-STAT signaling by *HOXB7* induced cellular transformation.

3.3. Overexpression of *HOXB7* induced tumorigenesis and metastasis.

To examine the oncogenic activities *in vivo*, we performed the subcutaneous tumorigenesis assay and the metastasis assay by orthotopic transplantation. As a result, NMuMG-HOXB7 cells formed tumors when subcutaneously injected into immunodeficient mice (Figure 4a). Tumor formation was also observed in orthotopic transplantation. Primary tumors at fatpad were formed in half of mice within one week and in almost all mice within ten days (Figure 4b). They continued to grow until sixteen weeks after transplantation (Figure 4c), suggesting that *HOXB7* induced oncogenic activities *in vivo*. Next, we examine the metastatic activity of NMuMG-luc-HOXB7 cells that express luciferase

gene with *HOXB7* gene. As a result of *in vivo* bioluminescence imaging with higher sensitivity,
bioluminescence from the lung was detected in one mouse of twelve at sixteen weeks after
transplantation (Figure 4d). To detect lung metastasis with more sensitivity, we excised the lungs and
performed *ex vivo* bioluminescence imaging. As a result, lung metastasis was detected in 4 of 12
mice (Figure 4e). Our results showed that *HOXB7* induced not only *in vivo* tumor formation but also
lung metastasis in orthotropic transplantation model.

To examine the contribution of STAT3 to *HOXB7*-mediated tumorigenicity *in vivo*, we
performed the knock down (KD) of STAT3 in NMuMG-luc-HOXB7 cells. STAT3 KD cells were
established using shRNA with retroviral vector (Figure 5a). As a result, STAT3 KD using shSTAT3#3
suppressed their focus-forming activity (Figure 5b). Next, STAT3 KD using shSTAT3#3 significantly
decreased tumorigenesis at orthotopic sites (Figures 5c and 5d). Our results indicated that STAT3
signaling contributed to tumorigenesis induced by *HOXB7*.

249
250
251

4. Discussion

In this study, we demonstrated that *HOXB7* induced cellular transformation and
tumorigenesis to metastasis in transplantation models through activation of the JAK-STAT pathway.
These results together with coamplification of *HOXB7* and *ERBB2* support its role in the poor
prognosis of HER2-positive breast cancer patients.

HOX family genes regulate the development and homeostasis as transcriptional factors
(Bhatlekar *et al.* 2014; Rezsohazy *et al.* 2015; Li *et al.* 2019). Several HOX genes contribute to
cancer malignancy. In fact, overexpression or knockdown experiments supported the idea of *HOXB7*
as an oncogene in the cases of breast, prostate, hepatocellular and gastric cancer cell lines (Errico
et al. 2016; He *et al.* 2017; Wang *et al.* 2017). *HOXB7* also induced EMT and migration/invasion
activity in MCF10A normal human mammary epithelial cells (Wu *et al.* 2006). *In vivo*, *HOXB7*
promoted malignant progression through activation of the TGFß signaling in a MMTV-Hoxb7/Her2
transgenic mouse model (Liu *et al.* 2015). In addition, *HOXB7* overexpression in MCF7 cells (luminal
A subtype) promoted acquired resistance to tamoxifen via activation of receptor tyrosine kinases and
ERα signaling (Jin *et al.* 2012; Jin *et al.* 2015). Previous analysis of microarray data showed that high
levels of *HOXB7* was associated with poor prognosis in HER2–positive, but not in HER2–negative
breast cancers (Chen *et al.* 2008). Consistently, our analysis with recent genomic and RNA-seq data
confirmed their study and further demonstrated that *HOXB7* was significantly co-amplified with
ERBB2. Taken together, coamplification of *HOXB7* with *ERBB2* may enhance the malignancy of
HER2-positive breast cancer.

Using an immortalized mammary gland epithelial cell NMuMG, *HOXB7* induced cellular
transformation *in vitro* and tumor formation *in vivo*. In this cellular context, *HOXB7* increased the

mRNAs of *Stat3* and protein levels of Jak2 and Stat3 with increased level of phosphorylated Stat3
and Stat5. Intriguingly, treatment with ruxolitinib, a JAK1/2 inhibitor, abolished cellular transformation
with inhibition of Stat3 and Stat5 phosphorylation, suggesting the involvement of the JAK-STAT
pathway in *HOXB7*-mediated transformation. Ruxolitinib binds intermediated-active kinase
conformation of Jak2 (Kesarwani *et al.* 2015). As observed in previous research, it causes
upregulation of Jak2 phosphorylation by ruxolitinib treatment via SOCS system (Lee *et al.* 2018).
This phenomenon was also observed in the treatment with pacritinib, another JAK2 inhibitor (Hart *et*
al. 2011). The JAK-STAT pathway is activated in several cancers and is believed to be involved in
tumorigenesis (Hu *et al.* 2021). For this reason, kinase inhibitors against JAK including ruxolitinib
have been developed. Even though *HOXB7* itself cannot be targeted, a crucial downstream
JAK-STAT pathway may be targeted against cancer with *HOXB7* aberrations (Qureshy *et al.* 2020).
In addition, since *Jak2* mRNA was not changed by *HOXB7* expression (Figure 2d), Jak2 is thought to
be regulated at posttranscriptional level. The further analysis for expressional mechanisms of
JAK-STAT signaling is needed for understanding the tumorigenesis induced by *HOXB7*.

Previous study demonstrated that *HOXB7* was not enough to induce tumor in
MMTV-Hoxb7 transgenic mice (Chen *et al.* 2008). In contrast, we showed that *HOXB7* alone was
enough to induce tumorigenesis and lung metastasis in the context of NMuMG cells. This
discrepancy cannot be explained, however, immortalization status of NMuMG may be involved.
Nevertheless, our observation that *HOXB7* enhances lung metastasis in orthotropic model underpin
the importance of *HOXB7* in lung metastasis, consistently with previous study of
M*MTV-Hoxb7/Erbb2* transgenic mice. The transplantation models using NMuMG-luc-HOXB7 cells
will be useful to discover the role of HOXB7 in cancer initiation and metastasis (Nakayama *et al.*
2021). Combined therapy against both *HOXB7* downstream JAK-STAT and ERBB2 may lead to
efficacious outcome in "*ERBB2*- and *HOXB7*-amplified" subtype of breast cancer.

298
299
300

Acknowledgments

We thank Dr. Y. Yamamoto (National Cancer Center Research Institute) and Dr. J. Fujimoto (JBic) for the meaningful comments and discussion, Prof. K. Miyazawa (University of Yamanashi, Yamanashi, Japan) for kindly providing the NMuMG cell line, and Prof. T. Kitamura (Institute of Medical Science, The University of Tokyo, Tokyo, Japan) for kindly providing the Plat-E cell line. This study was conducted in part as a program through the Fukushima Translational Research Project.

Funding

This study was supported by JSPS KAKENHI (grant no. 23241064, Grant-in-Aid for Scientific Research (A) to K.S.; grant no. 18K16269: Grant-in-Aid for Early-Career Scientist to J.N.; grant no. 20J01794, Grant in Aid for JSPS fellows to J.N.) and the Japan Biological Informatics Consortium (JBic).

Authors' contributions

J.N. and K.S. conceived and designed the study. K.A, M.S., S.K., A.M., and J.N. performed the experiments. K.A., M.S., K.N. and J.N. analyzed the data. N.G., S.W. and K.S. interpreted the data. J.N. wrote the manuscript. All authors reviewed and edited the manuscript.

Conflict of interest

The authors declare no conflict of interest.

References

Bhatlekar, S., Fields, J.Z. & Boman, B.M. (2014) HOX genes and their role in the development of human cancers. *J Mol Med (Berl)* **92**, 811-823.

Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A., Byrne, C.J., Heuer, M.L., Larsson, E., Antipin, Y., Reva, B., Goldberg, A.P., Sander, C. & Schultz, N. (2012) The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* **2**, 401-404.

Chen, H., Lee, J.S., Liang, X., Zhang, H., Zhu, T., Zhang, Z., Taylor, M.E., Zahnow, C., Feigenbaum, L., Rein, A. & Sukumar, S. (2008) Hoxb7 Inhibits Transgenic HER-2/neu–Induced Mouse Mammary Tumor Onset but Promotes Progression and Lung Metastasis. *Cancer Research* **68**, 3637-3644.

Errico, M.C., Jin, K., Sukumar, S. & Carè, A. (2016) The Widening Sphere of Influence of HOXB7 in Solid Tumors. *Cancer Res* **76**, 2857-2862.

Fazilaty, H. & Mehdipour, P. (2014) Genetics of breast cancer bone metastasis: a sequential multistep pattern. *Clinical & Experimental Metastasis* **31**, 595-612.

Han, Y., Azuma, K., Watanabe, S., Semba, K. & Nakayama, J. (2022) Metastatic profiling of HER2-positive breast cancer cell lines in xenograft models. *Clinical & Experimental Metastasis*.

Han, Y., Nakayama, J., Hayashi, Y., Jeong, S., Futakuchi, M., Ito, E., Watanabe, S. & Semba, K. (2020) Establishment and characterization of highly osteolytic luminal breast cancer cell lines by intracaudal arterial injection. *Genes Cells* **25**, 111-123.

Hanahan, D. & Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell* **144**, 646-674.

Hart, J.R., Liao, L., Yates, J.R., 3rd & Vogt, P.K. (2011) Essential role of Stat3 in PI3K-induced oncogenic transformation. *Proc Natl Acad Sci U S A* **108**, 13247-13252.

He, X., Liu, Z., Xia, Y. *et al.* (2017) HOXB7 overexpression promotes cell proliferation and correlates with poor prognosis in gastric cancer patients by inducing expression of both AKT and MARKs. *Oncotarget* **8**, 1247-1261.

Hu, X., li, J., Fu, M., Zhao, X. & Wang, W. (2021) The JAK/STAT signaling pathway: from bench to clinic. *Signal Transduction and Targeted Therapy* **6**, 402.

Ihara, T., Hosokawa, Y., Kumazawa, K., Ishikawa, K., Fujimoto, J., Yamamoto, M., Muramkami, T., Goshima, N., Ito, E., Watanabe, S. & Semba, K. (2017) An in vivo screening system to identify tumorigenic genes. *Oncogene* **36**, 2023-2029.

Jin, K., Kong, X., Shah, T. *et al.* (2012) The HOXB7 protein renders breast cancer cells resistant to tamoxifen through activation of the EGFR pathway. *Proc Natl Acad Sci U S A* **109**, 2736-2741.

Jin, K., Park, S., Teo, W.W. *et al.* (2015) HOXB7 Is an ERα Cofactor in the Activation of HER2 and Multiple ER Target Genes Leading to Endocrine Resistance. *Cancer Discov* **5**, 944-959.

Kesarwani, M., Huber, E., Kincaid, Z., Evelyn, C.R., Biesiada, J., Rance, M., Thapa, M.B., Shah, N.P., Meller, J., Zheng, Y. & Azam, M. (2015) Targeting substrate-site in Jak2 kinase prevents emergence of genetic resistance. *Scientific Reports* **5**, 14538.

Kolch, W., Halasz, M., Granovskaya, M. & Kholodenko, B.N. (2015) The dynamic control of signal transduction networks in cancer cells. *Nat Rev Cancer* **15**, 515-527.

Kuroiwa, Y., Nakayama, J., Adachi, C., Inoue, T., Watanabe, S. & Semba, K. (2020) Proliferative Classification of Intracranially Injected HER2-positive Breast Cancer Cell Lines. *Cancers (Basel)* **12**, 1811.

Lee, S., Shah, T., Yin, C., Hochberg, J., Ayello, J., Morris, E., van de Ven, C. & Cairo, M.S. (2018) Ruxolitinib significantly enhances in vitro apoptosis in Hodgkin lymphoma and primary mediastinal B-cell lymphoma and survival in a lymphoma xenograft murine model. *Oncotarget* **9**, 9776-9788.

Li, B., Huang, Q. & Wei, G.H. (2019) The Role of HOX Transcription Factors in Cancer Predisposition and Progression. *Cancers (Basel)* **11**.

Liu, S., Jin, K., Hui, Y., Fu, J., Jie, C., Feng, S., Reisman, D., Wang, Q., Fan, D., Sukumar, S. & Chen, H. (2015) HOXB7 promotes malignant progression by activating the TGFβ signaling pathway. *Cancer Res* **75**, 709-719.

Marotta, L.L., Almendro, V., Marusyk, A. *et al.* (2011) The JAK2/STAT3 signaling pathway is required for growth of CD44$^+$CD24$^-$ stem cell-like breast cancer cells in human tumors. *J Clin Invest* **121**, 2723-2735.

Matsui, A., Fujimoto, J., Ishikawa, K., Ito, E., Goshima, N., Watanabe, S. & Semba, K. (2016) Hepatocyte nuclear factor 1 beta induces transformation and epithelial-to-mesenchymal transition. *FEBS Lett* **590**, 1211-1221.

Nakayama, J., Han, Y., Kuroiwa, Y., Azuma, K., Yamamoto, Y. & Semba, K. (2021) The In Vivo Selection Method in Breast Cancer Metastasis. *Int J Mol Sci* **22**.

Nakayama, J., Ito, E., Fujimoto, J., Watanabe, S. & Semba, K. (2017) Comparative analysis of gene regulatory networks of highly metastatic breast cancer cells established by orthotopic transplantation and intra-circulation injection. *Int J Oncol* **50**, 497-504.

Nakayama, J., Matsunaga, H., Arikawa, K., Yoda, T., Hosokawa, M., Takeyama, H., Yamamoto, Y. & Semba, K. (2022) Identification of two cancer stem cell-like populations in triple-negative breast cancer xenografts. *Dis Model Mech* **15**.

Pereira, B., Chin, S.-F., Rueda, O.M. *et al.* (2016) The somatic mutation profiles of 2,433 breast cancers refine their genomic and transcriptomic landscapes. *Nat Commun* **7**, 11479.

Qureshy, Z., Johnson, D.E. & Grandis, J.R. (2020) Targeting the JAK/STAT pathway in solid tumors. *Journal of Cancer Metastasis and Treatment* **6**, 27.

Rezsohazy, R., Saurin, A.J., Maurel-Zaffran, C. & Graba, Y. (2015) Cellular and molecular insights into Hox protein action. *Development* **142**, 1212-1227.

Saito, M., Kato, Y., Ito, E. *et al.* (2012) Expression screening of 17q12-21 amplicon reveals GRB7 as an ERBB2-dependent oncogene. *FEBS Lett* **586**, 1708-1714.

Wang, T., Fahrmann, J.F., Lee, H. *et al.* (2018) JAK/STAT3-Regulated Fatty Acid β-Oxidation Is Critical for Breast Cancer Stem Cell Self-Renewal and Chemoresistance. *Cell Metab* **27**, 136-150.e135.

Wang, W.M., Xu, Y., Wang, Y.H., Sun, H.X., Sun, Y.F., He, Y.F., Zhu, Q.F., Hu, B., Zhang, X., Xia, J.L., Qiu, S.J., Zhou, J., Yang, X.R. & Fan, J. (2017) HOXB7 promotes tumor progression via bFGF-induced activation of MAPK/ERK pathway and indicated poor

408 prognosis in hepatocellular carcinoma. *Oncotarget* **8**, 47121-47135.
409 Wu, X., Chen, H., Parker, B., Rubin, E., Zhu, T., Lee, J.S., Argani, P. & Sukumar, S. (2006)
410 HOXB7, a homeodomain protein, is overexpressed in breast cancer and confers
411 epithelial-mesenchymal transition. *Cancer Res* **66**, 9527-9534.
412
413
414

　Figure legends

　Figure 1. High expression of HOXB7 correlated to poor prognosis in breast cancer patients.

　(a) Comparative expression analysis of *HOXB7* in breast cancer subtypes of TCGA. Luminal A (n =

　499), luminal B (n = 197), HER2-positive (n = 78), basal-like (n = 171), normal-like (n = 36),

　unidentified (n = 98). Welch's t-test (p < 0.005). (b) OncoPrint of *ERBB2* and *HOXB7* genomic status

　in TCGA. (c) Altered genetic profiles of *ERBB2* and *HOXB7*. Fisher's exact test (p < 0.05). (d)

　Comparative expression analysis of *HOXB7*. Welch's t-test (p < 0.05). (e left) Survival analysis of all

　METABRIC cohorts with *HOXB7* expression and the overall survival status shown in a Kaplan-Meier

　plot. Shading along the curves shows 95% confidential interval. Log-rank test (p = 0.0056) (e right)

　METABRIC cohorts with high HER2 expression (476 patients; Top 25% of all METABRIC cohorts)

　Log-rank test (p = 0.0088) (f) Survival analysis of other breast cancer cohorts (GSE1456; disease

　specific survival; GSE9893; overall survival). GSE1456, log-rank test (p = 0.00022); GSE9893,

　log-rank test (p < 0.0001).

　Figure 2. *HOXB7* expression induced transformation and activation of JAK-STAT signaling.

　(a) Bright imaging of *HOXB7*-expressing cells. (b) Focus formation assay in *HOXB7*-expressing

　NMuMG cells. *Venus*-expressing NMuMG cells were used as a negative control. Left: crystal violet

　staining. Right: bright field focus. Photos were taken at 10 days after the start of the experiment (n =

　1 biological replicates; 3 independent experiments). (c) Western blotting analysis of EMT-related

　genes in *HOXB7*-expressing NMuMG cells. Venus-expressing NMuMG cells were used as a

　negative control. (d) Western blotting of JAK-STAT and growth signaling pathway in

　HOXB7-expressing cells. (e) qRT-PCR of *Stat3* and *Jak2*. Welch's t-test (* p < 0.001, n = 3).

　Figure 3. Focus formation by *HOXB7* was regulated by Stat3 activation.

　(a) Western blotting of JAK-STAT signal pathway in *HOXB7*-expressing NMuMG cells with ruxolitinib

　treatment. (b) *In vitro* growth curve of *HOXB7*-expressing NMuMG cells treated with ruxolitinib. (n.s.:

　Not significant, n = 3 biological replicates) (c) Focus formation assay of *HOXB7*-expressing NMuMG

　cells treated with ruxolitinib. Foci were stained with crystal violet. Photos were taken at 10 days after

　the start of the experiment (n=3 biological replicates; 2 independent experiments).

　Figure 4. Overexpression of *HOXB7* induced tumorigenesis and lung metastasis.

　(a) Tumorigenesis assay of *HOXB7*-expressing NMuMG cells. *Venus*-expressing NMuMG cells were

　used as a negative control. Left: Tumor growth curve with volume. *HOXB7*-expressing NMuMG cells

　formed tumors following subcutaneously transplantation of *Rag2*$^{-/-}$ mice (n = 10). Right: Image of

　HOXB7-positve tumor. (b) Primary tumor formation curve by orthotopic transplantation.

　HOXB7-expressiong NMuMG-luc cells were transplanted to BALB/cSlc-nu/nu mice (n = 12). (c)

　Representative *in vivo* luminescence images of primary tumor in orthotopic transplantation. (d)

Representative *in vivo* luminescence image of lung metastasis in orthotopic transplantation. (e) *Ex vivo* luminescence images of lungs excised from orthotopically transplanted mice. Left: Lungs with no lung metastasis. Right: Lungs with lung metastasis.

Figure 5. STAT3 contributed to tumorigenicity of NMuMG-HOXB7 cells.

(a) Western blotting of STAT3 expression with shSTAT3 knockdown. (b) Focus formation assay of STAT3 knock down in NMuMG-HOXB7 cells. Foci were stained with crystal violet. Photos were taken at 10 days after the start of the experiment (n = 3 biological replicates; 2 independent experiments). (c) Tumor volume of primary tumor in orthotopic transplantation using NMuMG-luc-HOXB7-shGFP cells or NMuMG-luc-HOXB7-shSTAT3#3 cells. Student's t-test (p < 0.05, n = 6). (d) Representative *in vivo* luminescence images of primary tumor in orthotopic transplantation using NMuMG-luc-HOXB7-shGFP cells or NMuMG-luc-HOXB7-shSTAT3#3 cells to BALB/cSlc-nu/nu mice (n = 3).

Figure 1

Figure 2

Figure 3

(a) Cell: NMuMG

Ruxolitinib (µM)	Venus				HOXB7			
	0	0.1	0.3	1.0	0	0.1	0.3	1.0

p-Stat3 (Y705)

Stat3

p-Stat5 (Y694)

Stat5

p-Stat1 (S727)

Stat1

p-Jak1 (Y1034/1035)

Jak1

p-Jak2 (Y1007/1008)

Jak2

α-Tubulin

(b)

Cell number (×10⁵ cells) vs Time (hours)

- Venus
- Venus with Ruxolitinib
- HOXB7
- HOXB7 with Ruxolitinib

(c) Ruxolitinib (1µM)

(-) (+)

Venus

HOXB7

Figure 4

(a)

(b) **Tumor formation at fatpad**
Cells: NMuMG-HOXB7

(c) **Primary tumor**
Week 0　　Week 8　　Week 16

(d) **Lung metastasis**

(e) **Lung metastasis −**　　　**Lung metastasis +**

Figure 5

(a)

Cell: NMuMG-luc -HOXB7

shGFP shSTAT3#1 shSTAT3#2 shSTAT3#3

STAT3

p-STAT3 (Y705)

α-Tubulin

(b)

shGFP shSTAT3#3

(c)

Tumor volume (mm³)

p<0.05

shGFP shSTAT3#3

(d)

Week 0 Week 9

shGFP

shSTAT3#3

1.0
0.8
0.6
0.4
0.2
×10⁷

Radiance (p/sec/cm /sr)