A quantitative map of nuclear pore assembly reveals two distinct mechanisms

Shotaro Otsuka[1,3]*, Jeremy O. B. Tempkin[2], Antonio Z. Politi[1,4], Arina Rybina[1], M. Julius

Hossain[1], Moritz Kueblbeck[1], Andrea Callegari[1], Birgit Koch[1,5], Andrej Sali[2] and Jan Ellenberg[1]*

[1]Cell Biology and Biophysics Unit, European Molecular Biology Laboratory, Meyerhofstrasse 1,

69117 Heidelberg, Germany.

[2]Department of Bioengineering and Therapeutic Sciences, Department of Pharmaceutical

Chemistry, Quantitative Biosciences Institute, University of California, San Francisco, San

Francisco, CA 94143, USA.

[3]Present address: Max Perutz Labs, a joint venture of the University of Vienna and the Medical

University of Vienna, Vienna Biocenter (VBC), Dr. Bohr Gasse 9, 1030 Vienna, Austria.

[4]Present address: Max Planck Institute for Biophysical Chemistry, Göttingen, Germany.

[5]Present address: Max Planck Institute for Medical Research, Heidelberg, Germany

*Correspondence: shotaro.otsuka@univie.ac.at; jan.ellenberg@embl.de

Summary

Understanding how the nuclear pore complex (NPC) assembles is of fundamental importance to

grasp the mechanisms behind its essential function and understand its role during evolution of

eukaryotes[1–4]. While we know that at least two NPC assembly pathways exist, one during exit

from mitosis and one during nuclear growth in interphase, we currently lack a quantitative map of

their molecular events. Here, we use fluorescence correlation spectroscopy (FCS) calibrated live

imaging of endogenously fluorescently-tagged nucleoporins to map the changes in composition

and stoichiometry of seven major modules of the human NPC during its assembly in single

dividing cells. This systematic quantitative map reveals that the two assembly pathways employ

strikingly different molecular mechanisms, inverting the order of addition of two large structural

components, the central ring complex and nuclear filaments. Our dynamic stoichiometry data

allows us to perform the first computational simulation that predicts the structure of postmitotic

NPC assembly intermediates.

The nuclear pore complex (NPC) is the largest non-polymeric protein complex in eukaryotic cells.

It spans the double membrane of the nucleus (nuclear envelope; NE) to mediate the

macromolecular transport between the nucleus and the cytoplasm. To achieve this essential

function, the NPC forms an octameric proteinaceous channel composed of multiples of eight of

over 30 different nucleoporins (Nups) that form 6–8 protein modules, the NPC subcomplexes[1,2].

Therefore, more than 500 individual proteins have to come together to assemble one nuclear pore,

which has the mass of tens of ribosomes. NPCs are thought to represent a key step in the evolution

of endomembrane compartmentalization that allowed ancestral eukaryotes to separate their

genome from the cytoplasm[3,4].

In proliferating cells, there are two main pathways to assemble the NPCs. During nuclear

assembly after mitosis, the NPCs form together with nuclear membranes to rapidly build new

nuclei in the daughter cells (called postmitotic NPC assembly). During nuclear growth in

interphase, NPCs then continue to assemble continuously for homeostasis (referred to as

interphase assembly). Research over the last decade has revealed that postmitotic and interphase

NPC assembly possess distinct kinetic, molecular and structural features[5–12], suggesting that two

fundamentally different mechanisms build the same protein complex. In postmitotic assembly,

several thousand NPCs assemble within a few minutes during sealing of the initially fenestrated

nuclear membranes, whereas interphase NPC assembly occurs more sporadically, requires about

one hour, and involves a new discontinuity in the double membrane barrier of the NE. Studies

using molecular depletions have shown that the Nup ELYS is required for postmitotic assembly

but appears dispensable for interphase assembly[5], whereas the membrane curvature-sensing

domain of Nup133[6], Pom121 and Sun1[7,8], the import of Nup153 into the nucleus[9], and Torsins[10]

seem to be required only for interphase assembly. Recent studies correlating real time imaging

with three-dimensional electron microscopy have revealed that postmitotic NPC assembly

proceeds by radial dilation of small membrane openings[11], while in interphase, assembly induces

an asymmetric inside-out fusion of the inner and outer nuclear membranes[12].

However, how several hundred proteins self-organize to form the NPC channel *via* these two

distinct assembly pathways has remained largely enigmatic. It is technically challenging to locate

the transient and rare assembly events, which has prevented investigation of the structure of

assembly intermediates by either cryo-EM tomography or super-resolution microscopy. In

addition, the large number of building blocks and their cooperativity often leads to complex

nonlinear kinetics that can only be interpreted mechanistically using computational modelling of

the structures formed during assembly. While we have some information about the dynamic

addition of Nups after mitosis[13,14], only sparse dynamic data is available for interphase

assembly[15,16]. Importantly, these earlier studies could not distinguish postmitotic and interphase

assemblies, whose co-occurrence in different regions of the nucleus was only discovered later;

moreover, these studies provided only qualitative descriptions as they relied on ectopic expression

of fluorescently-tagged Nups. Kinetic data about NPC assembly that can distinguish between the

postmitotic and interphase pathways is required, including the copy numbers of Nups that

assemble into forming NPCs over time. Such data would enable modeling of the assembly process

and allow us to start to understand the two assembly mechanisms.

Quantitative live imaging of ten nucleoporins representing seven major NPC building blocks

To quantitatively analyze the changes in concentration of Nups at the NE during exit from mitosis

and nuclear growth in G1, we genome-edited HeLa cells, homozygously tagging the endogenous

genes for ten different Nups with mEGFP or mCherry. We chose proteins that represent the major building blocks of the fully assembled pore, including the nuclear filament protein Tpr, the nuclear Nup153, the Y-complex members Nup107 and Seh1, the central ring complex members Nup93 and Nup205, the central channel protein Nup62, the transmembrane protein Pom121 as well as the cytoplasmic filament proteins Nup214 and Nup358. Homozygous tagging was verified by careful quality control of the genome edited monoclonal cell lines[17], ensuring that the tagged subunit was expressed at physiological levels, localized to the NPC and that cell viability and mitotic progression were normal (Figs. 1 and 2). The fusion proteins are likely functional, given that most Nups show strong phenotypes upon knock-out or depletion[18].

To characterize the fluorescently-tagged Nups, we first performed super-resolution (STED) microscopy to determine NPC density in fully grown nuclei of the knock-in cell lines, showing that homozygous tagging had little effect on NPC density that was comparable within 15% between all cell lines with an average of 10.1 NPC per μm^2 (Fig. 1b, c), in good agreement with our previous estimates by electron microscopy of HeLa cells[12]. We then used fluorescence correlation spectroscopy (FCS) calibrated confocal microscopy[19,20] to determine the concentration and total number of the Nups at the NE in living cells (Extended Data Table 1, for details see Methods). Using the measured NPC density and the Nup concentration at the NE, we could calculate the average copy number of each Nup per NPC (Fig. 1d). This data showed that the investigated Nups were on average present in 16, 32, or 48 copies per pore, as expected from the eightfold symmetry of the complex and overall in good agreement with previous estimates by mass spectrometry[21]. This observation quantitatively validated that the homozygous tagging did not affect the incorporation of the Nups into the NPC. For Nup153 and Pom121 that exhibited lower concentrations likely due to subhomozygous tagging (Fig. 2a, Extended Data Fig. 1), we normalized their stoichiometry to the expected number of copies in the mature pore for comparison (Fig. 3a).

NPC assembly relies on and consumes almost half of the material inherited from the mother cell within one hour after mitosis

We then used our validated cell line resource to quantitatively image the Nups during both postmitotic and interphase NPC assembly, from metaphase until the end of the rapid nuclear growth phase in G1, two hours after anaphase onset. To this end, we performed systematic FCS-calibrated 3D confocal time-lapse microscopy[22] (Fig. 2a). Using the single molecule fluctuation calibration, the 4D imaging data could be converted into maps of subcellular protein concentration (Fig. 2a). Counterstaining live nuclei with SiR-DNA[23], enabled computational image segmentation to measure the soluble cytoplasmic pool and the NE associated pool over time[22]. Temporal alignment to anaphase onset then allowed us to compare the dynamic association of all Nups with the NE over time (Fig. 2a). Overall, the investigated Nups are present in 250,000 to 1,200,000 copies per human metaphase cell. After mitosis, this building material is split between the daughter cells with little detectable new protein synthesis in the first hour after anaphase onset (Fig. 2b). Notably, 34–53% of the soluble pool present in the cytoplasm in metaphase was rapidly re-localized to the NE during the first hour after exit from mitosis (43, 38, 47, 37, 41, 44, 53, and 34% for Nup107, Seh1, Nups205, 93, 62, 214, Tpr and Nup358, respectively) (Fig. 2b), indicating that NPC assembly initially relies almost entirely on the pool of building blocks inherited from the mother cell to form the first 4,000–5,000 NPCs[11,12].

Quantitative analysis of multimolecular kinetics of postmitotic and interphase NPC assembly

As we and others have demonstrated previously[12,24], postmitotic and interphase NPC assembly can be observed in the same living cell in different regions of the NE within the first two hours after mitosis. While postmitotic assembly dominates the peripheral "non-core" regions of the NE, the

central "inner core" area is only populated with NPCs after exit from mitosis when dense spindle

microtubules have been removed from the DNA surface[25]. Using computational segmentation and

assignment of the inner-core and non-core regions (Extended Data Fig. 2a), we measured the

concentration changes of the ten Nups in these two regions separately. A two-component model

of a fast (postmitotic) and a slow (interphase) assembly process fits the experimental data well,

allowing us to kinetically unmix the two assembly processes for each Nup (Extended Data Figs.

2b–d and 3). In this way, we could for the first time perform an integrated analysis of the real time

kinetics of absolute concentration changes of Nups in all major NPC modules during the two

assembly processes at the NE (Fig. 3a). This analysis immediately revealed that the overall

duration of the two processes is very different, with postmitotic assembly essentially complete 15

min after anaphase onset, while interphase assembly only reaches a plateau after 100 min,

consistent with our previous estimates based on live cell imaging[13,16] and correlative electron

microscopy[11,12]. Both processes reached the same final ratios between the different Nups and thus

presumably formed identical NPCs. However, the temporal order in which components were

added was distinct, including, for example, an earlier assembly of Pom121 relative to the Y-shaped

complex during interphase assembly, consistent with our previous observations[16].

To comprehensively investigate the molecular differences between the two assembly

processes, we relied on the constant NPC density and changes in nuclear surface area[12] to convert

the NE concentrations of all investigated Nups into their average copy number per assembling

NPC over time (Fig. 3a). This result in turn enabled us to estimate changes in subunit stoichiometry

of the complex during its assembly in living cells. To facilitate the comparative analysis of the

assembly kinetics between the two pathways, in which ten components assemble with different

speed and order, we first reduced the dimensionality of the kinetic data. To this end, we assigned

a single characteristic time-point of assembly to each Nup, by sigmoidal fitting of its full kinetic

signature (Extended Data Fig. 4a). Plotting the copy number *vs* the average time-point of assembly

provides an overview of the major molecular differences between the two assembly pathways (Fig.

3b).

While the Y-complex, Pom121, and Nup153 form a core of the first modules that assemble

almost simultaneously within one minute in postmitotic assembly, these components are stretched

out into a clear temporal order of first Nup153, second Pom121, and third the Y-complex (notably

with its two investigated subunits also assembling simultaneously in interphase) over more than

ten minutes in interphase assembly. The end of assembly on the other hand is marked for both

process by the addition of the large cytoplasmic filament protein Nup358. The major difference

was thus neither in initiation nor termination of assembly, but rather in the middle of the two

assembly pathways. During postmitotic assembly, the Y-complex is rapidly combined with

components of the central ring, building the inner core of the pore within only three minutes prior

to addition of either cytoplasmic or nuclear filament proteins, which follow later. In contrast,

during interphase assembly, the Y-complex is first combined with the nuclear filament protein Tpr

and the base of the cytoplasmic filament Nup214, while the central ring complex is added later.

This observation clearly shows that postmitotic and interphase NPC assembly not only proceed

with different speed but also follow a different molecular mechanism using an inverted molecular

order between the central ring and nuclear filaments (Fig. 3b, Extended Data Fig. 4b).

To obtain additional insight into the assembly mechanism, it is instructive to go beyond

examining the temporal order and examine our data for evidence of coupling between different

components of the NPC. Theoretically, for a sequential assembly mechanism where late steps

depend on early steps, the observed ensemble kinetics of a late binding protein must contain the

history of all previous events (see Methods for details). To test this prediction, we examined if one

or both of our assembly pathways exhibited a correlation between the average assembly time and

duration of the different components. We plotted the time needed for 90% of one protein to

accumulate in the population of assembling pores (the assembly duration) *vs* the time when this

accumulation is half maximal in the overall process (the average time-point of assembly) for all

analysed Nups (Fig. 3c, Extended Data Fig. 4a). Indeed, postmitotic assembly showed a strong

positive linear correlation (Fig. 3c), indicating a sequential assembly mechanism, which for

example implies that Nup62 is incorporated into the NPC before Tpr can bind. In contrast,

interphase assembly shows poor correlation between duration and time-point of assembly (Fig.

3c), indicating a less strictly sequential pathway, where proteins may assemble more

independently of each other.

Integrative modeling of the NPC assembly pathway

To obtain a more comprehensive mechanistic view, we computed a spatiotemporal model of the

macromolecular assembly pathway, based on our dynamic multimolecular stoichiometry data in

combination with the available ultrastructural data about NPC assembly[11] and the partial

pseudoatomic model of the mature NPC[26,27]. We modeled here only the postmitotic assembly, as

it showed the kinetic hallmarks of a sequential process and is known to proceed by dilating an

existing membrane pore with a smoothly growing proteinaceous density[11]. We focused on the

Nups contained in the structural model of the human NPC, including Nup107 and Seh1 for the Y-

complex as well as Nup93, Nup205, and Nup62 for the central ring/channel complex[26,27]. We

constrained their copy number by our stoichiometry data for the time points for which correlative

electron tomography data is available[11], in order to use the membrane shapes and associated

protein densities from the tomography to constrain Nup positions. To structurally model the

spatiotemporal assembly process, we generalized our integrative modeling method for determining

static structures of macromolecular assemblies[28,29]. In outline (see Methods for details), we first

model ensembles of structures at discrete time points, then connect them between time points into

assembly trajectories and finally rank the alternative trajectories by fit to our data.

We focus on the best fitting macromolecular assembly pathway that accounts for over 80% of

the posterior model density (the second-scoring accounts for only 19%). This trajectory starts by

formation of a single nuclear ring, composed of eight Y-complexes, concomitantly with an initial

accumulation of the FG-repeat protein Nup62 and Nup205 in the center of the membrane hole

(Fig. 4). The cytoplasmic Y-complex is then added on the cytoplasmic side, before the second set

of the Y-complex ring assembles on the nuclear side, again one eight-membered ring after another.

In the center of the pore, Nup62 dilates from an amorphous mass into a small ring and associates

with Nup205 to form the central ring complex. This "nuclear ring first" assembly mechanism is

consistent with the observation of an eight-fold symmetric protein density on the inner nuclear

membrane at early stages of assembly[11]. In addition, the early accumulation of Nup62 to the pore

center suggests a potential role of the hydrophobic FG-repeats in the initial dilation of the small

membrane hole into the larger, NPC sized channel.

Discussion

Our data revealed that the two assembly pathways employ strikingly different molecular

mechanisms. While it is at first glance surprising that the cell builds the same machine in two

fundamentally different ways, these distinct pathways can be rationalized by considering them in

the light of the rather different physiological requirements that the cell faces after mitosis and

during nuclear growth in interphase. Immediately after exit from mitosis, the nucleus is formed *de*

novo and the cell needs to rapidly seal its genome, exclude cytoplasm from it[30] and quickly restore

import of essential nuclear proteins. Since the new NE originates from highly fenestrated ER

sheets, nuclear transport channels can be built without a formal membrane fusion requirement[11].

The mechanism we observed for postmitotic assembly, appears well adapted to this context. The

rapid assembly of the nuclear Y-ring and the central ring could provide a fast way to prevent

complete fusion of the ER membrane holes and assemble a minimal, transport competent

proteinaceous phase in them. Assembly of this first "transport core" NPC is complete within less

than 10 min after mitosis, which precisely matches the time when transport competence is gained

by postmitotic nuclei[13,14].

By contrast in telo/G1 phase, the nucleus already has a sealed, double membraned NE with

about 2500 postmitotically assembled and transporting pores, but needs to expand to allow the

genome to take up its interphase functions. To support nuclear growth, the cell needs to maintain

a constant NPC density to ensure sufficient transport capacity and maintain homeostasis to provide

enough NPC building blocks for the next cell generation, which we have shown to initially rely

completely on material from the mother cell. Interphase NPC assembly requires a *de novo* fusion

between the inner and outer nuclear membrane but must not disrupt nuclear integrity. The

molecular pathway we observed for interphase assembly makes new predictions about its unique,

inside out evaginating, mechanism. The cell first not only builds the two nuclear Y-rings but also

accumulates the material for their cytoplasmic counterparts and then combines them with the

nuclear filament proteins, all prior to the time of membrane fusion[12]. This mechanism suggests

that the cytoplasmic Y-rings, including the base of the cytoplasmic filaments with Nup214, are

already "prebuilt" within the inner membrane evagination, where the small available volume

would predict that they must be present in a very different structure than in the fully mature pore

after fusion. In addition, the unexpectedly early presence of Tpr suggests a key role of this large

coiled-coil protein during interphase assembly, potentially in the membrane bending and/or fusion

process. Surprisingly, the central ring complex, which is a core structural element between the

nuclear and cytoplasmic Y-rings in the mature pore[27] is added only later, suggesting that during

interphase assembly, nuclear transport control may only be added after the membrane fusion step.

The order of Nup assembly for interphase assembly we observed here is consistent with the

recently reported order of NPC assembly in budding yeast that undergoes a closed mitosis[31], except

for the Tpr homologues Mlp1/2 which assemble late in yeast[31]. This difference could be due to the

fundamental differences in cell cycle remodeling of the NE or additional functions of Mlp proteins

that are believed to compensate for the absence of lamins in yeast.

Beyond the assembly mechanism the two pathways we have mapped here start to shed light

on the role of the NPC during evolution of the endomembrane system in eukaryotes[3,4]. We

speculate that the modern NPC combines ancient membrane bending (e.g. coiled-coil filaments,

such as Tpr) and membrane hole plugging (e.g. FG-repeat proteins of the central ring complex)

modules, that were potentially previously used separately for extruding the cell surface or keeping

transport channels open in the endomembranes around the genome. The key evolutionary

innovation might lie in combining and controlling these activities, potentially by the eight-

membered ring architecture of the nuclear Y-complex. In the future, comparing NPC assembly

pathways in different species on the eukaryotic evolutionary tree, might help us to understand how

the assembly of complex modern protein machines reflects their evolutionary origins.

Main references

1   Hampoelz, B., Andres-Pons, A., Kastritis, P. & Beck, M. Structure and Assembly of the

Nuclear Pore Complex. *Annu. Rev. Biophys.* **48**, 515–536 (2019).

2   Lin, D. H. & Hoelz, A. The Structure of the Nuclear Pore Complex (An Update). *Annu.*

Rev. Biochem. **88**, 725–783 (2019).

3   Baum, D. A. & Baum, B. An inside-out origin for the eukaryotic cell. *BMC Biol.* **12**, 76

(2014).

4   Otsuka, S. & Ellenberg, J. Mechanisms of nuclear pore complex assembly - two different

ways of building one molecular machine. *FEBS lett.* **592**, 475–488 (2018).

5   Franz, C. *et al.* MEL-28/ELYS is required for the recruitment of nucleoporins to chromatin

and postmitotic nuclear pore complex assembly. *EMBO Rep.* **8**, 165–172 (2007).

6   Doucet, C. M., Talamas, J. A. & Hetzer, M. W. Cell cycle-dependent differences in nuclear

pore complex assembly in metazoa. *Cell* **141**, 1030–1041 (2010).

7   Talamas, J. A. & Hetzer, M. W. POM121 and Sun1 play a role in early steps of interphase

NPC assembly. *J. Cell Biol.* **194**, 27–37 (2011).

8   Funakoshi, T., Clever, M., Watanabe, A. & Imamoto, N. Localization of Pom121 to the

inner nuclear membrane is required for an early step of interphase nuclear pore complex

assembly. *Mol. Biol. Cell* **22**, 1058–1069 (2011).

9   Vollmer, B. *et al.* Nup153 Recruits the Nup107-160 Complex to the Inner Nuclear

Membrane for Interphasic Nuclear Pore Complex Assembly. *Dev. Cell* **33**, 717-728 (2015).

10   Rampello, A. J. *et al.* Torsin ATPase deficiency leads to defects in nuclear pore biogenesis

and sequestration of MLF2. *J. Cell Biol.* **219** (2020).

11   Otsuka, S. *et al.* Postmitotic nuclear pore assembly proceeds by radial dilation of small

membrane openings. *Nat. Struct. Mol. Biol.* **25**, 21–28 (2018).

12   Otsuka, S. *et al.* Nuclear pore assembly proceeds by an inside-out extrusion of the nuclear

envelope. *Elife* **5** (2016).

13   Dultz, E. *et al.* Systematic kinetic analysis of mitotic dis- and reassembly of the nuclear

pore in living cells. *J. Cell Biol.* **180**, 857–865 (2008).

14   Otsuka, S., Szymborska, A. & Ellenberg, J. Imaging the assembly, structure, and function

of the nuclear pore inside cells. *Methods Cell Biol.* **122**, 219–238 (2014).

15   D'Angelo, M. A., Anderson, D. J., Richard, E. & Hetzer, M. W. Nuclear pores form de

novo from both sides of the nuclear envelope. *Science* **312**, 440–443 (2006).

16   Dultz, E. & Ellenberg, J. Live imaging of single nuclear pores reveals unique assembly

kinetics and mechanism in interphase. *J. Cell Biol.* **191**, 15–22 (2010).

17   Koch, B. *et al.* Generation and validation of homozygous fluorescent knock-in cells using

CRISPR-Cas9 genome editing. *Nat. Protoc.* **13**, 1465–1487 (2018).

18   Guglielmi, V., Sakuma, S. & D'Angelo, M. A. Nuclear pore complexes in development

and tissue homeostasis. *Development* **147** (2020).

19   Wachsmuth, M. *et al.* High-throughput fluorescence correlation spectroscopy enables

analysis of proteome dynamics in living cells. *Nat. Biotechnol.* **33**, 384–389 (2015).

20   Politi, A. Z. *et al.* Quantitative mapping of fluorescently tagged cellular proteins using

FCS-calibrated four-dimensional imaging. *Nat. Protoc.* **13**, 1445–1464 (2018).

21   Ori, A. *et al.* Cell type-specific nuclear pores: a case in point for context-dependent

stoichiometry of molecular machines. *Mol. Syst. Biol.* **9**, 648 (2013).

22   Cai, Y. *et al.* Experimental and computational framework for a dynamic protein atlas of

human cell division. *Nature* **561**, 411–415 (2018).

23   Lukinavicius, G. *et al.* SiR-Hoechst is a far-red DNA stain for live-cell nanoscopy. *Nat.*

Commun. **6**, 8497 (2015).

24   Maeshima, K. *et al.* Cell-cycle-dependent dynamics of nuclear pores: pore-free islands and

lamins. *J. Cell Sci.* **119**, 4442–4451 (2006).

25   Vietri, M. *et al.* Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear

envelope sealing. *Nature* **522**, 231–235 (2015).

26   von Appen, A. *et al.* In situ structural analysis of the human nuclear pore complex. *Nature*

526, 140–143 (2015).

27   Kosinski, J. *et al.* Molecular architecture of the inner ring scaffold of the human nuclear

pore complex. *Science* **352**, 363–365 (2016).

28   Kim, S. J. *et al.* Integrative structure and functional anatomy of a nuclear pore complex.

Nature **555**, 475–482 (2018).

29   Rout, M. P. & Sali, A. Principles for Integrative Structural Biology Studies. *Cell* **177**,

1384–1403 (2019).

30   Cuylen-Haering, S. *et al.* Chromosome clustering by Ki-67 excludes cytoplasm during

nuclear assembly. *Nature*, **587**, 285–290 (2020).

31   Onischenko, E. *et al.* Maturation Kinetics of a Multiprotein Complex Revealed by

Metabolic Labeling. *Cell* **183**, 1785–1800 (2020).

327
Fig. 1 | Quantitative imaging of GFP-knock-in nucleoporin (Nup) cell lines.
a, Genome-edited HeLa cells with homozygously mEGFP-tagged Nups observed by confocal
microscopy. Fluorescent intensity was converted into protein concentration by FCS-calibrated
imaging[20]. Images were filtered with a median filter (kernel size: 0.25 × 0.25 μm) for
presentation purposes. Scale bar, 10 μm. **b**, **c**, Stimulated emission depletion (STED)
microscopy. The genome-edited cells were stained with anti-Nup62 antibody and imaged (**b**),
and then the density of nuclear pores was quantified (**c**). The plot is from 6, 6, 6, 5, 6, 6, 6, and 4
cells for Nup107, Seh1, Nup205, Nup93, Nup62, Nup214, Tpr and Nup358, respectively. Scale
bar, 1 μm. **d**, Calculated copy number of Nups per nuclear pore. The plot is from 241, 37, 41, 20,
41, 26, 55, and 28 cells for Nup107, Seh1, Nup205, Nup93, Nup62, Nup214, Tpr and Nup358,
respectively. The median is depicted as a line.

339
340

Fig. 2 | Dynamic concentration maps of Nups after anaphase onset (AO).
a, HeLa cells whose Nups are endogenously tagged with mEGFP or mCherry were imaged every
30 sec by three-dimensional confocal microscopy. Single confocal sections are shown. Images
were calibrated by FCS to convert fluorescence intensities into cellular protein concentration.
*For Nup153 and Pom121, these Nups are not fully validated to be homozygously-tagged.
Images were filtered with a median filter (kernel size: 0.25 × 0.25 μm). Scale bar, 10 μm. **b**,
Dynamic change of Nup numbers during mitotic exit. The number of Nups in cytoplasm (dark
blue), nucleoplasm (medium dark blue) and nuclear envelope (light blue) are plotted against time
after anaphase onset. The plot is the mean of 15, 20, 13, 14, 22, 22, 19, and 24 cells for Nup107,
Seh1, Nup205, Nup93, Nup62, Nup214, Tpr and Nup358, respectively.

351
Fig. 3 | The molecular assembly order and maturation kinetics are distinct for postmitotic
and interphase assembly.
a, Plots of the average copy number per nuclear pore computed from mathematical modeling for
postmitotic (left) and interphase (right) assembly (see Methods and Extended Data Figs. 2 and 3
for details). For Nup153 and Pom121 (dashed lines), their absolute amount was estimated using
the copy number determined from the previous study (32 for Nup153 and 16 for Pom121)[21]. **b**,
The average copy number of individual Nups per nuclear pore are plotted along their average
time-points of assembly in postmitotic and interphase assembly pathways. **c**, Plots of Nup
assembly duration along the time-point of assembly for postmitotic and interphase assembly
pathways. The crosses indicate the 95% confidence intervals. The dots are the mean values.
Values are listed in Extended Data Table 2. The long straight line shows the result of a linear
regression to the mean values. The gray area is the 95% confidence interval of the linear
regression.

365
Fig. 4 | Integrative model of the postmitotic NPC assembly pathway.
a, Protein density (grey) overlaid with the NE surface (wireframe model, grey) at each time
point. C, cytoplasm; N, nucleoplasm. **b**, Copy numbers of Nups used to construct a static NPC
model at each time point. **c**, The best-scoring model of postmitotic assembly pathway (top and
side views). The uncertainty of each Nup localization is indicated by the density of the
corresponding color: Y-complex (green), Nup93-Nup188-Nup155 complex (blue), Nup205-
Nup93-Nup155 complex (cyan), Nup62-Nup54 complex (purple), and Nup155 (orange).

Methods

Cell culture

Wildtype HeLa kyoto cells (RRID: CVCL_1922) were kind gift from Prof. Narumiya in Kyoto University, and the genome was sequenced previously[32]. Cells were grown in high glucose Dulbecco's Modified Eagle's Medium (DMEM) containing 4.5 g/l D-glucose (Sigma Aldrich, St. Louis, MO) supplemented with 10% fetal calf serum (FCS), 2 mM l-glutamine, 1 mM sodium pyruvate, and 100 μg/ml penicillin and streptomycin at 37 °C and 5% CO_2. The mycoplasma contamination was inspected by PCR every 2 or 3 months and was always negative.

Genome editing

Monomeric enhanced GFP (mEGFP) and mCherry were inserted into the genome using zinc finger nucleases or CRISPR-Cas9 nickases[17]. The following six cell lines had been generated and published previously: Nup62-mEGFP[33], mEGFP-Nup107[12], mEGFP-Nup205[11], mEGFP-Nup214, mEGFP-Nup358 (also called RanBP2) and Tpr-mEGFP[22]. The following four cell lines were generated in this study: mEGFP-Seh1, Nup93-mEGFP, mEGFP-Nup153, and Pom121-mCherry. The gRNA sequences used for generating these cell lines are summarized in Extended Data Table 3. For the Nup93-mEGFP cell line, CRISPR-Cas9 nickases and the donor plasmid were transfected by electroporation (Neon Transfection System, Thermo Fisher Scientific, Waltham, MA) instead of a polymer-mediated transfection reagent.

FCS-calibrated live-cell imaging and estimation of Nup copy numbers per NPC

(i) Wild-type cells, (ii) wild-type cells transfected with mEGFP using Fugene6 (Promega, Madison, WI), (iii) mEGFP-Nup107 genome-edited cells, and (iv) the cells of another mEGFP-Nup genome-edited cell line, were seeded on each well of 8-well Lab-Tek Chambered Coverglass (Thermo Fisher Scientific). On the day of live-cell imaging, DMEM was replaced by imaging

medium: CO_2-independent medium without phenol red (Invitrogen) containing 20% FCS, 2 mM

l-glutamine, and 100 μg/ml penicillin and streptomycin. The imaging medium was supplemented

with 50 nM silicon–rhodamine (SiR) Hoechst[23]. Cells were incubated inside the microscope-body-

enclosing incubator at 37 °C for at least 30 min before imaging.

Calibrated imaging using fluorescence correlation spectroscopy (FCS) was carried out as

described in a previous report[20]. Briefly, the confocal volume was determined by performing FCS

using a dye with known diffusion coefficient and concentration (Alexa Fluor 488 NHS ester;

Thermo Fisher Scientific for mEGFP, Alexa Fluor 568 NHS ester; Thermo Fisher Scientific for

mCherry). To convert fluorescence intensity to the concentration, FCS was performed in the cells

that transiently-express mEGFP or mCherry alone. Then a calibration curve was obtained by

plotting the fluorescence intensity along the concentration. The background fluorescence signal

was measured in cells without expressing fluorescent proteins and subtracted.

To measure the concentration of Nups, mEGFP-Nup genome-edited cells in interphase were

imaged in 3D using a confocal microscope (LSM780; Carl Zeiss, Oberkochen, Germany) and a

40× 1.2 NA C-Apochromat water immersion objective (Carl Zeiss) at 37 °C in a microscope-body-

enclosing incubator, under the following conditions: 21 optical sections, section thickness of 2.0

μm, z-stacks of every 1.0 μm, and the xy pixel size of 0.25 μm. When the NE is not perpendicular

to the confocal plane of the 3D stacks, the fluorescence intensity at the NE is nonisotropic in the

point-spread function (PSF), which results in underestimation of the signal. To avoid such

underestimation, a single plane was selected that contains the largest nuclear area in which the NE

is perpendicular to the imaging plane and thus isotropic in the PSF. The fluorescence intensity of

Nups was quantified on this single plane using the NE mask with the width of three pixels (0.75

μm) that was generated from a SiR-DNA channel. Background fluorescence intensity was

measured in wild-type cells without expressing any fluorescent proteins and subtracted. The Nup

fluorescence intensity on the NE was converted to the concentration using the calibration curve

generated by FCS above. The number of Nups per square micro-meter was calculated from the concentration and then divided by the NPC density per square micro-meter measured by stimulated emission depletion (STED) microscopy. This absolute quantification of Nup copy number with FCS calibration was done using 47 mEGFP-Nup107 genome-edited cells in interphase. For other mEGFP-Nups genome-edited cells, their Nup fluorescent intensities on the NE were directly compared with the ones of mEGFP-Nup107 genome-edited cells on the same 8-well Lab-Tek Chambered Coverglass, and then their concentrations were determined using the intensity ratios to the mean intensity of mEGFP-Nup107 without using a FCS calibration curve. For Pom121-mCherry, the copy number was quantified independently.

Measurement of nuclear pore density by stimulated emission depletion (STED) microscopy

For STED super-resolution imaging, cells were fixed with 2.4% formaldehyde (Electron Microscopy Sciences, Hatfield, PA) in PBS for 10 min, extracted with 0.4% Triton X-100 (Sigma Aldrich) in PBS for 5 min, and blocked with 5% normal goat serum (Life Technologies, Carlsbad, CA) in PBS for 10 min at room temperature. Subsequently, the cells were incubated overnight at 4°C with a mouse anti-Nup62 (Cat. No. 610497; BD Biosciences, Franklin Lakes, NJ) antibody, and then with an Abberior® STAR RED-conjugated anti-mouse IgG (Cat. No. 2-0002-011-2, Abberior GmbH, Göttingen, Germany) for 30 min at room temperature. After multiple washes in PBS, cells were mounted in Vectashield (Cat. No. H-1500, Vector Laboratories Inc., Burlingame, CA). Super-resolution imaging was performed on a Leica SP8 3X STED microscope as described in a previous report[12]. The images were taken with a final optical pixel size of 20 nm, z-stacks of every 250 nm, and the optical section thickness of 550 nm. Images were filtered with a Gaussian filter (kernel size: 0.5×0.5 pixel) for presentation purposes. The shrinkage of the nucleus by formaldehyde fixation was quantified by comparing the volume of the nuclei of live cells with the ones of fixed cells. The shrinkage was $9.1 \pm 2.6\%$ (the average and standard error, N = 36 cells).

The NPC density was corrected for the nuclear shrinkage for the calculation of Nup copy number

per NPC in Fig. 1d.

To quantify NPC density, the raw STED data were processed in ImageJ

(http://rsbweb.nih.gov/ij/) with a mean filter (kernel size: 2 × 2 pixels) and a sliding paraboloid

(radius: 5 pixels) for background subtraction. Detection of central peak positions for individual

NPCs was carried out with the plugin TrackMate[34], using DoG detector and adjusting the detection

threshold as the spot diameter size. The resulting 3D NPC coordinates were used to visualize and

determine flat and curved regions of the nucleus. Using this map, circular and ellipsoidal ROIs

could then be selected in the flatter parts containing central NPC positions within the Z-depth of

approx. 500 nm, which corresponds to 2–3 microscopic slices in the images. The remaining signal

outside the ROIs, as in curved regions or cytoplasmic structures were discarded from further

analysis. NPC densities were calculated for each cell separately by dividing the number of NPCs

within the selected ROIs by the corresponding ROI areas. For each cell line, the values were

combined to calculate the mean and median NPC density values.

Quantification of Nup copy number in the cytoplasm and the nucleoplasm as well as in non-

core and core regions of the NE

Mitotic cells were imaged and monitored from anaphase onset for two hours in 3D by confocal

microscopy. The microscopy setup and the imaging conditions are described above. Time-lapse

imaging for mEGFP-tagged Nups was performed every 30 sec. Photobleaching was negligible and

thus not corrected. Time-lapse imaging for Pom121-mCherry was carried out every 60 sec, and

photobleaching was corrected by measuring a fluorescence signal decay in a neighboring cell in

the same field of view. Visualization of the chromosome surface in 3D was done in the Amira

software package[35].

To measure the Nup accumulation on the NE, single planes were selected that contain the

largest nuclear area at individual time points to avoid underestimation of the signal as mentioned

earlier. The Nup intensity was quantified on the NE mask with the width of 0.75 μm that was

generated from a SiR-DNA signal at each time point. Except for Nup107 and Seh1, the Nup signal

in the cytoplasm and nucleoplasm was measured and used as background. For Nup107 and Seh1,

only the cytoplasmic signal was used as background because of their localization at kinetochores.

These background values were quantified at individual time points and subtracted from the Nup

intensities on the NE. The measured Nup intensity was converted into the concentration and then

multiplied with nuclear surface area to calculate the total number of the Nups on the NE. For

Nup153 and Pom121, we did not convert the fluorescence intensity to the concentration as the cell

lines were not fully validate to be homozygously-tagged.

The Nup copy number was also calculated in the cytoplasm and the nucleoplasm during first

two hours after anaphase onset. The cytoplasm mask was created by subtracting a mask of the

nucleus generated from a SiR-DNA signal from a mask of the whole cell generated from a

mEGFP-Nup signal. The mask for the nucleoplasm was created by eroding three pixels of the

nuclear mask generated from a SiR-DNA signal. The Nup fluorescence intensity was quantified

on these cytoplasmic and nucleoplasmic masks over time and then converted into the

concentration. To calculate the total number of Nups in the cytoplasm and the nucleoplasm, the

measured concentration was multiplied by the volume of the respective compartments. For the

cytoplasm volume, we used the value that was measured previously using fluorescently-labelled

dextran[22]. The nucleoplasmic volume was quantified in each mEGFP-Nup knock-in cell line using

a SiR-DNA signal as described previously[12].

Core regions were predicted on the NE based on a previously described protocol using the

core marker Lap-2α[12]. Briefly, nuclear volume was segmented using SiR-DNA fluorescence

signals that were processed with a 3D Gaussian filter and a multi-level thresholding. Nuclear

volume was then divided into inner and outer volumes using the cutting plane that was constructed

from the largest eigenvector and the second one orthogonal to the first vector of the pixel

coordinates of the nuclear volume. Surface area of each nucleus was calculated and utilized to

adjust the size of the inner and outer core regions at individual time points. The previously defined

criteria for being core and non-core regions[12] was applied. The position of inner and outer core

was determined with respect to the intersection point of the largest eigenvector on the nuclear

surface.

Mathematical modeling for the nuclear pore assembly kinetics

Previous EM data showed that, within 2 hours after anaphase onset, postmitotic assembly is the

dominant process in the non-core region, whereas the slower interphase assembly prevails in the

core region[12]. Assuming that this relation is also reflected in the live-cell Nup dynamics, we

derived a mathematical model. We assumed that the observed total fluorescence intensity in the

non-core, $n(t)$, and core region, $c(t)$, at time-point t after anaphase onset is a linear combination

of the postmitotic and interphase assembly processes according to

$$n(\text{t}) = f_N pm(\text{t}) + (1 - f_N)ip(\text{t}) \qquad (1)$$

$$c(\text{t}) = f_C \, pm(\text{t}) + (1 - f_C) \, ip(\text{t}), \qquad (2)$$

512 where $pm(t)$ and $ip(t)$ are the postmitotic and interphase assembly, respectively. The fraction of

513 postmitotic assembly in the non-core and core regions are denoted f_n and f_c, respectively. To test

514 this assumption and obtain an estimate of the fractions, we used a phenomenological model that

515 accounts for the observed sigmoid-like kinetics

$$pm(t) = \frac{t^{n_p}}{t^{n_p} + K_p^{n_p}}, \qquad \text{for } t \geq d \qquad (3)$$

$$ip(t) = \frac{t^{n_i}}{t^{n_i} + K_i^{n_i}}, \qquad \text{for } t \geq d \qquad (4)$$

and $pm(t < d) = ip(t < d) = 0$. The parameters n_p, K_p and n_i, K_i characterize the postmitotic

and interphase kinetics, respectively. The average time-point where a Nup is recruited for the

interphase and postmitotic assembly is K_i and K_p, respectively. The parameter d accounts for an

additional delay in NPC core region assembly with respect to the non-core region due to the

presence of kinetochore microtubule fibers[25]. For the non-core region $d = 0$. For the core region d

was estimated from the data.

For each assembly mechanisms, we also computed the recruitment duration from Eqs. 3–4.

For instance, the time required to recruit 80% of the protein, i.e. from a fraction of $\alpha_1 = 0.1$ up

to $\alpha_2 = 0.9$, is given by (see also Extended Data Fig. 4a)

$$\Delta T_p = K_p \left(\left(\frac{\alpha_2}{1 - \alpha_2} \right)^{\frac{1}{n_p}} - \left(\frac{\alpha_1}{1 - \alpha_1} \right)^{\frac{1}{n_p}} \right) \tag{5}$$

$$\Delta T_i = K_i \left(\left(\frac{\alpha_2}{1 - \alpha_2} \right)^{\frac{1}{n_i}} - \left(\frac{\alpha_1}{1 - \alpha_1} \right)^{\frac{1}{n_i}} \right). \tag{6}$$

525 In the model, we subtracted a background computed from the average of the first 3 time points.

526 All data is subsequently normalized to the mean value between 100 and 120 min after anaphase

527 onset. This yields corrected fluorescence intensities, $N(t_j)$ and $C(t_j)$. To find the model

528 parameters we minimized the mean squared distance between data and model for all the time

529 points M

$$\chi^2 = \sum_{j=1}^{M} \left(\left(\frac{N(t_j) - n(t_j)}{\sigma_N(t_j)} \right)^2 + \left(\frac{C(t_j) - c(t_j)}{\sigma_C(t_j)} \right)^2 \right), \tag{7}$$

where $\sigma_N(t_j)$ and $\sigma_C(t_j)$ are the respective standard deviation in the non-core and core region.

The parameter estimation has been performed in two steps. First, for each protein/cell-line,

we estimated the postmitotic fraction in the core and non-core region and the kinetic parameters.

All in all we computed 60 parameters (6 parameters per protein) from 4446 data points. We found

that the model well agrees with the data (Extended Data Figs. 2 and 3, $R^2 > 0.99$). For the

postmitotic fractions, we obtained on average $f_n = 0.857$ [0.76, 0.95] and $f_c = 0.295$ [0.17, 0.4],

where the number in brackets indicates the 95% confidence interval as estimated using the profile

likelihood method[36]. Importantly, the obtained postmitotic fractions are well in agreement with

the previously reported estimates obtained from EM-data[12]. The delay in pore formation between

core and non-core region was estimated by systematically varying d from 0 to 6 min in steps of 1

minute. A value of $d = 2$ min, gave optimal result. In a second step we used the previously

estimated average postmitotic fractions and d and recomputed the kinetics parameters for each

protein. The model with reduced parameters gave very similar results. To verify if the choice of

common postmitotic fractions for all Nups is valid, we computed the Baysian information criterion

(BIC) according to the previous report[37]. The difference in BIC between the model with reduced

parameters, 42 parameters for all proteins, compared to the full model, 60 parameters, was -7,

indicating that the model with reduced parameters is justified. The obtained parameter values are

listed in Extended Data Table 2.

Our validated model can be used to derive the underlying post-mitotic and interphase

assembly kinetics. Using Eqs. 1–2 we obtain

$$ip(t) = \frac{f_n c(t) - f_c n(t)}{f_n - f_c}. \tag{8}$$

$$pm(t) = \frac{(1 - f_c)n(t) - (1 - f_n)c(t)}{f_n - f_c}. \tag{9}$$

These equations can directly be applied to the experimental data $N(t)$ and $C(t)$ using the previously

estimated fraction f_n and f_c.

In Fig. 3c we see a strong positive correlation for the average time point of assembly and the

duration of assembly. One can prove that such a relation is expected for an irreversible sequential

assembly mechanism with linear rate constants. Using the definition for average time and

duration[38], the average time-point of a protein binding at step i is

$$\tau_i = \sum_{j=1}^{i} 1/k_j, \tag{10}$$

556 where k_j is the binding rate constant of a protein binding at step j to the complex. The duration of

557 assembly of a protein binding at step i is given by

$$\theta_i = 2\sqrt{\sum_{j=1}^{i} 1/k_j^2}. \tag{11}$$

From Eqs 10–11, it is clear that $\tau_{i+1} > \tau_i$ and $\theta_{i+1} > \theta_i$ for any parameter combinations.

Therefore, late binding proteins show a longer duration of assembly.

Structural modeling of NPC assembly pathway

A model of the assembly pathway is defined by a series of static structures, including a static

structure at each sampled time point along the assembly process. Therefore, we model the NPC

assembly by first modeling static structures at each time point, independently from each other. We

then enumerate alternative assembly pathways and rank them based on the static structure scores

and plausibility of transitions between successive static structures.

Integrative modeling of static structures at each time point

The static structures are modeled by standard integrative structure modeling[29], as follows.

Representing a static structure model

The time points correspond to times with available ET protein densities[11]: 5min, 6 min, 8 min, 10

mins, and 15 min after anaphase onset. We divide the mature NPC structure (PDB 5a9q, 5ijo) into

eight spokes and further divide each spoke into a set of rigid subcomplexes, including the Y-

complex, the inner ring Nup205-Nup155-Nup93 subcomplex, the inner ring Nup93-Nup188-

Nup155 subcomplex, and the central channel Nup62-Nup58-Nup54 subcomplex. For each

domain, we coarse-grained the structure by grouping 10 consecutive amino acid residues into a

single bead at the center of mass of those residues. Each subcomplex is represented as a rigid body.

The NE is represented as a fixed toroid surface embedded in two parallel planes. Thus, the

variables of the model include the Euclidean coordinates of the Nup subcomplexes and the copy

number of each Nup subcomplex.

We set the inner pore diameter and minor radius of the pore at each time point to the mean of

previously determined NE cross sections[11] with a pore diameter of 51.5 nm, 58.4 nm, 72.7 nm,

84.6 nm, 79.8 nm, and 87 nm; and minor radius of 21.4 nm, 21.2 nm, 21.5, 20.3 nm, 17.1 nm and

15 nm for time points at 5min, 6min, 8min, 10min, 15min, and the mature pore, respectively.

Scoring a static structure model

The copy numbers of the NPC subcomplexes at each time point were restrained by a Gaussian

function with mean and variance determined by the single-cell traces presented in this study. The

relative likelihood of a set of copy numbers is proportional to the product of individual Gaussian

likelihoods.

Distances between pairs of Nups that are in contact with each other in the native NPC

structure[26,27] were restrained by a harmonic Gō-like model[39]. Inter-subcomplex contacts within 5

nm in the mature structure were restrained by a harmonic function (strength 0.01 kcal/mol Å).

Each Gō-like scoring term was scaled at each time point, from zero at the first time point to full

strength at the mature pore time point. Distances between all pairs of Nups were also restrained

by a harmonic excluded volume restraint (strength 0.01 kcal/mol Å). Proximity between Nup

domains containing a membrane interacting ALPS-motif and the NE was restrained by a harmonic

term (strength 0.1 kcal/mol Å), based on their sequences. Overlap between the Nups and NE

surface was avoided by imposing a harmonic repulsion between the Nups and NE surface (strength

of 0.01 kcal/mol Å).

The shape of a static structure was restrained by a correlation coefficient between the model

and ET protein density[11]. The forward model density was represented by fitting each Nup

subcomplex with a Gaussian mixture model of two components per subcomplex copy using the

gmconvert utility[40]. Similarly, the ET protein densities at each time point were represented with a

Gaussian mixture model with 150 components fit to the experimental density.

Sampling static structure models

A state of the NPC at any given time point is defined by the copy numbers and coordinates of its

components. Only copy number assignments and structures consistent with the C-8 symmetries

were sampled. In addition, we only sampled structures for the top 20-scoring Nup copy number

combinations. Each sampling started with the mature pore structure, followed by applying 10^6

Monte Carlo moves. These moves included rotational and translational perturbations to each Nup

subcomplex, drawn from a uniform distribution in the range from -0.04 to +0.04 radians and from

-4 to +4 Å, respectively.

Modeling the assembly pathway

With the static structure models in hand, we connect them into complete alternative assembly

pathways, as follows.

Each pathway is represented by a static structure at each sampled time point, starting with t =

5 min and culminating in the native structure; we do not model the completely disassembled NPC.

The score of a pathway is the sum of the scores for the static structures on the pathway (defined

above) and transitions between them. A transition score is uniform for all allowed transitions. A

transition between two successive static structures is allowed if the subcomplexes in the first

structure are included in the second structure. All possible pathways were enumerated, scored, and

ranked. The best-scoring pathways were extracted for further analysis (Fig. 4).

Sample size determination and statistical analysis

For quantitative imaging in Fig. 1a, d, the data were from 4, 4, 4, 2, 3, 2, 3, and 2 independent

experiments for Nup107, Seh1, Nup205, Nup93, Nup62, Nup214, Tpr and Nup358, respectively.

STED imaging in Fig. 1b, c was from one independent experiment. For dynamic quantitative

imaging in Fig. 2, the data were from 4, 4, 4, 2, 3, 4, 3, 2, 2, and 4 independent experiments for

Nup107, Seh1, Nup205, Nup93, Nup62, Nup214, Tpr, Nup358, Nup153, and Pom121,

respectively. Statistical analyses were performed only after all the data were taken. Sample sizes

for each experiment are indicated in figure legends. Sample sizes were based on pilot experiments

to determine the number of cells required to observe stable population averages with high

Pearson's correlation between replicates. Videos of dividing cells with rotating nuclei are removed

from the analysis, because we cannot properly assign the non-core and core regions.

Data and code availability

Fluorescence   images   will   be   available   in   the   Image   Data   Resource   (IDR;

https://idr.openmicroscopy.org/)   or   the   BioImage   Archive

(https://www.ebi.ac.uk/biostudies/BioImages/studies). Our integrative spatiotemporal model of

the postmitotic assembly of the human NPC is available in PDB-Dev (https://pdb-

dev.wwpdb.org/) under the accession code XX (PDB submission in progress). Integrative

Modeling Platform (IMP) is an open source program freely available under the LGPL license at

http://integrativemodeling.org; all input files, scripts, and output files are available at

http://integrativemodeling.org/npcassembly.

Method references

32   Landry, J. J. *et al.* The genomic and transcriptomic landscape of a HeLa cell line. *G3*

647   *(Bethesda)* **3**, 1213–1224 (2013).

648   33   Farrants, H. *et al.* Chemogenetic Control of Nanobodies. *Nat. Methods* **17**, 279–282

649   (2020).

650   34   Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle

651   tracking. *Methods* **115**, 80–90 (2017).

35   Pruggnaller, S., Mayr, M. & Frangakis, A. S. A visualization and segmentation toolbox for

electron microscopy. *J. Struct. Biol.* **164**, 161–165 (2008).

36   Raue, A. *et al.* Structural and practical identifiability analysis of partially observed

dynamical models by exploiting the profile likelihood. *Bioinformatics* **25**, 1923–1929

(2009).

37   Kass, R.E. & Raftery, A. E. Bayes Factors. *J. Am. Stat. Assoc.* **90**, 773–795 (1995).

38   Heinrich, R., Neel, B. G., & Rapoport, T. A. Mathematical models of protein kinase signal

transduction. *Mol. Cell* **9**, 957-70 (2002).

39   Go, N. & Abe, H. Noninteracting local-structure model of folding and unfolding transition

in globular proteins. I. Formulation. *Biopolymers* **20**, 991–1011 (1981).

40   Kawabata, T. Gaussian-input Gaussian mixture model for representing density maps and

atomic models. *J. Struct. Biol.* **203**, 1–16 (2018).

Acknowledgments

We thank the EMBL Advanced Light Microscopy Facility (ALMF) for their support in STED

super-resolution microscopy; Ben Webb for help with IMP and Josh Baker-Lepain for managing

the Wynton computer cluster at QBI@UCSF. This work was supported by grants from the Baden

Wuerttemberg Foundation (J.E. and A.S.), NIH/NIGMS R01GM083960 (A.S.), NIH/NIGMS

P41GM109824 (A.S.), NIH/NIGMS R01GM112108 (A.S.) and the European Molecular Biology

Laboratory (EMBL; S.O., A.Z.P., A.R., M.J.H., M.K., A.C., B.K., J.E.). S.O. and A.R. were

further supported by the EMBL Interdisciplinary Postdoc Programme (EIPOD) under Marie Curie

Actions COFUND. S.O. was additionally supported by a JSPS fellowship (The Japan Society for

the Promotion of Science, postdoctoral fellowship for research abroad).

Author contributions

SO, JOBT, AS, and JE designed the project. SO performed all the fluorescence microscopy
experiments and analyses. JOBT performed integrated structural modeling. AZP carried out
mathematical modeling for the nuclear pore assembly kinetics. AR developed an analysis pipeline
for nuclear pore density measurement. MJH established a computational image analysis pipeline
to quantify fluorescence intensities in non-core and core regions in 3D time-lapse images. M.K.,
A.C. and B.K. generated genome-edited cell lines. AS and JE supervised the work. SO, JOBT,
AS, and JE wrote the paper. All authors contributed to the analysis and interpretation of data and
provided input on the manuscript.

Competing interest declaration

The authors declare no competing interests.

Additional information

The manuscript contains supplementary material (Extended Data Figs. 1 to 4 and Extended Data
Tables 1 to 3).