1   **SARS-like coronaviruses in horseshoe bats (*Rhinolophus* spp.) in Russia, 2020.**

2   Sergey V. Alkhovsky, Sergey V. Lenshin, Alexey V. Romashin, Tatyana V.

3 Vishnevskaya, Oleg I. Vyshemirsky, Yulia I. Bulycheva, Dmitry K. Lvov, Asya K. Gitelman.

5   **Address for correspondence**: Sergey V. Alkhovsky, N.F Gamaleya National Research

6 Center on Epidemiology and Microbiology of Ministry of health of Russian Federation, 18

7 Gamaleya street, 123098 Moscow, Russia; email: salkh@ya.ru, s_alkhovsky@gamaleya.org.

9   **Keywords:** SARS-CoV, SARS-CoV-2, bat SARS-like coronaviruses, SARS-CoV-like

10 viruses, viral metagenomics, coronavirus, horseshoe bats, zoonotic viruses, Rhinolophus.

12   **Affiliations:**

13   D.I. Ivanovsky Institute of Virology of N.F. Gamleya National Center on Epidemiology

14 and Microbiology of Ministry of Health of Russian Federation, Moscow, Russia (S. Alkhovsky,

15 D. Lvov, T. Vishnevskaya, Yu. Bulycheva, A. Gitelman);

16   Reference center on coronavirus infection of N.F. Gamleya National Center on

17 Epidemiology and Microbiology of Ministry of Health of Russian Federation, Moscow, Russia

18 (S. Alkhovsky, T. Vishnevskaya);

19   Federal State Budgetary Scientific Institution «Scientific Research Institute of Medical

20 Primatology» of Russian Academy of Science, Adler, Russia (S. Lenshin, O. Vyshemirsky);

21   Federal State Budgetary Institution «Sochi National Park», Sochi, Russia (A. Romashin)

Abstract

We found and genetically described two novel SARS-like coronaviruses in feces and oral

swabs of the great (*R. ferrumequinum*) and the lesser (*R. hipposideros*) horseshoe bats in

southern region of Russia. The viruses, named Khosta-1 and Khosta-2, together with related

viruses from Bulgaria and Kenya, form a separate phylogenetic lineage. We found an evidence

of recombination events in evolutionary history of Khosta-1, which involved the acquisition of

structural proteins S, E, and M as well as nonstructural genes ORF3, ORF6, ORF7a, and ORF7b

from a virus that is closely related to Kenyan isolate BtKY72. Examination of bats by RT-PCR

revealed that 62,5% of great horseshoe bats in one of the caves were positive for Khosta-1 virus

while its overall prevalence was 14%. The prevalence of Khosta-2 was 1,75%. Our results show

that SARS-like coronaviruses circulate in horseshoe bats in the region and provide a new data on

their genetic diversity.

Introduction

Horseshoe bats (*Rhinolophidae: Rhinolophus*) are considered as a main natural reservoir

and source of zoonotic coronaviruses (CoV) which caused epidemic outbreak of severe acute

respiratory syndrome (SARS) and COVID-19 pandemic in 2002 and 2019, respectively (1,2).

These viruses, designated as SARS-CoV and SARS-CoV-2, together with related viruses found

in bats and other animals (SARS-like coronaviruses or SARS-CoV-like viruses) belong to the

subgenus *Sarbecovirus* of the genus *Betacoronavirus* of the family *Coronaviridae* (3). Horseshoe

bats are wide distributed in Asia, Europe, and North Africa. In East Asia (in particular, in

People's Republic of China (PRC)) SARS-CoV-like viruses circulate in multiple rhinolophid's

species, however the Chinese rufous (*R. sinicus*), the greater (*R. ferrumequinum*), the

intermediate (*R. affinis*), and the king (*R. rex*) horseshoe bats seem to be of major importance (4).

In Europe SARS-CoV-like viruses was found in the greater, the lesser (*R. hipposideros*), the

Miditerranean (*R. euryale*), Mehely's (*R. mehelyi*), and Blasius's (*R. blasii)* horseshoe bats (5–8).

The prevalence of SARS-like coronaviruses among bats in different caves-colonies can vary

from 0% to 60% (4,7,9,10). In Russia three species of horseshoe bats (the greater, the lesser, and

Miditerranean) are common in southern regions, lying below about 44º north latitude, mostly

including North Caucasus and Crimea. In present work we hypothesized that SARS-like

coronaviruses circulate in the region in local populations of horseshoe bats. To test this

hypothesis, we examined the colonies of bats located in the southern macroslope of the Greater

Caucasus on northern coast of the Black Sea in Russia. Using metagenomic analysis we found

and genetically described two new SARS-like coronaviruses in feces and oral swabs of the

greater and the lesser horseshoe bats. Further PCR analysis showed a high degree of infection of

bats with discovered viruses in some locations.

Materials and methods

The samples from bats were collected in Sochi National Park (Sochi-Adler, Krasnodar

krai, Russia) and surrounding areas in March-October 2020. The Sochi National Park is located

on southern macroslope of the Greater Caucasus descending to the northern coast of the Black

Sea (Figure 1). The park stuffs keep records of more than 300 karst formations (caves, breaks,

mines, clefts, etc.) that are natural refuges for bats and other troglophilous animal species. Bats

were caught by hand in eight locations including five caves as well as basements and attics of

houses (Table 1). The bats were caught in the frame of ongoing surveillance of bats populations

constantly carried out in the park. The species of the animals was determined based on their

morphological characters by an experienced park zoologist. The length of the forearm and

weight of animals were measured. To collect bat oral swabs (saliva and buccal cells) an

urological swab was placed in the bat's mouth for 10-15 sec and then placed in 250 µl of

phosphate-buffered saline (PBS). To collect feces, an animal was placed in a small white cotton

bag for 10-15 min. After that the animal was released, and the feces were collected from the

walls of the bag into cryovials. No bats were harmed during sample collection. A total of 120

samples of oral swabs and 77 samples of feces from five species of bats were collected (Table 1).

The samples were delivered to the laboratory on ice and stored at -70ºC until start of research.

Metagenomic analysis was conduct as described anywhere. Briefly, the feces were

suspended and homogenized in 0,5 ml of PBS and pooled by 0,1 ml in three samples (faces from

20-30 animals in the pool). Pooled samples were clarified by centrifugation (10 000 g, 15 min)

and treated with DNase I and RNase I_f (NEB, Great Britain) for removing of naked out-capsid

nucleic acid. The viral particles were sedimented from treated samples by ultracentrifugation

(30 000 g, 1 h) through 2 ml of 20% sucrose. Virus plaque was resuspended in 0,5 ml PBS. Total

RNA was isolated from 0,25 ml of obtained solution with TRIzol LS reagent (ThermoFisher

Scientific, USA).

Total RNA from oral swabs were isolated with TRIzol LS reagent from individual samples

and pooled by 20 µl in five pooled samples (20-25 samples in the pool). The pooled RNA was

precipitated by isopropyl alcohol with addition of glycogen followed by additional clarification

with RNeasy MinElute Cleanup kit (Qiagen, Germany). NEBNext rRNA Depletion kit (NEB)

was used to remove bacterial and eukaryotic rRNA from total RNA isolated from pooled

samples. Treated RNA was used for cDNA library preparation by NEBNext Ultra II RNA library

kit for Illumina (NEB).

The libraries were sequenced on a HiSeq4000 instrument (Illumina, USA) at the facility of

Resource Center "BioBank" of the Research Park of Saint Petersburg State University (Saint-

Petersburg, Russia). Reads were filtered by quality, trimmed to remove adapter's sequences, and

assembled *de novo* using CLC Genomics Workbench 7.0 software (Qiagen). Obtained contig

sequences were analyzed using blastx algorithm by DIAMOND software (11) against nr

'Viruses' database that included all reference viral sequences available in GenBank at December

2020. Nucleotide and deduced amino acid sequences were alignment by ClustalW implemented

in MEGAX software (12). The best substitution model was evaluated for each alignment by the

Model selection module in MEGAX software. Phylogenetic trees were inferred by 'maximum

likelihood' method using appropriate model with 1000 bootstrap replicates by MEGAX

software. Similarity plot analysis was conducted by SimPlot software (13). Possible

recombination was analyzed by RDP5 software (14).

Primers and probes for specific detection of discovered coronaviruses were developed

based on obtained sequences by Beacon 7.0 software (Premier Biosoft, USA). Real-time RT-

PCR was conducted with TaqPath 1-Step Multiplex Master Mix (ThermoFisher Scientific, USA)

and total RNA isolated by TRIzol LS reagent from individual oral swabs and faces samples.

Results

Results of sequencing of the samples. In total, 124 522 978 reads for three pooled fecal

samples and 170 112 341 reads for five pooled oral swab samples were obtained. The reads were

de novo assembled in contigs and analyzed by blastx algorithm for presence of viral sequences.

The search results revealed two extended contigs with a length of approximately 29 Kb with

open reading frames (ORFs) with similarity to members of the genus *Betacoronavirus* in the

pools one and three of fecal samples, respectively. With further analysis, a near-complete

genome of two novel SARS-like coronaviruses was sequenced. Matching contigs have also been

found in corresponding oral swab samples, but with smaller length and coverage. Two found

SARS-like coronaviruses were named BtCoV/Khosta-1/Rh/Russia/2020 and BtCoV/Khosta-

2/Rh/Russia/2020 and placed in GenBank by accession numbers MZ190137 and MZ190138,

respectively. Bellow they are referred to as Khosta-1 and Khosta-2, respectively.

Genetic and phylogenetic analysis. The genomic organization of Khosta-1 and Khosta-2

is similar to that of other SARS-like coronaviruses (Figure 2). Approximately two thirds of the

genome of coronaviruses is occupied by ORF1a and ORF1b genes which encode the proteins of

the replicative complex and translated as ORF1ab polyprotein due to ribosomal shifting. The rest

of the genome contains genes of structural proteins (S, E, M, and N), which form a virion, as

well as several non-structural proteins (ORF3, ORF6, ORF7, ORF8, ORF9, and ORFX), the

presence and structure of which varies in different viruses (3). Genome organization of Khosta-1

and Khosta-2 has the greatest similarity with BtCoV/BM48-31/2008 and BtKY72 viruses – two

SARS-like coronaviruses found in horseshoe bats in Bulgaria and Kenya in 2008 and 2007,

respectively (8,15). Their peculiarity is the absent of ORF8 gene which is common in bat SARS-

like coronaviruses from East and Southeast Asia (Figure 2).

Pairwise alignments of the deduced proteins of Khosta-1 and Khosta-2 virus with those of

other SARS-like coronaviruses showed its highest similarity with BtCoV/BM48-31/2008 and

BtKY72 viruses (Table 2). Khosta-1 is closest related to BtCoV/BM48-31/2008 with 92,5% aa

and 99% aa identity in the conservative ORF1a and ORF1b proteins, respectively. Similarity of

Khosta-1 with SARS-CoV and related viruses from China are on average 81,5% aa identity in

ORF1a protein and 96% in ORF1b protein. Comparison Khosta-1 with SARS-CoV-2 viruses

revealed 77,5% and 94,2% aa identity for ORF1a and ORF1b proteins, respectively. Despite the

high similarity of Khosta-1 and BtCoV/BM48-31/2008 in ORF1a and ORF1b proteins, structural

proteins S, E, and M of Khosta-1 are more similar to those of Kenyan virus BtKY72. Khosta-1

and BtKY72 share 89,1%, 98,7%, and 97,29% aa identity for S, E, and M proteins, whereas

these values for Khosta-1 and BtCoV/BM48-31/2008 are 84,37%, 89,47, and 95%, respectively.

N protein of Khosta-1 is more similar to that of BtCoV/BM48-31/2008 (96,64%aa identity) than

BtKY72 (92,6% aa identity).

In contrast, Khosta-2 does not have such an increased similarity with some group of

sarbecoviruses and has 79-81% aa identity with SARS-CoV viruses and 76-77% with SARS-

CoV-2 viruses in ORF1a protein. ORF1b protein of Khosta-2 has 93,5-95% aa identities with all

other bat SARS-like coronaviruses. Comparison of proteins of Khosta-1 and Khosta-2 showed

that these viruses differ from each other at about the same level as Khosta-2 differs from other

bat SARS-like coronaviruses (Table 2).

Recombination analysis. Genome sequence similarity between Khosta-1, Khosta-2,

BtCoV/BM48-31/2008, and BtKY72 was analyzed by Simplot and RDP5 software (Figure 2A-

2C). Simplot analysis showed high degree of similarity of Khosta-1 and BtCoV/BM48-31/2008

in the ORF1ab and N genes and a decrease in the similarity in the S-ORF7b region. Results of

analysis carried out by RDP5 software using different methods implemented in the program

showed clear signals of recombination events in evolutionary history of Khosta-1 (Figure 2C).

Recombination events presumably included acquisition of S-ORF7b region by ancestor of

Khosta-1 virus from a virus closely related to BtKY72.

Phylogenetic analysis. Phylogenetic analysis based on ORF1ab protein sequences showed

that Khosta-1, Khosta-2, BtCoV/BM48-31/2008, and BtKY72 form a monophyletic lineage

located between SARS-CoV and SARS-CoV-2 lineages of the *Sarbecovirus* subgenus (Figure

3A). The separate cluster this group of viruses also formed on phylogenetic tree based on

nucleotide sequences of S gene (Figure 3B). Topology of tree confirms the probable

recombination event in evolutionary history of Khosta-1. In ORF1ab tree Khosta-1 is grouped

together with BtCoV/BM48-31/2008, while in S gene tree with BtKY72.

Analysis of receptor binding motif (RBM) of S protein. Alignment of amino acid

sequences of RBM of Khosta-1 and Khosta-2 with certain sarbecoviruses are presented in

Figure 4. This is a highly variable region where multiple substitutions and deletions occur among

SARS-CoV related viruses. Khosta-1 and Khosta-2 as well as BtCoV/BM48-31/2008 have a

common deletion of four aa in the N-part of RBM. This deletion partially overlapping with the

deletion that is characteristic of HKU3-1 and related strains of bat SARS-CoV-like viruses that

unable to bind angiotensin-converting enzyme 2 (ACE2) receptor. We analyzed aa positions in

RBM which are thought to be crucial for binding of ACE2 receptor and, therefore, important for

adaptation of bat SARS-like coronaviruses to human (16,17). Only at position 442 Khosta-1 and

Khosta-2 share a common amino acid (L), which is also inherent in SARS-CoV-2 and related

viruses. Despite the significant genetic distance between Khosta-1 and BtKY72, crucial positions

in RBM and their context coincide between them. In contrast, position 479, 480, and 487 of

Khosta-2 coincides poorly with other groups of viruses (Figure 4).

PCR testing. We developed primers and probes for specific detection of Khosta-1 and

Kosta-2 viruses in faces and oral swabs by real time RT-PCR (Appendix Table). Results of PCR

testing of samples are presented in Table 1. RNA of Khosta-1 was detected mostly in the greater

horseshoe bats collected in Kolokolnaya cave. All four found positive oral swabs belonged to

animals with positive fecal samples. In other locations Khosta-1 virus was detected only in two

fecal samples – from the greater horseshoe bat from Khosta 1 cave and the lesser horseshoe bat

from Partizanskaya cave, respectively. Also note that RNA of Khosta-1 was detected in feces

much more often than in oral swabs and in a higher titer (based on Ct value, data not shown).

RNA of Khosta-2 virus was detected in two lesser horseshoe bats collected in basement of the

building at Research Institute of Medical Primatology. In one animal RNA of Khosta-2 was

detected in both faces and oral swabs, and in another only swab.

Discussion

The emergence of SARS-CoV and SARS-CoV-2 viruses is a result of adaptation of SARS-

CoV-like viruses, circulated in the horseshoe bats, to human (18,19). Horseshoe bats are

widespread and, presumably, SARS-like coronaviruses circulate in all parts of their range

including Asia, Europe and northern Africa. However, little information exists on the genetic

diversity of bat SARS-like coronaviruses in the regions outside East and Southeast Asia. We

described here two novel SARS-like coronaviruses circulated in the horseshoe bats in southern

region of Russia. Khosta-1 and Khosta-2 viruses are closely related to viruses recently described

in Bulgaria (strain BtCoV/BM48-31/2008) and Kenya (strain BtKY72) (8,15). Together they

form a separate "western" (as they are found in the western part of the range of horseshoe bats)

phylogenetic lineage of bat SARS-like coronaviruses. A feature of these viruses is the absence of

ORF8 gene, which is common in SARS-CoV, SARS-CoV-2, and most of bat SARS-like

coronaviruses of eastern lineages.

SASR-CoV and SARS-CoV-2 recognize host angiotensin-converting enzyme 2 (ACE2) as

its receptor. Crucial for binding of ACE2 receptor amino acids (442, 487, 479, 487, and 491) are

located in the RBM of the S protein (17,20). These amino acids and its context in RBM of

Khosta-1 and Khosta-2, like most other bat SARS-like coronaviruses, are quite different from

SASR-CoV and SARS-CoV-2 viruses. The most bat SARS-like coronaviruses unable to bind

ACE2 receptor of human and do not infect its cells (21). However, several strains of bat SARS-

like coronaviruses that can use ACE2 receptor and whose proteins are highly similar to SARS-

CoV or SARS-CoV-2 have been recently found in the Chinese rufous (*R. sinicus*) and the

intermediate (*R. affinis*) horseshoe bats in People's Republic of China (22–25). In this vein, it is

argued that direct progenitors of SARS-CoV and SARS-CoV-2 originated after sequential

recombination events between bat SARS-like coronaviruses. Since the recombination is of great

importance in the evolution of coronaviruses, we analyzed possible recombination events in the

western lineage of bat SARS-like coronaviruses. Despite the small number of known full-length

sequences (only four), we observe an evidence of recombination in evolutionary history of

Khosta-1. The alleged recombination event involved the acquisition of structural proteins S, E,

and M as well as nonstructural genes ORF3, ORF6, ORF7a, and ORF7b from a virus that is

closer to the Kenyan isolate BtKY72 than to the European strain BtCoV/BM48-31/2008. Based

on this, we can assume that genetic diversity of viruses in the region is significantly higher than

is known today and there is constant exchange of genes between them. These findings require

further investigation of the diversity of circulating variants, with particular emphasis on the

diversity of the S gene.

Using RT-PCR we showed that 14% of tested horseshoe bats were carriers for Khosta-1

virus and 1,75% for Khosta-2 virus. However, most of the Khosta-1 positive samples were found

in only one cave (Kolokolnaya cave) where infection rate of greater horseshoe bats reached

62,5%. This bias, together with the small number of samples from other locations, makes it

difficult to accurately estimate the prevalence of Khosta-1 in the region and requires further

research. The closest European region where such studies have been carried out is Bulgaria:

according to the data obtained by J. Drexler with coauthor (2010) SARS-like coronaviruses were

detected in 13,3% of greater horseshoe bat, 15,9% of Blasius's horseshoe bat, 30,8% of

Mehely's horseshoe bat, and 32,1% of Miditerranean horseshoe bat (8). Another studies found

38.8% positive lesser horseshoe bats in Slovenia and 37,9% positive greater horseshoe bats in

France (5,6). All this data show that the prevalence of SARS-like coronaviruses in horseshoe

bats in their western part of range can vary widely between different species, locations, and

possibly the time of year of observation.

In conclusion, we have shown that SARS-like coronaviruses circulate in horseshoe bats in

southern region of Russia and provide a new information on its genomics. Genetic diversity,

prevalence, host range, as well as potential threat to human of these viruses remain to be

determined.

Acknowledgments

The work was funded by Russian Foundation for Basic Research (RFBR), project

No**.** 20-04-60154.

Author Bio (first author only, unless there are only 2 authors)

Sergey V. Alkhovsky is a head of laboratory of biotechnology at D.I. Ivanovsky Institute

of Virology of N.F. Gamleya National Center on Epidemiology and Microbiology in Moscow,

Russia. His scientific interests are ecology, genomics, and evolution of zoonotic viruses with

special emphasizing to emerging and reemerging viruses that pose serious threat to public health.

It includes arboviruses as well as zoonotic viruses of rodents, bats, and birds, distributed in

Northern Eurasia.

References

1.   Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, Wong BH, et al. Severe acute respiratory

syndrome coronavirus-like virus in Chinese horseshoe bats. Proc Natl Acad Sci U S A.

2005;102(39):14040–5.

2.   Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, et al. Bats are natural reservoirs of

SARS-like coronaviruses. Science. 2005; 310(5748):676–9.

3.   de Groot RJ, Baker SC, Baric R, Enjuanes L, Gorbalenya AE, Holmes K V, et al. Family

Coronaviridae. In: King AM, Adams MJ, Carstens EB, Lefkowitz EJ, editors. Virus

taxonomy : classification and nomenclature of viruses : ninth report of the International

Committee on Taxonomy of Viruses. 1sted ed. London: Elsevier; 2012. p. 806–28.

4.   Fan Y, Zhao K, Shi Z-L, Zhou P. Bat Coronaviruses in China. Viruses. 2019;11(3):210.

5.   Rihtarič D, Hostnik P, Steyer A, Grom J, Toplak I. Identification of SARS-like

coronaviruses in horseshoe bats (Rhinolophus hipposideros) in Slovenia. Arch Virol.

2010;155(4):507–14.

6.   Ar Gouilh M, Puechmaille SJ, Diancourt L, Vandenbogaert M, Serra-Cobo J, Lopez Roïg

M, et al. SARS-CoV related Betacoronavirus and diverse Alphacoronavirus members

found in western old-world. Virology. 2018;517:88–97.

7.   Balboni A, Palladini A, Bogliani G, Battilani M. Detection of a virus related to

betacoronaviruses in Italian greater horseshoe bats. Epidemiol Infect. 2011;139(2):216–9.

8.   Drexler JF, Gloza-Rausch F, Glende J, Corman VM, Muth D, Goettsche M, et al.

Genomic Characterization of Severe Acute Respiratory Syndrome-Related Coronavirus in

European Bats and Classification of Coronaviruses Based on Partial RNA-Dependent

RNA Polymerase Gene Sequences. J Virol. 2010;84(21):11336–49.

9.   Wang LF, Shi Z, Zhang S, Field H, Daszak P, Eaton BT. Review of bats and SARS.

Emerg Infect Dis. 2006;12(12):1834–40.

10.   Drexler JF, Corman VM, Drosten C. Ecology, evolution and classification of bat

coronaviruses in the aftermath of SARS. Antiviral Res. 2014;101:45–56.

11.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.

Nature Methods. 2015.12(1):59–60.

12.   Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular

evolutionary genetics analysis using maximum likelihood, evolutionary distance, and

maximum parsimony methods. Mol Biol Evol. 2011;28(10):2731–9.

13.   Lole KS, Bollinger RC, Paranjape RS, Gadkari D, Kulkarni SS, Novak NG, et al. Full-

length human immunodeficiency virus type 1 genomes from subtype C-infected

seroconverters in India, with evidence of intersubtype recombination. J Virol.

1999;73(1):152–60.

14.   Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. RDP4: Detection and analysis of

recombination patterns in virus genomes. Virus Evol. 2015;1(1):vev003.

15.   Tong S, Conrardy C, Ruone S, Kuzmin I V, Guo X, Tao Y, et al. Detection of novel

SARS-like and other coronaviruses in bats from Kenya. Emerg Infect Dis.

2009;15(3):482–5.

16.   Li F. Receptor recognition and cross-species infections of SARS coronavirus.

2013;100(1):246-54.

17.   Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor Recognition by the Novel

300   Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of

301   SARS Coronavirus. J Virol. 2020;94(7):e00127.

302   18.   Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, et al. Bats are natural reservoirs of

303   SARS-like coronaviruses. Science. 2005;310(5748):676–9.

304   19.   Zhou P, Yang X Lou, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak

305   associated with a new coronavirus of probable bat origin. Nature. 2020; 579(7798):270-

306   273.

307   20.   Li W, Moore MJ, Vasllieva N, Sui J, Wong SK, Berne MA, et al. Angiotensin-converting

308   enzyme 2 is a functional receptor for the SARS coronavirus. Nature. 2003 Nov

309   27;426(6965):450–4.

310   21.   Ren W, Qu X, Li W, Han Z, Yu M, Zhou P, et al. Difference in Receptor Usage between

311   Severe Acute Respiratory Syndrome (SARS) Coronavirus and SARS-Like Coronavirus of

312   Bat Origin. J Virol. 2008 Feb 15;82(4):1899–907.

313   22.   Ge XY, Li JL, Yang X Lou, Chmura AA, Zhu G, Epstein JH, et al. Isolation and

314   characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. Nature.

315   2013;503(7477):535–8.

316   23.   Hu B, Zeng LP, Yang X Lou, Ge XY, Zhang W, Li B, et al. Discovery of a rich gene pool

317   of bat SARS-related coronaviruses provides new insights into the origin of SARS

318   coronavirus. PLoS Pathog. 2017; 13(11):e1006698.

319   24.   Lau SKP, Feng Y, Chen H, Luk HKH, Yang W-H, Li KSM, et al. Severe Acute

320   Respiratory Syndrome (SARS) Coronavirus ORF8 Protein Is Acquired from SARS-

321   Related Coronavirus from Greater Horseshoe Bats through Recombination. J Virol. 2015

Oct 15;89(20):10532–47.

25. Ge XY, Wang N, Zhang W, Hu B, Li B, Zhang YZ, et al. Coexistence of multiple

coronaviruses in several bat colonies in an abandoned mineshaft. Virol Sin. 2016 Feb

1;31(1):31–40.

Table 1. The bat samples collected and results of RT-PCR testing for Kosta-1 and

Khosta-2 viruses.

Location	Bat species	Number of samples collected		Khosta-1 virus positive samples (%*)		Khosta-2 virus positive samples (%*)	
		Oral swabs	Faces	Oral swabs	Faces	Oral swabs	Faces
Basement of the building at Research Institute of Medical Primatology (43°26'06.3"N 39°59'26.4"E)	Lesser horseshoe bat (*R. hipposideros*)	27	24	0	0	1 (3,7%)	2 (8,3%)
	Miditerranean horseshoe bat (*R. euryale*)	1	1	0	0	0	0
Museinaya cave (43°33'34.3"N 39°53'46.2"E)	Greater horseshoe bat (*R. ferrumequinum*)	4	2	0	0	0	0
	Lesser horseshoe bat (*R. hipposideros*)	3	2	0	0	0	0
Khosta 1 cave (43°33'49.5"N 39°53'57.2"E)	Greater horseshoe bat (*R. ferrumequinum*)	21	13	0	1 (7,7%)	0	0
	Common bent-wing bat *Miniopterus schreibersii*	3	1	0	0	0	0
Kolokolnaya cave (43°33'08.3"N, 39°56'02.4"E)	Greater horseshoe bat (*R. ferrumequinum*)	36	24	4 (11%)	15 (62,5%)	0	0

	Miditerranean horseshoe bat (*R. euryale*)	2	0	0	0	0	0
Partizanskaya cave (43°37'38.86"N, 39°54'46.06"E)	Greater horseshoe bat (*R. ferrumequinum*)	2	1	0	0	0	0
	Lesser horseshoe bat (*R. hipposideros*)	5	3	0	1 (33%)	0	0
Attic of house (44° 0'57.51"N, 39°15'3.63"E)	Lesser horseshoe bat (*R. hipposideros*)	6	4	0	0	0	0
Krasnoaleksandrovskaya cave (44° 0'57.21"N, 39°21'49.68"E)	Greater horseshoe bat (*R. ferrumequinum*)	1	0	0	0	0	0
	Lesser horseshoe bat (*R. hipposideros*)	6	0	0	0	0	0
	Myotis bat *Myotis* spp.	3	0	0	0	0	0
Attic of house, Izmaylovka village (43°37'51.72"N 39°49'45.38"E)	Lesser horseshoe bat (*R. hipposideros*)	0	2	0	0	0	0
Total		120	77	4 (4,6%**)	17 (14,9%**)	1 (0,89%**)	2 (1,75%**)

* the percentages quoted are indicative and not statistically reliable.

** value calculated only for horseshoe bats, common bent-wind bat and myotis bat were

excluded from the calculation.

Table 2. Similarity (%) of deduced amino acid sequences of proteins of Khosta-1 and

Khosta-2 viruses with certain representatives of *Sarbecovirus* subgenus (lineage B of

betacoronaviruses).

Protein	Viruses	Amino acid identity (%)								
		Bat SARS-CoV-	Bat SARS-CoV-	Bat SARS-CoV-like (China,	Civet SARS-CoV-	SARS-CoV	Bat SARS-CoV-2-	Pangolin SARS-	SARS-CoV-2	Khosta-1 vs Khosta-2

		like BGR/2008 (Bulgaria, 2008)	like BtKY72 (Kenya, 2007)	2005-2016)*	like SZ3 (China, 2003)	Urbani (2003)	like RatG13 (China, 2013)	CoV-2-like (China, 2017)	Wuhan-Hu-1 (2019)	
ORF1a	Khosta-1	92,95	84,6	81,53-81,6	81,67	81,53	77,2	77,89	77,32	82
	Khosta-2	81,1	80,9	79,4-79,6	79,5	79,4	76,3	77,1	76,45	
ORF1b	Khosta-1	99,07	96,3	95,82-96,3	96,15	96,15	94,22	94,22	94,21	94,75
	Khosta-2	94,7	93,7	94,9-95,17	95,02	94,9	93,44	93,47	93,5	
S	Khosta-1	84,37	89,11	75,5-76,2	75,7	75,7	73,0	72,4	72,22	82
	Khosta-2	79,54	79,7	73,03-73,9	73,2	73,0	72,5	71,74	72,54	
ORF3	Khosta-1	85,98	86,7	66,8-72,3	70,8	70,8	65,1	66,2	64,7	81,8
	Khosta-2	77,9	82,22	67,9-69,34	67,15	67,5	64,5	65,8	63,27	
E	Khosta-1	89,47	98,7	87,0	87	87	93,42	93,42	93,42	94,7
	Khosta-2	88,16	94,74	90,7	90,7	90,7	89,5	89,5	89,5	
M	Khosta-1	95,0	97,29	91,86-92,31	92,31	91,86	88,24	87,73	88,13	91
	Khosta-2	90,9	90,5	88,7-89,6	90,5	89,6	87,3	87,27	87,0	
ORF6	Khosta-1	68,25	63,0	49,21-52,38	49,21	49,21	50,82	50,82	50,82	58,73
	Khosta-2	58,1	58,1	44,4-47,6	46,03	46,03	46,7	46,7	46,7	
ORF7a	Khosta-1	69,7	70,6	58-59,7	61,34	61,34	58,5	59,32	58,5	73,5
	Khosta-2	63,25	70,34	58,2-59,26	60,0	60,0	60,0	58,3	59,13	
ORF7b	Khosta-1	86,05	81,4	71,8	71,8	71,8	61,5	71,8	74,4	70,7
	Khosta-2	71,4	73,1	64,2	64,2	64,2	64,2	64,2	64,2	
N	Khosta-1	96,64	92,6	88,36-88,9	89,1	89,1	87,9	87,6	87,4	91,85
	Khosta-2	91,13	90,21	85,75-86,73	86,5	86,5	85,5	86,4	85,24	

Figure 1. Map of the region where bat samples were collected. Location of Sochi

National Park and surrounding area is shown in grey.

Figure 2. Scheme of the genome of Khosta-1 and Khosta-2 viruses with the designation

of the main ORFs, and results of Simplot and recombination analysis. **A**. Khosta-1 was used as a

query sequence and Khosta-2, BM48-31/BGR/2008, and BtKy72 viruses were used as reference

sequences. **B**. Khosta-2 was used as a query sequence and Khosta-1, BM48-31/BGR/2008, and

BtKy72 were used as reference sequences. The analysis was performing with Kimura (2-

parametr) model, window size of 1000 bases and a step size of 100 bases. **C.** Results of bootstrap

analysis of recombination events in Khosta-1 genome by RDP5 software.

Figure 3. Phylogenetic trees inferred by maximum likelihood method based on analysis

of amino acid sequences of ORF1ab protein (**A**) and nucleotide sequences of S gene (**B**). HKU9-

like strain Bt-BetaCoV/GX2018, belonged to subgenus *Nobecovirus* (lineage D), was used as an

outer group for ORF1ab protein phylogeny. The viruses described in present work are marked by

black circle.

Figure 4. Alignment of receptor binding motif (RBM) of receptor binding domain (RBD)

of S protein of Khosta-1 and Khosta-2 with certain sarbecoviruses. Position (442, 472, 479, 480,

487, 491 numbering by SARS-CoV Urbani) in RBM which are thought to be important for

adaptation SARS-CoV-like viruses to human ACE2 receptor (16,17) are bolded. Bat SARS-

CoV-like viruses that are capable or not capable of utilizing ACE2 receptor are marked with

"ACE2(+)" or "ACE2(-)", respectively.

Figure 1.

Figure 2A

Figure 2B.

Figure 2C.

Figure 3A

377   **Figure 3B.**

Figure 4.