How to describe and measure phenology? An investigation on the diversity of metrics using phenology of births in large herbivores

Lucie Thel[1], Simon Chamaillé-Jammes[2, 3, 4] & Christophe Bonenfant[1, 4]

Authors details

[1] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Bâtiment 711, Université Lyon I, 43 Boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France.

[2] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier, France.

[3] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

[4] LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe.

Correspondence author: Lucie Thel, lucie.mc.thel@gmail.com.

Abstract

Proposed in 1849 by Charles Morren to depict periodical phenomena governed by seasons, the term "phenology" has spread in many fields of biology, from evolutionary to molecular biology. The wide adoption of the concept of phenology has been associated with the publication of a large number of metrics, with sometimes confusion about what they measure. Here, we aimed to link these metrics to four characteristics of phenology: timing, synchrony, rhythmicity and regularity. From 52 published metrics previously used to quantify phenology of births in large herbivores, we conducted a quantitative analysis based on simulations of normal and non-normal distributions of births (both intra- and inter-annual variations). We first compared and classified the metrics using a correlation matrix. Then, we evaluated the ability of each metric to capture the variation of the four phenology characteristics via a sensitivity analysis. We finally scored each metric according to eight criteria we thought were important to describe phenology correctly. The high correlation we found among the metrics corresponding to each of the four characteristics of phenology suggests that such diversity of metrics is not useful. Additionally, we show that the best metrics are not necessarily the most commonly used, and that simpler is often better. Mean vector orientation and mean vector length seem particularly suitable to describe the timing and synchrony (respectively) of births in a wide range of phenology patterns. The level of repeatability of both timing and synchrony is still poorly described, but Mood and Kolmogorov tests allow a first and easy quantification of this variability across the years. By identifying, amongst a large diversity of metrics, those that appear the most suitable, our study can facilitate individual and comparative studies of phenology, particularly important in the context of climate change.

Keywords: regularity, rhythmicity, seasonality, synchrony, timing, ungulate

Introduction

In 1849, Charles Morren coined the term "phenology" to describe how periodical phenomena such as plant growth and reproduction are governed by the course of seasons (Morren 1849, see also Demarée 2011). With his observations, he opened a new field of research, and almost two centuries later the concept of phenology has become a cornerstone of ecology (Begon *et al.* 1986), used in plant and animal ecology indifferently (Forrest and Miller-Rushing 2010). By describing when particular life-history events (*e.g.* flowering, parturition) occur in relation to the characteristics or states of the individual (*e.g.* size, age) as well as to external factors (*e.g.* photoperiod, predation risk) the concept of phenology is key to understanding the temporal cycles in the life cycle of species (Forrest and Miller-Rushing 2010). Nowadays the term phenology is commonly employed to describe the temporal occurrence of many aspects of a species biology (*e.g.* molting, migration, diapause in animals), but the phenology of reproduction (*e.g.* Sinclair *et al.* 2000, Rubenstein and Wikelski 2003, van den Hoff 2020) has attracted most interest. Reproductive phenology is an integral part of life history theory as it is at the heart of inter-generational trade-offs (*i.e.* between parents and offspring) and is a key factor of the reproductive success and fitness of the individuals (Stearns 1989, Forrest and Miller-Rushing 2010). On the one hand, the time of the year when most births occur is often linked to seasonal variations in food resources so that the flush of food resources matches the energetic needs of breeding, which ultimately improves the reproductive success of parents and the fitness of offspring (Plard *et al.* 2015). On the other hand, the spread of birth dates in a year is supposed to reflect anti-predator strategies to reduce the mortality associated with predation (Darling 1938, Gosling 1969), but also many other social and biological mechanisms (Ims 1990), such as avoidance of male harassment undergone by females (Boness *et al.* 1995) or intra-specific competition between offspring (Hodge *et al.* 2011).

In most ecological study cases, measurements and observations of phenology are

frequently performed at the population level by characterizing the temporal distribution of

biological events (Visser *et al.* 2010). Those rather complex and variable patterns are then

reduced to two main components: "timing", which is the date at which the event of interest

occurs, and "synchrony", which is the spread of the dates at which the event occurs, *i.e.* the

variability between individuals (Fig. 1). Besides, stimulated by research on the effects of

climate change on biodiversity (*e.g.* Crick and Sparks 1999, Parmesan 2007, Sarkar *et al.*

2019), the question of whether phenology is consistent or varies in time both at individual

and population levels has received increased interest in recent years (*e.g.* Renaud *et al.* 2019).

We therefore need to quantify underappreciated properties of phenology: the consistency of

the timing and synchrony (at the population scale) of the events from one reproductive season

to the next. As those characteristics of phenology are not described by specific words yet, we

suggest using "rhythmicity" and "regularity" to describe the consistency of timing and

synchrony respectively (Fig. 1), in line with Newstrom's terminology coined for tropical

plants (Newstrom *et al.* 1994).

Even if it is simple to grasp at first sight, the concept of phenology nevertheless

carries a lot of confusion in literature, both from a semantic and a descriptive point of view

(Visser *et al.* 2010). Past studies have explored phenology through the analysis of a huge

diversity of mathematical descriptors, many of which remaining specific to one single study.

This is problematic because well-defined, comparable and reliable descriptors of the temporal

distribution of biological events are key to achieving meaningful comparisons of phenology

patterns within or across species. For instance, English and colleagues reassessed the most

influential factors of reproductive synchrony in large herbivores from the existing literature,

but they had to narrow their original data set because there was no standardized way of

measuring and comparing synchrony across the studies (English *et al.* 2012). This large

diversity of metrics is further associated with a lack of generally accepted definitions of them

or even divergent definitions of the same word (see "seasonality" *sensu* Skinner *et al.* 2002

and Heideman and Utzurrum 2003), which really hampers our ability to make meaningful

comparisons (*e.g.* Ryan *et al.* 2007, Heldstab *et al.* 2018). Because experimental works are

logistically challenging or virtually impossible to conduct with large species, the comparison

of phenology patterns within a species living in contrasting environments, or across species

(Clauss *et al.* 2020) is of major importance to assess the role of explanatory factors

accounting for the often marked variability in phenology reported in empirical studies

(Rutberg 1987). Such comparative approaches (*sensu* Felsenstein 1985) shed light on the

ecological and evolutionary causes shaping the main stages of the life cycle of organisms

(Bronson 1989).

In spite of the diversity of approaches to describe phenology, we found only a few

attempts comparing phenology metrics to provide advice on which one should be used

preferentially according to the context of the study (Moussus *et al.* 2010, Landler *et al.* 2018).

Those initiatives are rare, and we currently lack a comprehensive comparison of metrics

previously used to characterize phenology and assess the extent to which the different metrics

actually capture the desired characteristics of the temporal distribution of events, nor do we

have a proper assessment of the sensitivity of those metrics to actual changes in phenology.

Here, we propose such a comparison of metrics based on a literature survey of reproductive

phenology in large herbivore species. We focus our overview on the taxonomic group of the

large herbivores because it has been studied in many species and at many locations (Rutberg

1987). As a consequence, we expect to find a wide variety of patterns of births, and a wide

diversity of metrics associated to describe them in literature. As hinted above, we first

clarified and formally defined four main terms describing phenology, built on our knowledge

of the literature: timing, synchrony, rhythmicity and regularity. We then conducted a

comparative analysis of 52 metrics that have been used to quantify the different

characteristics of phenology of births in large herbivores, highlighting their strengths and

weaknesses, and offering some guidance on which ones to use preferentially.

Materials and methods

We conducted a quantitative comparison of a wide range of metrics used to analyse

phenology in six steps. In Step 1, we recorded all metrics employed to measure phenology in

a selection of papers that we considered representative of the study of phenology of births in

large herbivores. In Step 2, we simulated contrasting phenology by varying independently the

four parameters that determine timing, synchrony, rhythmicity and regularity of phenology of

births (see details below). In Step 3, we calculated all metrics on the simulated phenology to

understand how they compare and what characteristics of phenology they actually measure.

In Step 4, we explored the similarities between metrics from a correlation matrix, and

identified categories of metrics capturing the same characteristic of phenology. In Step 5, we

evaluated the sensitivity of each metric to changes in the estimated parameter. In Step 6, we

ranked each metric based on eight criteria that we considered important to identify robust and

efficient metrics, but also meaningful from an ecological point of view (see Table 1 for a

description of each criterion).

Step 1: Retrieving and coding the different phenology metrics

We opportunistically searched the literature for articles focusing on the distribution of births

in large herbivores using keywords such as "phenology", "timing", "synchrony",

"seasonality", "period" or "season", and using various sources such as search engines and the

references in previously found articles. From these articles, published between 1966 and

2019, we recorded the metrics used to describe phenology of births at the population level.

We stopped our search once the rate at which we discovered new metrics with additional

papers became negligible.

We *a priori* classified each metric into one out of four categories, based on our

understanding of the original description and formula of the metric (Fig. 1): (1) *timing*

metrics, defining when within the year most births occur, (2) *synchrony* metrics, defining

whether females tend to give birth at the same time in a population in a given year, (3)

rhythmicity metrics, defining the consistency of timing between years, (4) *regularity* metrics,

defining the consistency of synchrony between years. In the literature, the term "seasonality"

can be used to describe the position of births in the year (*i.e.* timing, *e.g.* in Sinclair *et al.*

2000), the duration of birth period (*i.e.* synchrony, *e.g.* in Zerbe *et al.* 2012), and even the fact

that births occur at the same period of the year every year (*i.e.* rhythmicity and/or regularity,

e.g. in Heideman and Utzurrum 2003). However, this term is initially used to describe the

cyclical nature of the environment in a wider range than the study of birth phenology (Visser

et al. 2010). Thus, it should be used to describe organisms' phenology only when a direct

relationship between periodic environmental phenomena and the cycle of the organism at

stake has been demonstrated, which is not always the case in phenology studies. That is why

we suggest using the term "seasonality" only to describe the cyclicity of the environment and

to prefer neutral terms such as those we introduced in this paper to describe phenology of

births: rhythmicity and regularity.

Forty-seven articles (Supporting information 1) presented at least one mathematically-

defined phenology metric yielding overall 52 different metrics. In order to compare metrics

quantitatively, we had to slightly tweak some of them: when the metric was a boolean

(true/false) variable based on the significance of a statistical test ($n = 9$ metrics), we used the

value of the test statistics as output metric, thereby allowing us to investigate how the statistic

was influenced by the value of phenology parameters (see details in Supporting information

2). We could code all of them in R software (R Core Development Team 2019; except one in

Perl, www.perl.org).

Figure 1: Four characteristics of phenology of births can be explored to fully describe

phenology at the population scale: timing, synchrony, rhythmicity and regularity. Timing

describes when within the year most births occur, synchrony illustrates whether females tend

to give birth at the same time in a population in a given year, rhythmicity defines the

consistency of timing between years, regularity refers to the consistency of synchrony

between years. Green = timing, orange = synchrony, blue = rhythmicity, pink = regularity.

Step 2: Simulating phenology of births

We simulated phenology of births from statistical distributions with known parameters

(Supporting information 2) to assess what characteristic of phenology (timing, synchrony,

rhythmicity, regularity) each metric would capture, their sensitivity to changes into these four

key characteristics of interest, and the correlation between the 52 metrics. We simulated the

distributions of births over a year as most large herbivores breed once a year. This choice

does not limit the generality of our results: for species breeding more than once per year (*e.g.*

small species with short gestation length such as dikdik *Rynchotragus (Madoqua) kirki*,

Sinclair *et al.* 2000), the same metrics may be applied on sub-periods of time, each displaying

only one peak of births (*e.g.* Heideman and Utzurrum 2003 in bats).

Each simulated phenology was generated by randomly distributing births in time,

following a normal distribution. We distributed $n = 1000$ births within a year of 365 days,

repeated over 10 years (see why in "Material and Methods" section, Step 3). We changed

four parameters independently to modify the distribution of births: the mean day of birth for a

given year (*mean*), the standard deviation of the distribution of births for a given year (*sd*),

the range over which the mean birth date can vary across years (*Δmean*), and the range over

which the standard deviation can vary across years (*Δsd*). Each parameter varied in a range

from a minimum to a maximum value and was incremented with a constant step (Supporting

information 2). By choosing the value of these parameters, we could simulate changes in the

timing, synchrony, rhythmicity and regularity of the phenology of births respectively, and

independently. Because the simulated phenology of births relied on random draws, the actual

values of parameters in the simulated distribution of births could differ from the theoretical

values used in the simulation algorithm. We used the realized values of the distribution

parameters in the following analyses. Note that we replicated the same analyses using non-

normal distributions of births (*i.e.* skewed normal, bimodal, Cauchy, and random distribution)

to cover the variety of empirical distributions of births observed in *natura* and assess

robustness to non-normality (Supporting information 4). We performed all simulations using

the R software and made the code available on GitHub

(https://github.com/LucieThel/phenology-metrics).

Step 3: Computing the phenology metrics from simulated patterns of births

Among the 52 phenology metrics we analysed, most applied to a single year, but others

required two or more years of data to be computed (see the complete list in Supporting

information 3). Because we wanted to compute all 52 metrics, we chose to simulate annual

distributions of births over 10 consecutive years by default. For each simulation, we used data

from the first year to compute metrics requiring only one year of data ($n = 33$ metrics), data

from the first two years for metrics requiring two years of data ($n = 9$ metrics), and data from

the whole simulation for the other metrics ($n = 10$ metrics).

Step 4: Comparing the metrics

From the results of Step 3, we computed the global correlation matrix between all pairs of

metrics using Pearson correlations. We then identified groups of strongly correlated metrics

from the pairwise correlation coefficients and assigned each metric to one or several of the

four characteristics of phenology it was best related to. We compared this categorization with

our *a priori* classification of the metrics. This step enabled us to check our intuitive

classification of the metrics in addition to revealing whether some metrics could incidentally

capture several aspects of the distribution of births at once.

Step 5: Estimating the sensitivity of the metrics

For each metric, we performed a sensitivity analysis by quantifying the observed variation of

each metric with a fixed variation in the characteristic of phenology it was previously

associated with in Step 4. We did so by computing, for each possible pair of simulations

within the set of all simulations performed, the proportional difference between the realized

values of the phenology parameter of interest of the two simulations, and the proportional

difference between the values of the metric of interest of the same two simulations. In each

case the proportional difference was calculated as *[(Value$_{max}$ – Value$_{min}$) / Value$_{min}$] * 100*.

This formulation allowed us to work with positive values only, as we were interested in the

amplitude but not in the direction of the differences. We represented the scaled proportions of

variation by centering and scaling the proportions. We standardized the sensitivity of each

metric individually to prevent the representation of highly varying metrics from flattening the

representation of sparsely varying ones in the heat map representation.

Step 6: Scoring metrics

Finally, because there were too many different metrics, we could not discuss the pros and

cons for all of them. We chose instead to provide guidance about the usefulness of the

different metrics by scoring them according to a set of eight criteria that we considered as

important behaviour for a metric to be relevant (Table 1). We ranked the metrics from 0 (not

advised) to 8 (strongly advised) according to the number of criteria they fulfilled. Each

criterion could be evaluated independently from the others, except from the first four that

should all be met to consider that the metric describes the phenology characteristic *a minima*.

The proposed criteria (Table 1) consisted in verifying if 1) the metric varied according to the

phenology characteristic it was supposed to measure, 2) the variation of the metric according

to the phenology characteristic was monotonous, 3) the relationship with the characteristic of

phenology was strong (visual assessment of the association between the computed statistic

and the phenology characteristic), 4) the metric did not saturate within a biologically realistic

range of distributions of births. If those four criteria were satisfied, we then evaluated an

additional set of four criteria (see Table 1 for details). All criteria were scored from visual

inspection of the results by one of us (LT). Having systematic criteria helped us to minimize

the subjectivity of the scoring. We considered that metrics with scores ≤ 4 (for which the first

four essential criteria were not validated) should not be advised.

Table 1: Ordered list of the criteria used to evaluate the relevance of each metric describing

phenology of births. Each criterion can be individually fully (score of 1) or partially (score of

0.5) validated or no (score of 0) by each metric. The value for the first four criteria (in bold

type) should be > 0 to consider a metric to be possibly worthwhile and evaluate the

remaining criteria. The sum of the value obtained for each criterion gives the relevance index

of the metric (range between 0 and 8 points).

Criterion	Description	Score
goodness	Actually measures the parameter it is expected to measure	true = 1 false = 0
monotony	Varies monotonically with the value of the parameter it is expected to measure, *i.e.* the sign of the slope coefficient is constant	true = 1 false = 0
saturation	Does not saturate at the upper or lower boundary in a biological range of values (*e.g.* if a synchrony metric returned the same value when all births occurred during periods of various durations such as fifteen or thirty days, it was considered to saturate within a biologically realistic range of birth distributions because such distributions of births can be found in the wild)	true = 1 false = 0
strength[1]	Is characterized by a strong relationship with the parameter it is expected to measure, *i.e.* is the scatter plot not too dispersed around the general trend of the relationship between the metric and the phenology characteristic, as an empirical approach of the predictive power	high = 1 medium = 0.5 low = 0 strength of the association
normality	1- Does not assume normally distributed birth dates If false (assumes normality): 2- Is it robust to deviations to normality? *i.e.*, is the relationship between the metric and the parameter it is expected to measure conserved when births are not normally distributed	true = 1 false-true = 1 false-false = 0

origin	Does not depend on the temporal origin set by the investigator	true = 1 false = 0
linearity[2]	Is characterized by a linear relationship with the parameter it is expected to measure	type 1 = 1 type 2 and 3 = 0.5 type 4 = 0
unicity	Give a unique result	true = 1 false = 0

[1] high association: very small dispersion of points, medium association: small dispersion of

points that does not prevent from detecting a trend, low association: dispersion of points large

enough to prevent from detecting any trend, whatever the shape of the relationship (linear,

but also logistic or quadratic for instance). [2] type 1 is a linear relationship, type 2 is a

sigmoid-like relationship, type 3 is a quadratic-like relationship, type 4 is a binary

relationship.

Results

The mean number of metrics used in each paper was 3.8 ± 2.1 sd (*range* = 1-8). Eleven

metrics were *a priori* associated with timing, 25 with synchrony, 10 with rhythmicity and five

with regularity (we did not classify one metric because it could either be a rhythmicity or

regularity metric *a priori*). Those metrics were based on descriptive statistics, circular

statistics, statistical tests or statistical modelling such as general linear models. The metrics

were date, duration, count information (*e.g.* a number of births), binary classification (*i.e.* if a

given condition was satisfied or not), or unitless indices (Supporting information 3). We

identified 48 metrics linked to the parameter they were supposed to measure according to our

a priori classification. Among them, 8 metrics had a detectable relationship with at least one

of the three remaining phenology characteristics in addition to the relationship with the

phenology characteristic they were supposed to quantify (Supporting information 3 and 5).

The correlation matrix (Step 4) revealed obvious groups of metrics that were highly

correlated and thus reflected the same characteristic of phenology (Fig. 2). Five groups were

clearly identifiable, representing timing metrics (box 1), synchrony metrics (boxes 2 and 5),

rhythmicity metrics (box 3), and regularity metrics (box 4). The two groups of metrics

measuring synchrony (boxes 2 and 5) had highly but negatively correlated values (box 6).

This was indicative that all metrics of the two groups captured synchrony correctly, but in an

opposite way. Three metrics were singular and were associated with neither of the five

groups. One metric ("*splcomp*") seemed misclassified *a priori* as it correlated better with

synchrony than with regularity metrics. Another metric correlated well both with rhythmicity

and regularity metrics ("*diffbgper*").

Figure 2: Correlation matrix between all pairs of metrics, using Pearson correlations (n =

51, "rayleigh" removed because no variation observed). It was not possible to classify

"kolmomult" a priori in rhythmicity or regularity metrics, as it compares the complete

distribution of births between two years. Box 6 highlights the high but negative correlation

between the two groups of metrics measuring synchrony (boxes 2 and 5). Green = timing

metrics, orange = synchrony metrics, blue = rhythmicity metrics, pink = regularity metrics.

Note the high negative correlation between "compmean" and the other rhythmicity metrics,

highlighting that it is also a rhythmicity metric even if it is not located directly next to them.

       The sensitivity of the metrics to the simulated variation of the phenology

characteristics (Step 5) differed markedly between metrics, especially in synchrony and

regularity metrics (Fig. 3 and Supporting information 5). The scaled proportion of variation

of the metrics according to the proportion of variation of the associated parameters ranged

from -1.73 to 3.79 for timing metrics, from -3.81 to 2.01 for synchrony metrics, from -1.62 to

2.01 for rhythmicity metrics and from -1.60 to 1.91 for regularity metrics. The variation of

almost all timing, rhythmicity and regularity metrics according to variations of their

associated parameter was highly homogeneous. Synchrony metrics were less homogeneous,

certainly due to the fact that those metrics were the most numerous and based on more

diverse methods (proportion of variation, integrative indexes or moments of the distribution

of births, for instance). The metrics that were singular in the correlation matrix were clearly

visible in the heat maps, characterized by erratic or non-existent variations (*e.g.* "*skew*",

"*compmean*").

Figure 3: Heat maps representing the (scaled) proportion of variation of the metric in

relation to the proportion of variation of the parameter of phenology associated: a) timing

metrics according to the mean birth date for a given year (mean, n = 11), b) synchrony

metrics according to the standard deviation of the distribution of births for a given year (sd,

n = 25), c) rhythmicity metrics according to the range over which the mean birth date can

vary across years (Δmean, n = 10), d) regularity metrics according to the range over which

the standard deviation of the distribution of births can vary across years (Δsd, n = 6).

Colours in the heat maps reflect the amplitude of variation of each metric, scaled within the

metric. This scaling was done to prevent the large variation of some metrics to obscure the

smaller but meaningful variations of other metrics to be visible.

The same analyses conducted on the basis of non-normal distributions led to similar

observations in the case of asymmetric distributions (skewed normal, bimodal and Cauchy

distributions). The correlation matrices showed similar patterns of correlations between the

metrics, and the metrics varied analogously according to the variation of the *mean*, *sd*, *Δmean*

and *Δsd* of the distributions for normal and asymmetric distributions either (see Supporting

information 4 for a detailed analysis). Nevertheless, it is worth noting that a very limited

number of metrics depending on the skewness of the distribution did not perform as well with

the normal distribution than with asymmetric distributions. On the contrary, metrics

depending on the presence of a period without any birth did not perform as well with non-

normal distribution than with a normal distribution. In the case of a random distribution, no

clear correlations between metrics nor relationships between the metrics and the four

parameters of the distribution were detectable, except for some rare synchrony and timing

metrics (Supporting information 4).

The relevance score of the metrics (Step 6) varied between 0 and 8, covering the

complete range of variation possible (Fig. 4). All the timing metrics reached excellent scores

above 6. Nevertheless, the mean vector orientation ("*meanvo*") was the best metric, fulfilling

all our criteria with a score of 8 (Fig. 4). Three metrics provided a very good assessment of

the synchrony of births with a score of 7.5: the evenness index ("*pielou*"), the mean vector

length ("*meanvl*") and the comparison of the distribution of births to a uniform distribution

("*kolmouni*") (Fig. 4). The best metric to quantify rhythmicity measured the period length

between the median birth dates of two years ("*diffmed*"), with a score of 7 (Fig. 4). It is worth

noting that the non-parametric Mood test ("*mood*") provides a statistical assessment of

whether "*diffmed*" differs from 0. "*mood*" obtained a marginally lower score (6.5, Fig. 4) than

"*diffmed*" only because of a slight non-linearity in the relationship between simulation

parameter values and the metric's statistics. Altogether, we therefore considered that "*mood*"

could be very useful to measure rhythmicity. One metric quantifying regularity stood out

from the others according to our criteria: "*kolmomult*", which compares two birth

distributions (*score* = 7.5, Fig. 4). Our classification revealed what could be considered as

ineffective (*score* = 0, *n* = 4) and poor metrics (*score* ∈]0; 4], *n* = 14). Table 2 lists, for each

phenology characteristic, the metrics we identified as "best".

Figure 4: Score obtained by each phenology metric (n = 52) according to the eight criteria

used to assess the quality of the metrics (goodness, monotony, saturation, strength, normality,

origin, linearity and unicity, as defined in Table 1). Green = timing metrics, orange =

synchrony metrics, blue = rhythmicity metrics, pink = regularity metrics.

Table 2: List of the metrics considered as the best metric, for each characteristic of the

phenology of births (timing, synchrony, rhythmicity, regularity).

Phenology characteristic	Metric	Complete name	Description	Reference
Timing	*meanvo*	mean vector orientation	evaluate mean vector orientation of the birth distribution	Paré *et al.* 1996
Synchrony	*meanvl*	mean vector length	evaluate mean vector length of the birth distribution	Paré *et al.* 1996
Rhythmicity	*mood*	Mood test	compare median birth dates between two years	Berger and Cain 1999
Regularity	*kolmomult*	Kolmogorov Smirnov multi-year test	compare birth distribution between two years	Green and Rothstein 1993

Discussion

With more than fifty metrics used to describe and analyse the distribution of births in large

herbivores since 1966, our survey of the literature clearly illustrates the diversity of

approaches, even when focusing on a specific taxonomic group. Although the choice of a

metric is most of the time justified to answer a specific ecological question or on statistical

grounds, the lack of consensual methods to quantify phenology makes comparisons across

species or populations difficult at best, if possible at all. Our simulation study suggests that

such a diversity of metrics might be confusing and unnecessary because we could identify a

reduced set of simple metrics that work well to measure the different characteristics of

phenology. Moreover, we believe our work can also provide insights on how to analyse

phenology of other traits than births, such as migration dates of birds or flowering dates

(Moussus *et al.* 2010).

Many of the metrics we retrieved can be organised into four main categories, each one

capturing a particular characteristic of phenology: timing, synchrony, rhythmicity and

regularity. Of course, metrics belonging to the same category are not perfectly equivalent and

interchangeable (Fig. 2, see also a discussion comparing "*zerbe*" and "*rutberg*" metrics in

Zerbe *et al.* 2012). For instance, the correlations between timing metrics range between 0.68

and 1.00. The difference among metrics is more pronounced in the synchrony category with

correlations ranging from 0.05 to 1.00 (excluding "*kolmogau*" and "*skew*" metrics that appear

as singularities in the correlation matrix, Fig. 2). The way the different characteristics of

phenology are measured can lead to dependency between one another, and this could explain

the confusions noticed in literature between timing and synchrony through terms such as

"period" or "season" of births. Indeed, several metrics we tested vary not only according to

the phenology characteristic they were supposed to measure, but also according to other

characteristics of the phenology ($n = 8$ metrics). For instance, we show a strong correlation

between "*bgper*" and "*bgthper*" metrics, which evaluate the start of the birth period (*i.e.*

timing metrics), and the synchrony metrics. This association between different kinds of

metrics arises because the first births occur earlier when the standard deviation of the

distribution of births increases with all other parameters being equal (Fig. 2).

We attempted to identify what metrics could be the most suitable for measuring

timing, synchrony, regularity and rhythmicity by scoring them according to what we

subjectively considered as the main suitable properties. We considered for instance that a

good metric should not be restricted to one kind of pattern (*e.g.* unimodal), because the

distribution of births is not necessarily known a priori and may change between years due to

ecological factors (see Adams and Dale 1998 for instance). We actually found that only just

over 10 % of the metrics (based on the metrics for which this criterion was evaluated,

Supporting information 3) theoretically require normally distributed dates of birth to work

well. Additionally, all metrics assuming a normal distribution of births appeared robust to

deviations from normality. Thus, our criterion "normality" was actually not selective. The

metric should also not be sensitive to the temporal origin set by the investigator, as each

species (and even each population) has different favourable periods for its reproduction cycle

(*e.g.* mountain sheep from desert and alpine habitat, Bunnell 1982). Using the calendar year

would be biologically meaningless and will create artificial patterns of births by splitting the

distribution at the end of the year. Metrics that do not take into account the temporal origin of

the data prevent such limitations and allow an objective comparison. We identified only six

metrics ("*pielou*", "*mode*", "*meanvl*", "*meanvo*", "*kolmouni*" and "*kolmomult*") that meet this

assumption. Circular statistics could be prioritized to answer the difficulties linked to the

selection of temporal origin, as it is frequently done in primate literature (*e.g.* Di Bitetti and

Janson 2000). Notwithstanding such limitations, we could find several metrics that meet our

expectations of a good metric for each phenology characteristic (Table 2). Conversely, a few

metrics should not be recommended to describe phenology of births, such as the one

evaluating rhythmicity through the slope of the linear regression which describes the

evolution of the mean dates of births of several years ("*diffmean*"), or the one evaluating

synchrony through the duration of the period gathering at least a certain percent of births

("*nbtu*"). Those metrics do not capture the changes in the phenology parameter that they are

supposed to measure, or present undesirable statistical properties. The metric "*nbtu*" varied

non-monotonously with the level of synchrony of the birth phenology. The metric which

identifies the duration between first and last birth ("*per*"), a measure of synchrony, plateaued

for a range of biologically realistic values, limiting its use in a wide range of ecological

conditions (see Supporting information 3 for the score obtained by each metric for each

criterion, and Supporting information 4 and 5 for their detailed behaviour).

Overall, some phenology characteristics have been more consistently evaluated across

studies, a fact illustrated by the number of metrics of each category used in more than two

papers (Supporting information 6). If timing and synchrony of births are the easiest and most

frequent characteristics of phenology estimated and compared, only a handful of metrics

evaluates rhythmicity and regularity of the phenology of births across the years, and even

fewer are used more than twice (n = 2 and 0 metrics respectively). This gap might reflect the

difficulty in acquiring the adequate data. Indeed, robust analysis of rhythmicity and regularity

requires many years of data, which may not be available as phenology data collection is

costly and time-consuming (Kharouba and Wolkovich 2020). Moreover, scientists are less

interested in timing and synchrony consistency *per se* than in the relationship between timing

and synchrony, and ecological or environmental factors. Instead of measuring the consistency

of timing and synchrony *per se* across the years, they assess how the timing and synchrony

changes with temperature, rainfall or spring snow cover for instance (Paoli *et al.* 2018). In

any case, our study makes it clear that the currently available metrics are seldom and only

moderately correlated, particularly in the context of non-normal distributions of births

(Supporting information 4). Capturing the temporal variation of phenology across years

appears difficult and requires thoughtful selection and interpretation of the used metric. In the

context of climate change, there is an urgent need to develop statistical tools for measuring

regularity and rhythmicity, to test the hypothetical responses of phenology to changing

climate and predation regimes, for instance. Cosinor and Fourier analyses can help detecting

long-term trends and the phenology departure from such trends, but require a large amount of

data (Cancho-Candela *et al.* 2007).

We show that the assumption of a normal distribution or another bell-shaped

(asymmetric or not) distribution mimicking those found in *natura* (*e.g.* skewed normal,

bimodal or Cauchy distribution) has little consequences on our conclusions (Supporting

information 4). When there is no clear structure in the distribution of births, as often

encountered in some populations of large herbivores living in the southern hemisphere

(Sinclair *et al.* 2000), most metrics give inconsistent and unreliable results (Supporting

information 4). Describing random patterns using the metrics presented here is unlikely to be

useful, although evenness indexes such as "*pielou*" could at least provide a quantification of

the heterogeneity.

In conclusion, in the context of a comparative study with mainly non-random

distributions, we recommend using "*meanvo*" to describe timing and "*meanvl*" synchrony

respectively, because both are not influenced by temporal origin, a valued criterion. We

recommend using the underused "*mood*" (which statistically compares the median birth dates,

which are likely more robust measures than the date of the first birth or the mean birth date)

to describe rhythmicity and "*kolmomult*" to describe regularity (see Table 2 and Supporting

information 3 for a description of those metrics). Being non-parametric tests, they are

applicable in wide range of distributions as frequently observed in large herbivores

populations.

Declarations

Acknowledgments: This work was performed using the computing facilities of the CC

LBBE/PRABI and the IFB Cloud. The authors thank Simon Penel for his help with Perl

Programming Language, and Marcus Clauss for providing the code of one of the metrics. We

thank two anonymous reviewers and the subject editor form their comments on a previous

version of the manuscript. We also thank Marion Valeix for her comments on early drafts of

the manuscript, and Primaeɪlle Fusto for proofreading.

Funding: This work was supported by a grant from the "Ministère Francɪais de

l'Enseignement Supérieur, de la Recherche et de l'Innovation" through the "Ecole Doctorale

E2M2" of the "Université Claude Bernard Lyon 1".

Authors' contributions: LT, CB and SCJ conceived the ideas and designed the

methodology; LT performed data collection and analysed the data; LT wrote the first draft of

the manuscript, which was then edited by all authors. All authors gave final approval for

publication.

Data availability statement: Code available on GitHub:

https://github.com/LucieThel/phenology-metrics.

References

Adams, L. G. and Dale, B. W. 1998. Timing and synchrony of parturition in Alaskan caribou.

- J. Mammal. 79: 287–294.

Begon, M. *et al.* 1986. Ecology. Individuals, populations and communities. - Blackwell

scientific publications.

Boness, D. J. *et al.* 1995. Does male harassment of females contribute to reproductive

synchrony in the grey seal by affecting maternal performance? - Behav. Ecol. Sociobiol.

36: 1–10.

Bronson, F. H. 1989. Mammalian reproductive biology. - University of Chicago Press.

Bunnell, F. L. 1982. The lambing period of mountain sheep: synthesis, hypotheses, and tests.

- Can. J. Zool. 60: 1–14.

Cancho-Candela, R. *et al.* 2007. Decline and loss of birth seasonality in Spain: analysis of 33

421 731 births over 60 years. - J. Epidemiology Community Health 61: 713–718.

Clauss, M. *et al.* 2020. Basic considerations on seasonal breeding in mammals including their

testing by comparing natural habitats and zoos. - Mamm. Biol.: 1–14.

Crick, H. Q. P. and Sparks, T. H. 1999. Climate change related to egg-laying trends. - Nature

399: 423.

Darling, F. 1938. Bird flocks and the breeding cycle. - Cambridge University Press.

Demarée, G. R. 2011. From "Periodical Observations" to "Anthochronology" and

"Phenology"- the scientific debate between Adolphe Quetelet and Charles Morren on the

origin of the word "Phenology." - Int. J. Biometeorol. 55: 753–761.

Di Bitetti, M. S. and Janson, C. H. 2000. When will the stork arrive? Patterns of birth

seasonality in neotropical primates. - Am. J. Primatol. Off. J. Am. Soc. Primatol. 50: 109–

130.

English, A. K. *et al.* 2012. Reassessing the determinants of breeding synchrony in ungulates.

- PLoS One 7: e41444.

Felsenstein, J. 1985. Phylogenies and the comparative method. - Am. Nat. 125: 1–15.

Festa-Bianchet, M. 1988. Birthdate and survival in bighorn lambs (Ovis canadensis). - J.

Zool. 214: 653–661.

Forrest, J. and Miller-Rushing, A. J. 2010. Toward a synthetic understanding of the role of

phenology in ecology and evolution. - Philos. Trans. R. Soc. B Biol. Sci. 365: 3101–3112.

Gosling, L. M. 1969. Parturition and related behaviour in Coke's hartebeest, Alcelaphus

buselaphus cokei Günther. - J. Reprod. Fertil. Suppl. 6: 265–286.

Heideman, P. D. and Utzurrum, R. C. B. 2003. Seasonality and synchrony of reproduction in

three species of nectarivorous Philippines bats. - BMC Ecol. 3: 11.

Heldstab, S. A. *et al.* 2018. Geographical Origin, Delayed Implantation, and Induced

Ovulation Explain Reproductive Seasonality in the Carnivora. - J. Biol. Rhythms 33: 402–

419.

Hodge, S. *et al.* 2011. Reproductive competition and the evolution of extreme birth

synchrony in a cooperative mammal. - Biol. Lett. 7: 54-56.

Ims, R. A. 1990. The ecology and evolution of reproductive synchrony. - Trends Ecol. Evol.

5: 135–140.

Kharouba, H. M. and Wolkovich, E. M. 2020. Disconnects between ecological theory and

data in phenological mismatch research. - Nat. Clim. Chang.: 1–10.

Landler, L. *et al.* 2018. Circular data in biology: advice for effectively implementing

statistical procedures. - Behav. Ecol. Sociobiol. 72: 128.

Morren, C. 1849. Le globe, le temps et la vie. - Bulletins de l'Académie royale des Sciences,

des Lettres et des Beaux-Arts de Belgique 16: 660–684.

Moussus, J.-P. *et al.* 2010. Featuring 10 phenological estimators using simulated data. -

Methods Ecol. Evol. 1: 140–150.

Newstrom, L. E. *et al.* 1994. A new classification for plant phenology based on flowering

patterns in lowland tropical rain forest trees at La Selva, Costa Rica. - Biotropica: 141–

159.

Paoli, A. *et al.* 2018. Winter and spring climatic conditions influence timing and synchrony

of calving in reindeer. - PLoS One 13: e0195603.

Parmesan, C. 2007. Influences of species, latitudes and methodologies on estimates of

phenological response to global warming. - Glob. Chang. Biol. 13: 1860–1872.

Plard, F. *et al.* 2015. The influence of birth date via body mass on individual fitness in a long-

lived mammal. - Ecology 96: 1516–1528.

Renaud, L.-A. *et al.* 2019. Phenotypic plasticity in bighorn sheep reproductive phenology:

from individual to population. - Behav. Ecol. Sociobiol. 73: 50.

Rubenstein, D. R. and Wikelski, M. 2003. Seasonal changes in food quality: a proximate cue

for reproductive timing in marine iguanas. - Ecology 84: 3013–3023.

Rutberg, A. T. 1987. Adaptive hypotheses of birth synchrony in ruminants: an interspecific

test. - Am. Nat. 130: 692–710.

Ryan, S. J. *et al.* 2007. Ecological cues, gestation length, and birth timing in African buffalo

(Syncerus caffer). - Behav. Ecol. 18: 635–644.

Sarkar, U. K. *et al.* 2019. Climato-environmental influence on breeding phenology of native

catfishes in River Ganga and modeling species response to climatic variability for their

conservation. - Int. J. Biometeorol. 63: 991–1004.

Sinclair, A. R. E. *et al.* 2000. What determines phenology and synchrony of ungulate

breeding in Serengeti? - Ecology 81: 2100–2111.

Skinner, J. D. *et al.* 2002. Inherent seasonality in the breeding seasons of African mammals:

evidence from captive breeding. - Trans. R. Soc. South Africa 57: 25–34.

Stearns, S. C. 1989. Trade-offs in life-history evolution. - Funct. Ecol. 3: 259–268.

van den Hoff, J. 2020. Environmental constraints on the breeding phenology of Giant Petrels

Macronectes spp., with emphasis on Southern Giant Petrels M. giganteus. - Mar. Ornithol.

48: 33–40.

Visser, M. E. *et al.* 2010. Phenology, seasonal timing and circannual rhythms: towards a

unified framework. - Philos. Trans. R. Soc. B Biol. Sci. 365: 3113–3127.

Zerbe, P. *et al.* 2012. Reproductive seasonality in captive wild ruminants: implications for

biogeographical adaptation, photoperiodic control, and life history. - Biol. Rev. 87: 965–

990.