How to describe and measure phenology? An investigation on the diversity of metrics using phenology of births in large herbivores

Lucie Thel[1], Simon Chamaillé-Jammes[2, 3, 4] & Christophe Bonenfant[1, 4]

Authors details

[1] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Bâtiment 711, Université Lyon I, 43 Boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France.

[2] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier, France.

[3] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

[4] LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe.

Correspondence author: Lucie Thel, lucie.mc.thel@gmail.com.

Abstract

Proposed in 1849 by Charles Morren to depict periodical phenomena governed by seasons, the term "phenology" has spread in many fields of biology. With the wide adoption of the concept of phenology flourished a large number of metrics with different meaning and interpretation. Here, we first *a priori* classified 52 previously published metrics used to characterize the phenology of births in large herbivores according to four biological characteristics of interest: timing, synchrony, rhythmicity and regularity of births. We then applied each metric retrieved on simulation data, considering normal and non-normal distributions of births and varying distributions of births in time. We then evaluated the ability of each metric to capture the variation of the four phenology characteristics *via* a sensitivity analysis. Finally, we scored each metric according to eight criteria we considered important to describe phenology correctly. The high correlation we found among the many metrics we retrieved suggests that such diversity of metrics is unnecessary. We further show that the best metrics are not the most commonly used, and that simpler is often better. Circular statistics with the mean vector orientation and mean vector length seems, respectively, particularly suitable to describe the timing and synchrony of births in a wide range of phenology patterns. Tests designed to compare statistical distribution, like Mood and Kolmogorov-Smirnov tests, allow a first and easy quantification of rhythmicity and regularity of birth phenology respectively. By identifying the most relevant metrics our study should facilitate comparative studies of phenology of birth or of any other life-history event. For instance, comparative studies of the phenology of mating or migration dates are particularly important in the context of climate change.

Keywords: regularity, rhythmicity, seasonality, synchrony, timing, ungulate

Introduction

In 1849, Charles Morren coined the term "phenology" to describe how periodical phenomena such as plant growth and reproduction are governed by the course of seasons (Morren 1849, see also Demarée 2011). With his observations he opened a new field of research and almost two centuries later the concept of phenology has become a cornerstone of ecology (Begon *et al.* 1986), used in plant and animal ecology simultaneously (Forrest and Miller-Rushing 2010). By describing when particular life-history events (*e.g.* flowering, parturition) occur in relation to the characteristics or states of the individual (*e.g.* size, age) as well as to environmental factors (*e.g.* photoperiod, predation risk) the concept of phenology is key to understanding the temporal cycles in the life history of species (Forrest and Miller-Rushing 2010). Nowadays, the term phenology is commonly employed to describe the temporal occurrence of many aspects of a species biology (*e.g.* moulting, migration, diapause in animals), but the phenology of reproduction (*e.g.* Sinclair *et al.* 2000, Rubenstein and Wikelski 2003, van den Hoff 2020) has attracted most interest. Reproductive phenology is an integral part of life history theory as it is at the heart of inter-generational trade-offs (*i.e.* between parents and offspring) and is a key factor of the reproductive success and fitness of the individuals (Stearns 1989, Forrest and Miller-Rushing 2010). On the one hand, the time of the year when most births occur is often linked to seasonal variations in food resources so that the flush of food resources matches the energetic needs of breeding, which ultimately improves the reproductive success of parents and the fitness of offspring (Plard *et al.* 2015). While on the other hand, the spread of birth dates in a year is supposed to reflect anti-predator strategies to reduce the mortality associated with predation (Darling 1938, Gosling 1969), but also many other social and biological mechanisms (Ims 1990), such as avoidance of male harassment undergone by females (Boness *et al.* 1995) or intra-specific competition between offspring (Hodge *et al.* 2011).

In most ecological studies, measurements and observations of phenology are

frequently performed at the population level by characterizing the temporal distribution of

biological events (Visser *et al.* 2010). These rather complex and variable patterns are reduced

to two main components: "timing", the date at which the event of interest occurs, and

"synchrony", the spread of the dates at which the event occurs, *i.e.* the variability between

individuals (Fig. 1). Stimulated by research on the effects of climate change on biodiversity

(*e.g.* Crick and Sparks 1999, Parmesan 2007, Sarkar *et al.* 2019), the question of whether

phenology is consistent or varies in time, both at individual and population levels, has

received increased interest in recent years (*e.g.* Renaud *et al.* 2019). We therefore need to

quantify the underappreciated properties of phenology: the consistency of the timing and

synchrony (at the population scale) of the events from one reproductive season to the next. As

these characteristics of phenology are not described by specific words yet, we suggest using

"rhythmicity" and "regularity" to describe the consistency of timing and synchrony

respectively (Fig. 1), in line with Newstrom's terminology coined for tropical plants

(Newstrom *et al.* 1994).

Despite appearing simple, the concept of phenology carries a lot of confusion in

literature, both from a semantic and a descriptive point of view (Visser *et al.* 2010). Previous

studies have explored phenology using a vast diversity of mathematical descriptors, many of

which remain specific to a single study. This is problematic as well-defined, comparable and

reliable descriptors of the temporal distribution of biological events are key to achieving

meaningful comparisons of phenology patterns within or across species. English and

colleagues reassessed the most influential factors of reproductive synchrony in large

herbivores using the existing literature, but had to narrow their original data set because there

was no standardized way of measuring and comparing synchrony across the studies (English

et al. 2012). This large diversity of metrics is associated with a lack of widely accepted

definitions or divergent definitions for the same word (see "seasonality" *sensu* Skinner *et al.*

2002 and Heideman and Utzurrum 2003), which further limits our ability to make meaningful

comparisons (*e.g.* Ryan *et al.* 2007, Heldstab *et al.* 2018). As experimental studies are

logistically challenging or virtually impossible to conduct with large species, the comparison

of phenology patterns within a species living in contrasting environments or across species

(Clauss *et al.* 2020) is of major importance to assess the role of explanatory factors

accounting for the often marked variability in phenology reported in empirical studies

(Rutberg 1987). Such comparative approaches (*sensu* Felsenstein 1985) indeed shed light on

the ecological and evolutionary causes shaping the main stages of the life cycle of organisms

(Bronson 1989).

Despite the increasing diversity of approaches to describe phenology, we found only a

few attempts to compare phenology metrics and to provide advice on which one should be

used preferentially according to the context of the study (Moussus *et al.* 2010, Landler *et al.*

2018). These initiatives are rare and we currently lack a comprehensive comparison of the

metrics previously used to characterize phenology. The extent to which the different metrics

capture the desired characteristics of the temporal distribution of events, or the sensitivity of

those metrics to actual changes in phenology remains to be adequately assessed. Here, we

propose such a comparison of metrics based on a literature survey of reproductive phenology

in large herbivore species. We focus on the taxonomic group of the large herbivores as it has

been studied in a number of species and at different locations (Rutberg 1987). As a result, we

expect to find a wide variety of patterns of births and a wide diversity of metrics to describe

them. We first clarify and formally define the four main terms describing phenology: timing,

synchrony, rhythmicity and regularity, using our knowledge from the existing literature. We

then conduct a comparative analysis of 52 metrics that have been used to quantify the

different characteristics of phenology of births in large herbivores, highlighting their

strengths and weaknesses. To conclude, we recommend one metric for each of the four main

characteristics of phenology.

Materials and methods

We conducted a quantitative comparison of a wide range of metrics used to analyse

phenology in six steps. In Step 1, we recorded all metrics employed to measure phenology in

a selection of papers that we considered representative of the study of phenology of births in

large herbivores. In Step 2, we simulated contrasting phenology by varying independently the

four parameters that determine timing, synchrony, rhythmicity and regularity of phenology of

births (see details below). In Step 3, we calculated all metrics on the simulated phenology to

understand how they compare and what characteristics of phenology they actually measure.

In Step 4, we explored the similarities between metrics from a correlation matrix, and

identified categories of metrics capturing the same characteristic of phenology. In Step 5, we

evaluated the sensitivity of each metric to changes in the estimated parameter. In Step 6, we

ranked each metric based on eight criteria that we considered important to identify robust and

efficient metrics, but also meaningful from an ecological point of view (see Table 1 for a

description of each criterion).

Step 1: Retrieving and coding the different phenology metrics

We opportunistically searched the literature for articles focusing on the distribution of births

in large herbivores using keywords such as "phenology", "timing", "synchrony",

"seasonality", "period" or "season", and using various sources such as search engines and the

references in previously found articles. From these articles, published between 1966 and

2019, we recorded the metrics used to describe phenology of births at the population level.

We stopped our search once the rate at which we discovered new metrics with additional

papers became negligible.

We *a priori* classified each metric into one out of four categories based on our

understanding of the original description and formula of the metric (Fig. 1): (1) *timing*

metrics, defining when within the year most births occur, (2) *synchrony* metrics, defining

whether females tend to give birth at the same time in a population in a given year, (3)

rhythmicity metrics, defining the consistency of timing between years, (4) *regularity* metrics,

defining the consistency of synchrony between years. In the literature, the term "seasonality"

can be used to describe the location of births in the year (*i.e.* timing, *e.g.* in Sinclair *et al.*

2000), the duration of birth period (*i.e.* synchrony, *e.g.* in Zerbe *et al.* 2012), and even the fact

that births occur at the same period of the year every year (*i.e.* rhythmicity and/or regularity,

e.g. in Heideman and Utzurrum 2003). However, this term is initially used to describe the

cyclical nature of the environment in a wider range than the study of birth phenology (Visser

et al. 2010). Thus, it should be used to describe organisms' phenology only when a direct

relationship between periodic environmental phenomena and the cycle of the organism at

stake has been demonstrated, which is not always the case in phenology studies. For this

reason, we suggest using the term "seasonality" only to describe the cyclicity of the

environment and prefer the use of neutral terms such as those we introduced in this paper to

describe phenology of births: rhythmicity and regularity.

Forty-seven articles (Supporting information 1) presented at least one mathematically-

defined phenology metric yielding 52 different metrics. In order to compare metrics

quantitatively, we slightly tweaked some of them: when the metric was a boolean (true/false)

variable based on the significance of a statistical test ($n = 9$ metrics), we used the value of the

test statistics as output metric, thereby allowing us to investigate how the statistic was

influenced by the value of phenology parameters (see details in Supporting information 2).

All metrics could be coded in R software (R Core Development Team 2019) except one, for

which Perl was used (www.perl.org).

Figure 1: Four characteristics of phenology of births can be explored to fully describe phenology at

the population scale: timing, synchrony, rhythmicity and regularity. Timing describes when within the

year most births occur, synchrony illustrates whether females tend to give birth at the same time in a

population in a given year, rhythmicity defines the consistency of timing between years, regularity

refers to the consistency of synchrony between years. Green = timing, orange = synchrony, blue =

rhythmicity, pink = regularity.

Step 2: Simulating phenology of births

We simulated phenology of births from statistical distributions with known parameters

(Supporting information 2) to assess what characteristic of phenology (timing, synchrony,

rhythmicity, regularity) each metric would capture, their sensitivity to changes into these four

key characteristics of interest, and the correlation between the 52 metrics. We simulated the

distributions of births over a year as most large herbivores breed once a year. This choice

does not limit the generality of our results: for species breeding more than once per year (*e.g.*

small species with short gestation length such as dikdik *Rynchotragus (Madoqua) kirki*,

Sinclair *et al.* 2000), the same metrics may be applied on sub-periods of time, each displaying

only one birth peak (*e.g.* Heideman and Utzurrum 2003 in bats).

Each simulated phenology was generated by randomly distributing births in time,

following a normal distribution. We distributed $n = 1000$ births within a year of 365 days,

repeated over 10 years (see why in "Material and Methods" section, Step 3). We changed

four parameters independently to modify the distribution of births: the mean day of birth for a

given year (*mean*), the standard deviation of the distribution of births for a given year (*sd*),

the range over which the mean birth date can vary across years (*Δmean*), and the range over

which the standard deviation can vary across years (*Δsd*). Each parameter varied in a range

from a minimum to a maximum value and was incremented with a constant step (Supporting

information 2). Choosing the value of these parameters allowed us to simulate changes in the

timing, synchrony, rhythmicity and regularity of the phenology of births independently. As

the simulated phenology of births relied on random draws, the actual values of parameters in

the simulated distribution of births could differ from the theoretical values used in the

simulation algorithm. We used the realized values of the distribution parameters in the

following analyses. Note that we replicated the same analyses using non-normal distributions

of births (*i.e.* skewed normal, bimodal, Cauchy, and random distribution) to cover the variety

of empirical distributions of births observed in *natura* and assess robustness to non-normality

(Supporting information 4). We performed all simulations using the R software and made the

code available on GitHub (https://github.com/LucieThel/phenology-metrics).

Step 3: Computing the phenology metrics from simulated patterns of births

Among the 52 phenology metrics we analysed, most applied to a single year, but others

required two or more years of data to be computed (see the complete list in Supporting

information 3). As we aimed to compute all 52 metrics, we chose to simulate annual

distributions of births over 10 consecutive years by default. For each simulation, we used data

from the first year to compute metrics requiring only one year of data ($n = 33$ metrics), data

from the first two years for metrics requiring two years of data ($n = 9$ metrics), and data from

the whole simulation for the other metrics ($n = 10$ metrics).

Step 4: Comparing the metrics

With the results from Step 3, we computed the global correlation matrix between all pairs of

metrics using Pearson correlations. We then identified groups of strongly correlated metrics

from the pairwise correlation coefficients and assigned each metric to one or several of the

four characteristics of phenology it was best related to. We compared this categorization with

our *a priori* classification of the metrics. This step enabled us to check our intuitive

classification of the metrics in addition to revealing whether some metrics could incidentally

capture several aspects of the distribution of births at once.

Step 5: Estimating the sensitivity of the metrics

For each metric, we performed a sensitivity analysis by quantifying the observed variation of

each metric with a fixed variation in the characteristic of phenology it was previously

associated with in Step 4. We did this by computing for each possible pair of simulations

within the set of all simulations performed, the proportional difference between the realized

values of the phenology parameter of interest of the two simulations, and the proportional

difference between the values of the metric of interest of the same two simulations. In each

case the proportional difference was calculated as *[(Value$_{max}$ – Value$_{min}$) / Value$_{min}$] * 100*.

This formulation allowed us to work with positive values only as we were interested in the

amplitude but not in the direction of the differences.

Step 6: Scoring metrics

Finally, as there were too many different metrics, we were unable to discuss the pros and

cons for each of them. We chose instead to provide guidance about the usefulness of the

different metrics by scoring them according to a set of eight criteria that we considered as

important behaviour for a metric to be relevant (Table 1). Having systematic criteria helped

us to minimize the subjectivity of the scoring so we ranked the metrics from 0 (not advised)

to 8 (strongly advised) according to the number of criteria they fulfilled. The proposed

criteria (Table 1) consisted in verifying if 1) the metric varied according to the phenology

characteristic it was supposed to measure, 2) the variation of the metric according to the

phenology characteristic was monotonous, 3) the relationship with the characteristic of

phenology was strong (visual assessment of the association between the computed statistic

and the phenology characteristic), 4) the metric did not saturate within a biologically realistic

range of distributions of births. We considered that metrics with scores ≤ 4 (for which the

first four essential criteria were not validated) should not be advised. If those four criteria

were satisfied, we evaluated an additional set of four criteria (normality, independence of the

temporal origin, linearity and unicity of the output, see Table 1 for a detailed description). All

criteria were scored from visual inspection of the results by one of us (LT).

Table 1: Ordered list of the criteria used to evaluate the relevance of each metric describing

phenology of births. Each criterion can be individually fully (score of 1) or partially (score of 0.5)

validated or no (score of 0) by each metric. The value for the first four criteria (in bold type) should

be > 0 to consider a metric to be possibly worthwhile and evaluate the remaining criteria. The sum of

the value obtained for each criterion gives the relevance index of the metric (range between 0 and 8

points).

Criterion	Description	Score
goodness	Measures the parameter it is expected to measure	true = 1 false = 0
monotony	Varies monotonically with the value of the parameter it is expected to measure, *i.e.* the sign of the slope coefficient is constant	true = 1 false = 0
saturation	Does not saturate at the upper or lower boundary in a biological range of values (*e.g.* if a synchrony metric returned the same value when all births occurred during periods of various durations such as fifteen or thirty days, it was considered to saturate within a biologically realistic range of birth distributions because such distributions of births can be found in the wild)	true = 1 false = 0
strength[1]	Is characterized by a strong relationship with the parameter it is expected to measure, *i.e.* is the scatter plot not too dispersed around the general trend of the relationship between the metric and the phenology characteristic, as an empirical approach of the predictive power	high = 1 medium = 0.5 low = 0 strength of the association
normality	1- Does not assume normally distributed birth dates If false (assumes normality): 2- Is it robust to deviations to normality? *i.e.*, is the relationship between the metric and the parameter it is expected to measure conserved when births are not normally distributed	true = 1 false-true = 1 false-false = 0
origin	Does not depend on the temporal origin set by the investigator	true = 1 false = 0
linearity[2]	Is characterized by a linear relationship with the parameter it is expected to measure	type 1 = 1 type 2 and 3 = 0.5 type 4 = 0
unicity	Give a unique result	true = 1 false = 0

258   [1] *high association: very small dispersion of points, medium association: small dispersion of points*

259   *that does not prevent from detecting a trend, low association: dispersion of points large enough to*

260   *prevent from detecting any trend, whatever the shape of the relationship (linear, but also logistic or*

261   *quadratic for instance).* [2] *type 1 is a linear relationship, type 2 is a sigmoid-like relationship, type 3 is*

262   *a quadratic-like relationship, type 4 is a binary relationship.*

Results

The mean number of metrics used in each paper was 3.8 ± 2.1 sd (*range* = 1 - 8). Eleven

metrics were *a priori* associated with timing, 25 with synchrony, 10 with rhythmicity and five

with regularity. We did not classify one metric because it could either be a rhythmicity or

regularity metric *a priori*. Those metrics were based on descriptive statistics, circular

statistics, statistical tests or statistical modelling such as general linear models. The unit of the

metrics were date, duration, counts (*e.g.* a number of births), binary classification (*i.e.* if a

given condition was satisfied or not), or unitless indices (Supporting information 3).

The correlation matrix (Step 4) revealed groups of metrics that were highly correlated

and thus reflected the same characteristic of phenology (Fig. 2). Five groups were clearly

identifiable, representing timing metrics (Fig. 2 - box 1), synchrony metrics (Fig. 2 - boxes 2

and 5), rhythmicity metrics (Fig. 2 - box 3), and regularity metrics (Fig. 2 - box 4). The two

groups of metrics measuring synchrony had highly but negatively correlated values (Fig. 2 -

box 6). This indicated that all metrics of the two groups captured synchrony correctly,

however, in an opposing way. Three metrics were singular and were associated with neither

of the five groups. The metric which compares the slope coefficients of linear models

describing the log percent of cumulative births ("*splcomp*") should measure regularity, but it

rather correlated better with synchrony metrics. The metric which evaluates the duration

between the first birth dates of two reproductive cycles ("*diffbgper*"), an assessment of

rhythmicity, correlated well with both rhythmicity and regularity metrics. Seven other metrics

had a detectable relationship with at least one of the three remaining phenology

characteristics in addition to the relationship with the phenology characteristic they were

supposed to quantify (Supporting information 3 and 5).

Figure 2: Correlation matrix between all pairs of metrics, using Pearson correlations (n = 51,

"rayleigh" removed because no variation observed). It was not possible to classify "kolmomult" a

priori in rhythmicity or regularity metrics, as it compares the complete distribution of births between

two years. Box 6 highlights the high but negative correlation between the two groups of metrics

measuring synchrony (boxes 2 and 5). Green = timing metrics, orange = synchrony metrics, blue =

rhythmicity metrics, pink = regularity metrics. Note the high negative correlation between

"compmean" and the other rhythmicity metrics, highlighting that it is also a rhythmicity metric even

if it is not located directly next to them.

       The sensitivity of the metrics to the simulated variation of the phenology

characteristics (Step 5) differed markedly between metrics, especially in synchrony and

regularity metrics (Fig. 3 and Supporting information 5). The scaled proportion of variation

of the metrics according to the proportion of variation of the associated parameters ranged

from -1.73 to 3.79 for timing metrics, from -3.81 to 2.01 for synchrony metrics, from -1.62 to

2.01 for rhythmicity metrics and from -1.60 to 1.91 for regularity metrics. The variation of

almost all timing, rhythmicity and regularity metrics according to variations of their

associated parameter was highly homogeneous. Synchrony metrics were less homogeneous,

certainly due to the fact that those metrics were the most numerous and based on more

diverse methods (proportion of variation, integrative indexes or moments of the distribution

of births, for instance). The metrics that were singular in the correlation matrix were clearly

visible in the heat maps, characterized by erratic or non-existent variations (*e.g.* skewness of

the birth distribution "*skew*", and the comparison of mean date of births "*compmean*").

Figure 3: Heat maps representing the (scaled) proportion of variation of the metric in relation to the

proportion of variation of the parameter of phenology associated (sensitivity analysis): a) timing

metrics according to the mean birth date for a given year (mean, n = 11), b) synchrony metrics

according to the standard deviation of the distribution of births for a given year (sd, n = 25), c)

rhythmicity metrics according to the range over which the mean birth date can vary across years

(Δmean, n = 10), d) regularity metrics according to the range over which the standard deviation of

the distribution of births can vary across years (Δsd, n = 6).

The same analyses conducted on the basis of non-normal distributions led to similar

observations in the case of asymmetric distributions (skewed normal, bimodal and Cauchy

distributions). The correlation matrices showed similar patterns of correlations between the

metrics, and the metrics varied analogously according to the variation of the *mean*, *sd*, *Δmean*

and *Δsd* of the distributions for normal and asymmetric distributions either (see Supporting

information 4 for a detailed analysis). Nevertheless, it is worth noting that a very limited

number of metrics depending on the skewness of the distribution did not perform as well with

the normal distribution than with asymmetric distributions. On the contrary, metrics

depending on the presence of a period without any birth did not perform as well with non-

normal distribution than with a normal distribution. In the case of a random distribution, no

clear correlations between metrics nor relationships between the metrics and the four

parameters of the distribution were detectable, except for some rare synchrony and timing

metrics (Supporting information 4).

The relevance score of the metrics (Step 6) varied between 0 and 8, covering the

complete range of variation possible (Fig. 4) and we list, for each phenology characteristic,

the metrics we identified as "best" (Table 2). Our classification also revealed what could be

considered as ineffective (*score* = 0, *n* = 4) and poor metrics (*score* ∈]0; 4], *n* = 14). All the

timing metrics reached excellent scores above 6. Nevertheless, the mean vector orientation

("*meanvo*") was the best metric, fulfilling all our criteria with a score of 8 (Fig. 4). Three

metrics provided a very good assessment of the synchrony of births with a score of 7.5: the

evenness index ("*pielou*"), the mean vector length ("*meanvl*") and the comparison of the

distribution of births to a uniform distribution ("*kolmouni*") (Fig. 4). The best metric to

quantify rhythmicity measured the period length between the median birth dates of two years

("*diffmed*"), with a score of 7 (Fig. 4). It is worth noting that the non-parametric Mood test

("*mood*") provides a statistical assessment of whether "*diffmed*" differs from 0. The non-

parametric Mood test ("*mood*") obtained a marginally lower score (6.5, Fig. 4) than "*diffmed*"

only because of a slight non-linearity in the relationship between simulation parameter values

and the metric's statistics. Altogether, we therefore considered that "*mood*" could be very

useful to measure rhythmicity. One metric quantifying regularity stood out from the others

according to our criteria: the non-parametric Kolmogorov-Smirnov test ("*kolmomult*"), which

compares two birth distributions (*score* = 7.5, Fig. 4).

Figure 4: Score obtained by each phenology metric (n = 52) according to the eight criteria used to

assess the quality of the metrics (goodness, monotony, saturation, strength, normality, origin,

linearity and unicity, as defined in Table 1). Green = timing metrics, orange = synchrony metrics,

blue = rhythmicity metrics, pink = regularity metrics.

Table 2: List of the metrics considered as the best metric, for each characteristic of the phenology of births (timing, synchrony, rhythmicity, regularity).

Phenology characteristic	Metric	Complete name	Description	Reference
Timing	*meanvo*	mean vector orientation	evaluates mean vector orientation of the birth distribution	Paré *et al.* 1996
Synchrony	*meanvl*	mean vector length	evaluates mean vector length of the birth distribution	Paré *et al.* 1996
Rhythmicity	*mood*	Mood test	compares median birth dates between two years	Berger and Cain 1999
Regularity	*kolmomult*	Kolmogorov Smirnov multi-year test	compares birth distributions between two years	Green and Rothstein 1993

Discussion

With more than fifty metrics used to describe and analyse the distribution of births in large herbivores since 1966, our survey of the literature clearly illustrates the diversity of approaches, even when focusing on a specific taxonomic group. Although the choice of a metric is most of the time justified, either to answer a specific ecological question or on statistical grounds, the lack of consensual methods to quantify phenology makes comparisons across species or populations difficult at best, if possible at all. Our simulation study suggests that such a diversity of metrics may cause confusion and be unnecessary as we were able to identify a reduced set of simple metrics that works well to measure the different characteristics of phenology. Moreover, we believe our work can also provide insights into how to analyse phenology of other traits than births, such as migration dates of birds or flowering dates (Moussus *et al.* 2010).

Many of the metrics we retrieved can be organised into four main categories, each one capturing a particular characteristic of phenology: timing, synchrony, rhythmicity and regularity. Of course, metrics belonging to the same category are not perfectly equivalent and interchangeable (Fig. 2, see also a discussion comparing "*zerbe*" and "*rutberg*" metrics in

Zerbe *et al.* 2012). For instance, the correlations between timing metrics range between 0.68

and 1.00. The difference among metrics is more pronounced in the synchrony category with

correlations ranging from 0.05 to 1.00 (excluding "*kolmogau*" and "*skew*" metrics that appear

as singularities in the correlation matrix, Fig. 2). How different characteristics of phenology

are measured can lead to dependency between one another and this could explain the

confusions found in the literature between timing and synchrony through terms such as

"period" or "season" of births. Indeed, several of the metrics we tested vary not only

according to the phenology characteristic they were used to measure, but also according to

other characteristics of the phenology ($n = 8$ metrics). For instance, we show a strong

correlation between metrics that evaluate the start of the birth period (*i.e.* timing metrics

"*bgper*" and "*bgthper*") and the synchrony metrics in general. This association between

different types of metrics arises when the standard deviation of the simulated distributions of

births increases (while the mean is fixed), leading to earlier births (Fig. 2).

We attempted to identify what metrics could be the most suitable for measuring

timing, synchrony, regularity and rhythmicity of phenology by scoring them according to

what we subjectively considered as the main suitable properties. We considered that a good

metric should not be restricted to one kind of pattern (*e.g.* unimodal) as the distribution of

births is not necessarily known a priori and may change between years due to ecological

factors (see Adams and Dale 1998 for instance). Slightly more than 10 % of the metrics

theoretically require normally distributed dates of birth to work well (based on the metrics for

which this criterion was evaluated, Supporting information 3). We showed these metrics are

generally robust to deviations from normality so this assumption does not limit their

application to most data. The metrics should also be independent of the temporal origin set by

the investigator, as the favourable periods for reproduction cycle differ between species and

populations (*e.g.* mountain sheep *Ovis spp.* inhabiting desert and alpine ecosystems, Bunnell

1982). Using the calendar year would be biologically meaningless and will create artificial

patterns of births by splitting the distribution around the end of the year. We identified six

metrics independent of the temporal origin: the day with the highest number of births

("*mode*"), the evenness index ("*pielou*"), the mean vector orientation and length form the

circular statistics ("*meanvl*" and "*meanvo*" respectively), and the non-parametric

Kolmogorov-Smornov test comparing a birth distribution to a uniform distribution or another

birth distribution ("*kolmouni*" and "*kolmomult*" respectively). Circular statistics could be

favoured to answer the difficulties linked to the selection of temporal origin, as it is

frequently done in primate literature (*e.g.* Di Bitetti and Janson 2000). Notwithstanding such

limitations, we found several metrics that met our expectations of a good metric for each

phenology characteristic (Table 2 and Figure 4).

On the other side a few metrics should not be recommended to describe phenology of

births. The evaluation of rhythmicity describing the evolution of the mean dates of births of

several years with a linear regression ("*diffmean*"), or the quantification of synchrony through

the duration of the period gathering at least a certain percent of births ("*nbtu*") are not to be

advised. In addition to undesirable statistical properties, these metrics fail to capture the

changes in the phenology parameter adequately. The metric "*nbtu*" varied non-monotonously

with the level of synchrony of the birth phenology. Similarly, the duration between first and

last birth to measure synchrony ("*per*") plateaued for a range of biologically realistic values,

what limits its usability in a wide range of ecological conditions.

Overall, some phenology characteristics have been more consistently evaluated across

studies, a fact illustrated by the number of metrics of each category used in more than two

papers ($n = 5$, 7, 2 and 0 for timing, synchrony, rhythmicity and regularity respectively, see

Supporting information 6). If timing and synchrony of births are the easiest and most frequent

characteristics of phenology estimated and compared, only a handful of metrics evaluates

rhythmicity and regularity of the phenology of births across the years. Sound analysis of

rhythmicity and regularity indeed requires many years of data which may not be available as

such data is costly and time-consuming to collect (Kharouba and Wolkovich 2020).

Moreover, scientists are less interested in timing and synchrony consistency *per se* than in the

relationship between timing and synchrony, and ecological or environmental factors such as

temperature, rainfall or spring snow cover (Paoli *et al.* 2018). Our study shows that the

rhythmicity and regularity metrics currently available are only moderately correlated,

particularly when they are used to describe birth distributions that are not normally

distributed (Supporting information 4). Capturing the temporal variation of phenology across

years appears difficult and requires thoughtful selection and interpretation of the used metric.

Standardized and relevant statistical tools are needed to quantify regularity and rhythmicity of

phenology, and to test their hypothetical responses to global changes. This study should help

in this.

Although we show that the assumption of a normal distribution or another bell-shaped

(asymmetric or not) distribution mimicking those found in *natura* (*e.g.* skewed normal,

bimodal or Cauchy distribution) has no major consequences on our conclusions (Supporting

information 4), this is not true when there is no clear pattern in the distribution of births.

Indeed, most metrics give inconsistent and unreliable results when applied to birth dates

randomly distributed within the year (Supporting information 4), a pattern that has been

documented in some populations of large herbivores living in the southern hemisphere

(Sinclair *et al.* 2000). Describing random patterns using the metrics presented here is unlikely

to be useful because biologically meaningless: when births occur year-round, the timing and

rhythmicity are meaningless as they cannot reduce to one or two summarizing statistics.

Using evenness indexes such as "*pielou*" could at least provide a quantification of the

heterogeneity of the distribution of births.

In conclusion, we recommend using the circular mean vector orientation ("*meanvo*")

to describe timing and the circular mean vector length ("*meanvl*") to describe synchrony,

because both are not influenced by the temporal origin set by the investigator. We

recommend using the underused Mood test which statistically compares the median birth

dates ("*mood*") to describe rhythmicity and the Kolmogorov-Smirnov test which statistically

assesses if two birth distributions are similar to describe regularity ("*kolmomult*", see Table 2

and Supporting information 3 for a formal description of those metrics). Being non-

parametric tests, they are applicable in a wide range of distributions as frequently observed in

large herbivores populations.

Declarations

Acknowledgments: The authors thank Simon Penel for his help with Perl Programming

Language, and Marcus Clauss for providing the code of one of the metrics. The authors thank

two anonymous reviewers and the subject editor for their comments on a previous version of

the manuscript, and Marion Valeix for her comments on early drafts of the manuscript. They

deeply appreciated the insightful comments, editorial suggestions and proofreading by Anna

Cryer, Leif Egil Loe and Primaëlle Fusto. This work was performed using the computing

facilities of the CC LBBE/PRABI and the IFB Cloud.

Funding: This work was supported by a grant from the "Ministère Français de

l'Enseignement Supérieur, de la Recherche et de l'Innovation" through the "Ecole Doctorale

E2M2" of the "Université Claude Bernard Lyon 1".

Authors' contributions: LT, CB and SCJ conceived the ideas and designed the

methodology; LT performed data collection and analysed the data; LT wrote the first draft of

the manuscript, which was then edited by all authors. All authors gave final approval for

publication.

Data availability statement: Code available on GitHub:

https://github.com/LucieThel/phenology-metrics.

References

Adams, L. G. and Dale, B. W. 1998. Timing and synchrony of parturition in Alaskan caribou. - J. Mammal. 79: 287–294.

Begon, M. *et al.* 1986. Ecology. Individuals, populations and communities. - Blackwell scientific publications.

Boness, D. J. *et al.* 1995. Does male harassment of females contribute to reproductive synchrony in the grey seal by affecting maternal performance? - Behav. Ecol. Sociobiol. 36: 1–10.

Bronson, F. H. 1989. Mammalian reproductive biology. - University of Chicago Press.

Bunnell, F. L. 1982. The lambing period of mountain sheep: synthesis, hypotheses, and tests. - Can. J. Zool. 60: 1–14.

Clauss, M. *et al.* 2020. Basic considerations on seasonal breeding in mammals including their testing by comparing natural habitats and zoos. - Mamm. Biol.: 1–14.

Crick, H. Q. P. and Sparks, T. H. 1999. Climate change related to egg-laying trends. - Nature 399: 423.

Darling, F. 1938. Bird flocks and the breeding cycle. - Cambridge University Press.

Demarée, G. R. 2011. From "Periodical Observations" to "Anthochronology" and "Phenology"- the scientific debate between Adolphe Quetelet and Charles Morren on the origin of the word "Phenology." - Int. J. Biometeorol. 55: 753–761.

Di Bitetti, M. S. and Janson, C. H. 2000. When will the stork arrive? Patterns of birth seasonality in neotropical primates. - Am. J. Primatol. Off. J. Am. Soc. Primatol. 50: 109–130.

English, A. K. *et al.* 2012. Reassessing the determinants of breeding synchrony in ungulates. - PLoS One 7: e41444.

Felsenstein, J. 1985. Phylogenies and the comparative method. - Am. Nat. 125: 1–15.

Festa-Bianchet, M. 1988. Birthdate and survival in bighorn lambs (Ovis canadensis). - J.

Zool. 214: 653–661.

Forrest, J. and Miller-Rushing, A. J. 2010. Toward a synthetic understanding of the role of

phenology in ecology and evolution. - Philos. Trans. R. Soc. B Biol. Sci. 365: 3101–3112.

Gosling, L. M. 1969. Parturition and related behaviour in Coke's hartebeest, Alcelaphus

buselaphus cokei Günther. - J. Reprod. Fertil. Suppl. 6: 265–286.

Heideman, P. D. and Utzurrum, R. C. B. 2003. Seasonality and synchrony of reproduction in

three species of nectarivorous Philippines bats. - BMC Ecol. 3: 11.

Heldstab, S. A. *et al.* 2018. Geographical Origin, Delayed Implantation, and Induced

Ovulation Explain Reproductive Seasonality in the Carnivora. - J. Biol. Rhythms 33: 402–

419.

Hodge, S. *et al.* 2011. Reproductive competition and the evolution of extreme birth

synchrony in a cooperative mammal. - Biol. Lett. 7: 54-56.

Ims, R. A. 1990. The ecology and evolution of reproductive synchrony. - Trends Ecol. Evol.

5: 135–140.

Kharouba, H. M. and Wolkovich, E. M. 2020. Disconnects between ecological theory and

data in phenological mismatch research. - Nat. Clim. Chang.: 1–10.

Landler, L. *et al.* 2018. Circular data in biology: advice for effectively implementing

statistical procedures. - Behav. Ecol. Sociobiol. 72: 128.

Morren, C. 1849. Le globe, le temps et la vie. - Bulletins de l'Académie royale des Sciences,

des Lettres et des Beaux-Arts de Belgique 16: 660–684.

Moussus, J.-P. *et al.* 2010. Featuring 10 phenological estimators using simulated data. -

Methods Ecol. Evol. 1: 140–150.

Newstrom, L. E. *et al.* 1994. A new classification for plant phenology based on flowering

patterns in lowland tropical rain forest trees at La Selva, Costa Rica. - Biotropica: 141–

159.

Paoli, A. *et al.* 2018. Winter and spring climatic conditions influence timing and synchrony

of calving in reindeer. - PLoS One 13: e0195603.

Parmesan, C. 2007. Influences of species, latitudes and methodologies on estimates of

phenological response to global warming. - Glob. Chang. Biol. 13: 1860–1872.

Plard, F. *et al.* 2015. The influence of birth date via body mass on individual fitness in a long-

lived mammal. - Ecology 96: 1516–1528.

Renaud, L.-A. *et al.* 2019. Phenotypic plasticity in bighorn sheep reproductive phenology:

from individual to population. - Behav. Ecol. Sociobiol. 73: 50.

Rubenstein, D. R. and Wikelski, M. 2003. Seasonal changes in food quality: a proximate cue

for reproductive timing in marine iguanas. - Ecology 84: 3013–3023.

Rutberg, A. T. 1987. Adaptive hypotheses of birth synchrony in ruminants: an interspecific

test. - Am. Nat. 130: 692–710.

Ryan, S. J. *et al.* 2007. Ecological cues, gestation length, and birth timing in African buffalo

(Syncerus caffer). - Behav. Ecol. 18: 635–644.

Sarkar, U. K. *et al.* 2019. Climato-environmental influence on breeding phenology of native

catfishes in River Ganga and modeling species response to climatic variability for their

conservation. - Int. J. Biometeorol. 63: 991–1004.

Sinclair, A. R. E. *et al.* 2000. What determines phenology and synchrony of ungulate

breeding in Serengeti? - Ecology 81: 2100–2111.

Skinner, J. D. *et al.* 2002. Inherent seasonality in the breeding seasons of African mammals:

evidence from captive breeding. - Trans. R. Soc. South Africa 57: 25–34.

Stearns, S. C. 1989. Trade-offs in life-history evolution. - Funct. Ecol. 3: 259–268.

van den Hoff, J. 2020. Environmental constraints on the breeding phenology of Giant Petrels

Macronectes spp., with emphasis on Southern Giant Petrels M. giganteus. - Mar. Ornithol.

48: 33–40.

Visser, M. E. *et al.* 2010. Phenology, seasonal timing and circannual rhythms: towards a

unified framework. - Philos. Trans. R. Soc. B Biol. Sci. 365: 3113–3127.

Zerbe, P. *et al.* 2012. Reproductive seasonality in captive wild ruminants: implications for

biogeographical adaptation, photoperiodic control, and life history. - Biol. Rev. 87: 965–

990.