1

Mechanical demands of the two-handed hardstyle kettlebell swing in novice older adults: an exploratory profile

Neil J. Meigh[1], Wayne A. Hing[1], Ben Schram[1], Justin W.L. Keogh[1,2,3]

[1] Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Queensland, Australia
[2] Sports Performance Research Centre New Zealand, AUT University, Auckland, New Zealand
[3] Kasturba Medical College, Manipal Academy of Higher Education. Mangalore, Manipal, Karnataka, India

Corresponding Author:
Neil Meigh[1]
Bond Institute of Health & Sport, 2 Promethean Way, Robina, QLD, 4226, Australia
Email address: nmeigh@bond.edu.au

Abstract

Background. Understanding the mechanical demands of an exercise and its technique increases clinical confidence when assessing the benefits and risks of a prescribed exercise. This study profiles the mechanical demands of the hardstyle kettlebell swing in novice older adults and compares peak force with kettlebell deadlifts. These data will help therapists choose the most appropriate kettlebell exercise (deadlifts or swings) and weights for progressing kettlebell training for older adults.

Methods. Thirty-five insufficiently physically active, community-dwelling males and females (59-79 years) were recruited. Two-handed hardstyle swings were performed with 8-16 kg and deadlifts with 8-24 kg and 8-32 kg for females and males, respectively. Ground reaction forces (GRFs) were obtained from a floor-mounted force platform. Force-time curves (FTCs), peak force, forward force relative to vertical force, rate of force development (RFD) and swing cadence were investigated. Peak GRF was compared by exercise and by sex, with RFD reported for swings.

Results. For kettlebells weights up to 16 kg, paired samples T-tests show a large exercise effect ($\delta > 1.4$) with peak force significantly higher for swings than deadlifts. Data shows: (i) significantly higher peak force during swings than deadlifts ($\delta = 1.77$), reaching 4.5 (1.0) N.kg^{-1}, (ii) peak force during an 8 kg swing was greater than a 32kg deadlift, (iii) no significant difference in normalised peak force between males and females performing kettlebell swings, but a moderately large effect size during deadlifts (males > females, $\delta = 0.69$), (iv) mean RFD of 19.9 (4.7) N.s^{-1}.kg^{-1} with a very weak, positive correlation with kettlebell weight ($y = 14.4 + 0.32x$), and trivial or non-

significant effect of sex, (v) mean forward force equal to 5.5% of vertical force during swings,
increasing from 3.8 (1.6) % with 8 kg to 1.7 (2.6) % with 16kg.

Conclusion. Where GRF is a therapeutic target, kettlebell swings with an 8 kg kettlebell could
have similar effects to much heavier deadlifts (>24 kg). Compared to kettlebell deadlifts, the
performance of kettlebell swings may be an easier, more convenient, and more appealing option
for older adults in a primary care setting or at home. The hardstyle swing with 8 kg has the potential
to produce double bodyweight in GRF and might be a suitable exercise to improve lower limb
RFD and physical function in older adults. Findings from this study were used to inform the BELL
Trial, a pragmatic clinical trial of kettlebell training with older adults. www.anzctr.org.au
ACTRN12619001177145.

Introduction

Clinical trials of kettlebell training with older adults are limited, but results have been encouraging
(Chen et al. 2018; Marcelino 2017). Small and portable, kettlebells have been recommended for
their ease of use, affordability, and for being less intimidating than other equipment (Otto et al.
2012). These are attractive features for clinicians and researchers who have used kettlebells in
rehabilitative programs for older individuals with hip fracture (Soukkio et al. 2018) sarcopenia
(Chen et al. 2018) and Parkinson's disease (Marcelino 2017).

An important component of therapeutic exercise is the choice of load and knowledge of resultant
force acting on the body and its tissues. A successful clinical outcome of therapeutic exercise may
depend upon the correct manipulation of force to stimulate a desired adaptive response in a target
tissue (Khan & Scott 2009). In older adults, repetitive loading is used in the management of hip
and knee arthritis, back pain, and osteoporosis (Smidt et al. 2005; Teichtahl et al. 2015). Healthcare
providers must first do no harm, which necessitates avoiding exposure to inappropriate volume,
intensity, or load, which may result in acute injury, excessive muscle soreness, or irritate an already
painful condition such as osteoarthritis. Understanding the magnitude of force acting on the body
during these exercises is therefore a necessary starting point.

For an older population, GRF and RFD are important measures of functional decline (Hester et al.
2019; LaRoche et al. 2011). Ground reaction force is associated with strength and power at the hip
and knee (Tsuji et al. 2015), gait speed (LaRoche et al. 2011), falls risk (Kera et al. 2020), and may
be a better predictor of functional decline than both grip strength (Hester et al. 2019) and the five-
times Sit-to-Stand test (Tsuji et al. 2015). Movement patterns observed in a Sit-To-Stand, floor
transfer, and reaching overhead, are also seen in kettlebell training; the most widely recognised
type being the 'hardstyle' swing introduced by Pavel Tsatsouline (2006). A hardstyle swing is
believed to improve lower limb power and by association, the ability to apply GRF, therefore, the
kettlebell swing might positively influence lower limb function in older adults, reduce risk of falls,
and be an ideal tool to promote healthy aging (Meigh et al. 2019).

A recent review (Meigh et al. 2019) identified no standards or guidelines to inform healthcare providers of how to use kettlebells safely and confidently with older adults. Additionally, current resistance training guidelines for older adults (Borde et al. 2015; Fragala et al. 2019; Lopez et al. 2017) do not translate to hardstyle kettlebell practices. Kettlebell weights are not chosen based on a percentage of '1 repetition maximum', and simply adding a kettlebell to an exercise (Soukkio et al. 2018) is not representative of 'kettlebell training'. It is a safety concern for providers, to prescribing a ballistic, free-weight resistance training program for older adults, without guidelines and for which the mechanical demands, acute responses, and risks are unknown.

Hardstyle literature suggests that outcomes are dependent upon technique (Tsatsouline 2006; Tsatsouline 2013). Kinematic differences between novice and expert have been described (Back et al. 2016) with significant, large effect size differences in GRF between novice and Instructor (Meigh et al. 2021), however, these data are from healthy young adults. Data from young active populations are informative and provide important reference points (Bullock et al. 2017; Lake et al. 2014; Lake & Lauder 2012b; Levine et al. 2020; McGill & Marshall 2012; Ross et al. 2017), but these cannot be generalised to older adults, especially those who are sedentary, insufficiently active or living with chronic health conditions. Without population-specific data, healthcare providers are just as likely to over-dose as under-dose the person in their care, potentially causing harm, or having no positive effect at all. A profile of the kettlebell swing and kettlebell deadlift in older adults is therefore warranted.

Therefore, the primary aim of this study was to profile the mechanical demands (GRF, RFD, forward force, and swing cadence) of the 2-handed hardstyle kettlebell swing in novice older adults. A secondary aim of this study was to provide context for the GRF of a hardstyle swing by comparing it to a kettlebell deadlift. These data were used to inform exercise prescription for the BELL trial (www.anzctr.org.au ACTRN12619001177145).

Materials & Methods

Participants.

A total of 17 males and 18 females aged 58-79 years were recruited. None of the participants had previously used kettlebells. Thirty-two identified as sedentary, not having engaged in a structured exercise program for at least nine months, and few had experience with commercial resistance equipment. Participants were free from injury and did not disclose any health or medical conditions considered to be high risk (ESSA 2011). After a thorough explanation of the study aims, protocols, and potential risks, participants provided written informed consent. Ethical approval for this study was granted by Bond university Human Research Ethics committee (NM03279).

Protocol

Data were collected from the University biomechanics laboratory, each participant attending two 1-hour sessions on consecutive weeks. Two-handed kettlebell swings to chest-height were

performed on a floor-mounted force flatform (AMTI, Watertown, NY, USA) recording GRF at
1000 Hz using NetForce software (AMTI, USA). Participant body mass was captured by the force
plate from a period of quiet standing. Tri-plantar force variables were obtained from the floor-
mounted force platform. The variables of interest were peak GRF, dynamic RFD, and swing
cadence. Participants performed a single set of 12 repetitions with each kettlebell, with the middle
10 repetitions used for analysis. A custom program (Microsoft Excel, Version 2012) was used to
calculate peak force during each swing cycle of the set, with values manually assessed and verified
against the corresponding FTC. To obtain peak net force, system weight (body mass + kettlebell
weight) was subtracted from the square root of squared and summed data:

$\sqrt{F_z^2 + F_x^2 + F_y^2}$ (F_z = vertical force, F_y = horizontal forward force, F_x = medio-lateral force).

The back or bottom position of the swing was used as the start of each swing cycle. Dynamic RFD
(N.s^{-1}) during hip extension (propulsion) was calculated as the change in resultant GRF during
Phase 1, divided by elapsed time and normalised to body mass (N. s^{-1}.kg^{-1}), with this reported as
the mean of 10 swings. Cadence in swings per minute (SPM) was calculated from the average time
between peak force of hip extension in each swing cycle. Peak force was reported as resultant force
unless stated otherwise.

Figure 1. Key positions during the two-handed hardstyle kettlebell swing and kettlebell deadlift: dead-start / dead-
stop for a swing (A), start and end position of single swing cycle (B), mid-swing (C), dead-start for a kettlebell
deadlift (D), top position of the deadlift. Red tape marks the boundary of the force platform (E).

Procedure.
During the first session, participants received instruction on how to perform the hardstyle swing
as previously described (Meigh et al. 2021; Thomas 2014). The following drills served as a warm-
up; (1) unloaded hip-hinge - standing at a distance roughly equal to the length of their foot away
from a wall, participants used the wall as an external focus of attention (target for their buttocks to
touch) to practice the 'hinge' movement, before returning to an upright position. (2) performing
the 4 phases of a swing cycle (Meigh et al. 2021) without weight, led by the Instructor to a 4-count.
(3) double knee extension swings (Del Monte et al. 2020) with 8 kg to familiarise themselves with
the external load, bell movement through the legs, and weight shift. (4) towel-swings (Tsatsouline
2006) with 4-8 kg to differentiate a 'swing' from a 'lift' and (5) two to three sets of Instructor-led

swings performed at cadence of approximately 40 SPM, to illustrate the intended ballistic nature
of the hardstyle swing.
152
Force USA competition kettlebells from 8-32 kg of standardised dimensions were used, increasing
in 2 kg increments to 24 kg, then 4 kg increments to 32 kg. Target weights for swings and deadlifts
were pre-selected based on the lead investigator's experience as a kettlebell Instructor; 12 kg and
24 kg for females, and 16 kg and 32 kg for males, respectively. Sets of swings began and ended in
a 'dead-start' position (Fig. 1A). Females were given the opportunity to swing 14 kg if their 12kg
swing was performed confidently with acceptable technique. Males were not given the opportunity
to swing heavier than 16kg. An ideal hardstyle swing pattern was not required or expected.
Deadlifts were performed with feet positioned either side of the bell (Fig. 1D) as previously
described (Thomas et al. 2014). Only peak resultant force was analysed for deadlifts.
162

Statistical analyses.
Descriptive statistics were reported for normally distributed continuous variables. Measures of
centrality and dispersion are presented as mean (SD) and verified for normality using box-plots,
histograms, and Shapiro-Wilk tests. Differences between males and females, swings and deadlifts,
were analysed using independent samples T-tests and paired samples T-tests, respectively, subject
to the sampling distribution being normal for each group. Mean Difference (MD) is presented with
95% confidence interval (CI). Effect sizes were calculated and interpreted using Lenhard and
Lenhard (2016) and Magnusson (2021), quantified as trivial, small, moderate, large, very large,
and extremely large where effect size < 0.20, 0.20 - 0.59, 0.60 - 1.19, 1.20 - 1.99, 2.0 - 3.99 and $\geq$
4.0 respectively (Hopkins et al. 2009). Probability of superiority has been used to illustrate the
Cohen's d effect size, representing the chance that a person from group A will have a higher score
than a person picked at random from group B (Magnusson 2021). The relationships between peak
net ground reaction force and kettlebell weight for swings and deadlifts, and RFD and kettlebell
weight during swings, was investigated using simple linear regression. Preliminary analyses were
performed to ensure no violation of the assumptions of normality, linearity and homoscedasticity.
Statistical analyses were performed using SPSS (version 26.0; SPSS Inc., Chicago, IL, USA), and
$p < 0.05$ was used to indicate statistical significance.
180

181 ## Results
182 Participant demographics are presented in Table 1. All females performed swings with 8-12 kg
183 kettlebells. Nine females swung 14 kg and two swung 16 kg. All males performed swings with 8-
184 12 kg. Sixteen males swung 14 kg and 15 swung 16 kg.
185

 Table 1. Participant characteristics

Characteristic	Older Males	Older Females
Number	17	18
Age (years)	68.5 (4.2)	68.1 (4.0)
Height (cm)	176.8 (7.8)	163.9 (5.4)
Weight (kg)	90.2 (14.9)	71.7 (11.5)
BMI (kg/m^2)	29.1 (2.5)	26.6 (4.5)

Data are presented as mean (SD), unless otherwise specified.

Peak net ground reaction force.

There was a large ($\delta = 1.77$) significant difference in peak force between the kettlebell swing and kettlebell deadlift (Fig. 2 and Table 2), with the mean probability of superiority– for the kettlebell swing across loads of 8-16 kg being 89.5%. The relationship between peak force and kettlebell weight was investigated. In both cases, there was a very weak ($r < 0.30$) positive correlation between the two variables. Regression coefficients to predict peak force where x = kettlebell mass in kg.

Peak force - *swing* $y = 3.10 + 0.11x$ ($r = 0.26$, $p = 0.001$, CI [0.05, 0.81])
Peak force - *deadlift* $y = 2.63 + 0.02x$, ($r = 0.19$, $p = 0.001$, CI [0.01, 0.03])

Figure 2. Box-and-whisker plot of large significant difference in peak net ground reaction force between kettlebell swings and deadlifts Box represents median observations (horizontal rule) with 25th and 75th percentiles of observed data (top of bottom of box). Length of each whisker is 1.5 times interquartile range. Data points that are more than 1.5 x interquartile range are represented by "o" (Tukey's outlier detection method). * = $p < 0.05$

206 **Tables 2.** Peak net ground reaction force for swings and deadlifts (N.kg^{-1})

KB weight (kg)	Mean (SD)		MD (95% CI)	t	p	Cohen's	
	Swing	Deadlift				δ	Probability of superiority (%)
8	3.9 (1.0)	2.8 (0.6)	1.2 (0.8, 1.6)	29 (6.110)	<0.001	1.42	84.2
10	4.3 (1.0)	2.8 (0.7)	1.5 (1.2, 1.9)	29 (9.408)	<0.001	1.76	89.3
12	4.4 (0.8)	2.8 (0.6)	1.5 (1.2, 1.9)	29 (9.927)	<0.001	1.99	92.0
14	4.7 (1.0)	3.0 (0.6)	1.7 (1.3, 2.1)	22 (8.887)	<0.001	2.00	92.1
16	4.5 (1.0)	3.1 (0.6)	1.4 (0.8, 2.0)	15 (5.067)	<0.001	1.68	88.3

Only trivial-to-small non-significant effects were found between males and females performing
kettlebell swings up to 16 kg, however sex influenced peak force during kettlebell deadlifts. From
8-24 kg, peak force during deadlifts was moderately higher ($\delta = 0.69$) for males than females, with
a 68.7% probability of superiority. There was a statistically significant difference between males
and female using 8, 12 and 14 kg kettlebells, with the mean probability of superiority being 71.2%
($\delta = 0.79$) (Fig. 3 and Table 3).

215
216 **Figure 3.** Moderately large effect of sex on peak force during kettlebell deadlifts with 68.7% mean probability of
217 superiority (males > females) with 8-24 kg kettlebells. Data points that are more than 3 x interquartile range are
218 represented by "☆". * = p <0.05, ns = p >0.05

220   **Table 3.** Peak net ground reaction force for males and females during kettlebell deadlifts (N.kg⁻¹)

KB weight (kg)	Mean (SD)		MD [95% CI]	t	p	Cohen's	
	Male	Female				δ	Probability of superiority (%)
8	3.0 (0.7)	2.5 (0.4)	0.4 [<0.05, 0.87]	28 (2.112)	0.044	0.77	70.7
10	3.0 (0.7)	2.6 (0.6)	0.3 [-0.2, 0.8]	28 (1.318)	0.198	0.48	63.3
12	3.1 (0.7)	2.6 (0.4)	0.5 [<0.05, 0.9]	28 (2.242)	0.033	0.82	71.9
14	3.1 (0.6)	2.7 (0.6)	0.4 [<0.05, 0.9]	28 (1.970)	0.059	0.72	69.5
16	3.2 (0.6)	2.7 (0.6)	0.5 [<0.05, 0.9]	28 (2.114)	0.044	0.77	70.7
18	3.1 (0.6)	2.8 (0.6)	0.3 [-0.1, 0.7]	28 (1.417)	0.168	0.52	64.3
20	3.2 (0.6)	2.8 (0.6)	0.4 [<0.05, 0.8]	28 (1.899)	0.068	0.69	68.7
22	3.2 (0.6)	2.8 (0.5)	0.3 [<0.05, 0.7]	28 (1.797)	0.083	0.66	68.0
24	3.3 (0.6)	2.9 (0.6)	0.4 [<0.05, 0.8]	28 (1.861)	0.073	0.77	10.7
28	3.4 (0.7)	2.7 (0.2)	0.8 [-0.3, 1.8]	14 (1.537)	0.147	1.16	79.4
32	3.6 (1.0)	2.6 (0.5)	1.0 [-0.5, 2.5]	12 (1.408)	0.184	1.08	77.7

Rate of force development

Rate of force development was not significantly different between consecutive kettlebell weights.
There was a significant difference (δ = 0.5 - 0.7) in RFD between the 8 kg and 14 kg kettlebell,
and the 8kg and 16 kg kettlebell: MD = 2.2 (0.5, 3.9), t = (24) 2.721, p = 0.012, and MD = 3.89
(1.59, 6.20), t = (16) 3.577, p = 0.003, respectively (Fig. 4 and Table 4).

Analysis of the effect of sex on RFD revealed a significant difference between males and females,
but only with 8 kg (Fig. 4). Except for the n = 2 females who swung 16 kg, paired samples analysis
revealed no significant increase in RFD between the lightest and heaviest weights for males or
females. The relationship between RFD and kettlebell weight was investigated. Linear regression
shows a very weak, positive correlation between the two variables where y (RFD) = 14.4 + 0.32x
(r = 0.03, p = 0.032, CI [0.03, 0.61]) where x = kettlebell mass in kg.

Figure 4. Trivial effect of sex on RFD. * = p <0.05, ns = p >0.05. One outlier at 16 kg and one extreme outlier at 12
kg not shown.

Table 4. Rate of force development during kettlebell swings for males and females ($N.s^{-1}.kg^{-1}$)

KB weight (kg)	Mean (SD)		MD [95% CI]	t	p	Cohen's	
	Males	Females				δ	Probability of superiority (%)
8	15.5 (4.9)	18.7 (3.6)	3.3 [0.3, 6.6]	(33) 2.255	0.031	0.75	70.2
10	16.7 (4.7)	17.8 (3.7)	1.1 [-1.8, 4.0]	(33) 0.776	0.443	0.26	57.3
12	18.3 (6.5)	18.1 (3.2)	-0.2 [-3.7, 3.3]	(33) -0.111	0.913	0.04	51.1
14	18.4 (4.7)	18.8 (4.1)	0.5 [-3.4, 4.3]	(23) 2.47	0.807	0.09	52.5
16	19.3 (5.2)	21.8 (9.3)	2.5 [-6.5, 11.5]	(15) 0.560	0.560	0.45	62.5

Forward force

Mean forward force was 0.055 (5.5%) of the vertical GRF during the kettlebell swings. There was
a small (δ = 0.34) increase in the ratio of forward to vertical GRF between 8 and 16 kg, MD = 0.33
(0.21, 0.45), t = (14) 5.890, p = <0.001. The trivial to small differences between males and females
were not significant at any weight (Fig. 5). Between 8 and 14 kg, the absolute change in ratio of
horizontal to vertical force was 2.5% for females, increasing from 3.7% to 6.9%. For males, the
absolute change was 2.6%, from 3.8% with 8 kg to 7.3% with 16 kg.

Figure 5. No significant effect of sex on small increases in forward force with increasing kettlebell weight. A/V = ratio of *anterior* (forward) horizontal to *vertical* GRF.

Cadence

Mean swing cadence up to 16 kg for the kettlebell swings was 35.7 (2.3) SPM, range = 29.2 to 41.2. There was a statistically significant difference ($p < 0.05$) in cadence between 10 and 12 kg and 12 and 14 kg swings, however, means remained between 35 and 36 SPM. The effect of sex on cadence revealed significant differences between males and females at all weights (Fig. 6). Between 8 and 16 kg, there was a moderately large ($\delta = 0.9$) effect size, with a mean difference of 1.9 SPM and 73.8% probability of superiority (females > males).

Figure 6. Moderately large effect of sex on swing cadence. SPM = swings per minute. * = p <0.05

Force-time curve characteristics

Three distinct movement strategies emerged, which were inconsistent with a proficient swing
(Meigh et al. 2021). 1) lifting the bell with the upper limbs, 2) a whole-body effort to perform the
movement, 3) controlling the bells' decent from mid-swing, with little visual dissociation between
upper and lower limbs, in which the body appeared to collapse downward with the bell's vertical
displacement. Representative FTCs recorded from two participants which illustrate these
movement strategies, are shown in Fig. 7. A double-peak (Fig. 7A) is consistent with a sequential
movement, initially involving explosive hip extension, immediately followed by active shoulder
flexion to lift the bell to the desired height. A wider, multi-peaked profile (Fig. 7B) is consistent
with a more simultaneous movement, and lack of ballistic hip extension during propulsion.
Absence of a second smaller force peak (shown in Fig. 7A) is consistent with not allowing the bell
to drop from mid-swing. These profiles illustrate the effect of technique on force and cadence. The
ballistic pattern more consistent with a proficient hardstyle swing (A), resulted in higher peak net
force (368.2 N vs 257.1 N), higher force as a percentage of bodyweight (47.5 % vs 33.8 %), a
larger proportion of horizontal forward force (6.8% vs 2.9 %), and faster swing cadence (36.4 SPM
vs 34.1 SPM).

Figure 7. Representative force-time curves from two distinct movement strategies used by novice older adults to perform a two-handed hardstyle kettlebell swing.

Discussion

Knowledge of GRF and other kinetic variables enables healthcare providers to make informed decisions about the potential benefits and risks of an exercise. A properly designed resistance training program for older adults, includes multi-joint exercises such as the kettlebell swing and kettlebell deadlift, is individualised, periodised, progressive, and includes appropriate technique instruction (Fragala et al. 2019). A review (Meigh et al. 2019) was unable to identify descriptive data of kettlebell exercises from older adults for any of these variables. Results of the present study allow healthcare providers to make a more informed decision about whether, and how to use the kettlebell swing as a therapeutic exercise for older adults.

Ground reaction force

It was unsurprising that peak net GRF was significantly higher for swings than deadlifts. Force is a product of mass and acceleration and increases with speed in all environmental conditions (Heywood et al. 2019). Peak net ground reaction was 5% higher during swings with an 8 kg kettlebell than deadlifts with 32 kg, and 35% higher than deadlifts with 16 kg. If GRF is the desired

300 therapeutic stimulus, an 8 kg kettlebell swing could be a better option than a heavier deadlift. Other
301 factors such as reduced cost, ease of storage and transport, and availability in a variety of general-
302 purpose stores, could make use of lighter kettlebells for the swing less intimidating and more
303 attractive than the heavier kettlebells that would be required for the deadlift.
304
Ground reaction force during a proficient swing with an 8 kg kettlebell can exceed 2x body mass
(Meigh et al. 2021). For the novice older adults, net peak force during an 8kg kettlebell swing was
3.9 (1.0) $N.kg^{-1}$, compared to 8.85 $N.kg^{-1}$ for the younger Instructor. Whether force of this
magnitude can be attained by older adults using kettlebells of this mass remains to be seen, but it
is possible, and highlights opportunities for therapeutic exercise in this population. Lower limb
strengthening for example, is recommend for management of a host of chronic diseases including
symptomatic knee osteoarthritis (Goh et al. 2019; Turner et al. 2019). During the kettlebell swing,
knee flexion is limited (Back et al. 2016) and there are no large ground contact impacts, both
mechanisms which are commonly reported to irritate symptomatic arthritic knees. At a cadence of
40 swings per minute, a training load volume of 1,600 kg can be achieved in as little as five
minutes. Safe, easily achievable, and readily scalable, the swing could be an effective and efficient
option where time, motivation or desire might be limited. Ground reaction force relative to
bodyweight during an 8 kg swing is comparable to a bodyweight squat performed at a medium to
fast pace (Heywood et al. 2019), but the number of bodyweight squats a healthcare provider might
prescribe for an older adult, is likely to be far less than the number of swings that could be
performed in the same period of time. This may have significant implications for the prescription
of training load volume.
322
More than half of the female participants voluntarily swung a kettlebell heavier than the intended
12 kg maximum. It was also evident that some of the males had capacity to swing heavier than 16
kg. As these participants were all novices with respect to kettlebell training, this was very
encouraging to the potential feasibility of kettlebell swings in this population. Whether there is any
physical benefit from using heavier weights has yet to be determined, particularly as effort
increases disproportionately to kettlebell weight, and change in GRF is small (Meigh et al. 2021).
If the clinical benefits can be gained with a lighter rather than heavier weight, it appears the
potential risks of swinging a heavier bell could quickly outweigh the reward. To minimise the risk
of injury, it seems reasonable that 12-16 kg would be "sufficient" for most older adults, with other
program variables available to alter the physiological demand, such as the work:rest ratio, altering
technique and cadence, and option to use one-arm instead of two.
334
Males have a higher volume of lean muscle mass than females (Aguado-Henche et al. 2017;
Janssen et al. 2004; Pasco et al. 2020), so it was anticipated that peak force would be higher for
males. This was true for deadlifts but not for swings, which was surprising. Although the difference
in peak force for the deadlift was statistically significant for only three of the nine kettlebells up to
24 kg, there was a consistent moderate effect size with an almost 70% mean probability of

superiority of peak force being higher for males. There was a sex effect for deadlifts, however the
lack of such significance for swings is currently unclear.

There was a very large mean effect size (δ = 3.06) in peak force between the older adults in this
study and data reported from an Instructor (Meigh et al. 2021). For older adults, the average peak
force during swings was almost half that of the Instructor (52.1%), with a mean difference of 4.2
N.kg^{-1}, highlighting scope for potential improvement from training. Age has previously been
shown to have no effect on peak GRF during lifting (Lim et al. 2018b; Shin et al. 2006), but
moderate differences between sex have been observed during Sit-To-Stand tasks in community
dwelling older adults (males > females) (Tsuji et al. 2015). In the absence of sex-specific health
or medical conditions, these results suggest that choice of kettlebell weight should be influenced
by bodyweight, strength and/or stature rather than sex.

Novice starting weights for the kettlebell swing are commonly based upon recommendations
which are not specific to the swing or older adults (Tsatsouline 2006). Researchers have largely
followed these recommendations, however, some investigators have used weights as low as 3 kg
(Duncan et al. 2015; Fortner et al. 2014; Lim et al. 2018a; Rajala & Jensen 2015; Zin et al. 2018),
heavier weights up to 48 kg (Bullock et al. 2017; Del Monte et al. 2020; Falatic et al. 2015;
Holmstrup et al. 2016; Lake & Lauder 2012a; Oikarinen 2016; Santillo 2016; Smith et al. 2014),
or a percentage of bodyweight e.g. 10-20% (Mache & Hsieh 2016; Ros 2016). Results from this
study suggest that healthy novice females, regardless of age and training status, may be able to
perform swings safely a with 12-14 kg if they feel comfortable doing so.

Forward force
"Street wisdom" (McGill & Marshall 2012) suggests that most of the applied force is in the forward
horizontal direction. This belief has influenced training practices to include visualising the bell
being projected forward during the propulsion phase, and forcibly snapping the knees into full
extension. A recent kinetic analysis of a proficient hardstyle swing, revealed horizontal forward
force to be approximately 10-15% of the magnitude of vertical force, with kettlebell weights from
8kg to 32kg (Meigh et al. 2021). One of the most striking findings from this study, was that only
around 5% of the ground reaction of a swing was directly horizontally forward. Mean forward
force increased from just 3.8% with 8 kg to 7.1% with 16 kg. A small proportion of forward force
was expected, however, a mean of 5.5% with few swings exceeding 10% was surprisingly low.

There was a very large (δ = 2.05) average effect size in the magnitude of forward force between
older adults and Instructor (Meigh et al. 2021). Although the absolute mean difference was less
than 5%, the relative difference was just over half (55.1%). In some cases, horizontal force at peak
ground reaction was zero, and in a few instances, negative, albeit negligible in magnitude i.e.,
indicating that the kettlebell was in effect being pulled backward rather than projected forward at
the point of peak effort. This is an interesting biomechanical scenario given the kettlebells'

anteroposterior arc of motion and the street wisdom regarding forward force. Tsatsouline describes
the motion of the lower limbs during a kettlebell swing as akin to a standing vertical jump
(Tsatsouline 2006). Extending the knees as rapidly as possible with minimal force applied to the
ground, is likely counterproductive for achieving optimal performance during a kettlebell swing.
For older adults with painful arthritic knees, forcibly extending the joints may exacerbate
unwanted symptoms.
386
387 **Rate of force development**
388
Data from a proficient swing showed a strong negative correlation (r = -0.82) in RFD with
kettlebell weight (Meigh et al. 2021), so it was surprising to find a neutral correlation among the
novice older adults in this study (r = 0.03). The effect size between the older adults in this study
and Instructor was large (δ = 1.47), with a mean difference of 17.0 N.s^{-1}.kg^{-1}; almost half that of
the Instructor (54.2%). With large variation within the group, it is interesting to consider the
outliers. One male participant in the current study, who was still engaged in active employment,
recorded a higher RFD than the Instructor with the 12 kg kettlebell (δ = 0.25). The discrepancy in
this older male participant's RFD values and that of the other older adults in the present study is
interesting and may suggest that there exists the potential for some individuals to maintain high
levels of RFD into older age. Results from strength and power training in older adults (Caserotti
et al. 2008) indicate that it is feasible that kettlebell swings might increase RFD. Combined with
large improvements in dynamic single leg balance (Beltz 2013) and postural reaction time
(Manocchia et al. 2013), kettlebell training may mitigate age-related changes in RFD (Maffiuletti
et al. 2016) and reduce falls risk. Whether swings might provide superior clinical benefit in this
regard to a more basic deadlift alone remains to be seen.
404
405
406
Swing cadence
A swing cadence significantly lower than 40 SPM was expected. There was a very large (δ = 2.54)
average effect size between older adults and Instructor (Meigh et al. 2021), with the older adults
performing swings on average 6 SPM slower. Females however, having a cadence almost 2 SPM
higher than males was surprising. With kettlebell proficient females of comparable height
performing swings at 40 SPM (Wesley 2013), there is currently insufficient data to account for the
effect of sex with factors such as height and upper limb length i.e. a taller person with longer arms
will displace the kettlebell a greater distance. It was encouraging to see that cadence remained
relatively consistent throughout the group, indicating that the pre-selected weights were not 'too
heavy'. In current practice, it is unlikely that insufficiently physically active females over 60 years
of age would be asked to swing a 12-14 kg kettlebell, or men over 60 years asked to swing 16 kg.
These data could be used to challenge common practice regarding weight selection for older adults.
419

Force-time profile

The representative FTCs are more consistent with those of Mache and Hsieh (2016) and Lake and Lauder (2012b) than a proficient swing (Meigh et al. 2021). Our observations support the findings of Back and colleagues (2016) with one or both phases of 'lift' and 'float' frequently absent. Observations of technique and deviations from "ideal" could be correlated with changes in the corresponding FTC. A common strategy for some older adults was to actively flex the shoulders to lift the bell upward rather than swing it, typically corresponding with a double-peak FTC. A second smaller force peak seen in a proficient swing (Meigh et al. 2021) was likely absent if the person was observed to control the bells' decent from mid-swing, instead of allowing it to drop. Practically, this strategy shifts emphasis of the mechanical load of deceleration from the hips and lower limbs to the trunk and upper limbs.

Less visually apparent was the squatting motion previously observed in novices (Back et al. 2016). This relative lack of a squatting motion observed in the older participants in the current study was attributed to the pre-test drills encouraging the 'hip-hinge' pattern performed by the instructor with the participants in the current study. A wider, multi-peaked FTC and slower cadence was consistent with a more pronounced simultaneous movement pattern. Whether differences in swing pattern are clinically meaningful has not been investigated. It is possible that technique may influence performance improvement and rehabilitation conditions. An 'ideal' swing may not be achievable for some, and some between-participant variability in technique could be beneficial.

As anticipated, none of the participants were able to perform an ideal hardstyle swing consistently for all repetitions with all loads. There were, however, many examples where it was evident in the FTC, that participants had performed one or more swings within the set, which were approaching ideal (Supp. file A: M4, M6, M7, M9, M12, M13, M15, M16, M17, F1, F3, F5, F9, F11-15, F17, F18). Most participants had a force profile suggestive of a swing rather than lift, which was surprising and very encouraging. In addition to common drills, use of real-time biofeedback where available, and slow-motion video analysis, are likely to be beneficial for novices learning how to perform a hardstyle swing (Beerse et al. 2020).

The hardstyle swing may be considered to be a movement pattern that has some unusual characteristics and that may be challenging to learn for some individuals, especially older adults. While a range of hip hinge and ballistic drills can be effective for improving technique, some people can find them confusing and complicated to follow, a situation which can be compounded with multiple drills. Many of the participants were able to quickly improve their swing technique simply for focusing on increasing cadence. Swing cadence could be a useful measure to determine an appropriate upper limit for the most appropriate kettlebell weight, particularly in combination with an increasing rate of observed or reported effort. Achieving a consistent cadence of 40 SPM with a comfortable (lighter) kettlebell appears to be an appropriate strategy for the novice individual learning the swing. Health professionals should note that the cardiovascular demand of

460 kettlebell swings may be higher than walking (Thomas et al. 2014) and should anticipate a high
461 heart rate and cardiovascular response (Schnettler et al. 2009; Wong et al. 2017) and program for
462 the individual accordingly.
463
Our suggestion that changes in a FTC might be used to determine an optimal training weight
(Meigh et al. 2021), assumes a proficient swing and availability of real time biofeedback. So how
should optimal kettlebell weight be determined for novice older adults? For some participants,
increasing bell weight appeared to improve their technique, but for others, additional weight had
a detrimental effect. It remains to be seen whether monitoring mechanical demands to inform
weight selection provides any benefit beyond the existing pragmatic hardstyle approach:
Optimising starting weights and progression during the initial and early stages of learning a
hardstyle swing is where input from an appropriate experienced provider is likely to be most
beneficial.
473
The influence of confounding variables such technique, age, and training history on resultant force
are unclear. Pragmatically however, the combined testing loads for the deadlift —session were
1,728 kg and 2,448 kg for females and males, respectively. With many older adults routinely
under-dosed in resistance training programs (APTA 2014), these data might help change that trend.
Acknowledging that clinical reasoning should influence instruction around speed, movement,
treatment goals and individual preferences (Heywood et al. 2019), in the absence of identifiable
risks and within the constraints of a person's unique health history, clinical needs, and the desire
(or not) to engage in a program of resistance training, these findings might encourage providers to
broaden their perceived scope of therapeutic load.
483
Limitations. Results from predominantly insufficiently physically active older adults may not be
generalisable to all older adults, or those for whom a diagnosed health or medical conditions
presents an increased risk of harm from resistance exercise, or which effects their ability to perform
the exercise. Ground reaction force and technique among novices is likely to quickly improve with
practice and appropriate instruction, therefore, these data may not represent older adults who have
undergone a period of kettlebell exercise practice and training. Rate of force development should
be interpreted with caution. Specifically, in the absence of concurrent video analysis, calculation
of RFD from force-plate data of novices where the propulsive phase is not clear, is subject to
interpretation and reduces reliability. Ground reaction data from hardstyle swings cannot be
generalised to the double knee-bend swing (kettlebell Sport) or overhead (American) swing which
are kinematically different (Del Monte et al. 2020; Mitchell et al. 2015).
495

Conclusions

497 Results of this exploratory profiling study suggest that the prescription of exercise intensity
498 (weight) for kettlebell swings and kettlebell deadlifts for older adults should not be determined by
499 sex. All older females and males comfortably performed multiple deadlifts up to 24 kg and 32 kg,

respectively. Healthcare providers are encouraged to explore the potential of their older patients/clients in this regard. A two-handed hardstyle kettlebell swing with 8 kg resulted in greater ground reaction force than a 32 kg deadlift. Lighter weight bells may be a more cost-effective, convenient, and appealing option for therapists and patients than heavier alternatives. Where GRF is the clinical measure of interest, an 8 kg swing has the potential to provide similar acute kinetic demands as deadlifts with a much heavier bell, and likely to have a greater influence on increasing lower limb RFD. Further research is warranted to investigate the long-term effects of using only dynamic movements such as the swing compared with slower, heavier movements such as the deadlift. A target cadence of 40 SPM is likely to positively influence RFD and GRF in novices, but further research is required to establish the influence of hardstyle technique on clinical outcomes of interest in older adults.

Acknowledgements

The authors would like to acknowledge Mr Benjamin Hindle for his support in the calculation of rate of force development, and Mrs Evelyne Rathbone for her direction with statistical analysis.

Funding

This study was supported by an Australian Government Research Training Program Scholarship and will contribute towards a Higher Degree by Research Degree (Doctor of Philosophy).

Contributions

NM recruited the participants and conducted the study, curated and analysed the data, interpreted the results, conducted the formal analysis, and wrote the original draft. JK, BS and WH supported with ongoing consultation. JK and WH reviewed and provided revisions to earlier versions of the manuscript. All authors read and approved the final manuscript.

Ethics declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The primary author is a Physiotherapist and hardstyle kettlebell instructor, with an online presence as The Kettlebell Physio.

Rights and permissions

References

Aguado-Henche S, Carrascosa-Sánchez J, Cristóbal-Aguado S, and Clemente de Arriba C. 2017. Normative values
for body compartments of sedentary white people. *MOJ Anat Physiol* 4:217-225.
APTA. 2014. Don't prescribe under-dosed strength training programs for older adults. Instead, match the frequency,
intensity and duration of exercise to the individual's abilities and goals. *Available at*
http://ow.ly/O9C350E8Sb7 (accessed Nov 10, 2017 2017).
Back C-Y, Joo J-Y, and Kim Y-K. 2016. Kinematic comparisons of kettlebell two-arm swings by skill level. *Korean*
Journal of Sport Biomechanics 26:39-50.
Beerse M, Bigelow KE, and Barrios JA. 2020. The patterning of local variability during the acquisition of a novel
whole-body continuous motor skill in young adults. *Experimental Brain Research*.
Beltz. 2013. Effects of kettlebell training on aerobic capacity, muscular strength, balance, flexibility, and body
composition. . *Journal of Fitness Research* 2:4-13.
Borde R, Hortobagyi T, and Granacher U. 2015. Dose-Response Relationships of Resistance Training in Healthy
Old Adults: A Systematic Review and Meta-Analysis. *Sports Medicine* 45:1693-1720. 10.1007/s40279-
015-0385-9
Bullock GS, Schmitt AC, Shutt JM, Cook G, and Butler RJ. 2017. Kinematic and Kinetic Variables Differ Between
Kettlebell Swing Styles. *International Journal of Sports Physical Therapy* 12:324-332.
Caserotti P, Aagaard P, Buttrup Larsen J, and Puggaard L. 2008. Explosive heavy-resistance training in old and very
old adults: changes in rapid muscle force, strength and power. *Scandinavian Journal of Medicine & Science*
in Sports 18:773-782. 10.1111/j.1600-0838.2007.00732.x
Chen HT, Wu HJ, Chen YJ, Ho SY, and Chung YC. 2018. Effects of 8-week kettlebell training on body
composition, muscle strength, pulmonary function, and chronic low-grade inflammation in elderly women
with sarcopenia. *Experimental Gerontology* 112:112-118. 10.1016/j.exger.2018.09.015
Del Monte MJ, Opar DA, Timmins RG, Ross JA, Keogh JW, and Lorenzen C. 2020. Hamstring myoelectrical
activity during three different kettlebell swing exercises. *The Journal of Strength & Conditioning Research*
34:1953-1958.
Duncan M, Gibbard R, Raymond L, and Mundy P. 2015. The Effect of Kettlebell Swing Load and Cadence on
Physiological, Perceptual and Mechanical Variables. *Sports* 3:202-208. 10.3390/sports3030202
ESSA. 2011. Adult Pre-Exercise Screening System Tool (V1). *Available at http://ow.ly/Se9y50E8S9B2018*).
Falatic JA, Plato PA, Holder C, Finch D, Han K, and Cisar CJ. 2015. Effects of Kettlebell Training on Aerobic
Capacity. *Journal of Strength and Conditioning Research* 29:1943-1947. 10.1519/jsc.0000000000000845
Fortner HA, Salgado JM, Holmstrup AM, and Holmstrup ME. 2014. Cardiovascular and Metabolic Demands of the
Kettlebell Swing using Tabata Interval versus a Traditional Resistance Protocol. *Int J Exerc Sci* 7:179-185.
Fragala MS, Cadore EL, Dorgo S, Izquierdo M, Kraemer WJ, Peterson MD, and Ryan ED. 2019. Resistance
Training for Older Adults. *Journal of Strength and Conditioning Research* 33:2019-2052.
10.1519/jsc.0000000000003230
Goh S-L, Persson MSM, Stocks J, Hou Y, Welton NJ, Lin J, Hall MC, Doherty M, and Zhang W. 2019. Relative
Efficacy of Different Exercises for Pain, Function, Performance and Quality of Life in Knee and Hip
Osteoarthritis: Systematic Review and Network Meta-Analysis. *Sports medicine (Auckland)* 49:743-761.
10.1007/s40279-019-01082-0
Hester G, Ha P, Dalton B, VanDusseldorp T, Olmos A, Stratton M, Bailly A, and Vroman T. 2019. Rate of Force
Development as a Predictor of Mobility in Older Adults. *Journal of Geriatric Physical Therapy*.
10.1519/JPT.0000000000000258
Heywood S, McClelland J, Geigle P, Rahmann A, Villalta E, Mentiplay B, and Clark R. 2019. Force during
functional exercises on land and in water in older adults with and without knee osteoarthritis: Implications
for rehabilitation. *The Knee* 26:61-72. https://doi.org/10.1016/j.knee.2018.11.003

Holmstrup ME, Jensen BT, Evans WS, and Marshall EC. 2016. Eight Weeks of Kettlebell Swing Training Does not
Improve Sprint Performance in Recreationally Active Females. *Int J Exerc Sci* 9:437-444.
Hopkins WG, Marshall SW, Batterham AM, and Hanin J. 2009. Progressive statistics for studies in sports medicine
and exercise science. *Med Sci Sports Exerc* 41:3-13. 10.1249/MSS.0b013e31818cb278
Janssen I, Baumgartner RN, Ross R, Rosenberg IH, and Roubenoff R. 2004. Skeletal Muscle Cutpoints Associated
with Elevated Physical Disability Risk in Older Men and Women. *American Journal of Epidemiology*
159:413-421. 10.1093/aje/kwh058
Kera T, Kawai H, Takahashi J, Hirano H, Watanabe Y, Fujiwara Y, Ihara K, Kim H, and Obuchi S. 2020.
Association between ground reaction force in sit-to-stand motion and falls in community-dwelling older
Japanese individuals. *Archives of Gerontology and Geriatrics* 91:104221. 10.1016/j.archger.2020.104221
Khan KM, and Scott A. 2009. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue
repair. *British Journal of Sports Medicine* 43:247. 10.1136/bjsm.2008.054239
Lake JP, Hetzler BS, and Lauder MA. 2014. Magnitude and relative distribution of kettlebell snatch force-time
characteristics. *Journal of Strength and Conditioning Research* 28:3063-3072.
10.1519/jsc.0000000000000538
Lake JP, and Lauder MA. 2012a. Kettlebell swing training improves maximal and explosive strength. *Journal of*
Strength and Conditioning Research 26:2228-2233. 10.1519/JSC.0b013e31825c2c9b
Lake JP, and Lauder MA. 2012b. Mechanical demands of kettlebell swing exercise. *Journal of Strength and*
Conditioning Research 26:3209-3216. 10.1519/JSC.0b013e3182474280
LaRoche DP, Millett ED, and Kralian RJ. 2011. Low strength is related to diminished ground reaction forces and
walking performance in older women. *Gait and Posture* 33:668-672. 10.1016/j.gaitpost.2011.02.022
Lenhard W, and Lenhard A. 2016. Calculation of Effect Sizes. *Available at*
*http://www.psychometrica.de/effect_size.html*2020).
Levine NA, Hasan MB, Avalos MA, Lee S, Rigby BR, and Kwon Y-H. 2020. Effects of kettlebell mass on lower-
body joint kinetics during a kettlebell swing exercise. *Sports Biomechanics*:1-14.
10.1080/14763141.2020.1726442
Lim J, Kim M, Kim J, Hong J, Lee D, and Yu J. 2018a. Core and Lower limb Activations of the two-handed and
Single-Handed Kettlebell Exercises. *Research Journal of Pharmacy and Technology* 11:1101-1106.
Lim JY, Yi Y, Jung SW, and Park D-S. 2018b. Comparison of vertical ground reaction forces between female
elderly and young adults during sit-to-stand and gait using the Nintendo Wii Balance Board. *Physical*
Therapy Rehabilitation Science 7:179-185.
Lopez P, Pinto RS, Radaelli R, Rech A, Grazioli R, Izquierdo M, and Cadore EL. 2017. Benefits of resistance
training in physically frail elderly: a systematic review. *Aging Clinical and Experimental Research*:1-11.
10.1007/s40520-017-0863-z
Mache MA, and Hsieh C. 2016. A Temporal and Kinetic Comparison of The Kettlebell Swing and Maximal
Vertical Jump. ISBS-Conference Proceedings Archive.
Maffiuletti NA, Aagaard P, Blazevich AJ, Folland J, Tillin N, and Duchateau J. 2016. Rate of force development:
physiological and methodological considerations. *European Journal of Applied Physiology* 116:1091-1116.
10.1007/s00421-016-3346-6
Magnusson K. 2021. Interpreting Cohen's d effect size: An interactive visualization (Version 2.5.0) [Web App].
*Available at https://rpsychologist.com/cohend/*2021).
Manocchia P, Spierer DK, Lufkin AK, Minichiello J, and Castro J. 2013. Transference of kettlebell training to
strength, power, and endurance. *Journal of Strength and Conditioning Research* 27:477-484.
10.1519/JSC.0b013e31825770fe
Marcelino C. 2017. Effects of kettlebell training on functional performance, postural stability and isokinetic strength
of lower limbs in individuals with Parkinson's disease [masters]. *Brazil: Univeristy of Brazil*.
McGill SM, and Marshall LW. 2012. Kettlebell swing, snatch, and bottoms-up carry: back and hip muscle
activation, motion, and low back loads. *Journal of Strength and Conditioning Research* 26:16-27.
10.1519/JSC.0b013e31823a4063
Meigh NJ, Hing WA, Schram B, and Keogh JWL. 2021. Mechanical demands of the two-handed hardstyle kettlebell
swing performed by an RKC-certified Instructor. *bioRxiv*:2021.2005.2013.444085.
10.1101/2021.05.13.444085
Meigh NJ, Keogh JWL, Schram B, and Hing WA. 2019. Kettlebell training in clinical practice: a scoping review.
BMC Sports Science, Medicine and Rehabilitation 11:19. 10.1186/s13102-019-0130-z
Mitchell J, Johnson W, Riemann B, Krajewski K, and Coates C. 2015. *Biomechanical Loading of the American*
Kettlebell Swing.

Oikarinen S. 2016. American Kettlebell Swing and the Risk of Lumbar Spine Injury.
Otto WH, 3rd, Coburn JW, Brown LE, and Spiering BA. 2012. Effects of weightlifting vs. kettlebell training on
vertical jump, strength, and body composition. *Journal of Strength and Conditioning Research* 26:1199-
1202. 10.1519/JSC.0b013e31824f233e
Pasco JA, Stuart AL, Holloway-Kew KL, Tembo MC, Sui SX, Anderson KB, Hyde NK, Williams LJ, and
Kotowicz MA. 2020. Lower-limb muscle strength: normative data from an observational population-based
study. *BMC Musculoskeletal Disorders* 21:89. 10.1186/s12891-020-3098-7
Rajala BA, and Jensen RL. 2015. EMG of lower limb muscles during kettlebell exercises. ISBS-Conference
Proceedings Archive.
Ros T. 2016. Comparison of kettlebell jump squats vs kettlebell swings as postactivation potentiation on vertical
jumps. California State University, Fullerton.
Ross JA, Keogh JW, Wilson CJ, and Lorenzen C. 2017. External kinetics of the kettlebell snatch in amateur lifters.
PeerJ 5:e3111. 10.7717/peerj.3111
Santillo N. 2016. The Comparative Effect of Recovery Interventions on Blood Lactate Removal Post-High Intensity
Kettlebell Training Session. The William Paterson University of New Jersey.
Schnettler C, Porcari J, Foster C, and Anders M. 2009. Physiologic responses to a kettlebell workout. University of
Wisconsin–La Crosse.
Shin G, Nance M, and Mirka G. 2006. Differences in trunk kinematics and ground reaction forces between older and
younger adults during lifting. *International Journal of Industrial Ergonomics* 36:767-772.
10.1016/j.ergon.2006.05.008
Smidt N, de Vet HCW, Bouter LM, and Dekker J. 2005. Effectiveness of exercise therapy: A best-evidence
summary of systematic reviews. *Australian Journal of Physiotherapy* 51:71-85. 10.1016/S0004-
9514(05)70036-2
Smith CE, Lyons B, and Hannon JC. 2014. A Pilot Study Involving the Effect of Two Different Complex Training
Protocols on Lower Body Power. *Human Movement* 15. 10.1515/humo-2015-0003
Soukkio P, Suikkanen S, Kääriä S, Kautiainen H, Sipilä S, Kukkonen-Harjula K, and Hupli M. 2018. Effects of 12-
month home-based physiotherapy on duration of living at home and functional capacity among older
persons with signs of frailty or with a recent hip fracture - protocol of a randomized controlled trial
(HIPFRA study). *BMC Geriatrics* 18:232-232. 10.1186/s12877-018-0916-y
Teichtahl AJ, Wluka AE, Wijethilake P, Wang Y, Ghasem-Zadeh A, and Cicuttini FM. 2015. Wolff's law in action:
a mechanism for early knee osteoarthritis. *Arthritis research & therapy* 17:207. 10.1186/s13075-015-0738-
7
Thomas JF. 2014. *Comparison of interval kettlebell exercise and cycle ergometry: Effectiveness as a stimulus for
cardiorespiratory fitness*: Southeastern Louisiana University.
Thomas JF, Larson KL, Hollander DB, and Kraemer RR. 2014. Comparison of two-hand kettlebell exercise and
graded treadmill walking: effectiveness as a stimulus for cardiorespiratory fitness. *Journal of Strength and
Conditioning Research* 28:998-1006. 10.1519/jsc.0000000000000345
Tsatsouline P. 2006. *Enter The Kettlebell!: Strength Secret of the Soviet Supermen*. St. Paul, MN, USA: Dragon
Door Publications.
Tsatsouline P. 2013. *RKC Instructor Manual*. St. Paul, MN, USA: Dragon Door Publications Inc.
Tsuji T, Tsunoda K, Mitsuishi Y, and Okura T. 2015. Ground Reaction Force in Sit-to-stand Movement Reflects
Lower Limb Muscle Strength and Power in Community-dwelling Older Adults. *International Journal of
Gerontology* 9:111-118. 10.1016/j.ijge.2015.05.009
Turner MN, Hernandez DO, Cade W, Emerson CP, Reynolds JM, and Best TM. 2019. The Role of Resistance
Training Dosing on Pain and Physical Function in Individuals With Knee Osteoarthritis: A Systematic
Review. *Sports health* 12:200-206. 10.1177/1941738119887183
Wesley C. 2013. Effects of kettlebell mass & swing cadence on heart rate, blood lactate and ratings of perceived
exertion during an interval kettlebell swing protocol. ProQuest Dissertations Publishing.
Wong A, Nordvall M, Walters-Edwards M, Lastova K, Francavillo G, Tripken J, and Figueroa A. 2017. Cardiac
Autonomic and Blood Pressure Responses to an Acute Bout of Kettlebell Exercise. *Medicine and Science
in Sports and Exercise* 49:721-721. 10.1249/01.mss.0000518918.69080.ab
Zin MAM, Rambely AS, and Ariff NM. 2018. A preliminary study on effects of increment of loads to lower
extremity joints during kettlebell swing activity. AIP Conference Proceedings: AIP Publishing LLC. p
020120.
702