Targeted redesign of suramin analogs for novel antimicrobial lead development

Debayan Dey[1#], Suryanarayanarao Ramakumar[2], Graeme L. Conn[1,3#]

[1]Department of Biochemistry, Emory University School of Medicine, Atlanta, GA 30322, USA

[2]Department of Physics, Indian Institute of Science, Bangalore-560012, India

[3]Emory Antibiotic Resistance Center (ARC).

[#]**Corresponding author:** Debayan Dey, email: debayan.dey@emory.edu or Graeme L. Conn, email: gconn@emory.edu

Running title: Suramin analogs as antimicrobial leads

Keywords: Antiviral, antimicrobial, cheminformatics, virtual screening, precision docking, drug design strategies.

Author Contributions: DD, SR and GLC conceptualized the project. DD and GLC wrote the draft. All authors have given approval to the final version of the manuscript.

ABSTRACT

The emergence of new viral infections and drug resistant bacteria urgently necessitates expedient therapeutic development. Repurposing and redesign of existing drugs against different targets is one potential way in which to accelerate this process. Suramin was initially developed as a successful anti-parasitic drug, but has also shown promising antiviral and antibacterial activities. However, due to its high conformational flexibility and negative charge, suramin is considered quite promiscuous towards positively charged sites within nucleic acid binding proteins. Although some suramin analogs have been developed against specific targets, only limited structure activity relationship (SAR) studies were performed, and virtual screening has yet to be used to identify more specific inhibitor(s) based on its scaffold. Using available structures, we investigated suramin's target diversity, confirming that suramin preferentially binds to protein pockets which are both positively charged and enriched in aromatic or leucine residues. Further, suramin's high conformational flexibility allows adaptation to structurally diverse binding surfaces. From this platform, we developed a framework for structure- and docking-guided elaboration of suramin analog scaffolds using virtual screening of suramin and heparin analogs against a panel of diverse therapeutically relevant viral and bacterial protein targets. Use of this new framework to design potentially specific suramin analogs is exemplified using the SARS-CoV-2 RNA-dependent RNA polymerase (RdRp) and nucleocapsid protein, identifying leads that might inhibit a wide range of coronaviruses. The approach presented here establishes a new computational framework for designing suramin analogs against different bacterial and viral targets and repurposing existing drugs for more specific inhibitory activity.

INTRODUCTION

The rise of multi-drug resistant bacteria and emergence of new infectious viral pathogens, as exemplified by SARS-CoV-2 the cause of the COVID-19 pandemic, are dual major threats to human health world-wide. Effective new antimicrobials are thus urgently needed, but processes for their identification and development are often long and expensive. However, repurposing of existing, approved drugs and redesign of their activity against new targets represents one of the most promising opportunities to by-pass many of these hurdles (1).

The unique challenges posed by COVID-19 have resulted in testing of a large array of antivirals, anti-inflammatory drugs, and other potential inhibitors of receptors of viral attachment for their effectiveness against SARS-CoV-2. For example, the nucleoside analog remdesivir, originally developed for Ebola virus, was found to be effective against SARS-CoV-2 with high selectivity, leading to its use for treatment of COVID-19 in many countries (2). Other antivirals like favipiravir, ribavirin, darunavir, and lopinavir-ritonavir combination are either already in use or in clinical trials for SARS-CoV-2 treatment (3). In addition, non-antivirals drugs like dexamethasone, which is used commonly to treat inflammatory and autoimmune conditions (4), and the anti-parasitic ivermectin have shown potential as COVID-19 treatments (5, 6).

Suramin was among the first anti-infective agents developed in the early 1920s and has also recently been shown to inhibit the progression of SARS-CoV-2 infection in human airway epithelial cell culture model (7). Suramin appears on the World Health Organization's List of Essential Medicines and has a long history of clinical use, with application over the last century in treating African Trypanosomiasis (sleeping sickness) caused by *Trypanosoma brucei* (8), and other parasitic infections including onchocerciasis (African river blindness), leishmaniasis and malaria (9-11). With its long history of successful use as an anti-parasitic, suramin and its analogs

45 have also found use in wide variety of other treatments, such as anti-cancer agents, antivirals, and

46 venom antidotes (12).

47   Suramin was found to exhibit broad antiviral properties through distinct activities against

48 components of diverse viruses. For example, through binding to glycoprotein gp120 suramin

49 inhibits attachment of HIV to host T-cells (13). Suramin also limits host cell entry of several other

50 viruses, including herpes, dengue and hepatitis C virus (14-16) and also inhibits enteroviral

51 attachment to human host cells by binding to their nucleocapsid protein (17). Suramin also inhibits

52 some processes in viral replication, for example through binding to Zika virus NS2B/NS3

53 proteinase (18) or to the RNA-dependent RNA polymerase (RdRp) of norovirus, chikungunya

54 virus and SARS-CoV-2 (19-21). Suramin and related compounds thus hold promise as an initial

55 leads for multiple targets, including viral RdRp and nucleocapsid, as well as future structure-

56 activity relationship (SAR) studies or structure-guided design of suramin analogs as new

57 therapeutics.

58   Although suramin and its analogs have been studied for many decades, including some

59 SAR studies (22, 23), many fundamental questions remain regarding the origin of their

60 promiscuity, binding pocket preferences, and mode of on-target interaction. Suramin's

61 effectiveness in targeting viral nucleocapsid and RdRp has not yet been exploited to rationally

62 design more selective and less toxic analogs based on this scaffold. Moreover, to our knowledge,

63 no systematic high-throughput or virtual screening has been performed for suramin analogs for

64 any viral targets (e.g. RdRp or nucleocapsid) to explore potential lead molecules. Here, we

65 examine the structural basis of suramin's ability to bind to diverse proteins and use these insights

66 to develop a framework involving virtual screening, precision docking and docking-guided

67 elaboration of the suramin scaffold against different therapeutically important bacterial and viral

68 targets. Docking studies with a panel of nine diverse proteins reveals high-scoring suramin analogs

69 to be more conformationally selective than suramin and to have greater target specificity. Finally,

70 SARS-CoV-2 RNA-dependent RNA polymerase (RdRp) and nucleocapsid are used to exemplify

71 the process of suramin analog redesign to generate novel analogs with the potential to bind with

72 higher affinity and target specificity. Evolutionary analyses suggest that the optimized leads could

73 potentially inhibit a wide range of coronaviruses (24). This study thus sets out a computational

74 framework using structure- and docking-guided ligand elaboration that can support redesign and

75 repurposing existing antimicrobials for new, specific inhibitory activities.

76

77 **RESULTS**

78 **Structural basis for suramin protein target site diversity**

79 Suramin is a C2 symmetric molecule with each half comprised of a trisulfonated naphthyl ring

80 with negative charge at physiological pH, connected via amide linkages to toluene and phenyl

81 rings (**Fig. 1A**). Suramin is highly flexible with a total of 10 rotatable bonds (**Fig. 1A**; α_{1-5} and $\alpha_{1'-}$

82 $_{5'}$) and no substituent groups on the four central rings that might otherwise restrict its

83 conformational freedom. This flexibility, and thus conformational adaptability, is likely a key

84 factor in suramin's ability to interact with diverse target sites in many proteins, but also its high

85 off-target binding. Similarly, another polyanionic class of molecules similar to suramin, the

86 glycosaminoglycans (GAGs) such as heparin (**Fig. 1B**), can bind multiple diverse protein targets

87 but also suffer the limitation of high off-target binding.

88　　　　To begin defining the residue preferences in suramin binding pockets, we examined

89 structures of several suramin complexes with diverse proteins (**Table S1**). These structures

90 included the *Leishmania mexicana* pyruvate kinase (PDB 3PP7**; Fig. 1C**) with suramin bound in

its ATP binding site, snake venom protein myotoxin (PDB 1Y4L; **Fig. 1D**), and two human protein

targets for treatment of cancer, metabolic and neurological diseases, the NAD$^+$-dependent protein

deacetylase sirtuin 5 (SIRT5; PDB 2NYR; **Fig. 1E**) and the epigenetic reader chromobox homolog

7 (PDB 4X3U; **Fig. 1F**) (10, 25-27). A number of general features of the interaction networks

within the suramin binding pockets are immediately apparent. The trisulfo-naphthyl ring interacts

with positively charged residues (Arg, Lys, His) via its sulfate groups while the naphthyl ring

makes π-mediated interactions with aromatic residues, Phe, His, Tyr and Trp (**Fig. 1C-F**).

Aromatic residues with a polar ring atom may also make a hydrogen bonding interaction with a

sulfonate group, coordinating both features of the trisufonated naphthyl ring. The toluene and

central phenyl rings of suramin are additionally stabilized by networks of aromatic and

hydrophobic residues (**Fig. 1C-F**).

Suramin complexes with viral proteins have also been determined for norovirus RdRp

(PDB 3UR0) (19) and bunyavirus nucleocapsid (PDB 4J4V) (28), but these offer only partial

snapshots of favored suramin binding sites as just a portion of suramin could be modeled in each

crystal structure (**Fig. S1**). In the RNA binding channel of norovirus RdRp34, the trisulfo-naphthyl

ring of suramin is anchored by both positive (Lys171 and Arg392) and aromatic (Trp42) residues,

while the central region is stabilized by Gln66, Lys180, Lys181, Arg245 and Lys68. The

bunyavirus nucleocapsid forms a pentameric complex with an RNA binding cavity on the inner

edge of the ring-like assembly where suramin also interacts (28). Again, each of the modeled

suramin rings is stabilized by multiple ionic, hydrogen bonding and aromatic/ hydrophobic

stacking interactions, including residues Asn66, Lys67 and Arg95 with the trisulfo-naphthyl ring,

and Arg64, Thr63, Gly65, Phe177 and Pro127 the two central rings (**Fig. S1B**). Thus, visual

inspection of these viral protein structures reveals overall consistent trends in suramin's binding

pocket residue interaction preferences with other proteins of diverse origin and function.

More detailed analyses of available suramin-bound protein structures reveal consistent

features in the binding pocket physicochemical composition, including strong enrichment of

specific residues (**Fig. 1G**). Positively charged Lys and Arg are highly enriched in the pocket

compared to the full protein (25% and 15%, respectively, of all interacting residues), while His is

more modestly enriched (5%), but, as noted, can make both aromatic stacking and charged

interactions. The absence of negatively charged residues (~1.5% combined) also points towards

suramin's strong preference for positively charged binding regions, such as nucleic acid binding

sites. In total, polar residues are slightly less favored (~12 % total) as compared to aliphatic

hydrophobic residues (~14%), with a marked preference for Leu (10.6 %) among the latter group

(**Fig. 1G**). This may be due to leucine's aliphatic stacking potential with the aromatic rings of

suramin. Aromatic residues (Trp, Tyr and Phe; together ~25% of total) are also strongly enriched

in suramin binding sites.

Suramin's binding promiscuity for diverse protein targets is likely underpinned by its high

inherent conformational flexibility, as revealed by superposition of protein-bound suramin

molecules using one trisulfo-naphthyl ring (**Fig. 1H**). Suramin bound to thrombin (PDB 3BF6)

adopts the most extended conformation we identified (and is used hereafter as the reference), with

the trisulfo-naphthyl rings ~29 Å apart. Suramin bound to SIRT5 (PDB 2NYR) also has a linear

conformation but with differences in the α2, α3, α2', and α3' angles (i.e. the dihedrals between the

toluene and phenyl rings). In contrast, suramin bound to myotoxin I (PDB 6CE2) has its trisulfo-

naphthyl rings only ~14 Å apart and is observed in a highly bent conformation. Similarly, suramin

bound to myotoxin II (PDB 1Y4L) adopts a highly bent conformation, but distinguished by

additional changes in the α1 and α1' dihedral angles. To further assess the low energy

conformations of suramin and two analogs, NF449 and NF023, we used a computational *Monte*

Carlo conformational scan which revealed 116 potential low energy conformational states for

suramin, while NF449 and NF023 each have 143 and 80, respectively.

Collectively, these structural and computational conformation analyses confirm suramin to

be a highly flexible ligand, with multiple potential hinge points, that can adopt a large variety of

conformations to fit a given target protein binding pocket. Further, these favored binding pockets

are positively charged and enriched with aromatic and Leu residues, such as most commonly occur

in large interfaces or in protein regions which bind to nucleic acids or nucleotides.

146 Identification of suramin and heparin analogs targeting viral and bacterial proteins

Previous studies have identified inhibitory activity of suramin against bacterial and viral proteins,

but no computational or high-throughput screens have been conducted with suramin analogs

against such targets. Structure-guided analysis of suramin analogs bound to various viral or

bacterial targets using large-scale docking studies could, however, allow for downstream rational

design of novel analogs with improved affinity and specificity towards a desired target. We devised

a strategy to accomplish this goal using docking studies with suramin and heparin analogs and

using docking poses of the latter set of ligands to guide design of new suramin analogs with

substituents positioned to make additional protein target-specific interactions (**Fig. 2**).

Nucleic acid binding proteins in bacteria and viruses play crucial roles in the processing of

genetic information and thus represent important targets for therapeutic development. Due to

suramin's affinity for nucleic acid binding proteins, a test panel of diverse protein targets known

to interact with DNA or RNA was selected. Viral proteins, known to be bound by suramin,

included RdRp of Norovirus and SARS-CoV-2, which is involved in replication and transcription

160 of the viral RNA genome (19, 21); nucleocapsid of bunyavirus and SARS-CoV-2, which is

161 involved in packaging the viral RNA genome and virion assembly, and thus of critical importance

162 in viral infection (17, 28, 29); and, human immunodeficiency virus type 1 (HIV-1) protein gp120,

163 for which binding to galactosylceramide on human colon epithelial cells is blocked by suramin

164 (13, 30). To further diversify the protein target set, we included four bacterial proteins: DNA-

165 binding protein HU (HU), DNA recombination protein RecA (RecA), DNA-binding protein Fis

166 (Fis), and the thiostrepton-resistance methyltransferase (TsnR). HU and Fis are global regulators

167 which play important roles in bacterial gene regulation, biofilm development and maintaining

168 nucleoid architecture (31-34), while RecA is an essential DNA recombination and repair protein

169 known to be targeted by suramin (35). Finally, TsnR is a ribosomal RNA methyltransferase that

170 confers resistance to the antibiotic thiostrepton (36), and represents a wider group of RNA

171 modifying enzymes that are promising targets for novel drugs to break antibiotic resistance. To

172 generate the ligand sets, we next used a small panel of known biosimilars (**Table S2**) in a PubChem

173 search which identified ~2,000 and ~10,000 analogs of suramin and heparin, respectively (**Fig.**

174 **2A**).

175   Using the protein target panel and suramin/ heparin analog sets, high-throughput virtual

176 screening (HTVS; **Fig. 2B**) in the GLIDE module of the Schrödinger software produced a wide

177 range of binding scores, typically ranging from −1.0 to −9.0 kcal/mol, but with large differences

178 in distribution and average score for each protein target (**Fig. 3A**,**B**). Among the bacterial proteins,

179 suramin analogs bind on average more tightly to Fis and TsnR (mean binding score -5.7 and -5.9

180 kcal/mol, respectively) compared to HU and RecA (-5.1 and -4.6 kcal/mol, respectively; **Fig 3A**).

181 HIV protein gp120 was predicted to be bound less tightly with a mean score of -3.6 kcal/mol,

whereas both suramin and heparin analogs docked viral RdRp with comparatively higher score

(**Fig. 3A,B**).

Among the top 100 scoring suramin and heparin analogs identified by HTVS, more than

~50% of these ligands bind to only a single protein among the nine targets (**Fig. 3C**,**D**).

Approximately 30% of the remaining ligands in each set bind to only two or three receptors, with

just a small fraction showing greater promiscuity (high docking score for 4 or more targets). This

result indicates that virtual screening-based using suramin or heparin analogs can identify ligands

with greater selectivity for diverse protein targets compared to using either suramin or heparin

individually.

The top 100 suramin and heparin analogs from HTVS against each target protein were next

subject to more accurate docking pose and binding score calculation using Standard Precision (SP)

GLIDE (**Fig. 2B** and **Fig. 3E**,**F**). An improvement in the docking score was observed for many

analogs due to the ligand conformational sampling used in the SP GLIDE docking method as

compared to "rigid" ligand docking of HTVS. However, SP GLIDE docking had a very similar

trend in docking score distribution for each set of top 100 analogs (**Fig. 3E**,**F**). Heparin analogs

bind modestly better than those of suramin, likely due to their greater overall conformational

flexibility and number of hydrogen bond donors/acceptors, which allow them more readily adapt

to the wide variety of positively charged surfaces in these target proteins. The top scoring ligands

from SP docking (**Tables S3** and **S4**, and Supplementary File "Top 100 docking results") were

then selected for subsequent elaboration of the suramin scaffold and are hereafter named beginning

with "Sur" and "Hep" for suramin and heparin analogs, respectively.

Strategy for docking-guided elaboration of the suramin scaffold

To facilitate suramin analog lead redesign for increased target selectivity and affinity, the top

scoring suramin analogs for each protein target were visually inspected post-docking and then

superposed with the top scoring heparin analog identified in the same binding site. The heparin

analog set contains ligands that are typically more conformationally flexible and enriched in

potential hydrogen bond donors/ acceptors. While these features would make the heparin analogs

themselves poor leads for specific interactors, comparison of the bound analogs can direct

identification of sites within the corresponding suramin analog that can be further substituted to

make additional interactions with the target protein (**Fig. 2C**). As an example, structural analysis

of the top suramin and heparin analogs directed against HU and the resulting suramin analog

redesign strategy is described below.

HU interacts with DNA as a dimer via a positively charged binding cleft, lined with Lys

and Arg residues from both protomers. Sur-1 binds to HU in this cleft with the highest affinity of

the suramin analog set (-9.66 kcal/mol), placing two of its naphthyl rings over a surface formed by

Pro81 and the adjacent β-strand of both HU protomers at the center of the dimer interface (**Fig.**

4A). Sur-1 is composed of three sulfo-naphthyl rings (**Fig. 4B**), with both terminal rings substituted

with two sulfonyl groups, and a single sulfonate on the central ring. The majority of these groups

are coordinated by multiple Arg residues, including Arg53, Arg55 and Arg80 from both protomers,

and Arg58 and Lys 86 from a single protomer (**Fig. 4A**). Lys86 of the second protomer is also

positioned near (~4-6 Å) the carbonyl and sulfonyl groups of the central and terminal naphthyl

rings, respectively, exemplifying how the dimeric interface may be exploited by a pseudo-

symmetrical suramin analog. Additionally, Sur-1 and other top HU-binding suramin analogs (Sur-

2 and Sur-3; **Table S3**) are all markedly more conformationally rigid than suramin, due to the

diazene linker between the naphthyl rings. Sur-1 has only four rotatable bonds with a single low

228 energy conformation (**Fig. 4B**), compared to 10 rotatable bonds and 116 low energy conformations

229 in suramin. The ligand docking process thus identified suramin analogs with both improved

230 binding affinity to HU (based on docking score) and reduced conformational rigidity and net

231 charge. These features make Sur-1 a superior candidate compared to suramin for developing a

232 more specific inhibitor of HU.

233 Our docking procedure with the heparin analog set also identified Hep-41 as having high

234 affinity (-10.02 kcal/mol) for the same pocket as Sur-1 in the DNA binding cleft of HU (**Fig. 4C**).

235 Hep-41 is composed of sulfated iduronic acid and glucosamine with trioxidanylsulfanyl

236 substitution, which make extensive interactions with essentially the same group of residues in the

237 DNA binding cleft (i.e. Arg53, Arg55 Arg58, Arg80 and Lys86; **Fig. 4B**). However, while docking

238 of Sur-1 and Hep-41 are qualitatively similar, due to its larger number of hydrogen bond donors/

239 acceptors, docking of the latter ligand offers detailed structure-guided cues for placement of new

240 substituents in Sur-1. Specifically, superposition of the two analogs reveals sites on Sur-1 for

241 optimal geometric placement of additional polar groups to enhance interaction within the binding

242 site (**Fig. 4A,D**). For example, addition of polar groups on each naphthyl ring could increase target-

243 ligand interactions by further engaging Arg53 and Arg55 (from both protomers) via bifurcated

244 hydrogen bonding. These dual ligand docking comparisons can thus be used as a guide to improve

245 the design of suramin analogs to increase both their target specificity and binding affinity.

246 This same strategy was used for the other eight target proteins, with heparin analog

247 superposition identifying multiple sites in each top suramin analog where substitutions could be

248 made to improve affinity and target selectivity (marked * in Fig. S2-S5; also see Supplemental

249 Results for further details). Specifically, we identified sites for targeted redesign of Sur-6 (using

250 Hep-42) for binding to the positively charged cleft of TsnR in an extended conformation (**Fig. S2**

A-D), and Sur-13 (using Hep-45) and Sur-15 (using Hep-50) for binding with higher affinity and

specificity to Fis (**Fig. S2 E-H**) and RecA (**Fig. S3 A-D**), respectively. Among the viral proteins,

HIV gp120 (6IEQ) binds to smaller analogs than the other targets with the highest docking score

to Sur-22 and sites for targeted redesign identified by docking of Hep-51 (**Fig. S3E-H**). Similarly,

for norovirus RdRp (**Fig. S4A-D**) and bunyavirus nucleocapsid (**Fig. S4E, F**) the docking

identified Sur-24/ Hep-56 and Sur-27/ Hep-52 binding poses, respectively, for targeted redesign.

Corresponding results for SARS-CoV-2 RdRp and nucleocapsid are described in the next section

along with subsequent redesign and further docking analyses. In summary, virtual screening and

precision docking analyses using suramin and heparin analog chemical space has discovered new

lead analogs and a path to rational redesign for specific viral and bacterial protein targets.

Suramin analog redesign for specific, high-affinity lead inhibitors of SARS-CoV-2 RdRp and

nucleocapsid

SARS-CoV-2 RdRp is a large macromolecular complex composed of nsp12 catalytic subunit, with

an RNA binding tunnel that presents a potential site for drug targeting (**Fig. S5A,B**), and nsp7-

nsp8 cofactors (37). Virtual screening and docking analyses identified Sur-36 as binding with

highest docking score to SARS-CoV-2 RdRp, with Hep-58 also binding in the same region (**Fig.**

S5C). Sur-36 has a central biphenyl ring attached to terminal sulfo-naphthyl groups joined by

aconformationally restricted diazene linker (**Fig. 5B**). Two other similarly high-scoring ligands,

Sur-37 and Sur-38, were also identified (**Table S3**), but Sur-36 was selected as the lead candidate

due to the greater flexibility of these additional hits. Inspection of the RdRp binding pocket reveals

that each ring of Sur-36is stabilized by multiple interactions: the distal sulfo-naphthyl ring by

Lys551, Ser549 and Arg553, the central phenyl rings by Ile548 and Ala549, and the proximal

274 sulfo-naphthyl ring with Tyr546 and Asn497 (**Fig. 5A, B**).

275   Using the Hep-58 binding pose, four positions for additional substitutions in Sur-36 were

276 identified as described above for the other ligand/ protein target pairs (**Fig. S5D**). A process of

277 chemical enumeration, structure-guided design and iterative docking was then used to

278 systematically alter the functional groups of substituents at the identified attachment positions and

279 calculate the changes in the docking score (**Table S5**). The methoxy groups in the central phenyl

280 rings were first modified to carboxylate, phenyl-methanol, halides and other functional groups,

281 with docking analysis identifying the carboxylate groups (Sur-60; **Fig. S6A**) as conferring the

282 greatest increase in docking score, from -10.14 (Sur-36) to -11.24 kcal/mol (Sur-60). In Sur-60,

283 one phenyl ring carboxyl group engages Lys545, while the second carboxylate makes new contacts

284 with Arg555. Additional docking-guided modifications were then made in Sur-60 by

285 systematically testing additional functional groups in the terminal naphthyl rings and the linker

286 between the central phenyl rings. This optimization process resulted in substitution of one terminal

287 naphthyl group with a phenalene group, addition of a hydroxyl group to the remaining sulfo-

288 naphthyl group, and a carbonyl linker between the central rings to generate Sur-84 (-11.60

289 kcal/mol; **Fig. S6A**). Sur-84 makes additional contacts to His439 and Arg836 via one of the

290 carboxylate groups and a hydrophobic interaction between the phenalene group and Val557.

291 Finally, conversion of a central phenyl ring to naphthyl added hydrophobic contact with Tyr546

292 and generated Sur-85 (-11.96 kcal/mol; **Fig. 5C** and **S6A**).

293   The same process was also applied to the SARS-CoV-2 nucleocapsid RNA binding domain

294 (29) which has a shallow positively charged surface (**Fig. S5E-G**). Smaller suramin analogs with

295 phenyl, stilbene and naphthyl moieties typically docked with higher scores than longer suramin

296 analogs, with Sur-31 among the highest docking scores (-7.53 kcal/mol) and selected for further

297   redesign to improve its affinity. Sur-31 is composed of two terminal naphthyl groups with a central

298   unsubstituted phenyl ring linked connected by two conformationally restricted amide linkers (**Fig.**

299   **S5H**). Inspection of the RNA binding pocket reveals Sur-31 is stabilized by multiple interactions:

300   one sulfo-naphthyl ring with Arg93, Arg108 and Arg150, the central phenyl rings with Tyr110

301   and Tyr112, and the second sulfo-naphthyl ring with Tyr112, Thr92 and Arg89 (**Fig. 5D, E**). Our

302   heparin screen also identified Hep-59 as binding to the same region, revealing potential for

303   extending the Sur-31 scaffold at several sites (indicated * in **Fig. S5H**). Again using docking-

304   guided elaboration, one naphthyl ring was replaced by two sulfonyl-substituted phenyl rings

305   connected to the central phenyl ring via a conformationally restricted azide linker resulting in Sur-

306   114 with an improved of docking score of -8.79 kcal/mol (**Fig. S6B**). Sur-114 makes additional

307   interactions with backbone carbonyls of Ala157 and Ile158, along with an increased van der Waals

308   interaction with the protein. Further guided substitutions in the second naphthyl ring, adding

309   another disulfonyl-hydroxy naphthyl ring linked by a flexible linker, generated Sur-121 with a

310   further improvement in docking score to -9.30 kcal/mol (**Fig. S6B**). Final optimization of this

311   naphthyl ring with a sulfate group resulted in Sur-127, with a final docking score of -9.34 kcal/mol.

312   Sur-127 engages the overall same set of residues but with increased H-bonding and electrostatic

313   interactions compared to Sur-31 (**Fig. 5F** and **S6B**).

314   This process thus identified an initial suramin analog for both SARS-CoV-2 protein targets,

315   RdRp (Sur-36) and nucleocapsid (Sur-31), for which predicted affinity could be substantially

316   improved by heparin docking-guided redesign efforts, resulting in final analogs Sur-85 and Sur-

317   127, respectively. Overall, heparin docking- and structure-guided redesign of suramin analogs

318   resulted in a collection of potential leads with markedly improved docking scores and other

properties, compared to analogs identified in precision docking (compare green and purple scores,

respectively in **Fig. 5G**).

Finally, we investigated residue conservation to gain insights into the evolutionary

constrains on the potential inhibitor binding sites in SARS-CoV-2 RdRp and nucleocapsid

proteins. Across all coronaviruses the identified binding sites appear moderately and very highly

conserved in nucleocapsid and RdRp, respectively (**Fig. 6**). As such, the phylogenetic relationship

reveals the high similarity of both SARS-CoV-2 proteins to homologs in other β-coronavirus

family members, comprising predominantly bat-origin viruses such as MERS-CoV and HCoV-

HKU1 (**Fig. 6A, B; Fig S7**). The conservation of the RdRp binding site also extends to α-

coronavirus family members of bat origin, which originated the human pathogens HCoV-229E

and HCoV-NL63 (**Fig. 6A, B**). More broadly, the generally strong conservation in each binding

site implies that the designed suramin analogs, Sur-85 and Sur-127, would potentially be equally

effective in targeting these proteins in different coronaviruses. Thus, together the ligand docking/

elaboration strategies and evolutionary analyses provide a basis for designing specific inhibitors

with broad anti-coronaviral activity.

DISCUSSION

Suramin has garnered wide clinical interest for treating parasitic infections, as well as diseases

ranging from viral infections (e.g. HIV, Dengue and Ebola) to human cancers (12). Despite not

satisfying typical drug likeliness criteria, such as Lipinski's "rule of five", suramin and its analogs

have found success in clinical application. Suramin is negatively charged and binds predominantly

positively charged surfaces in proteins involved in DNA and RNA processing, e.g., DNA and RNA

polymerases, telomerase and chromodomain proteins, histone methyltransferases, and sirtuin

histone deacetylases (19, 21, 23, 27). Suramin also inhibits several membrane channels and

signaling proteins. Despite suramin having been used for more than 100 years and shown to inhibit

diverse protein families, there still exists a significant gap in our understanding of its promiscuous

antagonist properties which likely also underpin its off-target and side effects in clinical use.

346  Prior to the current work, the nature of preferred binding surfaces and molecular

determinants of suramin promiscuity were unclear. We reasoned that understanding these features

could facilitate structure-based design of new suramin analogs with increased target specificity.

Collectively, our analyses of available structures with suramin or suramin analogs revealed the

contributions of both the physicochemical nature of the target protein binding pocket and

conformational properties of suramin in making it a promiscuous ligand for diverse protein targets.

In particular, suramin's high flexibility, conferred by ten rotatable bonds, largely unsubstituted

phenyl groups, and numerous non-directional electrostatic interactions with positively charged

residues, collectively contribute to an ability to bind to structurally diverse nucleic acid interaction

surfaces. Additionally, our analyses revealed an enrichment of aromatic residues in favored

suramin binding sites and the role of various types of π-mediated interactions.

357  To date, suramin has been used as a lead in only a limited number of SAR studies. In one

case, suramin-derived fragments were designed, synthesized and modified to act as an antagonist

of the G protein-coupled receptor P_2Y_2 (38), which mediates microglial inflammation. In another

study, the toluene ring in suramin was substituted to improve suramin's binding to falcipain-2 (39),

a cysteine protease of the malaria parasite *Plasmodium falciparum*. Although both these

experimental studies used suramin as a lead, to our knowledge no high-throughput virtual

screening has been performed nor a structure-guided approach taken to improve the affinity and

selectivity of suramin based inhibitors. Here, we devised a virtual screening and precision docking

365 strategy using sets of suramin and heparin analogs, the latter of which were then used to guide the

366 redesign of top suramin analogs for enhanced target affinity and specificity. This redesign strategy

367 was fully exemplified using SARS-CoV-2 nucleocapsid and RdRp proteins in a proof-of-concept

368 redesign of suramin analogs as specific inhibitors of these viral proteins with potentially broad

369 anti-coronaviral activity.

370   Suramin has been explored for its inhibitory activity against both bacterial and viral

371 pathogens. Here, we demonstrated the potential utility of our virtual screening and ligand docking

372 enabled identification and redesign of suramin analogs to improve their affinity and specificity for

373 both known (RecA and HU) and new antibacterial targets (Fis and TsnR). For example, our study

374 produced suramin analogs which bind more tightly to HU, based on computational docking scores,

375 than those previously identified by screening inhibitors of HU (33). Similarly, our computational

376 strategy generated docking models for suramin and its analogs that can explain their previously

377 identified capacity to block the function of diverse viral proteins including HIV-1 gp120, norovirus

378 RdRp and bunyavirus nucleocapsid.

379   SARS-CoV-2 has also been shown to be inhibited by suramin (7) and, during preparation

380 of this manuscript, cryo-EM structural analysis and biochemical evidence revealed the molecular

381 basis of suramin binding to SARS-CoV-2 RdRp (21). Our docking study is consistent with these

382 structural insights in which multiple binding poses of suramin were found in the RNA binding

383 channel, one blocking the RNA template strand and the other binding to the primer strand in the

384 catalytic site. Suramin and its analogs have been shown to be 20-fold more potent than remdesivir

385 binding to the same site, and our results suggest that significant further improvement in suramin

386 analog design is possible. Specifically, we further computationally identified suramin analogs

387 which could potentially be more specific to this target and our docking guided elaboration of

suramin analogs showed improvement in the drug scaffold to make it a tighter binder.

Additionally, our studies identified the potential to target SARS-CoV-2 nucleocapsid with suramin

analogs and proposed analogs possessing enhanced affinity and specificity. This approach could

also be applied to the 3CL protease which was recently identified as another possible suramin

binding target (40). Our phylogenetic and evolutionary analyses also suggest that the suramin

binding pockets are evolutionary conserved in related coronavirus RdRp and nucleocapsid proteins

which could make these suramin analogs more broadly applicable therapeutics against this group

of viral pathogens.

In summary, this study has defined the basis of suramin promiscuity and the nature of its

preferred protein target binding sites and established a computational pipeline (**Fig. 2**) to rapidly

identify new suramin analogs targeting diverse viral and bacterial targets for novel antimicrobial

development. Suramin analogs screened in this study (e.g. Sur-8, Sur-23, Sur-25 and Sur-33) are

built from modular aromatic fragments which can be decorated with different polar or charged

substitutions and connected by various linkers (e.g. amide, sulfonamide and diazene; **Fig S8**).

These properties can be readily exploited for future fragment-based designs of suramin-like

inhibitors of viral and bacterial nucleic acid binding proteins. As such, our strategy offers the

possibility for expedient design of novel suramin analogs with improved properties and efficacy

for synthesis and experimental confirmation of their potential as novel lead antimicrobials.

MATERIALS AND METHODS

Virtual screening, docking and conformational analysis

Suramin and heparin analog sets were created using chemoinformatic guided searches in PubChem

with parameters: Tanimoto similarity of >90%, <40 rotatable bonds, and molecular weight < 2 kD.

The structures of compounds retrieved were processed with the LigPrep module in Schrödinger

Software. For conformational analysis of the ligands, the ConfGen module (41) of Schrödinger

Software was used with the "mix torsional low mode sampling" method, which is a combination

of *Monte Carlo* and systematic sampling of rotatable bonds to find low energy ligand

conformations.

To prepare each protein ("receptor") for ligand docking, hydrogen atoms were added, and

the structure energy minimized using the Protein Preparation Wizard in Schrödinger Software. The

docking grid location was determined by analysis of the nucleic acid binding region of each

protein, either by prediction using SiteMap (Schrödinger Software) or from published

experimental evidence for a given target protein. Docking grid size was determined based on the

nucleic acid binding pocket volume and designed to completely cover all residues and centered

symmetrically. Docking was performed using the virtual screening workflow of GLIDE (42) in

Schrödinger Software, in HTVS mode and then for precision docking in SP mode as used in our

previous studies (33, 43).

Ligand physicochemical properties and chemical similarity were calculated in the Canvas

module of Schrödinger Software (44). Ligand-protein nonbonded interactions were calculated

using BIOVIA DS Viewer for salt bridges, hydrogen bonds and other π mediated interactions.

PyMOL was used for the visual inspection of structures and generation of figures.

Phylogenetic analysis and conservation study

Homologous sequences of coronavirus RdRp and nucleocapsid were collected by BLAST search

in NCBI with ~10,000 and 5,000 sequences identified, respectively. The large number of

sequences arises due to the number of sequenced SARS-CoV-2 and related viruses that have >99%

identity in the dataset. Therefore, the CD-HIT server (45) was next used to remove redundant

sequences from the dataset using a sequence identity cut-off of 98%. Multiple sequence alignment

and evolutionary analyses were then performed using Clustal Omega and MEGA 7 (46),

respectively, with evolutionary history inferred using the Maximum Likelihood method with

bootstrap consensus tree inferred from 100 replicates. The fraction of replicate trees in which the

associated taxa clustered together in the bootstrap test (100 replicates) are shown next to the

branches. Evolutionary distances were computed using the JTT matrix-based method and are in

the units of number of amino acid substitutions per site. The rate variation among sites was

modeled with a gamma distribution (shape parameter 4). Finally, the phylogenetic tree was

visualized using FigTree (http://tree.bio.ed.ac.uk/software/figtree/). Conserved residues were

plotted on the SARS-CoV-2 RdRp (PDB 7BV1, chain A) and nucleocapsid (PDB 6M3M, chain

A) using ConSurf (47) from the pre-calculated multiple sequence alignment.

Acknowledgements

We thank Dr. William Wuest for comments on the manuscript and members of the Conn lab for

discussions and comment through the course of this work.

REFERENCES

1.   Guy RK, DiPaola RS, Romanelli F, Dutch RE. 2020. Rapid repurposing of drugs for
COVID-19. Science 368:829-830.

2.   Choy KT, Wong AY, Kaewpreedee P, Sia SF, Chen D, Hui KPY, Chu DKW, Chan MCW,
Cheung PP, Huang X, Peiris M, Yen HL. 2020. Remdesivir, lopinavir, emetine, and
homoharringtonine inhibit SARS-CoV-2 replication in vitro. Antiviral Res 178:104786.

3.   Costanzo M, De Giglio MAR, Roviello GN. 2020. SARS-CoV-2: Recent Reports on
Antiviral Therapies Based on Lopinavir/Ritonavir, Darunavir/Umifenovir,

Hydroxychloroquine, Remdesivir, Favipiravir and other Drugs for the Treatment of the New

Coronavirus. Curr Med Chem 27:4536-4541.

4. Abraham SM, Lawrence T, Kleiman A, Warden P, Medghalchi M, Tuckermann J,

Saklatvala J, Clark AR. 2006. Antiinflammatory effects of dexamethasone are partly

dependent on induction of dual specificity phosphatase 1. J Exp Med 203:1883-9.

5. Solinas C, Perra L, Aiello M, Migliori E, Petrosillo N. 2020. A critical evaluation of

glucocorticoids in the management of severe COVID-19. Cytokine Growth Factor Rev

doi:10.1016/j.cytogfr.2020.06.012.

6. Heidary F, Gharebaghi R. 2020. Ivermectin: a systematic review from antiviral effects to

COVID-19 complementary regimen. J Antibiot (Tokyo) doi:10.1038/s41429-020-0336-z.

7. da Silva CSB, Thaler M, Tas A, Ogando NS, Bredenbeek PJ, Ninaber DK, Wang Y,

Hiemstra PS, Snijder EJ, van Hemert MJ. 2020. Suramin inhibits SARS-CoV-2 infection in

cell culture by interfering with early steps of the replication cycle. Antimicrob Agents

Chemother doi:10.1128/AAC.00900-20.

8. Brun R, Blum J, Chappuis F, Burri C. 2010. Human African trypanosomiasis. Lancet

375:148-59.

9. Hawking F. 1978. Suramin: with special reference to onchocerciasis. Adv Pharmacol

Chemother 15:289-322.

10. Morgan HP, McNae IW, Nowicki MW, Zhong W, Michels PA, Auld DS, Fothergill-

Gilmore LA, Walkinshaw MD. 2011. The trypanocidal drug suramin and other trypan blue

mimetics are inhibitors of pyruvate kinases and bind to the adenosine site. J Biol Chem

286:31232-40.

11. Fleck SL, Birdsall B, Babon J, Dluzewski AR, Martin SR, Morgan WD, Angov E,

Kettleborough CA, Feeney J, Blackman MJ, Holder AA. 2003. Suramin and suramin

analogues inhibit merozoite surface protein-1 secondary processing and erythrocyte invasion

by the malaria parasite Plasmodium falciparum. J Biol Chem 278:47670-7.

12. Wiedemar N, Hauser DA, Maser P. 2020. 100 Years of Suramin. Antimicrob Agents

Chemother 64.

13. Fantini J, Hammache D, Delezay O, Yahi N, Andre-Barres C, Rico-Lattes I, Lattes A. 1997.
Synthetic soluble analogs of galactosylceramide (GalCer) bind to the V3 domain of HIV-1
gp120 and inhibit HIV-1-induced fusion and entry. J Biol Chem 272:7245-52.

14. Aguilar JS, Rice M, Wagner EK. 1999. The polysulfonated compound suramin blocks
adsorption and lateral difusion of herpes simplex virus type-1 in vero cells. Virology
258:141-51.

15. Garson JA, Lubach D, Passas J, Whitby K, Grant PR. 1999. Suramin blocks hepatitis C
binding to human hepatoma cells in vitro. J Med Virol 57:238-42.

16. Chen Y, Maguire T, Hileman RE, Fromm JR, Esko JD, Linhardt RJ, Marks RM. 1997.
Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate.
Nat Med 3:866-71.

17. Ren P, Zheng Y, Wang W, Hong L, Delpeyroux F, Arenzana-Seisdedos F, Altmeyer R.
2017. Suramin interacts with the positively charged region surrounding the 5-fold axis of the
EV-A71 capsid and inhibits multiple enterovirus A. Sci Rep 7:42902.

18. Albulescu IC, Kovacikova K, Tas A, Snijder EJ, van Hemert MJ. 2017. Suramin inhibits
Zika virus replication by interfering with virus attachment and release of infectious particles.
Antiviral Res 143:230-236.

19. Mastrangelo E, Pezzullo M, Tarantino D, Petazzi R, Germani F, Kramer D, Robel I,
Rohayem J, Bolognesi M, Milani M. 2012. Structure-based inhibition of Norovirus RNA-
dependent RNA polymerases. J Mol Biol 419:198-210.

20. Albulescu IC, van Hoolwerff M, Wolters LA, Bottaro E, Nastruzzi C, Yang SC, Tsay SC,
Hwu JR, Snijder EJ, van Hemert MJ. 2015. Suramin inhibits chikungunya virus replication
through multiple mechanisms. Antiviral Res 121:39-46.

21. Yin W, Luan X, Li Z, Zhou Z, Wang Q, Gao M, Wang X, Zhou F, Shi J, You E, Liu M,
Wang Q, Jiang Y, Jiang H, Xiao G, Zhang L, Yu X, Zhang S, Eric Xu H. 2021. Structural
basis for inhibition of the SARS-CoV-2 RNA polymerase by suramin. Nat Struct Mol Biol
28:319-325.

22. Ullmann H, Meis S, Hongwiset D, Marzian C, Wiese M, Nickel P, Communi D, Boeynaems

JM, Wolf C, Hausmann R, Schmalzing G, Kassack MU. 2005. Synthesis and structure-

activity relationships of suramin-derived P2Y11 receptor antagonists with nanomolar

potency. J Med Chem 48:7040-8.

23. Trapp J, Meier R, Hongwiset D, Kassack MU, Sippl W, Jung M. 2007. Structure-activity

studies on suramin analogues as inhibitors of NAD+-dependent histone deacetylases

(sirtuins). ChemMedChem 2:1419-31.

24. Kaina B. 2021. On the Origin of SARS-CoV-2: Did Cell Culture Experiments Lead to

Increased Virulence of the Progenitor Virus for Humans? In Vivo 35:1313-1326.

25. Murakami MT, Arruda EZ, Melo PA, Martinez AB, Calil-Elias S, Tomaz MA, Lomonte B,

Gutierrez JM, Arni RK. 2005. Inhibition of myotoxic activity of Bothrops asper myotoxin II

by the anti-trypanosomal drug suramin. J Mol Biol 350:416-26.

26. Schuetz A, Min J, Antoshenko T, Wang CL, Allali-Hassani A, Dong A, Loppnau P, Vedadi

M, Bochkarev A, Sternglanz R, Plotnikov AN. 2007. Structural basis of inhibition of the

human NAD+-dependent deacetylase SIRT5 by suramin. Structure 15:377-89.

27. Ren C, Morohashi K, Plotnikov AN, Jakoncic J, Smith SG, Li J, Zeng L, Rodriguez Y,

Stojanoff V, Walsh M, Zhou MM. 2015. Small-molecule modulators of methyl-lysine

binding for the CBX7 chromodomain. Chem Biol 22:161-8.

28. Jiao L, Ouyang S, Liang M, Niu F, Shaw N, Wu W, Ding W, Jin C, Peng Y, Zhu Y, Zhang

F, Wang T, Li C, Zuo X, Luan CH, Li D, Liu ZJ. 2013. Structure of severe fever with

thrombocytopenia syndrome virus nucleocapsid protein in complex with suramin reveals

therapeutic potential. J Virol 87:6829-39.

29. Kang S, Yang M, Hong Z, Zhang L, Huang Z, Chen X, He S, Zhou Z, Zhou Z, Chen Q, Yan

Y, Zhang C, Shan H, Chen S. 2020. Crystal structure of SARS-CoV-2 nucleocapsid protein

RNA binding domain reveals potential unique drug targeting sites. Acta Pharm Sin B

doi:10.1016/j.apsb.2020.04.009.

30. Yahi N, Sabatier JM, Nickel P, Mabrouk K, Gonzalez-Scarano F, Fantini J. 1994. Suramin

inhibits binding of the V3 region of HIV-1 envelope glycoprotein gp120 to

galactosylceramide, the receptor for HIV-1 gp120 on human colon epithelial cells. J Biol
Chem 269:24349-53.

31. Dillon SC, Dorman CJ. 2010. Bacterial nucleoid-associated proteins, nucleoid structure and
544
545 gene expression. Nat Rev Microbiol 8:185-95.

32. Stella S, Cascio D, Johnson RC. 2010. The shape of the DNA minor groove directs binding
546
547 by the DNA-bending protein Fis. Genes Dev 24:814-26.

33. Bhowmick T, Ghosh S, Dixit K, Ganesan V, Ramagopal UA, Dey D, Sarma SP,
548
549 Ramakumar S, Nagaraja V. 2014. Targeting Mycobacterium tuberculosis nucleoid-
550 associated protein HU with structure-based inhibitors. Nat Commun 5:4124.

34. Dey D, Nagaraja V, Ramakumar S. 2017. Structural and evolutionary analyses reveal
551
552 determinants of DNA binding specificities of nucleoid-associated proteins HU and IHF. Mol
553 Phylogenet Evol 107:356-366.

35. Nautiyal A, Patil KN, Muniyappa K. 2014. Suramin is a potent and selective inhibitor of
554
555 Mycobacterium tuberculosis RecA protein and the SOS response: RecA as a potential target
556 for antibacterial drug discovery. J Antimicrob Chemother 69:1834-43.

36. Dunstan MS, Hang PC, Zelinskaya NV, Honek JF, Conn GL. 2009. Structure of the
557
558 thiostrepton resistance methyltransferase.S-adenosyl-L-methionine complex and its
559 interaction with ribosomal RNA. J Biol Chem 284:17013-20.

37. Peng Q, Peng R, Yuan B, Zhao J, Wang M, Wang X, Wang Q, Sun Y, Fan Z, Qi J, Gao GF,
560
561 Shi Y. 2020. Structural and Biochemical Characterization of the nsp12-nsp7-nsp8 Core
562 Polymerase Complex from SARS-CoV-2. Cell Rep 31:107774.

38. Pillaiyar T, Funke M, Al-Hroub H, Weyler S, Ivanova S, Schlegel J, Abdelrahman A,
563
564 Muller CE. 2020. Design, synthesis and biological evaluation of suramin-derived dual
565 antagonists of the proinflammatory G protein-coupled receptors P2Y2 and GPR17. Eur J
566 Med Chem 186:111789.

39. Marques AF, Esser D, Rosenthal PJ, Kassack MU, Lima LM. 2013. Falcipain-2 inhibition
567
by suramin and suramin analogues. Bioorg Med Chem 21:3667-73.

40. Zhu W, Xu M, Chen CZ, Guo H, Shen M, Hu X, Shinn P, Klumpp-Thomas C, Michael SG,
Zheng W. 2020. Identification of SARS-CoV-2 3CL Protease Inhibitors by a Quantitative
High-throughput Screening. bioRxiv doi:10.1101/2020.07.17.207019.

41. Watts KS, Dalal P, Murphy RB, Sherman W, Friesner RA, Shelley JC. 2010. ConfGen: a
conformational search method for efficient generation of bioactive conformers. J Chem Inf
Model 50:534-46.

42. Friesner RA, Banks JL, Murphy RB, Halgren TA, Klicic JJ, Mainz DT, Repasky MP, Knoll
EH, Shelley M, Perry JK, Shaw DE, Francis P, Shenkin PS. 2004. Glide: a new approach for
rapid, accurate docking and scoring. 1. Method and assessment of docking accuracy. J Med
Chem 47:1739-49.

43. Dey D, Kavanaugh LG, Conn GL. 2020. Antibiotic Substrate Selectivity of Pseudomonas
aeruginosa MexY and MexB Efflux Systems Is Determined by a Goldilocks Affinity.
Antimicrob Agents Chemother 64.

44. Sastry M, Lowrie JF, Dixon SL, Sherman W. 2010. Large-scale systematic analysis of 2D
fingerprint methods and parameters to improve virtual screening enrichments. J Chem Inf
Model 50:771-84.

45. Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. Bioinformatics 28:3150-2.

46. Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. Mol Biol Evol 33:1870-4.

47. Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, Ben-Tal N. 2016. ConSurf
2016: an improved methodology to estimate and visualize evolutionary conservation in
macromolecules. Nucleic Acids Res 44:W344-50.

Figure 1. Structural basis for suramin protein target site diversity. Chemical structures of *A*,

suramin and *B*, heparin. Crystal structures of suramin (or partial fragment) bound to: *C*,

Leishmania mexicana pyruvate kinase (PDB 3PP7), *D*, snake venom protein myotoxin (PDB

1Y4L), *E*, NAD$^+$-dependent protein deacetylase SIRT5 (PDB 2NYR), and *F*, epigenetic reader

chromobox homolog 7 (PDB 4X3U). *H*, Residue propensities in the suramin binding pockets in

targets shown in *panels C-F* and **Figure S1**. *G*, Superposition via the boxed sulfo-naphthyl ring of

suramin molecules from co-crystal structures with diverse targets (PDB 1Y4L, 2NYR, 4X3U,

6CE2 and 3BF6).

Figure 2. Computational workflow used in this study. *A*, Suramin and heparin analog sets were

retrieved from PubChem with the indicated parameter filters and subject to conformer generation

in LigPrep (Schrödinger Software). *B,* Suramin and heparin analog sets were used for separate

HTVS and the resulting top 100 ligands then subject to higher precision (SP) docking in GLIDE

(Schrödinger Software). *C*, Following further shape and energetic analysis, chemoinformatics was

used to determine different fragments of top scoring suramin analogs. For heparin analogs, the top

poses corresponding to the top selected suramin analog for each target was next used for

pharmacophore-based identification of regions in the suramin analog which could be substituted

to improve its predicted affinity and target specificity. Identified novel analogs would next be used

as leads for experimental analysis and validation.

617

618 **Figure 3. Virtual screening and precision docking identify suramin analogs with increased**

619 **target specificity.** HTVS docking scores for *A,* suramin and *B,* heparin analogs against bacterial

620 targets HU, Fis, RecA and TsnR; and, viral targets gp120, SARS-CoV-2 and norovirus RdRp, and

621 SARS-CoV-2 and bunyavirus nucleocapsid (SARS-NC and BV-NC, respectively). A high

622 percentage of both *C*, suramin and *D*, heparin analogs bind to only one or two targets within the

623 protein panel used, suggesting that virtual screening has identified more specific leads as compared

624 to the parent molecules, suramin and heparin. *E*,*F*, SP GLIDE docking scores for the top 100

625 ligands from HTVS for suramin and heparin analogs, respectively.

Figure 4. Docking-guided elaboration of the top identified suramin analog against HU. *A*, SP GLIDE docking pose of suramin analog Sur-1 (purple) bound to the DNA interaction surface of HU (top, white surface; bottom, electrostatic potential surface). *B*, Comparison of the chemical structures of suramin and Sur-1, highlighting the reduced number of rotatable bonds in the analog. *C*, SP GLIDE docking pose of heparin analog Hep-1 (orange) bound to the same surface of HU. *D*, The overlapping binding poses of Hep-1 and Sur-1 which were used to identify sites in Sur-1 for redesign (indicated *) to improve binding affinity and specificity.

634

Figure 5. Design of suramin analogs specific to SARS-CoV-2 RdRp and nucleocapsid. *A*, Location of suramin analogs Sur-36 and Sur-84 in the RNA binding tunnel of SARS-CoV-2 RdRp. *B*, Zoomed in views of the boxed region in *panel A* of the SP GLIDE docking pose of Sur-36, an existing analog of suramin, bound to the RNA binding tunnel. *C*, Iterative rounds of chemical enumeration (**Figure S6**) and docking, resulted in the designed suramin analog Sur-84 shown in the same binding pocket. *D*, Location of suramin analogs Sur-31 and Sur-127 on the RNA binding surface of SARS-CoV-2 nucleocapsid. *E*,*F*, Zoomed in views of the boxed region in *panel D* of the SP GLIDE docking poses of Sur-31 and Sur-127 respectively. *G*, Comparison of SP GLIDE docking scores for the initial top 100 suramin analogs (purple) and analogs tested during the iterative chemical enumeration process (green), showing the improvement in predicted affinity. Suramin (red square), starting analogs (Sur-31 and Sur-36 for nucleocapsid (NC) and RdRp, respectively) and final redesigned analogs (Sur-127 and Sur-85) are highlighted.

Figure 6. Maximum likelihood phylogenetic tree of coronavirus RdRp and nucleocapsid. *A*, Phylogenetic tree of coronavirus capsid protein showing a separate clade for the β-coronavirus genus, with SARS-CoV-2 and MERS-CoV, which are mostly of bat origin. This clade is closely related to γ coronaviruses of avian origin while another bat origin α-coronavirus family forms a separate clade. Conservation of residues in coronavirus capsid proteins shows the binding pocket targeted by the designed suramin analog Sur-127 is evolutionarily conserved. *B*, As *panel A* but for SARS-CoV-2 RdRp and Sur-85