Runaway evolution from male-male competition

Allen J. Moore[1] | Joel W. McGlothlin[2] | Jason B. Wolf[3]

[1]Department of Entomology, University of Georgia, Athens, Georgia, United States

[2]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, United States

[3]Milner Centre for Evolution and Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom

Email: ajmoore@uga.edu, joelmcg@vt.edu, j.b.wolf@bath.ac.uk

Correspondence may be addressed to any author.

AUTHOR CONTRIBUTIONS

AJM, JWM, and JBW conceived the idea, developed the model, and wrote the paper.

KEYWORDS

aggression, honest signals, indirect genetic effects, male-male competition, quantitative genetics, runaway evolution, sexual selection, social evolution, weapons

Abstract

Wondrously elaborate weapons and displays that appear to be counter to ecological optima are widespread features of male contests for mates across the animal kingdom. To understand how such diverse traits evolve, here we develop a quantitative genetic model of sexual selection for a male signaling trait that mediates aggression in male-male contests and show that an honest indicator of aggression can generate selection on itself by altering the social environment. This can cause selection to accelerate as the trait is elaborated, leading to runaway evolution. Thus, an evolving source of selection provided by the social environment is the fundamental unifying feature of runaway sexual selection driven by either male-male competition or female mate choice. However, a key difference is that runaway driven by male-male competition requires signal honesty. Our model identifies simple conditions that provide clear, testable predictions for empirical studies using standard quantitative genetic methods.

INTRODUCTION

Darwin (1859, 1871) chronicled the amazing diversity of traits associated with success in

mating, proposing the theory of sexual selection to explain the diverse exaggerated,

spectacular, and bizarre structures and behaviors found in males of many species. Darwin

suggested that such traits evolve either because they enhance success in contests between

males for access to females or because they are preferred by females when choosing mates.

As Darwin (1871) wrote, "It is certain that amongst almost all animals there is a struggle

between the males for the possession of the female. This fact is so notorious that it would be

superfluous to give examples." In contrast, the ability of females to influence evolution

through choice of partners was almost immediately questioned and continued to be

controversial for decades after Darwin (Wallace 1889; Huxley 1938). However, theoretical

models of evolution via female choice (Lande 1981; Kirkpatrick 1982; Mead & Arnold 2004)

and empirical research documenting female preference in nature (Andersson 1982, 1994;

Rosenthal 2017) eventually led to mate choice becoming the dominant paradigm in studies of

sexual selection. The development of formal mathematical models showing that male traits

and female preferences coevolve in self-reinforcing fashion, an idea first proposed by Fisher

(1915, 1930), was particularly crucial to the acceptance of mate choice as an important

evolutionary mechanism. The key component of the Fisher process is that female preference

and a preferred male trait become genetically correlated as a result of assortative mating that

generates linkage disequilibrium between the preference and male trait. This can cause

sexually selected male traits to evolve at ever-increasing speed, a pattern that has been

referred to as an evolutionary "runaway" (Fisher 1930; Lande 1981; Bailey & Moore 2012).

Despite the current bias towards studies focused on mate choice, Darwin was not

wrong about male-male competition. Members of entire taxa are characterized by highly

modified sexually dimorphic structures that function only in male contests (e.g., Dermaptera,

Briceño & Eberhard 1995). Weapons can evolve to be massive and create real functional

constraints for the males that bear them, and such bizarrely elaborate and diverse structures

associated with duels are indeed found across the animal kingdom (Emlen 2008, 2014;

McCullough *et al.* 2016; O'Brien *et al.* 2018). In fact, male-male competition remains a more

common source of selection shaping male traits that influence mating success, and traits

expressed in male-male interactions can be as elaborate as those that are the target of female

preferences (Darwin 1871; Huxley 1938; Andersson 1994; Moore & Moore 2006; Emlen

2008, 2014; McCullough *et al.* 2016; O'Brien *et al.* 2018). However, we still lack robust

genetic models that generate testable predictions for the evolution of sexually selected traits

via male-male competition. Notably, the potential for male-male competition to result in a

runaway process that drives extreme trait elaboration remains unresolved.

Many elaborate male traits used in male-male contests, such as showy plumage

(Hagelin 2002), color (Seehausen & Schluter 2004), pheromones (Moore *et al.* 1997b), and

structures such as antlers (Wilkinson & Dodson 1997), horns (Emlen *et al.* 2005), forceps

(Briceño & Eberhard 1995), and claws (Sneddon *et al.* 1997) function as signals that may

provide information about some underlying qualities of the individuals, such as the

willingness or ability to fight (Parker 1974; Maynard Smith & Harper 1988; Maynard Smith

& Harper 2003; Emlen 2008, 2014). For example, there is often a positive association

between signals or weapons and other traits such as body size (McCullough *et al.* 2016;

O'Brien *et al.* 2018), making the signal or weapon an honest indicator of potential threat to an

opponent or a "badge of status" (Rohwer 1982). As such, males are expected to adaptively

modulate their behavior in response to these signaling traits, escalating contests they are more

likely to win and withdrawing from ones they are more likely to lose. Because the effect of

signaling traits inherently depends on social context, such traits serve as both targets and

sources of selection, potentially leading to self-reinforcing and accelerating selection as

occurs in the runaway process driven by female preference (Lande 1981; Bailey & Kölliker

2019). However, despite insights from game theory models (Parker 1974; Maynard Smith &

Brown 1986; Maynard Smith & Harper 1988; Maynard Smith & Harper 2003; Rutte *et al.*

2006), how this fundamental feature of extreme elaboration—an evolving source of

selection—may arise within male-male contests is unclear.

Here, we utilize a framework that explicitly incorporates socially contingent trait

expression and fitness (Moore *et al.* 1997a; Wolf *et al.* 1999; McGlothlin *et al.* 2010) to

model trait evolution arising from male-male competition. We show that when honest signals

are used to modulate the behavior of competitors, male-male competition leads to

evolutionary elaboration of male traits. We identify the necessary and sufficient conditions for

trait elaboration to become a runaway process and outline predictions that can be empirically

tested to evaluate this scenario in natural systems. We show that sexual selection by male-

male competition can have features that are analogous to those of runaway sexual selection by

female choice; just as in female mate choice, the social environment in male-male contests

may generate a self-reinforcing source of selection on the traits that mediate the interaction,

potentially leading to self-sustaining and escalating selection.

MODEL

To capture the influence of the social environment in a model of male-male competition, we

assume that individuals adjust their behavior in response to the signaling trait values of their

social partners, an assumption that is supported empirically and theoretically (Parker 1974;

West-Eberhard 1979; Maynard Smith 1982; West-Eberhard 1983, 1984; Moore *et al.* 1997a;

West-Eberhard 2003; O'Brien *et al.* 2018; Tinghitella *et al.* 2018; Rico-Guevara & Hurme

2019; Wiens & Tuschhoff 2020). Because the social context (i.e., the social environment) is

constructed from traits of conspecifics, this flexible response to social context provides the

opportunity for indirect genetic effects (Moore *et al.* 1997a), which allow the social

environment itself to evolve (Moore *et al.* 2002; Wolf 2003; Bijma & Wade 2008;

McGlothlin *et al.* 2010). Evolutionary changes in the social environment can lead to

concerted evolution because the social environment can be a source of selection on the traits

that themselves compose the social environment (West-Eberhard 1979; Wolf *et al.* 1999;

Bailey & Kölliker 2019; Araya-Ajoy *et al.* 2020; McGlothlin *et al.* 2021). Such "social

selection" (West-Eberhard 1979, 1983, 1984; Wolf *et al.* 1999; Bijma & Wade 2008;

McGlothlin *et al.* 2010) is expected to arise whenever traits act as both agents and targets of

selection.

Our model assumptions are based on common conditions observed in male-male

contests (Emlen 2008, 2014; Eberhard *et al.* 2018). Although we model the outcome of

pairwise duels between males drawn at random from the population, our model is easily

generalized to include multiple interactions between males (Supporting Appendix). First, we

assume that males possess a trait (designated by the subscript S) that is used as a signal

conveying potential threat in social contests. There are many diverse examples include traits

such as a plumage badges, exaggerated weapons, or vocal or chemical signals. Elaboration of

the signal may consist of an increase in size or complexity, although for heuristic simplicity,

we discuss the evolution of increased signal size. Second, we assume that males vary in the

underlying quality trait that reflects their fighting ability or some other aspect of their

phenotype that determines the potential interaction cost they represent to their opponent in a

contest. We describe this trait as body size (designated by the subscript B) for simplicity (see

the discussion of male quality in Eberhard *et al.* 2018). Finally, we assume males respond to

the assessment of the signal by modulating their behavioral response of aggression toward

their opponent (designated by the subscript A) within the contest because the signal provides

information on the likelihood that they would win an escalated contest (see below).

We assume that both signal size (z_S) and body size (z_B) are normally distributed

metric traits influenced by many loci of small effect. Expression of these traits can be

partitioned into heritable additive genetic effects (denoted a_S and a_B) and general non-

heritable (environmental and nonadditive genetic) effects (denoted e_S and e_B). We assume

that neither signal nor body size changes as a result of the social interaction. An individual's

total phenotypic value for each trait is then described by a simple sum of the heritable and

non-heritable components:

$$z_i = a_i + e_i, \tag{1}$$

where a_i is normally distributed with mean $\bar{a}_i$ and variance G_{ii} and e_i is normally distributed

with mean 0 and variance E_{ii}. We make the standard quantitative genetic assumption that

heritable and non-heritable components are uncorrelated.

We model male-male competition where the larger males will defeat smaller males in

a fight. We therefore further assume that the phenotypic value for aggressive behavior (z_A)

associated with a given genotype depends on social context, influenced by their rival's signal

size relative to their own, as suggested by West-Eberhard (1979, 1983, 1984). This effect is

captured in our model as a term in which aggression scales with the magnitude of the size

difference between opponents, which is supported by optimality models and empirical studies

(Huxley 1938; Parker 1974; Riechert 1984; Sneddon *et al.* 1997; Maynard Smith & Harper

2003; Emlen 2008, 2014). The phenotypic value of aggression can thus be written:

$$z_A = a_A + e_A + \delta_{AS}(z_S - z_S'), \tag{2}$$

where a_A and e_A represent standard additive genetic and uncorrelated non-heritable

components, respectively. Here and elsewhere, terms with primes indicate a value assigned to

the focal individual's opponent so z_S is the phenotypic value of the signal of the focal

individual and z_S' of their opponent. The coefficient δ_{AS} measures the influence of the

difference in signal size on the expression of aggressive behavior. Thus, δ_{AS} is analogous to

the ψ term in standard interacting phenotype models (Moore *et al.* 1997a), but differs because

it depends upon the value of an interactant's phenotype relative to the focal individual.

Because signal size is heritable, the phenotype value of aggression for the focal individual

includes modifications arising from both direct genetic effects of their own genotype ($\delta_{AS}a_S$)

and indirect genetic effects ($-\delta_{AS}a_S'$), which is defined as the effect of a social interactant's

genotype on the focal individual's phenotype (Moore *et al.* 1997a). We describe the

relationship between this model and the standard model of indirect genetic effects in the

Supporting Appendix.

The underlying genetic value of aggression is assumed to be genetically uncorrelated

to both that of signal size and body size (i.e., there is no direct pleiotropic relationship

between the traits such that genetic covariances $G_{SA} = G_{BA} = 0$). This is a conservative

assumption as a positive correlation would result in even faster evolution. However, the signal

may be genetically correlated to body size, providing signal honesty, which is quantified by

the covariance between signal size and body size (G_{SB}). Because the level of aggression

displayed is conditional on the social context, a correlation within the population is generated

if males with larger signals and/or larger body size are more aggressive on average (and vice

versa). Hence, aggression can be correlated to the signal and body size traits despite the

absence of a direct pleiotropic link (or linkage disequilibrium) because the flexible behavioral

response creates a relationship between these traits through the social interaction.

Selection imposed by male-male competition

In social interactions, associations between traits and fitness may cause selection via two pathways: nonsocial selection (quantified by the gradient β_N), which arises from effects of a focal individual's traits on its own fitness, and social selection (quantified by the gradient β_S), which arises from the effects of an opponent's traits on the fitness of a focal individual (Wolf *et al.* 1999). From Wolf *et al.* (1999), when both nonsocial and social selection are present, individual relative fitness can be written:

$$w = w_0 + \mathbf{z}^{\mathrm{T}}\boldsymbol{\beta_N} + \mathbf{z}'^{\mathrm{T}}\boldsymbol{\beta_S} + \varepsilon, \tag{3}$$

where w_0 is an intercept, $\mathbf{z}$ and $\mathbf{z}'$ are column vectors of focal and opponent traits, $\boldsymbol{\beta_N}$ and $\boldsymbol{\beta_S}$ are vectors of nonsocial and social selection gradients, ε is an uncorrelated error term, and the superscript T denotes transposition. Expressing relative fitness using Eq. 3 has two distinct advantages. First, selection gradients can be estimated in natural populations using multiple regression (Lande & Arnold 1983; Wolf *et al.* 1999; Formica *et al.* 2011; Fisher & Pruitt 2019), allowing our model to generate testable predictions. Second, selection gradients can be combined with genetic parameters to predict short-term evolutionary response to selection (Lande & Arnold 1983; Bijma & Wade 2008; McGlothlin *et al.* 2010).

To understand how these selection gradients arise from male-male contests with signaling, we can use evolutionary game theory (see Supporting Appendix) to write a mechanistic expression for relative fitness:

$$w = w_0 + b_A(z_A - z_A') - c_A z_A z_A' - c_B z_A' z_B' - c_S(\bar{z}_S - \theta_S)z_S + \varepsilon, \tag{4}$$

where terms including *b* represent fitness benefits and terms including *c* represent fitness costs. In Eq. 4, the benefit term and the first cost term derive from the hawk-dove model of evolutionary game theory (Appendix; Maynard Smith & Price 1973; Maynard Smith 1982;

180 McGlothlin *et al.* 2021). The coefficient b_A is the fitness benefit of winning a contest, which

181 we assume derives from greater access to females. In a contest, access to females is

182 determined by a focal individual's aggression relative to its opponent. Multiplying b_A by

183 $(z_A - z'_A)$ reflects the fact that the probability of winning a contest increases as a male can

184 become increasingly more aggressive than its opponent. This benefit, however, does not come

185 without a cost. The term $c_A z_A z'_A$ is the fitness cost of aggression associated with escalation of

186 encounters. Logically, an individual pays a cost for acting aggressive that depends on the

187 level of aggression shown by its opponent. In the hawk-dove model, this corresponds to the

188 cost associated with a hawk player facing a hawk opponent, which increases in likelihood as

189 both players act increasingly aggressive. A second fitness cost $(c_B z'_A z'_B)$ reflects the fact that

190 the fitness impact of aggression by an opponent (z'_A) depends on the size of the opponent (z'_B).

191 This cost, which we call the threat of the opponent, derives from the fact that larger males

192 impose a greater risk of harm than do smaller males. Finally, we assume that a third cost

193 $(c_S(\bar{z}_S - \theta_S)z_S)$ arises from natural selection favoring some optimal trait value (θ_S), which

194 therefore will oppose signal elaboration. Following a Gaussian model of selection (Lande

195 1976, 1979), selection against elaborate signals becomes stronger as the population mean of

196 the signal $(\bar{z}_S)$ becomes further away from its naturally selected optimum (θ_S). Although we

197 do not do so here, this term could be replaced with a multivariate Gaussian term (Lande 1979)

198 to add naturally selected optima for aggression and body size.

199  Taking partial derivatives of with respect to focal and opponent traits (evaluated at the

200 population mean) allows us to translate the fitness model in Eq. 4 into nonsocial and social

201 selection gradients (McGlothlin *et al.* 2021). The nonsocial gradients are:

202 $\beta_{N_S} = -c_S(\bar{z}_S - \theta_S)$            (5a)

203 $\beta_{N_B} = 0$                (5b)

204 $\beta_{N_A} = b_A - c_A\bar{z}_A$            (5c)

and social selection gradients:

$$\beta_{S_S} = 0 \tag{6a}$$

$$\beta_{S_B} = -c_B \bar{z}_A \tag{6b}$$

$$\beta_{S_A} = -b_A - c_A \bar{z}_A - c_B \bar{z}_B. \tag{6c}$$

Thus, males with large signals are selected against via nonsocial selection but interacting with

such males does not directly impose social selection. Body size is not under direct nonsocial

selection but imposes a fitness cost via social selection that increases with the population

mean of aggression. Nonsocial selection favors aggression until the benefits of aggression are

outweighed by the costs, while social selection imposed by opponent's aggression is always

negative, representing a net fitness cost. This gradient becomes increasingly negative as the

population mean aggression and body size increase. These selection gradients suggest that

signal size itself experiences no direct sexual selection. If signal size increases, it must do so

as an indirect response to selection on a correlated trait.

Evolutionary response to selection

Selection within a generation is translated into an evolutionary response across generations

through the association between the phenotype, upon which selection acts, and the genotype,

which contributes to the inheritance of the traits across generations. In quantitative genetics,

this genotype-phenotype relationship is most often summarized by the additive genetic

variance, which is used to predict evolutionary response to selection across generations

(Lande & Arnold 1983; Arnold 1994). However, for traits expressed in social interactions, we

must also consider social pathways to fitness, which arise from indirect genetic effects and

social selection, when calculating response to selection (Moore *et al.* 1997a; Bijma & Wade

2008; McGlothlin *et al.* 2010). Because the model of phenotypic modification described in

Eq. 2 deviates from the standard model of indirect genetic effects, we develop a general

equation for response to selection in the Supporting Appendix (Eq. A10). Using this equation,

the response to selection for the three traits in our model can in general be written:

$$\Delta \bar{z}_S = \frac{1}{2} G_{SS} \left(\beta_{N_S} + \delta_{AS} (\beta_{N_A} - \beta_{S_A}) \right) + \frac{1}{2} G_{SB} \beta_{N_B} \tag{7a}$$

$$\Delta \bar{z}_B = \frac{1}{2} G_{BB} \beta_{N_B} + \frac{1}{2} G_{SB} \left(\beta_{N_S} + \delta_{AS} (\beta_{N_A} - \beta_{S_A}) \right) \tag{7b}$$

$$\Delta \bar{z}_A = \frac{1}{2} G_{AA} \beta_{N_A}. \tag{7c}$$

The multiplier ½ in Eq. 7 arises because selection is acting only on males. Eq. 7a shows that

modification of aggressive behavior in response to the signaling trait (δ_{AS}) causes both

nonsocial and social selection gradients for aggression to contribute to signal evolution. This

behavioral modification also contributes to evolution of body size when the signal is honest,

which is captured by the covariance between signal size and body size (G_{SB}; Eq. 7b). This is

easily shown by setting the modification of aggression based on the signaling trait (δ_{AS}) to 0,

which recovers standard quantitative genetic expressions for evolution. In contrast, behavioral

modification never contributes to evolution of aggression (Eq. 7c).

By substituting Eqs. 5-6 into Eq. 7, we can predict evolutionary change using our

mechanistic fitness model (Eq. 4):

$$\Delta \bar{z}_S = \frac{1}{2} G_{SS} \left(\delta_{AS} (2 b_A + c_B \bar{z}_B) - c_S (\bar{z}_S - \theta_S) \right) \tag{8a}$$

$$\Delta \bar{z}_B = \frac{1}{2} G_{SB} \left(\delta_{AS} (2 b_A + c_B \bar{z}_B) - c_S (\bar{z}_S - \theta_S) \right) \tag{8b}$$

$$\Delta \bar{z}_A = \frac{1}{2} G_{AA} (b_A - c_A \bar{z}_A). \tag{8c}$$

Eq. 8a shows that when fitness is defined as in Eq. 4, evolution of the signaling trait beyond

its naturally selected optimum depends crucially on modification of aggression. If males do

not change their aggression to the signal (i.e., if $\delta_{AS} = 0$), the population mean of the

signaling trait cannot increase. From Eq. 8b, the evolution of body size depends on both δ_{AS}

and the presence of signal honesty (i.e., $G_{SB} > 0$). Eqs. 8a-b also show that evolution of the

signaling trait and of male body size is potentially open-ended because the evolutionary

response to selection for each trait becomes stronger as the population mean body size

increases. In contrast, from Eq. 8c, the evolution of aggression is self-limiting because

selection depends on the balance of the benefits and costs of aggression, the latter of which

become more intense as mean aggression intensifies. This observation suggests that both

signal size and body size may experience runaway evolution if the benefits of aggression and

the threat of the opponent are strong enough to outweigh natural selection against elaborate

signals, whereas aggression should always tend to quickly evolve to an equilibrium value.

To solve for equilibrium and to explore the conditions for such a runaway, we set Eqs.

8a-c equal to zero and solve for the equilibrium mean of each trait ($\hat{z}_i$):

$$\hat{z}_S = \theta_S + \delta_{AS} 2b_A + \delta_{AS} \frac{c_B}{c_S} \hat{z}_B \tag{9a}$$

$$\hat{z}_B = \frac{c_S}{\delta_{AS} c_B}(\hat{z}_S - \theta_S) - \frac{2b_A}{c_B} \tag{9b}$$

$$\hat{z}_A = \frac{b_A}{c_A}. \tag{9c}$$

As predicted, aggression will always reach a stable equilibrium whenever there is a cost of

aggression (Eq. 9c, Fig. 1). Eqs. 9a-b predicts a line of equilibria for signal size and body size,

because their evolutionary change is completely intertwined with the relationship $\Delta \bar{z}_B =$

$\frac{G_{SB}}{G_{SS}} \Delta \bar{z}_S$ (Fig. 1). The slope of the line of equilibria predicting mean signal size from mean

body size, and hence the evolutionary allometry of signal size, is $\delta_{AS} \frac{c_B}{c_S}$. This relationship

indicates that when comparing population means through time or across space, positive

allometry (i.e., a slope greater than unity) is predicted when the strength of behavioral

modification multiplied by the threat of the opponent ($\delta_{AS} c_B$) is greater than the strength of

natural selection on signal size (c_S). In general, when male behavior is more strongly

dependent on the signal of their opponent (i.e., when δ_{AS} is larger), more elaborate signals are

expected at equilibrium (Fig. 2).

Whether an evolving population will reach a predicted equilibrium (no runaway) or

overshoot it (runaway) also depends on the rate of evolution of body size versus natural

selection on signal size. Specifically, from Eq. 8a, for runaway evolution of signal size, body

size must evolve fast enough so that $\delta_{AS}(2b_A + c_B \bar{z}_B) > c_S(\bar{z}_S - \theta_S)$. Because b_A and θ_S are

constants, this occurs when $\delta_{AS} c_B \Delta \bar{z}_B > c_S \Delta \bar{z}_S$, or equivalently:

$\dfrac{G_{SB}}{G_{SS}} \delta_{AS} c_B > c_S.$ (10)

This result is also achievable by solving for the condition generating a negative eigenvalue of

the Jacobian of $\Delta \bar{z}$, which indicates an unstable equilibrium (Lande 1981; Bailey & Kölliker

2019).

The first term in Eq. 10 is the regression of body size on signal size, which is typically

large for weapons and signals (McCullough *et al.* 2016; Eberhard *et al.* 2018). As a regression

coefficient, this term measures the degree to which body size can be predicted from signal

size, which captures the logic of why the term measures signal honesty. In addition, Eq.10

indicates that runaway evolution of a signal is most likely to occur when three conditions

exist: the signal is honest (G_{SB} is large and positive), it modifies aggressive behavior of social

partners ($\delta_{AS} > 0$), and aggression imposes a fitness cost that increases when opponents are

larger (c_B). Fig. 1 illustrates a scenario in which the predicted outcome (equilibrium or

runaway) depends upon the value of the genetic covariance G_{SB}.

DISCUSSION

Our model provides explicit conditions for sexual selection arising from male-male

competition to result in elaborate signals and runaway evolution. We model the origin of costs

and benefits associated with male traits mediating male-male interactions using

considerations from evolutionary game theory, which allows us to derive expressions for

natural and social selection gradients that reflect the mechanistic properties of male contests.

We then incorporate these expressions for selection into a model of trait genetics based on the

interacting phenotypes framework, which accounts for the influence of indirect genetic effects

arising from interactions with an opponent. Elaboration of a signal occurs whenever males

adjust their level of aggression based on the signal of an opponent; i.e., $\delta_{AS} > 0$. This

elaboration becomes runaway evolution when the signal is honest and when the cost imposed

by aggression in an opponent increases with their body sizee (Eq. 10; Fig. 1, 2). In contrast,

aggression always reaches an equilibrium, both because the fitness benefit of aggression is

relative to that of the opponent and because of the fitness costs of escalated contests. Limits to

runaway evolution of the signaling trait depend on the strength of natural selection opposing

signal elaboration, which may arise through costs of producing or bearing the signal.

Our model does not specify the nature of the costs and benefits associated with

aggression, the signalling trait, and body size (condition). These are important variables,

likely ecologically contingent, and empirical work that quantifies these costs and benefits will

provide context for the generality of our model. However, one of the strengths of this

quantitative genetic modeling approach is that it provides predictions that are testable in

natural populations. Specifically, we expect the evolution of elaborate signaling traits that

resolve duels between males to evolve when three conditions are present. First, signals should

be reliable predictors of body size or some other proxy of fighting ability. Indeed, such signal

honesty, which is often characterized as positive allometry (McCullough *et al.* 2016; O'Brien

et al. 2018) or a positive genetic correlation between size and signal (Clark & Moore 1995;

McGlothlin *et al.* 2005; Laidre & Johnstone 2013), is a common feature of traits involved in

male-male competition. Second, males must modify their behavior in response to their

opponent's signal. We assume that males increase their aggression when encountering an

opponent with a smaller signal than their own and reduce their aggression when encountering

an opponent with a larger signal. Such adjustment is common in species that resolve contests

via limited fights or displays (Darwin 1871; West-Eberhard 1979, 1983; Emlen 2008, 2014).

In our model, this phenomenon alters the relationship between genotype and phenotype,

causing a net force of social selection to contribute to signal evolution (Eqs. 7a, 8a). Finally,

we expect social selection to be imposed via the aggression of opponents. This selection

becomes stronger as male body size or fighting ability evolves due to the threat of escalation

of fights with large opponents. Mean level of aggression need not change if the threat

escalates. Our model makes specific predictions for the signs of these gradients when

selection on signal size, body size, and aggression can all be measured (Eqs. 5-6). Most

crucially, our model predicts negative social selection gradients for both body size and

aggression, which reflect the costs of escalated contests. In populations that are experiencing

an evolutionary runaway, these gradients should become stronger as body size and signal size

coevolve. Although few studies have measured social selection gradients, the limited

evidence that exists supports the existence of negative social selection gradients imposed by

competitors (Formica *et al.* 2011; Fisher & Pruitt 2019).

Parallels to Lande's model of female choice

The results of our model are conceptually analogous to Lande's (1981) model of runaway

sexual selection via female choice, suggesting some key parallels between the processes. Both

our model and Lande's, which was the first formal model of Fisher's runaway process, result

in lines of equilibria that may be stable or unstable depending on the genetic parameters. For

the scenario of relative mate preference in Lande's model, the line of equilibria for a male

trait ($\hat{z}$) and a female preference ($\hat{y}$) can be written:

$$\hat{z} = \theta + \frac{b_{SS}}{c_{NS}}\hat{y} \qquad (11)$$

where θ is the naturally selected optimum, b_{SS} is the strength of sexual selection, and c_{NS} is

the strength of natural selection. Eq. 11 directly parallels Eq. 9a from our model and

emphasizes that in male-male competition, the force of sexual selection is provided not by

direct female choice but by male body size (or some other measure of willingness or ability to

engage in aggression). In male-male competition, the threat of the opponent (c_B) leads to

social selection, which is indirectly translated into evolutionary change in male signals via the

parameter δ_{AS}, measuring the dependence of aggression on relative signal size of two

competing males.

Similarly, the condition for runaway evolution of male traits and female preference

driven by mate choice in Lande's model can be written:

$$\frac{G_{mf}}{G_{mm}}b_{SS} > c_{NS} \qquad (12)$$

where G_{mf} represents the genetic covariance between male trait and female preference and

G_{mm} represents genetic variance of the male trait. The condition in Eq. 12 directly parallels

the condition in Eq. 10, emphasizing again that in male-male competition, $\delta_{AS}c_B$ provides the

force of social selection that indirectly leads to an evolutionary increase in male signal size.

Both types of runaway evolution are driven by genetic covariance. In mate choice, runaway is

driven by the covariance between the sexes that arises from choosy females mating with

attractive males, but in male-male competition, this effect arises directly from signal honesty,

i.e., the genetic covariance between a signaling trait and the threat it signals. Moreover, if the

mean level of aggression does not change, in male-male competition increasing costs are

associated only with the increasingly elaborated signal. This may occur when aggression is

not expressed outside of male-male competition and when limited fights settle contests

(Maynard Smith & Harper 1988; Maynard Smith & Harper 2003). These are common

conditions (Parker 1974; West-Eberhard 1983, 1984; Maynard Smith & Harper 1988;

Andersson 1994), suggesting that runaway from male-male competition may occur frequently

(McCullough *et al.* 2016; Rico-Guevara & Hurme 2019). Finally, the genetic covariance in

Lande's model arises from linkage disequilibrium that accumulates via nonrandom mating

whereas ours reflects pleiotropy between body size and signal size. Thus, the genetic

covariance driving runaway from male-male competition is likely to be much larger both

because recombination efficiently erodes linkage disequilibrium and positive allometry (e.g.,

signals and body size) is common and reflects pleiotropy. Indeed, in models of mate choice

where there is a pleiotropic relationship between direct benefits or good genes, runaway is

difficult. In our model, however, the pleotropic nature of the honest signal leads to runaway.

There are many other models of mate choice in the literature, and a full comparison to

them all is beyond the scope of this paper. Our goal here is simply to highlight potentially

common features of runaway evolution, the most important is that runaway sexual selection

by both male-male competition and female mate choice appears to be an evolving source of

selection provided by the social environment. A more expansive comparison may well

stimulate modifications or additions to the model we present here. In addition, there would be

much to gain by combining studies of female mate choice and male-male competition to

simultaneously test models of sexual selection (Hunt *et al.* 2009). This may be especially

enlightening when traits serve as both ornaments and arnaments (Berglund *et al.* 1996) or

when mate choice opposes male-male competition (Moore & Moore 1999).

Conclusion

Ritualized displays and elaborated signals associated with the potential for aggression are readily observed in nature and their importance often obvious and spectacular (Darwin 1871; Parker 1974; Maynard Smith & Harper 1988; Maynard Smith & Harper 2003; Emlen 2008, 2014). Yet the details of how these might evolve have been unclear. Previous game theory models have shown that overt aggression can be ameliorated by conventional signals (Parker 1974; Maynard Smith 1982; Maynard Smith & Harper 1988; Maynard Smith & Harper 2003; Rutte *et al.* 2006), and verbal models have proposed that signaling traits associated with male-male competition evolve exaggerated expression because social selection is intense (West-Eberhard 1979, 1983, 1984). Male-male competition may well result in intense selection (Maynard Smith & Brown 1986), as mating can be highly skewed toward one or a few males in a population (Darwin 1871; Andersson 1994; Shuster & Wade 2003), but this alone is insufficient to result in exaggerated traits. Our model shows that feedback between the behavioral and morphological traits mediating male-male competition create runaway evolution.

Sexual selection arising from male-male competition is prevalent and so the consequences of such selection important for understanding the generation of biological diversity. We hope our model stimulates empiricists in much the same way that the model of mate choice stimulated research on mating preferences. Ultimately, our understanding of the consequences of sexual selection arising from male-male competition will come from empirical research. Our hope is that this model helps direct and focus some of that research.

SUPPORTING APPENDIX

Fitness model for male-male interactions

To model the fitness consequences of an interaction between two males, we use a

modification of the classic hawk-dove game, which involves competition over a resource

(Maynard Smith & Price 1973, Maynard Smith 1982). In a traditional hawk-dove game, there

are two strategies. Hawks tend to start fights, while doves tend to flee. When two hawks meet,

a fight determines the outcome of the contest, with one hawk winning and receiving a

resource of value v, while the other hawk pays a cost c, which may be related to injury or

other costs of aggression. The average payoff for a hawk in a hawk-hawk encounter is thus

$\frac{v-c}{2}$. When two doves meet, they either divide the resource evenly or decide the contest

without aggression and its associated costs, leading to an average payoff for each dove of $\frac{v}{2}$.

When a hawk meets a dove, the hawk wins the contest, receiving a fitness payoff of v, while

the dove receives a payoff of 0, reflecting the fact that the dove neither wins the resource nor

pays a cost for being aggressive. If $z = 1$ represents playing hawk and $z = 0$ represents

playing dove, this fitness model can be written in terms of the phenotypes of two interactants

expressing traits z and z':

$$W = W_0 + \frac{v}{2}(1 + z - z') - \frac{c}{2}zz' + \varepsilon \qquad (A1)$$

Following McGlothlin et al. (2021), we generalize this equation to a continuous scale,

allowing z_A, representing aggression, to take on any non-negative value. This means that

individuals that are more aggressive (i.e., that employ a more hawkish strategy) will tend to

win interactions and that the fitness cost of being aggressive will increase as both individuals

become more aggressive (i.e., more hawkish). The fitness model in Eq. 4 differs from Eq. A1

by being expressed in terms of relative fitness (w) rather than absolute fitness (W). Thus, the

translations between the parameters in the general hawk-dove fitness model and the fitness

effects of aggression in Eq. (4) are

$w_0 = W_0 + \frac{v}{2}$ (A2a)

$b_A = \frac{v}{2}$ (A2b)

$c_A = \frac{c}{2}$ (A2c)

In words, b_A represents access to resources (here, mates) achieved by winning a contest, while

c_A represents the costs paid by the loser of a contest when both individuals are aggressive. We

also includes two additional unique costs. We assume that the cost of an encounter is

proportional to the body size (or fighting ability) of an opponent, resulting in a cost

proportional to c_B. Importantly, this cost is paid regardless of an individual's own behavior.

Second, bearing a signal larger than the naturally selected optimum imposes a cost

proportional to c_S, which is presumably paid outside the context of male-male encounters.

General equation for response to selection

Here, we develop a general model for evolution when trait expression depends upon the

difference between a focal individual's own traits and traits of another individual encountered

in the context of a social interaction. This model is directly applied to male-male contests in

the main text and may be useful for considering many other types of social interactions. First,

consider a vector of traits ($\mathbf{z}$) whose expression can be decomposed into three components: a

vector of additive genetic effects ($\mathbf{a}$), a vector of environmental effects ($\mathbf{e}$), and a social

response term that depends on the difference between the traits of the focal individual and

those of a social interactant ($\mathbf{z}'$):

$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\delta}(\mathbf{z} - \mathbf{z}').$ (A3)

The matrix $\boldsymbol{\delta}$ consists of components δ_{ij} that translate the effect of differences in trait j into

expression of trait i. Similarly, the phenotype of the social partner can be written as

$\mathbf{z}' = \mathbf{a}' + \mathbf{e}' + \boldsymbol{\delta}(\mathbf{z}' - \mathbf{z}).$ (A4)

As we show below, because the term $\boldsymbol{\delta}(\mathbf{z} - \mathbf{z}')$ in Eqs. A3-A4 contains phenotypes of both

individuals, it consists of a combination of direct and indirect genetic effects.

To calculate a response to selection for traits expressed as in Eq. A3, we first solve for

the multivariate phenotypic mean. Assuming that environmental effects have a mean of zero,

the trait mean is

$\bar{\mathbf{z}} = \bar{\mathbf{a}} + \boldsymbol{\delta}(\bar{\mathbf{z}} - \bar{\mathbf{z}}) = \bar{\mathbf{a}},$ (A5)

which means that the population trait mean will depend only on the mean additive genetic

value. The vector of total breeding values (**A**), which represents the genetic contribution to the

population mean and is used to calculate evolutionary responses to selection, is equivalent to

the vector of additive genetic effects (**a**). Next, to derive an explicit definition of the

phenotype, we first use substitution to write

$\boldsymbol{\delta}(\mathbf{z} - \mathbf{z}') = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}\boldsymbol{\delta}(\mathbf{a} + \mathbf{e} - \mathbf{a}' - \mathbf{e}')$ (A6)

where **I** is the identity matrix. After some algebra, Eq. A6 allows us to write explicit

definitions of the two phenotypes as

$\mathbf{z} = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a} + \mathbf{e}) - \boldsymbol{\delta}(\mathbf{a}' + \mathbf{e}'))$ (A7)

and

$\mathbf{z}' = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a}' + \mathbf{e}') - \boldsymbol{\delta}(\mathbf{a} + \mathbf{e})).$ (A8)

Response to selection can then be calculated following McGlothlin *et al.* (2010) as

$\Delta\bar{\mathbf{z}} = \text{Cov}(\mathbf{A}, \mathbf{z}^{\text{T}})\boldsymbol{\beta}_{\mathbf{N}} + \text{Cov}(\mathbf{A}, \mathbf{z}'^{\text{T}})\boldsymbol{\beta}_{\mathbf{S}}.$ (A9)

Substituting Eqs. A7-A8 into Eq. A9 and simplifying yields

$\Delta\bar{\mathbf{z}} = \mathbf{G}(\mathbf{I} - (1 + r)\boldsymbol{\delta}^{\text{T}})(\mathbf{I} - 2\boldsymbol{\delta}^{\text{T}})^{-1}\boldsymbol{\beta}_{\mathbf{N}} + \mathbf{G}(r\mathbf{I} - (1 + r)\boldsymbol{\delta}^{\text{T}})(\mathbf{I} - 2\boldsymbol{\delta}^{\text{T}})^{-1}\boldsymbol{\beta}_{\mathbf{S}},$ (A10)

where $\mathbf{G}$ is the additive genetic (co)variance matrix and r is relatedness. Eq. A10 can be used

to generate the specific model in the text by setting parameters as

$\mathbf{G} = \begin{bmatrix} G_{SS} & G_{SB} & 0 \\ G_{SB} & G_{BB} & 0 \\ 0 & 0 & G_{AA} \end{bmatrix}, \boldsymbol{\delta} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \delta_{AS} & 0 & 0 \end{bmatrix}, \boldsymbol{\beta}_{\mathbf{N}} = \begin{bmatrix} \beta_{N_S} \\ \beta_{N_B} \\ \beta_{N_A} \end{bmatrix}, \boldsymbol{\beta}_{\mathbf{S}} = \begin{bmatrix} \beta_{S_S} \\ \beta_{S_B} \\ \beta_{S_A} \end{bmatrix}, r = 0$ (A11)

and multiplying the result by $\frac{1}{2}$ to indicate selection acting only on males.

Relationship to indirect genetic effects models

The model described above differs from previous models of indirect genetic effects in that

phenotypes may be adjusted in relation to both the phenotypes of a social partner and other

phenotypes of the focal individual. The standard model of phenotypic expression used in trait-

based genetic effects models (Moore *et al.* 1997a) is:

$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\Psi}\mathbf{z}'.$ (A12)

The relationship between the two models can be seen by adding an additional term to the

standard model:

$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\xi}\mathbf{z} + \boldsymbol{\Psi}\mathbf{z}'.$ (A13)

The term $\boldsymbol{\xi}\mathbf{z}$, which is similar to developmental interaction effects (Wolf *et al.* 2001),

contains a conditional modification of the direct genetic effect in response to other phenotypic

traits of the same individual, while the term $\boldsymbol{\Psi}\mathbf{z}'$ contains indirect genetic effects (Moore *et*

al. 1997a). When $\boldsymbol{\xi} = \mathbf{0}$, Eq. 13 is equivalent to Eq. A12, and when $\boldsymbol{\xi} = -\boldsymbol{\Psi} = \boldsymbol{\delta}$, Eq. A13 is

equivalent to Eq. A3.

Incorporating conditional direct genetic effects is a way to mechanistically represent

genetic covariances in a quantitative genetic model (Wolf *et al.* 2001). Combining such

effects with indirect genetic effects allows exploration of a wide variety of models of

phenotypic adjustment, including behavioral modification, in evolutionary quantitative

genetic models. For full generality, we give the equation for multivariate response to selection

derived from Eq. A13. First, the vector of total breeding values derived from the trait mean is

$\mathbf{A} = (\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{a}.$ (A14)

By substituting Eqs. A13-A14 into Eq. A9,

$\Delta\bar{\mathbf{z}} = (\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{G}(\mathbf{I} + r(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}} - \boldsymbol{\Psi}^{\mathrm{T}}(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})^{-1}\boldsymbol{\beta}_{\mathbf{N}} +$

$(\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{G}(r\mathbf{I} + (\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}} - \boldsymbol{\Psi}^{\mathrm{T}}(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})^{-1}\boldsymbol{\beta}_{\mathbf{S}}.$ (A15)

This equation is unwieldy in its multivariate form, but one can follow the approach we take

here for male-male competition and use Eq. A15 to generate much simpler models of the

evolution of systems of responsive traits given assumptions about fitness and trait expression.

Extension to multiple opponents

Our results can be extended to incorporate interactions with multiple opponents. Suppose a

male encounters opponents sequentially. The phenotype of a focal individual averaged over

all encounters:

$\bar{\mathbf{z}} = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a} + \mathbf{e}) - \boldsymbol{\delta}(\bar{\mathbf{a}}' + \bar{\mathbf{e}}')),$ (A16)

and the average of his social partners:

$\bar{\mathbf{z}}' = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\bar{\mathbf{a}}' + \bar{\mathbf{e}}') - \boldsymbol{\delta}(\mathbf{a} + \mathbf{e})),$ (A17)

where an overbar is now taken to indicate an average over an individual male's encounters. If

we assume that the fitness effects of individual encounters accrue additively, then Eq. A10

may still be used to predict the evolution of the population mean, with $\boldsymbol{\beta_N}$ and $\boldsymbol{\beta_S}$ now

representing vectors of partial regression slopes of fitness on focal individual mean and social

group mean phenotypes, respectively. Specific fitness models may be substituted into Eq.

A10 in the same way as for models of single pairwise interactions.

ACKNOWLEDGMENTS

We thank our collaborators, students, and postdoctoral scientists who over the years provided discussions and comments on various earlier versions of this idea and model. This work was supported by grants from NERC (UK, NE/C510659/1 to JBW) and NSF (USA, IOS-1354358 to AJM, DEB-1457463 to JWM).

REFERENCES

Andersson, M. (1982). Female choice selects for extreme tail length in a widowbird. *Nature*, 299, 818-820.

Andersson, M. (1994). *Sexual Selection*. Princeton Univ. Press, Princeton.

Araya-Ajoy, Y.G., Westneat, D.F. & Wright, J. (2020). Pathways to social evolution and their evolutionary feedbacks. *Evolution*, 74, 1894-1907.

Arnold, S.J. (1994). Multivariate inheritance and evolution: a review of concepts. In: *Quantitative Genetic Studies of Behavioral Evolution* (ed. Boake, CR). Univ. of Chicago Press Chicago, pp. 17-48.

Bailey, N.W. & Kölliker, M. (2019). Social runaway: Fisherian elaboration (or reduction) of socially selected traits via indirect genetic effects. *Evolution*, 73, 1549-1563.

Bailey, N.W. & Moore, A.J. (2012). Runaway sexual selection without genetic correlations: social environments and flexible mate choice initiate and enhance the Fisher process. *Evolution*, 66, 2674-2684.

Berglund, A., Bisazza, A. & Pilastro, A. (1996). Armaments and ornaments: an evolutionary explanation of traits of dual utility. *Biol. J. Linn. Soc.*, 58, 385-399.

Bijma, P. & Wade, M.J. (2008). The joint effects of kin, multilevel selection and indirect genetic effects on response to genetic selection. *J. Evol. Biol.*, 21, 1175-1188.

Briceño, R.D. & Eberhard, W.G. (1995). *The functional morphology of male cerci and associated characters in 13 species of tropical earwigs (Dermaptera: Forficulidae, Labiidae, Carcinophoridae, Pygidicranidae)*. Smithsonian Institution Press, Washington, DC.

Clark, D.C. & Moore, A.J. (1995). Genetic aspects of communication during male-male
competition in the Madagascar hissing cockroach: honest signaling of size. *Heredity*,
75, 198-205.

Darwin, C. (1859). *The Origin of Species*. John Murray, London.

Darwin, C. (1871). *The Descent of Man, and Selection in Relation to Sex*. John Murray,
London.

Eberhard, W.G., Rodriguez, R.L., Huber, B.A., Speck, B., Miller, H., Buzatto, B.A. *et al.*
(2018). Sexual selection and static allometry: the importance of function. *Q. Rev.*
Biol., 93, 207-250.

Emlen, D.J. (2008). The Evolution of Animal Weapons. *Annu. Rev. Ecol. Evol. Syst.*, 39, 387-
413.

Emlen, D.J. (2014). *Animal Weapons: The Evolution of Battle*. Henry Holt & Co., New York.

Emlen, D.J., Marangelo, J., Ball, B. & Cunningham, C.W. (2005). Diversity in the weapons
of sexual selection: horn evolution in the beetle genus *Onthophagus* (Coleoptera :
Scarabaeidae). *Evolution*, 59, 1060-1084.

Fisher, D.N. & Pruitt, J.N. (2019). Opposite responses to selection and where to find them. *J.*
Evol. Biol., 32, 505-518.

Fisher, R.A. (1915). The evolution of sexual preference. *Eugen Rev*, 7, 184-192.

Fisher, R.A. (1930). *The Genetical Theory of Natural Selection*. Clarendon Press, Oxford.

Formica, V.A., McGlothlin, J.W., Wood, C.W., Augat, M.E., Butterfield, R.E., Barnard, M.E.
et al. (2011). Phenotypic assortment mediates the effect of social selection in a wild
beetle population. *Evolution*, 65, 2771-2781.

Hagelin, J.C. (2002). The kinds of traits involved in male-male competition: a comparison of
plumage, behavior, and body size in quail. *Behav. Ecol.*, 13, 32-41.

Hunt, J., Breuker, C.J., Sadowski, J.A. & Moore, A.J. (2009). Male-male competition, female
mate choice and their interaction: determining total sexual selection. *J. Evol. Biol.*, 22,
13-26.

Huxley, J.S. (1938). Darwin's theory of sexual selection and the data subsumed by it, in the
light of recent research. *Am. Nat.*, 72, 416-433.

Kirkpatrick, M. (1982). Sexual selection and the evolution of female mate choice. *Evolution*,
36, 1-12.

Laidre, M.E. & Johnstone, R.A. (2013). Animal signals. *Curr. Biol.*, 23, R829-R833.

Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution.
Evolution, 30, 314-334.

Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to
brain:body size allometry. *Evolution*, 33, 402-416.

Lande, R. (1981). Models of speciation by sexual selection on polygenic traits. *Proc. Natl.*
Acad. Sci. USA, 78, 3721-3725.

Lande, R. & Arnold, S.J. (1983). The measurement of selection on correlated characters.
Evolution, 37, 1210-1226.

Lucas, W.J. (1920). *A Monograph of the British Orthoptera*. The Ray Society, London.

Maynard Smith, J. (1982). *Evolution and the Theory of Games*. Cambridge Univ. Press,
Cambridge.

Maynard Smith, J. & Brown, R.L.W. (1986). Competition and body size. *Theor. Popul. Biol.*,
30, 166-179.

Maynard Smith, J. & Harper, D. (2003). *Animal Signals*. Oxford Univ. Press, Oxford.

Maynard Smith, J. & Harper, D.G.C. (1988). The evolution of aggression: can selection
generate variability? *Philos. Trans. R. Soc. Lond. B*, 319, 557-570.

Maynard Smith, J. & Price, G.R. (1973). The logic of animal conflict. *Nature*, 246, 15-18.

McCullough, E.L., Miller, C.W. & Emlen, D.J. (2016). Why sexually selected weapons are
not ornaments. *Trends Ecol. Evol.*, 31, 742-751.

McGlothlin, J.W., Akçay, E., Brodie, E.D., III, Moore, A.J. & Van Cleve, J. (2021). A
synthesis of evolutionary game theory and quantitative genetic models of social
evolution. *bioRxiv*, 10.1101/2021.03.27.437341.

McGlothlin, J.W., Moore, A.J., Wolf, J.B. & Brodie, E.D., III (2010). Interacting phenotypes
and the evolutionary process. III. Social evolution. *Evolution*, 64, 2558-2574.

McGlothlin, J.W., Parker, P.G., Nolan, V., Jr. & Ketterson, E.D. (2005). Correlational
selection leads to genetic integration of body size and an attractive plumage trait in
dark-eyed juncos. *Evolution*, 59, 658-671.

Mead, L.S. & Arnold, S.J. (2004). Quantitative genetic models of sexual selection. *Trends*
Ecol. Evol., 19, 264-271.

Moore, A.J., Brodie, E.D., III & Wolf, J.B. (1997a). Interacting phenotypes and the
evolutionary process: I. Direct and indirect genetic effects of social interactions
Evolution, 51, 1352-1362.

Moore, A.J., Haynes, K.F., Preziosi, R.F. & Moore, P.J. (2002). The evolution of interacting
phenotypes: genetics and evolution of social dominance. *Am. Nat.*, 160, S186-S197.

Moore, A.J. & Moore, P.J. (1999). Balancing sexual selection through opposing mate choice
and male competition. *Proc. R. Soc. Lond. B*, 266, 711-716.

Moore, A.J. & Moore, P.J. (2006). The genetics of sexual selection. In: *Evolutionary Genetics*
(eds. Fox, CW & Wolf, JB). Oxford Univ. Press Oxford, pp. 339-349.

Moore, P.J., Reagan-Wallin, N.L., Haynes, K.F. & Moore, A.J. (1997b). Odour conveys
status on cockroaches. *Nature*, 389, 25-25.

O'Brien, D.M., Allen, C.E., Van Kleeck, M.J., Hone, D., Knell, R., Knapp, A. *et al.* (2018).
On the evolution of extreme structures: static scaling and the function of sexually
selected signals. *Anim. Behav.*, 144, 95-108.

Parker, G.A. (1974). Assessment strategy and the evolution of fighting behaviour. *J. Theor.*
Biol., 47, 223-243.

Rico-Guevara, A. & Hurme, K.J. (2019). Intrasexually selected weapons. *Biol. Rev.*, 94, 60-
101.

Riechert, S.E. (1984). Games spiders play. III. Cues underlying context-associated changes in
agonistic behavior. *Anim. Behav.*, 32, 1-15.

Rohwer, S. (1982). The evolution of reliable and unreliable badges of fighting ability. *Am.*
Zool., 22, 531-546.

Rosenthal, G.G. (2017). *Mate Choice: The Evolution of Sexual Decision Making from*
Microbes to Humans. Princeton Univ. Press, Princeton.

Rutte, C., Taborsky, M. & Brinkhof, M.W.G. (2006). What sets the odds of winning and
losing? *Trends Ecol. Evol.*, 21, 16-21.

Seehausen, O. & Schluter, D. (2004). Male-male competition and nuptial-colour displacement
as a diversifying force in Lake Victoria cichlid fishes. *Proc. R. Soc. Lond. B*, 271,
1345-1353.

Shuster, S.M. & Wade, M.J. (2003). *Mating Systems and Strategies*. Princeton Univ. Press,
Princeton.

Sneddon, L.U., Huntingford, F.A. & Taylor, A.C. (1997). Weapon size versus body size as a
predictor of winning in fights between shore crabs, *Carcinus maenas* (L.). *Behav.*
Ecol. Sociobiol., 41, 237-242.

Tinghitella, R.M., Lackey, A.C.R., Martin, M., Dijkstra, P.D., Drury, J.P., Heathcote, R. *et al.*
(2018). On the role of male competition in speciation: a review and research agenda.
Behav. Ecol., 29, 783-797.

Wallace, A.R. (1889). *Darwinism*. Macmillan and Co., New York.

West-Eberhard, M.J. (1979). Sexual selection, social competition, and evolution. *Proc. Am.*
Phil. Soc., 123, 222-234.

West-Eberhard, M.J. (1983). Sexual selection, social competition, and speciation. *Q. Rev.*
Biol., 58, 155-183.

West-Eberhard, M.J. (1984). Sexual selection, competitive communication and species-
specific signals. In: *Insect Communication* (ed. Lewis, T). Academic Press New York,
pp. 283-324.

West-Eberhard, M.J. (2003). *Developmental Plasticity and Evolution*. Oxford Univ. Press,
Oxford.

Wiens, J.J. & Tuschhoff, E. (2020). Songs versus colours versus horns: what explains the
diversity of sexually selected traits? *Biol. Rev.*, 95, 847-864.

Wilkinson, G.S. & Dodson, G.N. (1997). Function and evolution of antlers and eye stalks in
flies. In: *The Evolution of Mating Systems in Insects and Arachnids* (eds. Choe, JC &
Crespi, BJ). Cambridge Univ. Press Cambridge.

Wolf, J.B. (2003). Genetic architecture and evolutionary constraint when the environment
contains genes. *Proc. Natl. Acad. Sci. USA*, 100, 4655-4660.

Wolf, J.B., Brodie, E.D., III & Moore, A.J. (1999). Interacting phenotypes and the
evolutionary process. II. Selection resulting from social interactions. *Am. Nat.*, 153,
254-266.

Wolf, J.B., Frankino, W.A., Agrawal, A.F., Brodie, E.D., III & Moore, A.J. (2001).
Developmental interactions and the constituents of quantitative variation. *Evolution*,
55, 232-245.

FIGURE 1 Evolution of a male signal, body size, and aggression in response to male-male competition. Panels A and B show evolutionary trajectories for each trait over 200 generations, and panels C and D show predicted lines of equilibria (heavy line) and their stability (gray arrows). In all panels, all three traits have the same genetic variance ($G = 1$), benefit ($b_A = 0.2$) and cost of aggression ($c_A = 0.05$), fitness cost deriving from the threat of a male's opponent ($c_B = 0.2$), cost of signal size ($c_S = 0.05$; with naturally selected optimum $\theta_S = 0$), and a responsiveness of aggression to body size ($\delta_{AS} = 0.4$). The line of equilibria is calculated from Eq. 9a using these values. In panels A and C, signal size is weakly correlated with body size ($G_{SB} = 0.4$), while in panels B and D, the two traits are more strongly correlated ($G_{SB} = 0.8$). When the genetic correlation between signal size and body size is weak, all three traits reach equilibria (A), with equilibrium aggression predicted solely by costs and benefits. Signal size and body size reach a point on the predicted line of equilibrium (C) that differs depending on their starting values. When the genetic correlation is strong, aggression still reaches an equilibrium, but signal size and body size run away together (B), overshooting the predicted line of equilibria (D). As in Fisherian selection from female mate choice (Lande 1981), male-male competition can drive traits to runaway elaboration or extinction when the line of equilibria is unstable (D).

FIGURE 2   Stronger dependence of male aggressive behavior leads to more elaborate traits at equilibrium. Panel A illustrates a relatively weak influence of opponent signal on male aggression ($\delta_{AS} = 0.4$), while panel B illustrates a stronger influence ($\delta_{AS} = 0.8$). In each panel, we use starting values for traits relevant to the highly sexually dimorphic earwig *Labidura riparia*, which uses its forceps as a signaling trait and is shown to the right of each panel (drawing modified from Lucas 1920). Other parameters are the same as Fig. 1A. When the influence of opponent signal is weak (A), both body size and signal show a moderate evolutionary increase. When the influence is stronger (B), both body size and signal increase more, but the final signal size is much larger relative to body size. The highly elaborate elongate forceps in panel B may be found in other earwig species like *Forcipula gariazzi*.