Runaway evolution from male-male competition

Allen J. Moore[a,1], Joel W. McGlothlin[b,2], and Jason B. Wolf[c,3]

[a]Department of Entomology, University of Georgia, Athens, GA 30602, USA

[b]Department of Biological Sciences, Virginia Tech, Blacksburg, VA 24061, USA

[c]Milner Centre for Evolution and Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK

[1]Correspondence may be addressed to any author. Email: ajmoore@uga.edu

[2]Email: joelmcg@vt.edu

[3]Email: j.b.wolf@bath.ac.uk

Author contributions: A.J.M., J.W.M., and J.B.W. designed research, performed research, and wrote the paper.

Keywords: aggression | quantitative genetics | runaway evolution | sexual selection | social signals

1   **Abstract**

Understanding why and how elaborated traits evolve remains a fascination and a challenge.

Darwin proposed both male-male competition and female mate choice as explanations for

elaboration because such traits are often mediators of social interactions that govern access

to mates. Although we have robust evolutionary quantitative genetic models for how mate

choice can lead to runaway evolution, we lack an equivalent framework for understanding

how male-male competition can drive extreme elaboration of traits. Here, we integrate the

logic of optimality models into the quantitative genetic framework of interacting phenotypes to

fill this gap. We assume that males modulate their aggression based on the relative size of a

trait that signals willingness and ability to fight and identify conditions where the signal

undergoes rapid and exponential evolution. Males receive fitness benefits from winning

contests, but they may accrue fitness costs due to threats imposed by their opponent. This

cost leads to a force of social selection that accelerates as the signaling trait is elaborated,

which may cause runaway evolution of the signal. Even when a runaway is checked by

natural selection, we find that signaling traits evolving by male-male competition can be

elaborated well beyond their naturally selected optimum. Our model identifies simple

conditions generating feedback between the behavioral and morphological traits mediating

male-male competition, providing clear testable predictions. We conclude that, like the well-

characterized case of female mate choice, male-male competition can provide a coevolving

source of selection that can drive a runaway process resulting in evolution of elaborate traits.

21   **Significance**

It has long been appreciated that female mate choice can drive elaboration of male displays,

but it has been less clear whether contests between males may lead to similar elaboration.

To address this fundamental question, we use a genetic model evolution via male-male

competition. We find that when a male trait is an honest indicator of aggression, it generates

selection on itself by altering the social environment. This process can cause the strength of

selection to increase as the trait becomes more elaborate, which can lead to runaway

evolution. Our results show that the key unifying feature of runaway sexual selection by both

male-male competition and female mate choice is an evolving source of selection provided

by the social environment.

Darwin (1, 2) suggested that exaggerated, spectacular, and bizarre structures and behaviors

evolve because they enhance success in male-male contests for mates or influence female's

choice of mates. However, most contemporary research directed at documenting and

understanding the processes driving such trait elaboration has focused on the role of mate

choice, largely driven by the success of the logic underlying the model of sexual selection

first proposed by Fisher (3-8). The Fisher process occurs when female preference and a

preferred male trait become genetically correlated, causing both the trait and the preference to

evolve in a self-sustaining fashion. In some cases, the Fisher process may cause sexually

selected male traits to evolve at ever-increasing speed, which has been termed an evolutionary

"runaway process" (3, 4, 9). However, as Darwin (2) noted, and has been recently emphasized

(7, 8, 10), mate choice alone is insufficient to explain the diversity of traits evolving by sexual

selection. Indeed, despite the bias in research toward female mate choice, male-male

competition is thought to be a more common source of selection shaping male traits (2, 7, 8,

10-14). Moreover, traits expressed in male-male interactions can be as elaborate as those that

are the target of female preferences (2, 7, 11-19). However, the process by which male-male

competition results in extreme elaboration remains unresolved.

Many elaborated traits function as weapons or signals of aggression mediating social

interactions among competing males outside of or in addition to their influence on female

choice (15, 17-20). These signals often act as conventions, providing information on

willingness and ability to fight (17, 18, 21-24). Such traits include showy plumage (25), color

(26), pheromones (27), and potential weapons such as antlers (28), horns (29), forceps (30),

and claws (31). A key feature is that these traits likely provide information about some

underlying qualities of the individuals, which can be used to adaptively modulate how males

behave in social interactions. For example, there is typically a positive association between

the signal or weapon and other traits such as body size (7, 19), which make the signal an

honest indicator of potential threat to an opponent. The role that these traits can play as honest

signals means that they can govern the outcome of male-male interactions, which allows such

traits to serve as both targets and sources of selection. Because social sources of selection can

themselves evolve, this may lead to a process where male-male interactions create trait

elaboration similar to the Fisherian runaway (4, 9, 32), but how this process might arise in

male-male contests has not been elucidated.

Population genetic models have been developed to understand how male-male

interactions can drive evolution and to investigate the conditions for evolutionary stability of

alternative strategies. For example, using simulations with diploid sexual inheritance at 10

unlinked loci, Maynard Smith and Brown (33) show that body size associated with aggression

has the potential to evolve indefinitely via competition, at least until alleles for large body

size become fixed. Optimality approaches to this problem have shown that systems that use

signals of potential aggression or resource holding potential to determine the outcome of

male-male contests can be evolutionary stable (21-24, 33, 34). While these models provide

essential insights into what can evolve through male-male interactions and a framework for

identifying evolutionary equilibria, they do not directly model the process by which the

fundamental feature of extreme elaboration—an evolving source of selection—arises. This

goal can be achieved by adapting the quantitative genetic "interacting phenotypes"

framework, which explicitly incorporates socially contingent trait expression and fitness into

models of the evolutionary process (9, 35-37), to build a model specific to understanding the

properties of male-male competition and how it can drive trait elaboration.

Here, we integrate the logic developed in optimality models of male-male competition

into the quantitative genetic framework of interacting phenotypes. We focus on the role of

social context in male-male competition, with the goal of exploring the potential for extreme

elaboration evolving through this process alone. We adopt the genetic framework of

interacting phenotypes because behavioral responses in contests between males are socially

contingent (38). When fitness is determined by male-male competition and when aggression

is potentially costly, males should be expected to evolve strategies wherein they modulate

their behavior in response to phenotypes of their competitors that signal willingness or ability

to fight. This socially contingent behavior should then shape how behavior generates selection

on the morphological traits that mediate the interaction. We identify the conditions where

such a system results in evolutionary elaboration of signaling traits and identify the necessary

and sufficient conditions for trait elaboration to become a runaway process. Finally, we

outline predictions that can be empirically tested to evaluate this model. We show that sexual

selection by male-male competition may have similar features to Fisherian runaway sexual

selection by female choice. Just as in female mate choice, social effects in male-male

competition may generate both genetic correlations between interacting individuals and

fitness effects that lead to self-sustaining and escalating selection, which can lead to runaway

evolution of sexually selected traits.

A Quantitative Genetic Model of Male-Male Competition

To capture the influence of the social environment in a model of male-male competition, we

assume that individuals adjust their behavior in response to the trait values of their social

partners, an assumption that is supported empirically and theoretically (8, 10, 12-14, 19, 21,

22, 36, 39, 40). Because the social context (i.e., the social environment) is constructed from

traits of conspecifics, this flexible response to social context provides the opportunity for

indirect genetic effects (36), which allow the social environment itself to evolve (35, 41-43).

Evolutionary changes in the social environment can lead to concerted evolution because the

social environment can be a source of selection on the traits that themselves compose the

social environment (12, 32, 37, 44, 45). Such "social selection" (12-14, 35, 37, 41) is

expected to arise whenever traits act as both the agents and the targets of selection and is

potentially a key to understanding how male-male competition drives the evolution of

elaborate male traits.

We assume that males use a trait (designated by the subscript S) to assess some

underlying potential threat of their opponents in social contests. Such a trait can be a weapon

or display that is used as a signal. Potential examples include traits such as a plumage badges,

exaggerated weapons, or vocal or chemical signals. Elaboration of the signal may consist of

an increase in size or complexity, although for heuristic simplicity, we discuss the evolution

of increased signal size. Males also vary in the underlying quality trait that represents an

index of their fighting ability or some other aspect of their phenotype that determines the

threat they represent to their opponent in a contest. We describe this trait as body size

(designated by the subscript B) for simplicity. As such, the signal may be genetically

correlated to body size, providing a degree of signal honesty. Males respond to the assessment

of the signal by modulating their behavior toward their opponent ("aggression," designated by

the subscript A) within the contest because the signal provides information on the quality of

their opponent, and hence provides information on the likelihood that they would win an

escalated contest (see below).

We assume that both signal size (z_S) and body size (z_B) are normally distributed

metric traits influenced by many loci of small effect. Expression of these traits can be

partitioned into heritable additive genetic effects (denoted a_S and a_B) and general non-

heritable (environmental and nonadditive genetic) effects (denoted e_S and e_B). We assume

that neither signal nor body size changes as a result of the social interaction. An individual's

total phenotypic value for each trait is then described by a simple sum of the heritable and

non-heritable components, or

$z_i = a_i + e_i,$ **[1]**

where a_i is normally distributed with mean $\bar{a}_i$ and variance G_{ii} and e_i is normally distributed

with mean 0 and variance E_{ii}. We make the standard quantitative genetic assumption that

heritable and non-heritable components are uncorrelated. Our model includes a genetic

covariance between signal size and body size (G_{SB}), which may be set to zero (indicating a

dishonest signal) or a positive value (indicating an honest signal).

For simplicity, we model the outcome of pairwise duels between males drawn at

random from the population. Our model is easily generalized to include multiple interactions

between males (see Materials and Methods). We assume that aggressive behavior (z_A) in

these male-male contests is context dependent and influenced by the relative signal size of the

competing males, as suggested by West-Eberhard (12-14). This assumption is well supported

by optimality models and empirical studies (11, 17, 18, 21, 24, 31, 46). The phenotypic value

of aggression can thus be written as

$$z_A = a_A + e_A + \delta_{AS}(z_S - z_S'), \qquad\qquad\qquad \textbf{[2]}$$

where a_A and e_A represent standard additive genetic and uncorrelated non-heritable

components, respectively, z_S is the signal of the focal individual, z_S' is the signal of the focal

individual's opponent. As for other traits in the model, a_A has mean $\bar{a}_A$ and variance G_{AA} and

e_A is normally distributed with mean 0 and variance E_{AA}. Here and elsewhere, terms with

primes indicate a value assigned to the focal individual's opponent. The coefficient δ_{AS}

measures the influence of the difference in signal size on the expression of aggressive

behavior. Thus, δ_{AS} is analogous to the ψ term in standard interacting phenotype models (36),

but differs because it depends upon the value of an interactant's phenotype relative to the

focal individual. Because signal size is heritable, the third term in Eq. **2** includes both a

modification of the direct genetic effect ($\delta_{AS}a_S$) and an indirect genetic effect ($-\delta_{AS}a_S'$),

which is defined as the effect of a social ineractant's genes on the expression of a focal

individual's phenotype (17). We describe the relationship between this model and the

standard model of indirect genetic effects in the Materials and Methods.

Because it is conditional and adjusted depending on the social context, the underlying

additive genetic value of aggression is assumed to be uncorrelated to both signal size and

body size ($G_{SA} = G_{BA} = 0$). However, the conditional expression of aggression leads males

with larger signals and/or larger body size to be more aggressive on average (and vice versa).

This can be viewed as the individual with the larger signal being more willing to escalate an

encounter, while the individual with the smaller signal tends to retreat from the interaction.

Selection Imposed by Male-Male Competition. In social interactions, associations

between traits and fitness may cause selection via two pathways, nonsocial selection

(quantified by the gradient β_N), which arises from effects of a focal individual's traits on its

own fitness, and social selection (quantified by the gradient β_S), which arises from the effects

of an opponent's traits on the fitness of a focal individual (37). When both are present,

individual relative fitness can be written as

$$w = w_0 + \mathbf{z}^T\boldsymbol{\beta_N} + \mathbf{z}'^T\boldsymbol{\beta_S} + \varepsilon, \qquad\qquad \textbf{[3]}$$

where w_0 is an intercept, $\mathbf{z}$ and $\mathbf{z}'$ are column vectors of focal and opponent traits, $\boldsymbol{\beta_N}$ and $\boldsymbol{\beta_S}$

are vectors of nonsocial and social selection gradients, ε is an uncorrelated error term, and the

superscript T denotes transposition (37). Expressing relative fitness using Eq. **3** has two

distinct advantages. First, selection gradients can be estimated in natural populations using

multiple regression (37, 47-49), allowing our model to generate testable predictions. Second,

selection gradients can be combined with genetic parameters to predict short-term

evolutionary response to selection (35, 41, 47).

While Eq. **3** provides a general framework for capturing sources of selection, to

understand the origin and properties (i.e., the magnitude and direction) of these selection

gradients, we use evolutionary game theory to write a mechanistic expression for relative

fitness based on the costs and benefits arising from male-male contests with signaling,

$$w = w_0 + b_A(z_A - z_A') - c_A z_A z_A' - c_B z_A' z_B' - c_S(\bar{z}_S - \theta_S)z_S + \varepsilon, \qquad \textbf{[4]}$$

where terms including b represent fitness benefits and terms including c represent fitness

costs. In Eq. **4**, the benefit term and the first cost term derive from the hawk-dove model of

evolutionary game theory (22, 44, 50). The coefficient b_A represents the fitness benefit of

winning a contest, which we assume derives from greater access to females. In a contest,

access to females is determined by a focal individual's aggression relative to its opponent.

Multiplying b_A by $(z_A - z_A')$ reflects the fact that the probability of winning a contest

increases as a male becomes increasingly more aggressive than its opponent. The term $c_A z_A z_A'$

represents the fitness cost of aggression associated with escalation of encounters. Logically,

an individual pays a cost for acting aggressive that depends on the level of aggression shown

by its opponent. A second fitness cost $(c_B z_A' z_B')$ reflects the fact that the fitness impact of

aggression by an opponent (z_A') depends on the size of the opponent (z_B'). This cost, which we

call the threat of the opponent, derives from the fact that larger males impose a greater risk of

harm by being aggressive than do smaller males. Finally, we assume that a third cost

$(c_S(\bar{z}_S - \theta_S)z_S)$ arises from natural selection favoring some optimal trait value (θ_S), which

therefore will oppose signal elaboration. Following a Gaussian model of selection (51, 52),

natural selection against elaborate signals becomes stronger as the population mean of the

signal $(\bar{z}_S)$ becomes further away from its naturally selected optimum (θ_S). Although we do

not do so here, this term could be replaced with a multivariate Gaussian term (52) to add

naturally selected optima for aggression and body size.

Because selection gradients are equivalent to partial regression slopes, we can use

partial derivatives of Eq. **4** with respect to focal and opponent traits (evaluated at the

population mean) to calculate nonsocial and social selection gradients (44). Taking these

derivatives, we find nonsocial selection gradients for each trait to be

$\beta_{N_S} = -c_S(\bar{z}_S - \theta_S)$ **[5a]**

$\beta_{N_B} = 0$ **[5b]**

$\beta_{N_A} = b_A - c_A\bar{z}_A$ **[5c]**

and social selection gradients to be

$\beta_{S_S} = 0$ **[6a]**

$\beta_{S_B} = -c_B\bar{z}_A$ **[6b]**

$\beta_{S_A} = -b_A - c_A\bar{z}_A - c_B\bar{z}_B.$ **[6c]**

In other words, males with large signals are selected against via nonsocial selection but

interacting with such males does not directly impose social selection. Body size is not under

direct nonsocial selection but imposes a fitness cost via social selection that increases with the

population mean of aggression. Nonsocial selection favors aggression until the benefits of

aggression are outweighed by the costs, while social selection imposed by opponent

aggression is always negative, representing a net fitness cost. This gradient becomes

increasingly negative as the population mean aggression and body size increase. Interestingly,

these selection gradients suggest that signal size itself experiences no direct sexual selection.

If signal size increases, it must do so as an indirect response to selection on a correlated trait.

Evolutionary Response to Selection. Selection within a generation is translated into an

evolutionary response across generations through the association between the phenotype,

upon which selection acts, and the genotype, which contributes to the inheritance of the traits

across generations. In quantitative genetics, this genotype-phenotype relationship is most

often summarized by the additive genetic variance, which is used to predict evolutionary

response to selection across generations (47, 53). However, for traits expressed in social

interactions, we must also consider social pathways to fitness, which arise from indirect

genetic effects and social selection, when calculating response to selection (35, 36, 41).

Because the model of phenotypic modification described in Eq. **2** deviates from the standard

model of indirect genetic effects, we develop a general equation for response to selection in

the Materials and Methods (Eq. **20**). Using this equation, the response to selection for the

three traits in our model can in general be written as

$\Delta \bar{z}_S = \frac{1}{2} G_{SS} \left(\beta_{N_S} + \delta_{AS} (\beta_{N_A} - \beta_{S_A}) \right) + \frac{1}{2} G_{SB} \beta_{N_B}$ **[7a]**

$\Delta \bar{z}_B = \frac{1}{2} G_{BB} \beta_{N_B} + \frac{1}{2} G_{SB} \left(\beta_{N_S} + \delta_{AS} (\beta_{N_A} - \beta_{S_A}) \right)$ **[7b]**

$\Delta \bar{z}_A = \frac{1}{2} G_{AA} \beta_{N_A}.$ **[7c]**

The multiplier $\frac{1}{2}$ in Eq. **7** arises because selection is acting only on males. Eq. **7a** shows that

modification of aggressive behavior in response to the signaling trait (δ_{AS}) causes both

nonsocial and social selection gradients for aggression to contribute to signal evolution. This

behavioral modification thus effectively causes both a direct genetic correlation between

signaling and aggression and a nonrandom association between the genetic effect on

aggression in the focal individual and the genetic effect on the signal in its opponent. This

behavioral modification also contributes to evolution of body size when the signal is honest

($G_{SB} > 0$; Eq. **7b**). In contrast, behavioral modification does not contribute to evolution of

aggression (Eq. **7c**).

By substituting Eqs. **5-6** into Eq. **7**, we can predict evolutionary change using our

mechanistic fitness model (Eq. **4**):

$\quad \Delta \bar{z}_S = \frac{1}{2} G_{SS} \big(\delta_{AS} (2b_A + c_B \bar{z}_B) - c_S (\bar{z}_S - \theta_S) \big)$ [8a]

$\quad \Delta \bar{z}_B = \frac{1}{2} G_{SB} \big(\delta_{AS} (2b_A + c_B \bar{z}_B) - c_S (\bar{z}_S - \theta_S) \big)$ [8b]

$\quad \Delta \bar{z}_A = \frac{1}{2} G_{AA} (b_A - c_A \bar{z}_A).$ [8c]

Eq. **8a** shows that when fitness is defined as in Eq. **4**, evolution of the signaling trait beyond

its naturally selected optimum depends crucially on behavioral modification. If males do not

change their behavior in response to the signal (i.e., if $\delta_{AS} = 0$), the population mean of the

signaling trait cannot increase. From Eq. **8b**, the evolution of body size depends on both δ_{AS}

and the presence of signal honesty (i.e., genetic covariance between signal and body size,

G_{SB}). Eqs. **8a-b** also show that evolution of the signaling trait and of male body size is

potentially open-ended, because the evolutionary response to selection for each trait becomes

stronger as the population mean body size increases. In contrast, from Eq. **8c**, the evolution of

aggression is self-limiting because selection depends on the balance of the benefits and costs

of aggression, the latter of which become more intense as mean aggression intensifies. This

observation suggests that both signal size and body size may experience runaway evolution if

the benefits of aggression and the threat of the opponent are strong enough to outweigh

natural selection against elaborate signals, whereas aggression should always tend to quickly

evolve to an equilibrium value.

$\quad\quad$ To solve for equilibrium and to explore the conditions for such a runaway, we set Eqs.

8a-c equal to zero and solve for the equilibrium mean of each trait ($\hat{z}_i$), yielding

$\quad \hat{z}_S = \theta_S + \delta_{AS} 2 b_A + \delta_{AS} \frac{c_B}{c_S} \hat{z}_B$ [9a]

$\quad \hat{z}_B = \frac{c_S}{\delta_{AS} c_B} (\hat{z}_S - \theta_S) - \frac{2 b_A}{c_B}$ [9b]

$\quad \hat{z}_A = \frac{b_A}{c_A}.$ [9c]

As predicted, aggression will always reach a stable equilibrium whenever there is a cost of

aggression (Eq. **9c**, Fig. 1). By Eqs. **9a-b**, there is a predicted line of equilibria for signal size

and body size, which occurs because by Eqs. **8a-b**, their evolutionary change is completely

intertwined with the relationship $\Delta \bar{z}_B = \frac{G_{SB}}{G_{SS}} \Delta \bar{z}_S$ (Fig. 1). The slope of the line of equilibria

predicting mean signal size from mean body size, and hence the evolutionary allometry of

signal size, is $\delta_{AS} \frac{c_B}{c_S}$. This relationship indicates that when comparing population means

through time or across space, positive allometry (i.e., a slope greater than unity) is predicted

when the strength of behavioral modification multiplied by the threat of the opponent ($\delta_{AS} c_B$)

is greater than the strength of natural selection on signal size (c_S).

Whether an evolving population will reach this line of equilibria or overshoot it also

depends crucially on the rate of evolution of body size versus natural selection on signal size.

Specifically, from Eq. **8a**, in order for runaway evolution to occur, body size must evolve fast

enough that $\delta_{AS}(2b_A + c_B \bar{z}_B) > c_S(\bar{z}_S - \theta_S)$ regardless of the population mean signal size.

Because b_A and θ_S are constants, this occurs when $\delta_{AS} c_B \Delta \bar{z}_B > c_S \Delta \bar{z}_S$, or equivalently, when

$\frac{G_{SB}}{G_{SS}} \delta_{AS} c_B > c_S.$ **[10]**

This result is also achievable by solving for the condition generating a negative eigenvalue of

the Jacobian of $\Delta \bar{\mathbf{z}}$, which indicates an unstable equilibrium (4, 32). Condition **10** indicates

that runaway evolution of a signal may occur when it is honest (G_{SB} is strong and positive),

when it modifies aggressive behavior of social partners ($\delta_{AS} > 0$), and when aggression

imposes a fitness cost that increases when opponents are larger (c_B). Fig. 1 illustrates a

scenario in which the predicted outcome (equilibrium or runaway) depends upon the value of

the degree of signal honesty (i.e., the genetic covariance G_{SB}).

Discussion

Our model provides explicit conditions for sexual selection arising from male-male competition to result in elaborate signals and runaway evolution. We model the origin of costs and benefits associated with male traits mediating male-male interactions using considerations from evolutionary game theory, which allows us to derive expressions for natural and social selection gradients that reflect the mechanistic properties of male contests. We then incorporate these expressions for selection into a model of trait genetics based on the interacting phenotypes framework, which accounts for the influence of indirect genetic effects arising from interactions with an opponent. Elaboration of a signal occurs whenever males adjust their level of aggression based on the signal of an opponent, and runaway evolution occurs when the genetic correlation between signal size and body size is strong (i.e., the signal is honest), when aggression in a social partner is modified based on the focal male signal, and when aggression is costly when the opponent is larger (Eqn. **10**; Fig 1). In contrast, aggression always reaches an equilibrium, both because the fitness benefit of aggression is relative to that of the opponent and because of the fitness costs of escalated contests. Limits to runaway evolution of the signaling trait depend on the strength of natural selection opposing signal elaboration, which may arise through costs of producing or bearing the signal.

Because our model is expressed in the language of evolutionary quantitative genetics, it provides predictions that are testable in natural populations. Specifically, we expect the evolution of elaborate signaling traits that resolve duels between males to evolve when three conditions are present. First, signals should be reliable predictors of body size or some other proxy of fighting ability. Indeed, such signal honesty, which is often characterized as positive allometry (7, 19) or a positive genetic correlation (54-56) between size and signal, is a common feature in species where male-male competition predominates. Second, males must

modify their behavior in response to their opponent's signal. We assume that males increase

their aggression when encountering an opponent with a smaller signal than their own and

reduce their aggression when encountering an opponent with a larger signal. Such adjustment

is common in species that resolve contests via limited fights or displays (2, 12, 13, 17, 18). In

our model, this phenomenon alters the relationship between genotype and phenotype, causing

a net force of social selection to contribute to signal evolution (Eqs. **7a**, **8a**). Finally, we

expect social selection to be imposed via the aggression of opponents. This selection should

become stronger as male body size or fighting ability evolves due to the threat of escalation of

fights with large opponents. Mean level of aggression need not change if the threat escalates.

Although our model could be tested by experimentally dissecting the fitness costs and

benefits of individual duels, it is perhaps most effectively tested by estimating nonsocial and

social selection gradients in wild populations (37). Our model makes specific predictions for

the signs of these gradients when selection on signal size, body size, and aggression can all be

measured (Eqs. **5-6**). Most crucially, our model predicts negative social selection gradients for

both body size and aggression, which indicate that the costs of escalated contests. In

populations that are experiencing an evolutionary runaway, these gradients should become

stronger as body size and signal size coevolve. Although few studies have measured social

selection gradients, the limited evidence that exists supports the existence of negative social

selection gradients imposed by competitors (48, 49).

Parallels to Models of Female Choice. The results of our model are conceptually

analogous to models of runaway sexual selection via female choice, suggesting some key

parallels between the processes. Both our model and Lande's (4) model of female choice

result in lines of equilibria. Under a model of relative mate preferences, the line of equilibria

for a male trait ($\bar{z}$) and a female preference ($\bar{y}$) can be written as

$$\bar{z} = \theta + \frac{b_{SS}}{c_{NS}} \bar{y} \qquad\qquad\qquad\qquad\qquad\qquad [11]$$

where θ is the natural selection optimum, b_{SS} is the strength of sexual selection, and c_{NS} is the

strength of natural selection (22). Eq. **11** directly parallels Eq. **9a** and emphasizes that, in

male-male competition, the force of sexual selection is provided not by direct female choice

but by male body size (or some other measure of willingness or ability to engage in

aggression). In male-male competition, the threat of the opponent (c_B) leads to social

selection, which is indirectly translated into evolutionary change in male signals via the

parameter δ_{AS}, measuring the dependence of aggression on relative signal size of two

competing males.

Similarly, the condition for runaway evolution of male traits and female preference,

driven by mate choice, can be written as

$$\frac{B}{G} b_{SS} > c_{NS}, \qquad\qquad\qquad\qquad\qquad\qquad [12]$$

where B represents the genetic covariance between male trait and female preference and G

represents genetic variance of the male trait (24). Condition **12** directly parallels condition **10**,

emphasizing again that in male-male competition, $\delta_{AS} c_B$ provides the force of social selection

that indirectly leads to an evolutionary increase in male signal size. Additionally, both types

of runaway evolution are driven by genetic covariance. In mate choice, the runaway effect is

driven by the covariance between the sexes that arises from choosy females mating with

attractive males, but in male-male competition, this effect arises directly from signal honesty,

i.e., the genetic covariance between a signaling trait and the threat it signals. Moreover, if

mean level of aggression does not change, in male-male competition increasing costs are

associated only with the increasingly elaborated signal. This may occur when aggression is

not expressed outside of male-male competition and when limited fights settle contests (23,

24). These may be common conditions (13-15, 21, 23), which suggests that runaway from

male-male competition may occur more frequently than runaway from mate choice (7, 40),

especially if increasing choosiness is associated with increasing fitness costs to females.

Conclusion. Ritualized displays and elaborated morphological signals associated with the

potential for aggression are readily observed in nature and their importance often obvious and

spectacular (2, 17, 18, 21, 23, 24). Yet the details of how these might evolve have been

unclear, despite the fascination of empirical research for years. Previous ESS models have

shown that overt aggression can be ameliorated by conventional signals (21-24, 34), and

verbal models have proposed that signaling traits associated with male-male competition

evolve exaggerated expression because social selection is intense (12-14). Male-male

competition may well result in intense selection (33) as mating can be highly skewed toward

one or a few males in a population (2, 15, 57), but this alone is insufficient to result in

exaggerated traits. Earlier considerations of mate choice models suggest that the potential for

runaway evolution is needed to explain the most exaggerated traits associated with this male-

male competition given that strong directional sexual selection is, presumably, easily checked

by counter-acting natural selection (6). The model we present here integrates costs and

benefits predicted by game theory and provides a genetic mechanism for the runaway

evolution of elaborated social signals associated with male-male competition. Our model

shows how simple conditions that generate feedback between the behavioral and

morphological traits mediating male-male competition create runaway evolution. Thus, the

unifying feature of runaway sexual selection by both male-male competition and female mate

choice is an evolving source of selection provided by the social environment.

Materials and Methods

General Equation for Response to Selection. Here, we develop a general model for

evolution when trait expression depends upon the difference between a focal individual's

own traits and traits of another individual encountered in the context of a social interaction.

This model is directly applied to male-male contests in the main text and may be useful for

considering many other types of social interactions. First, consider a vector of traits ($\mathbf{z}$)

whose expression can be decomposed into three components: a vector of additive genetic

effects ($\mathbf{a}$), a vector of environmental effects ($\mathbf{e}$), and a social response term that depends on

the difference between the traits of the focal individual and those of a social interactant ($\mathbf{z}'$):

$$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\delta}(\mathbf{z} - \mathbf{z}').$$ **[13]**

The matrix $\boldsymbol{\delta}$ consists of components δ_{ij} that translate the effect of differences in trait j into

expression of trait i. Similarly, the phenotype of the social partner can be written as

$$\mathbf{z} = \mathbf{a}' + \mathbf{e}' + \boldsymbol{\delta}(\mathbf{z}' - \mathbf{z}).$$ **[14]**

As we show below, because the term $\boldsymbol{\delta}(\mathbf{z} - \mathbf{z}')$ in Eqs. **13-14** contains phenotypes of both

individuals, it consists of a combination of direct and indirect genetic effects.

399   To calculate a response to selection for traits expressed as in Eq. **13**, we first solve

for the multivariate phenotypic mean. Assuming that environmental effects have a mean of

zero, the trait mean is

$$\bar{\mathbf{z}} = \bar{\mathbf{a}} + \boldsymbol{\delta}(\bar{\mathbf{z}} - \bar{\mathbf{z}}) = \bar{\mathbf{a}} \quad ,$$ **[15]**

which means that the population trait mean will depend only on the mean additive genetic

value. The vector of total breeding values ($\mathbf{A}$), which represents the genetic contribution to

the population mean and is used to calculate evolutionary responses to selection, is

equivalent to the vector of additive genetic effects ($\mathbf{a}$). Next, to derive an explicit definition of

the phenotype, we first use substitution to write

$$\boldsymbol{\delta}(\mathbf{z} - \mathbf{z}') = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}\boldsymbol{\delta}(\mathbf{a} + \mathbf{e} - \mathbf{a}' - \mathbf{e}')$$ **[16]**

where $\mathbf{I}$ is the identity matrix. After some algebra, Eq. **16** allows us to write explicit definitions

of the two phenotypes as

$$\mathbf{z} = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a} + \mathbf{e}) - \boldsymbol{\delta}(\mathbf{a}' + \mathbf{e}'))$$ **[17]**

and

$$\mathbf{z}' = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a}' + \mathbf{e}') - \boldsymbol{\delta}(\mathbf{a} + \mathbf{e})).$$ **[18]**

Response to selection can then be calculated following (35) as

$\Delta\bar{\mathbf{z}} = \text{Cov}(\mathbf{A}, \mathbf{z}^{\text{T}})\boldsymbol{\beta}_{\mathbf{N}} + \text{Cov}(\mathbf{A}, \mathbf{z}'^{\text{T}})\boldsymbol{\beta}_{\mathbf{S}}.$ **[19]**

Substituting Eqs. **17-18** into Eq. **19** and simplifying yields

$\Delta\bar{\mathbf{z}} = \mathbf{G}\big(\mathbf{I} - (1+r)\boldsymbol{\delta}^{\text{T}}\big)\big(\mathbf{I} - 2\boldsymbol{\delta}^{\text{T}}\big)^{-1}\boldsymbol{\beta}_{\mathbf{N}} + \mathbf{G}\big(r\mathbf{I} - (1+r)\boldsymbol{\delta}^{\text{T}}\big)\big(\mathbf{I} - 2\boldsymbol{\delta}^{\text{T}}\big)^{-1}\boldsymbol{\beta}_{\mathbf{S}},$ **[20]**

where $\mathbf{G}$ is the additive genetic (co)variance matrix and r is relatedness. Eq. **20** can be used
to generate the specific model in the text by setting parameters as

$\mathbf{G} = \begin{bmatrix} G_{SS} & G_{SB} & 0 \\ G_{SB} & G_{BB} & 0 \\ 0 & 0 & G_{AA} \end{bmatrix}, \boldsymbol{\delta} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \delta_{AS} & 0 & 0 \end{bmatrix}, \boldsymbol{\beta}_{\mathbf{N}} = \begin{bmatrix} \beta_{N_S} \\ \beta_{N_B} \\ \beta_{N_A} \end{bmatrix}, \boldsymbol{\beta}_{\mathbf{S}} = \begin{bmatrix} \beta_{S_S} \\ \beta_{S_B} \\ \beta_{S_A} \end{bmatrix}, r = 0$ **[21]**

and multiplying the result by $\frac{1}{2}$ to indicate selection acting only on males.

Relationship to Indirect Genetic Effects Models. The model described above differs from
previous models of indirect genetic effects in that phenotypes may be adjusted in relation to
both the phenotypes of a social partner and other phenotypes of the focal individual. The
standard model of phenotypic expression used in trait-based genetic effects models (36) is

$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\Psi}\mathbf{z}'.$ **[22]**

The relationship between the two models can be seen by adding an additional term to the
standard model:

$\mathbf{z} = \mathbf{a} + \mathbf{e} + \boldsymbol{\xi}\mathbf{z} + \boldsymbol{\Psi}\mathbf{z}'.$ **[23]**

The term $\boldsymbol{\xi}\mathbf{z}$, which is similar to the developmental interaction effects considered by Wolf *et*
al. (58), contains a conditional modification of the direct genetic effect in response to other
phenotypic traits, while the term $\boldsymbol{\Psi}\mathbf{z}'$ contains indirect genetic effects (36). When $\boldsymbol{\xi} = \mathbf{0}$,
model **23** is equivalent to model **22**, and when $\boldsymbol{\xi} = -\boldsymbol{\Psi} = \boldsymbol{\delta}$, model **23** is equivalent to model
13.
Incorporating conditional direct genetic effects is a way to mechanistically represent
genetic covariances in a quantitative genetic model (58). Combining such effects with indirect
genetic effects allows exploration of a wide variety of models of phenotypic adjustment,
including behavioral modification, in evolutionary quantitative genetic models. For full
generality, we give the equation for multivariate response to selection derived from model **23**.
First, the vector of total breeding values derived from the trait mean is

$\mathbf{A} = (\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{a}.$ [24]

By substitution of Eqs. **23-24** into Eq. **19**, we find

$\Delta\bar{\mathbf{z}} = (\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{G}\left(\mathbf{I} + r(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}}\right)\left(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}} - \boldsymbol{\Psi}^{\mathrm{T}}(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{\mathbf{N}} +$

$(\mathbf{I} - \boldsymbol{\xi} - \boldsymbol{\Psi})^{-1}\mathbf{G}(r\mathbf{I} + (\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}} - \boldsymbol{\Psi}^{\mathrm{T}}(\mathbf{I} - \boldsymbol{\xi}^{\mathrm{T}})^{-1}\boldsymbol{\Psi}^{\mathrm{T}})^{-1}\boldsymbol{\beta}_{\mathbf{S}}.$ [25]

This equation is unwieldy in its multivariate form, but one can follow the approach we take

here for male-male competition and use Eq. **25** to generate much simpler models of the

evolution of systems of responsive traits given a set of assumptions about fitness and trait

expression.

Extension to Multiple Opponents. Our results can be extended to incorporate interactions

with multiple opponents. Suppose a male encounters opponents sequentially. The phenotype

of a focal individual averaged over all encounters is then

$\bar{\mathbf{z}} = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\mathbf{a} + \mathbf{e}) - \boldsymbol{\delta}(\bar{\mathbf{a}}' + \bar{\mathbf{e}}')),$ [26]

and the average of his social partners is

$\bar{\mathbf{z}}' = (\mathbf{I} - 2\boldsymbol{\delta})^{-1}((\mathbf{I} - \boldsymbol{\delta})(\bar{\mathbf{a}}' + \bar{\mathbf{e}}') - \boldsymbol{\delta}(\mathbf{a} + \mathbf{e})),$ [27]

where an overbar is now taken to indicate an average over an individual male's encounters.

If we assume that the fitness effects of individual encounters accrue additively, then Eq. **20**

may still be used to predict the evolution of the population mean, with $\boldsymbol{\beta}_{\mathbf{N}}$ and $\boldsymbol{\beta}_{\mathbf{S}}$ now

representing vectors of partial regression slopes of fitness on focal individual mean and

social group mean phenotypes, respectively. Specific fitness models may be substituted into

Eq. **20** in the same way as for models of single pairwise interactions.

ACKNOWLEDGMENTS. We thank our collaborators, students, and postdoctoral scientists

over the years for discussions and comments on various earlier versions of the manuscript.

This work was supported by grants from NERC (UK, NE/C510659/1 to J.B.W.) and NSF

(USA, IOS-1354358 to A.J.M., DEB-1457463 to J.W.M.).

1.   C. Darwin, *The Origin of Species* (John Murray, London, 1859).

2.   C. Darwin, *The Descent of Man, and Selection in Relation to Sex* (John Murray,
London, 1871).

3.   R. A. Fisher, *The Genetical Theory of Natural Selection* (Clarendon Press, Oxford,
1930).

4.   R. Lande, Models of speciation by sexual selection on polygenic traits. *Proc. Natl.*
Acad. Sci. USA **78**, 3721-3725 (1981).

5.   M. Kirkpatrick, Sexual selection and the evolution of female mate choice. *Evolution*
36, 1-12 (1982).

6.   L. S. Mead, S. J. Arnold, Quantitative genetic models of sexual selection. *Trends*
Ecol. Evol. **19**, 264-271 (2004).

7.   E. L. McCullough, C. W. Miller, D. J. Emlen, Why sexually selected weapons are not
ornaments. *Trends Ecol. Evol.* **31**, 742-751 (2016).

8.   R. M. Tinghitella *et al.*, On the role of male competition in speciation: a review and
research agenda. *Behav. Ecol.* **29**, 783-797 (2018).

9.   N. W. Bailey, A. J. Moore, Runaway sexual selection without genetic correlations:
social environments and flexible mate choice initiate and enhance the Fisher process.
Evolution **66**, 2674-2684 (2012).

10.   J. J. Wiens, E. Tuschhoff, Songs versus colours versus horns: what explains the
diversity of sexually selected traits? *Biol. Rev.* **95**, 847-864 (2020).

11.   J. S. Huxley, Darwin's theory of sexual selection and the data subsumed by it, in the
light of recent research. *Am. Nat.* **72**, 416-433 (1938).

12.   M. J. West-Eberhard, Sexual selection, social competition, and evolution. *Proc. Am.*
Phil. Soc. **123**, 222-234 (1979).

13.   M. J. West-Eberhard, Sexual selection, social competition, and speciation. *Q. Rev.*
Biol. **58**, 155-183 (1983).

14.   M. J. West-Eberhard, "Sexual selection, competitive communication and species-
specific signals" in *Insect Communication,* T. Lewis, Ed. (Academic Press, New York,
1984), pp. 283-324.

15.   M. Andersson, *Sexual Selection* (Princeton Univ. Press, Princeton, 1994), pp. 599.

16.   A. J. Moore, P. J. Moore, "The genetics of sexual selection" in *Evolutionary Genetics,*
C. W. Fox, J. B. Wolf, Eds. (Oxford Univ. Press, Oxford, 2006), pp. 339-349.

17.   D. J. Emlen, The Evolution of Animal Weapons. *Annu. Rev. Ecol. Evol. Syst.* **39**, 387-
413 (2008).

18.   D. J. Emlen, *Animal Weapons: The Evolution of Battle* (Henry Holt & Co., New York,
2014).

19.   D. M. O'Brien *et al.*, On the evolution of extreme structures: static scaling and the
function of sexually selected signals. *Anim. Behav.* **144**, 95-108 (2018).

20.   J. Hunt, C. J. Breuker, J. A. Sadowski, A. J. Moore, Male-male competition, female
mate choice and their interaction: determining total sexual selection. *J. Evol. Biol.* **22**,
13-26 (2009).

21.   G. A. Parker, Assessment strategy and the evolution of fighting behaviour. *J. Theor.*
Biol. **47**, 223-243 (1974).

22.   J. Maynard Smith, *Evolution and the Theory of Games* (Cambridge Univ. Press,
Cambridge, 1982), pp. 226.

23.   J. Maynard Smith, D. G. C. Harper, The evolution of aggression: can selection
generate variability? *Philos. Trans. R. Soc. Lond. B* **319**, 557-570 (1988).

24.   J. Maynard Smith, D. Harper, *Animal Signals* (Oxford Univ. Press, Oxford, 2003), pp.
166.

25.   J. C. Hagelin, The kinds of traits involved in male-male competition: a comparison of
plumage, behavior, and body size in quail. *Behav. Ecol.* **13**, 32-41 (2002).

26.   O. Seehausen, D. Schluter, Male-male competition and nuptial-colour displacement
as a diversifying force in Lake Victoria cichlid fishes. *Proc. R. Soc. Lond. B* **271**,
1345-1353 (2004).

27.   P. J. Moore, N. L. Reagan-Wallin, K. F. Haynes, A. J. Moore, Odour conveys status
on cockroaches. *Nature* **389**, 25-25 (1997).

28.   G. S. Wilkinson, G. N. Dodson, "Function and evolution of antlers and eye stalks in
flies" in *The Evolution of Mating Systems in Insects and Arachnids,* J. C. Choe, B. J.
Crespi, Eds. (Cambridge Univ. Press, Cambridge, 1997).

29.   D. J. Emlen, J. Marangelo, B. Ball, C. W. Cunningham, Diversity in the weapons of
sexual selection: horn evolution in the beetle genus *Onthophagus* (Coleoptera :
Scarabaeidae). *Evolution* **59**, 1060-1084 (2005).

30.   R. D. Briceño, W. G. Eberhard, *The functional morphology of male cerci and*
associated characters in 13 species of tropical earwigs (Dermaptera: Forficulidae,
Labiidae, Carcinophoridae, Pygidicranidae), Smithsonian Contributions to Zoology
(Smithsonian Institution Press, Washington, DC, 1995).

31.   L. U. Sneddon, F. A. Huntingford, A. C. Taylor, Weapon size versus body size as a
predictor of winning in fights between shore crabs, *Carcinus maenas* (L.). *Behav.*
Ecol. Sociobiol. **41**, 237-242 (1997).

32.   N. W. Bailey, M. Kölliker, Social runaway: Fisherian elaboration (or reduction) of
socially selected traits via indirect genetic effects. *Evolution* **73**, 1549-1563 (2019).

33.   J. Maynard Smith, R. L. W. Brown, Competition and body size. *Theor. Popul. Biol.* **30**,
166-179 (1986).

34.   C. Rutte, M. Taborsky, M. W. G. Brinkhof, What sets the odds of winning and losing?
Trends Ecol. Evol. **21**, 16-21 (2006).

35.   J. W. McGlothlin, A. J. Moore, J. B. Wolf, E. D. Brodie III, Interacting phenotypes and
the evolutionary process. III. Social evolution. *Evolution* **64**, 2558-2574 (2010).

36.   A. J. Moore, E. D. Brodie III, J. B. Wolf, Interacting phenotypes and the evolutionary
process: I. Direct and indirect genetic effects of social interactions *Evolution* **51**, 1352-
1362 (1997).

37.   J. B. Wolf, E. D. Brodie III, A. J. Moore, Interacting phenotypes and the evolutionary
process. II. Selection resulting from social interactions. *Am. Nat.* **153**, 254-266 (1999).

38.   C. W. Miller, P. N. Joseph, Z. Emberts, Trade-offs between weapons and testes do
not manifest at high social densities. *J. Evol. Biol.* 10.1111/jeb.13790 (2021).

39.   M. J. West-Eberhard, *Developmental Plasticity and Evolution* (Oxford Univ. Press,
Oxford, 2003), pp. 816.

40.   A. Rico-Guevara, K. J. Hurme, Intrasexually selected weapons. *Biol. Rev.* **94**, 60-101
(2019).

41.   P. Bijma, M. J. Wade, The joint effects of kin, multilevel selection and indirect genetic
effects on response to genetic selection. *J. Evol. Biol.* **21**, 1175-1188 (2008).

42.   A. J. Moore, K. F. Haynes, R. F. Preziosi, P. J. Moore, The evolution of interacting
phenotypes: genetics and evolution of social dominance. *Am. Nat.* **160**, S186-S197
(2002).

43.   J. B. Wolf, Genetic architecture and evolutionary constraint when the environment
contains genes. *Proc. Natl. Acad. Sci. USA* **100**, 4655-4660 (2003).

44.   J. W. McGlothlin, E. Akçay, E. D. Brodie III, A. J. Moore, J. Van Cleve, A synthesis of
evolutionary game theory and quantitative genetic models of social evolution. *bioRxiv*
10.1101/2021.03.27.437341 (2021).

45.   Y. G. Araya-Ajoy, D. F. Westneat, J. Wright, Pathways to social evolution and their
evolutionary feedbacks. *Evolution* **74**, 1894-1907 (2020).

46.   S. E. Riechert, Games spiders play. III. Cues underlying context-associated changes
in agonistic behavior. *Anim. Behav.* **32**, 1-15 (1984).

47.   R. Lande, S. J. Arnold, The measurement of selection on correlated characters.
Evolution **37**, 1210-1226 (1983).

48.   V. A. Formica *et al.*, Phenotypic assortment mediates the effect of social selection in
a wild beetle population. *Evolution* **65**, 2771-2781 (2011).

49.   D. N. Fisher, J. N. Pruitt, Opposite responses to selection and where to find them. *J.*
Evol. Biol. **32**, 505-518 (2019).

50.   J. Maynard Smith, G. R. Price, The logic of animal conflict. *Nature* **246**, 15-18 (1973).

51.   R. Lande, Natural selection and random genetic drift in phenotypic evolution.
Evolution **30**, 314-334 (1976).
52.   R. Lande, Quantitative genetic analysis of multivariate evolution, applied to
brain:body size allometry. *Evolution* **33**, 402-416 (1979).
53.   S. J. Arnold, "Multivariate inheritance and evolution: a review of concepts" in
Quantitative Genetic Studies of Behavioral Evolution, C. R. Boake, Ed. (Univ. of
Chicago Press, Chicago, 1994), pp. 17-48.
54.   D. C. Clark, A. J. Moore, Genetic aspects of communication during male-male
competition in the Madagascar hissing cockroach: honest signaling of size. *Heredity*
75, 198-205 (1995).
55.   M. E. Laidre, R. A. Johnstone, Animal signals. *Curr. Biol.* **23**, R829-R833 (2013).
56.   J. W. McGlothlin, P. G. Parker, V. Nolan Jr., E. D. Ketterson, Correlational selection
leads to genetic integration of body size and an attractive plumage trait in dark-eyed
juncos. *Evolution* **59**, 658-671 (2005).
57.   S. M. Shuster, M. J. Wade, *Mating Systems and Strategies* (Princeton Univ. Press,
Princeton, 2003), pp. 520.
58.   J. B. Wolf, W. A. Frankino, A. F. Agrawal, E. D. Brodie III, A. J. Moore,
Developmental interactions and the constituents of quantitative variation. *Evolution*
55, 232-245 (2001).

Fig. 1. Evolution of a male signal, body size, and aggression in response to male-male competition. Panels *A* and *B* show evolutionary trajectories for each trait over 200 generations, and panels *C* and *D* show predicted lines of equilibria (heavy line) and their stability (gray arrows). In all panels, all three traits have the same genetic variance ($G = 1$), benefit ($b_A = 0.2$) and cost of aggression ($c_A = 0.05$), fitness cost deriving from the threat of a male's opponent ($c_B = 0.2$), cost of signal size ($c_s = 0.05$; with naturally selected optimum $\theta_S = 0$), and a responsiveness of aggression to body size ($\delta_{AS} = 0.4$). The line of equilibria is calculated from Eq. **9a** using these values. In panels *A* and *C*, signal size is weakly correlated with body size ($G_{SB} = 0.4$), while in panels *B* and *D*, the two traits are more strongly correlated ($G_{SB} = 0.8$). When the genetic correlation between signal size and body size is weak, all three traits reach equilibria (*A*), with equilibrium aggression predicted solely by costs and benefits. Signal size and body size reach a point on the predicted line of equilibrium (*C*) that differs depending on their starting values. When the genetic correlation is strong, aggression still reaches an equilibrium (*B*), but signal size and body size run away together (*B*), overshooting the predicted line of equilibria (*D*). As in Fisherian selection from female mate choice (4), male-male competition can drive traits to runaway elaboration or extinction when the line of equilibria is unstable (*D*).