On-pathway multivalency of a broadly neutralizing evolution drug candidate

Sourav Chowdhury[1], Daniel C. Zielinski[2], Christopher Dalldorf[2], Joao V Rodrigues[1], Bernhard O. Palsson[2, 3], Eugene I Shakhnovich[1*]

[1]Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA

[2]Department of Bioengineering, University of California, San Diego, La Jolla, CA, USA

[3]Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Kemitorvet, Building 220, 2800 Kongens Lyngby, Denmark

[*]Correspondence

Abstract:

A promising approach to prevent antibiotic resistance is design of multivalent antibiotics that constrain bacterial evolvability by blocking multiple escape routes. Previously we discovered that an "evolution drug" lead designed to broadly neutralize bacterial dihydrofolate reductase (DHFR) and its trimethoprim escape variants is multivalent in *E. coli* cells. In this work we identified the second target of this new lead. We analysed antibiotic induced global metabolic perturbations and deployed metabolic-network analysis and structural similarity search of the putative targets and narrowed down the search to a few enzymes. Further, combining *in vivo* and *in vitro* analyses, we identified HPPK (folK), an essential protein in bacterial folate-metabolism, as the non-DHFR target of the lead. We observed that compounds with interaction specificity against WT do not develop multivalency against related proteins. This multivalent-design strategy is effective in blocking the emergence of drug-resistant phenotypes.

Highlights:

- An evolution drug lead designed to block bacterial evolvability manifested behaviours suggesting an intra-cellular multivalent activity.
- A multi-layered metabolomic and structure guided approach unravelled the second intra-cellular target of the evolution-drug lead, showing folate-pathway inhibitions at two metabolic points.
- The on-pathway anti-folate activity is exhibited by inhibiting two essential proteins viz. DHFR and its upstream protein HPPK.
- Multivalent drug development schemes present a promising anti-evolution strategy to address the problem of "drug-resistance" by inhibiting more than one metabolic protein.

Introduction:

Antibiotic molecules are designed to selectively interact with and inhibit the action of essential bacterial proteins (Butler et al., 2013). While antibiotic drugs proved their efficacy in a multitude of applications, the emergence of resistant bacterial variants is becoming one of the biggest challenges to public health. Bacterial cells deploy a variety of complicated defence strategies to escape the action of antibiotics (Fajardo et al., 2008; Girgis et al., 2009; Shibl et al., 2001). A better understanding of the evolutionary escape strategies which bacterial cells deploy under the condition of the antibiotic selection can prove to be a valuable guide in designing drug-leads that put significant constraints on the bacterial evolvability.

A common mechanism of drug resistance involves mutations in the loci encoding the drug target protein which impair the protein-drug interaction. Combination drug-therapy aims at potentially circumventing this problem by targeting more than one intracellular molecular target. Even upon mutation in one of the target locus, combination drugs can still inhibit bacterial growth because of the inhibitory interaction with the non-mutated second target. There are multiple modalities of combination therapy and they are particularly effective in acute nosocomial infections caused by multi-drug resistant gram negative bacteria (Tamma et al., 2012). Bactrim which is a combination of Trimethoprim (TMP) and Sulfamethoxazole is another common example of combination-therapy which is used for treating a host of bacterial infections. But with more than one active drug component combination drugs present a different kind of a challenge in context to drug induced toxicity and a more complex pharmacokinetic and pharmacodynamic landscape (Tyers and Wright, 2019).

Development of multivalent drugs presents an alternative solution to this. These are essentially one molecule which can potentially interact with more than one molecular target. They present a potential avenue for deriving novel antibiotic compounds with enhanced efficacy and reduced bacterial resistance. Homodimers and oligomers of vancomycin (Griffin et al., 2003), paclitaxel (Rodrigues et al., 2003) and opioids (Portoghese, 2001) have all been reported to enhance binding relative to their constitutive monomeric components (Steinfeld et al., 2007).

Conventional therapeutic strategies essentially target the WT variant of the target protein. Drug design schemes which consider the WT variant and the most prominently appearing resistance-mutations can be promising strategies to delay the onset of resistance by constraining the most commonly accessed evolutionary escape routes.

In our previous work we deployed this strategy to block resistance to TMP (Zhang et al., 2021), a well-known antifolate antibiotic that inhibits bacterial growth by selectively blocking bacterial DHFR - a critical enzyme in bacterial one-carbon metabolism. We came up with a series of molecules which were either inhibitory to only WT variant of DHFR or WT along with three most prominently observed folA site TMP-escape mutants. We presented ` CD15-3 as an "evolution-drug" lead that significantly delayed the emergence of resistant phenotypes (Zhang et al., 2021). However, only partial growth rescue upon overexpression of DHFR potentially

indicated the presence of an additional non-DHFR intracellular target of CD15-3. This potential multivalent property puts additional constraint on bacterial evolvability and thereby makes it harder to emerge resistant phenotypes. In this respect, CD15-3 is not just a lead "evolution-drug" which delays drug resistance but is also a potential "monotherapy analogue" of combination drugs targeting more than one molecular target. To that end here we sought to discover alternative "hidden" targets of this compound.

In the current work we aim at understanding the intracellular mode of action of the new evolution drug lead to find its potential intracellular non-DHFR target. Comprehensive identification of antibiotic targets and its mechanism of action inside cells has been challenging (Silver, 2007). It is worth noting that even with traditional antibiotics with known specific molecular targets, understanding the series of intracellular molecular events and plausible off-target effects leading to bacterial growth inhibition has been a complex problem. The need for such understanding appears even more pressing as increasing numbers of novel antibiotic formulations are emerging to address the problem of drug resistance. Recent advances in systems biology tools are making intracellular drug target analysis an increasingly accessible task (Rabinowitz et al., 2011).

We deployed an integrated metabolomics guided multi-layer analysis pipeline to probe the intra-cellular mode of action and targeting of the new evolution drug CD15.3. To that end, we utilized untargeted global metabolomics and the metabolic network analysis to narrow down our search for the alternative CD15.3 target. Next, we carried out structure-based comparisons of proteins constituting the most perturbed metabolic pathway to further shortlist the plausible set of potential molecular targets. With overexpression studies of shortlisted proteins coupled with enzyme assays and microscopy, we identified and validated the intra-cellular non-DHFR target of CD15-3. This multivalent cellular targeting offers an explanation for the observed limited evolution of CD15.3 (Zhang et al., 2021) resistance and supports the potential of utilizing CD15.3 as a "monotherapy analogue" of combination drugs.

This work presents a case study for plausible strategy to develop multivalent evolution-drug leads using an integrated metabolomics and structure guided analysis pipeline which can effectively identify intra-cellular molecular targets of drug candidates.

Results:

An integrated metabolomics guided approach to discover new targets of a prospective evolution drug

We deployed an untargeted global metabolomics guided multi-layered analysis pipeline (Figure 1) to search for the intracellular mode of action of the evolution drug lead CD15-3 and determine its non-DHFR target

We analysed the metabolomics readout to identify differences in the global metabolome upon CD15-3 treatment and aimed to capture potential hotspots of metabolic perturbation. Using the findings from global metabolomics experiments as a guidance, we performed metabolic supplementation experiments and recovery assays. Altogether these experiments helped to identify which metabolic pathways are most perturbed upon CD15-3 treatment.

We then overexpressed the genes close to the points of metabolic perturbation and looked for signatures of growth recovery. We analysed these experimental data in the context of the metabolic network to narrow down the search options for the additional CD15-3 cellular target.

We carried out structural similarity analyses on the proteins constituting the most perturbed metabolic pathway and overexpressed an array of proteins based on the active-site similarity. This strategy helped to identify the potential non-DHFR target of CD15-3, which was further confirmed by carrying out imaging experiments and *in vitro* enzyme assays. We propose that this pipeline integrating multiple interconnected layers to identify the intracellular target can be potentially used to understand intracellular mode of action and molecular targets of other novel antibiotic formulations as well as other drug candidates with unknown target.

Figure 1: **Schematic representation of the integrated workflow deployed to search for the non-DHFR intracellular target of CD15-3.** *Untargeted global metabolomics identified metabolites which showed abundance differences upon CD15-3 treatment. This primary system-*

wide information was used for understanding the nature of metabolic perturbation upon CD15-3 treatment and identifying the putative non-DHFR target of CD15-3. Information from each layer is used in the subsequent layer of analysis which helped in narrowing down the potential target options and finally identifying the non-DHFR target of CD15-3

Untargeted Global Metabolomics reveals that CD15-3 induces metabolic stress with a strong impact on nucleotide metabolism

Figure 2. Major changes in the global metabolome under conditions of CD15-3 treatment. *Heat map describing the global metabolomic profile of untreated (WT) and treated (WT+CD15-3) cells. Color gradient indicates fold differences (in log scale) observed in the metabolite*

abundance levels in the CD15-3 treated set compared to the untreated WT cells. Horizontal lines in the heat map indicate metabolites. Color bar shows negative logarithm of p-values (Mann-Whitney Test) associated with abundance ratios (treated vs. control) of each metabolite. A higher intensity of the blue bar indicates more depletion while a higher intensity of the orange bar indicates more accumulation.

In a previous study, we showed that cellular overexpression of DHFR helped cells to partially recover from CD15-3-induced growth inhibition (Zhang et al., 2020). As full rescue of growth at all concentrations of CD15-3 was not observed with overexpression of DHFR, we went on to explore possible non-DHFR intracellular targets of CD15-3.

We first measured the metabolic perturbations upon CD15-3 treatment to obtain a metabolome-wide view of CD15-3 action inside the cell. To that end we carried out untargeted global metabolomics measurements to obtain the comparative global metabolome under untreated and CD15-3 treated conditions. Cells were grown in the presence and absence of CD15-3 for different lengths of time and were harvested for processing. Comparative metabolite abundances showed progressively increasing differences in cellular metabolism as the cells were exposed to CD15-3 for longer lengths of time. Figure 2 shows the differences reflected in metabolites involved in nucleotide metabolism, carbohydrate metabolism, cofactors, and peptides. As observed, a five hour exposure to CD15-3 significantly impacts the global metabolism of cells, with even higher differences being observed at 12 hours exposure.

Figure 3: Representative metabolites of nucleotide metabolism show significant fold changes upon CD15-3 treatment. *(A) Bar plot showing abundance fold differences (log scale) of selected metabolites as the cells grow in the presence of CD15-3. Metabolomics data for the treated and untreated control cells were gathered by harvesting cells at three different time points as shown in the figure. (B) Bar plot showing the metabolite abundance profile (log scale) of selected metabolites under the condition of the recovery experiment when cells pre-incubated with CD15-3 were grown in normal M9 media supplemented with 0.8g/L glucose. The fold differences in the metabolite abundances are the ratios of metabolite abundance observed in the (recovering) pre-exposed/CD15-3 treated cells to that of the naïve cells. (* indicates metabolite abundance ratios with p values ≤0.05, ** ≤0.001, *** ≤ 0.0001 as derived from a Mann-Whitney test). (C) Bar plot showing how metabolite supplementation impacts growth rate in the presence of CD15-3. Rescue from CD15-3-induced growth inhibition is observed with thymidine, serine, NAG and IMP supplementations. A fixed concentration of 200μM CD15-3 was used in these supplementation experiments. (D) Bar plot showing variation in lag-time during metabolite supplementation experiments. NAG and serine supplementation leads to a prolonged lag time compared to other metabolites.*

Thymidine, a constituent of the pyrimidine biosynthesis pathway shows a 15-fold drop in abundance after 5 hours of growth with CD15-3 treatment and a 17-fold drop after 12 hours of growth (Figure 3A). 4-aminobenzoate which is synthesized from chorismate and is an important metabolic intermediate leading to the biosynthesis of a host of crucial metabolites such as folates, shows 15-fold higher abundance at 5 hours of growth and 18-fold higher abundance at 12 hours of growth with CD15-3 treatment (Figure 3A). N10-formyl-THF, a precursor in the THF biosynthesis, showed 12-fold up regulation at 5 hours of growth with CD15-3 treatment and 15-fold higher abundance after 12 hours growth (Figure 3B). Folates are crucial for the biosynthesis of many important cellular metabolites, including glycine, methionine, formyl-methionine, thymidylate, pantothenate and purine nucleotides. Our comparative global metabolome showed significant fold differences in many of these metabolites indicating some plausible perturbation around the folate pathway and gross perturbation distributed throughout the overall nucleotide metabolism. For example, serine shows 11-fold lower abundance at 5 hours of growth and almost 20-fold lower abundance at 12 hours of growth under CD15-3 treatment (Figure3A). Likewise glycine also showed continuous cellular depletion upon CD15-3 treatment with 25-fold lower abundance after 12 hours of growth under CD15-3 treatment (Figure3A). Another interesting metabolic marker for perturbed purine metabolism is AICAR. AICAR is located upstream to serine and glycine biosynthesis steps and showed almost 8-fold higher abundance at 30 minutes of growth and 16-fold higher abundance at 5 hours of growth (Figure3A). Cellular buildup of AICAR at early stages of treatment indicates that purine metabolism gets disrupted quite early under CD15-3 treatment. UMP, a constituent of pyrimidine metabolism also showed cellular build up with 32-fold higher abundance after 12 hours of growth with treatment. Also, significant fold differences in the abundance levels of various peptides, co-factors and lipids were observed which too could be attributed to a CD15-3 induced metabolic stress response (Zhang et al., 2020). We observed significant fold differences in some metabolites constituting

carbohydrate metabolism. Pyruvate and citrate cellular build-up has been known to be metabolic stress response (Shimizu, 2014). Under CD15-3 treatment we observed 11 fold higher abundance of pyruvate and 12 fold higher abundance of citrate at 12 hours growth with CD15-3 treatment (Figure3A). Cellular buildup of citrate under CD15-3 treatment potentially indicates a possible slowdown of glycolysis and in turn energy metabolism.

Recovery experiments reveal the connection of metabolite perturbations to CD15-3 response

We went on to determine the abundance levels of the relevant metabolites during recovery following CD15-3 treatment. To determine, which metabolites display the most delayed recovery after CD15-3 treatment and hence are most impacted upon the treatment, we incubated WT cells in CD15-3 for 12 hours and subsequently transferred them to normal M9 media supplemented with 0.8g/L glucose. In a parallel control, WT cells were grown for 12 hours in M9 media and subsequently regrown in normal M9 media. Metabolite abundance levels were measured at four discrete time points (Supplementary Figure: 1) of cell harvesting. Notably, AICAR and several other metabolites that showed significant abundance changes in early stages of treatment gradually restored to normal levels during recovery (Figure3B). However, thymidine, IMP, and serine had significant fold differences until the pre-exponential phase 3. A similar trend was observed with AICAR, UMP, and N10 formyl THF, as these metabolites took a longer time to recover. N-acetylglutamate (NAG), which is a constituent of the ornithine biosynthesis pathway via the formation of N-acetyl ornithine, had significantly higher abundance upon treatment and was found to respond much earlier in the recovery experiment with abundance levels returning to normal. NAG build up also plausibly indicates some perturbation in the pyrimidine metabolism as ornithine participates in the biosynthesis of arginine, which interacts with pyrimidine metabolism through sharing carbamoyl phosphate as a precursor.

Metabolite abundance data and recovery experiments indicated multiple perturbations in nucleotide metabolism. This is consistent with the folate metabolism as intended target of CD15-3, with folate derivatives being a key cofactor in nucleotide biosynthesis. In our previous report we showed that CD15-3 interacts with DHFR and any impact on cellular DHFR would affect cellular tetrahydrofolate titer further perturbing protein and nucleotide biosynthesis (Rodrigues and Shakhnovich, 2019; Zhang et al., 2020). Further, this perturbation in nucleotide metabolism as captured by the global metabolome could suggest that CD15-3 primarily disrupts nucleotide metabolism.

Metabolic supplementation reveals a pathway-specific rescue from CD15-3 growth inhibition

The metabolite response indicates broad perturbations to metabolism across many distinct pathways. To narrow down which perturbations are most directly linked to CD15-3-dependent growth inhibition, we chose to use a subset of the identified metabolic markers for metabolic supplementation experiments to attempt a rescue from CD15-3-induced growth inhibition. To this end, wild type cells were grown in the presence of the externally supplemented metabolites under conditions of CD15-3 treatment.

In all these experiments, the external metabolite supplement concentration was kept at 0.5mM. Control experiments were performed in the absence of CD15-3 to check for intrinsic toxicity of these external metabolite supplementations (Supplementary Figure 2A). We observed that the metabolites selected for the metabolic supplementation experiments did not show toxic effect on the bacterial growth.

Also, we checked the possibility that these compounds might serve as alternate carbon sources. To that end we grew cells without CD15-3 in M9 minimal media without glucose. The growth profile was compared with bacterial growth in M9 minimal media with 0.8g/L glucose, which is the typical media composition used in the rest of the study. We observed that, apart from thymidine, no other metabolite served as an alternate carbon source (Supplementary Figure 2B). In the presence of thymidine, cells did show slow growth with a very long lag phase compared with the glucose control.

Supplementation with thymidine, NAG, serine and IMP showed growth rescue, as reflected in improved growth rates (Figure 3C) under the conditions of CD15-3 treatment. On the other hand, negligible or no effect on the growth rates were observed with external supplementation of aspartate, glycine, uridine, citrate, orotate and AMP. Thymidine and other metabolites which showed significant rescue of CD15-3 inhibition improved the growth rate to around 0.6 h^{-1} which is comparable to the growth rates observed in the WT cells in the absence of CD15-3.

It is interesting to note that metabolites whose external supplementation rescued growth rates did not significantly affect the lag time. Lag time in bacterial growth is a critical indicator of cellular adaptation in the ambient growing condition (Fridman et al., 2014). In our experimental condition a higher lag time in the presence of some external supplements and CD15-3 treatment reflects time of metabolic rewiring and adaptation in the supplement enriched growing condition. It is interesting to note that NAG and serine significantly prolonged the lag time (Figure 3D) under treatment conditions although both have shown to have positive impact in improving growth rates (Figure.1C). Thus, NAG and serine could be considered as partial rescuers from CD15-3 induced stress with improvement on only growth rate and worsening the lag time.

Figure 4 Effect of nutrient supplements on CD15-3-treated cells indicate broad inhibition of folate pathway. *A) Purine biosynthesis pathway. Both IMP and AMP supplements are downstream of folate-requiring steps. However, AMP is unable to be converted to dGTP due to an irreversible step in the pathway, while IMP can be converted to both dATP and dGTP. B) Pyrimidine biosynthesis pathway. The growth rescuing supplement, thymidine, enters the pathway downstream of the folate requiring step. Supplements that enter upstream of the folate requiring step, namely orotate and uridine, did not rescue growth. C) Serine and glycine biosynthesis. Serine, which produces the charged folate form 5,10-methylene tetrahydrofolate when converted to glycine, rescued growth, while glycine itself did not rescue growth.*

Metabolic pathway analysis of nutrient supplement rescue experiments

Metabolic supplementation demonstrated that diverse compounds were able to rescue inhibition by CD15-3, while other metabolites had no effect. To better understand the metabolic rationale for the supplementation rescue patterns of CD15-3, we utilized the most updated metabolic network reconstruction of *E. coli*, iML1515(Monk et al., 2017). We examined the trends in metabolite supplementation rescue experiments with respect to their location in corresponding metabolic pathways. We observed that the effectiveness of the supplement in rescuing growth was determined by the position of the supplement in the metabolic network viz a viz folate metabolism. Specifically, supplements that have the potential to mitigate folate deficiency, namely thymidine, IMP, and serine, were effective at rescuing growth. Supplementation with Thymidine and IMP bypass the folate-dependent step in pyrimidine and purine biosynthesis, respectively (Figure 4 A,B), while serine can contribute to folate production through its conversion to glycine (Figure 4C). Meanwhile, supplements that do not mitigate a folate deficiency do not rescue growth, even though these metabolites may be adjacent in the network to successful supplements, such as uridine, AMP, and glycine. Uridine is a pyrimidine precursor that is upstream of the folate-dependent biosynthetic step. Glycine has an unclear role, as metabolism through the glycine cleavage chain produces folate, while conversion to serine consumes folate. Thus, although DHFR may not be the primary target of CD15-3, the inhibitory activity of CD15-3 still appears to primarily work through folate limitation.

Figure 5: Protein structural similarity evidence supports other folate enzymes as possible targets of CD15-3. *A) Overview of workflow for structural analysis. B) Overview of folate pathway and protein structures extracted from PDB. C) Results for whole structure similarity analysis utilizing FATCAT. Low p-values reported by FATCAT indicate more similar structures. D) Results for active site analysis using isomif for comparison. Protein structure identifiers are taken from PDB.*

Structural analysis of possible alternate binding targets

Experimental evidence from metabolomics, supplementation, all suggested that folate perturbation is the primary mode of action of CD15-3. Thus, we hypothesized that the alternate target of CD15-3 also lies within the same metabolic pathway as DHFR. Notably, other enzymes in the folate biosynthetic pathway are also known to be drug targets, such as Dihydropteroate synthase, the target of sulfamethoxazole. To narrow down likely alternative targets of CD15-3, we employed a genome-scale reconstruction of enzyme protein structures, termed the GEM-PRO or Genome-scale Model with Protein Structures of *E. coli* (Monk et al., 2017) (Figure 5A, Figure

5B). Utilizing the FATCAT algorithm for structural alignment (Li et al., 2020), we found that DHFR did not have a high degree of structural similarity to any particular enzyme in the folate pathway based on whole chain alignment (Figure5C). However, protein structure analysis comparing the binding pocket of the intended target DHFR and other enzymes in the folate biosynthetic pathway via Tanimoto coefficients suggested possible alternative binding targets for CD15-3 (Figure 5D, Methods). Notable proteins with high binding pocket similarity scores included several enzymes involved in folate biosynthesis and folate interconversion, such as MTHFC, HPPK2, and DHPS. Thus, we decided to screen several candidate upstream enzymes in the folate biosynthetic pathway whose binding pockets have high similarity to DHFR's. To that end we evaluated whether overexpression of any of these enzymes rescues growth inhibition by CD15.3

Figure 6: Overexpression of folK (encoding HPPK) rescued cells from CD15-3 induced growth inhibition. (A) Overexpression was induced with 0.1% arabinose and folK was

expressed under pBAD promoter. folK rescued growth rate at all CD15-3 concentrations. (B) Overexpression of folA (encoding DHFR) was induced at 0.005% arabinose. folA overexpression had only partial effect in recovering growth rate in CD15-3 treated cells.(C) Distribution of cell lengths as derived from DIC imaging of WT and WT cells overexpressing folK under control and CD15-3 treatment conditions. Untreated WT and cells overexpressing folK had comparable cell lengths with median cell lengths of 1.06 μm. (n.s. indicates the distribution of cell lengths were not significantly different, Mann-Whitney test, p-value =0.952). Both WT and cells overexpressing folK were subjected to CD15-3 treatment at IC$_{50}$ concentrations. WT treated cells had a filamentous morphology and the median cell length (2.41 μm) double of that of the untreated WT set. WT cells overexpressing folK after CD15-3 treatment had a median cell length of 1.252 μm which is slightly higher than that of untreated set. However, the cell size distribution of the cells overexpressing folK showed much less change after CD15-3 treatment compared to that observed for the WT (* indicates the distribution of cell lengths were significantly different, Mann-Whitney test, p-value <0.001). (D) Inhibition of HPPK (encoded by folK) by CD15-3 in an *in vitro* assay (see text and Methods for details). Residual activity of HPPK at various concentrations of CD15-3 Inset shows the bar plot for the chemiluminescence signals at various concentrations of CD15-3 in the HPPK assay buffer.

Gene overexpression experiments indicate the plausible CD15-3 target inside cell

Regulated overexpression of candidate proteins viz., HPPK (encoded by gene folK), DHPS (encoded by gene folP), DHFS (encoded by gene folC), MTHFC (encoded by gene folD), MTHFR (encoded by gene metF) and ADCL (encoded by gene pabC) using the pBAD promoter with 0.1% arabinose induction was carried out to determine whether any of the overexpressed genes show recovery from CD15-3 induced growth inhibition and thus may be the non-DHFR target of CD15-3. In our previous study (Zhang et al., 2020) we showed that overexpression of folA (encoding DHFR) partially rescued CD15-3 induced toxicity at lower concentrations of CD15.3. However folA overexpression from pBAD was restricted to 0.005% arabinose induction to avoid overexpression induced toxicity (Bhattacharyya et al., 2016) which would complicate interpretation of the rescue experiments.

Of the assessed proteins only overexpression of folK showed clear sign of rescue from growth inhibition (Fig. 6A). folK encodes for 6-Hydroxymethyl-7,8-dihydropterin pyrophosphokinase (HPPK). HPPK is a key enzyme in the folate biosynthesis pathway catalyzing the pyrophosphoryl transfer from ATP to 6-hydroxymethyl-7,8-dihydropterin, is an attractive target for developing novel antimicrobial agents (Bermingham and Derrick, 2002; Chhabra et al., 2013; Chhabra et al., 2012). Upon folK overexpression cells did not show any change in growth rates in a broad range of concentrations of CD15-3 (Figure 6A). It stays close to 0.6 h^{-1} at all the concentrations of CD15-3, which is the typical growth rate of WT in the absence of CD15-3. On the other hand, overexpression of folA at 0.005% arabinose induction showed only partial rescue from CD15-3 inhibition (Figure 6B). Rescue in growth rate was more pronounced at lower CD15-3 concentration.

We further over-expressed folP (encoding DHPS) (Supplementary Figure 3A), folC (encoding DHFS) (Supplementary Figure 3B) and folD (encoding MTHFC) (Supplementary Figure 3C) to see any possible rescue effect on growth rates. Only at lower concentrations of CD15-3 (<50µM) overexpression of these genes partially rescued growth inhibition. Overexpression of folD (Supplementary Figure 3C) showed slight improvement in growth rate at mid concentration of CD15-3 (around 70 µM). *E. coli* folD gene encodes for a bifunctional enzyme having both methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities. The dehydrogenase and cyclohydrolase reversibly catalyze oxidation of N5,N10-methylenetetrahydrofolate to N5,N10-methenyltetrahydrofolate and the hydrolysis of N5,N10-methenyltetrahydrofolate to N10-formyltetrahydrofolate and play critical role in purine metabolism.

As an additional control, we overexpressed two more genes and found no recovery effect. The metF gene encodes for methylene THF-reductase (MTHFR). Overexpression of metF did not show any rescue in CD15-3 induced growth inhibition (Supplementary Figure 3D). Next, we overexpressed gene pabC which encodes for ADCL. pabC encoding aminodeoxychorismate lyase is involved in the biosynthesis of p-aminobenzoate (PABA) which is a precursor of tetrahydrofolate. ADCL converts 4-amino-4-deoxychorismate into 4-aminobenzoate (PABA) and pyruvate. Overexpression of pabC did not show any recovery of growth rates in CD15-3 treated cells (Supplementary Figure 3E).

Of all the candidate genes, only folK overexpression showed a clear rescue effect, with growth approaching WT level of around 0.6 hour^{-1} at all concentrations of CD15-3. This indicates strongly that CD15-3 interacts with cellular HPPK as its non-DHFR molecular target. The complete growth recovery observed with folK overexpression indicates that the growth-limiting folate perturbation originates in the folK mediated step, which in turn impacts rest of the folate pathway. In the Supporting Text (Supporting Information) we provide an explanation as to why folA overexpression leads only to partial rescue while overexpression of folK resulted in full recovery from CD15-3 induced inhibition. In short, the reason for the difference is in different expression levels of folA and folK in overexpression experiment. The Supporting Text presented quantitative analysis of the effects of inhibition of two proteins by a common inhibitor and the competing factors of inhibition of both enzymes and sequestration of the inhibitor by an overexpressed enzyme. Supplementary Figure 4 summarizes the effects of various metabolic supplements used and gene overexpression strategies deployed and show how CD15-3 has a folate-related mechanism of action.

Perturbation in the folate pathway leads to cellular filamentation and concomitant morphological changes (Ahmad et al., 1998; Bershtein et al., 2015; Bhattacharyya et al., 2021; Justice et al., 2008; Sangurdekar et al., 2011; Zaritsky et al., 2006). WT cells treated and untreated with CD15-3 were grown for 4 hours and subjected to DIC imaging. CD15-3 treated cells (Supplementary Figure 5B) showed a considerable extent of cellular filamentation as compared to untreated WT (Supplementary Figure 5A) cells grown for the same length of time.

A similar experiment was also done with WT cells overexpressing folK under pBAD promoter with 0.1% arabinose induction. WT cell sets overexpressing folK were grown for 4 hours under

control and CD15-3 treatment conditions. WT cells overexpressing folK did not show any visible change in cellular shape and size (Supplementary Figure 5D) compared to the untreated control (Supplementary Figure 5D). Upon comparison of the median cell lengths (Figure 6C), a slightly higher median cell length was observed in the folK overexpressing cells with CD15-3 treatment (median cell length = 1.252µm), as compared to untreated cells under control conditions (median cell length = 1.06µm). This slightly higher median cell length could be attributed to the fact that CD15-3 also targets cellular DHFR. Overexpression of folK although mostly reverses the effects of CD15-3 on cell shape; The overall pronounced rescue in cell length upon overexpressing folK further supports the conclusion that HPPK is the non-DHFR cellular target of CD15-3.

In vitro assay shows inhibition of HPPK (folK) by CD15-3

Next, we aimed to verify in an *in vitro* assay that CD15-3 indeed inhibits HPPK in a concentration dependent manner.

To that end we performed a KinaseGlo^{TM} assay to test for HPPK activity and its probable inhibition in presence of CD15-3. We derived HPPK by 1000 fold dilution of the cell lysates overexpressing HPPK under pBAD promoter at 0.1% arabinose induction. KinaseGlo^{TM} assay is based on chemiluminescence (Tanega et al., 2009). With higher ATP concentration in the assay buffer, luciferase leads to the conversion of beetle luciferin to oxy-luciferin with the emission of light. The HPPK mediated reaction utilizes ATP leading to ATP depletion and hence drop in the chemiluminescence signal. Any potential inhibition of HPPK would retain original concentration of ATP keeping the chemiluminescence signal intact or like the control (with no HPPK activity). We observed a marked drop in the absolute chemiluminescence signal intensity in the HPPK reaction set with no CD15-3 in the assay buffer (Figure 6D inset). Interestingly presence of CD15-3 led to enhanced absolute chemiluminescence signal intensity (Figure 6D inset) suggesting that CD15-3 does inhibit HPPK. The inhibitory effect appeared to be progressively higher upon increase of the CD15-3 concentration in the reaction-assay-buffer. For control we performed similar experiments with 400 µM of CD15-3 with one ATP dependent protein Adk and one ATP independent protein BSA to validate that the drop in signal we observe in presence of CD15-3 in the HPPK reaction-assay set is due to the specific inhibitory interaction between CD15-3 and HPPK (Supplementary Figure 6). We did not observe any drop in luminescence with BSA or ADK reaction set in the presence of CD15-3.

In our assay protocol, HPPK reaction was initiated with the introduction of the substrate, 6-Hydroxymethyl-7,8-dihydropterin in the assay buffer (Supplementary Figure 6). We used the absolute chemiluminescence intensity values to calculate % activity using the following relation:

$$\% \ Activity = \frac{Signal \ at \ inhibitor \ concentration - Signal \ at \ no \ reaction}{Signal \ at \ zero \ inhibitor \ concentration - signal \ at \ no \ reaction} \ X \ 100$$

where signal at inhibitor concentration is the chemiluminescence signal at any given concentration of CD15-3, signal at no reaction is the optical signal obtained when the substrate is not added and the reaction is not initiated.

Using this relation and plotting the absolute signal intensity values observed across the CD15-3 concentration gradient we found that HPPK retains 50% activity at 36.46 μM (IC$_{50}$) (Figure 6D) showing that HPPK is indeed an inhibitor of folk.

Figure 7: Antibacterial compounds specific only to WT DHFR do not exhibit multivalency with respect to other *E. coli* proteins. (A) Heatmap showing the in vitro interaction/inhibition profile of CD series compounds. The in vitro kinetic assay of DHFR catalytic activity was used to screen inhibitors against WT DHFR and three single mutants resistant to TMP (P21L, A26T and L28R). (B) CD20 was found to inhibit the growth of E.coli cells in M9 media (black trace) with an IC$_{50}$ of 36.4 μM (inset). Overexpression was induced with 0.005% arabinose and folA (encoding WT DHFR) was expressed under pBAD promoter. Recovery of growth rate was found to occur at all the CD20 working concentration (red trace). However CD20 was not found to inhibit the growth of E.coli cells with L28R variant of DHFR (blue trace). (C) Overexpression of folA (encoding WT DHFR) was induced at 0.005% arabinose. folA overexpression was found to have full recovery effect on growth inhibition for TMP treated cells (red trace).

Multivalency of CD15-3 stems from its design for promiscuity against resistant variants of DHFR

We hypothesized that multivalency of CD15-3 in *E. coli* is a side effect of its original design to simultaneously inhibit several TMP-resistant variants of DHFR (Zhang et al., 2021). To that end we assessed whether compounds of our designed CD series that specifically inhibit only WT DHFR would show similar multivalency in *E. coli* cells.

To assess multivalency we checked if inhibition by a CD series compound that inhibits only WT forms of DHFR can be fully rescued upon overexpression of folA gene alone. As a test case we chose CD20 from our CD series. As we reported in our previous work (Zhang et al., 2021) compound CD20 has similar scaffold to that of TMP and was not selected for our subsequent studies, including short term laboratory evolution experiment due to its lack of potency against the L28R variant of DHFR—one of the most prominently occurring escape mutants under TMP selection. As shown in Figure 7A CD20 has no in vitro inhibition effect on L28R in contrast to CD15 series which are potent inhibitors of all 3 resistant variants of DHFR, in addition to WT.

We performed growth assays and allowed *E.coli* cells to grow under the CD20 treatment using a concentration gradient. CD20 was found to inhibit bacterial growth (Figure 7B, black trace) with an IC_{50} of 36.4 µM (Figure 7B inset) but did not inhibit the growth of cells with L28R variant of DHFR (Figure 7B, blue trace). Interestingly overexpression of folA (encoding WT DHFR) led to full recovery from growth inhibition, as was observed over all the concentrations of CD20 tested (Figure 7B, red trace). We went on to carry out the same analysis with TMP,. Same protocols as used for CD20, were followed to allow the cells to grow in presence of TMP. Overexpression of folA (under pBAD promoter) was induced using 0.005% arabinose. As with CD20 folA overexpression relieves the TMP stress and recovers the cells from growth inhibition. These data support our hypothesis that multivalency of CD15-3 as an evolution drug is potentially linked to its ability to broadly neutralize escape variants of DHFR.

TMP in spite of its interaction specificity and potency is known to cause rapid emergence of resistance phenotypes, primarily by target modification (Oz et al., 2014) (Toprak et al., 2012) (Tamer et al., 2019). On the other hand evolution drug leads like CD15-3 lack TMP-like interaction specificity due to the intrinsic design to interact with modified active sites of DHFR However promiscuity of CD15-3 with respect to DHFR variants and alternative targets in *E. coli* allows to dramatically delay the onset of resistance phenotype (Zhang et al., 2021). For monovalent compounds like TMP or CD20 the first line of defence for bacterial cells is to fix promoter mutations leading to increased gene dosage with ensuing higher abundance of functional DHFR. Cellular overexpression induced by arabinose is a proxy for that. On the other hand, for multivalent leads like CD15-3 with more than one cellular target this is a not a viable drug-escape strategy. Indeed in our short term laboratory evolution experiment with CD15-3 we never encountered promoter mutations in any evolutionary trajectory (Zhang et al., 2021). Rather, a marginally resistant phenotype emerged by a more generic mechanism which involved duplication of a genomic segment having around 40 percent efflux genes (Zhang et al., 2021).

Discussion

Increasing incidences of antibiotic resistance present a major challenge to the existing pipelines of antibiotic design and synthesis. The complex "intrinsic-resistome" with constituent genes of diverse phylogenetic origins helps evolving bacterial cells adapt to a wide array of otherwise unfavourable ecological conditions and evade different types of selection pressures including antibiotic induced selection (Baquero et al., 2011). Thus, addressing the problem of antibiotic resistance or its more severe manifestation of "multi-drug resistance" is a complex problem and hence needs to consider the potential diversities in escape strategies which bacterial cells deploy. This is largely overlooked by the traditional drug discovery ventures.

In our earlier study we discovered an evolution-drug lead CD15-3 which markedly delays the emergence of antibiotic resistant bacterial phenotypes (Zhang et al., 2020). CD15-3 was designed to interact with WT as well as TMP resistant mutants which arise under TMP selection. Thus CD15-3 potentially blocks the evolutionary escape routes which *E. coli* traverses under TMP. Further, our experiments showed that CD15-3 has a non-DHFR intra-cellular target.

Blocking two cellular targets severely constraints evolution of antibiotic resistant *E. coli* phenotypes. Hence, it was important to determine the alternative target of CD15-3 in *E. coli* cells to better understand the mechanism of action of the new "evolution-drug" candidate inside bacterial cells.

A systematic understanding of the intra-cellular targeting is critical in modern drug discovery. Recent advances in untargeted global metabolomics are helping to quantitatively identify the metabolic cascades impacted by perturbations (Vincent et al., 2016; Wu et al., 2015; Zampieri et al., 2017). In the present study we deployed a series of global metabolomics guided methods to search for the cellular target of the novel "evolution-drug" lead CD15-3.

Using comparative global metabolomics and metabolic network analysis combined with structural comparison of folate pathway proteins, gene overexpression experiments, microscopy and *in vitro* enzyme assay we found HPPK as the non-DHFR target of CD15-3. Interestingly being an essential protein bacterial HPPK has been an attractive target for antibiotics design (Shaw et al., 2014). Unlike DHFR overexpression which only partially rescued CD15-3 induced growth inhibition (Zhang et al., 2021), HPPK overexpression showed clear recovery at all concentrations of CD15-3. In the Appendix we provide a plausible explanation as to why HPPK overexpression rescues bacterial growth in full compared to partial rescue observed with DHFR overexpression.

We showed how integrating multiscale systems-level approaches can reveal novel molecular targets of emerging lead antibiotics. Systematic exploration of system level approaches for drug target discovery and understanding their intracellular mechanism of action has potential broad application in pharmacology.

As CD15-3 was designed to interact with several variants of active site of DHFR (Zhang et al., 2021) and had to accommodate the active site conformational differences of four DHFR variants (including WT) it resulted in promiscuity in interaction. Inhibition of cellular HPPK with structurally similar (to DHFR) active site was the side effect of that. On the other hand, CD-20 with similar scaffold to that of TMP was specific to WT and did not show *in vitro* or *in vivo* interaction with escape variant L28R. Our growth experiment also suggested that CD-20 inhibits only WT with no inhibitory effect on the L28R folA variant and a full growth rescue was achieved with the overexpression of folA (encoding WT DHFR). Interestingly same behaviour was found with TMP inhibition where a full growth recovery was observed with the over-expression of folA. Overexpression experiments are a proxy for resistance mutations at the promoter site which lead to higher cellular abundance of target protein and relieve the action of the drug (Toprak et al., 2012). Thus, there appears to be a trade-off between specificity of drugs and their ability to constraint bacterial evolvability. Drug lead like CD15-3 with its ability to interact with modified active site of the target protein is multivalent by design. The by-product of single target multivalency is interaction promiscuity with other proteins on the folate pathway whose active site is structurally similar to that of DHFR resulting in concomitant inhibition of un-intentional target proteins.

With HPPK and DHFR variants as its intracellular targets, CD15-3 can be considered as the multivalent drug-lead, i.e., a "monotherapy-analog" of "combination therapy". We did not find any specific point mutations in either CD15-3 target folA (encoding DHFR) or folK (encoding HPPK) loci in short-term evolution experiment (Zhang et al., 2020). Future research could highlight similarities and differences between evolutionary escape routes involving multivalent monotherapy "evolution-drug" leads like CD15-3 and conventional combination therapies to gain better understanding of relative efficacy of these two therapeutic approaches in constraining evolution of resistance.

Unlike previous attempts to design poly/multi-valent drugs which were used in the development of anti-VRE(vancomycin resistant enterococci) antibiotics, multivalent influenza inhibitors, anti-HIV agents, antagonists of cholera toxin, and antitumor drugs we suggest an evolution guided design scheme (Song et al., 2013; Sun, 2007; Yuan et al., 2017; Zhang et al., 2002). Future studies in this direction should consider the negative design and screen the host cell proteome (for proteins with similar active site conformations) to avoid or minimize unintended interactions with host cellular proteins. Another exciting possibility would be to extend this approach to other critical bacterial pathways and check its efficacy on constraining bacterial evolvability. We propose the development of evolution informed multivalent antibiotic as a potential direction to supress the emergence of bacterial drug-resistant phenotypes.

Material and Methods

Antibacterial Growth Measurements and IC_{50} values

Bacterial cultures in M9 minimal medium were grown overnight at 37 °C and were then normalized to an OD of 0.1 using fresh medium. A new normalization to an OD=0.1 was conducted after additional secondary growth for ~4 hours. Thereafter the M9 medium and six different concentrations of the CD15-3 in the 96-well plates (1/5 dilution) were incubated. The incubation of the plates was performed at 37 °C and the orbital shacking and absorbance measurements at 600 nm were taken every 30 min during 15 h. Growth rate was calculated using logistic fitting on matlab.

Metabolomics analysis

Untargeted global metabolomics was performed to understand the global metabolome of the WT treated and control sets under different experimental contexts. In all the experiments cells were grown in M9 minimal media supplemented with 0.8g/L glucose in a 250 mL flask and temperature of 37^0C was maintained. Cells were pelleted with a brief 2 minutes pre-centrifugation incubation step on dry ice. After pelleting the cell pellets were mixed with 80% pre-chilled methanol Samples were thereafter vortexed and incubated in dry ice for 10 min followed by centrifugation at 4 °C for 10 min at maximum speed. The supernatant was collected, and the pellet was repeatedly processed by resorting to the above-mentioned procedures. Samples were stored at −80°C until analyzed by mass spectrometry. In all our experiments at least three independent biological replicates were analyzed. LC-MS analysis in the positive and

negative mode was performed as previously described (Bhattacharyya et al., 2016). A list of 48 experimentally measured retention times was used for initial calibration of the retention time predictions. We performed data analysis for untargeted metabolomics using the software the packages MzMatch (Scheltema et al., 2011) and IDEOM (Creek et al., 2012). In untargeted analysis for peak assignment IDEOM include both positive and negative peak M/Z values and predicted retention times calculated based on chemical descriptors. We followed the same method of analysis as we had in one of our earlier studies (Rodrigues and Shakhnovich, 2019).

Recovery Experiments

WT BW25113 cells were grown both in absence and presence of prospective compound CD15-3 in M9 media supplemented with $0.8gL^{-1}$ glucose. Both the naïve (which were not exposed to CD15-3 treatment) and pre-exposed cells (cells treated with CD15-3) were subjected to growth in normal M9 media and growth profiles were analyzed. The entire pre-exponential stage (lag phase) was grouped into equalt time frames and both naïve and pre-exposed cells were harvested for metabolomic studies. Same process was also executed for harvesting both naïve and pre-exposed cells at late log phase.

Overexpression Experiments

WT cells were transformed with blank vector plasmids (without inserts) as well as plasmids overexpressing genes viz. thyA, glyA, metF,purH, purC,folD,purD,adk and folK. The genes were under pBAD-promoter and the overexpression of the genes was induced using externally supplemented arabinose (0.1%).

Reporter metabolites analysis of metabolomics data

To analyze hotspots of metabolite concentration change, we utilized the reporter metabolites algorithm (Patil and Nielsen, 2005) implemented in MATLAB in the COBRA toolbox version 3.0 (Heirendt et al., 2019) and the latest metabolic network reconstruction of E. coli, termed iML1515 (Monk et al., 2017). The reporter metabolites method was modified to utilize metabolite concentration data as an input rather than gene expression data by transposing the stoichiometric matrix of the metabolic model. Default options were used for the reporterMets function. The resulting scores were reported for each reaction in the model.

Protein structural comparison

Proteins were selected that were nearby in metabolic maps in the pathway of interest as well as some unrelated metabolic proteins. Isomif was used to pull different binding clefts from each of the selected proteins and the 5 largest of each protein binding clefts were kept. The similarity (Tanimoto coefficient) between the clefts of the folA protein structure (PDB ID: 1DRA) and each of the selected clefts of the proteins using isomif which was then plotted to view which binding clefts were most similar. FATCAT was used to compare the overall structural similarity between all of the selected proteins and the results were plotted using a clustermap to look for similar proteins. Visualizations of the proteins were created using nglview.

Statistical and Python plots:

Statistical analyses of the data and their representation was carried out using Origin pro 8.1 package. Metabolomics data were processed using R based MS converting operation and IDEOM tool. For calibrating M/Z (mass to charge) values and retention times of the standard metabolites, XCalibur package was used. For the quantitative depictions of the metabolomics data statistically validated outputs were plotted using Python libraries of matplotlib and seaborn.

HPPK activity assay

HPPK activity assay was conducted with the help of KinaseGlo[TM] assay kit. In this assay, firefly luciferase utilizes the ATP remaining after HPPK catalysis to produce a luminescence signal that is directly proportional to ATP concentration; from this, the HPPK activity can be derived. The enzyme activity calculation and selection of optimum concentration was done following previously published methods (Chhabra et al., 2012). For kinetic measurements, an optimized HPPK concentration of 7 ng/50 µL assay volume was determined, which allowed for monitoring the first 10% of reactions turnover in a reasonable assay time period (20 min).

Measurements were performed in 96-well plates using assay buffer (100 mMTris-HCl/10 mM MgCl$_2$, pH 8.5, 0.01% (w/v) BSA, 0.01% (v/v) Tween 20 and 10 mM β-mercaptoethanol). Typically, 5 µl of test compound (dissolved in 50% DMSO) and 20 µl of enzyme were added to each well followed by 25 µl of assay buffer giving 0.3 µM pterin and 0.2 µM ATP in a total reaction volume 50 µl. After a 20 minute incubation at room temperature, the enzymatic reaction was stopped with 50 µl of KinaseGlo™ reagent. Luminescence was recorded after a further 10 min using the plate reader (Tecan Infinite M200 Pro).

Differential interference contrast (DIC)

WT cells were grown in M9 media supplemented with 0.8gL^{-1} glucose and casamino acids (mixtures of all amino acids except tryptophan) in absence and presence of CD15-3 at 42^{0}C for incubation and 300 rpm constant shaking. Drop in DHFR activity has been associated with cellular filamentation and similar phenotype is observed under TMP treatment(Bhattacharyya et al., 2021). Since CD15-3 targets intracellular DHFR and soluble fraction of cellular DHFR is lower at 42 degrees C we chose this temperature for our imaging studies (Bershtein et al., 2012).

Aliquots were taken from the growing culture and they were drop casted on agar bed/blocks. These blocks were taken further processed for differential inference contrast (DIC) imaging using Zeis Discovery imaging workstation. Multiple fields were observed and scanned for a single condition type and a minimum of three replicates were used for imaging studies. Similar methods for imaging were used for WT cell types overexpressing folK under conditions of absence and presence of CD15-3 compound. Intellesis Module was used for analyzing DIC images. On average, around 500 cells were analyzed for computing cell length. *E. coli* cell lengths in our imaging studies were not normally distributed. Nonparametric Mann-Whitney test was therefore used to determine if the cell length distributions were significantly different upon CD15-3 treatment.

Appendix: The case when 2 enzymes are inhibited by a common inhibitor.

Here we consider the situation when two enzymes are inhibited by common inhibitor. We use the following notations:

x_0 total concentration of the inhibitor (i.e. free + bound)

x_{free} concentration of free inhibitor in solution

E_1^0 total concentration of enzyme 1 in the cell

E_1^{free} concentration of free (unbound) enzyme 1 in the cell

$[E_1 x]$ concentration of bound enzyme 1-inhibitor complex in the cell

E_2^0 total concentration of enzyme 2 in the cell

E_2^{free} concentration of free (unbound) enzyme 2 in the cell

$[E_2 x]$ concentration of bound enzyme 2-inhibitor comnplex in the cell

K_{d1}, K_{d2} binding affinities of enzyme 1 and 2 respectively to the inhibitor

Using Law of Mass Action and conservation of mass for all three components (enzymes 1 and 2 and the inhibitor) we get for both enzymes:

$$[E_1 x] = \frac{E_1^{free} x_{free}}{K_{d1}}$$

$$[E_2 x] = \frac{E_2^{free} x_{free}}{K_{d2}}$$

$$[E_1 x] + [E_2 x] + x_{free} = x_0$$

$$[E_1 x] + E_1^{free} = E_1^0$$

$$[E_2 x] + E_2^{free} = E_2^0$$

$$E_1^0 - E_1^{free} = \frac{E_1^{free} x_{free}}{K_{d1}}; \quad E_1^{free} = \frac{E_1^0}{1 + \dfrac{x_{free}}{K_{d1}}}; \quad [E_1 x] = \frac{E_1^0 x_{free}}{x_{free} + K_{d1}}$$

$$E_2^0 - E_2^{free} = \frac{E_2^{free} x_{free}}{K_{d2}}; \quad E_2^{free} = \frac{E_2^0}{1 + \dfrac{x_{free}}{K_{d2}}}; \quad [E_2 x] = \frac{E_2^0 x_{free}}{x_{free} + K_{d2}}$$

$$\frac{E_1^0 x_{free}}{x_{free} + K_{d1}} + \frac{E_2^0 x_{free}}{x_{free} + K_{d2}} + x_{free} = x^0$$

$$x_{free}^3 + x_{free}^2 \left(E_1^0 + E_2^0 + K_{d1} + K_{d2} - x_0 \right) + x_{free} \left(E_1^0 K_{d2} + E_2^0 K_{d1} + K_{d1} K_{d2} - x_0 \left(K_{d1} + K_{d2} \right) \right) - x_0 K_{d1} K_{d2} = 0$$

Now consider the overexpression case $E_1^0 \gg E_2^0$; $K_{d1} \approx K_{d2}$ i.e. $\dfrac{E_1^0 x_{free}}{x_{free} + K_{d1}} \gg \dfrac{E_2^0 x_{free}}{x_{free} + K_{d2}}$

This inequality means that most of the inhibitor is bound to enzyme 1 (naturally as it is overexpressed). In this case the cubic equation for free concentration becomes quadratic because we neglect in the first approximation the inhibitor bound to enzyme 2

$$E_1^0 x_{free} + x_{free}^2 + K_{d1} x_{free} = x_0 x_{free} + K_{d1} x_0$$

$$x_{free}^2 + x_{free}\left(E_1^0 + K_{d1} - x_0\right) - K_{d1} x_0 = 0$$

$$x_{free} = \frac{-\left(E_1^0 + K_{d1} - x_0\right) + \sqrt{\left(E_1^0 + K_{d1} - x_0\right)^2 + 4 K_{d1} x_0}}{2}$$

assume high overexpression

$$E_1^0 \gg x_0, K_{d1}$$

$$x_{free} \approx \frac{x_0 K_{d1}}{E_1^0}$$

$$E_2^{free} = \frac{E_2^0}{1 + \dfrac{x_0}{E_1^0}\dfrac{K_{d1}}{K_{d2}}}$$

Thus the effect of overexpression of one enzyme affects the way how another enzyme is inhibited by the same inhibitor in the positive way: the more one enzyme is overexpressed the less other enzyme is inhibited because the inhibitor is sequestered into a binding complex with the overexpressed enzyme. However, with respect to increased expression of the inhibitor we can see 2 regimes:

when overexpression is high $\dfrac{K_{d2} x_0}{K_{d1} E_1^0} \ll 1$.

In this case $E_2^{free} = \dfrac{E_2^0}{1 + \dfrac{x_0 K_{d1}}{E_1^0 K_{d2}}} \approx E_2^0 \left(1 - \dfrac{x_0}{E_1^0}\dfrac{K_{d1}}{K_{d2}}\right) \approx E_2^0$

i.e. fitness does not depend on the inhibitor concentration because the overexpressed enzyme sequesters most of the inhibitor and the other target enzyme can function uninhibited.

In the opposite case of high concentration of the inhibitor relatively to the overexpressed enzyme or when binding affinities differ strongly between two enzymes, residual inhibitor can inhibit second (not overexpressed) enzyme.

This scenario corresponds to the case $\dfrac{K_{d2}x_0}{K_{d1}E_1^0} \square\ 1$ i.e. when there is plenty of inhibitor - more than overexpressed enzyme can bind - the approximation is different and the result is

$$x_{free} \approx x_0 - E_1^0$$

$$E_2^{free} = \frac{E_2^0}{1 + \dfrac{x_0 - E_1^0}{K_{d2}}} \approx \frac{E_2^0}{\dfrac{x_0 - E_1^0}{K_{d2}}}$$

We conclude that there can be an asymmetry between the effect of overexpression of HPPK and DHFR. If HPPK is highly overexpressed such that it sequesters most of the inhibitor, then fitness – concentration of the inhibitor curve will be flat. If DHFR is less expressed from the plasmid, i.e. at lower concentration of arabinose as in our experiment (0.005% for folA vs 0.1% for HPPK) then upon addition of inhibitor the inhibitor-growth will be flat at lower concentration of the inhibitor followed by loss of fitness at higher concentration of the inhibitor due to inhibition of HPPK by remaining inhibitor. Inhibition of growth occurs in this case when concentration of the inhibitor becomes greater or equal to concentration of the overexpressed protein. E.g. 1000 fold overexpression of DHFR equals to about 100 micromoles concentration in the cell – roughly similar to concentration of CD15.3 in the experiments.

References:

Ahmad, S., Kirk, S., and Eisenstark, A. (1998). Thymine metabolism and thymineless death in prokaryotes and eukaryotes. Annual review of microbiology *52*, 591-625.

Baquero, F., Coque, T.M., and De La Cruz, F. (2011). Ecology and evolution as targets: the need for novel eco-evo drugs and strategies to fight antibiotic resistance. Antimicrobial agents and chemotherapy *55*, 3649-3660.

Bermingham, A., and Derrick, J.P. (2002). The folic acid biosynthesis pathway in bacteria: evaluation of potential for antibacterial drug discovery. Bioessays *24*, 637-648.

Bershtein, S., Choi, J.M., Bhattacharyya, S., Budnik, B., and Shakhnovich, E. (2015). Systems-level response to point mutations in a core metabolic enzyme modulates genotype-phenotype relationship. Cell reports *11*, 645-656.

Bershtein, S., Mu, W., and Shakhnovich, E.I. (2012). Soluble oligomerization provides a beneficial fitness effect on destabilizing mutations. Proceedings of the National Academy of Sciences *109*, 4857-4862.

Bhattacharyya, S., Bershtein, S., Adkar, B.V., Woodard, J., and Shakhnovich, E.I. (2021). Metabolic response to point mutations reveals principles of modulation of in vivo enzyme activity and phenotype. Molecular systems biology *in press*, 738823.

Bhattacharyya, S., Bershtein, S., Yan, J., Argun, T., Gilson, A.I., Trauger, S.A., and Shakhnovich, E.I. (2016). Transient protein-protein interactions perturb E. coli metabolome and cause gene dosage toxicity. Elife *5*, e20309.

Butler, M.S., Blaskovich, M.A., and Cooper, M.A. (2013). Antibiotics in the clinical pipeline in 2013. The Journal of antibiotics *66*, 571-591.

Chhabra, S., Barlow, N., Dolezal, O., Hattarki, M.K., Newman, J., Peat, T.S., Graham, B., and Swarbrick, J.D. (2013). Exploring the chemical space around 8-mercaptoguanine as a route to new inhibitors of the folate biosynthesis enzyme HPPK. PloS one *8*, e59535.

Chhabra, S., Dolezal, O., Collins, B.M., Newman, J., Simpson, J.S., Macreadie, I.G., Fernley, R., Peat, T.S., and Swarbrick, J.D. (2012). Structure of S. aureus HPPK and the discovery of a new substrate site inhibitor. PloS one *7*, e29444.

Creek, D.J., Jankevics, A., Burgess, K.E., Breitling, R., and Barrett, M.P. (2012). IDEOM: an Excel interface for analysis of LC–MS-based metabolomics data. Bioinformatics *28*, 1048-1049.

Fajardo, A., Martinez-Martin, N., Mercadillo, M., Galán, J.C., Ghysels, B., Matthijs, S., Cornelis, P., Wiehlmann, L., Tümmler, B., and Baquero, F. (2008). The neglected intrinsic resistome of bacterial pathogens. PloS one *3*, e1619.

Fridman, O., Goldberg, A., Ronin, I., Shoresh, N., and Balaban, N.Q. (2014). Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. Nature *513*, 418-421.

Girgis, H.S., Hottes, A.K., and Tavazoie, S. (2009). Genetic architecture of intrinsic antibiotic susceptibility. PloS one *4*, e5629.

Griffin, J.H., Linsell, M.S., Nodwell, M.B., Chen, Q., Pace, J.L., Quast, K.L., Krause, K.M., Farrington, L., Wu, T.X., and Higgins, D.L. (2003). Multivalent drug design. Synthesis and in vitro analysis of an array of vancomycin dimers. Journal of the American Chemical Society *125*, 6517-6531.

Heirendt, L., Arreckx, S., Pfau, T., Mendoza, S.N., Richelle, A., Heinken, A., Haraldsdóttir, H.S., Wachowiak, J., Keating, S.M., and Vlasov, V. (2019). Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v. 3.0. Nature protocols *14*, 639-702.

Justice, S.S., Hunstad, D.A., Cegelski, L., and Hultgren, S.J. (2008). Morphological plasticity as a bacterial survival strategy. Nature Reviews Microbiology *6*, 162-168.

Li, Z., Jaroszewski, L., Iyer, M., Sedova, M., and Godzik, A. (2020). FATCAT 2.0: towards a better understanding of the structural diversity of proteins. Nucleic acids research.

Monk, J.M., Lloyd, C.J., Brunk, E., Mih, N., Sastry, A., King, Z., Takeuchi, R., Nomura, W., Zhang, Z., and Mori, H. (2017). i ML1515, a knowledgebase that computes Escherichia coli traits. Nature biotechnology *35*, 904-908.

Oz, T., Guvenek, A., Yildiz, S., Karaboga, E., Tamer, Y.T., Mumcuyan, N., Ozan, V.B., Senturk, G.H., Cokol, M., and Yeh, P. (2014). Strength of selection pressure is an important parameter contributing to the complexity of antibiotic resistance evolution. Molecular biology and evolution *31*, 2387-2401.

Patil, K.R., and Nielsen, J. (2005). Uncovering transcriptional regulation of metabolism by using metabolic network topology. Proceedings of the national academy of sciences *102*, 2685-2689.

Portoghese, P.S. (2001). From models to molecules: opioid receptor dimers, bivalent ligands, and selective opioid receptor probes (ACS Publications).

Rabinowitz, J., Purdy, J., Vastag, L., Shenk, T., and Koyuncu, E. (2011). Metabolomics in drug target discovery. Paper presented at: Cold Spring Harbor symposia on quantitative biology (Cold Spring Harbor Laboratory Press).

Rodrigues, J.V., and Shakhnovich, E.I. (2019). Adaptation to mutational inactivation of an essential gene converges to an accessible suboptimal fitness peak. Elife *8*, e50509.

Rodrigues, P.C., Scheuermann, K., Stockmar, C., Maier, G., Fiebig, H.H., Unger, C., Mülhaupt, R., and Kratz, F. (2003). Synthesis and in vitro efficacy of acid-sensitive poly (ethylene glycol) paclitaxel conjugates. Bioorganic & medicinal chemistry letters *13*, 355-360.

Sangurdekar, D.P., Zhang, Z., and Khodursky, A.B. (2011). The association of DNA damage response and nucleotide level modulation with the antibacterial mechanism of the anti-folate drug trimethoprim. BMC genomics *12*, 583.

Scheltema, R.A., Jankevics, A., Jansen, R.C., Swertz, M.A., and Breitling, R. (2011). PeakML/mzMatch: a file format, Java library, R library, and tool-chain for mass spectrometry data analysis. Analytical chemistry *83*, 2786-2793.

Shaw, G.X., Li, Y., Shi, G., Wu, Y., Cherry, S., Needle, D., Zhang, D., Tropea, J.E., Waugh, D.S., and Yan, H. (2014). Structural enzymology and inhibition of the bi-functional folate pathway enzyme HPPK–DHPS from the biowarfare agent F rancisella tularensis. The FEBS journal *281*, 4123-4137.

Shibl, A., Memish, Z., and Osoba, A. (2001). Antibiotic resistance in developing countries. Journal of chemotherapy *13*, 40-44.

Shimizu, K. (2014). Regulation systems of bacteria such as Escherichia coli in response to nutrient limitation and environmental stresses. Metabolites *4*, 1-35.

Silver, L.L. (2007). Multi-targeting by monotherapeutic antibacterials. Nature Reviews Drug Discovery *6*, 41-55.

Song, Y.n., Zhan, P., Li, X., Rai, D., De Clercq, E., and Liu, X. (2013). Multivalent agents: a novel concept and preliminary practice in Anti-HIV drug discovery. Current medicinal chemistry *20*, 815-832.

Steinfeld, T., Mammen, M., Smith, J.A., Wilson, R.D., and Jasper, J.R. (2007). A novel multivalent ligand that bridges the allosteric and orthosteric binding sites of the M2 muscarinic receptor. Molecular pharmacology *72*, 291-302.

Sun, X.-L. (2007). Recent anti-influenza strategies in multivalent sialyloligosaccharides and sialylmimetics approaches. Current medicinal chemistry *14*, 2304-2313.

Tamer, Y.T., Gaszek, I.K., Abdizadeh, H., Batur, T.A., Reynolds, K.A., Atilgan, A.R., Atilgan, C., and Toprak, E. (2019). High-order epistasis in catalytic power of dihydrofolate reductase gives rise to a rugged fitness landscape in the presence of trimethoprim selection. Molecular biology and evolution *36*, 1533-1550.

Tamma, P.D., Cosgrove, S.E., and Maragakis, L.L. (2012). Combination therapy for treatment of infections with gram-negative bacteria. Clinical microbiology reviews *25*, 450-470.

Tanega, C., Shen, M., Mott, B.T., Thomas, C.J., MacArthur, R., Inglese, J., and Auld, D.S. (2009). Comparison of bioluminescent kinase assays using substrate depletion and product formation. Assay and drug development technologies *7*, 606-614.

Toprak, E., Veres, A., Michel, J.-B., Chait, R., Hartl, D.L., and Kishony, R. (2012). Evolutionary paths to antibiotic resistance under dynamically sustained drug selection. Nature genetics *44*, 101-105.

Tyers, M., and Wright, G.D. (2019). Drug combinations: a strategy to extend the life of antibiotics in the 21st century. Nature Reviews Microbiology *17*, 141-155.

Vincent, I.M., Ehmann, D.E., Mills, S.D., Perros, M., and Barrett, M.P. (2016). Untargeted metabolomics to ascertain antibiotic modes of action. Antimicrobial agents and chemotherapy *60*, 2281-2291.

Wu, C., Kim, H.K., van Wezel, G.P., and Choi, Y.H. (2015). Metabolomics in the natural products field–a gateway to novel antibiotics. Drug Discovery Today: Technologies *13*, 11-17.

Yuan, H., Jiang, W., Von Roemeling, C.A., Qie, Y., Liu, X., Chen, Y., Wang, Y., Wharen, R.E., Yun, K., and Bu, G. (2017). Multivalent bi-specific nanobioconjugate engager for targeted cancer immunotherapy. Nature nanotechnology *12*, 763-769.

Zampieri, M., Zimmermann, M., Claassen, M., and Sauer, U. (2017). Nontargeted metabolomics reveals the multilevel response to antibiotic perturbations. Cell reports *19*, 1214-1228.

Zaritsky, A., Woldringh, C.L., Einav, M., and Alexeeva, S. (2006). Use of thymine limitation and thymine starvation to study bacterial physiology and cytology. Journal of bacteriology *188*, 1667-1679.

Zhang, Y., Choudhury, S., Rodrigues, J., and Shakhnovich, E. (2020). Development of antibacterial compounds that block evolutionary pathways to resistance. arXiv preprint arXiv:201101098.

Zhang, Y., Chowdhury, S., Rodrigues, J.V., and Shakhnovich, E. (2021). Development of antibacterial compounds that constrain evolutionary pathways to resistance. Elife *10*, e64518.

Zhang, Z., Merritt, E.A., Ahn, M., Roach, C., Hou, Z., Verlinde, C.L., Hol, W.G., and Fan, E. (2002). Solution and crystallographic studies of branched multivalent ligands that inhibit the receptor-binding of cholera toxin. Journal of the American Chemical Society *124*, 12991-12998.

Supplementary figures:

Supplementary Figure 1: Growth curves for WT (naïve in red) and WT pre-exposed cells (CD15.3 recovered in black). PE1, PE2 and PE3 are respectively pre-exponential stages 1, 2 and 3 of growth curve. Stages (PE1, PE2, PE3 and Late Log) marked in red are for naïve WT cells and those marked in black refer to pre-exposed WT. For the recovery assay experiments, naïve and CD15.3 pre-exposed cells were harvested at their respective pre-exponential and late log phases of growth.

Supplementary Figure 2: Metabolites selected for external supplementation experiments did not show any toxic effect on the growth profiles of the cells. (A) Bar plot showing growth rates of WT cells grown in the presence of metabolites as external supplements in M9 media. (B) Growth curve shows that these external metabolites did not show any potential to function as an alternate C-source in the growth media.

Supplementary Figure 3: Overexpression of (A) folP (encoding DHPS) (B) folC (encoding DHFS) and (C) fol D (encoding MTHFC) showed slight improvement in the growth rates of the CD15-3 treated cells only at lower concentrations (<50 μM) of CD15-3. folP, folC and folD genes were overexpressed at 0.1% arabinose induction under pBAD promoter. (D) Overexpression of metF (encoding MTHFR) under pBAD promoter with 0.1% arabinose induction did not lead to recovery from CD15.3 induced growth inhibition. (E) Overexpression of ADCL (encoded by gene pabC) under pBAD promoter with 0.1% arabinose induction was found to have no recovery effect on CD15.3 treated cells

Supplementary Figure 4: Summary of selected supplement and overexpression evidence for a folate-related mechanism of action of CD15. A) De novo folate biosynthesis. Three enzymes were overexpressed. Overexpression of the intended targeted, folA, partially restored growth. Overexpression of pabC (adcl) did not restore growth. B) Purine biosynthesis. IMP and AMP metabolic supplementation was done. IMP supplementation rescued growth while AMP had no effect in rescuing CD15.3 induced growth inhibition. C) Pyrimidine biosynthesis. Thymidine supplementation led to growth rescue from CD15.3 induced inhibition and Orotate and uridine had no effect in recovery. D) Serine biosynthesis. Metabolic supplementation of serine resulted in growth rescue. E) Folate interconversion.

Supplementary Figure 5: Overexpression of folK rescues cells from CD15-3 induced morphological changes. DIC image of WT cells under (A) control (no CD15-3 treatment) and (B) treated (CD15-3 treatment) conditions. CD15-3 treated cells shows visible signs of cellular filamentation. (C) DIC image of WT E. coli cells overexpressing folK under control (no CD15-3 treatment) and (D) CD15-3 treated condition. Unlike WT cells (B) showing cellular filamentation in presence of CD15-3, folK overexpressing cells upon CD15-3 treatment did not manifest visible signs of filamentation.

Supplementary Figure 6: Bar plot showing the range of chemiluminescence signal at zero substrate (no reaction) condition and the end of reaction. Absolute chemiluminescence signal intensity as observed under different control conditions. 1000 fold diluted cell pellets (same as is used for folK overexpressing cells) obtained from cells not induced with 0.1% arabinose did not lead to any drop in the chemiluminescence signal. Similar readout was also found with BSA (a non-ATP using protein) and Adk (an ATP dependent protein) in presence of CD15-3 in the assay buffer.

Acknowledgement: This work is supported by NIH RO1 068670 (to E.S.). We are grateful to Amir Bitran for helping in the omics data analysis, Sanchari Bhattachrya for providing some of the plasmid constructs and Bharat V. Adkar for his help in the *in vitro* enzyme assay optimization.