Succession comprises a sequence of threshold-induced community assembly

processes towards multidiversity

Maximilian Hanusch[1], Xie He[1], Victoria Ruiz-Hernández[1], Robert R. Junker[1,2*]

[1]Department of Biosciences, Paris Lodron University Salzburg, 5020 Salzburg, Austria

[2]Evolutionary Ecology of Plants, Department of Biology, Philipps-University Marburg, 35043

Marburg, Germany

*Corresponding author: robert.junker@uni-marburg.de

Keywords: succession, community assembly, ecological threshold, multidiversity, regime-

shift, multi-betadiversity, biotic interactions

Word count: 4,946; Abstract: 151; Introduction: 1069; Materials and Methods: 1,845; Results:

914; Discussion: 967

Number of References: 50; Number of Figures: 4; Supporting information: 5 Tables.

Abstract

Research on successions and community assembly both address the same processes such as dispersal, species sorting, and biotic interactions but lack unifying concepts. Recent theoretical advances integrated both research lines proposing a sequence of stochastic and niche-based processes along successional gradients. Shifts in assembly processes are predicted to occur abruptly once abiotic and biotic factors dominate over dispersal as main driver. Considering the multidiversity composed of five organismal groups including plants, animals, and microbes, we empirically show that stochastic dispersal-dominated community assembly is replaced by environmental filters and biotic interactions after around 60 years of succession in a glacier forefield. The niche-based character of later successional processes is further supported by a decline in multi-beta-diversity. Our results may update concepts of community assembly by considering multiple taxa, help to bridge the gap between research on successions and community assembly, and provide new insights into the emergence of multidiverse and complex ecosystems.

Introduction

Succession and community assembly research are neighboring fields in ecology, each with a long history of constructive debates and a large body of literature. Both research lines are founded on overlapping concepts considering processes such as dispersal, environmental filtering, biotic interactions, and stochasticity as structuring elements of local communities and ecosystems. Nevertheless, succession and community assembly research focus on different spatial and temporal scales, which may have hindered a mutual exchange of ideas (Chang and HilleRisLambers, 2016). Succession research relates to the initial development of ecosystems and communities over time. Community assembly studies, on the other hand, try to elucidate the past drivers of local and recent diversity patterns regardless of temporal components. Attempts to synthesize the two fields led to the development of an integrated conceptual framework of succession and community assembly dynamics. Here, drivers of community structure are considered from an explicitly temporal perspective, assuming a temporal sequence of assembly processes shifting from dispersal to niche-based and interaction-mediated processes along the course of succession (Chang and HilleRisLambers, 2016). The proposed framework also emphasizes the importance of threshold dynamics, predicting that gradual changes in the abiotic and biotic environment are followed by rapid shifts in the predominant processes that determine community assembly (Suding and Hobbs, 2009; Kadowaki *et al.*, 2018). These shifts are expected to be associated with various community level responses such as changes in diversity, species composition and functionality (Scheffer *et al.*, 2001; Suding and Hobbs, 2009; Chang and HilleRisLambers, 2016). Recent research suggests that sudden ecosystem shifts are not restricted to changes in species composition of single organismal groups but that local diversity patterns emerge from on-site interactions between multiple taxa (Lu and Hedin, 2019; Berdugo *et al.*, 2020). Multidiversity, an aggregate measure of biodiversity that integrates the standardized diversities of multiple taxa (Allan *et al.*, 2014), has been shown to reflect ecosystem functionality (Delgado-Baquerizo *et al.*, 2020), species composition (Felipe-Lucia *et al.*, 2020), and multitrophic interactions (Manning *et al.*, 2015) more accurately than diversity measures considering only one taxon.

Additionally, the responses of different organismal groups to successional gradients may differ

either due to stochastic or deterministic drivers (Kaufmann, 2001; Blaalid *et al.*, 2012; Fenton

and Bergeron, 2013; Dini-Andreote *et al.*, 2015; Måren *et al.*, 2018; Junker *et al.*, 2021) and

thus conclusions about the primary assembly processes may be biased in studies considering

a single or few taxa only (Pulsford, Lindenmayer and Driscoll, 2016).

The assumed changes of assembly processes along successions, a decrease in the

importance of dispersal and a simultaneous increase in niche-based and interaction-based

processes, have consequences for the taxonomic, functional and phylogentic composition of

communities and thus beta diversity (Odum, 1969; Chang and HilleRisLambers, 2016). The

framework by Chang & HilleRisLambers (2016) predicts an increase in functional and

phylogentic diversity of local single-taxon communities in later successions due to niche

differentiation and intensified biotic interactions. In a multidiversity context where multiple taxa

are considered, increased interactions and thus potentially strong dependencies between taxa

may result in aggregated co-occurrence patterns (Ohlmann *et al.*, 2018), which may have

consequences for the beta-diversity between local communities – and not necessarily for the

functional and phylogenetic diversity within a community. For instance, interactions between

plants and soil inhabiting microorganisms intensify with successional age (Chen *et al.*, 2019)

and regulate plant-animal interactions (Howard, Kao-Kniffin and Kessler, 2020), which is

reflected in the co-occurence of the partners involved in these interactions. Such interactions

may thus result in reduced species turnover across local assemblages in older successional

stages. Yet, it is an ongoing debate whether ecological communities generally converge

towards a core set of species over the course of succession (Lepš *et al.*, 1991; Fukami *et al.*,

2005) and studies on successional convergence have mostly been focused on singular taxa

neglecting potential interactions and co-occurrences across organismal groups (Brown and

Jumpponen, 2014; Castle *et al.*, 2016; Chang *et al.*, 2019). To specifically test the effect of

interactions across taxa on species turnover, we introduce multi-betadiversity, an aggregated

measure of taxonomic community dissimilarity. This index allows to test whether the expected

increase in biotic interactions is reflected in a reduced compositional variation (i.e., reduced

multi-betadiversity regarding several taxa) of the multidiverse community once niche-based

and interaction-mediated assembly processes dominate over dispersal as main mechanisms

shaping multidiversity.

Assuming that multidiversity more precisely reflects ecosystem responses and functions, we

tested predictions deduced from contemporary succession hypotheses using a dataset on

multidiversity comprising inventories of plants, animals, and microorganisms along a

successional gradient. Most studies on multidiversity have been undertaken in well-developed

ecosystems that have undergone severe anthropogenic alterations in the past and focused on

the drivers of biodiversity decline (Allan *et al.*, 2014; Manning *et al.*, 2015; Felipe-Lucia *et al.*,

2020). The mechanisms behind the successional emergence of multidiversity and ecosystem

complexity under natural conditions, however, are yet poorly understood. Thus, considering

multidiversity within a successional framework will help to gain new insights into the temporal

dynamics of the processes that shape the initial emergence of biodiversity in natural

ecosystems that comprise multiple organismal groups with unique and complementary

ecological functions. We adopted the concept of multidiversity to empirically test predictions of

the integrated framework of succession and community assembly dynamics on an ecological

gradient of primary succession following glacial retreat in the Austrian Alps (Junker *et al.*,

2020). We assessed the multidiversity and multi-betadiversity of five organismal groups,

including vascular plants, bryophytes, invertebrates, and soil inhabiting fungi and bacteria on

110 plots spanning 170 years of primary succession. We estimated multidiversity by the sum

of ranked normalized Shannon-diversities and calculated multi-betadiversity as the mean Bray-

Curtis dissimilarities between plots regarding the composition of individual taxa (see methods).

According to the conceptual framework by Chang & HilleRisLambers (2016), we predicted that

community assembly will undergo a rapid shift that is induced by a sudden change of the biotic

and abiotic environment along the successional gradient. We expect this shift to be associated

with increased biotic interactions that result in a reduced compositional variation between plots

of the multidiverse community.

Accordingly, we identified a threshold in multidiversity development (i.e., a shift from an

increase of multidiversity with time to a rather stationary phase) and estimated the relative

importance of dispersal, environmental and biotic drivers on the assembly of the multidiverse

community before and after the threshold using path analysis. We further highlighted the more

structured character of community assembly after the shift by comparing multi-betadiversity

estimates of the early and late successional communities.

Results

Breaking point in multidiversity. Multidiversity follows a non-linear trajectory along

the successional gradient. We screened the gradient for the predicted shift from a developing

(characterized by an increase in multidiversity over time) to a more stationary ecosystem

(characterized by non-monotonic variation in multidiversity over time) using breaking point

analysis, which detects changes in the slope of associations. According to Groffman (2006)

breaking points represent ecological thresholds, which is characterized by a rapid change of

the slope in the association between time since deglaciation as explanatory variable and

multidiversity as dependent variable in our study. We found such a breaking point by cross-

validating a broken-stick model to alternative models that either did not include a breaking point

or included a disjunct breaking point using Bayesian inference (Lindeløv, 2020). We then used

Bayes Factors analysis to validate the exact location of the breaking point at plot number 44

which reflects about 60 years of succession (see Methods, Fig. 1, Supplementary Table 1).

Resulting piecewise linear models showed a significant linear increase in multidiversity during

the early successional stage and a stationary multidiversity (non-significant relationship with

time) for the late successional stage (early: $t_{42} = 5.77$, $p < 0.001$, $r^2 = 0.44$; late: $t_{64} = -0.55$, $p =$

0.58, $r^2 = 0.004$; Fig.1). To assess whether the change in assembly processes is also reflected

in the composition of the multidiverse community, we calculated the relative abundance of

each taxon (see Methods) along the successional gradient and visualized it in a temporally

ordered multi-taxa community table of species occurrences (Fig. 2). The threshold became

also evident in the change of community composition with different sets of taxa before and

after the breaking point. Prior to the threshold, communities mainly consist of species that

reach their abundance optima early in succession. These pioneering species were soon

accompanied by taxa with no clear preference of early or late successional stages leading to

an increase in multidiversity over time. After the threshold, specialists for early successions

are replaced by specialists for late successional stages, consequently multidiversity remained

stationary over time (Fig. 2).

Drivers of multidiversity. To quantify the relative importance of dispersal and

environmental drivers, we specified a path model in which time since deglaciation (as a proxy

for dispersal events that occur at a random frequency as main driver (Lowe and McPeek,

2014)) and mean seasonal soil temperature (as an environmental factor that is important for

diversity of several trophic levels (Ohler, Lechleitner and Junker, 2020)) act as exogenous

variables and directly affect diversities of single organismal groups. The total effects of the

dispersal and environmental drivers on multidiversity were estimated as indirect effects

mediated through the organismal groups. We also tested the effects of other environmental

variables, such as soil nutrient content and soil-pH value but could not detect any significant

effects and thus removed these variables from the final model (see Methods and

Supplementary tables 2-4). Biotic interactions were modeled as the residual covariance among

the diversity values of all organismal groups. Community covariances can be interpreted as

biotic interactions and especially positive covariances are indicative of facilitative effects within

the community (Houlahan *et al.*, 2008; Ranta *et al.*, 2008). The path analysis provided good

model fit (Model fit: $p_{chi\text{-}square}$ > 0.05; CFI >0.95; TLI ≥ 0.9; SRMR < 0.09; $RMSEA_{early}$ < 0.05;

$RMSEA_{late}$ = 0.066 with lower bound of confidence interval = 0) and revealed different

influences of the exogenous variables between the early and late successional environments

(Figure 3, Supplementary Table 5). During early succession, time since deglaciation had a

strong positive effect on the diversity of bryophytes (β = 0.65, p < 0.01), arthropods (β = 0.61,

p < 0.01), and bacteria (β = 0.31, p = 0.04), whereas there was no significant effect of

temperature on any organismal group. Time since deglaciation also had a significant positive

indirect effect on total multidiversity (β = 0.74, p < 0.01) during early succession, whereas

temperature did not affect multidiversity (β = 0.01, p = 0.93, Figure 3a). Furthermore, diversities

of the five organismal groups varied independently of each other suggesting little or no mutual

influences. In late succession, mean soil temperature had a significant positive direct effect on

the diversity of bryophytes ($\beta = 0.30$, $p = 0.01$), yet there was a significantly positive indirect

effect on multidiversity mediated through all organismal groups ($\beta = 0.29$, $p = 0.01$). Plot age

had no longer an effect on multidiversity ($\beta = -0.01$, $p = 0.89$) and we detected positive residual

covariances between vascular plant ($\beta = 0.26$, $p = 0.04$), invertebrate ($\beta = 0.25$, $p = 0.05$) and

bacterial ($\beta = 0.28$, $p = 0.02$) diversities with fungal diversity, suggesting mutual influences

between taxa (Figure 3b).

Multibetadiversity. We predicted that the shift from dispersal dominated stochastic

events in early successional stages to niche-based processes in late successional stages is

associated with lower beta-diversity among communities after the threshold. To test this

prediction, we calculated a measure of total community dissimilarity, multi-betadiversity (mbD)

for both successional stages by bootstrapping Bray-Curtis-similarities of the multidiverse

community (see Methods). Despite different trajectories of betadiversity for different

organismal groups - an increase in dissimilarity for bryophyte, and a decrease for those groups

that show positive community covariances in the path model in the old successional stage

(bacteria, fungi, and vascular plant), and no detectable change for invertebrate communities -

multibetadiversity was significantly lower in the late successional stage (mean $\pm$ SD; mbD_{early}

$= 0.73 \pm 0.14$; $mbD_{late} = 0.71 \pm 0.13$; Fig. 4).

Discussion

Our study is an empirical test of the conceptual framework integrating succession and

community assembly dynamics proposed by Chang & HilleRisLambers (2016)[1] and modified

here in the context of multidiverse communities. We showed that threshold dynamics play an

important role in the assembly of multidiverse communities under natural conditions and that

these thresholds are associated with substantial changes in the community assembly

processes. So far, the importance of threshold dynamics has been recognized for

anthropogenically altered ecosystems either as catastrophic shifts with respect to climate

change (Scheffer *et al.*, 2001; Berdugo *et al.*, 2020) or for restoration efforts of
anthropogenically altered landscapes (Suding and Hobbs, 2009; Kadowaki *et al.*, 2018). We
revealed a threshold after about 60 years of ecosystem development when multidiversity
became stationary. Significant thresholds that are marked by a steep increase followed by
stationarity after 40 to 60 years of succession appear to be a generalizable pattern that occurs
in various aspects of ecosystem development, such as plant, invertebrate, and microbial
diversities and functionality in glacier forefields in Europe and Northern America (Kaufmann,
2001; Tscherko *et al.*, 2003; Raffl *et al.*, 2006; Schütte *et al.*, 2010; Dong *et al.*, 2016). Thus,
acknowledging the effects of threshold dynamics during the development of natural
multidiverse ecosystems, and their universality in ecological systems provides valuable
insights into the patterns and processes of initial ecosystem development.

Our results indicate a strongly dispersal driven assembly in the first 60 years of succession
and a pronounced influence of the environment as well as increased biotic interactions
afterwards. Although different organismal groups follow individual assembly trajectories during
primary succession, multidiversity is mainly promoted by stochastic drivers during the initial
phase of ecosystem development, most assumably by heterogenous dispersal events. Over
the course of succession, stochasticity is replaced by environmental filtering and biotic
interactions as the structuring mechanism of multidiversity. Our data do not allow to directly
test for mutual influences between organismal groups on the taxon level, but we find a strong
pattern of positive community covariances after accounting for environmental variation that is
indicative of facilitative processes and biotic dependencies within the community (Houlahan *et*
al., 2007; Ranta *et al.*, 2008). A previous study in the same glacier forefield showed that after
more than 50 years of ecosystem development, the soil microbial community is mainly
supported by carbon from recent plant production, whereas during the initial stage of
ecosystem development, the microbial community is mainly sustained by ancient carbon
sources (Bardgett *et al.*, 2007). The time point of the shift in the main carbon source largely
coincides with our delineated threshold in multidiversity development. The increased
interdependence of the soil microbial and plant community during the later successional stage

is also reflected in the positive covariances of vascular plant, fungal and bacterial diversities in

our path model. Although bryophytes have been shown to closely interact with epiphytic

microorganisms after glacial retreat (Arróniz-Crespo *et al.*, 2014), potential interactions of

bryophytes with microorganisms may not be detectable in our path-model as we sampled soil

microbial communities and not all substrate types colonized by bryophytes in our study site

(e.g. rocks, scree and litter).

The threshold-mediated character of successional processes is further supported by a

pronounced decline in multibeta-diversity that is indicative of an increased dependence

structure among the organismal groups. In line with that assumption, we could only detect a

lower beta-diversity in those organismal groups that also show positive community covariances

in the path model. Although co-occurrence patterns must not necessarily reflect biotic

interactions (Blanchet, Cazelles and Gravel, 2020), this finding strengthens the assumption of

biotic interactions as an underlying cause for a reduced beta-diversity between plots in later

successional stages, which finds support in studies on plant community assembly (Fukami and

Nakajima, 2013; Martínez-García *et al.*, 2015). The literature suggests the increasing

importance of interactions in community development trajectories. One explanation put forward

is that with increasing niche differentiation, energy and nutrient flows in an ecosystem follow

more complex biochemical pathways that necessitate biotic interactions to retain nutrients in

the ecosystem through closed mineral cycles (Odum, 1969; Chang and HilleRisLambers,

2016). Our results support the increasing importance of biotic interactions as in the late

successional stage, our path model revealed positive community covariances between the soil

microbial, vascular plant, and arthropod communities that may represent such complex

nutritional cycles - comprising primary producers, heterotrophic organisms, and decomposers

- that are limited by energy influx (reflected as soil temperature in our model). For the

organismal groups forming these putative cycles, we find a reduced betadiversity between the

plots that indicates a strong interrelatedness of individual taxa after the threshold, which may

lead to re-occurring core assemblages of species of several taxa. These assumptions and the

threshold suggested by our data after 60 years are further supported by the study of Bardgett

et. al (2007) that revealed a high dependence of heterotrophic microorganisms on plant communities in the Ödenwinkel forefield after a similar number of years. We thus recommend extending the integrative framework of succession and community assembly for multiple interacting taxa that mutually shape their diversity and composition and thus cause a reduction of betadiversity.

Understanding the relative importance and temporal dynamics of deterministic and stochastic processes is a key challenge in community ecology, especially in natural systems and has been in the center of broad debates among ecologists. Using the multidiversity and multi-betadiversity approach allows us to comprehensively understand the processes leading to stable, resilient, and complex ecosystems, which may remain vague in single-taxon approaches. Thus, our study contributes to a synthesis of community ecological theories into succession research, acknowledging the fundamental importance of abrupt state shifts in natural ecosystems. These results are not only a proof of concept but also emphasize that succession is a multi-faceted rather than a linear process that comprises a sequence of assembly processes towards multidiversity and ecosystem complexity.

Methods

Study Design. The study was conducted in the long-term ecological research platform Ödenwinkel which was established in 2019 in the Hohe Tauern National Park, Austria (Dynamic Ecological Information Management System – site and dataset registry: https://deims.org/activity/fefd07db-2f16-46eb-8883-f10fbc9d13a3, last access: March 2021). A total of $n = 135$ permanent plots was established within the glacier forefield of the Ödenwinkelkees which was covered by ice at the latest glacial maximum in the Little Ice Age around 1850. For this study, we used a subset of $n = 110$ plots with complete datasets with all biotic and abiotic variables available. The plots represent a successional gradient spanning over 1.7km in length and were evenly distributed within the glacier forefield at an altitude ranging from 2070 to 2170 m a.s.l. Plots were defined as squares with an area of 1 m² and were all oriented in the same cardinal direction. Further details on the design of the research

platform, exact plot positions and details on the surrounding environment, as well as on the historical glacial extent can be found in Junker et al. (2020).

Sampling. In 2019, we identified all vascular plant and bryophyte species present on the plots and estimated their cover with a resolution of 0.1 %. We sampled above-ground arthropod diversity by installing two pitfall traps on each plot. Traps were set active for a total of $n = 7$ days. The abundance of all arthropods, excluding Collembola and Acari, larger than 3 mm was counted. The abundance of Collembola and Acari and of animals smaller than 3mm was estimated based on random samples of aliquots of the total sample. All arthropods and other animals are identified to the order level. Soil inhabiting bacteria and fungi were sampled from soil cores from an approximate depth of 3 cm, as soil development in the proglacial study area was limited to the top layers of the pedosphere. Soil-microbiome samples were analyzed by next-generation sequencing and microbiome profiling of isolated DNA was performed by Eurofins Genomics (Ebersberg, Germany). Prior to the statistical analysis of microbial communities, we performed a cumulative sum scaling (CSS) normalization (R package "metagenomeSeq" v1.28.2, Paulson *et al.* 2013) on the count data to account for differences in sequencing depth among samples. Detailed information on the sampling strategies of all organismal groups can be found in Junker et al. (2020). Temperature measurements were done by installing temperature loggers (MF1921G iButton, Fuchs Elektronik, Weinheim, Germany) 10 cm north of each plot centre, at the same depth of 3 cm at which the microbial samples were taken. Mean temperature of the growing season was calculated based on the recordings ranging from 26th of June to 16th of September representing the period in which the plots were free of permanent snow cover before and after the winter 2019/2020. In 2020, soil samples were taken and soil nutrients (Ca, P, K, Mg and total N_2) as well as soil pH were measured on all plots by AGROLAB Agrar und Umwelt GmbH (Sarstedt, Germany). Soil nutrient analysis was performed according to the Ö-Norm Boden: L 1087: 2012-12 (K and P - mg/1000g), L 1093: 2010-12 (Mg - mg/1000g) and L 1083: 2006-04 (pH). Total N2 (%) was determined according to the DIN EN 16168: 2012-11. Detailed information on the sampling strategy can be found in Junker et. al (2020).

Calculation of multidiversity. Multidiversity is defined as the cumulative diversity of a

number of taxonomic groups (Allan *et al.*, 2014). The multidiversity of the $n = 110$ plots

composed of the diversities of vascular plants, bryophytes, invertebrates, fungi, and bacteria

present in each plot and was defined as follows: First, we calculated the Shannon diversities

(H) of each of the taxonomic groups in each plot. Second, we ranked the plots by increasing

diversity for each of the five taxonomic groups individually, i.e., for each plot we received $n =$

5 ranks. The mean rank of each plot was defined as multidiversity *mD*. For better interpretability

of the index, we then normalized the mean ranks as x' = ((x - min(x)) / (max(x) - min(x)) to

scale them between zero and one. Ranks were used to give the same weighting to each

taxonomic group despite deviations in the absolute values of Shannon diversity and to reduce

the impact of outliers in *mD* values. To account for differences in the sequencing depth of the

microbial raw dataset, we performed multiple rarefactioning prior to the calculation of Shannon

diversity by averaging the results of $n = 999$ iterations (R package "rtk" v0.2.5.7; Saary *et al.*,

2017) instead of using the CSS-normalized dataset.

Breaking point analysis. According to the hypothesis of community assembly dynamics

(Chang & HilleRisLambers 2016) and by visual inspection of the relationship between

multidiversity and successional age of the plots we expected two different stages of community

establishment along the successional gradient. These stages are separatable by the transition

from a developing (characterized by an increase in multidiversity over time) to a more

stationary ecosystem (characterized by non-monotonic variation in multidiversity over time).

Such non-linear associations are suggestive of regime-shifts of the ecosystem and can be

interpreted as ecological thresholds (Groffman *et al.*, 2006). Piecewise regression models

have been shown to be the most suitable tool for the detection of ecological thresholds in

natural systems, as they are able to correctly estimate the probability, as well as the number

and position of ecological thresholds (Ficetola and Denoël, 2009). We screened the

successional gradient for the existence of a threshold by comparing four different models with

mD as the dependent and plot index number (surrogate for plot age) as the independent

variable using the R-package "mcp" v0.3.0.9 (Lindeløv, 2020). The mcp-method fits piecewise

regression models with a pre-defined number of breaking points and is based on Bayesian

inference. We specified a base model m1 (*mD*-values remain constant along the gradient) and

compared it to three alternative models (m2 = constant linear increase of *mD*-values without a

breaking point, m3 = one breaking point in *mD*-values with a segregated slope (abrupt-

threshold-model), m4 = one breaking point in *mD*-values with a joined slope ("broken-stick" or

smooth-threshold-model). For each model, we separately ran three Markov Chain Monte Carlo

estimators with a uniform prior for a total of n = 11000 generations while discarding the first n

= 1000 generations as burn-in. Model convergence was estimated by visually inspecting the

trace plots and checking that all model parameters reached stationarity. We then compared

the predictive performance of the four models using leave-one-out-cross-validation and

confirmed the exact position of the breaking point using Bayes Factors. Both validation

methods can be applied for a robust and accurate testing of competing hypotheses in

ecological datasets (Aho, Derryberry and Peterson, 2014; Conn *et al.*, 2018; Hanusch *et al.*,

2020).

Community taxa occurrence. To visualize the distribution of individual taxa and thus

the community wide taxonomic turnover along the successional gradient, we estimated the

abundance optimum of each taxon that occurred on at least three plots. Abundance optima

were estimated by calculating the mean plot of occurrence weighted by the abundance of each

taxon (i.e., cover for vascular plants and bryophytes, individual count for invertebrates and

CSS-normalized read number for microorganisms). We calculated the relative abundance per

plot for each taxon by using z-scores of abundances. For each taxon, the mean abundance

was set as 0 with one standard deviation as +1 or -1. Resulting z-scores were then rescaled

between -1 and 1. Range size was estimated by calculating the variance of occurrence plots

(i.e., the span of plots on which a taxon was found along the gradient) weighted by the

abundance of the taxon on the respective plots.

Path analysis. We used path-analysis i) to model the influence of the abiotic

environment on the diversities of all taxonomic groups individually ii) to estimate the strength

of community covariances between the diversities of those groups iii) to infer the effect sizes

of the diversity of each group on total *mD* and iv) estimate the strength of indirect effects of the

exogenous variables that are mediated through the organismal groups on *mD.* We built

separate models with identical structure for the early (*n* = 44 plots) and late (*n* = 66 plots)

successional stages. The stages were delineated by the threshold identified in the breakpoint

analysis. Exogenous variables in the model were time since deglaciation and mean soil

temperature of the growing season. Time since deglaciation reflects the plot age and can be

seen as a proxy for the increasing chance of heterogenous dispersal events that occur over

time. The mean temperature of the growing season is an estimate for environmental

heterogeneity and has been shown to affect diversity directly and indirectly on various trophic

levels (Ohler, Lechleitner and Junker, 2020). All variables were scaled by subtracting the mean

and dividing through the standard deviation. We first ran a full model that included soil pH and

soil nutrient content as additional exogenous variables. None of the two variables showed a

significant direct effect on any organismal group or a significant indirect effect on multidiversity

during either the young or late successional stage. Accordingly, we stepwise removed the two

variables from the model while cross-checking whether the effect strength of one variable

became significant in the absence of the other variable. As no significant effects occurred or

the model fit significantly decreased, we decided to remove both variables from the final model.

Within the final model, the strength of residual covariance between the diversities of all

taxonomic groups was estimated while accounting for influences of time since deglaciation and

temperature. If corrected for an underlying common cause, such as environmental

autocorrelation, strong covariances between members of a community can be interpreted as

biotic interactions and especially positive community covariances are indicative of facilitative

effects within the community (Houlahan *et al.*, 2008; Ranta *et al.*, 2008). All path-models were

estimated using the R-package "lavaan" v0.6-7 (Rosseel, 2012).

Multibetadiversity. We defined Multibetadiversity (*mbD*) as the cumulative averaged

pairwise-dissimilarity across a number of organismal groups. First, we split the total community

composition data table (sites x species, *n* = 110 plots) for each organismal group into two

tables containing the plots before (*n* = 44 plots) and after (*n* = 66 plots) the threshold. Second,

we performed Bootstrapping (BS) on the subset tables for each of the $n = 5$ organismal groups

individually. BS was done for each organismal group on both successional stages separately

and was necessary to account for differences in sample sizes (i.e., number of plots) of the

successional stages. Further, BS provides more robust inferential results and allows to obtain

confidence intervals for the resulting beta diversity estimates. Each BS-replicate consisted of

the following steps: For the early and late successional stage separately, we generated $n =$

1000 new community tables per organismal group by randomly sampling the communities of

$n = 44$ plots (number of plots in the early successional stage) allowing replacement draws. We

then calculated the mean of the pairwise Bray-Curtis dissimilarities for each of the

bootstrapped community tables. Subsequently, we calculated *mbD* as the mean of the $n = 5$

dissimilarity values per BS-replicate.

Figures

Figure 1: Multidiversity along the successional gradient. Multidiversity increases during the early successional stage and remains stationary as succession continues. The vertical dashed line indicates the estimated threshold in multidiversity development, which is the threshold in community assembly as predicted by the framework of Chang & HilleRisLambers (2016). The grey regression line indicates the linear correlation between multidiversity and time since deglaciation using a broken-stick-model. Blue dashed lines indicate the 2.5% and 97.5% quantiles of fitted values. The yellow curve on the x-axis resembles the posterior distribution for the estimate of the breaking point i.e., higher density of the frequency indicates a higher probability for the breaking point to be located at a given value.

Figure 2: Single taxa occurrence along the successional gradient for each taxon that occurs on at least three plots along the gradient. Tiles indicate the relative abundance of each taxon at a given plot relative to the mean abundance of the taxon along the gradient. Taxa are ordered by the weighted average plot of their occurrence, i.e. taxa that reach their abundance

optima early in succession appear in the top left and late successional specialist in the bottom

right. Relative taxon abundance is color-coded and reflects z-scores with the mean abundance

as 0 and one standard deviation as +1 or -1, rescaled between -1 and 1. Multidiversity is

indicated as light pink density plot on top of the community composition plot. Grey density plot

on the left side indicates the range size of the taxa, i.e whether their occurrence is restricted

to a narrow range of the successional gradient or whether taxa occur along the whole gradient.

Black bars on the right represent the organismal group of the taxa in each row, symbols of

organismal groups are ordered by decreasing frequency; white dashed horizontal line marks

the threshold.

Figure 3: Path analysis regarding the effects of stochastic and deterministic drivers on the

multidiverse community for the early and late successional stages. Time since deglaciation

and mean seasonal temperature act as exogenous variables with directed effects on the

diversities of vascular plants, bryophyte diversity, invertebrates, and soil inhabiting fungi and

bacteria. Indirect effects of exogenous variables on multidiversity were calculated by mediation

analysis through the direct effects of the organismal groups on multidiversity. Biotic interactions

are modeled as covariances between organismal groups. Numbers represent standardized

path coefficients and are given for significant paths only.

Figure 4: Betadiversity values for each organismal group and community-wide multi-

betadiversity values for the early and late successional stages. Mean Bray-Curtis dissimilarities

were used to calculate betadiversity and multi-betadiversity values. Green violin plots resemble

the dissimilarity values of the early successional stage (n=44 plots), rose violin plots indicate

the late successional stage (n=66 plots). Violin plots indicate the distribution of the mean

dissimilarities for n=1000 bootstrap draws. Boxplots indicate the upper and lower quartile of

the estimates, and the size of the notch represents the confidence interval of the median. A)

mean of all organismal groups, i.e., multi-betadiversity B) bacteria C) fungi D) vascular plants

E) bryopyhtes F) invertebrates

References

Aho, K., Derryberry, D. and Peterson, T. (2014) 'Model selection for ecologists: The worldviews

of AIC and BIC', *Ecology*, 95(3), pp. 631–636. doi: 10.1890/13-1452.1.

Allan, E. *et al.* (2014) 'Interannual variation in land-use intensity enhances grassland

multidiversity', *Proceedings of the National Academy of Sciences of the United States of*

America, 111(1), pp. 308–313. doi: 10.1073/pnas.1312213111.

Arróniz-Crespo, M. *et al.* (2014) 'Bryophyte-cyanobacteria associations during primary

succession in recently deglaciated areas of Tierra del Fuego (Chile)', *PLoS ONE*, 9(5), pp. 15–

17. doi: 10.1371/journal.pone.0096081.

Bardgett, R. D. *et al.* (2007) 'Heterotrophic microbial communities use ancient carbon following

glacial retreat', *Biology Letters*, 3(5), pp. 487–490. doi: 10.1098/rsbl.2007.0242.

Berdugo, M. *et al.* (2020) 'Global ecosystem thresholds driven by aridity', *Science*, 367(6479),

pp. 787–790. doi: 10.1126/science.aay5958.

Blaalid, R. *et al.* (2012) 'Changes in the root-associated fungal communities along a primary

succession gradient analysed by 454 pyrosequencing', *Molecular Ecology*, 21(8), pp. 1897–

1908. doi: 10.1111/j.1365-294X.2011.05214.x.

Blanchet, F. G., Cazelles, K. and Gravel, D. (2020) 'Co-occurrence is not evidence of

ecological interactions', *Ecology Letters*. Edited by E. Jeffers, 23(7), p. ele.13525. doi:

10.1111/ele.13525.

Brown, S. P. and Jumpponen, A. (2014) 'Contrasting primary successional trajectories of fungi

and bacteria in retreating glacier soils', *Molecular Ecology*, 23(2), pp. 481–497. doi:

10.1111/mec.12487.

Castle, S. C. *et al.* (2016) 'Biogeochemical drivers of microbial community convergence across

actively retreating glaciers', *Soil Biology and Biochemistry*, 101, pp. 74–84. doi:

10.1016/j.soilbio.2016.07.010.

Chang, C. C. *et al.* (2019) 'Testing conceptual models of early plant succession across a

disturbance gradient', *Journal of Ecology*, 107(2), pp. 517–530. doi: 10.1111/1365-

2745.13120.

Chang, C. and HilleRisLambers, J. (2016) 'Integrating succession and community assembly

perspectives', *F1000Research*, 5, pp. 1–10. doi: 10.12688/F1000RESEARCH.8973.1.

Chen, C. *et al.* (2019) 'Meta-analysis shows positive effects of plant diversity on microbial

biomass and respiration', *Nature Communications*, 10(1), pp. 1–10. doi: 10.1038/s41467-019-

09258-y.

Conn, P. B. *et al.* (2018) 'A guide to Bayesian model checking for ecologists', *Ecological*

Monographs, 88(4), pp. 526–542. doi: 10.1002/ecm.1314.

Delgado-Baquerizo, M. *et al.* (2020) 'Multiple elements of soil biodiversity drive ecosystem

functions across biomes', *Nature Ecology and Evolution*, 4(2), pp. 210–220. doi:

10.1038/s41559-019-1084-y.

Dini-Andreote, F. *et al.* (2015) 'Disentangling mechanisms that mediate the balance between

stochastic and deterministic processes in microbial succession', *Proceedings of the National*

Academy of Sciences of the United States of America, 112(11), pp. E1326–E1332. doi:

10.1073/pnas.1414261112.

Dong, K. *et al.* (2016) 'Soil fungal community development in a high Arctic glacier foreland

follows a directional replacement model, with a mid-successional diversity maximum',

Scientific Reports, 6. doi: 10.1038/srep26360.

Felipe-Lucia, M. R. *et al.* (2020) 'Land-use intensity alters networks between biodiversity,

ecosystem functions, and services', *Proceedings of the National Academy of Sciences of the*

United States of America, 117(45), pp. 28140–28149. doi: 10.1073/pnas.2016210117.

Fenton, N. J. and Bergeron, Y. (2013) 'Stochastic processes dominate during boreal bryophyte

community assembly', *Ecology*, 94(9), pp. 1993–2006. doi: 10.1890/12-1944.1.

Ficetola, G. F. and Denoël, M. (2009) 'Ecological thresholds: An assessment of methods to

identify abrupt changes in species-habitat relationships', *Ecography*, 32(6), pp. 1075–1084.

doi: 10.1111/j.1600-0587.2009.05571.x.

Fukami, T. *et al.* (2005) 'Species divergence and trait convergence in experimental plant

community assembly', *Ecology Letters*, 8(12), pp. 1283–1290. doi: 10.1111/j.1461-

0248.2005.00829.x.

Fukami, T. and Nakajima, M. (2013) 'Complex plant-soil interactions enhance plant species

diversity by delaying community convergence', *Journal of Ecology*, 101(2), pp. 316–324. doi:

10.1111/1365-2745.12048.

Groffman, P. M. *et al.* (2006) 'Ecological thresholds: The key to successful environmental

management or an important concept with no practical application?', *Ecosystems*, 9(1), pp. 1–

13. doi: 10.1007/s10021-003-0142-z.

Hanusch, M. *et al.* (2020) 'Biogeography and integrative taxonomy of Epipterygium (Mniaceae,

Bryophyta)', *Taxon*, 69(6), pp. 1150–1171. doi: 10.1002/tax.12324.

Houlahan, J. E. *et al.* (2007) 'Compensatory dynamics are rare in natural ecological

communities', *Proceedings of the National Academy of Sciences of the United States of

America*, 104(9), pp. 3273–3277. doi: 10.1073/pnas.0603798104.

Houlahan, J. E. *et al.* (2008) 'The utility of covariances: A response to Ranta et al.', *Oikos*,

117(12), pp. 1912–1913. doi: 10.1111/j.1600-0706.2008.17078.x.

Howard, M. M., Kao-Kniffin, J. and Kessler, A. (2020) 'Shifts in plant–microbe interactions over

community succession and their effects on plant resistance to herbivores', *New Phytologist*,

226(4), pp. 1144–1157. doi: 10.1111/nph.16430.

Junker, R. R. *et al.* (2020) 'Ödenwinkel: An Alpine platform for observational and experimental

research on the emergence of multidiversity and ecosystem complexity', *Web Ecology*, 20(2),

pp. 95–106. doi: 10.5194/we-20-95-2020.

Junker, R. R. *et al.* (2021) 'Divergent assembly processes? A comparison of the plant and soil

microbiome with plant communities in a glacier forefield', *FEMS Microbiology Ecology*, 97(10),

pp. 1–10. doi: 10.1093/femsec/fiab135.

Kadowaki, K. *et al.* (2018) 'Merging community assembly into the regime-shift approach for

informing ecological restoration', *Ecological Indicators*, 85(December 2017), pp. 991–998. doi:

10.1016/j.ecolind.2017.11.035.

Kaufmann, R. (2001) 'Invertebrate succession on an alpine glacier foreland', *Ecology*, 82(8),

pp. 2261–2278. doi: 10.1890/0012-9658(2001)082[2261:ISOAAG]2.0.CO;2.

Lepš, J. *et al.* (1991) 'Convergence or Divergence: What Should We Expect from Vegetation

Succession?', *Oikos*, 62(2), p. 261. doi: 10.2307/3545274.

Lindeløv, J. K. (2020) 'mcp: An R Package for Regression With Multiple Change Points',

Journal of Statistical Software. doi: 10.31219/osf.io/fzqxv.

Lowe, W. H. and McPeek, M. A. (2014) 'Is dispersal neutral?', *Trends in Ecology and Evolution*,

29(8), pp. 444–450. doi: 10.1016/j.tree.2014.05.009.

Lu, M. and Hedin, L. O. (2019) 'Global plant–symbiont organization and emergence of

biogeochemical cycles resolved by evolution-based trait modelling', *Nature Ecology and*

Evolution, 3(2), pp. 239–250. doi: 10.1038/s41559-018-0759-0.

Manning, P. *et al.* (2015) 'Grassland management intensification weakens the associations

among the diversities of multiple plant and animal taxa', *Ecology*, 96(6), pp. 1492–1501. doi:

10.1890/14-1307.1.

Måren, I. E. *et al.* (2018) 'Changing contributions of stochastic and deterministic processes in

community assembly over a successional gradient', *Ecology*, 99(1), pp. 148–157. doi:

10.1002/ecy.2052.

Martínez-García, L. B. *et al.* (2015) 'Host identity is a dominant driver of mycorrhizal fungal

community composition during ecosystem development', *New Phytologist*, 205(4), pp. 1565–

1576. doi: 10.1111/nph.13226.

Odum, E. P. (1969) 'The strategy of ecosystem development', *Science*, 164(3877), pp. 262–270. doi: 10.1126/science.164.3877.262.

Ohler, L. M., Lechleitner, M. and Junker, R. R. (2020) 'Microclimatic effects on alpine plant communities and flower-visitor interactions', *Scientific Reports*, 10(1), pp. 1–9. doi: 10.1038/s41598-020-58388-7.

Ohlmann, M. *et al.* (2018) 'Mapping the imprint of biotic interactions on β-diversity', *Ecology Letters*, 21(11), pp. 1660–1669. doi: 10.1111/ele.13143.

Paulson, J. (2014) 'metagenomeSeq: Statistical analysis for sparse high-throughput sequencing', *Bioconductor.Jp*, pp. 1–20. Available at: http://www.cbcb.umd.edu/software/metagenomeSeq.

Pulsford, S. A., Lindenmayer, D. B. and Driscoll, D. A. (2016) 'A succession of theories: Purging redundancy from disturbance theory', *Biological Reviews*, 91(1), pp. 148–167. doi: 10.1111/brv.12163.

Raffl, C. *et al.* (2006) 'Vegetation Succession Pattern and Diversity Changes in a Glacier Valley, Central Alps, Austria', *Arctic, Antarctic, and Alpine Research*, 38(3), pp. 421–428. doi: 10.1657/1523-0430(2006)38[421:vspadc]2.0.co;2.

Ranta, E. *et al.* (2008) 'Detecting compensatory dynamics in competitive communities under environmental forcing', *Oikos*, 117(12), pp. 1907–1911. doi: 10.1111/j.1600-0706.2008.16614.x.

Rosseel, Y. (2012) '{lavaan}: An {R} Package for Structural Equation Modeling', *Journal of Statistical Software*, 48(2), pp. 1–36. Available at: https://www.jstatsoft.org/v48/i02/.

Saary, P. *et al.* (2017) 'RTK: Efficient rarefaction analysis of large datasets', *Bioinformatics*, 33(16), pp. 2594–2595. doi: 10.1093/bioinformatics/btx206.

Scheffer, M. *et al.* (2001) 'Catastrophic shifts in ecosystems', *Nature*, 413(6856), pp. 591–596.

doi: 10.1038/35098000.

Schütte, U. M. E. *et al.* (2010) 'Bacterial diversity in a glacier foreland of the high Arctic',

Molecular Ecology, 19(SUPPL. 1), pp. 54–66. doi: 10.1111/j.1365-294X.2009.04479.x.

Suding, K. N. and Hobbs, R. J. (2009) 'Threshold models in restoration and conservation: a

developing framework', *Trends in Ecology and Evolution*, 24(5), pp. 271–279. doi:

10.1016/j.tree.2008.11.012.

Tscherko, D. *et al.* (2003) 'Functional diversity of the soil microflora in primary succession

across two glacier forelands in the Central Alps', *European Journal of Soil Science*, 54(4), pp.

685–696. doi: 10.1046/j.1351-0754.2003.0570.x.

Contributions

R.R.J. conceived and initiated the study. M.H., X.H., V.R.H. and R.R.J designed the study and

conducted field work. M.H. performed the processing and analysis of the data with main inputs

from R.R.J. and X.H.. M.H. and R.R.J. drafted the initial version of the manuscript. All authors

contributed critically to the interpretation of the results, revising, and approving the final version

of the manuscript.

Data availability

Raw sequences of next-generation 16S rRNA gene amplicon sequencing are available at the

NCBI Sequence Read Archive (SRA) under the BioProject accession PRJNA701884 and

PRJNA701890. Raw floristic and zoological data will be made available before publication.

Code availability

R scripts will be made available before publication.

Acknowledgements

We thank the Hohe Tauern National Park Salzburg administration and the Rudolfshütte for

organizational and logistic support, the governing authority Land Salzburg for the permit to

conduct our research (permit no. 20507-96/45/7-2019), Jan-Christoph Otto, Tobias Seifert,

and Anna Vojtkó for help in the field. Hamed Azarbad, Lisa-Maria Ohler and Verena Zieschank

provided valuable comments to improve the study. This research has been supported by the

Austrian Science Fund (FWF), which provided funding to Robert R. Junker (grant no. Y1102).

Competing interests

602 The authors declare no competing interests.

Corresponding author

604 Correspondence and requests for materials should be addressed to Robert R. Junker.