Genomic evidence for inbreeding depression and purging of deleterious

genetic variation in Indian tigers

Anubhab Khan[1]*, Kaushalkumar Patel[1], Harsh Shukla[1], Ashwin Viswanathan[1,2], Tom

van der Valk [3], Udayan Borthakur[4], Parag Nigam[5], Arun Zachariah[6], Yadavendradev

Jhala[5], Marty Kardos[7]*, Uma Ramakrishnan[1,8]*

[1] National Centre for Biological Sciences, TIFR, Bangalore, India

[2] Nature Conservation Foundation, Mysore, India

[3] Swedish Museum of Natural History, Stockholm, Sweden

[4] Aranyak, Guwahati, India

[5] Wildlife Institute of India, Dehradun, India

[6] Kerala Forest Department, Wayanad, India

[7] NOAA, National Marine Fisheries Services, Seattle, USA

[8] Senior Fellow, DBT Wellcome Trust India Alliance

* Corresponding authors, anubhabkhan@gmail.com, martykardos@gmail.com,

uramakri@ncbs.res.in

Classifications: Biological sciences, Genetics, Ecology, Evolution

Keywords : large carnivore, purifying selection, natural selection, conservation

genomics, small population size, whole genome sequencing, genetic load,

endangered species

Abstract

Increasing habitat fragmentation leads to wild populations becoming small, isolated, and threatened by inbreeding depression. However, small populations may be able to purge recessive deleterious alleles as they become expressed in homozygotes, thus reducing inbreeding depression and increasing population viability. We used whole genomes sequencing from 57 tigers to estimate individual inbreeding and mutation loads in a small-isolated, and two large-connected populations in India. As expected, the small-isolated population had substantially higher average genomic inbreeding (F_{ROH}=0.57) than the large-connected (F_{ROH}=0.35 and F_{ROH}=0.46) populations. The small-isolated population had the lowest loss-of-function mutation load, likely due to purging of highly deleterious recessive mutations. The large populations had lower missense mutation loads than the small-isolated population, but were not identical, possibly due to different demographic histories. While the number of the loss-of-function alleles in the small-isolated population was lower, these alleles were at high frequencies and homozygosity than in the large populations. Together, our data and analyses provide evidence of (a) high mutation load; (b) purging and (c) the highest predicted inbreeding depression, despite purging, in the small-isolated population. Frequency distributions of damaging and neutral alleles uncover genomic evidence that purifying selection has removed part of the mutation load across Indian tiger populations. These results provide genomic evidence for purifying selection in both small and large populations, but also suggest that the remaining deleterious alleles may have inbreeding associated fitness costs. We suggest that genetic rescue from sources selected based on genome-wide differentiation should offset any possible impacts of inbreeding depression.

Significance statement

Habitat fragmentation is sequestering species into small and isolated populations with high chances of extinction. But are small and isolated populations at risk of extinction from inbreeding depression, or does inbreeding and purging of deleterious alleles reduce such threat? Using whole genomes from several wild Indian tiger populations, we provide evidence supporting purging of highly deleterious variants in a small isolated population. However, our analyses also indicate that the remaining highly deleterious alleles are at high frequencies, suggesting continued inbreeding depression despite some successful purging. We discuss the implications of our results for conservation, including possible genomics-informed genetic rescue strategies.

Introduction

A large proportion of the earth's biodiversity persists in small and isolated populations in today's anthropogenically modified world[1]. Such populations may suffer from decreased genetic variation and increased inbreeding[2] which together lead to decreased fitness and increased extinction risk[3]. Several theoretical[4,5,6,7] experimental[8] and empirical studies (for example 9) reveal that species surviving in small and isolated populations are at the greatest risk of extinction.

While species exist in nature along a continuum from small to large populations with different levels of isolation, populations of endangered species often tend to be small and isolated. African wild dog, Ethiopian wolf, and great Indian bustard are examples of species where all populations are small and isolated[10,11,12]. The 'small population paradigm' of conservation biology suggests that such smaller and more isolated populations are at a higher risk of extinction due to inbreeding depression and demographic stochasticity[13,14,15].

Inbred individuals express deleterious, partially recessive alleles that are inherited identical-by-descent (IBD) from related parents, leading to inbreeding depression[16]. Such inbreeding depression can reduce the average fitness of a population,

eventually leading to reduced population size, and possibly extinction[17]. A commonly

adopted strategy to conserve inbred populations is genetic rescue[18], which aims to

increase average fitness by decreasing the frequency of deleterious mutations and

increase heterozygosity at loci harboring deleterious alleles, via translocations of

individuals from genetically differentiated populations. A meta-analysis of empirical

data from wild populations showed broadly consistent positive effects of genetic

rescue on fitness[15,19].

Population genetic theory[20,21,22] predicts that purifying selection can reduce

inbreeding depression by purging deleterious alleles from inbred populations in the

absence of immigration. Whether isolated populations are likely to purge a

substantial fraction of the mutation load has been of longstanding interest in

evolutionary biology and conservation. Early empirical data from pedigreed captive

populations suggested that purging was either absent, or resulted only in slight

decreases in inbreeding depression (e.g., 23,24). However, several experimental

studies based on model organisms reveal substantial purging and significant

reduction in inbreeding depression in small populations25,26,27. Recent molecular

and population genetic studies have found genomic evidence for purging in wild

populations (e.g., 28,29,30). Despite broad empirical support for the efficacy of

genetic rescue[15,19], genomic evidence for purging and the long term persistence of

some small isolated populations have been cited to question the small population

paradigm, and to argue that standard genetic rescue practices are likely to be

counterproductive[29,31,32,33]. Whether purging removes enough deleterious alleles to

improve the viability of small, isolated populations (contradicting the small population

paradigm) remains an open question. We address this question by contrasting

genomic inbreeding and mutation load in small isolated versus large-connected

populations of wild tigers.

We use Bengal tigers (*Panthera tigris tigris*) from India as a model to investigate

levels of inbreeding and relative mutation loads in small-isolated and large-

connected populations, and examine the potential for genetic rescue. Tigers are

large, endangered carnivores, but Bengal tigers have high genetic variation

compared to other sub-species, with some sub-populations showing high inbreeding

indicating isolation[34]. All Bengal tiger populations have been through historic

bottlenecks, but inbreeding and genetic variation vary among populations. Also,

some populations are relatively large and connected while others are small and

isolated from both genetic[35] and demographic perspectives[36,37], making them an

ideal system to investigate inbreeding, mutation load and possible genetic rescue

strategies.

We use genomic data to measure the impact of historically declining population sizes

and connectivity on inbreeding and mutation load in three wild Bengal tiger

(*Panthera tigris tigris*), populations that are small-isolated (SI) and large-connected.

The small-isolated population is from north-western India and the large-connected

populations are from south India (s-LC) and central India (c-LC). We expected large-

connected populations to be the least inbred and to have the lowest mutation load.

Alternatively, purging could result in lower mutation load in this small-isolated

population. We also explore strategies for genetic rescue that might effectively

decrease inbreeding depression. For tigers, we specifically suggest strategies for

identifying populations that may benefit from genetic rescue, and how such

strategies may be effectively implemented.

Results and Discussion

Genomic estimates of inbreeding

Tigers exist as several genetic population clusters within India with the total

population size of these clusters ranging from 60 to 1000 individuals. Each genetic

cluster consists of several protected areas, with varying connectivity with other such

protected areas and clusters[35,36,38]. For the purpose of this paper, we defined

populations, and whether they are 'isolated' or 'connected' based on population

genetic and gene flow analyses using markers across the genome[34,35,39]. Here, the

'large' populations currently have hundreds of tigers (minimum 300 tigers), while

'small' populations currently have below 100 tigers[36]. While both large populations

are part of connected landscapes, they do not have identical histories of connectivity.

The southern large-connected population is disconnected from other tiger population

genetic clusters in India (e.g. from Central India and Northwest), while central large-

connected population (Central Indian landscape) was connected by gene flow to

other tiger genetic clusters until recently (see 39,35). SFS-based demographic

history models in Armstrong et al.[34] suggest that most Indian tigers diverged from

each other recently. Increased agriculture, bounty hunting and illegal poaching,

habitat loss and fragmentation together have led to local extinctions of tiger

populations[40,41]. The small-isolated population we studied here has less than one

hundred individuals, but was connected (until very recently) to the central large-

connected population[42] and potentially to the now extinct tiger population in

Afghanistan[43].

We quantify inbreeding by identifying long stretches of the genome that are

homozygous and identical by descent (i.e., runs-of-homozygosity, ROH). The

summed length of ROH divided by total autosomal size is an accurate genomic

measure of inbreeding (F_{ROH}) (e.g., 44). We found that the mean F_{ROH}, measured

using ROH longer than 100Kb ($F_{ROH>100Kb}$), 1Mb ($F_{ROH>1Mb}$) and 5Mb ($F_{ROH>5Mb}$) of all

individuals in our dataset (regardless of population) was above 0.48, 0.31 and 0.23

respectively. Tigers from the small-isolated population were most inbred with an

average $F_{ROH>100Kb}$ of 0.57, while average inbreeding due to recent ancestors (on

average up to 5 generations ago, contingent on assumed recombination rate from

domestic cat[45]) was above $F_{ROH>5Mb}$= 0.37 (Figure 1a). The F_{ROH} was proportional to

effective population size (N_e), with the small isolated population having lower N_e at

all time intervals compared to both large and connected populations. This level of

inbreeding is empirically associated with high risk of extinction[46,47,48]. However,

shorter ROH could be due to shared demographic history of bottlenecks (e.g.

F_{ROH}=100kb) and inbreeding depression may be observed at higher F values for

large carnivores[49]. Pedigree based studies[50] have reported that an inbreeding

coefficient of 0.47 is lethal for tigers.

We show that inbreeding was low in the large-connected populations compared to

the small-isolated population (Figure 1a,b). We compared our estimates for zoo

individuals with their known pedigree values to authenticate these results and found

that estimates of inbreeding indeed correspond with pedigree values[51]

(Supplementary Table 1). Overall, the long ROH of ≥10Mb, signifying inbreeding due to ancestors as recent as 3 generations, are rare in large populations but frequent in the small-isolated population. The mean inbreeding due to recent ancestors in the small-isolated population ($F_{ROH>10Mb}$) was above 0.28, which could lead to several negative effects[46,48]. We found that the large connected populations showed inbreeding due to recent ancestors (on average up to 5 generations ago) of less than $F_{ROH>5Mb}$ = 0.1. The large southern population has more inbred individuals than the large central population, potentially due to its geographical placement, historic size (Figure 1d) and connectivity[35,36,39]. Overall, although the theoretical expectation is that populations with N_e < 1,000 are unlikely to persist[7,30,52,53], this result indicates that a population with a census size of ~600 tigers may be sufficiently large to buffer short term inbreeding effects with some geneflow. The long term viability of these populations remains to be tested. While there are instances of small tiger (and other carnivore) populations persisting[36,37], this could possibly be due to their ability to purge some deleterious alleles. This implies that other large felid species, like lions[54], snow leopards[55,56] and jaguars[57] that often persist in population sizes close to 600 may not necessarily experience the detrimental effects of inbreeding depression over the short term. We caution that this is likely to depend on founder effects and chance variation in the mutation load among populations (but see 30), and does not address the likelihood for fitness decline over the long term[7].

Figure 1: Inbreeding and shared ancestry in Indian tigers. Here we show cumulative inbreeding due to parental shared ancestry due to ancestors at different historical time ranges as estimated by different lengths of ROH, that is, ROH ≥ 10 Mb indicate shared ancestry on average up to 3 generations ago, 5 Mb stretches indicate on average up to 5 generations, 1 Mb stretches indicate on average up to 26 generations and 0.1Mb stretches indicate on average up to 263 generations ago. The figure shows cumulative inbreeding in each population due to shared ancestry between parents at each time range. Error bars are 95% CI. (a), inbreeding in individuals due to shared ancestry 26 generation ago (b), the proportion of the genome shared in in 1 Mb or longer tracts between pairs of individual from a population of individuals belonging to central large connected, southern large connected and small isolated populations. Pan India comparisons in (c) include all pairwise comparisons of wild individuals. Error bars are standard errors. N_e over time as estimated from F_{ROH} for the different populations. Error bars are standard deviations (d).

In addition to having the highest F_{ROH}, we found that pairs of individuals from the

small-isolated population also shared large tracts of genome that were IBD (Figure

1c). This could be a worrying sign for its long-term future as the pairwise IBD sharing

translates into inbreeding in the offspring. On average, pairs of individuals from the

small-isolated population shared about 40% of their genome (as more than 1Mb long

IBD stretches), while pairs from the southern large-connected and the central large-

connected populations share about 25% and 15% respectively, slightly higher than

the corresponding mean F_{ROH} values (Figure 1a). As with inbreeding in offspring,

pairwise IBD sharing is expected to be influenced by founding bottlenecks in addition

to recent small N_e (Figure 1d). The small-isolated population had the lowest

mitochondrial haplotype diversity and the lowest number of mitochondrial haplotypes

(Supplementary Figure 1). This lack of diverse lineages, large tracts of shared IBD

genome between individuals, and the recent small population size may therefore

require continued immigration of individuals to sustain genetic variation[58], or in the

absence of natural immigration, may require genetic rescue.

Mutation loads in small-isolated and large-connected tiger populations

We evaluated the relationships between F_{ROH} and the frequency and genomic

distribution of putatively deleterious genotypes identified using the Ensembl Variant

Effect Predictor[59] (VEP). Under the assumption that most deleterious alleles are

partially recessive[28,60,61], the number of homozygous damaging alleles is likely to be

informative of the fitness cost of individual inbreeding. First, we found that the

number of homozygous putatively damaging alleles (loss-of-function (LOF) and

missense alleles together) was proportional to F_{ROH} in all populations. The number of

putatively damaging homozygotes increased from <1500 for individuals with $F_{ROH} \approx$

0.3 to approximately 2200 for individuals with $F_{ROH} \approx 0.6$ ($P < 0.01$ for each

population, linear regressions; Figure 2). This genomic evidence suggests a fitness

cost associated with increased inbreeding. Additionally, the burden of homozygous

putatively damaging mutations was substantially larger on average in the small-

isolated population (mean $\approx$ 2,000 per individual) than in the central large-connected

population (mean < 1,600 per individual, $P < 0.0001$ randomization test) or the

southern large-connected population (mean $>$ 1,800 per individual, $P = 0.0025$,

randomization test, Figure 2). This is likely due to higher N_e in the central large-

connected population as compared to the southern large-connected population

(Figure 1d).

We further investigated differential mutation load between populations for LOF and

missense mutations identified with VEP, and the R_{XY} method[28,62] to estimate relative

excess of number of putatively deleterious alleles in one population compared to

another. R_{XY} compares the number of derived alleles of a particular category (LOF,

missense, or putatively neutral) found in one population relative to the other. If R_{XY} is

one, both populations X and Y have the same load of derived alleles, whereas if R_{XY}

> 1 then population X has more derived alleles than Y and vice versa if R_{XY} < 1 [28,62].

As above, we found a higher load associated with missense mutations in the small-

isolated population (Figure 2c). This is likely explained by more efficient purifying

selection against missense mutations of relatively small effect, which are typically

only slightly recessive[61], due to larger N_e in the large populations than in the small

population. We also found a lower load of LOF mutations in the small-isolated

population compared to the large populations (Figure 2c). This suggests that higher

inbreeding in the small-isolated population facilitated purging of relatively large-effect

LOF mutations – which are more likely to be highly recessive than small-effect

deleterious mutations[61,63]. Together, these different approaches suggest a higher

putatively damaging allele load in small isolated population coupled with purging of

loss-of-function alleles in the small isolated (and inbred) tiger population.

Despite this evidence for purging of LOF alleles in the small-isolated population, it

appears that high inbreeding lead to individuals in this population having a higher

number of homozygous LOF alleles on average (N=250) compared to the southern

large-connected (N=238) and central large-connected (N=218) populations (Figure

2b). High homozygosity for remaining LOF alleles suggests that the fitness cost due

to LOF mutations may be higher in the small-isolated population compared to the

large populations despite partial purging. Importantly, while it is impossible to know

the realized fitness effects of homozygosity for LOF and missense mutations, this

analysis suggests that purging was not sufficient to result in a smaller fitness cost of

inbreeding in the small-isolated population compared to the larger populations. We

observe similar results when we used a Genomic Evolutionary Rate Profiling

(GERP)[65] analysis to identify potentially highly deleterious alleles by keeping loci

with GERP scores in the top 0.1 percent (Supplementary Figure 3). For this analysis,

and those below, it important to recognize that our measure of mutation load is not a

direct measure of fitness. Rather, our inferences rest on the assumption that

missense and LOF mutations, and mutations at conserved sites are deleterious on

average, which is well supported empirically[65,66]. Definitively determining whether the

bottlenecks faced by these populations[40,41] and the additional bottleneck faced by

north-west Indian population three generations ago in 2004[67] purged part of the

mutation load would require historical genetic data to quantify pre- versus post-

bottleneck mutation loads. Additionally, better estimates of very recent demographic

history would help to predict the potential for purging during the bottlenecks.

Figure 2 Relationships between inbreeding and number of homozygous damaging alleles (LOF plus missense) **(a)**, and homozygous LOF mutations **(b)**, and relative mutation loads in populations of various sizes and levels of isolation **(c)**. Filled circles represent the values of F_{ROH} (x-axis) for different individuals, and the number of homozygous damaging mutations (y-axis), with each population represented by color as indicated in the legend. Open circles and squares represent the mean F_{ROH} and mean number of homozygous damaging mutations, respectively, for each population, with the error bars representing 95% percentile bootstrap confidence intervals. R_{XY} indicates excess of deleterious alleles in population X relative to Y. All

populations had similar numbers of neutral (intergenic) mutations, and error bars are

standard errors from jacknife in **(c)**.

Annotated sets of deleterious variants are rarely available for endangered species.

We compared our missense mutations to annotated sets of domestic cat mutations

(Supplementary Table 2). The missense mutations we identified mapped to genes

responsible for disease states such as hypertrophic cardiomyopathy, progressive

retinal atrophy, polycystic kidney disease, cystinuria, gangliosidosis, hyperoxaluria,

hypothyroidism, mucolipidosis and mucopolysaccharidosis. Further studies are

needed to functionally validate the effect of these mutations in tigers. Studies on

inbred/small populations of Asiatic lions revealed several abnormalities[68], and

genomic studies here might provide further insights on deleterious mutations in large

cats. In the small-isolated population at least one tiger has a potential eye condition

(Supplementary Figure 2), although no skeletal defects were observed in an earlier

study[69]. Additional quantitative genetic studies on inbred zoo individuals can aid in

understanding the possible phenotypic effects of inbreeding and inbreeding

depression for endangered species.

Site-frequency spectra (SFS) revealed that nearly 14% of both putatively damaging

and neutral derived alleles were fixed in the small-isolated population (Figure 3). A

smaller proportion of derived alleles (both damaging and neutral) were fixed in the

large populations (< 0.06). However, there were slightly more fixed derived alleles in

the southern large connected population than in the central large-connected

population. The higher abundance of fixed derived alleles (both damaging and

neutral) in the small-isolated population, and to a lesser extent in the southern large-

connected population, suggests that genetic drift (e.g., due to historical population

bottlenecks and founder effects) drove previously rare alleles to fixation in these

populations (Figure 3)[70]. The fixation of deleterious alleles is consistent with

theoretical predictions for very small populations[4,7] and empirical observations from

wolves[32], Apennine brown bears[71] and other species[31]. In the Isle Royale wolves,

high frequency deleterious alleles were consistent with frequent bone deformities in

the population[32,72]. The large fraction of fixed putatively damaging mutations in the

small isolated population suggests that partial purging of the mutation load due to

LOF mutations (Figure 2c) was not sufficient to erase a substantial fraction of the
mutation load in this population. While the SFS was essentially flat for polymorphic
loci (i.e., excluding loci with derived allele frequencies of 0 or 1) in the small isolated
population (as expected following a population bottleneck), there was a larger
fraction of rare derived alleles in the large populations, as expected for large
populations at mutation drift equilibrium[70]. A larger proportion of putatively damaging
than neutral alleles were at or near fixation in each of the large populations, and
across all of India (Figure 3), possibly due to positive selection driving selective
sweeps on a small fraction of non-synonymous mutations. The hump-shaped SFS
are likely the result of a complex demographic history for Indian tigers, including
strong population structure, and historical population bottlenecks[70,73,74].

Lastly, the frequency distributions of putatively deleterious and neutral derived alleles
reveals signatures of purging via natural selection. Some previously rare, deleterious
alleles are expected to be removed from a population by random genetic drift alone,
during periods of small N_e (e.g., population bottlenecks or founding events). Effective
natural selection through purging is expected to eliminate deleterious derived alleles
more frequently than neutral alleles, or to keep them at lower frequencies than
neutral derived alleles. We expected putatively deleterious derived alleles to either
be absent, or to be less frequent (on average) than putatively neutral derived alleles
if purging has occurred in our study populations. This prediction was supported via a
randomization test in each population and for all populations combined ($P < 2e-4$, 2-
sided randomization test). This suggests that purifying selection has successfully
removed a small fraction of the mutation load in each of the three study populations
and across India. The higher proportion of fixed and very high frequency derived
putatively damaging alleles in the large populations and across Pan India may be
caused by historical selective sweeps on a small fraction of positively selected
nonsynonymous mutations, the details of which we leave to future research.

Figure 3. Site frequency spectrum for damaging (missense and LOF mutations combined, orange) and neutral mutations (blue) for each population. The proportion of loci (y-axis) is shown for each possible derived allele frequency (x-axis). Sample sizes were equalized among the three populations by randomly subsampling 16 non-missing alleles for each locus in each of the three populations. Fixed (frequency=1) and missing (frequency=0) alleles are included in the SFS for the three individual populations. We subsampled 47 non-missing alleles at each locus to calculate the SFS for pan India to account for missing genotypes.

Interestingly, both inbreeding and mutation load for the southern large connected

population (from Western ghats in India) are higher than the central large connected

population (from Central India, Figures 1 and 2). This could reflect the central

location of this population (see 35,38,39). Additionally, the central large connected

population may have had higher historical connectivity (with other population

clusters), a two dimensional network of local populations (versus a linear array in

southern large connected populations) and other ecological factors that allow higher

population densities to be achieved in all habitats of central large connected

population compared to southern large connected populations.

Genetic rescue strategies

Our data suggest that genomic inbreeding (F_{ROH} up to 0.6 in the small isolated

population) is comparable to populations experiencing inbreeding depression (e.g.

Florida puma, F_{ROH} up to 0.6[75] and Isle Royale wolves, F_{ROH} between 0.1 and 0.5[32]).

While we also demonstrate the effects of purifying selection, homozygous LOF

alleles are more frequent, and mildly deleterious missense mutations are in excess

in the small isolated population.

Gene flow is the proposed strategy to sustain small and isolated

populations[18]. Geneflow that results in masking of deleterious alleles responsible for

genetic load, leading to increased population growth rate is genetic rescue.

Evolutionary rescue is geneflow that increases genetic variation[76] and thus adaptive

potential. High F_{ST} at loci with high frequency damaging alleles in the receiving

population (frequency > 0.9) should be indicative of good source populations for

genetic rescue. In practice, for both evolutionary rescue and genetic rescue, source

populations are identified as those that have high genome-wide genetic

differentiation with the receiving population (to maximize the heterozygosity of

admixed offspring (reviewed in 19)). Geneflow from populations with high genome-

wide F_{ST} in theory should result in masking deleterious alleles, allowing both genetic

rescue and evolutionary rescue. However, the smaller number of loci with high-

frequency damaging alleles compared to all polymorphic loci could lead to

discordance between these two proxies for genetic and evolutionary rescue (i.e.,

genetic differentiation at loci with putatively damaging alleles may not be a good

predictor of genome-wide genetic differentiation).

We investigated whether evolutionary rescue is a good proxy for genetic

rescue. We propose that certain source populations may be ideal if they result in

masking of damaging alleles and increased genome-wide heterozygosity (top right

quadrant of Figure 4a). Alternatively, other source populations (bottom right

quadrant) will increase heterozygosity but not mask damaging alleles, while those in

the top left quadrant mask damaging alleles but may not increase heterozygosity

substantially genome-wide (Figure 4a). Our data and estimates reveal a strong and

positive correlation between the two proxies (Figure 4b, adjusted r^2 = 0.91), re-

enforcing that selecting source populations based on high genome wide F_{ST} would

result in both genetic and evolutionary rescue. For the small isolated population, our

data suggest that Kaziranga (proportion of damaging loci rescued = 0.74; Genome-

wide F_{ST} = 0.26, damaging loci F_{ST} =0.26) is the best source population, while

ecologically more similar populations like Kanha (proportion of damaging loci

rescued = 0.69; Genome-wide F_{st} = 0.22; damaging loci F_{ST} =0.23) and Corbett

(proportion of damaging loci rescued = 0.65; Genome-wide F_{st} = 0.26; damaging loci

F_{ST} =0.28) may be more practical, and almost as effective at genetic rescue. For the

large isolated population in Southern India, both proxies are lower on an average

(average proportion of damaging loci rescued = 0.65; average genome-wide F_{ST} =

0.195, average damaging allele F_{ST} = 0.21, across source populations). Note that

high genome-wide F_{ST} between a donor and recipient population could also be due

to intense bottlenecks and resulting drift in endangered species, and such population

pairs may not result in effective genetic rescue. We caution that these inferences are

subject to assumed fitness effects of damaging alleles. Most genome-wide

predictions of deleterious alleles in wild species await annotation with phenotypic

data[77], such a strategy could be implemented when the loci affecting fitness are

identified as in the case of Tasmaninan devils (*Sarcophilus harrisii*[78]) and Soay

sheep[79].

Figure 4. (a) A schematic showing the possible distribution of genome-wide F_{ST} and (b) the observed values for our data, assuming two target populations, Ranthambore Tiger Reserve (in red circles, the small isolated population) and Wayanad Tiger Reserve (blue squares, a protected area within the Southern large connected population). F_{ST} on the y-axis is for loci where the putatively damaging allele is at a frequency >0.9 in the target population, and for all loci across the genome on the x-axis.

Whether there is ever a substantial advantage to selecting source populations for genetic rescue attempts based on differentiation for putatively damaging loci (versus across the whole genome) remains to be tested empirically. Genome wide surveys would help in identifying source populations that best mask damaging alleles and increase genome-wide heterozygosity; however there are other considerations (as highlighted in 33) that could impact selection of source populations. A denser sampling of tigers across the Indian tiger range is necessary to predict best rescue individuals for particular target populations. Such dense sampling of genomes can benefit from sequencing of non-invasive samples since acquiring blood samples from endangered species can be difficult[80]. Methods involving sequencing genomes from shed hair[81] or fecal samples[82] will be useful in creating such translocation networks and monitoring the populations for gene flow.

 Implications for conservation of endangered species

We find that small and isolated populations, in this case tigers of Ranthambore Tiger

Reserve, have high inbreeding arising from both recent ancestors and ancestors in

deep history. Our analyses revealed a signature of purging for putatively large effect

(LOF) deleterious alleles in this population. However, this small and isolated

population also had the highest average number of homozygous damaging alleles of

the three study populations, suggesting potential fitness cost of inbreeding in this

population despite purging of some large effect mutations, and compared to the

large and connected populations. Together, these results demonstrate that purging

(as inferred from genomic signatures) does not eliminate all damaging alleles, and

hence does not contradict the small population paradigm of conservation biology[19].

Careful predictions of putative phenotypic effects of the existing deleterious alleles at

high frequencies might be instructive to understand future inbreeding depression,

and these alleles could potentially be used as early warning signs for population

decline. However, caution is warranted in this regard because it is still unclear how

effectively molecular predictions (i.e., identification of loss of function or derived

alleles at conserved sites) can translate into predictions of fitness differences among

extant individuals and populations.

Historical isolation and bottlenecks due to overharvest appear to have a significant

effect on missense (potentially mild effect) mutation load. Our analyses suggest that

purging can also occur in large-connected populations in the context of endangered

species (Figure 3). This could be because all tiger populations are in general small,

and the range of population sizes is not very wide[36]. Continued isolation of these

populations will increase mutation load in the future. Thus maximizing connectivity

might be the best strategy to minimize extinction (from an ecological perspective),

increase genome-wide genetic variation, and also minimize mutation load in the

future. Maximizing connectivity in the wild would require increasing population

abundance since large populations are sources of dispersing individuals, but also

planning for habitat corridors, since demographics of tiger landscapes are dominated

by extinction and recolonization dynamics (e.g. see 83, 84). Such strategies may be

relevant to several endangered species with wide ranges but fragmented
populations, like elephants, lions and wild dogs.

Critical to providing recommendations for management is understanding ongoing
evolutionary trajectories for endangered species. Given the large number of frequent
deleterious alleles in the small isolated population, management could aim to
decrease the number of close to fixation damaging alleles, potentially maximizing
mean absolute fitness. Our data reveal that source populations with high genome-
wide differentiation (with the target population) would adequately mask deleterious
alleles and increase heterozygosity. Gene flow from even the relatively proximate
large-connected population in Central India would decrease the frequency of
damaging alleles, and allow genetic rescue. We caution that such gene flow could
also introduce other damaging alleles (e.g. in wolves, see 85).

Overall, genomic data and analyses provide richer, more nuanced ways to address
inbreeding and genetic rescue of small and isolated populations of endangered
species (i.e., the small population paradigm in conservation). We demonstrate
effective purging of some loss-of-function alleles, but not mutations with smaller
effects, highlighting the need to characterize detrimental mutations and how they will
be addressed through genetic rescue in future conservation efforts. Conservation
management based on predictive models for fitness effects in the context of
observed mutations will aid long-term persistence of populations in the wild.

Methods

Sample collection

We collected tissue from tranquilized or dead tigers from several tiger reserves (14
protected areas across India); zoo individuals from Sagar et al.[51] were also analyzed
(Figure 5, Supplementary Table 1). Individuals belonged to one of five population
clusters: 1) **Small-isolated population** from north-western India (NW) (n=19) **-** we
obtained tissues from 15 NW individuals and used data from two NW individuals in
Armstrong et al.[34] and two NW individuals in Khan et al.[81]. Our NW samples consist

only of individuals that were born in Ranthambore Tiger Reserve (RTR) but could have been translocated to other protected areas (for example Sariska Tiger Reserve) as per forest department records. 2) **Central Large-connected population** from central India (C) (n=9) **-** we obtained tissue from four Kanha Tiger Reserve (KTR) individuals and used data from Armstrong et al.[34] for five other KTR individuals. 3) **Southern Large-connected population** from southern India (n=11) **-** we obtained tissue from two Wayanad Wildlife Sanctuary (WWS) individuals and two Bandipur Tiger Reserve (BTR) individuals. We additionally used data from Armstrong et al.[34] for five WWS individuals and Liu et. al.[86] for two Nagarhole Tiger Reserve (NTR) individuals. Since WWS, BTR and NTR populations are effectively contiguous, we refer to them together as Wayanad (WAY) tigers. 4) **Nandankanan Zoo**, Odisha, India – Sagar et al.[51] obtained five blood samples from inbred individuals from Nandankanan Zoo, and these genomes were used as a control to test our methods. The pedigree based inbreeding coefficient of these individuals vary from 0.21 (n=1), 0.26 (n=1) and 0.28 (n=3) (Supplementary Table 1). Such pedigree-based estimates are expected to be an underestimate of the actual inbreeding due to the low depth of pedigrees[44]. 5) **Other protected areas -** we also sampled one individual tiger from Lalgarh Division Forest, two individuals from Corbett Tiger Reserve (COR), one from Chitwan National Park, four from Kaziranga Tiger Reserve, one from Periyar Tiger Reserve, one from Bor Tiger Reserve, one from Chandrapur district in Maharashtra and two from Sundarban Tiger Reserve. These individuals combined with others are used for drawing pan-India inferences. The sampling locations are depicted in Figure 6.

Figure 5. Sample locations and populations. The numbers in the figures represent the protected area where, North-west population: 1=Ranthambore Tiger Reserve (n=11), 2=Sariska Tiger Reserve (n=7), South India population: 3=Bandipur Tiger Reserve (n=3), 4=Wayanad Wildlife Sanctuary (n=6), 12=Nagarhole Tiger Reserve (n=2), 5=Periyar Tiger Reserve (n=1), Central Cluster: 6=Kanha Tiger Reserve (n=9), 7=Corbett Tiger Reserve (n=2), 8=Lalgarh Range (n=1), 10=Sunderban Tiger Reserve (n=2), 11=Bor Tiger Reserve (n=1), 13= Chitwan National Park (n=1), 14=Chandrapur (n=1), North East population: 9=Kaziranga Tiger Reserve (n=4) , Zoo: Zoo=Nanadankanan Zoo (n=5), Sagar et al. 2021.

DNA extraction and sequencing

We extracted DNA using Qiagen blood and tissue extraction kit (Cat. 69504) as per the manufacturer's instructions. We prepared DNA whole genome sequencing libraries using NEBNext UltraTM II DNA Library Prep Kit (Cat. E7645L, NEB Inc.). We quantified DNA on QubitTM 3.0 fluorometer using Qubit High sensitivity dsDNA Assay (#Q32854, Thermo Fisher Scientific). We then fragmented the quantified DNA by sonication using Covaris LE220 ultrasonicator and Covaris microTUBE (#520053, Covaris® Inc.) to obtain a final insert size of 250–350 bp. Next, we performed end-repair on the DNA fragments where blunt ends are created on either side of the fragments. We added a single "A" nucleotide on the 3′ ends of the fragments to facilitate the ligation of NEB stem-loop adapters. We cleaned the ligated products and size selected using Agencourt AMPure XP beads (#A63882, Beckman Coulter). We amplified these size selected products using an eight-cycle PCR during which indices (Barcodes) and flow-cell binding sequences were added. After a final clean-up with Agencourt AMPure XP beads, we quantified the libraries using Qubit DNA Assay and assessed the fragments using DNA TapeStation D1000 Screen Tape (#5067-5582,5583, Agilent Technologies). We finally clonally amplified the quantified libraries on a cBOT and sequenced them on the HiSeq X with 150bp paired end chemistry.

Whole Genome Re-sequencing and variant discovery

We trimmed the raw reads using TRIMMOMATIC[87] to have an mean PHRED-scaled quality of 30 in a sliding window of 15 bp, and removed any read that was shorter than 36 bp after trimming from further analysis. We aligned these reads to a Bengal tiger reference genome (JAHFZI000000000) using BOWTIE2[88]. The alignments were then saved in a binary format (BAM) using SAMTOOLS1.9[89]. We marked duplicate reads with the Picard Tools `MarkDuplicates` command (http://broadinstitute.github.io/picard). We called variants from the BAM files using Strelka with default options[90]. The variants were filtered with VCFtools[91] to retain biallelic sites with a minimum minor allele count of 3, remove indels and loci with mean depth across individuals below 2.5 percentile and above 97.5 percentile

across all loci. We removed all sites with missingness > 20% after removing

genotypes with genotype quality (GQ) less than 30. We removed the X chromosome

from the analysis. 1601148 SNPs with mean read depth = 17.2 (+/- 1.8) remained for

analyses after these filtering steps.

Runs of Homozygosity

We identified ROH using a sliding window, likelihood ratio method[92,93]. For this

analysis we used SNP data without the minimum genotype quality filter described

above, and then used genotype likelihoods, rather than called genotypes, as input in

order to maximize the genomic information available for detecting ROH. We split

each chromosome into sliding windows of 200 SNPs and step size of 10 SNPs. For

each individual *i* and window *j*, we calculated the probability of the observed

genotype likelihood at each SNP *k* (G_k) assuming the two alleles were IBD, and also

assuming they were non-IBD (Wang et al. 2009). We calculated the LOD score for

each window as

$LOD(j,i) = \sum_{k=1}^{k_i} log_{10}\left(\frac{Pr(G_k|IBD)}{Pr(G_k|non-BD)}\right),$

LOD scores across all windows and individuals form a bimodal distribution. Windows

with LOD score ≥ 0 were called IBD, while windows in the left hand side of the

distribution were called non-IBD as in Kardos et al.[44,94]. Consecutive IBD windows

were joined to map the starting and stop positions of ROH for each individual. F_{ROH}

was calculated as the fraction of the autosomal genome in ROH for example,

$$F_{ROH>100Kb} = \frac{\sum ROH_{>100Kb}}{Total\ autosomal\ size}$$

We calculated the percent of the genome in ROH above a particular size class to

estimate inbreeding arising from ancestors in different historical time periods. The

coalescent times of ROH were estimated as *g=100/(2rL)* where *g* is the expected

time (in generations) back to the parental common ancestor where the IBD

haplotypes forming an ROH coalesce[95], r is the recombination rate (here, 1.9cM per

Mb based on domestic cat[45]) and L is the length of the ROH in Mb. For example, a

100Kb ROH are estimated to arise from a single haplotype in an ancestor 263

generations ago on average. An individual's genomic inbreeding arising from

ancestors up to 263 generations ago is then estimated using only ROH longer than

100 kb. Similar calculations were performed for other ROH lengths. Table 1 lists the

lengths of ROH and corresponding average estimated coalescent times for the IBD

haplotypes that form the ROH.

Table 1. ROH lengths and the corresponding expected mean time of their origin due

to inbreeding where number of years assumes a generation length of 5 years.

ROH length(s)	generations since parents shared ancestor	years since parents shared ancestor
<100Kb	More than 263	More than 1315
>100Kb	Up to 263	Up to 1315
>1Mb	Up to 26	Up to 130
>5Mb	Up to 5	Up to 25
>10Mb	Up to 3	Up to 15

We constructed percentile bootstrap confidence intervals for mean F_{ROH} in each

population[96]. We randomly resampled individuals within a population with

replacement 10,000 times, each time calculating the mean F_{ROH} as described above.

The 0.025 and 0.975 quantiles defined the 95% confidence interval for mean F_{ROH} in

each population. We used a randomization procedure to test for a difference in the

mean F_{ROH} for each pair of populations as follows. First, we randomized the

population affiliation for each individual 10,000 times. For each randomization

replicate, we calculated the mean F_{ROH} for each of the two populations. We

calculated the P-value as the proportion of 10,000 randomization replicates where

the difference in F_{ROH} was at least as large as the empirical difference in F_{ROH}.

IBD stretches of genome shared between pairs of individuals

We estimated stretches of genome identical by descent (IBD) between pairs of

individuals using *IBDseq* default parameters[97]. We filtered for shared stretches

longer than 1Mb and estimated sum of shared lengths of more than 1Mb per

autosomal genome size. We estimated this for all pairs of individuals within a

population and all pairs across populations.

Estimating effective population size from ROH

We used the estimates of F_{ROH} using ROH arising from ancestors in 4 different

estimated time frames (Figure 1) to estimate N_e over each respective time frame. If

the estimated coalescent times for ROH are unbiased, then the average F_{ROH} based

on ROH with estimated maximum coalescent times less than *t* generations back in

time is an estimator of the inbreeding accumulated in the population from the time of

sampling back to *t* generations ago. We estimated N_e from the following expression

of mean expected individual inbreeding as a function of N_e over *t* generations

$$F_{ROH,t} = 1 - \left(1 - \frac{1}{2N_e}\right)^t.$$

This approach assumes that the probability of inferring an ROH segment due to

several smaller homozygous segments is very low. We calculated 95% confidence

intervals for N_e by bootstrapping over individuals within each population as described

above for our estimates of mean F_{ROH}.

Mitochondrial haplotype diversity

We used the reads aligned to the mitochondrial scaffold of BenTig1.0. We called

consensus fasta files for each individual from this as described in Khan et al.[81].

These multiple consensus fasta files were aligned using clustal omega

(https://www.ebi.ac.uk/Tools/msa/clustalo/) and written to nexus output files. We then

used DnaSP 6[98] to estimate haplotype diversity.

Identifying effect of mutations

We re-filtered the SNPs as described above with VCFtools[91], except here we

removed genotypes with phred-scaled genotype quality <30, and then removed loci

that were monomorphic, had >20% missing data, or read depth in the bottom and

top 2.5% quantiles for mean read depth across individuals. We also removed loci

and with extreme values of F_{IS} (F_{IS}<-0.5 or F_{IS} >0.95) estimated across all Indian

tigers to remove loci likely frequent genotyping or read mapping errors; this

expectedly resulted in mean F_{IS}>0 (0.14).

We used Ensembl Variant Effect Predictor[59] on SNPs to identify consequence of a

mutation and classify them as missense, loss-of-function (LOF) or intergenic as

described by Xue et al.[28]. We used annotation files generated for the Bengal tiger

assembly using RNA seq.

Mutation load

We used the R_{XY} method described by Do et al.[62] and implemented by Xue et al.[28] to

estimate relative excess of mutation loads in a population X with respect to another

population Y. We randomly selected 7 individuals from each population and

estimated the relative mutation load for each of the three possible pairwise

combinations of the S-I, s-L-C, c-L-C populations. Standard deviations were obtained

by 100 rounds of jackknife. For each round of jackknife for missense and LOF

mutations, we randomly excluded 10-15% of loci and repeated the estimation. For

intergenic variants, 100,000 loci were randomly excluded for each round.

To infer the ancestral state of each site in the tiger reference we converted the

references for the Domestic cat (GenBank: GCA_000181335.4), Cheetah

(GCA_003709585.1) and Lion (GCA_008795835.1) into FASTQ reads by sliding

across the genome in non-overlapping windows of 100 base pairs and transforming

each window into a separate FASTQ read. The resulting FASTQ reads were then

mapped to the tiger reference genome with bwa mem v0.7.17, slightly lowering the

mismatch penalty (*-B* 3) and removing reads that mapped to multiple regions.

Mapped reads were realigned around indels using GATK *IndelRealigner*. Next, we

converted the mapped reads into a haploid FASTA consensus sequence, excluding

all sites with depth above one (as such sites contain at least one mismapped read)

using ANGSD *-dofasta*. The ancestral allele at a locus was then determined as the

majority allele found in genomic alignment of the domestic cat, cheetah and lion.

Sites where the ancestral allele could not be identified (e.g., where the ancestral

identified in the cat-cheetah-lion alignment was not present in our sample of tigers)

were excluded.

We counted the number of homozygous deleterious alleles per individual as number

of derived missense and LOF mutations. We quantified uncertainty in the mean

number of homozygous deleterious alleles for each population using percentile

bootstrap confidence intervals as described above. We used randomization tests as

described above to test for differences in the mean number of homozygous

damaging and LOF alleles per individual for each pair of populations.

Site-frequency spectrum

We equalized the sample sizes across the populations, and across loci within each

population by randomly subsampling 16 non-missing alleles from each locus in each

population population before estimating the derived neutral and derived damaging

allele frequencies. For the SFS analysis of all populations combined, we subsampled

88 non-missing alleles from each locus. Repeating the subsampling process and

analysis had no effect on the results.

We used randomization tests to determine if the mean putatively damaging derived

allele frequency was statistically significantly different from the mean putatively

neutral derived allele frequency in each population, and for all populations combined

(Figure 3). We randomly reassigned the neutral versus damaging status for each

allele frequency 5,000 times. For each of the 5,000 replicates, we recalculated the

randomized mean derived neutral and damaging allele frequency. A *P*-value was

calculated as the proportion of the 5,000 permutations that resulted in a difference in

the mean neutral versus damaging derived allele frequency that was larger than

observed in the non-randomized data.

Genetic rescue and F_{ST}

We estimated genome wide genetic differentiation as F_{ST} with VCFtools. For loci with

high frequency (>0.85, closer to fixation) deleterious (missense and LOF) alleles in a

population, we listed all the loci in each target population (population in need of

rescue). For every possible source population, we estimated the proportion of these

loci with frequency < 0.85. We then calculated F_{ST} at only these loci to estimate

damaging loci F_{ST}. The proportion or loci rescued was calculated as the fraction of

loci that had < 0.85 frequency in the possible source population.

We investigate this for five possible source populations, including tigers from Kanha

Tiger Reserve (KTR), Wayanad Wildlife Sanctuary (WAY), Ranthambore Tiger

Reserve (RTR), Corbett Tiger Reserve (COR) and Kaziranga Tiger Reserve (KAZ).

Acknowledgements

This work was supported primarily by a DBT/Wellcome Trust India Alliance

Fellowship [IA/S/16/2/502714] awarded to UR. AK received support from SciGenome

Research Foundation. Permissions for sample collection were granted in letters F-19

()CWLW/2018-19/653-55 dated 15-06-2018; E2/PCCF/WL/CR/40/2013-14 and

PCCF(WL)/E2/CR/28/2017-18. Zoo samples were collected as described in Sagar et

al., 2021, under CZA permit - 9-3/2005-CZA(Vol lll)(D)/694/2017. Leslie Lyons

collected and kindly shared the list of cat mutations. Gratitude to NCBS IT team

support during data analysis; NCBS data cluster used is supported under project no.

12-R&D-TFR-5.04-0900, Department of Atomic Energy, Government of India). Byron

Weckworth, Meghana Natesh, Mousumi Ghosh, Megan Aylward, Chris Kyriazis,

Ullas Karanth and Phil Hedrick provided helpful discussions and comments on drafts

of the manuscript. Gratitude to field assistants for their continuous presence in the

field and assisting in sample collection for this work. We thank the editor and two

anonymous reviewers for their comments that helped improve the manuscript.

Data availability
All sequencing data have been deposited in BioProject PRJNA728665 in NCBI

References

1) N. M. Haddad, *et al.*, Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**, 1500052 (2015).

2) D. R. Schlaepfer, B. Braschler, H.-P. Rusterholz, B. Baur, Genetic effects of anthropogenic habitat fragmentation on remnant animal and plant populations: a meta-analysis. *Ecosphere* **9**, e02488 (2018).

3) D. Spielman, B. W. Brook, R. Frankham, Most species are not driven to extinction before genetic factors impact them. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 15261–15264 (2004).

4) R. Lande, Risk of population extinction from fixation of new deleterious mutations. *Evolution* **48**, 1460–1469 (1994).

5) R. Lande, Risks of Population Extinction from Demographic and Environmental Stochasticity and Random Catastrophes. *Am. Nat.* **142**, 911–927 (1993).

6) R. Lande, G. Barrowclough, Effective population size, genetic variation, and their use in population. *Viable populations for conservation*, 87–123 (1987).

7) M. Lynch, J. Conery, R. Burger, Mutation accumulation and the extinction of small populations. *Am. Nat.* **146**, 489–518 (1995).

8) D. Newman, D. Pilson, Increased probability of extinction due to decreased genetic effective population size: experimental populations of clarkia pulchella. *Evolution* **51**, 354–362 (1997).

9) R. Burrows, H. Hofer, M. L. East, Demography, extinction and intervention in a small population: the case of the Serengeti wild dogs. *Proc. Biol. Sci.* **256**, 281–292 (1994).

10) R. Woodroffe, C. Sillero-Zubiri, Lycaon pictus (amended version of 2012 assessment). *The IUCN Red List of Threatened Species* **2020** (2020).

11) J. Marino, C. Sillero-Zubiri, Canis simensis. The IUCN Red List of Threatened Species.

12) S. Dutta, A. R. Rahmani, Y. V. Jhala, Running out of time? The great Indian bustard Ardeotis nigriceps—status, viability, and conservation strategies. *Eur. J. Wildl. Res.* **57**, 615–625 (2011).

13) G. Caughley, Directions in Conservation Biology. *J. Anim. Ecol.* **63**, 215 (1994).

14)P. W. Hedrick, R. C. Lacy, F. W. Allendorf, M. E. Soule, Directions in
conservation biology: Comments on caughley. *Conserv. Biol.* **10**, 1312–1320
(1996).

15)K. Ralls, P. Sunnucks, R. C. Lacy, R. Frankham, Genetic rescue: A critique of
the evidence supports maximizing genetic diversity rather than minimizing the
introduction of putatively harmful genetic variation. *Biol. Conserv.* **251**, 108784
(2020).

16)D. Charlesworth, B. Charlesworth, Inbreeding Depression and its Evolutionary
Consequences. *Annu. Rev. Ecol. Syst.* **18**, 237–268 (1987).

17)L. F. Keller, D. M. Waller, Inbreeding effects in wild populations. *Trends Ecol.*
Evol. **17**, 230–241 (2002).

18)A. R. Whiteley, S. W. Fitzpatrick, W. C. Funk, D. A. Tallmon, Genetic rescue
to the rescue. *Trends Ecol. Evol.* **30**, 42–49 (2015).

19)R. Frankham, Genetic rescue of small inbred populations: meta-analysis
reveals large and consistent benefits of gene flow. *Mol. Ecol.* **24**, 2610–2618
(2015).

20)P. W. Hedrick, Purging inbreeding depression and the probability of extinction:
full-sib mating. *Heredity* **73 (Pt 4)**, 363–372 (1994).

21)P. W. Hedrick, A. Garcia-Dorado, Understanding Inbreeding Depression,
Purging, and Genetic Rescue. *Trends Ecol. Evol.* **31**, 940–952 (2016).

22)A. García-Dorado, Understanding and predicting the fitness decline of shrunk
populations: inbreeding, purging, mutation, and standard selection. *Genetics*
190, 1461–1476 (2012).

23)J. D. Ballou, Ancestral inbreeding only minimally affects inbreeding
depression in mammalian populations. *J. Hered.* **88**, 169–178 (1997).

24)E. H. Boakes, J. Wang, W. Amos, An investigation of inbreeding depression
and purging in captive pedigreed populations. *Heredity* **98**, 172–182 (2007).

25)V. Avila, C. Amador, A. García-Dorado, The purge of genetic load through
restricted panmixia in a Drosophila experiment. *J. Evol. Biol.* **23**, 1937–1946
(2010).

26)N. Pekkala, K. E. Knott, J. S. Kotiaho, K. Nissinen, M. Puurtinen, The benefits
of interpopulation hybridization diminish with increasing divergence of small
populations. *J. Evol. Biol.* **25**, 2181–2183 (2012).

27) E. López-Cortegano, A. Vilas, A. Caballero, A. García-Dorado, Estimation of
genetic purging under competitive conditions. *Evolution* **70**, 1856–1870
(2016).

28) Y. Xue, *et al.*, Mountain gorilla genomes reveal the impact of long-term
population decline and inbreeding. *Science* **348**, 242–245 (2015).

29) J. A. Robinson, C. Brown, B. Y. Kim, K. E. Lohmueller, R. K. Wayne, Purging
of Strongly Deleterious Mutations Explains Long-Term Persistence and
Absence of Inbreeding Depression in Island Foxes. *Curr. Biol.* **28**, 3487–
3494.e4 (2018).

30) C. Grossen, F. Guillaume, L. F. Keller, D. Croll, Purging of highly deleterious
mutations through severe bottlenecks in Alpine ibex. *Nat. Commun.* **11**, 1001
(2020).

31) J. A. Robinson, *et al.*, Genomic Flatlining in the Endangered Island Fox. *Curr.*
Biol. **26**, 1183–1189 (2016).

32) J. A. Robinson, *et al.*, Genomic signatures of extensive inbreeding in Isle
Royale wolves, a population on the threshold of extinction. *Sci Adv* **5**,
eaau0757 (2019).

33) C. C. Kyriazis, R. K. Wayne, K. E. Lohmueller, Strongly deleterious mutations
are a primary determinant of extinction risk due to inbreeding depression. *Evol*
Lett **5**, 33–47 (2021).

34) E. E. Armstrong, *et al.*, Recent Evolutionary History of Tigers Highlights
Contrasting Roles of Genetic Drift and Selection. *Mol. Biol. Evol.* **38**, 2366–
2379 (2021).

35) N. Alcala, A. Goldberg, U. Ramakrishnan, N. A. Rosenberg, Coalescent
theory of migration network motifs. *Mol. Biol. Evol.* **36**, 2358–2374 (2019).

36) Jhala, Y.V., Qureshi, Q. and Nayak, A.K., *Status of tigers, copredators and*
prey in India (National Tiger Conservation Authority, Government of India,
New Delhi, and Wildlife Institute of India, Dehradun) (2020).

37) Q. Qureshi, *et al.*, *Connecting tiger populations for long-term conservation*
(National Tiger Conservation Authority & Wildlife Institute of India Dehradun)
(2014).

38) V. Kolipakam, S. Singh, B. Pant, Q. Qureshi, Y. V. Jhala, Genetic structure of
tigers (Panthera tigris tigris) in India and its implications for conservation.
Global Ecology and Conservation **20**, e00710 (2019).

39) M. Natesh, *et al.*, Conservation priorities for endangered Indian tigers through
a genomic lens. *Sci. Rep.* **7**, 9614 (2017).

40) M. Rangarajan, *India's wildlife history: an introduction* (Orient Blackswan)
(2005).

41) J. Sramek, Face Him Like a Briton": Tiger Hunting, Imperialism, and British
Masculinity in Colonial India, 1800-1875. *Vic. Stud.* **48**, 659–680 (2006).

42) P. Singh, G. V. Reddy, *Lost Tigers Plundered Forests: A report tracing the*
decline of the Tiger across the state of Rajasthan (1900 to Present (WWF
India) (2016).

43) S. Mondol, M. W. Bruford, U. Ramakrishnan, Demographic loss, genetic
structure and the conservation implications for Indian tigers. *Proc. Biol. Sci.*
280, 20130496 (2013).

44) M. Kardos, *et al.*, Genomic consequences of intensive inbreeding in an
isolated wolf population. *Nat Ecol Evol* **2**, 124–131 (2018).

45) G. Li, *et al.*, A High-Resolution SNP Array-Based Linkage Map Anchors a
New Domestic Cat Draft Genome Assembly and Provides Detailed Patterns
of Recombination. *G3* **6**, 1607–1616 (2016).

46) K. Ralls, K. Brugger, J. Ballou, Inbreeding and juvenile mortality in small
populations of ungulates. *Science* **206**, 1101–1103 (1979).

47) K. Ralls, J. D. Ballou, A. Templeton, Estimates of lethal equivalents and the
cost of inbreeding in mammals. *Conserv. Biol.* **2**, 185–193 (1988).

48) L. Laikre, N. Ryman, Inbreeding Depression in a Captive Wolf (Canis lupus)
Population. *Conserv. Biol.* **5**, 33–40 (1991).

49) T. van der Valk, M. Manuel, T. Marques-Bonet, K. Guschanski, Estimates of
genetic load in small populations suggest extensive purging of deleterious
alleles. *bioRxiv*, 696831 (2019).

50) A. K. Roychoudhury, K. S. Sankhala, Inbreeding in white tigers. *Proc. Indian*
Acad. Sci., Anim. Sci. **88**, 311–323 (1979).

51) V. Sagar, *et al.*, High frequency of an otherwise rare phenotype in a small and
isolated tiger population. *PNAS* (in press manuscript ID 2020-25273).

52) R. Frankham, C. J. A. Bradshaw, B. W. Brook, Genetics in conservation
management: Revised recommendations for the 50/500 rules, Red List
criteria and population viability analyses. *Biol. Conserv.* **170**, 56–63 (2014).

53) I. G. Jamieson, F. W. Allendorf, How does the 50/500 rule apply to MVPs?
Trends Ecol. Evol. **27**, 578–584 (2012).

54) H. Bauer, S. Van Der Merwe, Inventory of free-ranging lions Panthera leo in
Africa. *Oryx* **38**, 26–31 (2004).

55) J. L. Fox, S. P. Sinha, R. S. Chundawat, P. K. Das, Status of the snow
leopard Panthera uncia in Northwest India. *Biol. Conserv.* **55**, 283–298
(1991).

56) R. M. Jackson, J. D. Roe, R. Wangchuk, D. O. Hunter, Estimating snow
leopard population abundance using photography and capture–recapture
techniques. *Wildl. Soc. Bull.* **34**, 772–781 (2006).

57) S. C. Silver, *et al.*, The use of camera traps for estimating jaguar Panthera
onca abundance and density using capture/recapture analysis. *Oryx* **38**, 148–
154 (2004).

58) J. R. Adams, L. M. Vucetich, P. W. Hedrick, R. O. Peterson, J. A. Vucetich,
Genomic sweep and potential genetic rescue during limiting environmental
conditions in an isolated wolf population. *Proceedings of the Royal Society B:*
Biological Sciences **278**, 3336–3344 (2011).

59) W. McLaren, *et al.*, The Ensembl Variant Effect Predictor. *Genome Biol.* **17**,
122 (2016).

60) A. García-Dorado, A. Caballero, On the average coefficient of dominance of
deleterious spontaneous mutations. *Genetics* **155**, 1991–2001 (2000).

61) A. F. Agrawal, M. C. Whitlock, Inferences about the distribution of dominance
drawn from yeast gene knockout data. *Genetics* **187**, 553–566 (2011).

62) R. Do, *et al.*, No evidence that selection has been less effective at removing
deleterious mutations in Europeans than in Africans. *Nat. Genet.* **47**, 126–131
(2015).

63) B. E. Housden, *et al.*, Loss-of-function genetic tools for animal models: cross-
species and cross-platform differences. *Nat. Rev. Genet.* **18**, 24–40 (2017).

64) G. M. Cooper, *et al.*, Distribution and intensity of constraint in mammalian
genomic sequence. *Genome Res.* **15**, 901–913 (2005).

65) C. D. Huber, B. Y. Kim, C. D. Marsden, K. E. Lohmueller, Determining the
factors driving selective effects of new nonsynonymous mutations. *Proc. Natl.*
Acad. Sci. U. S. A. **114**, 4465–4470 (2017).

66) A. Eyre-Walker, P. D. Keightley, The distribution of fitness effects of new
mutations. *Nat. Rev. Genet.* **8**, 610–618 (2007).

67) S. Narain, H. S. Panwar, M. Gadgil, V. Thapar, S. Singh, *Joining the dots: The*
report of the Tiger Task Force. Project Tiger Directorate (Union Ministry of
Environment, Government of India) (2005).

68) Y. V. Jhala, *et al.*, Asiatic Lion: Ecology, Economics, and Politics of
Conservation. *Frontiers in Ecology and Evolution* **7**, 312 (2019).

69) A. Sadhu, *et al.*, Demography of a small, isolated tiger (Panthera tigris tigris)
population in a semi-arid region of western India. *BMC Zool.* **2**, 16 (2017).

70) G. Luikart, F. W. Allendorf, J. M. Cornuet, W. B. Sherwin, Distortion of allele
frequency distributions provides a test for recent population bottlenecks. *J.*
Hered. **89**, 238–247 (1998).

71) A. Benazzo, *et al.*, Survival and divergence in a small group: The
extraordinary genomic history of the endangered Apennine brown bear
stragglers. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E9589–E9597 (2017).

72) J. Räikkönen, J. A. Vucetich, R. O. Peterson, M. P. Nelson, Congenital bone
deformities and the inbred wolves (Canis lupus) of Isle Royale. *Biol. Conserv.*
142, 1025–1031 (2009).

73) A. Keinan, J. C. Mullikin, N. Patterson, D. Reich, Measurement of the human
allele frequency spectrum demonstrates greater genetic drift in East Asians
than in Europeans. *Nat. Genet.* **39**, 1251–1255 (2007).

74) G. V. Barroso, A. F. Moutinho, J. Y. Dutheil, "A Population Genomics Lexicon"
in *Statistical Population Genomics*, pp. 3–17 (2020).

75) N. F. Saremi, *et al.*, Author Correction: Puma genomes from North and South
America provide insights into the genomic consequences of inbreeding. *Nat.*
Commun. **10**, 5276 (2019).

76) A. A. Hoffmann, A. D. Miller, A. R. Weeks, Genetic mixing for population
management: From genetic rescue to provenancing. *Evol. Appl.* **14**, 634–652
(2021).

77) M. Kardos, A. B. A. Shafer, The Peril of Gene-Targeted Conservation. *Trends*
Ecol. Evol. **33**, 827–839 (2018).

78) P. A. Hohenlohe, *et al.*, Conserving adaptive potential: lessons from
Tasmanian devils and their transmissible cancer. *Conserv. Genet.* **20**, 81–87
(2019).

79) M. A. Stoffel, S. E. Johnston, J. G. Pilkington, J. M. Pemberton, Genetic
architecture and lifetime dynamics of inbreeding depression in a wild
mammal. *Nat. Commun.* **12**, 2972 (2021).

80) A. Khan, A. Tyagi, Considerations for initiating a wildlife genomics research
project in south and south-east Asia. *J. Indian Inst. Sci.* (2021)
https:/doi.org/10.1007/s41745-021-00243-3.

81) A. Khan, *et al.*, Are shed hair genomes the most effective noninvasive
resource for estimating relationships in the wild? *Ecol. Evol.* **10**, 4583–4594
(2020).

82) K. L. Chiou, C. M. Bergey, Methylation-based enrichment facilitates low-cost,
noninvasive genomic scale sequencing of populations from feces. *Sci. Rep.* **8**,
1975 (2018).

83) P. Thatte, A. Joshi, S. Vaidyanathan, E. Landguth, U. Ramakrishnan,
Maintaining tiger connectivity and minimizing extinction into the next century:
Insights from landscape genetics and spatially-explicit simulations. *Biol.*
Conserv. **218**, 181–191 (2018).

84) B. Yumnam, *et al.*, Prioritizing tiger conservation through landscape genetics
and habitat linkages. *PLoS One* **9**, e111207 (2014).

85) P. W. Hedrick, J. A. Robinson, R. O. Peterson, J. A. Vucetich, Genetics and
extinction and the example of Isle Royale wolves. *Anim. Conserv.* **22**, 302–
309 (2019).

86) Y.C. Liu, *et al.*, Genome-Wide Evolutionary Analysis of Natural History and
Adaptation in the World's Tigers. *Current Biology* **28**, 3840–3849.e6 (2018).

87) A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

88) B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**, 357–359 (2012).

89) H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078–2079 (2009).

90) C. T. Saunders, *et al.*, Strelka: accurate somatic small-variant calling from
sequenced tumor-normal sample pairs. *Bioinformatics* **28**, 1811–1817 (2012).

91) P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**,
2156–2158 (2011).

92) S. Wang, C. Haynes, F. Barany, Genome-wide autozygosity mapping in
human populations. *Genetic Epidemiology: The* **33**, 172–180 (2009).
93) T. J. Pemberton, *et al.*, Genomic patterns of homozygosity in worldwide
human populations. *Am. J. Hum. Genet.* **91**, 275–292 (2012).
94) M. Kardos, A. Qvarnström, H. Ellegren, Inferring Individual Inbreeding and
Demographic History from Segments of Identity by Descent in Ficedula
Flycatcher Genome Sequences. *Genetics* **205**, 1319–1334 (2017).
95) E. A. Thompson, Identity by descent: variation in meiosis, across genomes,
and in populations. *Genetics* **194**, 301–326 (2013).
96) B. Efron, R. J. Tibshirani, *An introduction to the bootstrap* (CRC press) (1994).
97) B. L. Browning, S. R. Browning, Detecting identity by descent and estimating
genotype error rates in sequence data. *Am. J. Hum. Genet.* **93**, 840–851
(2013).
98) J. Rozas, *et al.*, DnaSP 6: DNA Sequence Polymorphism Analysis of Large
Data Sets. *Mol. Biol. Evol.* **34**, 3299–3302 (2017).
966
967