Differential Activity-Dependent Increase in Synaptic Inhibition and Parvalbumin Interneuron Recruitment in Dentate Granule Cells and Semilunar Granule Cells

Milad Afrasiabi[1], Akshay Gupta[1,2], Huaying Xu[3], Bogumila Swietek[1] and Vijayalakshmi Santhakumar[1,2]

[1]Department of Pharmacology, Physiology and Neuroscience, Rutgers New Jersey Medical School, Newark, New Jersey 07103, [2]Department of Molecular, Cell and Systems Biology, [2]Department of Statistics, University of California Riverside, Riverside, California 92521.

Abbreviated title: Feedback inhibition of dentate output neuron subtypes

Word count for Abstract: 207   Introduction: 652   Discussion: 1532

Pages: 40   Number of figures: 9   Tables: 1

Competing Interests: The authors declare no competing interests.

Acknowledgements: We thank Dr. Luke Fritzky at the Rutgers Imaging Core for help with imaging, Dr. Archana Proddutur for immunostaining validation of transgenic mouse lines and Dr. Deepak Subramanian for developing schematics. The project was supported by NIH/NINDS R01 NS069861, R01NS097750 and NJCBIR CBIR16IRG017 to V.S., NJCBIR CBIR15FEL011 to M.A and CBIR11FEL003 to A.G.

Correspondence:

Vijayalakshmi Santhakumar, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
3401 Watkins Drive (Rm 1308 Spieth Hall) Riverside, CA 92521
E-mail: vijayas@ucr.edu

Abstract:

Strong inhibitory synaptic gating of dentate gyrus granule cells (GCs), attributed largely to fast-

spiking parvalbumin interneurons (PV-INs), is essential to maintain sparse network activity

needed for dentate dependent behaviors. However, the contribution of PV-INs to basal and input

driven sustained synaptic inhibition in GCs and semilunar granule cells (SGCs), a sparse

morphologically distinct dentate projection neuron subtype are currently unknown. In studies

conducted in hippocampal slices from mice, we find that although basal inhibitory postsynaptic

currents (IPSCs) are more frequent in SGCs and optical activation of PV-INs elicited IPSCs in

both GCs and SGCs, optical suppression of PV-INs failed to reduce IPSC frequency in either cell

type. Amplitude and kinetics of IPSCs evoked by perforant path activation were not different

between GCs and SGCs. However, the robust increase in sustained polysynaptic IPSCs elicited

by paired afferent stimulation was lower in SGCs than in simultaneously recorded GCs. Optical

suppression of PV-IN selectively reduced sustained IPSCs in SGCs but not in GCs. These results

demonstrate that PV-INs, while contributing minimally to basal synaptic inhibition in both GCs

and SGCs in slices, mediate sustained feedback inhibition selectively in SGCs. The temporally

selective blunting of activity-driven sustained inhibitory gating of SGCs could support their

preferential and persistent recruitment during behavioral tasks.

Significance Statement: Our study identifies that feedback inhibitory regulation of dentate

semilunar granule cells, a sparse and functionally distinct class of projection neurons, differs

from that of the classical projection neurons, granule cells. Notably, we demonstrate relatively

lower activity dependent increase in sustained feedback inhibitory synaptic inputs to semilunar

granule cells when compared to granule cells which would facilitate their persistent activity and

preferential recruitment as part of memory ensembles. Since dentate granule cell activity levels

during memory processing are heavily shaped by basal and feedback inhibition, the fundamental

differences in basal and evoked sustained inhibition between semilunar granule cells and granule

cells characterized here provide a framework to reorganize current understanding of the dentate

circuit processing.

Keywords: parvalbumin, dentate gyrus, GABA, feedback inhibition

Introduction:

The hippocampal dentate gyrus is known for its uniquely sparse activity, essential for its function in pattern separation, a process of disambiguating similar inputs into distinct patterns of neural activity and memory representations (Bekinschtein et al., 2013; GoodSmith et al., 2017). The characteristically sparse dentate activity is maintained by the relatively hyperpolarized resting membrane potential and low input resistance of the projection neurons and their powerful inhibitory regulation (Heinemann et al., 1992; Lothman et al., 1992). Dentate inhibition includes sustained tonic/extrasynaptic GABA currents as well as distinct phases of synaptic inhibition, which shape basal and afferent-evoked activity (Stell et al., 2003; Ewell and Jones, 2010). Apart from the classical basal spontaneous synaptic inhibition and afferent evoked feedforward and feedback inhibition, dentate granule cells (GCs) have been proposed to receive "sustained" increase in feedback synaptic inhibition following paired stimulation of perforant path (PP) inputs (Larimer and Strowbridge, 2010). This sustained feedback inhibition has been suggested as a mechanism for surround inhibition needed to facilitate pattern separation (Larimer and Strowbridge, 2010; Walker et al., 2010). However, the reliability, cell-specificity and temporal structure of this sustained feedback inhibition is not known and the identity of inhibitory neurons recruited during this prolonged feedback inhibition has not been examined.

Although dentate GCs have, for long, been considered a structurally and functionally uniform class of projection neurons, detailed morphometric and physiological analysis in rats have revealed a sparse, structurally distinct population of neurons with axonal projections to CA3 described as semilunar granule cells (SGCs) by Ramón y Cajal (1995) (Williams et al., 2007; Gupta et al., 2012; Save et al., 2018). Functionally, SGCs have sustained firing in response to paired PP stimulation, which distinguish them from GCs (Larimer and Strowbridge, 2010).

While SGCs, characterized by wider dendritic span, have been reported in mice (Save et al.,

2018; Erwin et al., 2020), detailed morphometric analysis and functional characterization of

SGCs in mice is lacking. Of particular relevance to dentate inhibitory regulation, whether SGCs

receive similar early (feedforward or disynaptic feedback) as well as sustained polysynaptic

feedback inhibition as GCs or may escape this inhibitory gate is currently unknown. Studies in

adolescent rats show that SGCs have higher tonic GABA currents and receive more frequent

synaptic inhibitory inputs than GCs (Gupta et al., 2012; Gupta et al., 2020). Uniquely, SGCs

show reduction in both tonic GABA currents and frequency of synaptic inhibitory inputs after

brain injury rather than the increase observed in GCs (Gupta et al., 2012). While these findings

suggest different interneuron subtypes may contribute to synaptic inhibition in GCs and SGCs,

which class of dentate interneurons underlie this differential control, is not known. Parvalbumin-

interneurons (PV-INs) are a major class of GABAergic neurons which are classified by

expression of the calcium binding protein parvalbumin and their fast, non-adapting firing pattern.

PV-INs include the perisomatically projecting basket cells which contribute to feedforward and

feedback inhibition of GCs (Kraushaar and Jonas, 2000; Hefft and Jonas, 2005; Ewell and Jones,

2010) and axo-axonic cells which target granule cell axon-initial segments (Sik et al., 1997;

Howard et al., 2005). A single basket cell can project to as many as 10,000 GCs (Freund and

Buzsaki, 1996; Santhakumar, 2008) and is an ideal candidate to mediate surround inhibition.

However, whether PV-INs innervate SGCs with somata located in the inner molecular layer,

outside the dense axon collaterals of parvalbumin positive basket cells has not been tested.

Recent studies have indicated that SGCs may be disproportionately recruited to memory engrams

(Erwin et al., 2020) making it important to understand whether inhibitory regulation of SGCs

differs from that of GCs. This study was conducted to systematically examine the cell-type

specific differences in distinct phases of synaptic inhibition; basal spontaneous synaptic

inhibition, PP evoked, early feedforward, and feedback inhibition as well as sustained feedback

inhibition, between the two classes of dentate projection neurons and to determine the

contribution of PV-INs to specific phases of inhibition in GCs and SGCs.

Materials and Methods

Animals: All experiments were performed in accordance with IACUC protocols approved by Rutgers-NJMS, Newark, NJ, and the University of California at Riverside, CA and in keeping with the ARRIVE guidelines. The study included male and female wild-type C57BL/6J (WT) or appropriately validated optogenetic mice (4-8 weeks old, Jackson Laboratory, Bar Harbor, ME). Parvalbumin (PV)-Cre mice (B6;129P2-Pvalbtm1(cre)Arbr/J; JAX #8069) were crossed with either of the two floxed lines (Chr2-YFP: B6;129S-Gt(ROSA)26Sortm32(CAG-COP4*H134R/EYFP)Hze/J; JAX#12569 or NpHR3-YFP: Rosa-CAG-LSL-eNpHR3.0-EYFP-WPRE; JAX #14539) to generate experimental PV-Chr2 and PV-NpHR3 mice. Immunofluorescence labeling for PV and YFP in sections from PV-YFP mice identified that over 80% of PV labeled cells expressed YFP, and all YFP positive cells expressed PV validating the specificity of the PV-Cre transgenic lines. Mice were housed with littermates (4 mice per cage) in a 12-h light-dark cycle. Food and water were provided ad libitum.

Slice Physiology: Animals were anesthetized under 3% isoflurane and decapitated. Whole brains were extracted and horizontal brain slices (350 µm) were prepared using Leica VT1000 or VT1200S Vibratomes (Wetzlar, Germany) in ice-cold high Sucrose-aCSF containing (in mM) 85 NaCl, 75 sucrose, 24 $NaHCO_3$, 25 glucose, 4 $MgCl_2$, 2.5 KCl, 1.25 NaH_2PO_4, and 0.5 $CaCl_2$. Slices were bisected and incubated at $32 \pm 1°C$ for a minimum of 30 min in a submerged holding chamber containing recording aCSF and subsequently held at room temperature (RT, 22-23°C). The recording aCSF contained (in mM) 126 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 1.25 NaH_2PO_4, 26 $NaHCO_3$ and 10 D-glucose. All solutions were saturated with 95% O_2 and 5% CO_2 and maintained at a pH of 7.4 for 2-6 h (Gupta et al., 2012; Yu et al., 2016).

Slices (350 µm) were transferred to a submerged recording chamber and perfused with

oxygenated aCSF at 33 ± 1°C. Whole-cell voltage- and current-clamp recordings from GCs,

SGCs and presumed interneurons at the border of the hilus and GCL were performed under IR-

DIC visualization with a Nikon Eclipse FN-1 (Nikon Corporation, Shinagawa, Tokyo, Japan) or

Olympus BX50 (Olympus Corporation, Shinjuku, Tokyo, Japan) microscope, using 40X water-

immersion objectives. Recordings were obtained using Axon Instruments Axopatch 200B or

MultiClamp 700B amplifiers (Molecular Devices, Sunnyvale, CA). Data were obtained using

glass microelectrodes (3–5 MΩ) pulled using a Narishige PC-10 puller (Narishige Japan, Tokyo,

Japan) or Sutter P-1000 Glass puller (Sutter Instruments, Novato, CA) and were low-pass filtered

at 3 kHz, digitized using DigiData 1440A and acquired using pClamp10 at 10-kHz sampling

frequency with gains of 0.1 and 1. Recordings were performed using high chloride internal

solution containing (in mM) 125 KCl, 10 K-gluconate, 10 HEPES, 2 $MgCl_2$, 0.2 EGTA, 2 Na-

ATP, 0.5 Na-GTP and 10 PO Creatine titrated to a pH 7.25 with KOH (pH 7.25; 270–290

mOsm) or cesium methane sulfonate ($CsMeSO_4$) internal solution containing (in mM) 140

cesium methane sulfonate, 10 HEPES, 5 NaCl, 0.2 EGTA, 2 Mg-ATP, 0.2 Na-GTP (pH 7.25;

270–290 mOsm). Biocytin (0.3%) was included in the internal solution for post-hoc cell

identification (Gupta et al., 2012; Yu et al., 2015). In experiments with KCl based internal,

synaptic GABA currents were recorded in perfusing aCSF containing the glutamate receptor

antagonist kynurenic acid (3 mM, Tocris, Ellisville, MO). Recorded neurons were initially held

at -70 mV for interneurons and -60 mV for GCs and SGCs and the response to 500 milli-second

positive and negative current injections were examined to determine active and passive

characteristics. Voltage Clamp recordings of IPSCs (spontaneous and evoked) were obtained

from a holding potential of -70 mV for interneurons and -60 mV for GCs and SGCs (Yu et al.,

2015; Gupta et al., 2020). CsMeSO$_4$ based internal was used to record the IPSCs as outward

currents from a holding potential of at 0 mV in the absence of glutamate receptor antagonists. In

some experiments tetrodotoxin (TTX, 1 µM) or gabazine (10 µM) were used to isolate action

potential independent miniature IPSCs or block IPSCs, respectively. Post-hoc biocytin

immunostaining and morphological analysis was used to definitively identify interneurons, GCs

and SGCs included in this study. Morphologies were reconstructed using Neurolucida360

(Microbrightfield) for further analysis (See Morphometry below).

Evoked IPSCs were elicited by single or paired stimulation of the PP using bipolar concentric

stimulating electrodes placed at the junction of the dorsal blade and the crest, just outside the

fissure (Korgaonkar et al., 2020). Constant current stimuli (0.5 µA-5 mA) were applied using a

high-voltage stimulus isolator (A365R, World Precision Instruments, Sarasota, FL). The evoked

mono and polysynaptic IPSCs were recorded simultaneously in GCs and SGC pairs. In some

experiments, slices from were exposed to wide-field blue (λ=470nm) or amber (λ=589nm, 10

mW/mm^2) illumination using Lambda DG-4 Plus (Sutter Instruments, Novato, CA) or Thorlabs

4-Wavelength High-Power LED Sources (Thorlabs, Newton, NJ).

Immunohistochemistry: Following physiological recordings, slices were fixed in

0.1mM phosphate buffer containing 4% paraformaldehyde at 4°C. For *post-*

hoc immunohistochemistry, slices were incubated overnight at 4°C with Alexa Fluor 594-

conjugated streptavidin in 0.3% Triton X-100 and 2% normal goat serum containing PBS. Slices

were mounted on glass slides using vectashield. Sections were visualized and imaged using a

Zeiss LSM 510 confocal microscope with a 0.5 numerical aperture 20× objective.

Morphometry: Stained hippocampal sections were visualized and imaged using a Nikon A1R

laser scanning confocal microscope with a 20X, 0.5 NA objective lens. Cell reconstructions,

from confocal image stacks, were performed using the directional kernels user-guided

reconstruction algorithm in Neurolucida 360 (MBF Bioscience) followed by manual correction

and validation in 3D. About 10-20 percent of each dendritic arbor was reconstructed manually

(Gupta et al., 2020). Algorithms in Neurolucida Explorer (MBF Biosciences) were used to

extract nonnominal or nonordinal somatodendritic morphological quantitative parameters for use

in statistical comparisons and hierarchical cluster analysis. A total of 17 projection neurons in

which the dendritic arbors were fully reconstructed were analyzed. A total of 42 somato-

dendritic parameters (Defined in Gupta et al., 2020) from 17 morphologically reconstructed

neurons, including both features measured in Neurolucida Explorer from the 3D reconstructions

and parameters measured manually in 2D rendering (Neurolucida 360, MicroBrightfield) were

analyzed.

Quantification and Statistical Analysis: Data were tested for uniform distribution and each

quantified variable was fit to the sum of two or more Gaussian functions and quality of fit

determined using maximum likelihood analysis (MLA; v2test) to assess normal distribution of

parameters within each cell type. Variables with a nonuniform distribution were used for

subsequent cluster analysis. Hierarchical cluster analysis on principal components (HCPC) of

morphological parameters was conducted using R version 3.5.0, using R package FactoMineR

by an investigator (H. X.) blinded to cell types and age groups. Hierarchical clustering on the

selected principal components was performed using Ward's criterion with Euclidean distance to

generate the dendrogram. The clustering partition was obtained from hierarchical clustering and

improved with K-means method (Husson et al., 2010).

Individual spontaneous, miniature and evoked IPSCs were detected using custom software in

IgorPro7.0 (WaveMetrics, Lake Oswego, OR) and different parameters were analyzed (Gupta et

al., 2012). Events were visualized, and any "noise" that spuriously met trigger specifications was

rejected. EPSCs were detected and analyzed using event detection in Clampfit. Kinetics and

charge transfer were calculated from the averaged trace of all accepted sIPSC events. Rise time

was measured as the time for amplitude to change from 20 to 80% of peak. Amplitude weighted

τ_{decay} was calculated from a two-exponential fit to the IPSC decay. sIPSC charge transfer was

calculated as the area under the curve of the baseline adjusted average sIPSC trace. Intrinsic

properties were extracted by analyzing the IV traces in Clampfit. Input resistance was calculated

in response to -100 pA current injection. Spike frequency adaptation ratio was calculated as

interval between the first two spikes/interval between the two last spikes in response to +70 pA

current injection. Sample sizes were not predetermined and conformed with those employed in

the field. Significance was set to $p < 0.05$, subject to appropriate Bonferroni correction. Statistical

analysis was performed by paired and unpaired Student's *t* test (Microsoft Excel 2007) or Two-

way repeated measures ANOVA, One-way ANOVA or One-way ANOVA on Ranks followed

by Pairwise Multiple Comparison using Holm-Sidak method or Dunn's method (Sigma Plot

12.3) as appropriate. Data that passed normality test are shown as mean±SEM (standard error of

the mean). Data that were not distributed normally are reported as median and interquartile range

(IQR) as appropriate.

Results

Quantitative morphometry resolves GCs and SGCs in mice into distinct groups.

Recordings were obtained from neurons in the inner molecular layer (IML) to target putative

SGCs and granule cell layer to target GCs followed by *post hoc* recovery of cell morphologies

from biocytin fills (Fig. 1A-E). Unsupervised clustering of morphometric data obtained from

reconstruction of neurons in which dendritic arbors were fully recovered was used to verify

whether the somato-dendritic parameters distinguishing GCs from SGCs in mice are similar to

those characterized in rats (Williams et al., 2007; Gupta et al., 2020). Projection neurons, which

included GCs and SGCs, were identified based on the presence of spiny dendrites in the

molecular layer and axons with mossy fiber boutons entering the hilus and targeting CA3 (Gupta

et al., 2020). Morphometric parameters of the cells reconstructed in 3D were obtained using

automated algorithms in Neurolucida Explorer (Define in Gupta et al., 2020). Principal

Component Analysis (PCA) of 42 morphometric parameters from 17 cells revealed that the first

two principal components (PCs) explained more than 50% of the total variance in the data, while

the first seven components explained 90% of the variance. Hierarchical cluster analysis on PCs

(HCPC) on the first seven PCs suggested partitioning of cells into three clusters (Fig. 1F) as

illustrated on the factor map produced by the first seven PCs (Fig. 1G). In spite of the exclusion

of categorical variables such as soma location from HCPC, cells classified as GCs and SGCs by

investigator (M.A.) segregated into two distinct clusters based on Dimension1 (Dim1).

Interestingly, cells classified as GCs further segregated into two clusters (Clusters #1 and #2)

based on Dim2. Variables more than 70% correlated with the 1st PC were identified as most

responsible for PC formation. Evaluation of the quantitative variables underlying the clusters

revealed that cluster #3, comprised of putative SGCs, was distinguished by a higher number of

primary dendrites, greater dendritic angle, soma width, and cell surface area and lower dendritic

complexity (Fig. 1H-J). While cells in Cluster #1 had fewer, typically one, primary dendrite,

lower dendritic angle, and higher dendritic complexity (Fig. 1H-J), Cluster #2 segregated based

on the presence of more fourth order segments, ends and intersections. Interestingly, dendritic

angle was different between groups, increasing from Cluster #1 to Cluster #3 (Fig 1H). The

number of primary dendrites in Cluster #3 (putative SGCs) was significantly higher than in both

Clusters #1 and #2 (Fig 1I). Similarly, Cluster #3 (putative SGCs) had the least dendritic

complexity which was significantly lower than in Cluster #1 while differences in dendritic

complexity between Clusters #1 and #2 failed to reach statistical significance (Fig 1J).

Morphometric parameters from putative GCs (Clusters #1 and #2) were pooled together and

compared with Cluster #3 (putative SGCs) for further analysis. Like GCs, SGC axons extended

to CA3 and showed mossy fiber terminals in the hilus (Fig. 2A-B), consistent with features of a

projection neuron. Similar to findings in rats (Gupta et al., 2020), GCs and SGCs differed in

dendritic angle, number of primary dendrites, soma aspect ratio and dendritic complexity (Fig.

2C-F, Table 1). In contrast, certain global parameters such as total number of dendritic ends and

total length of dendrites were not different between the cell types (Fig. 2G-H, Table 1). Together

these data provide a comprehensive morphometric comparison of GCs and SGCs in mice. Our

results confirm that the presence of 3 or more primary dendrites and wide dendritic angle reliably

distinguishes SGCs from GCs as reported in recent studies (Save et al., 2018; Gupta et al., 2020;

Rovira-Esteban et al., 2020).

SGCs differ from GCs in intrinsic physiology and frequency of basal synaptic inhibition.

Examination of active and passive responses to graded current injections revealed no difference

in firing frequency between the two cell types (Fig. 2I-K). However, input resistance was

significantly lower in SGCs (Fig. 2K). Spike frequency adaptation ratio (ratio of inter-spike

interval between first 2 and last 2 spikes during a +70 pA current injection) in SGCs was lower

than in GCs (Fig. 2L). These findings are consistent with differences in intrinsic properties

identified between the SGCs and GCs (Williams et al., 2007; Gupta et al., 2012; Save et al.,

2018).

Spontaneous inhibitory postsynaptic currents (sIPSCs), recorded from holding potential of -60

mV in the presence of glutamate blockers, were more frequent in SGCs (Fig. 3A, C: sIPSC

frequency in Hz, GCs: 8.38±1.14, N=18/12 mice, SGCs: 13.48±1.07, N=19/ 13 mice, p=0.003, t-

test). However, sIPSCs amplitude (Fig. 3D: sIPSC amplitude in pA, GCs: 42.98±3.35, N=18,

SGCs: 41.52±3.01, N=19, p=0.75, t-test), amplitude weighted decay time constant ($\tau_{decay-WT}$ in

ms, GCs: 3.31±0.22, N=18, SGCs: 3.2±0.2, N=19, p=0.72, t-test) and 20-80% rise times (in ms,

GCs: 0.22±0.01, N=18, SGCs: 0.20±0.0, N=19, p=0.13, t-test) were not different between GCs

and SGCs (Fig. 3B, D-F). These findings demonstrate that SGCs in mice show the same

structural and functional distinctions from GCs identified in prior studies in rats (Gupta et al.,

2012; Gupta et al., 2020). These data validate the morphological criteria which we used to

distinguish GCs and SGCs in the subsequent experiments and indicate differences in inhibitory

inputs between the cell types.

Limited role of parvalbumin interneurons in steady-state synaptic inhibition of GCs and

SGCs in slices.

To determine whether differential inputs from parvalbumin expressing interneurons (PV-INs),

contribute to the higher action potential-dependent sIPSCs frequency in SGCs we examined if

optical suppression of PV-INs in mice expressing the inhibitory opsin, halorhodropsin (NpHR3),

under control of the PV promotor reduced sIPSC frequency. Unlike archaerhodopsin and

chloride conducting channelrhodopsins which can depolarize terminals and induce synaptic

release, NpHR3 has been shown to hyperpolarize neurons without triggering transmitter release

making it the ideal choice to examine synaptic release (Mahn et al., 2016). Validity of the PV-

Cre mouse line was confirmed by colocalization of parvalbumin and YFP using immunostaining

(not shown) and by fast, non-adapting firing in recordings from YFP positive neurons (Fig. 4A).

Since activation of NpHR3 can reduce the chloride gradient (Alfonsa et al., 2015), light exposure

was restricted to 2s periods. Recordings from YFP labeled neurons established that the 2s amber

light illumination reliably and reversibly hyperpolarized YFP positive fast spiking neurons (n>10

cells) and suppressed action potential firing (Fig. 4A). Spontaneous IPSCs were recorded from

GCs and SGCs for a total of seven minutes: Recordings started with a two-minute baseline

followed by five one-minute cycles with a 2s exposure to amber light at the beginning of each

cycle followed by time for recovery of the chloride gradient between light exposure (Fig. 4B).

Spontaneous IPSC parameters measured during the 2s prior to light exposure were averaged and

compared to the parameters averaged over the periods during the five cycles of light exposure.

Contrary to expectations, optical suppression of PV-INs failed to reduce both frequency (in Hz,

GC no-light: 14.19 ± 2.5; GC Amber light: 13.6 ± 2.4, N=5/3 mice, p=0.61, paired t-test) and

amplitude (in pA, GC no-light: 46.65 ± 9.62; GC Amber light: 45.13 ± 5.94, N=5/3 mice, p=0.42,

paired t-test) of GC sIPSCs (Fig. 4 C-E). Similarly, optical suppression of PV-IN firing failed to

reduce sIPSC frequency (in Hz, SGC no-light: 15.62 ± 2.6; SGC Amber light: $14.69.6\pm2.0$, N=5/3

mice, p=0.79, paired t-test) and amplitude (in Hz, GC no-light: 43.18 ± 5.1; GC Amber light:

40.95 ± 3.2, N=5/4 mice, p=0.42, paired t-test) in SGCs as well (Fig. 4C-E). These data raise a

surprising possibility that PV-INs may not contribute to basal sIPSCs in either GCs or SGCs in

acute slice preparations.

Given the unexpected outcome, and since the optogenetic system was functionally validated

(Fig. 4A), we considered whether use of glutamate receptor antagonist to isolate IPSCs may have

reduced PV-IN firing and underestimated their contribution to sIPSCs. To address this

possibility, IPSCs were recorded as outward currents from cells held at 0 mV (reversal for

glutamatergic currents) using a low chloride internal. As expected, sIPSC frequency recorded in

the absence of glutamate blockers was higher (Fig 4F-H) than that recorded in glutamate receptor

blockers (Fig 3 and 4C-D) in both cell types. Nevertheless, sIPSC frequency and amplitude in

GCs were not reduced by amber light suppression of PV-INs (Fig. 4F-H: frequency in Hz, no-

light: 26.80±3.78, light: 28.32±2.41, N=13/ 8 mice, p=0.49, paired t-test; amplitude in pA, no-

light: 26.43±2.88, light: 25.46±2.10, N=13/ 8 mice, p=0.76, paired t-test). Similarly, sIPSC

parameters in SGCs were also unaltered during amber light exposure (Fig. 4F-H: frequency in

Hz, no-light: 35.77±5.44, light: 36.81±4.48, N=11/ 8 mice, p=0.66, paired t-test; amplitude in

pA, no-light: 28.96±3.72, light: 26.74±3.17, N=11/ 8 mice, p=0.25, paired t-test). Together, these

data obtained under two different recording conditions support the surprising conclusion that PV-

INs contribute minimally to basal sIPSC frequency in both dentate projection neuron subtypes.

While unlikely, a potential reason for the above outcome is that a majority of the sIPSCs in our

recordings were action potential independent miniature events, which would not be modified by

suppressing PV-IN firing. To exclude this possibility, we tested the ability of the sodium channel

antagonist tetrodotoxin (TTX, 1μM) to reduce frequency of IPSCs indicating the presence of

action potential driven events. As expected based on prior studies (Otis et al., 1991; Goswami et

al., 2012), TTX significantly reduced IPSC frequency in both GCs (frequency in Hz, before

ACSF: 11.58±1.58, TTX: 1.4±0.44 N=7, p=0.0003, paired t-test) and SGCs (frequency in Hz,

before ACSF: 10.83±2.09, TTX: 1.96±0.9 N=5, p=0.001, paired t-test). TTX also reduced IPSC

amplitude in both cell types (not shown). Overall, the ability of TTX to suppress IPSC frequency

indicates that majority of sIPSCs in both SGCs and GCs are driven by interneuron firing. Taken

together, these data support the conclusion that PV-INs do not contribute significantly to action

potential driven basal sIPSCs in both GCs and SGCs in slice preparations.

Activation of parvalbumin interneurons drives synaptic inhibition in both GCs and SGC.

Since sIPSCs represent synaptic inhibition in the absence of afferent inputs, we examined

whether direct optical activation of PV-IN evoked IPSCs (oeIPSCs) in GCs and SGCs in slices

from mice expressing channelrhodopsin (ChR2) in PV-INs. Following baseline recordings of

sIPSCs in glutamate antagonist without light, recordings were continued during optical activation

of ChR2 in PV-IN using blue light (λ=473nm, 10 s). Optical activation of PV-INs elicited a

significant increase in IPSC frequency in GCs (Fig. 5A-C; frequency in Hz, no-light: 8.45±1.31,

light: 52.11±5.92, N=14/ 8 mice, p=2E-06, paired t-test) confirming the robust PV-IN input to

GCs. PV-IN activation increased IPSC frequency in SGCs as well (Fig. 5A-C: frequency in Hz,

no-light: 13.68±1.20, light: 59.49±4.62, N=14/ 9 mice, p=2E-07, paired t-test), demonstrating

functional PV-IN innervation of SGCs. However, IPSC peak amplitude was not increased by

optical activation of PV-INs in either GCs or SGCs (Fig. 5D-F). These data demonstrate that

SGCs, like GCs, receive PV-IN inputs and that direct activation of PV-INs leads to an increase in

synaptic inhibitory events in both GCs and SGCs. Since PV-INs are known to be recruited

during PP activation and support feedforward and feedback GC inhibition (Ewell and Jones,

2010), we examined whether recruitment of inhibition by afferent stimulation differed between

the cell types. Single PP stimulation at increasing intensities evoked IPSCs in both SGCs and

GCs. The peak amplitude of afferent evoked IPSCs was not different between cell types (Fig.

5E-F, N=12 cells each from 8 mice, F(1,10)=0.874, p=0.37 by Two-Way Repeated Measures

ANOVA), demonstrating that both GCs and SGCs receive similar early feedforward/feedback

inhibition upon single PP stimulation.

334 Afferent stimulation drives greater enhancement of sustained feedback inhibition in

335 granule cells.

A functional characteristic of SGCs, not observed in GCs, is their unique ability for sustained

firing in response to paired afferent stimuli in rat hippocampal slices, which has been proposed to

drive feedback inhibition and sculpt pattern separation (Larimer and Strowbridge, 2010). We

examined the reliability and reproducibility of SGC sustained firing in SGCs in mice. As is

typical for GCs, a single PP stimulus evoked a single action potential in GCs (3/3 cells tested) as

well as in SGCs (4/4 cells tested) but failed to elicit sustained spiking (not shown). However,

when the PP was stimulated with a pair of suprathreshold stimuli at 10 ms interval, consistent

with theta frequency inputs (Larimer and Strowbridge, 2010; Braganza et al., 2020), a subset of

SGCs showed prolonged spiking activity. Sustained firing was observed in 51.51% of the trials

(Fig 6A-B, 17/33 of trials in 2/4 SGCs). The sustained firing was initiated between 200ms to 2s

after the paired stimulation, although the latency to sustained firing was over 6s in a few trials

indicating that the sustained firing was mediated by network effects (Fig. 6C, median 1.14s IQR:

0.61-1.9s). This evoked sustained firing in SGC lasted several seconds, at times until the end of

the recording 20s after stimulation (Fig. 6D). On average, the sustained firing lasted 15s, which

is likely an underestimate due to the termination of recordings. None of the GCs tested exhibited

firing beyond the early evoked firing in any of the trials (0/12 trials, 3 GCs).

352 Sustained firing in SGCs has been correlated with hilar mossy cell and interneuron firing as well

353 as reduced GC excitability (Larimer and Strowbridge, 2010). However, whether GCs receive

354 enhanced synaptic inhibition during this period and whether SGCs undergo a parallel sustained

355 increase in feedback synaptic inhibition is not known. Using simultaneous recordings in GC-

356 SGC pairs (Fig 6E), we directly compared recruitment of polysynaptic sustained inhibition in

357 GCs and SGCs following paired PP stimulation. Representative image of cells filled during

358 paired recordings show the GC with soma located in the cell layer and SGC somata in the inner

359 molecular layer (Fig. 6G). Paired PP stimuli elicited a large early eIPSC peak (Fig. $6E_1$ insets)

360 followed by a prolonged barrage of "evoked-sustained" IPSCs (esIPSCs) lasting over 20s in both

361 GCs and SGCs (Fig. $6E_1$). To better characterize the evoked sustained activity, we binned IPSCs

362 into 5s segments before the stimulus and 1s segments 500ms-10.5s after the stimulation, with

363 additional 5s segments 15-20s and 25-30s post-stimulus (Fig. 6F). The 500ms immediately

364 following stimulation was not included in the analysis in order to avoid the period of maximum

365 evoked IPSC conductance which included summated events and obscured individual events.

366 The peak IPSC frequency following paired PP stimulation (500-1500 ms) showed high

367 variability and was not different between cell types (in Hz, GC: 64.7±4.3; SGC: 72.9±5.3, n=12

368 each p>0.05 by unpaired Student's t-test). Although the sIPSC frequency prior to stimulation

369 was stable within each cell, there was considerable cell-to-cell variability. To normalize for the

370 variability and assess relative change in esIPSC frequency, the difference in IPSC frequency

371 following stimulation was normalized to the pre-stimulus sIPSC frequency averaged over 3 X 5-

372 sec pre-stimulus bins ((post-pre)/pre*100). Both GCs and SGCs showed a significant post-

373 stimulus increase in esIPSC frequency lasting over 20s. In addition to the significant effect of

374 time, the difference between cell types was statistically significant (Fig. 6F, $F_{(1,11)}$=9.4, for effect

of time and $F_{(1,11)}=4.9$ for cell-type, p<0.05 by Two-way RM ANOVA). Pairwise comparison

between cell types, revealed that the increase in esIPSC frequency normalized to pre-stimulus

frequency was significantly greater in GCs than in SGCs. Both GCs and SGCs showed increase

in esIPSC frequency for up to 10s post stimulus reaching similar maximum frequency and

declining back to pre-stimulus levels by 20-30s. However, the relative increase in esIPSC

frequency (normalized to pre-stimulus frequency) quantified over the first 500-1500ms (1s) after

stimulation was significantly higher in GCs than in SGCs (p<0.05, by TW-RM ANOVA

followed by Pairwise Multiple Comparison using Holm-Sidak method).

Interestingly, not all cells showed robust esIPSC barrages even within the same slice, suggesting

cell specific recruitment of feedback inhibition. We found that 16.67% (2/12) GCs and 33.33%

(4/12) SGCs failed to show esIPSC barrage, defined as doubling in basal frequency in the first 3

post-stimulus bins examined (3s). Additional recordings conducted in the presence of the

GABA$_A$ receptor antagonist, gabazine (10 µM) confirmed that, like eIPSC, esIPSCs were also

eliminated by gabazine (not shown). These results demonstrate that paired PP stimulation results

in prolonged changes to dentate inhibition and SGC excitability. Crucially, the data reveal that

the relative increase in sustained feedback inhibition in SGCs is less than in GCs and suggest that

in spite of their higher basal synaptic inhibitory tone, SGC show lower increase in inhibitory

gating than GCs following activation of inputs.

Role of PV-INs in sustained feedback inhibition of dentate projection neurons following

paired afferent stimulation.

Since PV-INs are known to mediate strong granule cell feedforward and feedback inhibition

(Hefft et al., 2002), we examined whether they exhibit sustained firing in response to paired PP

stimuli coinciding with increased esIPSCs in GCs. PV-INs in the hilar granule cell border were

identified based on YFP labeling and confirmed by the high frequency, non-adapting firing (Fig.

7A$_1$, insets). PV-IN voltage traces in responses to paired PP stimulation (Fig. 7A) are illustrated

alongside the time course of esIPSC in a GC (Fig. 7D) for reference. A subset of PV-INs (4/11

cells, 36.4%) exhibited spontaneous action potential firing at rest under our recording conditions.

Interestingly, only 54.5% (6/11 cells) PV-INs showed sustained increase in firing upon paired

PP stimulation. An additional 18.2% (2/11 cells) PV-INs received increased EPSC drive during

this period suggesting that they may be driven by SGC firing (Fig. 7B). However, 27.3% (3/11)

PV-INs showed no apparent sustained change in EPSC frequency upon PP stimulation in spite of

the early evoked firing. The prolonged PV-IN firing followed the initial stimulation with a

median latency of 1.02s (IQR 0.62-2.03s) which was similar to the latency to evoked SGC firing

(Fig. 7C). Since SGCs and not GCs show evoked sustained firing following paired perforant path

stimulation, these data suggest that the sustained firing in SGCs likely drives PV-IN firing.

These results raise the possibility that PV-IN firing contributes to a certain component of

esIPSCs in GCs and SGCs.

To verify whether halorhodopsin suppression could reduce PP-evoked PV-IN firing, we

performed control recordings from YFP-labeled neurons in PV-NpHR3 mice. While optical

suppression failed to fully eliminate firing of PV-INs in response to paired PP stimulation it

consistently reduced firing without altering firing in hilar neurons lacking YFP expression (Fig.

8A-D). Additionally, we verified that optical suppression of PV-INs did not induce persistent

firing in GC or eliminate sustained firing in SGCs (Fig. 8E).

To directly test the contribution of PV-INs to esIPSC barrages in GCs and SGCs, we recorded

esIPSCs in putative GC/SGC pairs in PV-NpHR3 mice in the absence or presence of amber light

(2s exposure starting 100ms after the onset of PP stimulation) (Fig. 9A-C). Light-on and light-off

conditions randomized (Fig. 9, trials with red lined in Fig. 9D indicate light-on first recordings).

Surprisingly, esIPSC frequency, normalized to basal sIPSC frequency prior to stimulation, was

not reduced by optical suppression of PV-INs (Fig. 9D: esIPSC frequency as % of pre-stimulus

frequency, no-light: 264.59±49.76, light: 225.85±43.08, N=8, p>0.05, by paired t-test). In

contrast, recordings from SGCs showed a significant reduction in esIPSC frequency during

suppression of PV-INs with amber-light (Fig. 8D: esIPSC frequency as % of pre-stimulus

frequency, no-light: 209.76±57.92, light: 127.86±41.15, N=8, p= 0.015 by paired t-test). Thus,

optical suppression of PV-INs selectively reduced esIPSC frequency in SGCs to approximately

half of that observed in the absence of light demonstrating fundamental differences in inhibitory

regulation of GCs and SGCs during evoked activity.

Discussion:

Here we identify that lateral sustained feedback inhibition, which has been proposed to shape dentate pattern separation and memory engram formation, is significantly lower in dentate SGCs than in GCs. We further show that parvalbumin interneurons selectively support activity-driven sustained lateral inhibition of SGCs, but not GCs. The difference in local circuit feedback inhibitory regulation of SGCs and GCs during afferent activation could shape their distinct role in dentate processing.

SGCs, originally defined by Ramón y Cajal (1995) are a sparse, functionally distinct subpopulation of dentate output neurons with CA3 projections (Williams et al., 2007; Larimer and Strowbridge, 2010; Gupta et al., 2012; Save et al., 2018). Temporally structured inhibition by diverse interneurons projecting across the somato-dendritic axis help maintain the sparse granule cell activity needed for pattern separation (Arima-Yoshida et al., 2011; Dengler and Coulter, 2016; Hainmueller and Bartos, 2020). Granule cells receive distinct phases of synaptic inhibition including steady-state spontaneous synaptic inhibition, afferent evoked mono- and disynaptic feedforward and feedback inhibition, as well as extrasynaptic inhibition (Soltesz et al., 1995; Kraushaar and Jonas, 2000; Stell et al., 2003). Additionally, afferent stimulation has been proposed to recruit sustained polysynaptic feedback GC inhibition (Larimer and Strowbridge, 2010). Focusing on synaptic inhibition, we demonstrate three functionally distinct phases of synaptic inhibition, namely, basal, early evoked, and sustained feedback synaptic inhibition in both GCs and SGCs and characterize the temporal profile and reliability of dentate evoked-sustained inhibition. We demonstrate that evoked-sustained feedback inhibition is not unique to GC but is present in SGCs as well. However, given the time dependent changes in evoked-sustained inhibition and robust summation of robust IPSC summation, we did not evaluate

activity dependent changes in extrasynaptic GABA currents which are best examined in steady

state. Although SGCs receive more frequent basal sIPSCs, GCs show greater enhancement of

sustained feedback IPSCs upon afferent activation. These data identify that compared to GCs,

SGCs exhibit blunted elevation in inhibitory gating following input activation, which likely

facilitates their ability to sustain prolonged firing in response to afferent activation.

Although GCs and SGCs differ in their development, structure and excitability (Williams et al.,

2007; Gupta et al., 2012; Save et al., 2018; Gupta et al., 2020), neurochemical markers to

distinguish the cell types are lacking and morphology, characterized in rats, is used as to classify

SGCs and GCs. Whether the neuropeptide proenkephalin (PENK), expressed in some prox-1

positive dentate neurons tagged based on increases in the immediate early gene FOS (Erwin et

al., 2020), is selective for SGCs remines to be determined. However, our clustering analysis

based on morphometric parameters, and the PCA of underlying features validate the use of

number of primary dendrites (>3), dendritic angle, and soma-aspect ratio to distinguish SGCs

from GCs. Moreover, dendritic complexity, which could influence intrinsic physiology and input

integration (Gulledge et al., 2005; van Elburg and van Ooyen, 2010; van der Velden et al., 2012),

was significantly lower in SGCs than in GCs, as reported in rats (Gupta et al., 2020). Curiously,

putative GCs segregated into two subclusters, based on dendritic complexity, dendritic angle,

and number of primary dendrites raising the possibility that physiology of dentate projection

neuron subtypes scales with number of primary dendrites and dendritic complexity.

Nevertheless, the physiological differences between GCs and SGCs identified here and in earlier

studies (Williams et al., 2007; Larimer and Strowbridge, 2010; Gupta et al., 2012; Save et al.,

2018) justify examination of their role in the dentate circuit.

Consistent with prior findings in rats (Gupta et al., 2012; Gupta et al., 2020), SGCs in mice

receive more frequent sIPSCs than GCs. Since SGC somata and dendrites are located in the inner

molecular layer, outside the dense axonal arbor of PV-INs in the granule cell layer, we

anticipated fewer PV-INs inputs to SGCs than to GCs. Surprisingly, halorhodopsin suppression

of PV-INs failed to reduce sIPSC frequency in both GCs and SGCs in spite of reliably

hyperpolarizing (>5 mV) and reducing firing in YFP positive PV-INs. Moreover,

channelrhodopsin activation of PV-INs evoked robust increases in GC and SGC IPSCs

demonstrating viable, synaptically connected PV-INs. Furthermore, the same amber light

protocol reduced evoked sustained IPSCs in SGCs, confirming the ability of halorhodopsin to

suppress of PV-IN firing and synaptic release (Mahn et al., 2016). Why, then, is PV-IN

suppression unable to reduce action potential dependent sIPSCs in GCs and SGCs even in the

absence of glutamatergic blockers? One possibility is that PV-IN suppression disinhibits and

increases firing of a subset of non-PV interneurons which compensate for the loss of PV-IN

mediated inhibition. In this regard, the reciprocal activity states of PV and CCK neurons,

recently reported in CA1, suggests that such a compensation is possible (Dudok et al., 2021).

However, whether such a mechanism would exactly compensate for the reduction in PV-IN

mediated IPSCs in a slice is debatable and cannot be resolved by using chemogenetic strategies

which would potentially suffer from similar network level changes. The inability of

optical/chemogenetic silencing to disambiguate direct from circuit-effects need to be considered

while using cell-specific silencing strategies (Bernard, 2020). Interestingly, a prior study found

limited contribution of PV-INs to action potential independent miniature IPSCs in granule cells

(Goswami et al., 2012) and mutations impairing PV expression were shown to alter granule cell

evoked IPSCs without impacting sIPSC frequency (Lucas et al., 2010). These findings are

consistent with our results and raise the possibility that diverse populations of hilar and

molecular layer interneurons may contribute substantively more than PV-INs to basal sIPSCs in

GCs and SGCs. While our findings do not imply lack of PV-INs contribution to basal inhibition

in vivo, the most parsimonious conclusion from our data is that PV-INs have limited role in basal

sIPSCs in GCs and SGCs in slices.

PV-INs mediate robust and precise feedforward and feedback inhibition of GCs (Andersen et al.,

1966; Hefft and Jonas, 2005; Ewell and Jones, 2010). We find that optical activation of PV-INs

robustly increases IPSC frequency in both SGCs and GCs demonstrating that, despite their

location in the inner molecular layer, SGCs receive inhibitory synapses from PV-INs. Moreover,

afferent evoked IPSC peak amplitude and kinetics of were not different between GCs and SGCs

indicating that, unlike spontaneous IPSCs, inhibition evoked by a single perforant path stimulus

is comparable between the cell types.

Finally, we examined sustained feedback inhibition, previously found to correlate with SGC

firing and advanced as a mechanism for lateral inhibition (Larimer and Strowbridge, 2010). We

confirm that paired suprathreshold PP stimulation elicits persistent SGC firing in about 50% of

trials and was accompanied by a two-fold increase in IPSC frequency in both GCs and SGCs.

Sustained SGC activity followed the early evoke firing by over 200ms and persisted for over 10s.

Interestingly, although individual SGC firing was observed only in half the trials, GCs and/or

SGCs showed evoked sustained IPSCs in all slices examined, suggesting that activation of a

subset of SGCs may be sufficient to sustain feedback inhibition. We report that about half the

PV-INs examined showed increased firing following paired PP stimulation with latency and

duration corresponding to that of evoked sustained inhibition in GCs and SGCs. Notably, even

PV-INs lacking sustained firing showed increased EPSP frequency following paired PP

stimulation indicating enhanced glutamatergic drive. Since SGC axons selectively innervate PV-

IN somata (Rovira-Esteban et al., 2020), our findings are consistent with sustained firing in

SGCs, which does not occur in GCs, driving sustained PV-IN activity.

Although both SGCs and GCs show sustained increase in IPSCs following paired afferent

stimulation, the relative increase in IPSC frequency in SGCs is significantly lower than in GCs.

Thus, SGCs which receive more robust polysynaptic NMDA currents than GCs (Larimer and

Strowbridge, 2010) are, yet, subject to less gating feedback synaptic inhibition placing them in a

privileged position to support persistent firing. This enhanced firing likely promotes expression

of activity dependent genes and recruitment of SGCs as part of cellular ensembles during

behavioral tasks (Erwin et al., 2020). Our analyses identify selective contribution of PV-INs to

sustained feedback inhibition in SGCs but not in GCs. These results are paradoxical since wide

field optical activation of PV-INs increases IPSCs in both SGCs and GCs and raises an

intriguing possibility of target specific differences in PV-IN synaptic release and/or short-term

plasticity similar to those reported in glutamatergic terminals (Sylwestrak and Ghosh, 2012;

Blackman et al., 2013; Larsen and Sjostrom, 2015; Eltes et al., 2017) or selective reciprocal

connectivity between SGCs and a subset of PV-INs.

In conclusion, our study demonstrates an input strength dependent shift in three distinct temporal

phases of synaptic inhibition between GCs and SGCs. During steady-state basal activity, SGCs

receive greater synaptic inhibitory drive than GCs. In response to afferent activation, evoked

synaptic inhibition in SGCs appears comparable to GCs, which, given the greater glutamatergic

inputs reported in SGCs (Larimer and Strowbridge, 2010), represents a relative reduction in

scaling of inhibition to excitation in SGCs. With further increase in drive during paired

stimulation, the increase in sustained feedback inhibition in SGCs is lower than in GCs. This

activity dependent blunting of synaptic inhibitory inputs in SGCs compared to GCs indicates that

SGCs, by having reduced activity dependent inhibitory gating, may support a distinct role in

dentate processing by integrating, rather than sparsifying, inputs and driving lateral inhibition

and selectivity of GCs.

Author Contributions: M.A., B.S performed experiments; M.A., A.G. and H.X. analyzed data;

M.A., H.X., and V.S. interpreted results of experiments; M.A., prepared figures; M.A. and V.S.

conceived of and designed research; M.A., and V.S. drafted manuscript.

Reference List

Alfonsa H, Merricks EM, Codadu NK, Cunningham MO, Deisseroth K, Racca C, Trevelyan AJ

(2015) The contribution of raised intraneuronal chloride to epileptic network activity. J Neurosci

35:7715-7726.

Andersen P, Holmqvist B, Voorhoeve PE (1966) Entorhinal activation of dentate granule cells.

Acta Physiol Scand 66:448-460.

Arima-Yoshida F, Watabe AM, Manabe T (2011) The mechanisms of the strong inhibitory

modulation of long-term potentiation in the rat dentate gyrus. Eur J Neurosci 33:1637-1646.

Bekinschtein P, Kent BA, Oomen CA, Clemenson GD, Gage FH, Saksida LM, Bussey TJ (2013)

BDNF in the dentate gyrus is required for consolidation of "pattern-separated" memories. Cell

reports 5:759-768.

Bernard C (2020) Optogenetics: Keep Interpretations Light. eNeuro 7.

Blackman AV, Abrahamsson T, Costa RP, Lalanne T, Sjostrom PJ (2013) Target-cell-specific

short-term plasticity in local circuits. Front Synaptic Neurosci 5:11.

Braganza O, Mueller-Komorowska D, Kelly T, Beck H (2020) Quantitative properties of a

feedback circuit predict frequency-dependent pattern separation. eLife 9.

Dengler CG, Coulter DA (2016) Normal and epilepsy-associated pathologic function of the

dentate gyrus. Prog Brain Res 226:155-178.

Dudok B, Klein PM, Hwaun E, Lee BR, Yao Z, Fong O, Bowler JC, Terada S, Sparks FT, Szabo

GG, Farrell JS, Berg J, Daigle TL, Tasic B, Dimidschstein J, Fishell G, Losonczy A, Zeng H,

Soltesz I (2021) Alternating sources of perisomatic inhibition during behavior. Neuron 109:997-

1012 e1019.

Eltes T, Kirizs T, Nusser Z, Holderith N (2017) Target Cell Type-Dependent Differences in

Ca(2+) Channel Function Underlie Distinct Release Probabilities at Hippocampal Glutamatergic

Terminals. J Neurosci 37:1910-1924.

Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2020) A Sparse,

Spatially Biased Subtype of Mature Granule Cell Dominates Recruitment in Hippocampal-

Associated Behaviors. Cell reports 31:107551.

Ewell LA, Jones MV (2010) Frequency-tuned distribution of inhibition in the dentate gyrus. The

Journal of neuroscience : the official journal of the Society for Neuroscience 30:12597-12607.

Freund TF, Buzsaki G (1996) Interneurons of the hippocampus. Hippocampus 6:347-470.

GoodSmith D, Chen X, Wang C, Kim SH, Song H, Burgalossi A, Christian KM, Knierim JJ

(2017) Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. Neuron

93:677-690 e675.

Goswami SP, Bucurenciu I, Jonas P (2012) Miniature IPSCs in hippocampal granule cells are

triggered by voltage-gated Ca2+ channels via microdomain coupling. J Neurosci 32:14294-

14304.

Gulledge AT, Kampa BM, Stuart GJ (2005) Synaptic integration in dendritic trees. J Neurobiol

64:75-90.

Gupta A, Elgammal FS, Proddutur A, Shah S, Santhakumar V (2012) Decrease in tonic

inhibition contributes to increase in dentate semilunar granule cell excitability after brain injury.

J Neurosci 32:2523-2537.

Gupta A, Proddutur A, Chang YJ, Raturi V, Guevarra J, Shah Y, Elgammal FS, Santhakumar V

(2020) Dendritic morphology and inhibitory regulation distinguish dentate semilunar granule

cells from granule cells through distinct stages of postnatal development. Brain structure &

function 225:2841-2855.

Hainmueller T, Bartos M (2020) Dentate gyrus circuits for encoding, retrieval and discrimination

of episodic memories. Nat Rev Neurosci 21:153-168.

Hefft S, Jonas P (2005) Asynchronous GABA release generates long-lasting inhibition at a

hippocampal interneuron-principal neuron synapse. NatNeurosci 8:1319-1328.

Hefft S, Kraushaar U, Geiger JR, Jonas P (2002) Presynaptic short-term depression is maintained

during regulation of transmitter release at a GABAergic synapse in rat hippocampus. J Physiol

539:201-208.

Heinemann U, Beck H, Dreier JP, Ficker E, Stabel J, Zhang CL (1992) The dentate gyrus as a

regulated gate for the propagation of epileptiform activity. Epilepsy ResSuppl 7:273-280.

Howard A, Tamas G, Soltesz I (2005) Lighting the chandelier: new vistas for axo-axonic cells.

Trends Neurosci 28:310-316.

Husson F, Josse J, Pages J (2010) Principal component methods - hierarchical clustering -

partitional clustering: why would we need to choose for visualizing data? In: Technical Report –

Agrocampus.

Korgaonkar AA, Li Y, Sekhar D, Subramanian D, Guevarra J, Swietek B, Pallottie A, Singh S,

Kella K, Elkabes S, Santhakumar V (2020) Toll-like Receptor 4 Signaling in Neurons Enhances

Calcium-Permeable alpha-Amino-3-Hydroxy-5-Methyl-4-Isoxazolepropionic Acid Receptor

Currents and Drives Post-Traumatic Epileptogenesis. Ann Neurol 87:497-515.

Kraushaar U, Jonas P (2000) Efficacy and stability of quantal GABA release at a hippocampal

interneuron-principal neuron synapse. J Neurosci 20:5594-5607.

Larimer P, Strowbridge BW (2010) Representing information in cell assemblies: persistent

activity mediated by semilunar granule cells. Nature neuroscience 13:213-222.

Larsen RS, Sjostrom PJ (2015) Synapse-type-specific plasticity in local circuits. Curr Opin

Neurobiol 35:127-135.

Lothman EW, Stringer JL, Bertram EH (1992) The dentate gyrus as a control point for seizures

in the hippocampus and beyond. Epilepsy ResSuppl 7:301-313.

Lucas EK, Markwardt SJ, Gupta S, Meador-Woodruff JH, Lin JD, Overstreet-Wadiche L,

Cowell RM (2010) Parvalbumin deficiency and GABAergic dysfunction in mice lacking PGC-

1alpha. J Neurosci 30:7227-7235.

Mahn M, Prigge M, Ron S, Levy R, Yizhar O (2016) Biophysical constraints of optogenetic

inhibition at presynaptic terminals. Nat Neurosci 19:554-556.

Otis TS, Staley KJ, Mody I (1991) Perpetual inhibitory activity in mammalian brain slices

generated by spontaneous GABA release. Brain Res 545:142-150.

Ramón y Cajal S (1995) Histology of the nervous system of man and vertebrates. New York:

Oxford University Press.

Rovira-Esteban L, Hajos N, Nagy GA, Crespo C, Nacher J, Varea E, Blasco-Ibanez JM (2020)

Semilunar Granule Cells Are the Primary Source of the Perisomatic Excitatory Innervation onto

Parvalbumin-Expressing Interneurons in the Dentate Gyrus. eNeuro 7.

Santhakumar V (2008) Modeling mossy cell loss and mossy fiber sprouting in epilepsy. In:

Computational Neuroscience in Epilepsy (Soltesz I, Staley KJ, eds), pp 89-111: Academic Press.

Save L, Baude A, Cossart R (2018) Temporal Embryonic Origin Critically Determines Cellular

Physiology in the Dentate Gyrus. Cereb Cortex 10.1093/cercor/bhy132.

Sik A, Penttonen M, Buzsaki G (1997) Interneurons in the hippocampal dentate gyrus: an in vivo

intracellular study. European Journal of Neuroscience 9:573-588.

Soltesz I, Smetters DK, Mody I (1995) Tonic inhibition originates from synapses close to the

soma. Neuron 14:1273-1283.

Stell BM, Brickley SG, Tang CY, Farrant M, Mody I (2003) Neuroactive steroids reduce

neuronal excitability by selectively enhancing tonic inhibition mediated by delta subunit-

containing GABAA receptors. Proc Natl Acad Sci U S A 100:14439-14444.

Sylwestrak EL, Ghosh A (2012) Elfn1 regulates target-specific release probability at CA1-

interneuron synapses. Science 338:536-540.

van der Velden L, van Hooft JA, Chameau P (2012) Altered dendritic complexity affects firing

properties of cortical layer 2/3 pyramidal neurons in mice lacking the 5-HT3A receptor. J

Neurophysiol 108:1521-1528.

van Elburg RA, van Ooyen A (2010) Impact of dendritic size and dendritic topology on burst

firing in pyramidal cells. PLoS computational biology 6:e1000781.

Walker MC, Pavlov I, Kullmann DM (2010) A 'sustain pedal' in the hippocampus? Nature

neuroscience 13:146-148.

Williams PA, Larimer P, Gao Y, Strowbridge BW (2007) Semilunar granule cells: glutamatergic

neurons in the rat dentate gyrus with axon collaterals in the inner molecular layer. The Journal of

neuroscience : the official journal of the Society for Neuroscience 27:13756-13761.

Yu J, Swietek B, Proddutur A, Santhakumar V (2015) Dentate total molecular layer interneurons

mediate cannabinoid-sensitive inhibition. Hippocampus 25:884-889.

Yu J, Proddutur A, Swietek B, Elgammal FS, Santhakumar V (2016) Functional Reduction in

Cannabinoid-Sensitive Heterotypic Inhibition of Dentate Basket Cells in Epilepsy: Impact on

Network Rhythms. Cereb Cortex 26:4229-4314.

Figure Legends

Figure 1: GCs and SGCs cluster into distinct subtypes on the basis of morphology. A-D**.**

Example images of GCs (A-B) and SGCs (C-D) illustrate the somatic location and dendritic

spread. Note that reconstruction of neuron in A is presented in Fig 2A. Scale bar: 100µm; E.

Schematic of the somatic location of the cells included in the cluster analysis. The GCL is

illustrated as a green box and IML as an orange box with somata illustrated in blue and GCs in

green. All images were obtained a 20X magnification. GCL: Granule Cell Layer, IML: Inner

Molecular Layer. F. Dendrogram generated by Hierarchical Clustering on Principal Components

(HCPC) based on 42 morphometric parameters (Defined in Gupta et al., 2020) using Ward's

method with Euclidean distance suggests three putative clusters. G. Factor map base on first 7

PCs. H-J. Summary data of dendritic angle (H), number of primary dendrites (I) and Dendritic

Complexity (J) for cells in the three clusters. Data are presented as mean±sem. * indicates

$p<0.05$ by One-way ANOVA followed by pairwise comparison with Holm-Sidak method for H

and $p<0.05$ by One-way ANOVA on Ranks followed by pairwise comparison with Dunn's

method for I and J where the data failed Shapiro-Wilk Normality Test. A. U: Arbitrary Units.

Figure 2. Morphological and physiological characteristics of GCs and SGCs in mice. A-B.

Overlay of confocal image and Neurolucida 360 reconstruction of a biocytin filled GC (A) and

SGC (B) show dendritic arbors and axon projection towards CA3. Cyan: axons; Orange, Green,

Yellow: dendritic trees. C-H. Summary plots of morphological parameters including number of

primary dendrites (C), dendritic angle (D), soma aspect ratio (E), dendritic complexity (F),

dendritic total length (G) and total number of dendritic ends (H) between GCs (n=10) and SGCs

(n=8). I. Voltage traces in response to +70 pA and -100 pA current injections in a GC (left) and

SGC (right) illustrate firing pattern and passive properties. J. Summary plot of firing frequency

in response to increasing current injections in GCs (n=8) and SGCs (n=10). K-L. Summary

histogram of input resistance (measured from response to -100 pA), (K) and spike frequency

adaptation ratio (L) in GCs (n=18) and SGCs (n=19). Data presented as mean±sem. * indicates

$p<0.05$ by independent t-test.

Figure 3. SGCs receive more spontaneous inhibitory synaptic events than GCs. A.

Representative current traces from a holding potential of -60 mV illustrate sIPSCs in a GC

(above) and SGC (below). Text in gray box indicates recording condition. B. Example of

average sIPSC trace from a GC (above) and SGC (below). C-F. Summary histogram of sIPSC

frequency (C), amplitude (D), amplitude weighted decay time constant (E) and 20-80% rise time

(F) in GCs (n=18) and SGCs (n=19). Data presented as mean±sem. * indicates $p<0.05$ by

independent t-test.

Figure 4. Optical suppression of PV-INs fails to reduce sIPSC frequency in both GCs and

SGCs. A. Representative membrane voltage trace (above) from a holding potential of -70 mV in

a YFP-positive PV interneuron expressing halorhodopsin in the granule cell-hilar border

illustrates the high frequency firing during depolarizing current injection (100 pA for 10 sec) and

the ability of amber light (bar above) to suppress firing. Representative trace from 19 trials in 4

cells. Membrane voltage trace from an NpHR3-YFP expressing PV-IN shows rapid and

reversible hyperpolarization (below) during the 2s amber light stimulation. Schematic to the right

illustrates recording condition and optical stimulation. B. Schematic of the experimental

paradigm illustrates amber light stimulation of NpHR3-YFP expressing PV-IN while recording

from GC/SGC (above) and the protocol for light OFF and light ON condition below. C. Current

traces from a GC (above) and SGC (below) recorded from a holding potential of -60 mV using

high-chloride internal shows sIPSCs as inward currents before and during NpHR3 activation

(amber bar) in PV-INs. IPSCs were isolated in kynurenic acid (3 mM). D-E. Plots summarize

effect of optical suppression of PV-IN on sIPSC frequency (D) and amplitude (E) (n=5 GCs and

SGC). F. Representative traces illustrate sIPSCs in a GC (above) and SGC (below) before and

during activation amber light exposure (amber bar). Recordings were obtained using a low-

chloride internal from a holding potential of 0 mV without synaptic blockers. G-H. Summary

plots of effect amber light on sIPSC frequency (G) and amplitude (H) in GCs (n=13) and SGCs

(n=11). Data presented as mean±sem. Text in gray box indicates recording condition.

Figure 5. Activation of PV-INs and afferent stimulation evoke inhibitory synaptic currents

in both dentate projection neuron subtypes. A. Schematic illustrates blue light stimulation of

ChR2-YFP expressing PV-IN while recording from GC/SGC (above) and light OFF and light

ON protocol below. B) Example recordings from a GC (above) and SGC (below) illustrate

sIPSCs before and during optical activation of PV-INs. Cells were held at -60 mV and

kynurenic acid (3 mM) was used to isolate IPSCs. C. Summary plots of sIPSC frequency before

and during blue light exposure in GCs (n=14) and SGCs (n=14). D. Schematic of the

experimental paradigm illustrates PP stimulation while recording from GC/SGC. E. Perforant

path evoked IPSC in GC and SGC recorded as outward currents using a low-chloride internal

and a holding potential of 0 mV. F. Summary plot of eIPSC peak amplitude in GCs and SGCs in

response to stimulation of the PP at increasing current intensities. Data presented as mean±sem.

* indicates $p<0.05$ by paired t-test (in panel C) or unpaired t-test (in panel D). Text in gray box

indicates recording condition.

Figure 6. Afferent evoked prolonged SGC firing is associated with sustained increase in

sIPSC frequency in both GC and SGC. A. Membrane voltage traces in a GC (above) and SGC

(below) evoked by a pair of suprathreshold stimuli to the PP (denoted by 2 arrows). Note

prolonged firing in the SGC while none of the GCs (n=3 cells, 12 trails) showed more than 1-3

spikes. B. Plot shows the proportion of trials that evoked sustained firing in SGCs (n= 4 cells, 33

trials). C-D. Plots illustrate the distribution of latency from afferent stimulation to the start of

sustained firing (C) and duration of sustained firing in SGCs recorded for a maximum of 20s

after stimulation. E. Schematic of paired GC-SGC recordings in response to paired PP

stimulation (E_2). Current traces of simultaneously recorded GC-SGC pair illustrate the sustained

increase in IPSC frequency compared to pre-stimulus baseline (E_1). The eIPSC peak is truncated

to better illustrate the sustained change in IPSCs over 20s. E_1 Insets: traces at a different scale

illustrate the peak eIPSC amplitude. F. Summary plot of sIPSC frequency in GCs and SGCs (n=

12 each) prior to stimulation (binned over 5s) and evoked sustained IPSC frequency (1s bins for

the first 10s and 5sec bins at 20 and 30s) for success rate in induction of prolonged spiking

activity in GCs (12 trials, N=3) and SGCs (33 trials, N=5). G. Maximum intensity projection of a

confocal image stack of a GC-SGC pair filled during recordings (presented as grey scale and

inverted) illustrate dendric arbors and hilar axon (arrows). GCL: granule cell layer; ML:

molecular layer, GC: granule cell; SGC: semilunar granule cell. Data presented as median

(interquartile range) for in C and mean±sem in D and F. * indicates p<0.05 for effect of cell-type

(F(1,11)=4.9) by TW-RM ANOVA. Text in gray box indicates recording condition.

Figure 7. Increase in PV-IN excitability during evoked sustained sIPSCs in dentate

projection neurons. A. Example traces show increased firing (above), enhanced EPSP barrages

(middle) and unaltered EPSP frequency (below) in YFP-positive PV-INs (A$_1$) in response to the

paired PP stimulation paradigm (denoted by 2 arrows) illustrated in A$_2$. Insets in middle and

lower traces illustrate high frequency firing in recorded PV-IN. B. Pie chart summarizes the

proportion of PV-INs that showed evoked sustained firing, EPSP increase or no change in

response to stimulation (n= 11 cells, 33 trials). C. Plot illustrates the distribution of latency of

enhanced PV-IN firing following paired PP stimulation. D. Current traces from a GC showing

sustained IPSCs is included for comparison. Text in gray box indicates recording condition.

Figure 8: Cell specificity of halorhodopsin suppression of PV neuron firing. A.

Representative membrane voltage traces from a YFP-positive PV interneuron expressing

halorhodopsin in the granule cell-hilar border in response to a pair of suprathreshold perforant

path stimuli (denoted by 2 arrows) at 10 ms interval illustrated the early evoked firing in PV

neurons (above). Effect of amber light exposure on perforant path evoked PV-IN firing is

illustrated (below). B. Example voltage traces from a YFP-negative hilar neuron in response to a

pair of suprathreshold perforant path stimuli at 10 ms interval (above). Effect of amber light

exposure on perforant path hilar neuron firing is illustrated (below). C. Summary data of number

of spikes elicited following the first and second stimuli and the effect of halorhodopsin activation

on perforant path evoked firing in PV-IN. D. Summary plot compares the effect of amber light

on afferent evoke firing in halorhodopsin expressing PV-IN and YFP-negative hilar neurons. *

Indicates p<0.05 by paired t-test. Data presented as mean±sem. E. Representative traces show

lack of sustained firing in GCs and sustained firing in SGCs following paired suprathreshold

perforant path stimulation. Text in gray box indicates recording condition.

Figure 9. Optical suppression of PV-INs selectively reduces evoked sustained IPSCs in

SGCs. A_1. Schematic illustrating the simultaneous dual GC-SGC recordings in response to

paired stimulation of the PP and amber light stimulation (above). GC current trace illustrates the

sustained increase in IPSC frequency compared to pre-stimulus baseline (below). A_2. GC trace

shows that evoked sustained IPSCs in response to paired PP stimuli (denoted by 2 arrows)

appears unchanged by amber light suppression (amber bar) of PV-INs. B_1. SGC current trace

illustrates the sustained increase in IPSC frequency compared to pre-stimulus baseline. B_2. SGC

trace shows that evoked sustained IPSCs in response to paired PP stimuli are suppressed by

amber light. Note that the eIPSC peak is truncated to better illustrate the sustained change in

IPSCs over 20s. C. Maximum intensity projection of a confocal image stack of a GC-SGC pair

filled during recordings illustrated the soma location dendric arbors and hilar axon (arrows) of

the cells. The image is presented as grey scale and inverted to better illustrate the axon. GCL:

granule cell layer; ML: molecular layer, GC: granule cell; SGC: semilunar granule cell. D.

Summary plots of effect of amber light suppression of PV-INs on evoked sustained IPSC

frequency in GCs and SGCs (n=8 each). Red lines indicate trials with light-on first recordings.

Data presented as mean±sem. * Indicates p<0.05 by paired t-test. Text in gray box indicates

recording condition.

Tables

Table 1: Summary of morphometric and physiology data in GC and SGCs.

A PV-IN

V = -70 mV

20 mV

500 mS

V = -70 mV

PV-IN

5 mV

2 S

B

Baseline   2 S 50 S 2 S 50 S 2 S 50 S 2 S 50 S 2 S

C High Chloride Internal

3mM Kynurenic Acid

GC V_H = -60 mV

SGC V_H = -60 mV

20 pA

500 mS

D

E

F Low Chloride Internal

No Blockers

GC V_H = 0 mV

20 pA

500 mS

SGC V_H = 0 mV

G

H

A

Baseline   10 S

B

High Chloride Internal
3mM Kynurenic Acid

GC
$V_H = -60$

SGC
$V_H = -60$ mV

20 pA
500 mS

C

sIPSC Frequency (Hz)

No Light   Light   No Light   Light
      GC   SGC

*   *

D

Perforant path

E

Low Chloride Internal
No Blockers

GC   SGC

$V_H = 0$ mV $V_H = 0$ mV   200 pA
                                                    100 mS

F

eIPSC Amplitude (pA)

Stimulus Intensity (µA)

GC
SGC

A₁

GC

V_H = 0 mV

Low Chloride Internal
No Blockers

Perforant path

SGC

GC

ML

GCL

PV+

Hil

50 pA | 2 S

A₂

GC

V_H = 0 mV

50 pA | 2 S

C

ML

GCL

SGC

Hilus

GC

100 µM

B₁

SGC

V_H = 0 mV

100 pA | 2 S

B₂

SGC

V_H = 0 mV

100 pA | 2 S

D

IPSC Frequency Change (%)

*

No Light | Light | No Light | Light

GC | SGC

Table 1: Summary of morphometric and physiology data in GC and SGCs.

Fig #	Parameter (unit)	Cluster 1 (5 cells/ 8 mice)	Cluster 2 (4 cells /8 mice)	Cluster 3 (8 cells/ 8 mice)	F and P Value
1H	Dendritic Angle (Degree)	56.28±2.84	80.21±10.82	105.12±6.17	$F(2,14)=16.13$ p <0.001 by One Way ANOVA Clu1 v Clu2 =0.038 Clu1 v Clu3<0.001 Clu2 v Clu3=0.032
1I	# of Primary Dendrites (Count)	1.00±0.00	2.00±0.00	4.60±0.40	<0.001 by Kruskal-Wallis ANOVA on Ranks Clu1 v Clu2 =0.038 Clu1 v Clu3<0.001 Clu2 v Clu3=0.032
1J	Dendritic Complexity (A.U.)	204147±27920	61393±11452	30435±4609	0.002 by Kruskal-Wallis ANOVA on Ranks Clu1 v Clu2 >0.05 Clu1 v Clu3<0.05 Clu2 v Clu3>0.05
Fig #	**Parameter (unit)**	**GC**		**SGC**	**P Value by Independent Student's t-test**
2C	# of Primary Dendrites (Count)	1.50±0.17		4.5±0.33	1.90E-07
2D	Dendritic Angle (Degree)	69.00±5.98		106.23±5.71	0.0004
2E	Soma Aspect Ratio	0.60±0.05		1.08±0.13	0.001
2F	Dendritic Complexity (A.U.)	143653.30±26506.76		30290.86±3515.42	0.002
2G	Total Dendritic Length (µM)	2147.37±128.20		2269.97±126.30	0.512
2H	Total Dendritic Ends (Count)	16.90±1.13		16.87±0.81	0.986
2K	Input Resistance (MΩ)	309.89±19.58		229.33±29	0.031
2L	Spike Frequency Adaptation Ratio	0.45±0.05		0.68±0.07	0.023