Tetraspanin Cd9b and Cxcl12a/Cxcr4b have a synergistic effect on the control of collective cell migration

Marsay KS [1,3,4¶], Greaves S[2¶], Mahabaleshwar H[5], Ho CM[5], Roehl H[2*], Monk PN[3], Carney T. J.[4,5] and Partridge LJ[1]

[1]Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, UK

[2]Department of Biomedical Science University of Sheffield, Sheffield, UK

[3]Department of Infection, Immunity and Cardiovascular Science, University of Sheffield, Sheffield, UK.

[4]Institute of Molecular and Cell Biology, Agency for Science, Technology and Research, 61 Biopolis Drive, 138673, Singapore

[5]Lee Kong Chian School of Medicine, Experimental Medicine Building, Yunnan Garden Campus, 59 Nanyang Drive, Nanyang Technological University, 636921, Singapore

*Corresponding authors

Email: h.roehl@sheffield.ac.uk (HR)

¶ KM and SG are joint first authors and contributed equally to this work.

Abstract

Collective cell migration is essential for embryonic development and

homeostatic processes. During zebrafish development, the posterior lateral line

primordium (pLLP) navigates along the embryo flank by collective cell migration. The

chemokine receptors, Cxcr4b and Cxcr7b, as well as their cognate ligand, Cxcl12a,

are essential for this process. We corroborate that knockdown of the zebrafish *cd9*

tetraspanin orthologue, *cd9b*, results in mild pLL abnormalities. Through generation

of CRISPR and TALEN mutants, we show that *cd9a* and *cd9b* function partially

redundantly in pLLP migration, which is delayed in the *cd9b* single and *cd9a; cd9b*

double mutants. This delay led to a transient reduction in neuromast numbers. Loss

of both Cd9a and Cd9b sensitized embryos to reduced Cxcr4b and Cxcl12a levels.

Together these results provide evidence that Cd9 modulates collective cell migration

of the pLLP during zebrafish development. One interpretation of these observations

is that Cd9 contributes to more effective chemokine signalling.

Introduction

Cells can migrate individually or in groups, the latter is known as collective

cell migration. During this process, cells exhibit coordinated behaviour, group

polarisation and maintain cell-cell contacts [1]. This mode of migration is employed

during embryonic development in the morphogenesis of multiple organ systems and

is also important for effective immune responses. This mechanism contributes to

several diseases including metastatic cancer and rheumatoid arthritis [2].

The zebrafish lateral line consists of a series of mechanosensory organs

(neuromasts), which are distributed along the lateral surface of the zebrafish body

and connected to the central nervous system by afferent axons [3]. This arrangement

of innervated neuromasts is achieved by the collective migration of a placodal

primordium of approximately 100 cells, arising just posterior to the otic placode [4].

This posterior lateral line primordium (pLLP), then migrates along the horizontal

myoseptum to the tip of the tail. During migration, clusters of cells are deposited from

the trailing end of the pLLP, which ultimately differentiate into neuromasts.

Primordium migration is directed by expression of the chemokine Cxcl12a

along the horizontal myoseptum, which is received by two chemokine receptors

expressed within the primordium [5–7]. Cxcr4b is expressed in the leading two thirds

of the primordium, where ligand binding induces $G\beta1$ signalling and actin

polymerisation to promote a migratory phenotype [8]. Cxcr7 is expressed in the

anterior-most third and functions as a ligand sink, allowing the formation of a local

Cxcl12a gradient. This results in directed migration of the primordium along the

Cxcl12a pathway towards the caudal fin [9,10]. There is evidence that the membrane

environment of chemokine receptors, including Cxcr4 and Cxcr7, strongly influences

their signalling properties. This includes homodimerisation, oligomerisation and

heteromerisation with other membrane receptors [11]. In addition, chemokine

receptors have been linked to membrane microdomains where the lipid environment

strongly modulates function [12].

Tetraspanins are a large family of small integral membrane proteins, which

have been shown to organise neighbouring membrane proteins into complexes

called tetraspanin enriched microdomains. This is often referred to as the tetraspanin

web, as different interactions build to form dynamic signalling networks that often

induce similar functional outcomes [13]. Thus, tetraspanins are associated with a

wide variety of cellular functions including signalling and cell migration [14]. For

example, the tetraspanin CD9 is downregulated in many human cancers including

lung, breast and ovarian, and reduced CD9 expression is related to colon cancer

metastasis [15]. In particular, CD9 has been shown to regulate, and be regulated by,

CXCL12-CXCR4 signalling [16–18]. CD9 was found in close proximity to CXCR4 on

the membrane of B acute lymphoblastic leukaemia cells in vitro, and enhanced their

CXCL12 dependent migration [18]. If regulation of CXCL12/CXCR4 signalling by

CD9 is conserved, we hypothesised Cd9 would be strongly expressed in the

zebrafish pLLP, where it might modulate Cxcr4 signalling and thus pLLP migration.

Indeed microarray experiments have localised the zebrafish *cd9b* paralogue to the

migrating primordium, whilst a morpholino (MO) experiment indicated loss of *cd9b*

altered primordium structure at 36 hours post fertilisation (hpf) and reduced

neuromast number at 48 hpf [2].

In this study we aimed to investigate further the role of zebrafish CD9

orthologues in Cxcr4b-mediated pLLP migration, through use of genetic knockouts.

We show expression of both *cd9a* and *cd9b* paralogues in the primordium and

confirmed the abnormal pLL phenotype seen previously with *cd9b* knockdown. The

cd9b mutants showed delayed pLLP migration but did not fully replicate *cd9b*

morphant pLL phenotypes. We therefore generated *cd9a; cd9b* double knockouts

(*cd9* dKOs). These showed both reduced migration of the pLLP and increased

sensitivity to reduced levels of Cxcr4b and Cxcl12a, supporting a role for CD9 in

regulation of Cxcr4 signalling.

Methods

Zebrafish maintenance

Adult wild-type zebrafish (WT), *Tg(-8.0cldnb:LY-EGFP)*[zf106] and *cd9a/b*

mutants were housed and bred in a regulated 14:10 hour light: dark cycle under UK

Home Office project licence 40/3459 in the Bateson Centre aquaria at the University

of Sheffield or in the Singapore IMCB zebrafish facility under the Biological Resource

Centre oversight with project license IACUC 140924. Zebrafish were raised under the

standard conditions at 28°C [19]. Ages are expressed as hours (hpf) or days (dpf)

post fertilisation.

Morpholino injection

Antisense morpholino oligonucleotides (MO) were obtained from GeneTools

LLC and re-suspended in MilliQ H_2O to give a stock concentration of 1 mM and

injected in one cell stage embryos. A Flaming/Brown micropipette puller was used

to create micro-injection needles from borosilicate glass capillary tubes (0.5 mm inner

diameter, Sutter). The PV800 Pneumatic PicoPump, as part of the micro-injection jig,

was set up to release the required amount of injection material by adjusting the air

pressure and air expulsion time. For the knockdown of *cd9b* two MOs were designed,

a translation blocker with the following sequence (*cd9b* MO1): 5'-

tttatgaggagaaacccaagactga-3' and a splice site blocker (*cd9b* i2e3) with the following

sequence: 5'-aacccctgaacacagagaaacaaca-3', whilst the published mismatch MO

was used 5'- tttccctgctgcttatacagcgatg -3' [20]. For knockdown of *cxcr4b* and

cxcl12a, the following sequences were used respectively, 5'-

aatgatgctatcgtaaaattccat-3'and 5'-ttgagatccatgtttgcagtgtgaa-3' [21].

Zebrafish mutant production

cd9b mutants were created from WT embryos using transcription activator-

like effector nucleases (TALEN) and maintained on an WT background. TALEN

(ZGene Biotech Inc., Taiwan) were provided in a pZGB4L vector, targeting the *cd9b*

sequence 5'-ttgctctttatcttca-3'. Injected embryos were outcrossed and sequenced to

identify mutations. A frameshifting deletion mutation (c.42_49del) was selected that

caused premature termination within the first transmembrane domain (*cd9b*[pg15]allele).

CRISPR-Cas9 was used to create *cd9a* mutants using the gRNA sequence 5'-

gagtgtatatcctcattgcgg-3', which targeted the 3rd exon encoding part of the second

transmembrane domain. *cd9* dKO mutants were created by injecting the above *cd9a*

gRNA and Cas9 RNA into *cd9b^{pg15}* embryos. These fish were screened for germline

transmission by sequencing and backcrossed to *cd9b^{pg15}* mutants. An indel mutation

deleting 4bp and inserting 8bp (c.180_187delinsTCGCTATTGTAT; *cd9a^{la61}*)

generated a frameshift mutation resulting in a premature stop codon in exon 3, which

was predicted to truncate the protein before the large extracellular domain.

Heterozygous fish of the same genotype were incrossed and adult F2 fish were

genotyped to identify homozygous *cd9b^{pg15}; cd9a^{la61}* (*cd9* dKO).

In situ hybridisation

Embryos were raised at 28°C in petri dishes containing E3 solution. The E3

was changed daily, and any dead embryos removed. At 30-32 hpf, embryos were

anaesthetised using tricaine and dechorionated before returned to fresh E3 solution.

At the relevant timepoint embryos were fixed overnight at 4°C using 4 % (w/v)

paraformaldehyde (Sigma-Aldrich, UK) in phosphate buffered saline (PBS). Embryos

were washed in PBS/0.05 % (v/v) Tween 20 (PBST), then put through a

methanol/PBS series using 30 %, 60 % and 100 % (v/v) methanol before being

stored in 100% methanol (Sigma-Aldrich) at -20°C. In situ hybridisation (ISH) was

carried out as described by [22], except for the embryo digestion with proteinase K,

for which 30-32 hpf embryos were digested with 10 mg/ml proteinase K at 20°C for

22 min. Primers used for PCR generation of the in situ probes are given in Table 1

below. The protocol was conducted with the embryos in 1.5 ml microfuge tubes for

the first two days, after which they were held in 12-well plates for staining before

transferring back to microfuge tubes for storage. Stained embryos were stored in the

dark in 80% (v/v) glycerol.

Table 1. Primers for making ISH probes.

Probe	Primer name	Sequence (5'-3)
claudin b	Claudinb F	aaacgaaaaagcatggcatc
claudin b	Claudinb R	gaggctgtttcaaacgtggt
cd9a	CD9a F	gtcatattcgcggtcgaagt
cd9a	CD9a R	ctgcgagaacaacaaagcaa
cd9b	CD9b F	gttcgccacaagtgcctgat
cd9b	CD9b R	tacatgttactttctctccaaacaat

Time lapse imaging

Time-lapse recording was performed using an inverted Zeiss LSM700

Confocal microscope. All larvae were anaesthetised using 0.02% tricaine and then

embedded on their side in 1% low melting agarose (Lonza) on a glass bottom dish

(MalTEK Corporation) and covered with E3 supplemented with 0.02% tricaine.

Separate Z-stack images covering the depth of the horizontal myoseptum were taken

at specified intervals over a specified period and assembled into a final movie at a

specified frame rate.

Statistics

Data distribution was first assessed for normality using a D'Agostino-Pearson

omnibus K2 normality test on the experimental residuals, as well as creating a

histogram of residuals. For normally distributed data, an ANOVA with Dunnet's or

Holms-Sidak multiple comparisons tests were used. For non-normally distributed

data non-parametric tests, the Mann-Whitney U test or Kruskal-Wallis with Dunn's

multiple comparisons test, were used.

Results

Both *cd9* zebrafish paralogues are expressed in the lateral line

Zebrafish possess two Cd9 paralogues, Cd9a [NP_997784] and Cd9b [NP_998593] which show 60% and 59% identity to human CD9 respectively and 63% identity to each other using multiple sequence comparison by log‑expectation (MUSCLE) [23]. Whole mount in situ hybridisation (WISH) using probes against *cd9a* and *cd9b* demonstrated expression of both paralogues in the migrating primordium at 36 hpf (Figs 1A-D). Expression of both *cd9a* and *cd9b* was also observed in recently deposited neuromasts and was retained in neuromasts until at least 5 dpf (Figs 1G-L).

Fig 1. *cd9a* and *cd9b* are expressed in the pLLP during zebrafish development. A-F: Micrographs of WISH of *cd9a* and *cd9b* at 36 hpf with sense and anti-sense probes. (a-b) Overview shows staining in pLL; (c-d) higher resolution images show expression in the primordium; (e-f) sense probes show no staining. G-J: WISH of *cd9a* and *cd9b* at 48 hpf; (g-h) overview shows staining in pLL; (i-j) higher resolution images show expression in a neuromast. K-L: WISH of *cd9a* and *cd9b* at 5 dpf in a neuromast. Arrows indicate primordium (black) and neuromasts (red). Scale bar indicates 50 μm (white).

cd9b morphant phenotype

To evaluate if there was a role for either paralogue in pLLP migration, we initially focused on *cd9b* as its role in the primordium had been previously reported [2]. Two different MOs were designed to target *cd9b*; *cd9b* MO1 was a translation blocking MO which targeted the 5' UTR of *cd9b* RNA and *cd9b* i2e3 was a splice

blocking MO designed against the intron 2 - exon 3 splice site. These MOs were

injected independently into 1-cell stage embryos to ensure that they produced the

same phenotype. Embryos injected with either *cd9b* MO1 or *cd9b* i2e3, but not the

mismatch or uninjected embryos, showed a significant decrease in neuromasts

deposited (Fig 2). For embryos injected with *cd9b* MO1, there was a decrease in the

percentage of trunk length between the first and last neuromasts (Fig 2I). This

suggests that the primordium is stalling prematurely or migrating more slowly. This

recapitulates and expands on the work by Gallardo et al., 2010 which suggested a

role for Cd9b in lateral line development [2].

Fig 2. *cd9b* MO caused a decrease in neuromasts deposited at 52 hpf. A: Intron

exon diagram of the *cd9b* gene with MO target sites. B-G: Representative images of

WISH of *claudin b* at 52 hpf of (b) uninjected embryos, (c) embryos with 0.5 µM

Mismatch, (d) 0.1 µM *cd9b* MO1, (e) 0.1 µM *cd9b* i2e3 MO1, (f) 0.2 µM *cd9b* MO1,

(g) 0.2 µM *cd9b* i2e3 MO. H: Graph showing the number of neuromasts deposited at

52 hpf is significantly decreased after injection with both *cd9b* MO1 and *cd9b* i2e3

MO, with as low as 0.1 µM MO. I: Graph showing the percentage of trunk between

the first and last deposited neuromast is decreased in *cd9b* MO1 morphants but not

cd9b i2e3 morphants. ANOVA with Dunnett's multiple comparisons test was used.

n= minimum 12 lateral lines analysed per condition. ****p=<0.05.

cd9b mutant phenotype

To confirm this MO result, we created a TALEN mediated knockout of *cd9b.* A

cd9b TALEN pair was designed by ZgeneBio using the program "TEL Effector

Nucleotide Targeter 2.0". The TALEN pair was designed to target Exon 1 and

predicted to cut in the 1st transmembrane domain (Fig 3A). The TALEN pair was

injected and after 72 hpf a proportion of embryos were genotyped. Injected embryos

showed clear mosaicism around the TALEN cut site so embryos were raised to

create mosaic adult F0s. F0s transmitting a mutation to offspring were out-crossed

with WT fish and the resulting F1 offspring were raised. To create a *cd9b*

homozygous mutant line with a single mutant allele, F1s with a c.42_49del mutation

were selected, incrossed and the resulting offspring raised to adulthood. This allele

(*cd9b^{pg15}*) was used as it had the largest deletion and caused the earliest nonsense

stop codon. The 8bp deletion in the *cd9b* allele leads to a frameshift in exon 1,

changing codon 15 from TTT (Phe) to CAA (Glu), then 22 aberrant amino acids

followed by a stop codon (p.Phe15GlufsTer22) (Fig 3B). *cd9b^{pg15}* homozygous

mutants are viable and showed normal development. Loss of *cd9b* in situ signal in

the *cd9b* mutants suggested strong nonsense mediated decay (NMD) of the mutated

allele (S1 Fig). Due to the phenotypes seen in *cd9b* morphants, it was expected that

a lateral line phenotype would be seen in *cd9b* mutants with fewer neuromasts

deposited. However, no significant difference was found in the number of neuromasts

at 52 hpf (Fig 3). As *cd9b* is expressed throughout pLL development, migration of the

primordium as well as lateral line structure was assessed at 36 hpf. Although the pLL

shows no structural abnormalities in *cd9b* mutants at 52 hpf, it is possible that *cd9b*

mutants show a lateral line phenotype earlier in development and have recovered by

52 hpf. At 36 hpf, *cd9b* mutants show delayed primordium migration, with the

percentage of trunk traversed by the primordium reduced in mutants (Fig 3).

Fig 3. ***cd9b* mutant does not recapitulate morphant phenotype, although**

primordium migration is delayed at 36 hpf. A: Nature of the *cd9b* mutant allele

showing TALEN site location within the intron-exon structure of the gene. B: The

TALEN target sequence in exon 1 is shown in blue; the 8bp deletion in the *cd9b*

allele is indicated under the WT sequence as dashes. This leads to a frameshift

changing codon 15 from TTT (Phe) to CAA (Glu), then 22 aberrant amino acids (red

lettering) followed by a stop codon (*). C: Schematic of the Cd9b protein with location

of premature stop codon given by red arrow. The disulfide bonds between the

conserved CCG motif and conserved cysteines are indicated by the dashed lines.

EC1/2= Extracellular domain 1/2, aa= amino acid. D-E: Sequence chromatograms of

genomic DNA from (d) WT and (e) *cd9b^{pg15}* alleles. Deleted base pairs are

underlined in red. F-I: Representative images of WISH of c*laudin b* ISH on (f-g) WT

and (h-i) *cd9b* mutants at time shown. J-K: Graphs quantifying pLLP measurements

in WT and *cd9b* KOs; (j) the migration of the *cd9b* KO primordium at 36 hpf is

significantly delayed compared to WT. (k) There is no significance in number of

neuromasts deposited at 52 hpf. Unpaired T test on untransformed data. n=

minimum 20. ****=p=<0.05.

Generation of a *cd9a* mutant

To ascertain a role for Cd9a, we targeted the *cd9a* gene using CRISPR/Cas9.

A gRNA was designed to target *cd9a* in exon 3 (Fig 4A) and injected along with RNA

encoding Cas9 into WT embryos to generate *cd9a* KOs. After screening for

mosaicism, the CRISPR/Cas9 injected embryos were raised to maturity and

screened for germline transmission of the *cd9a* mutation. F0s were outcrossed with

WT fish to generate F1 *cd9a* single heterozygous mutants carrying an indel mutant

allele (c.183_186delinsAT; *cd9a^{ic62}*). This indel led to a frameshift mutation, changing

codon 62 from ATT (Iso) to TGC (Cys), then generating 54 aberrant amino acids

followed by a premature stop codon before the EC2 domain (p.Iso62Cysfs54Ter) (Fig

4B-C). Heterozygous *cd9a^{+/ic62}* adults were in-crossed to produce the F2 generation.

Adult F2s were viable and fertile and were genotyped to identify homozygous *cd9a^{ic62}*

KOs. Loss of *cd9a* in situ signal in the *cd9a* mutants suggested strong NMD of the

mutated allele (S2 Fig). The effect of *cd9a* KO on pLL development was investigated

but overall, there were no major defects. No significant difference was found in the

number of neuromasts deposited at 48 hpf or primordium migration at 36 hpf (Fig 4F-

K).

Fig 4. **Generation of *cd9a* mutant does not show any abnormal pLLP**

development. A: Nature of the *cd9a* mutant allele showing CRISPR site location

within the intron-exon structure of the gene. B: The CRISPR target sequence in exon

3 is shown in blue; the 2bp deletion in the *cd9a* allele is indicated under the WT

sequence as dashes. This leads to a frameshift changing codon 62 from ATT (Iso) to

TGC (Cys), then 54 aberrant amino acids (red lettering) followed by a stop codon. C:

Schematic of the Cd9a protein with location of premature stop codon, indicated by

red arrow, at a predicted 115 aa. EC1/2= Extracellular domain 1/2 D-E: Sequence

chromatograms of genomic DNA from (d) WT and (e) *cd9a^{ic62}* allele. Deleted base

pairs are underlined in red. F-G: *claudin b* expression in WT at (f) 30 hpf and (g) 48

hpf. H-I: *claudin b* expression in *cd9a* mutants at (h) 30 hpf and (i) 48 hpf. J-K:

Graphs quantifying pLLP measurements in WT and *cd9a* KOs; (j) migration of the

cd9a KO primordium at 36 hpf is similar to WT. (k) There is no significance in number

of neuromasts deposited. Significance was assessed using an unpaired t test. N=

minimum 7. Bars show mean +/- SD.

Generation of a *cd9a; cd9b* double mutants

Due to the redundant nature of tetraspanins, and *cd9a* having a similar mRNA

expression pattern to *cd9b,* it was speculated that functional redundancy between

Cd9a and Cd9b might be masking stronger phenotypes in the single mutants. Hence,

we generated *cd9a; cd9b* double mutants. To do this, the same *cd9a* gRNA as above

(Fig 4) and *cas9* mRNA were injected into *cd9b^pg15^* embryos. These fish were

screened for mutation of *cd9a* as above and then backcrossed to *cd9b^pg15^* mutants.

An indel mutation in exon 3 was detected (c.180_184delinsTCGCTATTGTAT) that

was predicted to lead to a frameshift mutation and an early stop codon, which

truncated the protein before the EC2 domain (p.Leu61Alafs9Ter) (Fig 5A-D).

Transheterozygous fish were crossed and adult F2s were genotyped to identify

doubly homozygous *cd9b^pg15^; cd9a^la61^* individuals, from now on referred to as *cd9*

dKOs. Adult *cd9* dKOs were viable and fertile and loss of *cd9a* and *cd9b* in situ signal

suggested strong NMD of the mutated alleles (S3 Fig). It was expected that with both

paralogues knocked out, the lateral line phenotype would now be as severe as *cd9b*

morphants with fewer neuromasts deposited. However, no significant difference was

found in the number of neuromasts at 48 hpf (Fig 5). Primordium migration was

significantly delayed in *cd9* dKO embryos at 36 hpf (Fig 5). This matches the

phenotype shown by the *cd9b^pg15^* single KOs. To see if the knockout of *cd9a^la61^* had

an additional effect on the primordium delay, embryos from *cd9b^pg15^* KOs were

compared to those from *cd9* dKOs (S4 Fig). The distance the primordium had

migrated at 36 hpf was measured and no difference was seen between *cd9b^pg15^*

single KO and *cd9b* dKO.

Fig 5. Generation of *cd9a* mutant in the *cd9b^pg15^* KO background to create a double *cd9* KO

mutant. A: *cd9a* mutant allele sequence from exon 3, showing CRISPR target

sequence in blue. The 4bp deletion (dashes) and 12bp insertion (lowercase) results

in an 8bp insertion as indicated under the WT sequence. This leads to a frameshift

changing codon 61 from CTC (Leu) to GCG (Ala), then 9 aberrant amino acids (red

lettering) followed by a stop codon. B: Schematic of the Cd9 protein with location of

premature stop codon for Cd9a (red arrow) and Cd9b (green arrow) at a predicted 70

aa and 36 aa respectively. EC1/2= Extracellular domain 1/2. C-D: Sequence

chromatograms of genomic DNA from (c) WT and (d) *cd9a^la61^* allele. Deleted base

pairs are underlined in red and inserted base pairs in blue. E-F: *claudin b* expression

in WT at (e) 30 hpf and (f) 48 hpf. G-H: *claudin b* expression in *cd9* dKO mutants at

(g) 30 hpf and (h) 48 hpf. I-J: Graphs showing pLLP measurements in WT and *cd9*

dKOs; (i) migration of the *cd9* dKO primordium at 36 hpf is significantly delayed

compared to WT. (j) There is no significance in number of neuromasts deposited.

Significance was assessed using an unpaired t test. N= minimum 9, p= <0.001, n1=

neuromast 1 etc. Bars show mean +/- SD.

In order to further investigate the migration and organisation of the

primordium in *cd9* dKO embryos they were crossed into the *cldnb:gfp* transgenic line.

Under the *claudin b* promoter GFP is expressed in the lateral line primordium and

newly deposited neuromasts [5]. This allowed observation of the primordium

migration in real time. Primordium migration appeared normal in the *cd9* dKO

embryos (Fig 6, Videos S1, S2). At the leading edge of the primordium filopodia

could be seen as well as proliferating cells. Within the migrating primordium

rearrangements occurred and rosettes were formed, increasing the length of the

primordium. As the trailing cells decelerated and deposited cells as neuromasts, the

primordium reduced in size. Quantification of various aspects of the primordium

during deposition revealed similar results in WT and mutants (S5 Fig).

Fig 6. **Primordium organisation appears normal in *cd9* dKO(*cldnb:gfp)* embryos**

from 30 hpf. A-B: Still images from a time-lapse recording of a migrating primordium

in (a) WT and (b) *cd9* dKOs. 0 minutes shows initial deposition as a proneuromast

becomes distinct from the primordium and then 2 sequential images show (a'-b') 20

minutes and (a''-b'') 60 minutes later. In the primordium of both WT and *cd9* dKOs,

filopodia can be seen at the leading edge (white arrow) and the formation of rosettes

in the trailing edge (white dashed circle). Representative images from two videos that

included two depositions each. Scale bar: 20 μm.

Cd9 interacts with the Cxcr4b/Cxcl12a pathway in the

migrating zebrafish lateral line primordium

To investigate if *cd9* had an effect on *cxcr4b* expression in the primordium,

cxcr4b WISH was performed but no perturbation of *cxcr4b* expression was seen in

the *cd9b* KOs (S6 Fig). Quantification showed no significant difference between the

expression of *cxcr4b* in WT and *cd9b* KOs. To investigate if Cd9 was affecting

migration in the zebrafish primordium through the Cxcr4b/Cxcl12a pathway, an

experiment using MOs was conducted. *Cxcr4b* and *cxcl12a* MOs induce premature

stalling of the primordium at concentrations of 1.5 mM and 0.5 mM respectively [21].

100 µM of *cxcr4b* MO was found to be the highest concentration injected into WT

embryos that did not induce a phenotype. However, this concentration of MO

resulted in a further delay of the primordium in the *cd9* dKOs (Fig 7A). For *cxcl12a*

the highest concentration of MO that did not induce a phenotype in WT embryos was

determined to be 12.5 µM but this had no further effect on the primordium migration

in *cd9* dKOs (S7 Fig). If the *cxcl12a* MO concentration was raised to 25 µM then the

primordium in both WT and *cd9* dKOs was delayed although the primordium in *cd9*

dKOs was significantly more delayed (Fig 7B).

Fig 7. *cxcr4b* and *cxcl12a* MOs further delay primordium migration at 36 hpf in

***cd9* dKO embryos.** A-B: Distance migrated by the primordium (labelled by *claudin b*

ISH) were recorded for (a) injection of 100 µM *cxcr4b* MO, and (b) 25 µM *cxcl12a*

MO. Significance was assessed using one-way ANOVA, *= $p < 0.05$, **= $p < 0.01$, ***=

p<0.001 ****= p<0.0001. N= minimum 15. Box extends from the 25th to 75th

percentile and whiskers from 10th to 90th.

Discussion

In this study we investigated the role of Cd9 in the migration of the pLLP in

zebrafish development. We first verified previous data showing *cd9b* expression in

the primordium at 30 hpf [2]. We also noticed *cd9b* was expressed throughout

recently deposited neuromasts at 48 hpf and 5 dpf. Our MO knockdown recapitulated

previous results from Gallardo et al., 2010. *cd9b* morphants showed fewer

neuromasts deposited, which suggested a role for Cd9b in the development of the

lateral line. A homozygous *cd9b* mutant was created using TALENs but surprisingly,

the morphant phenotype was not recapitulated in *cd9b* homozygous mutants. This

lack of phenotype could be due to several reasons. Firstly, the phenotype seen in

cd9b morphants could be due to off target effects of the MOs and not due to *cd9b*

knockdown. However, the fact that two different MOs and researchers induced the

same phenotype suggests it was not the lack of specificity [2]. Secondly, the Cd9b

truncated protein could have some residual function, although ISH results showed a

downregulation of mRNA suggesting NMD was occurring. Also, the *cd9b* mutation

aborts the normal sequence after aa15, before the EC2 domain, so any translated

protein would be expected to have minimal function. Finally, the *cd9b* morphant

phenotypes may be specific but tetraspanin redundancy could rescue the phenotype

in *cd9b* mutants. Tetraspanins are well known for their redundancy within the

tetraspanin family and mouse knockouts of single tetraspanins often appear healthy

and viable with mild phenotypes, whereas double tetraspanin knockout mice often

show increased numbers and severity of phenotypes [24–31]. In zebrafish, this

redundancy may be amplified due to the occurrence of a fish-specific whole genome

duplication in teleost fish. This means that many tetraspanins, for which there is one

mammalian ortholog, have two paralogs in zebrafish [32–36]. This is true for CD9, as

mammals have a single CD9 whereas zebrafish have Cd9a and Cd9b. We

demonstrated *cd9a* to also be expressed in the pLL in a similar pattern to *cd9b.*

Furthermore, NMD of mutated genes, as we have seen in both our mutants, has

been shown to invoke an upregulation of closely related genes as a genetic

compensation mechanism [37].

Following this, tetraspanin *cd9a* was targeted for knockout using CRISPR

technology. C*d9a^{ic62}* homozygous mutants and *cd9b^{pg15} cd9a^{la61}* double homozygous

mutants were generated. *Cd9a* KO embryos displayed a normal lateral line

phenotype with normal primordium migration and neuromast deposition. Initial

analysis of *cd9* dKO embryos using the same measurements revealed that the

number of neuromasts deposited and spacing were normal yet the distance of

primordium migration at 36 hpf was significantly delayed. It was expected that

knockout of *cd9a* would result in a stronger pLL phenotype but *cd9* dKOs showed a

similar phenotype to the *cd9b* KOs. Furthermore, when primordium migration of *cd9b*

single KO was compared to that of the *cd9* dKOs there was no difference. These

results suggest that the different phenotypes seen between the *cd9b* morphants and

mutants are not due to compensation by *cd9a*. It remains possible that there is

compensation by another tetraspanin [38]. Close relations to CD9 are the

tetraspanins CD81 and CD63, which have been shown previously to substitute for

CD9 in some circumstances. Overexpression of CD81 partially rescued the infertility

phenotype seen in CD9 KO mice [39] and it was recently demonstrated that CD9

deletion in human melanoma cells was quickly compensated by CD63 expression

upregulation [40]. It would be interesting to perform RNA-seq or SWATH-MS on the

cd9 dKOs to identify upregulated RNA and proteins respectively [41].

Videos generated of *cd9* dKOs with fluorescent *claudin b* showed primordium

migration and internal organisation were normal. Measurements of the primordium

after DAPI stain also revealed the primordium were similar sizes between mutants

and WT at 36 hpf. Overall, it seemed the delay in migration at 36 hpf in the *cd9* dKO

embryos was not due to alterations in the primordium organisation. Therefore, the

cd9 dKO phenotype appeared to be a migratory defect that could either be due to a

change in the onset of migration, a change in the speed of migration or a

combination of the two. CD9 has previously been reported to promote

CXCR4b/CXCL12a signalling in mammalian cells so we decided to investigate a

potential interference with this chemokine signalling pathway [16–18]. Results from

the WISH analysis and quantification showed no obvious perturbation of *cxcr4b*

expression in the *cd9* KOs. To postulate if there was some interaction between Cd9

and the Cxcr4b/Cxcl12a signalling pathway a MO experiment was performed. It was

theorised that a primordium lacking Cd9 would be more susceptible to disruption by

sub-functional doses of either *cxcr4b* or *cxcl12a* MO than WT. Injection with a sub-

functional concentration of 100 µM *cxcr4b* MO demonstrated that the *cd9* dKO larvae

were more sensitive to MO treatment compared to WT with significant retardation in

primordium migration in the dKO. This suggests that there is some interaction

between Cd9 and Cxcr4b within the primordium that promotes migration. Injection

with *cxcl12a* MO was more ambiguous as at a sub-functional concentration of 12.5

µM neither the WT nor the *cd9* dKO primordium migration was perturbed. When the

dose of *cxcl12a* MO was increased to 25 µM a migratory delay was induced in both

genotypes but the delay seen in the *cd9* dKO primordium was significantly worse.

Despite disruption in the presence of the MO, the primordium was still able to

eventually migrate the full length in the *cd9* dKO embryos, suggesting that Cd9 is not

essential for maintaining Cxcr4b/Cxcl12a signalling but may play a regulatory or

buffering role. A further experiment to check for interaction between Cd9 and

Cxcl12a/Cxcr4b in the primordium would be to use immunohistochemistry to

investigate co-localisation. Unfortunately, there are no tetraspanin, Cxcl12a or

Cxcr4b antibodies available in zebrafish. However, two plasmids are available that

encode *cxcr4b-egfp* or *cxcl12a-venus* constructs [42]. An interesting experiment

would be to inject embryos with these plasmids and visualise the distribution of

fluorescent versions of the proteins in the *cd9* dKOs.

We mentioned that the migratory defect could be due to a change in the onset

of migration. This could be further investigated by performing *claudin b* ISHs at the

onset and earliest stages of development of the posterior lateral line. On the other

hand, to try and determine if Cd9b has a role in the speed of migration, ISHs could

also be performed at various time points throughout primordium migration and speed

could be calculated in a similar method to previous studies [43,44].

We focused on a connection between Cd9 and Cxcr4b/Cxcl12 signalling in

this paper because of previous research linking these proteins. However, Cd9, like

other tetraspanins, contributes to cellular processes by organising molecules within

the plasma membrane and there are many other signalling pathways involved in

primordium development [45,46]. The fundamental four are FGF, Wnt, Notch and

chemokine signalling pathways and their specific and coordinated expression

ensures the regular morphogenesis and migration of the pLL [47]. Even though the

organisation of the primordium at high resolution appears normal, analysis of the

expression of these crucial signalling molecules, through WISH or

immunohistochemistry, during primordium development could reveal subtle changes

in their organisation that may be affecting migration. Whilst Fgf signalling functions to

determine rosette formation and deposition, Wnt signalling drives proliferation in the

leading zone [46]. Proliferation analysis with BrdU incorporation could show if there

are any changes in cell proliferation within the primordium, specifically in the leading

edge. CD9 has also been demonstrated to downregulate expression of Wnt

signalling pathways in cell culture [48,49]. Another important component of collective

cell migration is cell adhesion so migratory cells can pull insensitive cells in the same

direction. Within the primordium cadherins mediate cell-cell adhesion between

primordium cells and are required for directed and robust migration [50]. Cadherin-

dependent intercellular adhesion regulated by tetraspanins plays an important role in

tumour invasion and metastasis [51,52]. CD9 has been associated with keratinocyte

motility by regulating E-cadherin-mediated cell-cell contacts [53].

The use of computational models has been a valued tool to integrate analysis

of the lateral line primordium acquired through microscopic, cellular, molecular and

genetic analysis [47,54]. These models allow researchers to edit parameters and

observe their influences on the migrating primordium. This could help us choose the

most appropriate avenue to explore next by checking what modifications are able to

regenerate the phenotype we observe in the *cd9* dKO embryos.

In conclusion, we propose that within the primordium Cd9b is functioning to

compartmentalise signalling molecules, like Cxcr4b, to orchestrate and amplify

signalling upon ligand binding but is not essential. This would explain why only subtle

differences are seen between WT and *cd9* dKOs.

489 Acknowledgements

494 References

1. Olson HM, Nechiporuk AV. Using Zebrafish to Study Collective Cell Migration

in Development and Disease. Front cell Dev Biol. 2018 Aug 17;6:83.

2. Gallardo VE, Liang J, Behra M, Elkahloun A, Villablanca EJ, Russo V, et al.

Molecular dissection of the migrating posterior lateral line primordium during

early development in zebrafish. BMC Dev Biol. 2010;10(1):120.

3.   Dijkgraaf BYS. The functioning and significance of the lateral-line organs. Biol

Rev Camb Philos Soc. 1963 Feb;38:51–105.

4.   Metcalfe WK, Kimmel CB, Schabtach E. Anatomy of the posterior lateral line

system in young larvae of the zebrafish. J Comp Neurol. 1985 Mar

15;233(3):377–89.

5.   Haas P, Gilmour D. Chemokine Signaling Mediates Self-Organizing Tissue

Migration in the Zebrafish Lateral Line. Dev Cell. 2006 May;10(5):673–80.

6.   Dambly-Chaudière C, Cubedo N, Ghysen A. Control of cell migration in the

development of the posterior lateral line: antagonistic interactions between the

chemokine receptors CXCR4 and CXCR7/RDC1. BMC Dev Biol.

2007;7(1):23.

7.   Valentin G, Haas P, Gilmour D. The Chemokine SDF1a Coordinates Tissue

Migration through the Spatially Restricted Activation of Cxcr7 and Cxcr4b. Curr

Biol. 2007 Jun;17(12):1026–31.

8.   Xu H, Ye D, Behra M, Burgess S, Chen S, Lin F. GBeta1 controls collective

cell migration by regulating the protrusive activity of leader cells in the

posterior lateral line primordium. Dev Biol. 2014;385(2):316–27.

9.   Venkiteswaran G, Lewellis SW, Wang J, Reynolds E, Nicholson C, Knaut H.

Generation and Dynamics of an Endogenous, Self-Generated Signaling

Gradient across a Migrating Tissue. Cell. 2013 Oct;155(3):674–87.

10.   Donà E, Barry JD, Valentin G, Quirin C, Khmelinskii A, Kunze A, et al.

Directional tissue migration through a self-generated chemokine gradient.

Nature. 2013 Nov 25;503(7475):285–9.

11.   Stephens B, Handel TM. Chemokine Receptor Oligomerization and Allostery.

524   In: Progress in molecular biology and translational science. Prog Mol Biol

525   Transl Sci; 2013. p. 375–420.

526   12.   Thelen M, Legler DF. Membrane lipid environment: Potential modulation of

527   chemokine receptor function. Cytokine. 2018 Sep;109:72–5.

528   13.   Tarrant JM, Robb L, van Spriel AB, Wright MD. Tetraspanins: molecular

529   organisers of the leukocyte surface. Trends Immunol. 2003 Nov;24(11):610–7.

530   14.   Yáñez-Mó M, Barreiro O, Gordon-Alonso M, Sala-Valdés M, Sánchez-Madrid

531   F. Tetraspanin-enriched microdomains: a functional unit in cell plasma

532   membranes. Trends Cell Biol. 2009 Sep;19(9):434–46.

533   15.   Ovalle S, Gutiérrez-López MD, Olmo N, Turnay J, Lizarbe MA, Majano P, et

534   al. The tetraspanin CD9 inhibits the proliferation and tumorigenicity of human

535   colon carcinoma cells. Int J cancer. 2007 Nov 15;121(10):2140–52.

536   16.   Leung KT, Chan KYY, Ng PC, Lau TK, Chiu WM, Tsang KS, et al. The

537   tetraspanin CD9 regulates migration, adhesion, and homing of human cord

538   blood CD34+ hematopoietic stem and progenitor cells. Blood. 2011 Feb

539   10;117(6):1840–50.

540   17.   Brzoska E, Kowalski K, Markowska-Zagrajek A, Kowalewska M, Archacki R,

541   Plaskota I, et al. Sdf-1 (CXCL12) induces CD9 expression in stem cells

542   engaged in muscle regeneration. Stem Cell Res Ther. 2015 Mar 24;6(1):46.

543   18.   Arnaud MP, Vallée A, Robert G, Bonneau J, Leroy C, Varin-Blank N, et al.

544   CD9, a key actor in the dissemination of lymphoblastic leukemia, modulating

545   CXCR4-mediated migration via RAC1 signaling. Blood. 2015 Oct

546   8;126(15):1802–12.

19.   Brand MM, Granato M, Nüsslein-Volhard C, Dahm R. Zebrafish: A practical

approach. Keeping and raising zebrafish. 2002;7:39.

20. Trikić MZ, Monk P, Roehl H, Partridge LJ. Regulation of Zebrafish Hatching by

Tetraspanin cd63. Kanellopoulos J, editor. PLoS One. 2011 May

19;6(5):e19683.

21. David NB, Sapède D, Saint-Etienne L, Thisse C, Thisse B, Dambly-Chaudière

C, et al. Molecular basis of cell migration in the fish lateral line: role of the

chemokine receptor CXCR4 and of its ligand, SDF1. Proc Natl Acad Sci U S

A. 2002;99(25):16297–302.

22. Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount

zebrafish embryos. Nat Protoc. 2008 Jan 20;3(1):59–69.

23. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res. 2004 Mar 8;32(5):1792–7.

24. Knobeloch K-P, Wright MD, Ochsenbein AF, Liesenfeld O, Lohler J,

Zinkernagel RM, et al. Targeted Inactivation of the Tetraspanin CD37 Impairs

T-Cell-Dependent B-Cell Response under Suboptimal Costimulatory

Conditions. Mol Cell Biol. 2000;20(15):5363–9.

25. Takeda Y, Tachibana I, Miyado K, Kobayashi M, Miyazaki T, Funakoshi T, et

al. Tetraspanins CD9 and CD81 function to prevent the fusion of mononuclear

phagocytes. J Cell Biol. 2003 Jun 9;161(5):945–56.

26. Takeda Y, He P, Tachibana I, Zhou B, Miyado K, Kaneko H, et al. Double

Deficiency of Tetraspanins CD9 and CD81 Alters Cell Motility and Protease

Production of Macrophages and Causes Chronic Obstructive Pulmonary

Disease-like Phenotype in Mice. J Biol Chem. 2008 Sep;283(38):26089–97.

27. Wright MD, Geary SM, Fitter S, Gregory W, Lau L, Sheng K, et al.

Characterization of Mice Lacking the Tetraspanin Superfamily Member CD151

Characterization of Mice Lacking the Tetraspanin Superfamily Member CD151. Mol Cell Biol. 2004;24(13):5978–88.

28. Hemler ME. Tetraspanin Proteins Mediate Cellular Penetration, Invasion, and Fusion Events and Define a Novel Type of Membrane Microdomain. Annu Rev Cell Dev Biol. 2003;19(1):397–422.

29. Hemler ME. Targeting of tetraspanin proteins — potential benefits and strategies. Nat Rev Drug Discov. 2008 Sep;7(9):747–58.

30. Hassuna N, Monk PN, Moseley GW, Partridge LJ. Strategies for Targeting Tetraspanin Proteins. BioDrugs. 2009 Dec;23(6):341–59.

31. Gartlan KH, Belz GT, Tarrant JM, Minigo G, Katsara M, Sheng K-C, et al. A Complementary Role for the Tetraspanins CD37 and Tssc6 in Cellular Immunity. J Immunol. 2010 Sep 15;185(6):3158–66.

32. Meyer A, Van de Peer Y. From 2R to 3R: evidence for a fish-specific genome duplication (FSGD). BioEssays. 2005 Sep;27(9):937–45.

33. Brunet FG, Crollius HR, Paris M, Aury J-M, Gibert P, Jaillon O, et al. Gene Loss and Evolutionary Rates Following Whole-Genome Duplication in Teleost Fishes. Mol Biol Evol. 2006 Sep 1;23(9):1808–16.

34. Garcia-España A, Chung P-J, Sarkar IN, Stiner E, Sun T-T, DeSalle R. Appearance of new tetraspanin genes during vertebrate evolution. Genomics. 2008 Apr;91(4):326–34.

35. Huang S, Tian H, Chen Z, Yu T, Xu A. The evolution of vertebrate tetraspanins: gene loss, retention, and massive positive selection after whole genome duplications. BMC Evol Biol. 2010;10(1):306.

36. Howe DG, Bradford YM, Conlin T, Eagle AE, Fashena D, Frazer K, et al. ZFIN, the Zebrafish Model Organism Database: increased support for mutants

and transgenics. Nucleic Acids Res. 2012 Oct 15;41(D1):D854–60.

37.   El-Brolosy MA, Kontarakis Z, Rossi A, Kuenne C, Günther S, Fukuda N, et al.

Genetic compensation triggered by mutant mRNA degradation Europe PMC

Funders Group. Nature. 2019;568(7751):193–7.

38.   Huang S, Yuan S, Dong M, Su J, Yu C, Shen Y, et al. The phylogenetic

analysis of tetraspanins projects the evolution of cell–cell interactions from

unicellular to multicellular organisms. Genomics. 2005 Dec;86(6):674–84.

39.   Kaji K, Oda S, Miyazaki S, Kudo A. Infertility of CD9-Deficient Mouse Eggs Is

Reversed by Mouse CD9, Human CD9, or Mouse CD81; Polyadenylated

mRNA Injection Developed for Molecular Analysis of Sperm–Egg Fusion. Dev

Biol. 2002 Jul;247(2):327–34.

40.   Suárez H, Andreu Z, Mazzeo C, Toribio V, Pérez☐Rivera AE, López☐Martín

S, et al. CD9 inhibition reveals a functional connection of extracellular vesicle

secretion with mitophagy in melanoma cells. J Extracell Vesicles. 2021 May

12;10(7).

41.   Lin Q, Low LWL, Lau A, Chua EWL, Matsuoka Y, Lian Y, et al. Tracking

genome-editing and associated molecular perturbations by SWATH mass

spectrometry. Sci Rep. 2019 Dec 23;9(1):15240.

42.   Minina S, Reichman-Fried M, Raz E. Control of Receptor Internalization,

Signaling Level, and Precise Arrival at the Target in Guided Cell Migration.

Curr Biol. 2007 Jul;17(13):1164–72.

43.   Matsuda M, Nogare DD, Somers K, Martin K, Wang C, Chitnis AB. Lef1

regulates Dusp6 to influence neuromast formation and spacing in the

zebrafish posterior lateral line primordium. Development. 2013;140(11):2387–

97.

44.   Valdivia LE, Young RM, Hawkins TA, Stickney HL, Cavodeassi F, Schwarz Q,

624   et al. Lef1-dependent Wnt/ -catenin signalling drives the proliferative engine

625   that maintains tissue homeostasis during lateral line development.

626   Development. 2011;138(18):3931–41.

627   45.   Yauch RL, Hemler ME. Specific interactions among transmembrane 4

628   superfamily (TM4SF) proteins and phosphoinositide 4-kinase. Biochem J.

629   2000;351 Pt 3:629–37.

630   46.   Thomas ED, Cruz IA, Hailey DW, Raible DW. There and back again:

631   development and regeneration of the zebrafish lateral line system. Wiley

632   Interdiscip Rev Dev Biol. 2015 Jan;4(1):1–16.

633   47.   Dalle Nogare D, Chitnis AB. A framework for understanding morphogenesis

634   and migration of the zebrafish posterior Lateral Line primordium. Mech Dev.

635   2017;148:69–78.

636   48.   Lee JH, Bae JA, Lee JH, Seo Y-W, Kho DH, Sun EG, et al. Glycoprotein 90K,

637   downregulated in advanced colorectal cancer tissues, interacts with

638   CD9/CD82 and suppresses the Wnt/beta-catenin signal via ISGylation of beta-

639   catenin. Gut. 2010 Jul 1;59(7):907–17.

640   49.   Huang C, Liu D, Masuya D, Kameyama K, Nakashima T, Yokomise H, et al.

641   MRP-1/CD9 gene transduction downregulates Wnt signal pathways.

642   Oncogene. 2004 Sep 30;23(45):7475–83.

643   50.   Colak-Champollion T, Lan L, Jadhav AR, Yamaguchi N, Venkiteswaran G,

644   Patel H, et al. Cadherin-Mediated Cell Coupling Coordinates Chemokine

645   Sensing across Collectively Migrating Cells. Curr Biol. 2019 Aug;29(15):2570-

646   2579.e7.

51.   Johnson JL, Winterwood N, DeMali KA, Stipp CS. Tetraspanin CD151

regulates RhoA activation and the dynamic stability of carcinoma cell-cell

contacts. J Cell Sci. 2009 Jul 1;122(13):2263–73.

52.   Mohan A, Nalini V, Mallikarjuna K, Jyotirmay B, Krishnakumar S. Expression

of motility-related protein MRP1/CD9, N-cadherin, E-cadherin, α-catenin and

β-catenin in retinoblastoma. Exp Eye Res. 2007;84(4):781–9.

53.   Jiang X, Teng M, Ji R, Zhang D, Zhang Z, Lv Y, et al. CD9 regulates

keratinocyte differentiation and motility by recruiting E-cadherin to the plasma

membrane and activating the PI3K/Akt pathway. Biochim Biophys Acta - Mol

Cell Res. 2020;1867(2):118574.

54.   Dalle Nogare D, Chitnis AB. NetLogo agent-based models as tools for

understanding the self-organization of cell fate, morphogenesis and collective

migration of the zebrafish posterior Lateral Line primordium. Semin Cell Dev

Biol. 2020;100(December 2019):186–98.

Supporting Information

S1 Fig. ***cd9b* is significantly decreased in *cd9b* KO embryos.** A-B: Representative images of *cd9b* WISH at 36 hpf in (a) WT and (b) *cd9b* homozygous embryos.

S2 Fig. ***cd9a* is significantly decreased in *cd9a* KO embryos.** A-D: Representative images of *cd9a* WISH on (a-b) WT and (c-d) *cd9a* homozygous embryos at time shown.

S3 Fig. ***cd9a* and *cd9b* are both significantly decreased in *cd9* dKO embryos.** A-B: Representative images of *cd9a* ISH on (a) WT and (b) *cd9* dKO mutants at time shown. C-D: Representative images of *cd9b* ISH on (c) WT and (d) *cd9* dKO mutants at time shown.

S4 Fig. *cd9a* is not compensating for *cd9b* in the primordium. Graph showing the distance migrated by the primordium at 36 hpf is the same in *cd9b* KO and *cd9* dKO embryos. Significance was assessed using an unpaired t test, N= minimum 17. Bars show mean +/- SD.

S5 Fig. Primordium shape is normal in the *cd9* dKO embryos at 36 hpf. A: Representative image of a DAPI-stained primordium from a WT zebrafish at 36 hpf with red outline to show measured area. B-E: Graphs showing measurements of the primordium (b) height, (c) width, (d) width to height ratio, and (e) area in WT and *cd9* KO embryos. The height was measured three times at equal points along the primordium and then averaged. The width of primordium was measured from the two furthest points along the middle of the primordium. Width was then divided by height to generate a ratio. The area was circled using the freehand selection and measured. Cells were counted using the multipoint tool on Image J software. Significance was

assessed using an unpaired t test, p= <0.05, N= minimum 11. Bars show mean +/-

SD.

S6 Fig. *cxcr4b* expression is not altered in the pLLP of *cd9* dKO embryos at

indicated stages. A-D: Representative images of *cxcr4b* ISH in the primordium of

(a,c) WT and (b,d) *cd9* dKO mutants at time shown. Scale bar: 50 μm. E-F: Graphs

showing measurements of (e) length and (f) height of *cxcr4b* expression in the

primordium of WT and *cd9* KO embryos at indicated stages. Length measurements

were taken along the middle of the embryo between the two furthest points of

expression within the primordium. Height measurements were taken between the two

highest points of expression within the primordium. Significance was assessed using

an unpaired T test. N= minimum 13 for 24 hpf and 11 for 30 hpf. Bars show mean +/-

SD.

S7 Fig. 12.5 μM of *cxcl12a* MO does not further delay primordium migration at

36 hpf in the *cd9* dKO embryos. Distance migrated by the primordium (labelled by

claudin b ISH) was recorded for injection of 12.5 μM *cxcl12a* MO. Significance was

assessed using one-way ANOVA. N= minimum 13. Box extends from the 25th to

75th percentile and whiskers from 10[th] to 90[th].

S1 Video. WT primordium migration. *WT*(*cldnb:gfp)* from 30-38 hpf. Images taken

every 1 minute with a 20x objective. Movie length: 420 min.

S2 Video. *cd9* dKO primordium migration. *cd9* dKO(*cldnb:gfp)* from 30-36 hpf.

Images taken every 1 min with a 20x objective. Movie length: 360 min.

A Somite position of primordium at 36 hpf after injection with *cxcr4b* MO

Legend:
- Uninjected WT
- Injected with 100 μM *cxcr4b* MO
- Uninjected *cd9* dKO
- *cd9* dKO injected with 100 μM *cxcr4b* MO

B Somite position of primordium at 36 hpf after injection with cxcl*12a* MO

Legend:
- Uninjected WT
- WT injected with 25 μM *cxcl12a* MO
- Uninjected *cd9* dKO
- *cd9* dKO injected with 25 μM *cxcl12a* MO

A

cd9b MO1   *cd9b* i2e3

↳ *cd9b*

B Uninjected

C 0.5 μM Mismatch MO

D 0.1 μM MO1

E 0.1 μM i2e3 MO

F 0.2 μM MO1

G 0.2 μM i2e3 MO

H

Number of neuromasts

ns   **** **** **** **** **** ****

Uninjected
Mismatch MO 0.5μM
CD9b MO1 0.1μM
CD9b MO1 0.2μM
CD9b MO1 0.4μM
CD9b i2e3 0.1μM
CD9b i2e3 0.2μM
CD9b i2e3 0.4μM

I

Percentage of trunk length

ns   **** **** **** ns   ns   ns

Uninjected
Mismatch MO 0.5μM
CD9b MO1 0.1μM
CD9b MO1 0.2μM
CD9b MO1 0.4μM
CD9b i2e3 0.1μM
CD9b i2e3 0.2μM
CD9b i2e3 0.4μM

A *cd9b* TALEN

→ *cd9b*

B

TALEN site

wt GTACT TGCTCTTTATCTTC AA
cd9b^{pg15} GTACTTGCT________TCAA

wt MAAGGGLQCIKYLLLSSTSSS
GSPVQEFWLLVGCASMLKL...
cd9b^{pg15} MAAGGGLQCIKYLL QLHLLAR
RYRSFGCWSVAALRC*

C

EC2

EC1

Extracellular space
Plasma membrane
Cytoplasm

D *wildtype*

AGTACTTGCTCTTTATCTTCAACT

E *cd9b*^{pg15}

GTACTTGCTTCAACTT

F 36 hpf

cd9b^{+/+}

G 52 hpf

cd9b^{+/+}

H

cd9b^{-/-}

I

cd9b^{-/-}

J

Percentage

****

wildtype *cd9b -/-*

K

Number of neuromasts

ns

WT siblings *cd9b -/-*

A *cd9a* CRISPR

cd9a

B
CRISPR site
wt GAGTGTATATCCTCATTGCG
cd9a^{ic62} GAGTGTATATCCT_AT_GCG

wt PYVFYTGVYILIAAGALMM
VVGFFCC...
cd9a^{ic62} PYVFYTGVYILCGRCAHDG
GGIFRMLR...QGY*

C
EC2
EC1
36 54 111 195 Extracellular space
Plasma membrane
12 81 85 221 Cytoplasm

D *wildtype*
NFFTCCT(ATT)CGGCCGGT

E *cd9a*^{ic62}
TFTCCTATCGGCCGGTC

30 hpf	48 hpf

F *cd9a*$^{+/+}$ **G** *cd9a*$^{+/+}$

H *cd9a*$^{-/-}$ **I** *cd9a*$^{-/-}$

J Somite position of primordium at 36 HPF
ns
wildtype *cd9a -/-*
Somite number

K Neuromasts deposited at 48 HPF
ns
wildtype cd9a -/-
Number of neuromasts

A

CRISPR site
wt GAGTGTATATCCTCATTGCG
cd9a^{la61} GAGTGTATAT____tgcgtattgtatATT

wt PYVFYTGVYILIAAGALMM
cd9a^{la61} PYVFYTGVYIAYCILRPVRS*

B

EC2

EC1

Extracellular space

36   54   111   195
              Plasma membrane
12   81   85   221
              Cytoplasm
1

C *wildtype*

AGGAGTGTATATCCTCATTGCGGCCGG

D *cd9a^{la61}*

AGGAGTGTATATTGCGTANTGTATTGCGGCC

30 hpf

E *cd9a^{+/+} cd9b^{+/+}*

G *cd9a^{-/-} cd9b^{-/-}*

48 hpf

F *cd9a^{+/+} cd9b^{+/+}*

H *cd9a^{-/-} cd9b^{-/-}*

I Somite position of primordium at 36 HPF

Somite number

***

wildtype   *cd9 dKO*

J Number of neuromasts deposited at 48 HPF

Number of neuromasts

ns

wildtype   *cd9 dKO*