The cellular architecture of microvessels, pericytes and neuronal cell types in organizing regional brain energy homeostasis in mice

Yuan-ting Wu[1,*], Hannah C. Bennett[1,*], Uree Chon[1], Daniel J. Vanselow[2], Qingguang Zhang[3], Rodrigo Muñoz-Castañeda[4], Keith C. Cheng[3], Pavel Osten[4], Patrick J. Drew[3,5], Yongsoo Kim[1]

1 Department of Neural and Behavioral Sciences, The Pennsylvania State University, Hershey, PA, USA

2 Department of Pathology, The Pennsylvania State University, Hershey, PA, USA

3 Center for Neural Engineering, Department of Engineering Science and Mechanics, The Pennsylvania State University, University Park, PA, USA.
4 Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA
5 Biomedical Engineering, and Neurosurgery, The Pennsylvania State University, University Park, PA, USA

*Co-First author

Corresponding author:

Yongsoo Kim

Email: yuk17@psu.edu

Keywords: Cerebral vasculature, microvessel, pericyte, brain mapping, hemodynamics, parvalbumin, nNOS, vGlut1

Abstract

The cerebrovascular network and its mural cells must meet the dynamic energy demands of different neuronal cell types across the brain, but their spatial relationship among these cells is largely unknown. Here, we apply brain-wide mapping methods to create a comprehensive cellular-resolution resource comprising the distribution of and quantitative relationship between microvessels, pericytes, and glutamatergic and GABAergic neurons including neuronal nitric oxide synthase-positive (nNOS+) neurons and their subtypes in mice. Leveraging these data, we discovered region-specific signatures of vasculature and cell type compositions across cortical and subcortical areas, including strikingly contrasting correlations between the density of vasculature, pericytes, glutamatergic neurons and parvalbumin-positive interneurons versus nNOS+ neurons in the isocortex. We also found surprisingly low vasculature and pericyte density in the hippocampus, and distinctly high pericyte to vasculature ratio in the hypothalamus. These findings suggest that vascular density and mural cell composition is finely tuned to maintain regional energy homeostasis.

Introduction

The brain is the most energy-demanding organ per gram and is powered by an intricate web of vascular and mural cells that dynamically supply blood and clear metabolic waste (Hartmann et al., 2015; Sweeney et al., 2016; van Veluw et al., 2020; Vergara et al., 2019; Zhao et al., 2015). Pericytes, a key mural cell type, wrap around microvessels and are proposed to regulate blood flow and vascular permeability (Hall et al., 2014; Hartmann et al., 2021; Hill et al., 2015; Howarth et al., 2021; Nikolakopoulou et al., 2019; Rungta et al., 2018; Sweeney et al., 2016). Moreover, neuronal activity is known to regulate vascular diameter directly or indirectly (via astrocytes), which is referred to as a neurovascular coupling (Attwell et al., 2010; Kaplan et al., 2020). Neurons, unlike muscle, strictly rely on aerobic metabolism, thus neuronal functions critically depend on efficient vasculature support (Hall et al., 2012; Vergara et al., 2019). Not surprisingly, impairment of the cerebrovasculature, pericytes, and neurovascular coupling has been widely implicated in many neurological disorders such as stroke and neurodegenerative disorders (Lecrux and Hamel, 2011; Zhao et al., 2015). Yet, despite its significance, we have limited knowledge on the cellular architecture of vasculature and pericytes, especially with respect to their quantitative relationship with neuronal cell types across the whole brain. This relationship is likely of critical importance for the heterogeneous coupling of neural activity to blood flow described across different brain regions (Boido et al., 2019; Devonshire et al., 2012; Huo et al., 2014; Shih et al., 2009; Zhang et al., 2019a).

The generation of action potentials and synaptic transmission are energetically demanding (Howarth et al., 2012) and accordingly energy consumption is proposed to be linearly correlated with the number of neurons across different animal species including humans (Herculano-Houzel, 2011). However, neurons comprise highly distinct subtypes with different morphological, electrophysiological, and molecular characteristics (Kepecs and Fishell, 2014; Tasic et al., 2018). For example, the major classes of GABAergic neurons in the cortex include parvalbumin (PV), somatostatin (SST), and vasoactive intestinal peptide (VIP) expressing neurons, each of which make distinct synaptic connections with pyramidal neurons and each other (Kepecs and Fishell, 2014). Moreover, these neuronal cell types are expressed at different densities across cortical areas: PV interneurons have high density in sensory cortices and low density in association cortices, while SST neurons showed the opposite density pattern in mice (Kim et al., 2017). Different neuronal subtypes also have differential energy demands. For instance, the fast spiking PV neurons are among the highest energy demanding neuronal types (Inan et al., 2016). On the other hand, another neuronal type – the neuronal nitric oxide synthase (nNOS) expressing neurons – can actively regulate blood supply by causing vasodilation (Echagarruga et al., 2020; Krawchuk et al., 2020; Lee et al., 2020). Taken together, these data suggest that determining specific spatial relationships between neuronal cell types and the vascular network is critically important for understanding the demand for and the mechanism of distinct blood flow regulation across different brain regions.

It has been technically challenging to comprehensively examine mammalian cerebrovasculature because of its fine and complex branching across the entire brain, with the diameter of small capillaries being only ~5μm. However, recent progress in meso-scale mapping methods began elucidating the structural organization of the cerebrovasculature and its regional heterogeneity across the mouse brain at unprecedented detail (Ji et al., 2021; Kirst et al., 2020; Todorov et al., 2020; Xiong et al., 2017). However, while these studies provide a detailed description of the

general organization of mouse brain cerebrovasculature, the question of how cerebrovasculature and its mural cells are structurally organized to support different neuronal cell types across the whole brain remains largely unanswered. To address the relationship between cerebrovasculature and mural and neuronal cell types, we have devised a serial two-photon tomography (STPT)-based imaging approach with newly developed analytical tools to derive the first cellular architecture atlas containing cerebrovasculature, capillary pericytes, and several major neuronal cell types, including PV interneurons and vasomotor nNOS neurons in the adult mouse brain.

Leveraging this data resource, we asked the following questions about the quantitative relationship of vasculature and associated cell types in different brain regions, with emphasis on the isocortex due to the well-established functional roles of cortical neuronal subtypes. (1) Are densities of microvessels and pericytes uniform across the entire cortex or are there distinct spatial patterns that may be linked to the densities of neuronal cell types, such as the energy demanding PV neurons and vasomotor nNOS neurons? And (2) does blood perfusion via microvessels occur evenly in 3D across the cortex, or are there directional preferences based on cortical layers or different cortical areas? Answering these two questions will help to elucidate local cerebrovascular organization principles and shed light on how distinct cortical domains maintain balance between energy demands and supplies. Lastly, we also asked whether (3) there are areas with distinct vascularization and pericyte coverage that may predispose certain brain areas to pathological conditions. For instance, impairment of the hippocampus and association cortical areas are frequently associated with neurodegenerative disorders such as Alzheimer's disease (Ballinger et al., 2016; Sweeney et al., 2018). Comparing vascular networks in these regions to other areas may help to explain brain regional vulnerabilities.

132   **Results**
133
Comprehensive vascular mapping in the intact mouse brain
Our first goal is to examine the spatial arrangement of the cerebrovasculature in an intact whole
mouse brain to understand the anatomical basis of the vascular network. We filled microvessels
from 2-month-old C57BL/6 mice with a cardiac perfusion of FITC-conjugated albumin gel
(Figure 1A)(Blinder et al., 2013; Tsai et al., 2009) and used STPT imaging in combination with
two-photon optical scans and serial sectioning to image the whole mouse brain at 1x1x5μm
(*x,y,z;* media-lateral, dorsal-ventral, rostral-caudal) (Figure 1B). Since the vascular tracing
requires precise stitching across X-Y tiles throughout z stacks, we developed a new stitching
algorithm to correct optical aberrations, bleaching in overlapped tile areas, and z stack
alignments (Figure 1C-D; Figure S1). To detect and analyze cerebrovascular arrangement, we
developed a computational pipeline that binarizes and traces the original image of the FITC-
filled vasculature, as well as quantifies the diameter of each vessel and its branching points
across the whole brain (Figure 1E-G; See Methods for more details). Individual brains were then
registered to the Allen Common Coordinate Framework (CCF) to quantify signals across
different anatomical areas (Wang et al., 2020) (Figure 1H-N; Table S1; MovieS1). Although we
observed near-complete vasculature labeling with our FITC-based filling approach, we
implemented an additional quality control step to reject data from areas with potentially
incomplete labeling or imaging artifacts (Figure S1). Moreover, we confirmed that our approach
closely reflects vasculature *in vivo* by directly comparing STPT results with *in vivo* two-photon
measurements of the same vasculature in same mice (Figure S2).
154   Initial analysis of this data revealed that for vascular density, there is up to a three-fold
difference between highly (e.g., cerebellar nuclei) and sparsely vascularized regions (e.g., medial
amygdala)(Figure 1K-N). For example, the cerebellum, midbrain, and thalamic areas have high
vascular density while the cortex, hippocampus, amygdala, and hypothalamic regions show
overall low vascular density (Figure 1K-L). These results are consistent with recent studies to
quantify vascular densities in the mouse brain, further confirming that there are vascular density
differences across the brain based on regionally heterogenous metabolic needs (Ji et al., 2021;
Kirst et al., 2020).
162   Since vascular branching represents the degree of connectivity in the vasculature, we ask
how vascular branching density is quantitatively related to vascular length density across brain
regions (Figure 1M). If the vascular density increases while maintaining their connectivity, we
can expect that branching density will proportionally increase to the power of 1.5. Our result
shows that the branching density increases to the power of 1.183, indicating that vascular
branches have lower connectivity as vascular density increases (Figure 1M). In contrast, vascular
radii do not show significant correlation with vascular density (Figure 1N).
169
Vasculature across the isocortex is not uniform
The mouse isocortex, particularly the primary somatosensory cortex, has been a major area of
focus for *in vivo* studies examining the cellular mechanisms of neurovascular coupling
(Echagarruga et al., 2020; Hill et al., 2015; Krawchuk et al., 2020; Vazquez et al., 2018). Yet, it
is not known whether the vasculature is anatomically organized in a similar way across the entire
cortex or whether different cortical areas may have unique vascular organization signatures and
consequently different neurovascular coupling (Huo et al., 2014; Zhang et al., 2019a).

177   We found highly regionally heterogeneous vascular distribution in the isocortex, with
178 unique patterns for distinct functional cortical areas (Figure 2). Sensory motor areas such as
179 somatosensory (SS) and auditory cortices (AUD) showed overall high vascular density,
180 branching density, and radii while medial prefrontal areas including prelimbic (PL), medial
181 orbital (ORBm), and infralimbic (ILA) were low on all three measurements (Figure 2A-B). In
182 contrast, many lateral association areas including the posterior agranular insular (AIp), ECT, and
183 PERI showed high vascular diameter with relatively lower vascular density (Figure 2A-B).
184 Noticeably, the retrosplenial cortex (RSP), linked with spatial navigation and memory processing
185 (Mitchell et al., 2018), has low vascular diameter despite high vascular density (Figure 2A-B).
186 To examine the spatial distribution intuitively while maintaining high resolution information, we
187 devised an isocortical flatmap based on Laplace's equation (Figure 2C-F; see Methods for more
188 details). When averaged vessel length density is plotted in the cortical flatmap, it is clear that
189 densely vascularized areas are tightly aligned with anatomical borders of the SS, AUD, visual
190 (VIS), and RSP cortices (Figure 2G, arrowheads). Cortical layer-specific maximum projection of
191 the length density shows that sharp boundaries between cortical areas are strongly driven by the
192 layer 2/3/4 vascular distribution (Figure 2H). Interestingly, one measure of the cortical
193 vasculature that did not show strong regional differences is the vessel radius, which is relatively
194 similar across cortical areas with the exception of the RSP that comprises vessels with low
195 vascular diameter (Figure 2G).
196   Taken together, these data provide strong evidence that cortical vascularization is not
197 uniform but is distinctly organized in functionally different cortical areas.
198
199

200 **Quantification of permeability reveals spatial heterogeneity of the vasculature network**
201 Since microvessels provide a large surface area to exchange oxygen and glucose, we examined
202 the link between microvessel structure and its influence on blood perfusion in the brain by
203 applying a mathematical approach to infer permeability to blood flow and directionality of the
204 microvascular network based on our vascular measurements (Figure 3A; Movie S2; Table S2).
205 To limit our simulation to small vessels, we removed large vessels with radius > 4 μm. The
206 permeability was calculated through solving the Hagen–Poiseuille equation system of the entire
207 network in the control volume (Figure 3A), taking into account vascular density, connectivity,
208 and network topology (e.g., tortuosity). For example, a network twice as dense with identical
209 topology will have doubled permeability (Figure 3B; length density), while doubling the vascular
210 radius of an identical network gives a sixteen-fold increase in permeability (Figure 3B; vessel
211 radius). With respect to vascular topology, a bias in the orientation of the vasculature will lead to
212 a higher permeability tensor in parallel to the vessel, but a lower permeability in orthogonal
213 directions (Figure 3B; directionality), and a highly twisted network can significantly reduce the
214 permeability (Figure 3B; tortuosity). After probing the permeability of a given microvessel in all
215 directions, we spherically integrated the permeability tensor, yielding a scalar as final
216 permeability output (Figure 3B) representing how well blood can pass through microvessels at a
217 given input from the arteries.
218   We present two examples from the isocortex, one is the primary somatosensory cortex
219 barrel field (SSp-bfd) with relatively higher permeability, and the other one is the ORBm in the
220 medial prefrontal cortex with poor permeability (Figure 3C). Higher permeability of the SSp-bfd
221 occurs due to the high density of vasculature that is arranged in all directions despite higher
222 vascular tortuosity (Figure 3C, left side). In contrast, the lower permeability of the ORBm is

largely driven by low vascular density despite its relatively lower tortuosity (Figure 3C, right side). Across isocortical areas, sensory-motor and RSP areas have overall higher permeability compared to medial prefrontal and lateral association areas (Figure 3D; Table S2).

Since the geometry of microvascular networks can influence directionality of blood flow, we examined the microvessel anisotropy using the permeability tensor in the isocortex (Figure 3E-I). We used three axes according to the cortical column direction: penetrating (P) axis along the cortical column, anterior-posterior (AP), and medial-lateral (ML) (Figure 3G). Our analysis showed that microvessels oriented in the P-axis dominated in the anterior (e.g., motor areas) and posterior cortical areas (e.g., visual area) while mid-cortical areas (e.g., somatosensory area) showed vasculature orientation in the P and AP axes dominating the superficial layer and the deep layers, respectively (Figure 3E-I). For instance, the secondary motor cortex (MOs) shows dominant P-axis vasculature (magenta) while the SSp-bfd shows clear switch to AP axis (cyan) vasculature preferentially between layers 4 – 6 (Figure 3E-F). Collectively, our data demonstrate that the isocortex contains different cortical domains with area and cortical layer specific microvascular perfusion patterns.

When we applied the permeability measurement across the whole brain, we found that the thalamic area (e.g., VENT), cortical subplate areas (e.g., BLA), selected midbrain areas including the inferior colliculus (IC) show higher permeability (Figure 3J; Figure S3; Table S2; Movie S2). In contrast, many hippocampus areas including the dentate gyrus (DG) and the subiculum (SUB) showed low permeability (Figure 3J; Figure S3; Table S2). This baseline difference of microvascular permeability can help to explain regional vulnerabilities of hippocampal areas in association with many neurodegenerative disorders (Ballinger et al., 2016; Sweeney et al., 2018).

Pericyte density mapping reveals differential pericyte coverage between cortical and subcortical areas

Pericytes encapsulate the microvasculature and are thought to actively regulate microvascular diameter and permeability (Attwell et al., 2016; Bennett and Kim, 2021; Hartmann et al., 2021; Nelson et al., 2020; Nikolakopoulou et al., 2019). Here, we ask whether pericyte density in different brain areas parallels that of cortical vasculature and whether the ratios of pericyte to vascular density may differ by brain area. To map pericyte distribution, we crossed PDGFRβ-Cre mice with Ai14 reporter mice (PDGFRβ-Cre:Ai14 mice) to fluorescently label vascular mural cells (Cuttler et al., 2011; Hartmann et al., 2015) and imaged their whole brain distribution by STPT (Figure 4A-B). To conduct whole brain quantification of this cell population, we developed a new machine learning algorithm to specifically count capillary pericytes from fluorescently-labeled cells in PDGFRβ-Cre:Ai14 mice (Figure 4C;Figure S4; Table S3; Movie S3; see the methods section for more details). Furthermore, we used our previously established qBrain mapping pipeline to map the signals onto the adult CCF (Kim et al., 2017; Wang et al., 2020) and quantify the 3D density of the pericyte distribution across the whole brain (Figure 4C-H; Table S4; see the methods section for more details).

When we examined the regions of the isocortex, we found that cortical pericyte density showed a similar distribution pattern to the vascular density (Figure 4C-D; Tables S4). While the sensory motor areas, including the somatosensory and auditory regions as well as the RSP, contain a higher density of pericytes, the medial prefrontal and lateral association cortices show lower pericyte density (Figure 4C-D; Table S4). Moreover, when comparing cortical layers, layers 4 and 5 contain the highest density of pericytes across the whole isocortex, while layer 1

contains the lowest pericyte density (Figure 4D; Table S4). This is particularly apparent in
regions of high pericyte density, such as the somatosensory and auditory cortices. Overall,
cortical pericyte density shows strong positive correlation with vascular length density (Figure
4E).
For the rest of the brain, we found pericyte density varied by a factor of two across
different brain areas and showed significantly positive correlation with vascular density across
the brain (Figure 4F-H; Table S4). Noticeably, cortical sensory circuit pathways including
sensory thalamic nuclei (e.g., ventral group of dorsal thalamus; VENT, Dorsal lateral geniculate
nucleus; LGd) are among the high-density groups while memory circuit pathways including the
DG of the hippocampus and mammillary body (MBO) are among the low-density group (Figure
4F-H; Table S4). Furthermore, many hypothalamic areas (e.g., anterior hypothalamic nucleus;
AHN) have relatively high pericyte density despite low vascular density (Figure 4G).
In summary, our pericyte mapping results reveal positive correlations with microvascular density
but with extensive regional heterogeneity across the whole brain.
283
Cell type mapping with genetic intersection approach reveals regionally distinct nNOS
subtypes expression
Next, we examined the relationship between the cerebrovascular network with pericytes and the
distribution of nNOS-expressing neurons that are known to control neurovascular coupling in the
brain (Echagarruga et al., 2020; Krawchuk et al., 2020; Lee et al., 2020). We employed a genetic
labeling method using nNOS-CreER mice with postnatal tamoxifen induction (Figure
5A)(Taniguchi et al., 2011). Since nNOS neurons also contain different subtypes co-expressing
SST, NPY, PV, or VIP, which are linked with distinct functions including regulating vascular
diameter (Perrenoud et al., 2012; Tricoire and Vitalis, 2012; Williams et al., 2017), we utilized
genetic intersection methods using Cre and Flp dependent reporter mice (Ai65) crossed with
nNOS-CreER and one of the subtype markers (NPY-, SST-, PV-, or VIP-Flp), in combination
with tamoxifen induction (Figure 5A)(He et al., 2016). Then, we used the qBrain mapping
method with new nNOS detection algorithms to examine reporter gene expression from these
different mouse lines (Figure 5B-E; Figure 3S; Movie S4;Table S5).
Our results indicate that the total nNOS neuronal density is highest in the accessory
olfactory bulb, the striatum-like amygdala (e.g., the medial amygdala), the hypothalamus, and
the cerebellum (Figure 5F-H). In contrast, the isocortex, the hippocampus, and the thalamus
showed relatively lower overall nNOS neuronal density (Figure 5F-H). The nNOS/NPY neurons
represent the majority of nNOS subtypes in cerebral cortical and cerebral nuclei areas, including
hippocampal areas (Figure 5F-H). The nNOS/SST subtype showed similar density compared to
the nNOS/NPY subtype overall, but with lower density in hippocampal regions (Figure 5F-H).
The nNOS/PV subtype showed overall low density across the whole brain, except for very high
density in the cerebellum (Figure 5F-H). Lastly, the nNOS/VIP subtype had the lowest density
compared to the other nNOS[+] subtypes, except in a few areas, such as the subiculum of the
hippocampus, where the density was modest (Figure 5F-H). Noticeably, many amygdala and
hypothalamic areas as well as the accessory olfactory bulb showed high nNOS density that was
not reflected in the nNOS interneuron subtype populations, suggesting that nNOS neurons in
these areas may represent different nNOS subtypes (Chachlaki et al., 2017).
312
Cortical nNOS neurons are negatively correlated with vascular and pericyte densities

Since the vasomotor function of cortical nNOS neurons is well-established (Echagarruga et al.,
2020; Lourenço et al., 2017; Perrenoud et al., 2012), we examined whether the density
distribution of nNOS neurons in the isocortex is significantly correlated with the density of
cerebrovasculature and pericytes. Overall, we observed up to two-fold differences in nNOS
neuronal density across isocortical brain regions (Figure 5I-K; Table S5). Surprisingly, nNOS
neurons showed higher expression in the medial and lateral association cortices (e.g., the
agranular insular cortex), while sensory-motor areas as well as the RSP showed lower density,
which is the opposite of the vascular density pattern (Figure 5I,K). The highest density of nNOS
neurons is found in layer 6 in all cortical areas (Figure 5I). For nNOS subtypes, nNOS/NPY and
nNOS/SST subtypes showed similar density patterns with the total nNOS neurons (Figure 5J-K).
In contrast, nNOS/PV and nNOS/VIP subtypes, despite much lower density, showed relatively
higher expression in the RSP and the lateral cortex, respectively (Figure 5J-K). When we
performed correlation analysis for the relationship between nNOS density and vascular density
across different areas, we found a significant negative correlation in the isocortex (Figure 5L).
All subtypes except nNOS/PV showed a similar negative correlation with vascular density
(Figure 5M). Similarly, nNOS neurons, including nNOS/NPY and nNOS/VIP subtypes, showed
significant negative correlation with pericyte density (Figure 5M). In contrast, nNOS neurons
and all of their subtypes did not show any correlation with average vasculature radius (Figure
5M). This data suggests that cortical nNOS neurons have overall stronger vasomotor regulation
in high association cortices than in sensory cortices.
334
Cortical PV interneurons and glutamatergic neurons show positive correlation with the
vascular network
Glutamatergic and GABAergic neuronal cell types have different energy consumption and
metabolic costs (Buzsáki et al., 2007). Here, we examined whether specific neuronal subtypes
show any significant correlation with vascular and pericyte distribution in the isocortex (Figure
6; Table 1; Table S6). We used pan-glutamatergic (vGlut1) and pan-GABAergic (Gad2) Cre
driver lines crossed with conditional nuclear tdTomato reporter mice (Ai75) and used the qBrain
mapping method to quantify each cell type signal and map the distribution onto our isocortical
flatmap (Figure 6; Movie S5). For non-overlapping GABAergic interneuron subtypes PV, SST,
and VIP interneurons, we used our previously mapped data based on the cell type specific Cre
drivers crossed with conditional nuclear GFP reporter mice (Kim et al., 2017)(Figure 6A-C).
Density plotting using our isocortical flatmap demonstrates the different neuronal cell type
distributions and localizations across the isocortex, and shows a clear pattern when compared to
the vessel length and pericyte densities (Figure 6C). Among GABAergic cell types, PV
interneurons show a strikingly significant positive correlation with the vascular length density
(Figure 6C-E). All sensory cortical areas, including the primary somatosensory cortex, showed
high PV interneuron and vascular density (Figure 6B-E). In contrast, the other interneuron
subtypes (SST and VIP) and pan-GABA (Gad2) did not show a significant correlation with
vascular density (Figure 6C-E). Lastly, pan-glutamatergic neurons (vGlut1) showed significant
positive correlation with vascular density (Figure 6C,E). As expected, the pericyte distribution
mirrored the same correlation patterns as the vessel length density with all of the neuronal
subtypes studied (Figure 6E). Importantly, cortical PV neurons are involved in the generation of
gamma-band oscillations (Cardin et al., 2009; Sohal et al., 2009), which are linked with
increased vasodilation and blood flow in the brain (Drew et al., 2020). Thus, our results suggest

that neurovascular and pericyte density are proportionally distributed to the PV neurons in order
to support local neural activity in sensory cortices.
361
362

Web visualization
Ease of access and intuitive visualization is a key when examining large scale imaging datasets.
Toward these goals, we created a web-based resource (https://kimlab.io/brain-map/nvu/) that
displays navigable z-stacks of full-resolution images for our STPT datasets including FITC filled
vasculature, PDGFRβ-Cre:Ai14 for pericytes, nNOS-Cre:Ai14 for total nNOS neurons, and
nNOS-Cre:Ai65:(NPY, SST, PV, or VIP-Flp) for nNOS subtypes. This web-based resource also
provides interactive 3D visualizations, allowing users to navigate our quantitative vascular and
cell type measurements registered in the Allen CCF.
371

Discussion

The structural organization of regional vascular networks is crucial to understand their function as well as susceptibility to pathology. Here we present cellular resolution maps of cerebral vasculature, pericytes, and neuronal subtypes in the mouse brain. Our cerebrovascular map, in combination with fluid dynamic simulations, reveals differential coverages of microvessels and pericytes as well as their relationships with neuronal cell types, highlighting heterogenous blood perfusion and potential differences in regulation across different brain regions as exemplified in Figure 7. Most notably, we observed strikingly strong correlations between vascular density and parvalbumin interneuron and vasomotor nNOS neuron density in the isocortex, suggesting that regional differences in neuronal cell type composition linked with different energy demands are met by region-specific patterns of cortical vascularization. In combination with web visualization, our maps serve as a comprehensive resource to examine microvasculature and associated cell types.

Cortical neuronal cell types and the vascular network

A prevailing theory of cortical organization is that the cortex is composed of repeating cortical columns with a common microcircuit motif (Douglas and Martin, 2004). For example, excitatory and inhibitory neuronal subtypes make stereotypic local connection patterns, called canonical cortical circuits, that are similarly observed across cortical areas (Packer and Yuste, 2011; Pi et al., 2013). However, this view has been challenged by recent data that different cortical domains show distinct cell type compositions and hemodynamic responses (Kim et al., 2017; Zhang et al., 2019a). Results from the current study, as summarized in Figure 7 and Table 1, provide further evidence that vascular networks, including pericytes and vasomotor neurons, are organized differently to meet energy demand from sensory and association cortices (Howarth et al., 2012; Vergara et al., 2019).

Sensory signals require precise temporal and spatial information processing in sensory cortices such as the somatosensory cortex barrel field representing whisker-evoked stimuli in rodents. In contrast, association cortices integrate information from broader areas with slower temporal kinetics. We previously identified a higher density of PV neurons in sensory cortices compared to association cortices (Kim et al., 2017). Cortical PV neurons are fast spiking interneurons that participate in generating gamma oscillations and are one of the most energy demanding neurons (Cardin et al., 2009; Hu and Jonas, 2014; Inan et al., 2016; Kann, 2016). Thus, our current results suggest that a high density of microvessels and capillary pericytes in the sensory cortices provide an efficient energy support system for PV dominated local circuits to accommodate high energy consumption and handle high speed sensory processing. In contrast, association cortices contain relatively high densities of nNOS neurons despite low vascular, pericyte, and PV densities. Although nNOS interneurons represent only about 2% of cortical neurons, activation of nNOS neurons robustly dilates cerebral arterioles to generate increases in cerebral blood flow (Echagarruga et al., 2020; Hosford and Gourine, 2019; Krawchuk et al., 2020; Lee et al., 2020). Moreover, type I nNOS neurons co-expressing NPY and SST in the deep cortical layer are a rare cortical GABAergic type with long-range projection to other cortical areas, which can help to synchronize activities of target areas (Melzer and Monyer, 2020; Tomioka and Rockland, 2007; Tomioka et al., 2005). Thus, the relatively higher density of nNOS/SST neurons in association areas suggests that this cell type can exert more powerful vasodilation in larger areas to compensate for a lower vascular density in these high cognitive areas, while also orchestrating vasodilation in other cortical areas.

Microvascular map to understand regional blood flow

419
We used serial two-photon tomography to visualize the whole cerebrovasculature at single
capillary resolution from intact mouse brains that closely represents physiological conditions,
confirmed by *in vivo* two-photon microscopy. Although recent approaches using light sheet
microscopy to examine fine cerebrovascular structure provide advantages in rapid data
acquisition as well as 3D immunolabeling to mark different vascular compartments (Kirst et al.,
2020; Todorov et al., 2020), the required tissue clearing methods can introduce microscopic
volume distortions, which can lead to inconsistent measurements (Ji et al., 2021). Indeed, our
vascular measurements (e.g., length density) are consistent with other recent data that utilized
serial two-photon microcopy (Ji et al., 2021).
Our detailed geometric analysis of microvessels enables us to quantify vascular
permeability across different brain areas as a first step to building a more comprehensive model
of cerebral perfusion. Our permeability measurements represent potential blood flow efficiency
based on structural arrangements of microvessels upon a given input. The frontal cortical and
hippocampal areas are highly vulnerable in normal aging and neurodegenerative disorders such
as Alzheimer's disease (Sengillo et al., 2013; Zhang et al., 2019b). Our results suggest that these
areas have overall low microvessel permeability with low vascular densities, even at the baseline
condition, providing further evidence of the regional vulnerability to microvascular insults. Our
results also highlight area and layer specific vascular anisotropy in the isocortex, which is
consistent with recent studies (Ji et al., 2021; Kirst et al., 2020). However, the functional
significance of this vascular anisotropy remains unclear. Future works including computational
modeling considering additional information (e.g., blood pressure and viscosity) can help to gain
a more complete understanding of brain blood perfusion (Balogh and Bagchi, 2019; Blinder et
al., 2010).

Brain-wide pericyte map

444
Pericytes actively regulate the diameter and permeability of microvessels (Hartmann et al., 2021;
Nikolakopoulou et al., 2019). For example, optogenetic stimulation of capillary pericytes in the
cortex induces vasoconstriction, reducing red blood cell flux up to 50% (Hartmann et al., 2021;
Nelson et al., 2020). Our results complement this work by providing pericyte density across the
brain. We observed a strong positive relationship between pericyte and vascular density in the
cortex, suggesting that pericyte coverage per microvessel remains similar across different
cortical areas in the normal adult mouse brain. Moreover, subcortical mapping results indicate
that thalamic areas have overall higher pericyte density. Interestingly, thalamic pericytes showed
resistance to disrupted PDGFRβ signaling, while cortical and striatal pericytes were vulnerable
(Nikolakopoulou et al., 2017). The combination of high density and cellular resilience may
confer extra protection to maintain vascular integrity in the thalamus. Conversely, relatively low
pericyte density in the hippocampal areas and association cortices can make these areas more
vulnerable to pathological conditions (Montagne et al., 2015; Sengillo et al., 2013; Zhao et al.,
2015).

In summary, our quantitative information on cerebrovasculature and associated cell types can
help to gain a cellular architectural basis of how energy demand and supply maintain balance in a
normal brain and how this homeostatic mechanism changes under pathological conditions in the
future.

Material and Methods

Animals

Animal experiments were approved by the Institutional Animal Care and Use Committee at Penn State University and Cold Spring Harbor Laboratory. For all genotypes in this study, both adult male and female mice were used. Adult 2-month-old C57BL/6 mice were bred from C57BL/6 mice directly obtained from the Jackson Laboratory and used for vascular tracing experiments with FITC filling (N=4). For pericyte specific experiments, male PDGFRβ-Cre mice (Cuttler et al., 2011) were crossed with female Ai14 mice (Jax: Stock No: 007914) as previously described (Hartmann et al., 2015). These PDGFRβ-Cre:Ai14 mice exhibit PDGFRβ-driven tdTomato expression in two distinct vascular cell types, pericytes and vascular smooth muscle cells (vSMCs). For isocortical cell types, vGlut1-Cre (Jax: 023527) and Gad2-Cre (Jax: 010802) mice were crossed with Ai75 reporter mice (Jax: 025106). nNOS-CreER mice were used to label nNOS neurons (Jax: Stock No: 014541)(Taniguchi et al., 2011). After nNOS-CreER mice were crossed with Ai14 mice, the nNOS-CreER:Ai14 offspring were administered with an intraperitoneal (i.p.) tamoxifen (Sigma, cat.no. T5648-1G) injection (100mg/kg) at P16. Similarly, for nNOS-subtypes, nNOS-CreER mice were initially crossed with Ai65 mice (Jax; Stock No: 021875), which were further crossed with PV-flp (Jax Stock No: 022730), SST-flp (Jax Stock No: 028579), NPY-flp (Jax Stock No: 030211), or VIP-flp (Jax Stock No: 028578) mouse lines, to generate triple transgenic mice which allowed for tdTomato fluorescent labeling of nNOS expression within these interneuron populations. To allow for postnatal specific expression of tdTomato in nNOS+ subtype populations, tamoxifen injections dosed at 75mg/kg were given at P10, P12, and P14 timepoints. We used 10 animals for each PDGFRβ:Ai14, nNOS:Ai14, nNOS:VIP:Ai65, 9 animals for nNOS:NPY:Ai65 and nNOS:PV:Ai65, 8 animals for nNOS:SST:Ai65 from both males and females as well as 7 animals (all males) for vGlut1:Ai75, and 9 animals (all males) for Gad2:Ai75. We used tail genomic DNA with PCR for genotyping. Brain samples were collected at 2 months old age for all mouse lines.

Perfusion and tissue processing for STPT imaging

Animals were deeply anesthetized with a ketamine-xylazine mixture (100 mg/kg ketamine, 10 mg/kg xylazine, i.p. injection) for both regular perfusion and vascular labeling. Transcardiac perfusion with a peristaltic pump (Ismatec, cat.no.: EW-78018-02) was used with 1X PBS followed by 4% paraformaldehyde, both injected through a small incision in the left ventricle, in order to wash out blood and allow for tissue fixation, respectively. Brains were dissected carefully in order to preserve all structures. For vessel labeling, transcardiac perfusion with a peristaltic pump (Welch, Model 3100) was used with 1X PBS followed by 4% paraformaldehyde at 0.3 ml/min, in order to wash out blood and for tissue fixation, respectively. To ensure that the large surface vessels would remain filled with the gel perfusate, the body of the mouse was tilted by 30° before gel perfusion (with the head tilted down), as previously described (Tsai et al., 2009). Following the fixative perfusion, the mouse was perfused at 0.6 ml/min with 5 ml of a 0.1% (w/v) fluorescein isothiocyanate (FITC) conjugated albumin (Sigma-Aldrich, cat.no.: A9771-1G) in a 2% (w/v) solution of porcine skin gelatin (Sigma-Aldrich, cat.no: G1890-500G) in 1X PBS. Immediately after perfusion, the heart, ascending and descending aorta as well as the superior vena cava, were all clamped with a hemostat (while the butterfly needle was simultaneously removed from the left ventricle). This served to prevent any pressure changes in or gel leakage from the brain vasculature. Next, the entire mouse body was submerged in an ice

510 bath to rapidly solidify the gel in the vessels. Then, the head was fixed in 4% PFA for one week,
511 followed by careful dissection of the brain to avoid damages to pial vessels. After fixation and
512 dissection, the brain was placed in 0.05M PB until imaging. Any animals that had poor perfusion
513 and/or possible air bubbles interfering with the gel perfusion were excluded from imaging and
514 any further analysis.
515
Serial two photon tomography (STPT) imaging.
Prior to imaging, the brain sample was embedded in oxidized agarose and cross-linked in 0.05M
sodium borohydrate at 4C for at least 2 days ahead of imaging (Kim et al., 2017; Newmaster et
al., 2019). This procedure allows for seamless cutting at 50μm thick sections, while also
preventing any tearing of the brain surface. The embedded brain sample was then glued to the
sample holder and fully submerged in 0.05M PB in an imaging chamber. For STPT imaging
(TissueCyte), we used 910nm excitation using a femtosecond laser (Coherent Ultra II) for all
samples. Signals in the green and red spectrum were simultaneously collected using a 560 nm
dichroic mirror (Chroma). For pericyte and neuronal subtypes, STPT imaging was conducted
with 1x1 μm (x,y) resolution in every 50 μm (z), with the imaging plane set at 40μm deep from
the surface, as previously described (Kim et al., 2017; Newmaster et al., 2019). For vascular
imaging, optical imaging (5 μm z step, 10 steps to cover 50μm in z) was added in the imaging,
producing 1x1x5 μm (x,y,z) resolution beginning at 20μm deep from the surface. Due to length
of imaging time required for vascular imaging, each brain sample was imaged through multiple
imaging runs to adjust the imaging window size in order to reduce overall imaging time.
531
Computational: STPT Image reconstruction
To measure and fix an optical aberration from an objective lens (Figure S1), we imaged a 25 μm
EM-grid (SPI supplies, cat.no.: 2145C) to represent the ground truth spatial data (Han et al.,
2018). We annotated all cross points of the grid and computed the B-spline transformation
profile from the grid image to the orthogonal coordinate sets using ImageJ (NIH). The pre-
scripted program then corrected every image tile by calling the ImageJ deformation function
using that profile. Afterwards, we used the entire set of imaged tiles (full mouse brain in this
case) to map out the tile-wise illumination profile. The images were grouped according to the
stage movement, which affects the photo-bleaching profile. The program avoids using pixels that
are considered empty background or dura artifacts using preset thresholds. Using those averaged
profile tiles, the program normalized all the tile images. Please note, this profile is unique for
each sample. Finally, the program picked 16 coronal slices (out of the nearly 2,000) with equal
spacing and utilized ImageJ's grid/collection stitching plugin to computationally stitch those 16
slides. The program then combined the transformation profiles from center to outer edge
according to the calculated pairwise shifting distance. It used a tile-intensity weighted average to
ensure the empty tiles did not contribute to the final profile. This approach significantly reduced
the computational time and allowed parallelization with no communication overhead. The
program automatically performed the aforementioned alignment and stitched the image set
together. The program finally aligned the image sets if the sample was imaged through multiple
runs during imaging acquisition.
552
Computational: Vessel Digitization/Tracing
We started with interpolating the data into 1x1x1 μm resolution with cubic interpolation then
subtracted the signal color channel (green) with the background color channel (red) to remove

auto-fluorescent backgrounds. Next we performed a voxel binarization. The voxel with at least
one of the following conditions passed as the foreground signal (vasculature), **a**. the voxel passed
a fixed threshold (6x that of the non-empty space average) or **b**. passed a threshold (2.4x that of
the non-empty space average) after subtracting a circular 35% local ranking filter. The binarized
image was then skeletonized using 26-neighbor rule (Kollmannsberger et al., 2017). The code
then reconnected lose ends that were within 10 μm distance and removed all the short stem/furs
shorter than 50 μm starting with the short ones and iterated until no more fur artifacts were found
(Figure S1F). By using the binary image and the skeleton (center-line), the radius for each
skeleton pixel can be measured. The code then grouped all the skeleton pixels into segments with
the branching nodes, and all the segments shorter than 2x radius were further cleaned up with
shortest graph path (Figure S1G). ROIs with poorly connected (<250 μm/node) were excluded in
further analysis as shown Figure S1H. Finally, the code documented and traced all the segments
and nodes with their connectivity, length, averaged radius, and raw skeleton locations. The full
pipeline here is programed to be fully automatic and the code was fully vectorized and
parallelized with reasonable memory consumption per thread (~8GB).
571
Computational: Fluid Dynamic Simulation
The goal of calculating and visualizing a permeability tensor is to illustrate how well fluid can
flow through the local microvasculature of a given volume in a given direction. Since the
direction distribution of the microvasculature can be anisotropic, the fluid flow can move with a
direction that is different from the pressure gradient direction, thus making the permeability in a
tensor form. Such a tensor can illustrate the local microvascular performance and its directional
characteristic.
The equation of permeability tensor is given by:

$$\overline{\overline{k}} \cdot \nabla P = \boldsymbol{Q}$$

or

$$\begin{bmatrix} k_{xx} & k_{yx} & k_{zx} \\ k_{xy} & k_{yy} & k_{zy} \\ k_{xz} & k_{yz} & k_{zz} \end{bmatrix} \begin{bmatrix} P_{,x} \\ P_{,y} \\ P_{,z} \end{bmatrix} = \begin{bmatrix} Q_x \\ Q_y \\ Q_z \end{bmatrix}$$

where k is the permeability tensor, P is the pressure, Q is the fluid flux, the subscript index is the
Cartesian coordinate direction, and the comma is partial differentiation. We chose a size of 400 x
400 x 400 μm as the local representative control volume. We then probed the system with three
∇P that are equal to three unit-vector on the Cartesian coordinate. The pressure profile was
applied on the surface of the cubical control volume, then the network flow profile was
calculated by solving the system of equations of the Hagen–Poiseuille equation (with the
viscosity set to unity for normalization) and conservation of flux. We chose the center cut plan to
measure the directional flux and consequently, the permeability. Finally, to illustrate the vascular
directionality of the isocortex, we projected the tensor onto penetrating, anterior-posterior,
medial-lateral vectors according to their location within the isocortex using the equation $k_{pj} =$
$\left| \overline{\overline{k}} \cdot \boldsymbol{n}_{pj} \right|$, where subscript pj indicates the direction of the projecting vessels.
594
595   **Computational: Deep Learning Neural Network (DLNN) pericyte counting**
596   We used a deep learning neural network (DLNN) to detect and classify cells. Instead of using a
597   fully convoluted neural network like Unet, we chose to use per-cell multi-resolution-hybrid
598   ResNet classification with potential cell locations. This makes the AI compute time significantly
599   shorter. The potential cell locations were identified with local maximum within a radius of r = 8

μm. The image around the potential cell locations was fed to the network with two different resolutions. One is 101 x101 μm (101 x 101 pixel) and the other one is 501 x 501 μm (201 x 201 pixel). The two-window system allows the network to capture characteristics from two zoom scales at the same time. In order to use global maximum at the end of the network, we stacked an empty (value zero) image onto the 3D direction of each image, which made them 101 x 101 x 2 and 201 x 201 x 2 pixels. We then assigned value 1 to the location of the potential cell, in this case, the center. At the end of the two networks of those two images, the intermediate images were flattened and concatenated into one. The classification was done with two bins, 'pericytes' and 'everything else.' The detailed schematic describing the network is in Figure S4. We deployed two human annotators with the same training to annotate the data, and only used the mutually agreed data to train the AI to eliminate human error and bias. We used a strict set of criteria to include only capillary pericytes that have two subtypes (junctional and helical pericytes). Cells were counted only when the cell body was in the imaging plane and clear pericyte cell morphology could be detected. Cells associated with larger vessels, often with vascular smooth muscle morphology, were not counted to prevent the inclusion of erroneous cell types. We also excluded transitional cell types often referred to as either ensheathing pericytes or precapillary arteriolar smooth muscle cells, due to the controversy in the field as to whether this should be included as a pericyte subtype (Attwell et al., 2016; Grant et al., 2019; Hartmann et al., 2015). A total 12,000 potential cell locations from multiple anatomical regions across 4 different brains were annotated by both annotators. 90% of the data selected at random was used to train the AI and the remaining 10% was used for validation. The 90% of the data taken for training was further truncated down to 3,400 potential cell locations with half positive and half negative for training. The positive cell selections in the raw data were around 19.6% (annotator #1) to 21.1% (annotator #2). The validation set was not truncated to represent actual performance. The performance can be found in Table S3.

Computational: Deep Learning Neural Network (DLNN) nNOS neuron counting
The morphology and size of tdTomato positive cells in the granular layer of the cerebellum from nNOS-CreER:Ai14 mice differs significantly from other tdTomato positive nNOS neurons in other brain regions. Thus, we developed new DLNN AI algorithms to consider not only cell morphology but also the location of cells by putting additional zoomed-out, low-resolution images of whole coronal sections. The network set-up is similar to the pericyte classification with one more image containing the coronal section with the cell location. The inputs are 101 x101 μm (101 x 101 pixel), 501 x 501 μm (201 x 201 pixel), and the full frame low resolution 12 x 8 mm (201 x 201 pixels). Similar to the pericyte network, we made images 101 x 101 x 2, 202 x 202 x 2, and 202 x 202 x 2 pixels with the cell location marked as value 1. At the end, those three sub-networks were flattened and concatenated into one. The classification is done with three bins, nNOS neurons, cerebellar granular nNOS neurons, and everything else. One human user created 10,000 annotations from 5 nNOS and 5 nNOS subtype brains. 5,000 cells from 5 brains were initially used to train the AI. Another 5,000 cells from 5 new brains were used to evaluate the AI performance. The AI reached an F1 score = 0.96, which is comparable to human performance. The details for the network are in Figure S4. The performance can be found in Table S3.

Isocortical flatmap

We started with Allen CCF annotation images to solve the Laplace equation by setting the
surface of cortical layer 1 as potential '1,' the surface of layer 6b as '0,' and the surface of
everything else as flux '0' (Wang et al., 2020). We used the potential map to find the gradient
direction as the projecting direction. The projection was first traced to the cortical surface and
then flattened at the Anterior-Posterior (A-P) tangential plane, which later preserved the A-P
coordinate on the flat map. The flattened map has the y-axis mapped as the original A-P
coordinate at the surface, and the x-axis was adjusted to represent the surface arc (azimuth)
length to the reference X-zero. The reference X-zero was defined on the cortical ridge in the
dorsal direction (maximum Y point in 3D) with a straight cut in the A-P direction. Finally, the
projection profile was saved at two resolutions, 10 x 10 x 10 μm^3 and 20 x 20 x 20 μm^3. We
created a Matlab script that can map any signals (previously registered to the Allen CCF) into a
3D projected isocortical flatmap.
657
Conversion of 2D based counting to 3D cell density
STPT imaging has very accurate cutting and stage depth movement, which allows us to convert
the 2D cell counting to 3D cell density. We used previously calculated 3D conversion factors for
cytoplasmic (factor = 1.4) and nuclear signals (factor = 1.5) to generate density estimates of
nNOS neurons and other neuronal cell type datasets (Kim et al., 2017). To estimate the 3D
conversion factor for pericytes, we imaged one PDGFRβ-Cre:Ai14 mouse brain with 1 x 1 x 5
μm, as done with vascular imaging (Figure 1B). Then, we applied our pericyte cell counting in
all images from this densely imaged dataset. Since the DLNN cell counting was done in 2D, the
same cell can be counted more than once in adjacent z stack images. We resolved this issue by
compiling the counting into a 5 x 5 x 5 μm^3 resolution 3D array, dilating the mask by 1 pixel,
and counting the connected pieces, which avoids over counting any cell that is within 25 μm
proximity. This 3D counting result was then compared to the 2D counting from the 5[th] section
from every ten z optical image stack (Figure 1B). The calculated 3D/2D ratio was 3.33 for
pericytes, which was applied as a conversion factor to estimate pericyte numbers in 3D.
To estimate the anatomical volume from each sample, the Allen CCF was registered to
individual samples first using Elastix (Klein et al., 2010). Anatomical labels were
transformed based on the registration parameters and the number of voxels associated with
specific anatomical IDs were used to estimate the 3D volume of each anatomical area (Kim et
al., 2017).
677
In vivo two-photon recording and comparison with STPT vascular measurement
Surgery. All surgeries were performed under isoflurane anesthesia (in oxygen, 5% for induction
and 1.5-2% for maintenance). A custom-machined titanium head bolt was attached to the skull
with cyanoacrylate glue (#32002, Vibra-tite). The head bolt was positioned along the midline
and just posterior to the lambda cranial suture. Two self-tapping 3/32" #000 screws (J.I. Morris)
were implanted into the skull contralateral to the measurement sites over the frontal lobe and
parietal lobe. For measurements using two-photon laser scanning microscopy (2PLSM), a
polished and reinforced thin-skull (PoRTS) window was made covering the right somatosensory
cortex as described previously (Drew et al., 2010; Zhang et al., 2019a). Following the surgery,
mice were returned to their home cage for recovery for at least one week, and then started
habituation on experimental apparatus. Habituation sessions were performed 2-4 times over the
course of one week, with the duration increasing from 5 min to 45 min.

Measurements using two-photon laser scanning microscopy (2PLSM). Mice were briefly
anesthetized with isoflurane (5% in oxygen) and retro-orbitally injected with 50 µL 5%
(weight/volume in saline) fluorescein-conjugated dextran (70 kDa, Sigma-Aldrich, cat.no.:
46945), and then fixed on a spherical treadmill. Imaging was done on a Sutter Movable
Objective Microscope with a 20X, 1.0 NA water dipping objective (Olympus, XLUMPlanFLN).
A MaiTai HP (Spectra-Physics, Santa Clara, CA) laser tuned to 800 µm was used for
fluorophore excitation. All imaging with the water-immersion lens was done with room
temperature distilled water between the PoRTS window and the objective. All the 2PLSM
measurements were started at least 20 minutes after isoflurane exposure to reduce the disruption
of physiological signals due to anesthetics (Gao et al., 2017). High-resolution image stacks of the
vasculature were collected across a 500 by 500 µm field and up to a depth of 250 um from the
pial surface. All the images were acquired with increasing laser power up to 100 mW at a depth
of ~200 µm. Lateral sampling was 0.64 um per pixel and axial sampling was at 1 um steps
between frames. Shortly (within 20 minutes) after the imaging, the mouse was perfused with
FITC filling for STPT based *ex vivo* vasculature imaging.
705
In vivo and ex vivo comparison.
In order to compare our measurements for vessel radii in STPT imaging datasets to vessel
parameters measured *in vivo*, the same animals that were used for 2PLSM (See **In vivo two-**
photon recording and comparison with STPT vascular measurement) underwent the FITC-
fill perfusion and STPT imaging steps described above. However, STPT imaging was only
conducted on the cortical hemisphere used for 2PLSM, with imaging spanning from prefrontal
regions to visual cortex regions, in order to appropriately capture the primary somatosensory
cortex limb region. Following stitching and tracing of the images, the raw imaging data was 3D
reconstructed in order to visualize the cortical surface. To find the 2PLSM imaging window,
vessel landmarks used to reimage in 2PLSM were again used to identify the same landmark
vessels in the STPT imaging dataset (Figure S2). The region of interest was further confirmed by
anatomical landmarks (proximity to bregma, surface vessels, etc.) through overlay of STPT and
2PLSM imaging window regions. Next, within a STPT imaging z stack, borders were inserted
using ImageJ software to further outline the in vivo imaging window region in the 3D data. Then
the in vivo imaging z stack data were used to identify branch points along the penetrating vessel
tracked during 2PLSM. This provided identifiable characteristics to further locate the same
vessel in the STPT imaging dataset. Once the exact vessel was identified in the STPT images, the
precise 3D coordinates were tracked to accurately obtain the radii measurements from the traced
vessel data, see the **Computational: Vessel Digitization/Tracing section** for details. In 2PLSM
images, vessel diameter measurements were manually taken with adjusted pixel/micron distances
using the straight-line function in ImageJ. These vessel diameter measurements accounted for the
lumen of the vessel, at half of the maximum fluorescence intensity profile and were adjusted for
pixelation of 2PLSM data. These measurements have been further refined through VasoMetrics
ImageJ macro (McDowell et al., 2021). To identify the radii and diameter measurements from
the STPT imaging data, exact vessel coordinates were used to retrieve the associated vessel radii
measurements using custom MATLAB code. A minimum of 10 vessel diameter measurements
were taken per imaging window (each animal contained 2 imaging regions of interest) per
animal.
734
735

Statistical analysis

All statistical analysis, including multi-region of interest (ROI) correlation analysis, was done in Matlab (Mathworks). We used an averaged value of the experimented animals while treating each ROI as an individual data point to calculate the correlation coefficient R between vascular and cell density measurements. The p value was calculated based on the null hypothesis that the two groups have no correlation; the values were adjusted with the Bonferroni correction for multiple comparison correction.

Acknowledgments

This publication was made possible by an NIH grant R01NS108407 to Y.K., R01NS078168 and R01NS101353 to P.J.D. Its contents are solely the responsibility of the authors and do not necessarily represent the views of the funding agency. We thank Dr. Volkhard Linder for kindly sharing PDGFRβ-Cre transgenic mice and Rebecca Betty for assistance in editing the manuscript. We acknowledge use of computational resources in the High Performance Computing cluster at the Penn State College of Medicine.

Contributions

Conceptualization, Y.K.; Data Collection, H.C.B, U.C., Y.K.; Developing Computational Analysis, Y.W.; Data Analysis, Y.W., H.C.B., U.C.; In vivo two-photon imaging, Q.Z., P.J.D.;Neuronal subtypes STPT data collection; R.M., P.O.; Web visualization, D.J.V., K.C.; Manuscript preparation: Y.K., Y.W., H.C.B. with help from other authors.

Competing Interests

The authors declare no competing interests.

Data Sharing Plan

High-resolution serial two-photon tomography images can be found at https://kimlab.io/brain-map/nvu/

Custom-built codes including isocortical flatmaps are available as Supplementary data (Code S1-S3)

Figure 1. High resolution 3D mapping of the entire vasculature in the mouse brain.
A. Fluorescent dye (FITC)-conjugated albumin gel perfusing the mouse brain through the heart
to label cerebrovasculature. **B**. Combination of physical sectioning (vibratome cutting) and
optical sectioning to achieve lossless imaging of a sample. Left: black-plane indicates the
physical cuts, yellow-zones indicates an optical sectioning imaging plane. Right: 10 optical
imaging sections per one physical sectioning. **C-D**. Stitching with optical aberration and tile line
correction (D) from uncorrected images (C). **E-G**. Example outputs from each stage of the
analysis pipeline. Top row: 100 μm thick 3D volume from the white box area from the middle
row, Middle row: An example coronal section, Bottom row: The whole brain results. **E.** The raw

image volume of FITC labeled vasculature. **F**. The binarized vasculature. **G**. The traced vasculature. Large (radius > 5 μm) and small vessels are colored as red and green, respectively in the top and middle image. The bottom image shows the vasculature density. **H-J**. The averaged vasculature length density (H), branching density (I), and radii (J) from four brains are registered to the Allen CCF and displayed with heatmap in 8 major areas across the whole mouse brain. **K**. Average vessel density, radius, and branching density across the whole brain. Individual animal data that pass the tracing quality control as shown in Figure S1 are shown as X. See also Table S1. **L**. Examples of areas with different vasculature density and radii. **M-N**. The correlation between vessel density and branching density (M) and the correlation between vessel density and the averaged radius (N). Size of each ROI is displayed according to relative volume of the area. Note strong correlation between vessel density with branching density, but not with vascular radii.

Figure 2. Heterogeneous vascular arrangements in the isocortex

A. Examples of cortical areas with different vasculature structures. **B**. Average vessel density, radius, and branching density in the isocortex. See also Table S1. **C-E.** Creating an isocortical flatmap. **C.** Anatomical borderlines of the Allen CCF. **D**. Gradient vectors from solving the Laplace equation by setting cortical layer 1 and layer 6 as end points. **E.** The flattened projected-profile. Each line is 0.2 mm apart. The four arrowheads indicate the flattening direction while maintaining the original A-P coordinates. **F**. The cortical flatmap with Allen CCF border line. Y axis: Bregma anterior-posterior (A-P) coordinates, X axis: Azimuth coordinate represents the physical distance by tracing the cortical surface on the coronal cut. **G**. The averaged vasculature length and branching density as well as vessel radius plotted onto the cortical flat map. Note the high density of vasculature in the somatosensory (gray arrowhead), auditory (white arrowhead),

809　　and retrosplenial (black arrowhead) cortex, while there is a low density in lateral association
810　　cortex (white arrow). **H**. Cortical layer specific max projection of vasculature length density.
811
812

Figure 3. Anisotropy of the cerebral microvascular network and its impact on permeability

A. The flow chart of the permeability fluid dynamic simulation. From left to right: Original

traced data, applying pressure profile on the surface of the control volume with a gradient profile

and solving the coincide flux equation set for the permeability tensor, the annotation rule of the

permeability tensor, and the sampling dots of the numerical spherical integration. Equation

symbols: l: perfusing length, P: pressure, Q: Blood flux, R: resistance, μ: viscosity, k:

permeability, Δ: changing of the quantity, $\cdot$: dot product, ∇: gradient, $\oiint s$: spherical integral, S:

spherical surface, n: normal direction, bold-font: vector, double-top-bar: tensor. **B**. Examples

illustrating how the structure of the vasculature network impacts the permeability tensor. **C**. Two

examples of the real network with its permeability tensor from primary somatosensory, barrel

field (SSp-bfd), and medial orbital cortex (ORBm). **D**. Spherically integrated permeability results

in the cortical flatmap. **E-F**. Microvessel anisotropy measurement in the isocortex. Examples

from the motor cortex (E) and somatosensory (F) cortices. **G.** Microvessels in (E-F) are colored

based on its directionality; magenta for penetrating (P), cyan for anterior-posterior (AP), yellow

for medial-lateral (ML). Directionality axis is based on cortical column angles in each area. **H**.
Permeability measurement of the secondary motor cortex (MOs) and the SSp-bfd in the three
directionalities. **I**. Microvessel directionality in cortical layers. Only the dominant direction is
displayed for simplicity. **J.** Permeability results of the whole brain. See also Table S2.
833
834

835
836
Figure 4: Pericyte density mapping across the whole mouse brain
A. Schematic depicting genetic construct for PDGFRβ-Cre:Ai14 mouse model. **B.**
Representative coronal images and a reconstructed brain from the STPT imaging of a PDGFRβ-
Cre:Ai14 mouse. **C.** Example images of areas showing variability in pericyte density. High
pericyte density in the auditory (top left) and somatosensory cortex (top right) while low density
in the prelimbic (bottom left) and retrosplenial cortex (bottom right). **D.** Cortical flatmap of
averaged pericyte density across the isocortex by cortical layer. **E.** Scatter plot demonstrating
significantly positive correlation between pericyte density and vascular length density in
isocortical regions (R=0.859, p value =1.86×10^{-12}). **F.** Heat maps of pericyte density distribution
across the adult brain. **G.** Scatterplot comparison of average vessel length density and pericyte
density across all brain regions shows significant correlation (R=0.4818, p value = 2.65×10^{-6}).
H. Quantifications of pericyte density across all brain regions. Brain regions are color coded to
match the regions represented in (G). See also Table S4.

Figure 5: Brain-wide density map of nNOS neurons and their subtypes
A. Cell type-specific transgenic mice with tamoxifen treatments to label either total nNOS neurons or nNOS subtypes co-expressing NPY, SST, PV, or VIP. **B.** Coronal section examples from STPT Imaging of a nNOS:Ai14 mouse. **C-E.** AI based detection of nNOS cells with two distinct shapes. High resolution image (C) showing nNOS neurons in the molecular (green) and granular cell layer (red) of the cerebellum (D). **E.** Representative image of total nNOS population throughout the brain. **F.** Heat maps demonstrating the distribution of total nNOS and nNOS subtype populations. See also Movie S4 and Table S5. **G.** Representative raw images of nNOS, nNOS/NPY, nNOS/SST, nNOS/PV and nNOS/VIP neurons in the olfactory bulb, hippocampus, medial amygdala (MEA), dorsomedial hypothalamus (DMH) and cerebellum. Reference atlas images included on nNOS/VIP images show the area displayed for each region of interest. Main olfactory bulb (MOB), Ammon's horn (CA1), dentate gyrus (DG), optic tract (opt), fornix (fx), granular (gr), molecular (mo). **H.** Graph showing nNOS density by brain region for the total nNOS neurons and their subtypes. Size of circle corresponds to density as shown in the key at the bottom. **I-J.** Isocortical flatmap for averaged density of total nNOS neurons by layer (I) and overall density of nNOS subtypes in the isocortex (J). Note higher nNOS density in the medial prefrontal and lateral association areas. **K.** Representative STPT images of nNOS cell types from the primary somatosensory cortex and the agranular insular cortex. **L.** Scatter plot showing significant negative correlation between total nNOS density and vessel length density in the isocortex (R value =-0.806, p value = 6.9×10^{-7}). **M.** Correlation matrix between nNOS cell types and vascular/pericyte measurements. *p<0.05, **p<0.005 after the Bonferroni correction. (-) denote negative correlation.

Figure 6: Cortical parvalbumin and vGlut1 neurons positively correlated with vasculature density

A. Cell type-specific labeling of neurons to visualize cortical inhibitory interneuron and
excitatory neuron populations. **B.** Distribution of inhibitory interneurons, excitatory neurons, and
the vasculature and its tracing result (large vessels=red, microvasculature=green) from the
densely vascularized primary somatosensory and sparsely vascularized infralimbic cortices. **C.**
Cortical flatmap showing density distributions of neuronal subtypes as well as vessel length and
pericyte densities. **D.** Correlation between vascular density and neuronal subtypes. Note very
strong positive correlation with PV density (R = 0.968, p value = 8.5×10^{-22}) and positive
correlation with vGlut1 excitatory neuronal density (R= 0.552, p value = 5.9×10^{-3}). **E.**
Correlation matrix between neuronal subtypes, vessel length density and pericyte density.
*p<0.05, **p<0.005 after the Bonferroni correction.
889
890

891
892
893 **Figure 7: Cortical organization of the vascular/pericyte network and neuronal cell types**
894 Sensory cortices including the somatosensory cortex are characterized by relatively high density
895 of vessels, pericytes, PV interneurons, and vGlut1 excitatory neurons, and a low density of
896 nNOS neurons. In contrast, association cortices show the opposite pattern.
897
898

Region of Interest	Vessel length density (m/mm^3)	Pericyte (cell/mm^3)	nNOS neurons (cell/mm^3)	PV neurons (cell/mm^3)	vGlut neurons (cell/mm^3)
Motor	0.81 ± 0.05	6,603 ± 489	104 ± 17	2,678 ± 203	35,566 ± 823
Somatosensory	0.91 ± 0.07	8,208 ± 739	108 ± 16	3,664 ± 281	45,022 ± 838
Auditory	0.86 ± 0.06	8,006 ± 848	121 ± 16	2,965 ± 295	42,178 ± 1,676
Visual	0.82 ± 0.07	6,930 ± 697	112 ± 22	2,927 ± 352	49,329 ± 1,132
Retrosplenial	0.92 ± 0.07	7,290 ± 869	103 ± 20	3,330 ± 387	44,631 ± 2,494
Posterior Parietal	0.87 ± 0.07	7,428 ± 641	91 ± 17	3,220 ± 255	47,030 ± 875
Orbital	0.75 ± 0.08	6,338 ± 891	106 ± 17	2,290 ± 320	38,014 ± 1,559
Anterior Cingulate	0.8 ± 0.05	6,856 ± 555	154 ± 32	2,371 ± 213	41,208 ± 1,039
Prelimbic	0.65 ± 0.04	5,645 ± 726	171 ± 34	1,199 ± 171	39,270 ± 1,098
Infralimbic	0.64 ± 0.02	6,139 ± 602	232 ± 38	503 ± 120	43,679 ± 1,973
Visceral	0.74 ± 0.04	6,837 ± 689	177 ± 33	1,846 ± 204	37,081 ± 1,546
Gustatory	0.73 ± 0.03	7,421 ± 581	194 ± 34	1,652 ± 212	40,783 ± 1,467
Agranular Insular	0.65 ± 0.05	5,966 ± 500	207 ± 41	1,154 ± 118	31,290 ± 1,023
Perirhinal	0.63 ± 0.03	5,232 ± 615	265 ± 69	979 ± 142	24,789 ± 3,196
Temporal Association	0.7 ± 0.04	6,596 ± 639	187 ± 37	1,189 ± 210	42,407 ± 1,907
Ectorhinal	0.61 ± 0.02	5,967 ± 510	283 ± 63	623 ± 117	38,578 ± 3,400

Table 1: Density of the cerebrovasculature, pericytes, and neuronal subtypes in the isocortex

Number = average ± standard deviation, see Table S1, S3, S5, S6 for full dataset.

References

Attwell, D., Buchan, A.M., Charpak, S., Lauritzen, M., Macvicar, B.A., and Newman, E.A. (2010). Glial and neuronal control of brain blood flow. Nature *468*, 232–243.

Attwell, D., Mishra, A., Hall, C.N., O'Farrell, F.M., and Dalkara, T. (2016). What is a pericyte? J Cereb Blood Flow Metab *36*, 451–455.

Ballinger, E.C., Ananth, M., Talmage, D.A., and Role, L.W. (2016). Basal Forebrain Cholinergic Circuits and Signaling in Cognition and Cognitive Decline. Neuron *91*, 1199–1218.

Balogh, P., and Bagchi, P. (2019). Three-dimensional distribution of wall shear stress and its gradient in red cell-resolved computational modeling of blood flow in in vivo-like microvascular networks. Physiological Reports *7*, e14067.

Bennett, H.C., and Kim, Y. (2021). Pericytes Across the Lifetime in the Central Nervous System. Front. Cell. Neurosci. *15*.

Blinder, P., Shih, A.Y., Rafie, C., and Kleinfeld, D. (2010). Topological basis for the robust distribution of blood to rodent neocortex. Proc Natl Acad Sci U S A *107*, 12670–12675.

Blinder, P., Tsai, P.S., Kaufhold, J.P., Knutsen, P.M., Suhl, H., and Kleinfeld, D. (2013). The cortical angiome: an interconnected vascular network with noncolumnar patterns of blood flow. Nature Neuroscience *16*, 889–897.

Boido, D., Rungta, R.L., Osmanski, B.-F., Roche, M., Tsurugizawa, T., Le Bihan, D., Ciobanu, L., and Charpak, S. (2019). Mesoscopic and microscopic imaging of sensory responses in the same animal. Nat Commun *10*, 1110.

Buzsáki, G., Kaila, K., and Raichle, M. (2007). Inhibition and brain work. Neuron *56*, 771–783.

Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., and Moore, C.I. (2009). Driving fast-spiking cells induces gamma rhythm and controls sensory responses. Nature *459*, 663–667.

Chachlaki, K., Malone, S.A., Qualls-Creekmore, E., Hrabovszky, E., Münzberg, H., Giacobini, P., Ango, F., and Prevot, V. (2017). Phenotyping of nNOS neurons in the postnatal and adult female mouse hypothalamus. J. Comp. Neurol. *525*, 3177–3189.

Cuttler, A.S., LeClair, R.J., Stohn, J.P., Wang, Q., Sorenson, C.M., Liaw, L., and Lindner, V. (2011). Characterization of Pdgfrb-Cre transgenic mice reveals reduction of ROSA26 reporter activity in remodeling arteries. Genesis (New York, N.Y. : 2000) *49*, 673–680.

Devonshire, I.M., Papadakis, N.G., Port, M., Berwick, J., Kennerley, A.J., Mayhew, J.E.W., and Overton, P.G. (2012). Neurovascular coupling is brain region-dependent. Neuroimage *59*, 1997–2006.

Douglas, R.J., and Martin, K.A.C. (2004). Neuronal circuits of the neocortex. Annu Rev
Neurosci *27*, 419–451.

Drew, P.J., Shih, A.Y., Driscoll, J.D., Knutsen, P.M., Blinder, P., Davalos, D., Akassoglou, K.,
Tsai, P.S., and Kleinfeld, D. (2010). Chronic optical access through a polished and reinforced
thinned skull. Nat Methods *7*, 981–984.

Drew, P.J., Mateo, C., Turner, K.L., Yu, X., and Kleinfeld, D. (2020). Ultra-slow Oscillations in
fMRI and Resting-State Connectivity: Neuronal and Vascular Contributions and Technical
Confounds. Neuron *107*, 782–804.

Echagarruga, C.T., Gheres, K.W., Norwood, J.N., and Drew, P.J. (2020). nNOS-expressing
interneurons control basal and behaviorally evoked arterial dilation in somatosensory cortex of
mice. Elife *9*.

Gao, Y.-R., Ma, Y., Zhang, Q., Winder, A.T., Liang, Z., Antinori, L., Drew, P.J., and Zhang, N.
(2017). Time to wake up: Studying neurovascular coupling and brain-wide circuit function in the
un-anesthetized animal. NeuroImage *153*, 382–398.

Grant, R.I., Hartmann, D.A., Underly, R.G., Berthiaume, A.-A., Bhat, N.R., and Shih, A.Y.
(2019). Organizational hierarchy and structural diversity of microvascular pericytes in adult
mouse cortex. Journal of Cerebral Blood Flow and Metabolism : Official Journal of the
International Society of Cerebral Blood Flow and Metabolism *39*, 411–425.

Hall, C.N., Klein-Flügge, M.C., Howarth, C., and Attwell, D. (2012). Oxidative phosphorylation,
not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information
processing. J Neurosci *32*, 8940–8951.

Hall, C.N., Reynell, C., Gesslein, B., Hamilton, N.B., Mishra, A., Sutherland, B.A., O'Farrell,
F.M., Buchan, A.M., Lauritzen, M., and Attwell, D. (2014). Capillary pericytes regulate cerebral
blood flow in health and disease. Nature *508*, 55–60.

Han, Y., Kebschull, J.M., Campbell, R.A.A., Cowan, D., Imhof, F., Zador, A.M., and Mrsic-
Flogel, T.D. (2018). The logic of single-cell projections from visual cortex. Nature *556*, 51–56.

Hartmann, D.A., Underly, R.G., Grant, R.I., Watson, A.N., Lindner, V., and Shih, A.Y. (2015).
Pericyte structure and distribution in the cerebral cortex revealed by high-resolution imaging of
transgenic mice. Neurophotonics *2*, 041402.

Hartmann, D.A., Berthiaume, A.-A., Grant, R.I., Harrill, S.A., Koski, T., Tieu, T., McDowell,
K.P., Faino, A.V., Kelly, A.L., and Shih, A.Y. (2021). Brain capillary pericytes exert a
substantial but slow influence on blood flow. Nat Neurosci.

He, M., Tucciarone, J., Lee, S., Nigro, M.J., Kim, Y., Levine, J.M., Kelly, S.M., Krugikov, I.,
Wu, P., Chen, Y., et al. (2016). Strategies and Tools for Combinatorial Targeting of GABAergic
Neurons in Mouse Cerebral Cortex. Neuron.

Herculano-Houzel, S. (2011). Scaling of Brain Metabolism with a Fixed Energy Budget per
Neuron: Implications for Neuronal Activity, Plasticity and Evolution. PLOS ONE *6*, e17514.

Hill, R.A., Tong, L., Yuan, P., Murikinati, S., Gupta, S., and Grutzendler, J. (2015). Regional
Blood Flow in the Normal and Ischemic Brain Is Controlled by Arteriolar Smooth Muscle Cell
Contractility and Not by Capillary Pericytes. Neuron *87*, 95–110.

Hosford, P.S., and Gourine, A.V. (2019). What is the key mediator of the neurovascular coupling
response? Neurosci Biobehav Rev *96*, 174–181.

Howarth, C., Gleeson, P., and Attwell, D. (2012). Updated energy budgets for neural
computation in the neocortex and cerebellum. J Cereb Blood Flow Metab *32*, 1222–1232.

Howarth, C., Mishra, A., and Hall, C.N. (2021). More than just summed neuronal activity: how
multiple cell types shape the BOLD response. Philos Trans R Soc Lond B Biol Sci *376*,
20190630.

Hu, H., and Jonas, P. (2014). A supercritical density of Na + channels ensures fast signaling in
GABAergic interneuron axons. Nature Neuroscience *17*, 686–693.

Huo, B.-X., Smith, J.B., and Drew, P.J. (2014). Neurovascular coupling and decoupling in the
cortex during voluntary locomotion. Journal of Neuroscience *34*, 10975–10981.

Inan, M., Zhao, M., Manuszak, M., Karakaya, C., Rajadhyaksha, A.M., Pickel, V.M., Schwartz,
T.H., Goldstein, P.A., and Manfredi, G. (2016). Energy deficit in parvalbumin neurons leads to
circuit dysfunction, impaired sensory gating and social disability. Neurobiol Dis *93*, 35–46.

Ji, X., Ferreira, T., Friedman, B., Liu, R., Liechty, H., Bas, E., Chandrashekar, J., and Kleinfeld,
D. (2021). Brain microvasculature has a common topology with local differences in geometry
that match metabolic load. Neuron.

Kann, O. (2016). The interneuron energy hypothesis: Implications for brain disease. Neurobiol
Dis *90*, 75–85.

Kaplan, L., Chow, B.W., and Gu, C. (2020). Neuronal regulation of the blood-brain barrier and
neurovascular coupling. Nat Rev Neurosci *21*, 416–432.

Kepecs, A., and Fishell, G. (2014). Interneuron cell types are fit to function. Nature *505*, 318–
326.

Kim, Y., Yang, G.R., Pradhan, K., Venkataraju, K.U., Bota, M., García Del Molino, L.C.,
Fitzgerald, G., Ram, K., He, M., Levine, J.M., et al. (2017). Brain-wide Maps Reveal
Stereotyped Cell-Type-Based Cortical Architecture and Subcortical Sexual Dimorphism. Cell
171, 456-469.e22.

Kirst, C., Skriabine, S., Vieites-Prado, A., Topilko, T., Bertin, P., Gerschenfeld, G., Verny, F.,
Topilko, P., Michalski, N., Tessier-Lavigne, M., et al. (2020). Mapping the Fine-Scale
Organization and Plasticity of the Brain Vasculature. Cell *0*.

Klein, S., Staring, M., Murphy, K., Viergever, M.A., and Pluim, J.P.W. (2010). elastix: a toolbox
for intensity-based medical image registration. IEEE Transactions on Medical Imaging *29*, 196–
205.

Kollmannsberger, P., Kerschnitzki, M., Repp, F., Wagermaier, W., Weinkamer, R., and Fratzl, P.
(2017). The small world of osteocytes: connectomics of the lacuno-canalicular network in bone.
New J. Phys. *19*, 073019.

Krawchuk, M.B., Ruff, C.F., Yang, X., Ross, S.E., and Vazquez, A.L. (2020). Optogenetic
assessment of VIP, PV, SOM and NOS inhibitory neuron activity and cerebral blood flow
regulation in mouse somato-sensory cortex. J Cereb Blood Flow Metab *40*, 1427–1440.

Lecrux, C., and Hamel, E. (2011). The neurovascular unit in brain function and disease. Acta
Physiologica (Oxford, England) *203*, 47–59.

Lee, L., Boorman, L., Glendenning, E., Christmas, C., Sharp, P., Redgrave, P., Shabir, O.,
Bracci, E., Berwick, J., and Howarth, C. (2020). Key Aspects of Neurovascular Control
Mediated by Specific Populations of Inhibitory Cortical Interneurons. Cereb. Cortex *30*, 2452–
2464.

Lourenço, C.F., Ledo, A., Barbosa, R.M., and Laranjinha, J. (2017). Neurovascular-
neuroenergetic coupling axis in the brain: master regulation by nitric oxide and consequences in
aging and neurodegeneration. Free Radical Biology and Medicine *108*, 668–682.

McDowell, K.P., Berthiaume, A.-A., Tieu, T., Hartmann, D.A., and Shih, A.Y. (2021).
VasoMetrics: unbiased spatiotemporal analysis of microvascular diameter in multi-photon
imaging applications. Quant Imaging Med Surg *11*, 969–982.

Melzer, S., and Monyer, H. (2020). Diversity and function of corticopetal and corticofugal
GABAergic projection neurons. Nature Reviews Neuroscience *21*, 499–515.

Mitchell, A.S., Czajkowski, R., Zhang, N., Jeffery, K., and Nelson, A.J.D. (2018). Retrosplenial
cortex and its role in spatial cognition. Brain Neurosci Adv *2*.

Montagne, A., Barnes, S.R., Sweeney, M.D., Halliday, M.R., Sagare, A.P., Zhao, Z., Toga,
A.W., Jacobs, R.E., Liu, C.Y., Amezcua, L., et al. (2015). Blood-brain barrier breakdown in the
aging human hippocampus. Neuron *85*, 296–302.

Nelson, A.R., Sagare, M.A., Wang, Y., Kisler, K., Zhao, Z., and Zlokovic, B.V. (2020).
Channelrhodopsin Excitation Contracts Brain Pericytes and Reduces Blood Flow in the Aging
Mouse Brain in vivo. Front Aging Neurosci *12*, 108.

Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M., Hidema, S.,
Nishimori, K., Hammock, E.A.D., and Kim, Y. (2019). Quantitative Cellular-Resolution Map of
the Oxytocin Receptor in Postnatally Developing Mouse Brains. BioRxiv 719229.

Nikolakopoulou, A.M., Zhao, Z., Montagne, A., and Zlokovic, B.V. (2017). Regional early and
progressive loss of brain pericytes but not vascular smooth muscle cells in adult mice with
disrupted platelet-derived growth factor receptor-β signaling. PLoS One *12*, e0176225.

Nikolakopoulou, A.M., Montagne, A., Kisler, K., Dai, Z., Wang, Y., Huuskonen, M.T., Sagare,
A.P., Lazic, D., Sweeney, M.D., Kong, P., et al. (2019). Pericyte loss leads to circulatory failure
and pleiotrophin depletion causing neuron loss. Nature Neuroscience *22*, 1089–1098.

Packer, A.M., and Yuste, R. (2011). zhang. J. Neurosci. *31*, 13260–13271.

Perrenoud, Q., Geoffroy, H., Gauthier, B., Rancillac, A., Alfonsi, F., Kessaris, N., Rossier, J.,
Vitalis, T., and Gallopin, T. (2012). Characterization of Type I and Type II nNOS-Expressing
Interneurons in the Barrel Cortex of Mouse. Frontiers in Neural Circuits *6*, 36.

Pi, H.-J., Hangya, B., Kvitsiani, D., Sanders, J.I., Huang, Z.J., and Kepecs, A. (2013). Cortical
interneurons that specialize in disinhibitory control. Nature *503*, 521–524.

Rungta, R.L., Chaigneau, E., Osmanski, B.-F., and Charpak, S. (2018). Vascular
Compartmentalization of Functional Hyperemia from the Synapse to the Pia. Neuron *99*, 362-
375.e4.

Sengillo, J.D., Winkler, E.A., Walker, C.T., Sullivan, J.S., Johnson, M., and Zlokovic, B.V.
(2013). Deficiency in Mural Vascular Cells Coincides with Blood–Brain Barrier Disruption in
Alzheimer's Disease. Brain Pathology *23*, 303–310.

Shih, Y.-Y.I., Chen, C.-C.V., Shyu, B.-C., Lin, Z.-J., Chiang, Y.-C., Jaw, F.-S., Chen, Y.-Y., and
Chang, C. (2009). A new scenario for negative functional magnetic resonance imaging signals:
endogenous neurotransmission. J Neurosci *29*, 3036–3044.

Sohal, V.S., Zhang, F., Yizhar, O., and Deisseroth, K. (2009). Parvalbumin neurons and gamma
rhythms enhance cortical circuit performance. Nature *459*, 698–702.

Sweeney, M.D., Ayyadurai, S., and Zlokovic, B.V. (2016). Pericytes of the neurovascular unit:
key functions and signaling pathways. Nature Neuroscience *19*, 771–783.

Sweeney, M.D., Kisler, K., Montagne, A., Toga, A.W., and Zlokovic, B.V. (2018). The role of
brain vasculature in neurodegenerative disorders. Nature Neuroscience *21*, 1318–1331.

Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., Kvitsani, D., Fu, Y., Lu, J., Lin, Y.,
et al. (2011). A Resource of Cre Driver Lines for Genetic Targeting of GABAergic Neurons in
Cerebral Cortex. Neuron *71*, 995–1013.

Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,
Garren, E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct transcriptomic
cell types across neocortical areas. Nature *563*, 72–78.

Todorov, M.I., Paetzold, J.C., Schoppe, O., Tetteh, G., Shit, S., Efremov, V., Todorov-Völgyi,
K., Düring, M., Dichgans, M., Piraud, M., et al. (2020). Machine learning analysis of whole
mouse brain vasculature. Nature Methods *17*, 442–449.

Tomioka, R., and Rockland, K.S. (2007). Long-distance corticocortical GABAergic neurons in
the adult monkey white and gray matter. J Comp Neurol *505*, 526–538.

Tomioka, R., Okamoto, K., Furuta, T., Fujiyama, F., Iwasato, T., Yanagawa, Y., Obata, K.,
Kaneko, T., and Tamamaki, N. (2005). Demonstration of long-range GABAergic connections
distributed throughout the mouse neocortex. Eur J Neurosci *21*, 1587–1600.

Tricoire, L., and Vitalis, T. (2012). Neuronal nitric oxide synthase expressing neurons: a journey
from birth to neuronal circuits. Frontiers in Neural Circuits *6*, 82.

Tsai, P.S., Kaufhold, J.P., Blinder, P., Friedman, B., Drew, P.J., Karten, H.J., Lyden, P.D., and
Kleinfeld, D. (2009). Correlations of neuronal and microvascular densities in murine cortex
revealed by direct counting and colocalization of nuclei and vessels. The Journal of
Neuroscience : The Official Journal of the Society for Neuroscience *29*, 14553–14570.

Vazquez, A.L., Fukuda, M., and Kim, S.-G. (2018). Inhibitory Neuron Activity Contributions to
Hemodynamic Responses and Metabolic Load Examined Using an Inhibitory Optogenetic
Mouse Model. Cereb Cortex *28*, 4105–4119.

van Veluw, S.J., Hou, S.S., Calvo-Rodriguez, M., Arbel-Ornath, M., Snyder, A.C., Frosch, M.P.,
Greenberg, S.M., and Bacskai, B.J. (2020). Vasomotion as a Driving Force for Paravascular
Clearance in the Awake Mouse Brain. Neuron *105*, 549-561.e5.

Vergara, R.C., Jaramillo-Riveri, S., Luarte, A., Moënne-Loccoz, C., Fuentes, R., Couve, A., and
Maldonado, P.E. (2019). The Energy Homeostasis Principle: Neuronal Energy Regulation Drives
Local Network Dynamics Generating Behavior. Front Comput Neurosci *13*.

Wang, Q., Ding, S.-L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M., Facer,
B., Ho, A., et al. (2020). The Allen Mouse Brain Common Coordinate Framework: A 3D
Reference Atlas. Cell *181*, 936-953.e20.

Williams, R.H., Vazquez-DeRose, J., Thomas, A.M., Piquet, J., Cauli, B., and Kilduff, T.S.
(2017). Cortical nNOS/NK1 Receptor Neurons are Regulated by Cholinergic Projections From
the Basal Forebrain. Cerebral Cortex (New York, NY : 1991) 1–21.

Xiong, B., Li, A., Lou, Y., Chen, S., Long, B., Peng, J., Yang, Z., Xu, T., Yang, X., Li, X., et al.
(2017). Precise Cerebral Vascular Atlas in Stereotaxic Coordinates of Whole Mouse Brain.
Frontiers in Neuroanatomy *11*, 128.

Zhang, Q., Roche, M., Gheres, K.W., Chaigneau, E., Kedarasetti, R.T., Haselden, W.D.,
Charpak, S., and Drew, P.J. (2019a). Cerebral oxygenation during locomotion is modulated by
respiration. Nature Communications *10*, 1–15.

Zhang, X., Yin, X., Zhang, J., Li, A., Gong, H., Luo, Q., Zhang, H., Gao, Z., and Jiang, H.
(2019b). High-resolution mapping of brain vasculature and its impairment in the hippocampus of
Alzheimer's disease mice. Natl Sci Rev *6*, 1223–1238.

Zhao, Z., Nelson, A.R., Betsholtz, C., and Zlokovic, B.V. (2015). Establishment and Dysfunction
of the Blood-Brain Barrier. Cell *163*, 1064–1078.

1114