A community-science approach identifies genetic variants associated with three color morphs in ball pythons (*Python regius*)

Autumn R. Brown[1], Kaylee Comai[1], Dominic Mannino[1], Haily McCullough[1], Yamini Donekal[1], Hunter C. Meyers[1], The BIO306W Consortium[^1], Chiron W. Graves[1*], and Hannah S. Seidel[1*]

[1]Department of Biology, Eastern Michigan University, Ypsilanti, MI, USA

* Corresponding author:

E-mail: hseidel@emich.edu (HSS), cgraves6@emich.edu (CWG)

[^]The BIO306W Consortium comprises students in an undergraduate laboratory course at Eastern Michigan University. Students are listed in the Acknowledgments.

Abstract

Color morphs in ball pythons (*Python regius*) provide a unique and largely untapped resource for understanding the genetics of coloration in reptiles. Here we use a community-science approach to investigate the genetics of three color morphs affecting production of the pigment melanin. These morphs—Albino, Lavender Albino, and Ultramel—show a loss of melanin in the skin and eyes, ranging from severe (Albino) to moderate (Lavender Albino) to mild (Ultramel). To identify genetic variants causing each morph, we recruited shed skins of pet ball pythons via social media, extracted DNA from the skins, and searched for putative loss-of-function variants in homologs of genes controlling melanin production in other vertebrates. We report that the Albino morph is associated with missense and non-coding variants in the gene *TYR*. The Lavender Albino morph is associated with a deletion in the gene *OCA2*. The Ultramel morph is associated with a missense variant and a putative deletion in the gene *TYRP1*. Our study is one of the first to identify genetic variants associated with color morphs in ball pythons and shows that pet samples recruited from the community can provide a resource for genetic studies in this species.

Data availability: All relevant data are within the paper and its Supporting Information files.

DNA sequences are available from GenBank, under accession numbers MZ269492-MZ269502.

Funding information: This work was supported by a Faculty Research Fellowship and James

H. Brickley Award from Eastern Michigan University to HSS and an Undergraduate Research

Stimulus Program Award and a Don Brown and Meta Hellwig Undergraduate Research Award

from Eastern Michigan University to ARB. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Color patterns are distinctive and beautiful features of many animal species. Among the many functions of color are to camouflage animals to their surroundings, protect tissues from ultraviolet radiation, warn predators of poisons, and provide signals for mating and social communication [1]. Color patterns have also been targeted by artificial selection to create novel and fanciful color patterns in domestic animals [2].

Colors are produced through a combination of chemical pigments and physical structures (structural colors). Pigments absorb light, whereas structural colors are created when light is reflected by the nanoscale geometry of a tissue. Common pigments include brown-to-black melanin, present throughout the animal kingdom, and yellow-to-red carotenoids and pteridines, common in birds, reptiles, and lower vertebrates [3]. Color-producing structures include ordered keratin matrices in bird feathers [4], chitin layers in butterfly wings [5], and purine crystals in the skin of fish, amphibians, and reptiles [6,7]. These color-producing structures can produce a variety of colors, including iridescent colors, depending on the wavelengths of light they reflect [8].

The genetics and development of color patterns in vertebrates have been studied most extensively in mammals. Mammals largely rely on a single type of pigment (melanin), produced in the skin by a single type of cell (melanocyte) [9]. This system provides mammals with skin and hair colors ranging from black to reddish-brown, depending on the chemical structure of the melanin [10].

Skin colors outside mammals are more diverse and include bright colors of all hues. This increased complexity is produced through a combination of structural colors, melanin pigments, and non-melanin pigments, and it relies on multiple types of specialized color-producing cells in the skin [11–15]. Color cell development has been characterized to some extent in fish [16–18], and genes controlling the use of non-melanin pigments have been identified in a few species of

fish and birds [19–25]. The genetics and development of non-mammalian color patterns are less

well understood in other vertebrates, particularly in reptiles [although see 26–29].

A unique resource for understanding the genetics of color patterns in reptiles is the ball

python (*P. regius*). Ball pythons are native to sub-Saharan Africa, but have become common as

pets in the United States [30]. Wild ball pythons exhibit a mottled color pattern, consisting of

brown-to-black melanin and red-to-yellow (non-melanin) pigments in the skin (Fig 1). Captive-

bred ball pythons, by contrast, include many variants of the normal color pattern [26,31–33].

These variants, referred to as 'color morphs', include animals having reduced melanin (e.g.

Albino), increased melanin (e.g. Cinnamon), reduced red-to-yellow coloration (e.g. Axanthic), or

complex changes in the placement of color patches on the skin (e.g. Spider, Clown, Genetic

Stripe, and Enchi). Many color morphs are heritable and show simple dominant or recessive

patterns of inheritance. These inheritance patterns are consistent with single-gene causality, but

only a single genetic variant associated with a ball python color morph (Piebald) has been

identified to date [26]. Ball pythons therefore represent a tractable yet largely untapped resource

for understanding the genetics of coloration in reptiles. An additional feature of ball pythons

convenient for genetic studies is that DNA samples can be obtained non-invasively from shed

skin [34].

The goal of the current study was to perform proof-of-concept experiments to identify the

genetic causes of color morphs in ball pythons, using shed skins of pet ball pythons recruited

via social media. We focused on three morphs for which candidate genes could be readily

identified: Albino, Lavender Albino, and Ultramel. These morphs show a loss of brown-to-black

coloration in the skin and eyes, characteristic of a defect in melanin production (Fig 1). These

morphs are recessive and non-allelic (i.e. crosses between these morphs yield offspring with

normal coloration). Their loss of melanin ranges from severe (Albino) to moderate (Lavender

Albino) to mild (Ultramel). This range of phenotypes mirrors the range of phenotypes observed

for loss-of-function variants in genes required for melanin production in other vertebrates (Table

1). This similarity provided a list of candidate genes that we predicted might harbor loss-of-

function variants causing the Albino, Lavender Albino, and Ultramel color morphs in ball

pythons. Our study demonstrates the feasibility of using community-sourced samples for genetic

studies in ball pythons and lays the groundwork for future investigations of reptile-specific color

patterns in this species.

Fig 1. The Albino, Lavender Albino, and Ultramel color morphs have reduced brown-to-

black coloration, characteristic of a loss of melanin. Red-to-yellow coloration is unaffected

in these morphs. Hypothesized causative genes represent genes in which loss-of-function

variants in other vertebrates produce similar phenotypes (Table 1). (A) Wildtype. (B) Ultramel.

(C) Lavender Albino. (D-F) Albino. (D) Albino animal described as an Alb^{Albino} homozygote. (E)

Albino animal described as an Alb^{Candy} homozygote. (F) Albino animal described as an Alb^{Toffee}

homozygote. Photo credits, Ryan Young of Molecular Reptile, Chiron Graves, Phil Barclay,

Michael Freedman of The Florida Reptile Ranch.

103 **Table 1. Loss-of-function phenotypes of melanogenesis genes across vertebrates.**

Gene	Species with identified loss-of-function variants		Role of protein	Loss-of-function phenotype
TYR	Fish	Japanese carp [35], Japanese rice fish [36]	Rate-limiting enzyme in melanin synthesis pathway	Severe loss of melanin
	Frogs and snakes	Japanese wild frogs (three species) [37], Japanese rat snake [38]		
	Birds	Chicken [39–41]		
	Ungulates	Water buffalo [42], Cattle [43], Red deer [44], Asinara donkey [45]		
	Rodents	House mouse [46], Wistar rat [47], Domestic guinea pig [48]		
	Primates	Crab-eating macaque [49], Capuchin monkey [50], Hamadryas baboon [51], Human [reviewed in 52]		
	Other mammals	Humpback whale [53], Ferret [54], American mink [55], Domestic cat [56–58], Silver fox [59]		
OCA2	Fish	Mexican cavefish [60], Lake Malawi cichlid [61]	Cl⁻ channel regulating melanosome pH	Moderate loss of melanin
	Snakes	Corn snake [62]		
	Mammals	Domestic dog [63], Bama miniature pig [64], House mouse [65], Human [reviewed in 52]		
TYRP1	Fish	Zebrafish [66]	Enzymes contributing to melanin synthesis	Mild loss of melanin
	Birds	Saker falcon [67]		
	Mammals	Chinese indigenous pig [68], Liangshan pig [69], Domestic cat [58,70], Soay sheep [71], Valais Red sheep [72], American mink [73], Human [reviewed in 52]		
TYRP2	Mammals	House mouse [74,75]		
SLC7A11	Mammals	House mouse [76]	Solute transporters	
SLC24A5	Fish	Zebrafish [77]		
	Mammals	Domestic dog [78], Horse [79], Human [reviewed in 52]		
SLC45A2	Fish	Japanese rice fish [80]		
	Birds	Chicken and Japanese quail [81]		
	Mammals	Horse [82], House mouse [83], Bengal tiger [84], Western lowland gorilla [85], Human [reviewed in 52]		

Results

Obtaining a reference sequence for melanogenesis genes in a wildtype ball python

Genes required for melanin production have been identified in humans and other

vertebrates (Table 1). These genes encode enzymes that synthesize melanin (*TYR*, *TYRP1*,

TYRP2) [86], a chloride channel required for maintaining the pH of melanosomes (*OCA2*) [87],

and transporters thought to import solutes into the cell or into organelles (*SLC7A11*, *SLC24A5*,

and *SLC45A2*) [76,88,89]. These genes are highly conserved among vertebrates and occur in

single copy in most vertebrate genomes. Loss-of-function variants in these genes in humans

cause a genetic disorder known as oculocutaneous albinism, which is characterized by loss of

melanin in the skin, hair, and eyes [52]. This loss of melanin ranges from severe to mild,

depending on the causative gene and genetic variants therein [52]. Similar phenotypes occur in

other animals, where the loss of melanin extends to feathers and scales (Table 1).

To obtain a reference sequence for melanogenesis genes in ball python, we amplified

and sequenced the coding regions of *TYR*, *TYRP1*, *TYRP2*, *OCA2*, *SLC7A11*, *SLC24A5*, and

SLC45A2 from a single ball python having normal coloration (henceforth 'wildtype'). Primers for

amplification were designed against the genome of Burmese python (*Python bivittatus*), the

closest relative of ball python for which genome sequence was available [90]. Comparison of

ball python sequences to sequences from Burmese python revealed 97.2-99.2% nucleotide

identity within coding regions and 96.9-98.5% nucleotide identity within flanking non-coding

regions (Table 2). This analysis provided a reference sequence for genes in ball python that

might harbor loss-of-function variants causing the Albino, Lavender Albino, and Ultramel color

morphs.

Table 2. Comparison of ball python and Burmese python genomic and protein sequences
for melanogenesis genes.

Region	Gene	Total nucleotide sequence aligned (bp)	Nucleotide identity (%)	Amino acid identity (%)
Coding	*TYR*	1,591	98.5	99.6
	TYRP1	1,584	99.2	99.6
	TYRP2	1,473	98.3	99.1
	OCA2	2,577	98.9	99.7
	SLC7A11	1,506	98.7	99.7
	SLC24A5	1,379	98.2	99.5
	SLC45A2	1,557	97.2	98.9
Non-coding	*TYR*	577	96.9	(not applicable)
	TYRP1	1,160	97.0	
	TYRP2	1,060	97.9	
	OCA2	2,154	98.5	
	SLC7A11	3,407	97.9	
	SLC24A5	1,335	97.9	
	SLC45A2	999	97.0	

The Albino color morph is associated with three haplotypes of *TYR*

The Albino color morph in ball pythons is characterized by an absence or near absence of melanin in the skin and eyes. This phenotype resembles the phenotype caused by loss of function of *TYR* in other vertebrates. *TYR* encodes the enzyme catalyzing the rate-limiting step of melanin production [86]. Loss of this enzymatic activity in other vertebrates causes a severe loss of melanin, which is typically more severe than the loss of melanin caused by loss of function of other melanogenesis genes (Table 1). We therefore hypothesized that the Albino color morph in ball pythons was caused by loss-of-function variants in *TYR*.

To test whether the Albino color morph was associated with variants in *TYR*, we performed a small-scale association study using 50 Albinos and 56 Non-Albinos, recruited from a total of 18 states in the United States. Polymorphic sites were identified through pilot sequencing of a subset of animals (see Methods). Six polymorphic sites were selected for the association study, two near the 5' end of *TYR* and four near the 3' end (Fig 2A). Each animal was genotyped at each site, and haplotypes were reconstructed using PHASE [91,92]. Association between haplotype and color morph was tested using the case-control test of PHASE. As a negative control, we also tested for association between haplotype and color morph at *TYRP1*, *TYRP2*, and *OCA2*, using two polymorphic sites per gene (S1 Table). This analysis revealed a significant association between haplotype and color morph at *TYR* (p = 0.04, Bonferroni corrected), but no association for the other three genes (p > 0.05, Bonferroni corrected). The association for TYR was driven by two features of the haplotype distribution: (i) diversity of *TYR* haplotypes was reduced among Albinos compared to Non-Albinos (Fig 2A), and (ii) two of the three *TYR* haplotypes found in Albinos were rare among Non-Albinos (Fig 2A). These results demonstrate an association between the Albino color morph and variants in

TYR; furthermore, these results show that all Albino animals were homozygous or compound

heterozygous for any of three haplotypes of *TYR*.

Fig 2. The Albino color morph is associated with three haplotypes of *TYR*. (A) *TYR*

haplotype frequencies among Albinos and Non-Albinos. (B) Alignment of *TYR* protein

sequences surrounding the missense variants *D394G* and *P384L*. (C) *TYR* genotypes and

breeder designations of animals used in the association study. **+**, any *TYR* haplotype found

exclusively among Non-Albinos. (D) PCR amplification of a genomic fragment containing the

variable number tandem repeat (VNTR) shown in E. Non-Albinos are examples of Non-Albinos

homozygous for the longer allele of the VNTR. (E) Schematic of the three *TYR* haplotypes

found in Albinos. The TYR^{Albino} haplotype contains no coding variants and no splice-site variants

compared to wildtype. (A, E) Hash mark, discontinuity in the Burmese python reference

genome.

We hypothesized that each of the three haplotypes of *TYR* found among Albinos might

carry a distinct loss-of-function variant in the gene. To search for such variants, we selected one

Albino animal homozygous for each haplotype and sequenced the *TYR* coding regions and

adjacent splice sites in these animals. We found that one of these animals was homozygous for

an aspartic acid-to-glycine variant (*D394G*) in the third coding region of *TYR*. A second animal

was homozygous for a proline-to-leucine variant (*P384L*), also in the third coding region of *TYR*.

Both variants alter conserved residues (Fig 2B), and the *P384L* variant occurs at the same site

as a similar variant (*P384A*) associated with oculocutaneous albinism in humans [93]. The third

animal carried no coding variants and no splice-site variants compared to wildtype. These

results show that two of the three *TYR* haplotypes found among Albinos carried missense

variants that are likely disruptive for TYR protein function. We term these *TYR* haplotypes

TYR^{D394G} and TYR^{P384L}. We term the third Albino haplotype, which lacked coding or splice-site

variants compared to wildtype, TYR^{Albino}.

We hypothesized that *D394G*, *P384L*, and an unidentified variant on the *TYR^Albino^*

haplotype were causative for the Albino color morph. This hypothesis predicts that animals

homozygous or compound heterozygous for these variants will be Albino. Heterozygotes and

non-carriers are predicted to be Non-Albino. To test this prediction, we genotyped our full panel

of 50 Albinos and 56 Non-Albinos for the missense variants *D394G* and *P384L*. Consistent with

our prediction, we found that animals homozygous or compound heterozygous for *D394G*,

P384L, or the *TYR^Albino^* haplotype were exclusively Albino (Fig 2C). Heterozygotes and non-

carriers were exclusively Non-Albino (Fig 2C). The most common haplotype among Albinos was

*TYR^Albino^* (haplotype frequency of 71%), followed by *TYR^P384L^* (16%) and *TYR^D394G^* (13%). We

conclude that *D394G*, *P384L*, and an unidentified variant on the *TYR^Albino^* haplotype are likely

causative for the Albino color morph. Any combination of these variants produces the Albino

phenotype.

The *TYR^Albino^* haplotype lacks an obvious loss-of-function variant

The lack of coding or splice-site variants on the *TYR^Albino^* haplotype led us to hypothesize

that this haplotype carried a loss-of-function variant in a non-coding region. To search for such

variants, we examined the *TYR* promoter. We sequenced ~2 kb immediately upstream of the

TYR start codon in a *TYR^Albino^* homozygote and compared this sequence to wildtype. We found

that the *TYR^Albino^* haplotype differed from wildtype at a total three sites (three single-base

substitutions). In all three cases, the *TYR^Albino^* haplotype was homozygous for the allele shared

with Burmese python (i.e. the ancestral allele). This result suggests that these variants are not

causative for the Albino color morph. We conclude that the causative variant in the *TYR^Albino^*

haplotype does not reside within the sequenced region, ~2 kb upstream of the *TYR* start codon.

We hypothesized that the TYR^{Albino} haplotype might contain a large insertion or deletion

(indel) in an intron that disrupts gene function. Intronic indels can disrupt slicing and have been

found to disrupt the function of melanogenesis genes in other species [e.g. 62,73]. To search for

large intronic indels, we tiled PCR amplicons across TYR introns and compared amplicon sizes

between a TYR^{Albino} homozygote and a wildtype animal. These amplicons tiled across a total of

~38 kb of intronic sequence, which included part of intron 1 and all of introns 2, 3, and 4 (Fig

2E). (Tiling across intron 1 was incomplete because this intron contains a discontinuity in the

Burmese python reference genome. Amplification across this discontinuity was unsuccessful in

ball python, even in wildtype.) Size differences between the TYR^{Albino} homozygote and wildtype

were assessed via standard agarose gel electrophoresis, which we estimate was sensitive

enough to reveal indels larger than ~50-100 bp, depending on amplicon size (which ranged

from ~0.7-3.5 kb). This analysis revealed one size difference (Fig 2D). Further sequencing

revealed that this size difference was caused by a variable number tandem repeat (VNTR) that

was ~400 bp larger in the TYR^{Albino} homozygote than in wildtype. Genotyping of the 56 Non-

Albinos for this indel revealed that eight of these animals were homozygous for the larger allele

of the VNTR (Fig 2D). This finding suggests that the larger allele of the VNTR is not causative

for the Albino color morph. We propose that the TYR^{Albino} haplotype harbors a loss-of-function

variant other than a large intronic indel or outside the genomic regions analyzed here.

TYR^{P384L} is associated with reduced phenotype severity

compared to TYR^{D394G} and TYR^{Albino}

Ball python breeders describe the Albino color morph as having three alleles: Alb^{Albino},

Alb^{Candy}, and Alb^{Toffee}. Evidence for this view comes in part from variation in the coloration of

Albino animals. In some Albino animals, the brown-to-black coloration observed in wildtype is

entirely absent, and skin patches appear white (Fig 1D). These Albinos are typically considered

to be Alb^{Albino} homozygotes. Other Albinos show a less severe phenotype, where skin patches

are faintly beige or lavender instead of white (Figs 1E and 1F). These Albinos are typically

considered to carry one or more copies of Alb^{Candy} or Alb^{Toffee}.

We hypothesized that the Albino alleles recognized by breeders might correspond to the

TYR haplotypes identified through sequencing (TYR^{D394G}, TYR^{P384L}, and TYR^{Albino}). To test this

hypothesis, we examined the breeder designations of the 50 Albinos in our panel. We found that

the Alb^{Candy} or Alb^{Toffee} designations typically corresponded the TYR^{P384L} haplotype (Fig 2C). The

Alb^{Albino} designation typically corresponded to the other two haplotypes (TYR^{D394G} and TYR^{Albino},

Fig 2C). This correspondence was imperfect, and exceptions existed (e.g. three animals

designated as Alb^{Albino} / Alb^{Candy} compound heterozygotes did not carry the TYR^{P384L} haplotype,

Fig 2C). We conclude that the Alb^{Candy} and Alb^{Toffee} designations typically (but not exclusively)

represent the TYR^{P384L} haplotype. The Alb^{Albino} designation typically (but not exclusively)

represents the TYR^{D394G} or TYR^{Albino} haplotype. The association between the Alb^{Candy} and

Alb^{Toffee} designations and the TYR^{P384L} haplotype suggests that this haplotype may confer a

slightly less severe phenotype than do TYR^{D394G} and TYR^{Albino}.

242 The Lavender Albino color morph is associated with a
243 deletion in *OCA2*

The Lavender Albino color morph is characterized by skin patches that are lavender

instead of brown or black. This phenotype is thought to arise from melanin levels that are

dramatically reduced but not entirely eliminated. Similar phenotypes have been observed in

other vertebrates upon loss of function of *OCA2* (Table 1). *OCA2* encodes a chloride channel

required for maintaining the pH of melanosomes [87,94]. When the OCA2 protein is absent or

non-functional, the enzymes that synthesize melanin are less active, and only small amounts of

melanin are produced [95,96]. The resulting phenotype is typically less severe than the loss-of-

function phenotype of *TYR*, but more severe than the loss-of-function phenotypes of other

melanogenesis genes (Table 1). We therefore hypothesized that the Lavender Albino color

morph was caused by loss-of-function variants in *OCA2*.

To search for loss-of-function variants in *OCA2*, we selected a single Lavender Albino

animal and sequenced 23 of the 24 coding regions of *OCA2* in this animal. Comparison of these

sequences to wildtype revealed no coding variants and no splice-site variants. We attempted to

repeat this analysis for the remaining coding region (coding region 18), but we were unable to

amplify this coding region from the Lavender Albino animal. Further test amplifications revealed

that the Lavender Albino animal was homozygous for a 1,514-bp deletion spanning coding

region 18 (Figs 3A and 3B). This deletion removes 36 amino acids from the protein and likely

introduces a frameshift into the transcript, given that the coding regions downstream of the

deletion are out of frame compared to coding regions upstream of the deletion (Fig 3E). This

frameshifted transcript is predicted to produce a protein lacking six of the 12 transmembrane

helices present in wildtype OCA2. Truncations occurring at similar positions in the OCA2 protein

have been identified in humans and are associated with oculocutaneous albinism [97]. We

conclude that the deletion of *OCA2* coding region 18 likely disrupts protein function and is a

strong candidate for the cause of the Lavender Albino color morph.

Fig 3. The Lavender Albino color morph is associated with a deletion in *OCA2*. (A)

Schematic of the *OCA2* gene. (B) PCR amplification demonstrating the deletion in Lavender

Albino. (C) PCR amplification used for genotyping the deletion in the animals in panel D. (D)

Genotypes of 14 Lavender Albinos, 76 Non-Lavender Albinos, and one animal described as

heterozygous for Lavender Albino. This set of animals includes the original Lavender Albino

animal in which the OCA2 deletion was identified. (E) Schematic of the *OCA2* haplotype found

in Lavender Albinos. (A, E) Hash mark, discontinuity in the Burmese python reference genome.

The Lavender Albino color morph is considered by breeders to have a single allele. We

therefore predicted that the *OCA2* deletion would be shared by other Lavender Albino animals.

We predicted that Non-Lavender Albinos would be heterozygous for the deletion or non-carriers.

To test this prediction, we genotyped the *OCA2* deletion in 13 additional Lavender Albino

animals. We also genotyped 76 Non-Lavender Albinos and one animal described as

heterozygous for the Lavender Albino color morph. We found that all 13 Lavender Albinos were

homozygous for the deletion (Figs 3C and 3D). All 76 Non-Lavender Albinos were non-carriers.

The animal reported to be heterozygous for Lavender Albino was heterozygous for the deletion.

These findings support the conclusion that the *OCA2* deletion is causative for the Lavender

Albino color morph (Fig 3E).

The Ultramel color morph is associated with a missense variant and a putative deletion in *TYRP1*

The Ultramel color morph is characterized by skin patches that are tan or light brown,

rather than dark brown or black. This phenotype suggests a mild loss of melanin. Similar

phenotypes in other vertebrates have been observed upon loss of function of any five genes:

TYRP1, *TYRP2*, *SLC7A11*, *SLC24A5*, and *SLC45A2* (Table 1). *TYRP1* and *TYRP2* encode

enzymes involved in synthesizing melanin [86]. Their enzymatic activities are partially reductant

with other enzymes in the melanin synthesis pathway, thus explaining their milder loss-of-

function phenotypes compared to *TYR*. *SLC7A11* encodes a transporter responsible for

importing cystine into the cell [98]. Cystine is a precursor to some forms of melanin; hence

absence of *SLC7A11* alters melanin levels [76]. *SLC24A5* and *SLC45A2* encode a K^+-

dependent Na^+-Ca^{2+} exchanger [88] and a putative sugar transporter [89], respectively. Loss of

either of these proteins causes a reduction in melanin, through mechanisms that may involve

defects in regulation of melanosome pH [83,88,99]. We hypothesized that the Ultramel color

morph was caused by loss-of-function variants in one of these genes.

To search for loss-of-function variants in *TYRP1*, *TYRP2*, *SLC7A11*, *SLC24A5*, and

SLC45A2, we selected a single Ultramel animal and sequenced the coding regions of each of

these genes in this animal. Comparison of these sequences to wildtype revealed a single coding

variant: an arginine-to-histidine substitution (*R305H*) in coding region 4 of *TYRP1*. The Ultramel

animal was homozygous for this variant. The arginine residue at this site is conserved across

vertebrates (Fig 4A) and is also conserved in *TYR*, which is paralogous to *TYRP1* [100]. An

arginine-to-histidine substitution at the homologous site in *TYR* has been reported in humans

and is associated with oculocutaneous albinism [101–103]; thus, histidine at this site is likely

disruptive to protein function. The *R305H* variant is therefore a good candidate for the cause of

the Ultramel color morph.

Fig 4. The Ultramel color morph is associated with a missense variant and a putative

deletion in *TYRP1*. (A) Alignment of *TYRP1* protein sequence surrounding missense variant

R305H. (B) Genotypes of 11 Ultramels and 78 Non-Ultramels. This set of animals includes the

original Ultramel animal in which the *R305H* variant was identified. Ultramels heterozygous for

R305H are presumed to be heterozygous for the putative deletion of coding regions 6 and 7;

testing these animals for the putative deletion was not possible because the putative deletion

cannot be detected when heterozygous. (C) Top, schematic of the *TYRP1* gene. Bottom left,

PCR amplifications demonstrating the putative deletion in the Ultramels lacking *R305H*. Bottom

right, alignment of PCR amplicons to the *TYRP1* gene. (D) Schematic of the two *TYRP1*

haplotypes found in Ultramels. (A, D) Hash mark, discontinuity in the Burmese python reference

genome.

The Ultramel color morph is considered by breeders to have a single allele. We therefore

predicted that the *R305H* variant would be shared by other Ultramels. We predicted that Non-

Ultramels would be heterozygous or non-carriers. To test this prediction, we genotyped the

R305H variant in 10 additional Ultramels and 78 Non-Ultramels. We found that five of the 10

Ultramels were homozygous for the *R305H* variant (Fig 4B). Of the remaining Ultramels, two

were heterozygous for the *R305H* variant, and three did not carry it. By contrast, none of the 78

Non-Ultramels carried the *R305H* variant. This pattern is consistent with the *R305H* variant

causing the Ultramel phenotype in some Ultramels but not others.

We hypothesized that the *R305H* variant represents one of two loss-of-function alleles of

TYRP1. Under this scenario, Ultramels lacking the *R305H* variant are predicted to be

homozygous for an alternate loss-of-function allele. Ultramels heterozygous for *R305H* are

predicted to be compound heterozygotes. To test this prediction, we amplified and sequenced

coding regions 1 through 5 of *TYRP1* in one of the three Ultramels lacking the *R305H* variant.

We found no coding variants and no splice-site variants compared to wildtype. We attempted to

repeat this analysis for coding regions 6 and 7 of *TYRP1*, but we were unable to amplify these

coding regions from this animal (Fig 4C). Identical results were observed for the other two

Ultramels lacking the *R305H* variant (S5 Table). Further test amplifications indicated that coding

regions 6 and 7 were missing from the genomes of all three animals (Fig 4C, S5 Table). Loss of

these coding regions was specific to Ultramels lacking the *R305H* variant and did not occur in

any of the Ultramels carrying the *R305H* variant, nor in any of the 78 Non-Ultramels (Fig 4B). A

simple explanation of this pattern is that the Ultramels lacking the *R305H* variant were

homozygous for a deletion spanning coding regions 6 and 7 of *TYRP1*. This putative deletion

removes at least 2,208 bp from the genome and 117 amino acids from the TYRP1 protein,

including the second of two zinc-binding domains. (The exact size and breakpoints of the

putative deletion remain undetermined because the deletion extends into a discontinuity in the

Burmese python reference genome.) Truncations occurring at similar positions in the TYRP1

protein have been identified in humans and are associated with oculocutaneous albinism [104].

We conclude that the Ultramel color morph is likely caused by variants in *TYRP1* and has two

alleles: missense variant *R305H* and a putative deletion of coding regions 6 and 7 (Fig 4D).

Discussion

The goal of our study was to use pet samples recruited from the community to identify

the genetic causes of the Albino, Lavender Albino, and Ultramel color morphs. We succeeded in

recruiting 11 or more animals for each morph, along with a larger number of animals not

belonging to these morphs. This sample size, albeit small, was sufficient to identify putatively

causal variants for each morph (Fig 5). The Albino color morph was associated with three

alleles of TYR: missense variant *D394G*, missense variant *P384L*, and haplotype TYR^{Albino},

which lacks coding or splice-site variants compared to wildtype. The Lavender Albino color

morph was associated with a single allele of *OCA2*, a 1,514-bp deletion that removes *OCA2*

coding region 18. The Ultramel color morph was associated with two alleles of *TYRP1*:

missense variant *R305H* and a putative deletion that removes *TYRP1* coding regions 6 and 7.

Due to the small sample size of our study, we cannot exclude the possibility of additional loss-

of-function alleles of these genes segregating in the ball python population. However, such

alleles are expected to be at low frequency, given their absence in our sample. These findings

are consistent with genetic data from other vertebrates (Table 1), indicating that the loss-of-

function phenotypes of these genes range from severe (*TYR*) to moderate (*OCA2*) to mild

(*TYPR1*). Our study demonstrates that pet ball pythons are a tractable resource for genetic

analysis of coloration in reptiles, at least for color morphs having obvious candidate genes.

Fig 5. Summary of genetic variants associated with the Albino, Lavender Albino, and

Ultramel color morphs. Photo credits, Ryan Young of Molecular Reptile, Chiron Graves, Phil

Barclay, Michael Freedman of The Florida Reptile Ranch.

Molecular functions of genetic variants associated with color morphs

The missense variants and deletions found in *TYR*, *TYRP1*, and *OCA2* are likely hypomorphic or null alleles. The deletion in *OCA2* likely results in a frameshifted transcript that, if not degraded by nonsense-mediated decay, encodes a protein lacking six of the 12 transmembrane helices present in wildtype OCA2 [94]. This truncated protein is therefore unlikely to retain the Cl⁻ channel activity observed for wildtype OCA2 [87]. *TYR* and *TYPR1* encode globular enzymes that require copper or zinc as co-factors [86]. Variants *P384L* and *D394G* in *TYR* reside in the second of TYR's two copper-binding domains and may therefore alter the ability of the TYR to bind copper. Variant *R305H* in *TYRP1* occurs at a site that normally forms salt bridges with residues located elsewhere in the peptide chain [86]. An arginine-to-histidine substitution at this site likely disrupts these salt bridges and may therefore interfere with proper protein folding. The putative deletion in *TYRP1* removes the C-terminal ~20% of the protein. This deleted region includes the second of TYRP1's two zinc-binding domains and several alpha helices residing to the enzyme's hydrophobic core [86]. Truncated TYRP1 protein is therefore unlikely to fold properly, nor is it likely to properly associate with zinc.

The TYR^{Albino} haplotype is the most common *TYR* haplotype found among Albinos, but it lacks an obvious loss-of-function variant. This haplotype contains no coding variants, no splice-site variants, no derived variants within 2 kb upstream of the start codon, and no large indels in three of four introns (with the exception of a longer allele of a VNTR, which is not specific to the TYR^{Albino} haplotype). We propose that the TYR^{Albino} haplotype contains a loss-of-function variant not detectable by methods used in the current study. Examples include regulatory variants farther upstream of the start codon, substitutions in introns that disrupt slicing, or large indels or rearrangements involving the first intron. Cryptic loss-of-function variants in melanogenesis

395   genes are thought to be relatively common in humans, where ~10-20% of oculocutaneous

396   albinism patients are heterozygous for pathogenic variants in a known melanogenesis gene but

397   lack a coding or splice-site variant on the opposite allele [105–107].

Multiple alleles of the Albino color morph

Prior to the current study, the Albino color morph was recognized by breeders as having

three alleles: Alb^{Albino}, Alb^{Candy}, and Alb^{Toffee}. Alb^{Albino} was thought to confer a more severe

phenotype than Alb^{Candy} and Alb^{Toffee} (Figs 1D-F). Our results confirm that the Albino color morph

has three molecular alleles (TYR^{D394G}, TYR^{P384L}, and TYR^{Albino}). Yet these alleles do not perfectly

correspond to the alleles recognized by breeders. The molecular allele TYR^{P384L} typically

corresponds to the breeder designations Alb^{Candy} and Alb^{Toffee}. The molecular alleles TYR^{D394G}

and TYR^{Albino} typically correspond to the breeder designation Alb^{Albino} (Fig 2C). This

correspondence is imperfect and exceptions exist (Fig 2C).

We speculate that the TYR^{P384L} haplotype was discovered twice and was named Alb^{Candy}

by one breeder and Alb^{Toffee} by another breeder. We speculate that TYR^{D394G} and TYR^{Albino}

confer similar phenotypes and were not previously recognized by breeders as distinct. The less

severe phenotype associated with TYR^{P384L} may reflect the $P384L$ variant being less disruptive

to gene function compared with variants on the other two alleles. Alternatively, the less severe

phenotype associated with TYR^{P384L} may reflect genetic linkage to modifiers in other genes. In

either case, our study demonstrates that the Albino allele designations current in use by

breeders do not accurately reflect molecular genotype. Renaming of the Albino alleles is

warranted, although owners and breeders may be resistant to renaming due to cultural

attachment to existing allele names.

Multiple alleles of the Ultramel color morph

Our finding of multiple *TYRP1* alleles for the Ultramel color morph was unexpected. This

morph was not previously described as having multiple alleles. We speculate that one of the two

alleles associated with the Ultramel color morph may represent an allele originally associated

with a morph known as Caramel Albino. Multiple lineages of Caramel Albinos have been

described, and their coloration similar to Ultramels. Caramel Albinos have been disfavored

among owners and breeders because of spinal kinking and reduced female fertility. Caramel

Albinos were not included in our study because we were unable to recruit any Caramel Albino

samples. We speculate that the Caramel Albino morph may be allelic with Ultramel, and that

one of two *TYRP1* alleles associated with Ultramel may represent an allele originally described

as Caramel Albino. Alternately, Ultramel may be distinct from Caramel Albino, and the two

TYRP1 alleles may represent two distinct origins of the Ultramel morph.

Prospects for genetic testing

Many breeders of ball pythons wish to identify heterozygotes of recessive color morphs.

Currently the only tool for identifying heterozygotes is test breeding, which is slow and laborious.

The results of the current study will enable simple genetic testing for Albino, Lavender Albino,

and Ultramel, which will allow heterozygotes to be identified prior to reproductive maturity.

Testing for Lavender Albino can be performed by genotyping the *OCA2* deletion. Testing for

Albino can be performed by sequencing the *TYR* missense variants *D394G* and *P384L*, and by

genotyping variants that distinguish the TYR^{Albino} haplotype from other *TYR* haplotypes.

Detection of *D394G* or *P384L* can be considered diagnostic for Albino because these variants

are likely causative for the Albino color morph. Detection of the TYR^{Albino} haplotype is less

diagnostic because the causative variant on TYR^{Albino} haplotype remains unknown. Testing for

one of the two Ultramel alleles can be performed by sequencing the *TYRP1* missense variant

R305H. The other Ultramel allele—the putative deletion of *TYRP1* coding regions 6 and 7—

cannot currently be detected in heterozygotes using simple methods because the breakpoints of

this putative deletion are unknown.

Use of Burmese python reference genome

One challenge for genetic studies in ball python is the absence of a reference genome.

To fill this gap, we relied on the genome of Burmese python [90]. This genome is a scaffold-

level assembly, and some genes are fragmented across scaffolds. This fragmentation limited

our analysis of the *TYR^Albino^* haplotype and the putative deletion in *TYRP1*. For *TYR^Albino^*, a

discontinuity in the first intron of *TYR* in the Burmese python genome prevented us from

assessing whether the *TYR^Albino^* haplotype contained a large indel or rearrangement in this

intron. For the putative deletion in *TYRP1*, a discontinuity immediately downstream of *TYRP1*

prevented us from mapping the exact breakpoints of the deletion. As genomic resources for

Burmese python and ball python expand, we expect to identify a putatively causative variant on

the *TYR^Albino^* haplotype and to map the breakpoints of the putative deletion in *TYRP1.* This

information will increase the prospects for genetic testing for these alleles.

Ball pythons as a genetic system

The market for pet ball pythons is huge, and many owners prefer pets with novel color

patterns. This demand has led breeders to propagate genetic variants affecting coloration

[31,33]. This effort has fortuitously created a collection of "mutants" useful for understanding the

genetics of coloration in reptiles. Examples include ball python morphs with altered red-to-yellow

coloration (e.g. Axanthic) and morphs in which the normal mottled color pattern is converted to

dorsal strips (e.g. Clown, Genetic Stripe, Super Stripe). Red-to-yellow pigments are largely

uncharacterized in reptiles [although see 28,29], and morphs affecting these pigments may

provide insight into the metabolism and storage of these pigments. Morphs with stripes are

reminiscent of evolutionary changes in color patterning across snake species [108] and may

provide insights into the developmental mechanisms by which these changes evolve. Future

studies of ball python color morphs will be aided by our groundwork showing that ball python

samples can be recruited effectively from pet owners, and that genetic analyses in ball python

can be scaffolded using the genome of Burmese python as a reference. Similar groundwork has

also been provided by a recent study characterizing the ball python morph known as Piebald

[26]. We expect that a continued community-science approach will be effective in developing

ball pythons into a resource for understanding the genetics of reptile coloration.

Materials and methods

Recruitment of ball python sheds

Ball python sheds were recruited from pet owners and breeders by placing

announcements in Twitter, Reddit, Instagram, and Facebook, and by contacting sellers having

active listings on Morph Market (www.morphmarket.com). Contributors were instructed to allow

sheds to dry (if wet) and to ship sheds via standard first-class mail. Contributors sending

multiple sheds were instructed to package sheds individually in plastic bags during shipping.

Contributors were not provided monetary compensation for sheds, although some contributors

were given pre-paid shipping envelopes to cover shipping costs. Contributors were thanked via

social media whenever possible. Upon receipt, sheds were stored at -20°C to kill any insect

larvae infesting the sheds.

To maximize genetic diversity within each category of morph, we limited our sample of

animals of any one morph to animals contributed by different contributors. Exceptions were

made if animals contributed by the same contributor had been obtained from different breeders.

The goal of this strategy was to reduce the number of animals that were close relatives (e.g.

siblings or parent-offspring pairs). Obtaining full pedigree information was not possible because

most owners lacked this information (e.g. "I bought my animal at a pet store" or "I got my animal

from a friend who was moving away"). The idea that animals in our sample were derived from

multiple lineages is supported by our discovery of multiple alleles for the Albino and Ultramel

color morphs supports. However, we cannot exclude the possibility that some animals in our

sample may have been close relatives.

The total set of animals comprised 50 Albinos, 14 Lavender Albinos, 11 Ultramels, one

animal described as heterozygous for Lavender Albino, and 46 animals described as having

normal coloration (i.e. wildtype) or belonging to color morphs other than Albino, Lavender

Albino, or Ultramel. Phenotypes of Albinos, Lavender Albinos, and Ultramels were confirmed by

examining shed skins for a reduction of brown-to-black coloration. The single animal described

as heterozygous for Lavender Albino was included to demonstrate the three-primer assay for

genotyping the OCA2 deletion. Heterozygotes for Albino or Ultramel and additional

heterozygous for Lavender Albino were excluded from our study because we have found that

animals described as heterozygous for recessive traits by breeders and owners are sometimes

not heterozygous for these traits (e.g. due to mis-attribution of paternity).

Performing experiments in an undergraduate laboratory course

The majority of experiments and analyses described in this study were performed by

undergraduate students as part of a laboratory course at Eastern Michigan University

(BIO306W). This practice required that our experimental design rely on simple techniques,

namely PCR and Sanger sequencing. To avoid student errors in these techniques, we

implemented the following precautions. First, students never extracted DNA from more than one

animal within the same laboratory period. Second, students performed negative and positive

control reactions for all PCR amplifications. Data from students having incorrect controls were

excluded from analysis. Third, all sequence analyses were performed independently by three or

more students. When the results of all students did not all agree, sequences were re-analyzed

by the instructor (HSS).

Annotation of melanogenesis genes in Burmese python

Our analyses in ball python used the genome of Burmese python as a reference. We

therefore required high-confidence gene annotations in Burmese python for the genes used in

our study (*TYR*, *TYRP1*, *TYRP2*, *OCA2*, *SLC7A11*, *SLC24A5*, and *SLC45A2*). Preliminary

inspection of existing gene annotations in Burmese python suggested that many gene

annotations contained errors (e.g. missing one or more coding regions). A main cause of these

errors was that the Burmese python genome (*Python_molurus_bivittatus-5.0.2*) is a scaffold-

level assembly, and many genes were split across scaffolds. We therefore curated new gene

annotations in Burmese python, using conservation of gene structure across species. Alignment

of protein sequences for each gene from corn snake, anole lizard, chicken, and/or mouse

revealed that gene structure was highly conserved for all seven genes: Genes in each species

contained the same number of coding regions, and the coding-region boundaries relative to

protein sequences were perfectly conserved across species, with the exception of one slight

difference in mouse for one boundary of *SLC45A2* (S2 Table). We therefore felt confident that

conservation of coding-region boundaries could be used to curate new gene annotations in

Burmese python. While it remains theoretically possible that our new gene annotations contain

minor errors, our confidence in these gene annotations is high, given the high conservation of

gene structure across species.

To curate new gene annotations, we performed TBLASTN searches of the Burmese

python genome (*Python_molurus_bivittatus-5.0.2*) using protein sequences from anole lizard or

corn snake as the query. Hits were examined manually. Gene annotations were built to match

the coding-region boundaries conserved across species. Details for each gene annotation are

given below.

TYR

The Burmese python genome was queried using TYR protein sequence from anole

lizard (*XP_003219419.1*). The N-terminus of this query (coding region 1) hit Burmese python

transcript *XM_007438960.1* on scaffold 4418. The C-terminus of this query (coding regions 2 to

5) hit Burmese python transcript *XM_007436041.2* on scaffold 3103. TYR was annotated as the

union of these transcripts. The 3' boundary of coding region 1 was adjusted to match the

boundary conserved across species (…TTC TCT TCA TGG CAA-3').

TYRP1

The Burmese python genome was queried using TYRP1 protein sequence from corn

snake (*XP_034266320.1*). The N-terminus of this query (coding regions 1 to 5) hit Burmese

python transcript *XM_007426971.3* on scaffold 801. The C-terminus of this query (coding

regions 6 and 7) hit unannotated regions on this same scaffold. *TYRP1* was annotated as the

union of transcript *XM_007426971.3* and coding regions 6 and 7. The boundaries of coding

regions 6 and 7 were built to match the boundaries conserved across species (coding region 6,

5'-ATA TCT CAA CAT ACC....AAG TTC AGT GGC CAT-3'; coding region 7, 5'-CAC AAG CTC

TCC ATG....CAG TCA GAT GTG TGA-3').

TYRP2

The Burmese python genome was queried using TYRP2 protein sequence from corn

snake (*XP_034273310.1*). The N-terminus of this query (coding regions 1 to 3) hit unannotated

regions on Burmese python scaffold 4970. The C-terminus of this query (coding regions 4 to 8)

hit Burmese python transcript *XM_025174219.1* on this same scaffold. *TYRP2* was annotated

as the union of coding regions 1 to 3 and C-terminal five coding regions of transcript

XM_025174219.1. Transcript *XM_025174219.1* contains a sixth coding region that does not

match *TYRP2* and was therefore excluded from the gene annotation. The boundaries of coding

regions 1 to 3 were built to match the boundaries conserved across species (coding region 1, 5'-

ATG GCC TTC CTG CTG....GTT GCC AAT GCA CAG-3'; coding region 2, 5'-GAC ATT TTG

CTG GCT....GAG ATA CTC TAT TAG-3'; coding region 3, 5'- GAC CAG GCC GTC CCT....GAA

AGA GAT CTG CAG-3').

OCA2

The Burmese python genome was queried using OCA2 protein sequence from corn

snake (*XP_034287267.1*). The N-terminus of this query (coding regions 1 to 9) hit Burmese

python transcript *XM_025173964.1* on scaffold 4704. The C-terminus of this query (coding

regions 10 to 24) hit Burmese python transcript *XM_007433276.1* on scaffold 2194. *OCA2* was

annotated as the union of these transcripts. The 5' boundary of coding region 10 was adjusted

to match the boundary conserved across species (5'-ATT GTC CAC AGG ACA…).

SLC7A11

575   The Burmese python genome was queried using SLC7A11 protein sequence from corn

576   snake (*XP_034257397.1*). All regions of the query hit Burmese python transcript

577   *XM_007430906.3* on scaffold 1543. *SLC7A11* was annotated as transcript *XM_007430906.3*,

578   with no further adjustments.

SLC24A5

The Burmese python genome was queried using SLC24A5 protein sequence from corn

snake (*XP_034290605.1*) extended at its N-terminus to an upstream start codon present in the

parent transcript (*XM_034434714.1*). The N-terminus of the query (coding region 1) hit an

unannotated region on Burmese python scaffold 3984. The C-terminus of this query (coding

regions 2 to 9) hit Burmese python transcript *XM_007438007.2* on this same scaffold. *SLC24A5*

was annotated as the union of coding region 1 and transcript *XM_007438007.2*. The boundaries

of coding region 1 were built to match the boundaries conserved across species (5'-ATG CAG

CCT GCC GAG....TCC GCG AGG ATC CCG -3'). The 5' boundary of coding region 2 was

adjusted to match the boundary conserved across species (5'-AGA ACG AAA CCC GCT...).

SLC45A2

The Burmese python genome was queried using *SLC45A2* protein sequence from corn

snake (*XP_034298386.1*). The N-terminal region of this query (coding regions 1 and 2) hit

Burmese python transcript *XM_007432459.2* on scaffold 1939. The C-terminal region of this

query (coding regions 5 to 7) hit Burmese python transcript *XM_007437721.2* on scaffold 3858.

SLC45A2 was annotated as the union of these transcripts.

DNA extraction

Sheds were rinsed in tap water to remove dirt and debris. Sheds were air dried and

lysed overnight at ~60°C in ~1 ml lysis buffer (100 mM Tris-HCl pH 8.0, 100 mM EDTA, 2%

sodium dodecyl sulfate, 3 mM $CaCl_2$, 2 mg/ml Proteinase K) per ~8 cm^2 piece of shed. Lysate

was separated from visible fragments of undigested shed and further cleared by centrifugation

at 13,000 x g for 2 min. To precipitate protein, ammonium acetate was added to supernatant to

a final concentration of 1.875 M. Samples were incubated on ice for 5-10 min and centrifuged at

13,000 x g for 3-5 min at 4°C. Supernatant was mixed with an equal volume of magnetic bead

mixture (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 1.6 M NaCl, 0.2% Tween-20, 11% polyethylene

glycol, 0.04% washed SpeedBeads [Sigma #GE45152105050250]), and samples shaken for 5-

10 min. Beads were separated from supernatant using a magnet, washed twice in 0.2 ml 70%

ethanol for 2 min, and air dried for ~1 min. DNA was eluted from beads in TE buffer (10 mM

Tris-HCl pH 8.0, 1 mM EDTA) at 65°C for >5 min.

Primer design and PCR

Primers were designed against the genome of Burmese python or against genomic sequences from ball python obtained in an earlier step of the study. Primers were designed using Primer3 [109], using default parameters and a target annealing temperature of 60°C. Amplification was first tested at 57°C, to allow for occasional divergence between ball python and Burmese python genomic sequences. In some cases, annealing temperatures were later adjusted to 52°C, 60°C, or 61°C, to obtain stronger product or to eliminate background bands.

Genomic fragments were amplified using OneTaq polymerase (NEB #M0480) or Q5 polymerase (NEB #M0491). Genotyping assays described below used OneTaq, unless otherwise specified. OneTaq reactions consisted of 1X OneTaq Standard Reaction Buffer, 200 µM dNTPs, 0.2 µM of each primer, and 0.025 U/µl OneTaq polymerase. OneTaq thermocycling conditions were as follows: 94°C for 2 min; 30-35 cycles of 94°C for 30 sec, 52-61°C for 30 sec, and 68°C for 1-4 min; and 68°C for 5 min. Q5 reactions consisted of 1X Q5 Reaction Buffer, 200 µM dNTPs, 0.5 µM of each primer, and 0.02 U/µl Q5 polymerase. Q5 thermocycling conditions were as follows: 98°C for 30 sec; 30-35 cycles of 98°C for 10 sec, 58-61°C for 15 sec, and 72°C for 1.5-3 min; and 72°C for 5 min. Reactions used 10-100 ng template DNA per 20 µl volume.

Sanger sequencing

PCR products were purified for Sanger sequencing using magnetic beads or gel extraction. For magnetic-bead purification, PCR reactions were mixed with three volumes of magnetic-bead mixture (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 1.6-2.5 M NaCl, 0.2% Tween-20, 11-20% polyethylene glycol, 0.04% washed SpeedBeads [Sigma #GE45152105050250]), and agitated for 5 min. Beads were separated from supernatant using a magnet, washed twice in 0.2 ml 80% ethanol for >30 sec, and air-dried for 30 sec. PCR products were eluted from beads in 10 mM Tris-HCl pH 8.0 for >3 min at 65°C. Gel extraction was performed using QIAquick Gel

632   Extraction Kit (Qiagen #28704), according to the manufacturer guidelines. Sanger sequencing

633   was performed by Eton Bioscience Inc (etonbio.com).

Sequencing of coding regions and comparison to Burmese python

Coding regions of melanogenesis genes were amplified and sequenced using primers

given in S3 Table. Chromatograms were trimmed using SnapGene Viewer

(snapgene.com/snapgene-viewer) and aligned to one another or to genomic sequences from

Burmese python using ApE (jorgensen.biology.utah.edu/wayned/ape). Alignments were

examined manually to identify divergent and polymorphic sites. Sequence identity between ball

python and Burmese python was calculated across alignable sequence, excluding indels.

Association study genotyping

Variants for the association study were identified through amplification and pilot

sequencing of genomic fragments from five animals (four Non-Albinos and one Albino). For

TYR, *TYRP1*, and *TYRP2*, we identified one or more genomic fragments containing multiple

polymorphic sites within the same amplicon. These genomic fragments were selected for the

association study. For *OCA2*, we did not identify any genomic fragments containing multiple

polymorphic sites; we therefore selected two genomic fragments each containing a single

polymorphic site. Divergence between the Albino animal and the other four animals was not a

criterion for inclusion of variants in the association study. Variants were genotyped by amplifying

and sequencing genomic fragments containing the variants. Locations of variants and primers

used for amplifying and sequencing variants are given in S1 Table. Genotypes are provided in

S6 Table.

Haplotype reconstruction

Haplotypes were reconstructed using PHASE version 2.1 [91,92]. Parameters were set to 200 iterations, a thinning interval of 2, and a burn-in parameter of 100. These settings produced identical or nearly identical output for seven runs seeded with different random numbers; thus, these settings met the criteria for effective choice of parameter settings, according to the PHASE documentation. The case-control permutation test was performed by comparing Albinos to Non-Albinos.

Genotyping assays for *TYR*

TYR missense variants *D394G* and *P384L* were genotyped by amplifying and sequencing a genomic fragment containing *TYR* coding region 3. This fragment was amplified using primers 13F (5'-ACT TTC AGG TGG GCA GCA G-3') and 13R (5'-GCT GAC AAC TAA AAT CTC TGC AA-3') and an annealing temperature of 52°C, or using primers 242F (5'-GCC ATT GTA GCT TCT TAC CAC TC-3') and 242R (5'-TTC CAG TCC ATA TAC AAG ATA TCC AA-3') and an annealing temperature of 57°C. The amplicon was sequenced using primer 13F or 242F. Genotypes are provided in S6 Table.

TYR promoter and intronic regions are amplified using primers given in S4 Table. Promoter fragments were sequenced in full, and intronic fragments were sequenced from one end, to confirm that the correct region of the genome had been amplified. Sequencing primers are given in S4 Table. Size differences among intronic fragments were assessed by separating fragments on a 1.25% agarose gel. Gels were run long enough for the shortest ladder band (100 bp) to migrate ~8 cm from its starting position.

Allele sizes of the variable number tandem repeat (VNTR) in *TYR* were genotyped by amplifying a genomic fragment located in *TYR* intron 3, using primers 214F (5'-TCT CAC CTG

677 ATG GCA CAT TC-3') and 209R (5'-GTG CCC ACC CTG ATG TTA TT-3') and an annealing

678 temperature of 60°C. Amplicon sizes were analyzed as for intronic fragments, described above.

Genotyping assays for *OCA2*

The *OCA2* deletion was initially identified by amplifying and sequencing a genomic

fragment spanning *OCA2* coding region 18. This fragment was amplified using primers 218F (5'-

ACC CCG TAG CCT CTT CAA AT-3') and 166R (5'-TGG GTG GCA AAC AAT CAT AA-3'), an

annealing temperature of 60°C, and Q5 polymerase. Amplicons were sequenced using both

primers.

The *OCA2* deletion was genotyped after its initial identification using a three-primer PCR

assay. This assayed used one forward primer and two reverse primers. The forward primer and

one of the reverse primers were located outside the deletion: 217F (5'-GGA GAG AGA ATC

CAA CCC TTG -3') and 166R (5'-TGG GTG GCA AAC AAT CAT AA-3'). The second reverse

primer was located within the deletion: 188R (5'-CAA AGA CCA TTG TCC ATT TCC-3'). The

annealing temperature was 57°C. This assay produces a 429-bp product for the wildtype allele

and a 349-bp product for the deletion allele. Heterozygotes produce both products. Genotypes

are provided in S6 Table.

Genotyping assays for *TYRP1*

TYRP1 missense variant *R305H* was genotyped by amplifying and sequencing a

genomic fragment spanning *TYRP1* coding region 4. This fragment was amplified using primers

18F (5'-GCT CTT TTC TCT AAG TCT GAC CTC -3') and 18R (5'-TCT TGT CCC ACA AAA

GGA TTT -3') and an annealing temperature of 57°C. The amplicon was sequenced using

primer 18F.

The putative deletion of *TYRP1* coding regions 6 and 7 was identified using primers

given in S5 Table. The putative deletion was genotyped after its initial identification by

amplifying a genomic fragment of *TYRP1* spanning coding regions 6 and 7. This fragment was

amplified using primers 20F (5'-GCA TTG TTT TAT CAG CCA TGA A-3') and 21R (5'-GGA ATT

GAG ACA AAT CCT TGG-3') and an annealing temperature of 57°C. Genotypes are provided in

S6 Table.

Protein sequence alignment

Protein sequences were aligned using Clustal Omega [110], using default parameters.

Acknowledgements

We thank Matt Rockman and Katy Greenwald for advice on haplotype reconstruction; Bob Winning, Anne Casper, David Kass, and two anonymous reviewers for comments on the manuscript; and the Educational Course Support program of New England BioLabs for reagents used in undergraduate teaching labs. We thank the following individuals for contributing ball python sheds: Adam and Nicole Schmid; Alycia Butler; Amanda Hall; Andelyn Czajka; Brad Carter of Driftless Reptiles; Bryan Rivera; Chiron Graves; Chun Ku of Dynasty Reptiles; Dale Porcher; Daniel Ross; David Wolf of Tornado Alley Reptiles; David Burstein; Dawn and Kelsi Greene of Super Natural Balls; Dayna Plehn; Debby Brauer; Epic Vibrant Balls; Eric Chung of Chung Reptiles; Erin Burt; George Straub; Haily McCullough; Jaden Christensen; Jake Lewis; Jamie Palazzo of New Day Reptiles; Jeff Kearns; Jeff Linton; Jodi Wilkowski; Joe Myers; John Cordone of Blue Water Reptiles; Jordan Noland; Justin Kobylka of J. Kobylka Reptiles; Lindsay VanOrman; Lisa Huis; Manuel San Juan; Mark Bilger; Lynnet Melton; Maryann Barbon; Mia Hynes; Michael, Lisa, and Bodie Cole of Ballroom Pythons South; Morgan Evans and Michael Kitto of MK Pythons; Morgan Shelton; Paul and Amber Fiorito of Vivid Scales; Pets 'n' Things of Saline, MI; Rachel Voyt; Royal Black Balls; Ryan Boyd and Brittney Delacruz; Ryan Young of Molecular Reptile; Sergio McDole; Stephanie S. Crisp; Steve P. of Prime Pets; Zac Parpart; and several anonymous contributors. We thank the following individuals for providing images of color morphs: Beth Woodyard; Cat Church; Cormier Jason; Christine Miller; Darin Taylor; Daniel Hatcher; David C. Callahan; Donald Grinstead; Elijah Snyder; Innovative Ectotherms; Jake Lewis; Jessica Allison; Jessica Van Riper; Justin Kobylka; James Thompson; Justin Revington; Mariette van Vuuren; Mark Hopkins; Mark Smith; Matthew Lopez; Michael Freedman of The Florida Reptile Ranch; Morgan Evans; Phil Barclay, Robert Cooper; Ron Heisenberg; Ryan Young; Selectively Bred Serpents; Seb Des Légendes Celtiques; and Thananan Jivaramonaikul.

Most of the data in this study were collected by undergraduates enrolled in a laboratory

course at Eastern Michigan University. These students constitute the BIO306W Consortium.

These students were Alexandra Ernst, Alia Frederick, Alissa Zoltowski, Amber Northcutt,

Andraya Ackerman, Anna Pathammavong, Annette Miller, Ashly Matzek, Asra Akhlaq, Aubrey

Martin, Bailey Knight, Benjamin Huff, Beth Wasserman, Brian Donald Condron, Caleb Sommer,

Cassandra Rigor, Charles Southwell, Chase Chitwood, Chelsea Brown, Christina Roka, Ciarra

Womack, Clay McKenzie, Daniela Nappo, Darby Fracassa, Deirdre McCarter, Dhruvalkumar

Patel, Dominic Paoletti, Drake Dzierwa, Erica R. Geml, Erin Bissett, Ezekiel Butcher, Garrett

Chance, Garry Lewis, Genesis Garmendia, Geo Pullockaran, Hajer Musa Abuzir, Haley Praski,

Hanan Alroaini, Iqra Akhlaq, Ismael Yasin, Janelle Aethyr, Janelle Janisse, Jayce Alee

Perysian, Jemar Rooks, Jonathan Chang, Jonathan Harris, Joseph H. Oberlitner, Joshua

Mason, Juwan Taylor, Kailynn Sparks, Karissa Urban, Karli Siefman, Kealy Szymanski, Kelsy

Roque, Keyan Marshall, Khaled Ali, Karleigh Hassenzahl, Kylie Powrie, Lauren Colone, Lissette

Rosas, Manoj Perumallapalli, Mariam Samir, Maryam Nimer, Maya Mackey, Megan McNulty,

Mel Roberts, Micaela Schempf, Molly Cook, Myah Kelly, Nahiel Sukar, Natalie Diaz, Natasha

MacKay, Nathan Barnett, Nathaniel Gonzalez, Noura Taybeh, Pablo De la Vega, Rida Ali,

Ronnie Bryans, Ryan Elliott, Saja Hussein, Samantha Glowacki, Samuel Teener, Sarah

Holtzen, Sarah Schmidt, Shanti Bernstein, Shelan Mizuree, Smarpita Singh, Stevie Zabrosky,

Taia Broadbent, Tommiea Robertson, Tyler Schallhorn, Verginio Persicone, William Soder,

Wolfgang Ebersole, and Yvette Campbell.

References

1.   Cuthill IC, Allen WL, Arbuckle K, Caspers B, Chaplin G, Hauber ME, et al. The biology of
color. Science. 2017;357: eaan0221. doi:10.1126/science.aan0221

2.   Cieslak M, Reissmann M, Hofreiter M, Ludwig A. Colours of domestication. Biol Rev.
2011;86: 885–899. doi:10.1111/j.1469-185X.2011.00177.x

3.   Toews DPL, Hofmeister NR, Taylor SA. The Evolution and Genetics of Carotenoid
Processing in Animals. Trends Genet. 2017;33: 171–182. doi:10.1016/j.tig.2017.01.002

4.   Prum RO. Anatomy, physics, and evolution of avian structural colors. Bird Coloration.
Cambridge, MA: Harvard University Press; 2006.

5.   Ghiradella H. Structure and Development of Iridescent Lepidopteran Scales - the
Papilionidae as a Showcase Family. Ann Entomol Soc Am. 1985;78: 252–264.
doi:10.1093/aesa/78.2.252

6.   Bagnara JT, Taylor JD, Hadley ME. The dermal chromatophore unit. J Cell Biol. 1968;38:
67–79. doi:10.1083/jcb.38.1.67

7.   Saenko SV, Teyssier J, van der Marel D, Milinkovitch MC. Precise colocalization of
interacting structural and pigmentary elements generates extensive color pattern variation
in Phelsuma lizards. BMC Biol. 2013;11: 105. doi:10.1186/1741-7007-11-105

8.   Kinoshita S, Yoshioka S. Structural colors in nature: The role of regularity and irregularity
in the structure. ChemPhysChem. 2005;6: 1442–1459. doi:10.1002/cphc.200500007

9.   Slominski A, Tobin DJ, Shibahara S, Wortsman J. Melanin pigmentation in mammalian
skin and its hormonal regulation. Physiol Rev. 2004;84: 1155–1228.
doi:10.1152/physrev.00044.2003

10. Meredith P, Sarna T. The physical and chemical properties of eumelanin. Pigm Cell Res.
2006;19: 572–594. doi:10.1111/j.1600-0749.2006.00345.x

11. Bagnara JT, Fernandez PJ, Fujii R. On the blue coloration of vertebrates. Pigm Cell Res.
2007;20: 14–26. doi:10.1111/j.1600-0749.2006.00360.x

12. Olsson M, Stuart-Fox D, Ballen C. Genetics and evolution of colour patterns in reptiles.
Semin Cell Dev Biol. 2013;24: 529–541. doi:10.1016/j.semcdb.2013.04.001

13. Shawkey MD, D'Alba L. Interactions between colour-producing mechanisms and their
effects on the integumentary colour palette. Philos Trans R Soc B-Biol Sci. 2017;372:
20160536. doi:10.1098/rstb.2016.0536

14. Bagnara JT, Matsumoto J. Comparative Anatomy and Physiology of Pigment Cells in
Nonmammalian Tissues. The Pigmentary System. John Wiley & Sons, Ltd; 2006. pp. 11–
59. doi:10.1002/9780470987100.ch2

15. Singh AP, Nuesslein-Volhard C. Zebrafish Stripes as a Model for Vertebrate Colour
Pattern Formation. Curr Biol. 2015;25: R81–R92. doi:10.1016/j.cub.2014.11.013

16. Huang D, Lewis VM, Foster TN, Toomey MB, Corbo JC, Parichy DM. Development and
genetics of red coloration in the zebrafish relative Danio albolineatus. eLife. 2021;10:
e70253. doi:10.7554/eLife.70253

17. Kimura T, Nagao Y, Hashimoto H, Yamamoto-Shiraishi Y, Yamamoto S, Yabe T, et al.
Leucophores are similar to xanthophores in their specification and differentiation
processes in medaka. Proc Natl Acad Sci U S A. 2014;111: 7343–7348.
doi:10.1073/pnas.1311254111

18. Nusslein-Volhard C, Singh AP. How fish color their skin: A paradigm for development and
evolution of adult patterns Multipotency, plasticity, and cell competition regulate
proliferation and spreading of pigment cells in Zebrafish coloration. Bioessays. 2017;39:
1600231. doi:10.1002/bies.201600231

19. Cooke TF, Fischer CR, Wu P, Jiang T-X, Xie KT, Kuo J, et al. Genetic Mapping and
Biochemical Basis of Yellow Feather Pigmentation in Budgerigars. Cell. 2017;171: 427-+.
doi:10.1016/j.cell.2017.08.016

20. Gazda MA, Araujo PM, Lopes RJ, Toomey MB, Andrade P, Afonso S, et al. A genetic
mechanism for sexual dichromatism in birds. Science. 2020;368: 1270-+.
doi:10.1126/science.aba0803

21. Kwon YM, Vranken N, Hoge C, Lichak MR, Francis KX, Camacho-Garcia J, et al.
Genomic consequences of domestication of the Siamese fighting fish. 2021 Apr p.
2021.04.29.442030. doi:10.1101/2021.04.29.442030

22. Lopes RJ, Johnson JD, Toomey MB, Ferreira MS, Araujo PM, Melo-Ferreira J, et al.
Genetic Basis for Red Coloration in Birds. Curr Biol. 2016;26: 1427–1434.
doi:10.1016/j.cub.2016.03.076

23. Mundy NI, Stapley J, Bennison C, Tucker R, Twyman H, Kim K-W, et al. Red Carotenoid
Coloration in the Zebra Finch Is Controlled by a Cytochrome P450 Gene Cluster. Curr
Biol. 2016;26: 1435–1440. doi:10.1016/j.cub.2016.04.047

24. Toomey MB, Lopes RJ, Araujo PM, Johnson JD, Gazda MA, Afonso S, et al. High-density
lipoprotein receptor SCARB1 is required for carotenoid coloration in birds. Proc Natl Acad
Sci U S A. 2017;114: 5219–5224. doi:10.1073/pnas.1700751114

25. Zhang W, Wang H, Brandt DYC, Hu B, Sheng J, Wang M, et al. The genetic architecture
of phenotypic diversity in the betta fish (Betta splendens). bioRxiv. 2021;
2021.05.10.443352. doi:10.1101/2021.05.10.443352

26. Garcia-Elfring A, Roffey HL, Hendry AP, Barrett RDH. A nonsense mutation in TFEC is
the likely cause of the recessive piebald phenotype in ball pythons (Python regius). 2020
Nov p. 2020.10.30.362970. doi:10.1101/2020.10.30.362970

27.   Guo L, Bloom J, Sykes S, Huang E, Kashif Z, Pham E, et al. Genetics of white color and
824   iridophoroma in "Lemon Frost" leopard geckos. PLoS Genet. 2021;17: e1009580.
825   doi:10.1371/journal.pgen.1009580

826   28.   McLean CA, Lutz A, Rankin KJ, Stuart-Fox D, Moussalli A. Revealing the Biochemical
827   and Genetic Basis of Color Variation in a Polymorphic Lizard. Mol Biol Evol. 2017;34:
828   1924–1935. doi:10.1093/molbev/msx136

829   29.   Ullate-Agote A, Burgelin I, Debry A, Langrez C, Montange F, Peraldi R, et al. Genome
830   mapping of a LYST mutation in corn snakes indicates that vertebrate chromatophore
831   vesicles are lysosome-related organelles. Proc Natl Acad Sci U S A. 2020;117: 26307–
832   26317. doi:10.1073/pnas.2003724117

833   30.   Bale R. Ball python exports raise concerns as demand for the popular pet grows. National
834   Geographic. 2020.

835   31.   Broghammer S. Python regius: Atlas of Colour Morphs Keeping and Breeding. NTV Natur
836   und Tier-Verlag; 2019.

837   32.   Irizarry KJL, Bryden RL. In Silico Analysis of Gene Expression Network Components
838   Underlying Pigmentation Phenotypes in the Python Identified Evolutionarily Conserved
839   Clusters of Transcription Factor Binding Sites. Adv Bioinformatics. 2016;2016: 1286510.
840   doi:10.1155/2016/1286510

841   33.   McCurley K. Complete Ball Python, A Comprehensive Guide to Care, Breeding, and
842   Genetic Mutations: Kevin McCurley: 9780976733409: Amazon.com: Books. ECO /
843   Serpent's Tale NHBD; 2005.

844   34.   Burbrink FT, Castoe TA. Molecular Phylogeography of Snakes. Snakes: Ecology and
845   Conservation. Cornell University Press; 2011. pp. 38–77. doi:10.7591/9780801459092-
846   006

847   35.   Liu Q, Qi Y, Liang Q, Song J, Liu J, Li W, et al. Targeted disruption of tyrosinase causes
848   melanin reduction in Carassius auratus cuvieri and its hybrid progeny. Sci China-Life Sci.
849   2019;62: 1194–1202. doi:10.1007/s11427-018-9404-7

850   36.   Koga A, Wakamatsu Y, Kurosawa J, Hori H. Oculocutaneous albinism in the i(6) mutant
851   of the medaka fish is associated with a deletion in the tyrosinase gene. Pigm Cell Res.
852   1999;12: 252–258. doi:10.1111/j.1600-0749.1999.tb00758.x

853   37.   Miura I, Tagami M, Fujitani T, Ogata M. Spontaneous tyrosinase mutations identified in
854   albinos of three wild frog species. Genes Genet Syst. 2017;92: 189–196.
855   doi:10.1266/ggs.16-00061

856   38.   Iwanishi S, Zaitsu S, Shibata H, Nitasaka E. An albino mutant of the Japanese rat snake
857   (Elaphe climacophora) carries a nonsense mutation in the tyrosinase gene. Genes Genet
858   Syst. 2018;93: 163–167. doi:10.1266/ggs.18-00021

859   39.   Tobita-Teramoto T, Jang GY, Kino K, Salter DW, Brumbaugh J, Akiyama T. Autosomal
860   albino chicken mutation (ca/ca) deletes hexanucleotide (-deltaGACTGG817) at a copper-
861   binding site of the tyrosinase gene. Poult Sci. 2000;79: 46–50. doi:10.1093/ps/79.1.46

40. Chang C-M, Coville J-L, Coquerelle G, Gourichon D, Oulmouden A, Tixier-Boichard M.
Complete association between a retroviral insertion in the tyrosinase gene and the
recessive white mutation in chickens. BMC Genomics. 2006;7: 19. doi:10.1186/1471-
2164-7-19

41. Chang CM, Furet JP, Coville JL, Coquerelle G, Gourichon D, Tixier-Boichard M.
Quantitative effects of an intronic retroviral insertion on the transcription of the tyrosinase
gene in recessive white chickens. Anim Genet. 2007;38: 162–167. doi:10.1111/j.1365-
2052.2007.01581.x

42. Florisbal Dame MC, Xavier GM, Oliveira-Filho JP, Borges AS, Oliveira HN, Riet-Correa F,
et al. A nonsense mutation in the tyrosinase gene causes albinism in water buffalo. BMC
Genet. 2012;13: 62. doi:10.1186/1471-2156-13-62

43. Schmutz SM, Berryere TG, Ciobanu DC, Mileham AJ, Schmidtz BH, Fredholm M. A form
of albinism in cattle is caused by a tyrosinase frameshift mutation. Mamm Genome.
2004;15: 62–67. doi:10.1007/s00335-002-2249-5

44. Reiner G, Tramberend K, Nietfeld F, Volmer K, Wurmser C, Fries R, et al. A genome-
wide scan study identifies a single nucleotide substitution in the tyrosinase gene
associated with white coat colour in a red deer (Cervus elaphus) population. BMC Genet.
2020;21. doi:10.1186/s12863-020-0814-0

45. Utzeri VJ, Bertolini F, Ribani A, Schiavo G, Dall'Olio S, Fontanesi L. The albinism of the
feral Asinara white donkeys (Equus asinus) is determined by a missense mutation in a
highly conserved position of the tyrosinase (TYR) gene deduced protein. Anim Genet.
2016;47: 120–124. doi:10.1111/age.12386

46. Yokoyama T, Silversides D, Waymire K, Kwon B, Takeuchi T, Overbeek P. Conserved
Cysteine to Serine Mutation in Tyrosinase Is Responsible for the Classical Albino
Mutation in Laboratory Mice. Nucleic Acids Res. 1990;18: 7293–7298.
doi:10.1093/nar/18.24.7293

47. Blaszczyk WM, Arning L, Hoffmann KP, Epplen JT. A Tyrosinase missense mutation
causes albinism in the Wistar rat. Pigm Cell Res. 2005;18: 144–145. doi:10.1111/j.1600-
0749.2005.00227.x

48. Yu F, Jiao S, Lai W, Liu Z, Zhu M, Zhu W, et al. Conserved aspartate-to-glycine mutation
in tyrosinase is associated with albino phenotype in domestic guinea pigs (Cavia
porcellus). Anim Genet. 2018;49: 354–355. doi:10.1111/age.12683

49. Kim Y-H, Park S-J, Choe S-H, Lee J-R, Cho H-M, Kim S-U, et al. Identification and
characterization of the tyrosinase gene (TYR) and its transcript variants (TYR_1 and
TYR_2) in the crab-eating macaque (Macaca fascicularis). Gene. 2017;630: 21–27.
doi:10.1016/j.gene.2017.07.047

50. Galante Rocha de Vasconcelos FT, Hauzman E, Henriques LD, Kilpp Goulart PR, Galvao
O de F, Sano RY, et al. A novel nonsense mutation in the tyrosinase gene is related to
the albinism in a capuchin monkey (Sapajus apella). BMC Genet. 2017;18: 39.
doi:10.1186/s12863-017-0504-8

51.   Koga A, Hisakawa C, Yoshizawa M. Baboon bearing resemblance in pigmentation
pattern to Siamese cat carries a missense mutation in the tyrosinase gene. Genome.
2020;63. doi:10.1139/gen-2020-0003

52.   Montoliu L, Gronskov K, Wei A-H, Martinez-Garcia M, Fernandez A, Arveiler B, et al.
Increasing the complexity: new genes and new types of albinism. Pigment Cell Melanoma
Res. 2014;27. doi:10.1111/pcmr.12167

53.   Polanowski AM, Robinson-Laverick SM, Paton D, Jarman SN. Variation in the Tyrosinase
Gene Associated with a White Humpback Whale (Megaptera novaeangliae). J Hered.
2012;103: 130–133. doi:10.1093/jhered/esr108

54.   Blaszczyk WM, Distler C, Dekomien G, Arning L, Hoffmann K-P, Epplen JT. Identification
of a tyrosinase (TYR) exon 4 deletion in albino ferrets (Mustela putorius furo). Anim
Genet. 2007;38: 421–423. doi:10.1111/j.1365-2052.2007.01619.x

55.   Anistoroaei R, Fredholm M, Christensen K, Leeb T. Albinism in the American mink
(Neovison vison) is associated with a tyrosinase nonsense mutation. Anim Genet.
2008;39: 645–648. doi:10.1111/j.1365-2052.2008.01788.x

56.   Imes DL, Geary LA, Grahn RA, Lyons LA. Albinism in the domestic cat (Felis catus) is
associated with a tyrosinase (TYR) mutation. Anim Genet. 2006;37: 175–178.
doi:10.1111/j.1365-2052.2005.01409.x

57.   Lyons LA, Imes DL, Rah HC, Grahn RA. Tyrosinase mutations associated with Siamese
and Burmese patterns in the domestic cat (Felis catus). Anim Genet. 2005;36: 119–126.
doi:10.1111/j.1365-2052.2005.01253.x

58.   Schmidt-Kuntzel A, Eizirik E, O'Brien SJ, Menotti-Raymond M. Tyrosinase and tyrosinase
related protein 1 alleles specify domestic cat coat color phenotypes of the albino and
brown loci. J Hered. 2005;96: 289–301. doi:10.1093/jhered/esi066

59.   Yan S, Zhao D, Hu M, Tan X, Lai W, Kang J, et al. A single base insertion in the
tyrosinase gene is associated with albino phenotype in silver foxes (Vulpes vulpes). Anim
Genet. 2019;50: 550–550. doi:10.1111/age.12816

60.   Gross JB, Wilkens H. Albinism in phylogenetically and geographically distinct populations
of Astyanax cavefish arises through the same loss-of-function Oca2 allele. Heredity.
2013;111: 122–130. doi:10.1038/hdy.2013.26

61.   Kratochwil CF, Urban S, Meyer A. Genome of the Malawi golden cichlid fish
(Melanochromis auratus) reveals exon loss of oca2 in an amelanistic morph. Pigment Cell
Melanoma Res. 2019;32: 719–723. doi:10.1111/pcmr.12799

62.   Saenko SV, Lamichhaney S, Barrio AM, Rafati N, Andersson L, Milinkovitch MC.
Amelanism in the corn snake is associated with the insertion of an LTR-retrotransposon
in the OCA2 gene. Sci Rep. 2015;5: 17118. doi:10.1038/srep17118

63.   Caduff M, Bauer A, Jagannathan V, Leeb T. OCA2 splice site variant in German Spitz
dogs with oculocutaneous albinism. PLoS One. 2017;12: e0185944.
doi:10.1371/journal.pone.0185944

64. Zhang Y, Hong Q, Cao C, Yang L, Li Y, Hai T, et al. A novel porcine model reproduces
human oculocutaneous albinism type. Cell Discov. 2019;5: 48. doi:10.1038/s41421-019-
0117-7

65. Shoji H, Kiniwa Y, Okuyama R, Yang M, Higuchi K, Mori M. A nonsense nucleotide
substitution in the oculocutaneous albinism II gene underlies the original pink-eyed
dilution allele (Oca2(p)) in mice. Exp Anim. 2015;64: 171–179. doi:10.1538/expanim.14-
0075

66. Krauss J, Geiger-Rudolph S, Koch I, Nuesslein-Volhard C, Irion U. A dominant mutation
in tyrp1A leads to melanophore death in zebrafish. Pigment Cell Melanoma Res. 2014;27:
827–830. doi:10.1111/pcmr.12272

67. Cortimiglia C, Castiglioni B, Pizzi F, Stella A, Capra E. Involvement of tyrosinase-related
protein 1 gene in the light brown plumage phenotype of Falco cherrug. Anim Genet.
2017;48: 125–126. doi:10.1111/age.12506

68. Ren J, Mao H, Zhang Z, Xiao S, Ding N, Huang L. A 6-bp deletion in the TYRP1 gene
causes the brown colouration phenotype in Chinese indigenous pigs. Heredity. 2011;106:
862–868. doi:10.1038/hdy.2010.129

69. Wu X, Zhang Y, Shen L, Du J, Luo J, Liu C, et al. A 6-bp deletion in exon 8 and two
mutations in introns of TYRP1 are associated with blond coat color in Liangshan pigs.
Gene. 2016;578: 132–136. doi:10.1016/j.gene.2015.12.011

70. Lyons LA, Foe IT, Rah HC, Grahn RA. Chocolate coated cats: TYRP1 mutations for
brown color domestic cats. Mamm Genome. 2005;16: 356–366. doi:10.1007/s00335-004-
2455-4

71. Gratten J, Beraldi D, Lowder BV, McRae AF, Visscher PM, Pemberton JM, et al.
Compelling evidence that a single nucleotide substitution in TYRP1 is responsible for
coat-colour polymorphism in a free-living population of Soay sheep. Proc R Soc B-Biol
Sci. 2007;274: 619–626. doi:10.1098/rspb.2006.3762

72. Paris JM, Letko A, Hafliger IM, Ammann P, Flury C, Drogemuller C. Identification of two
TYRP1 loss-of-function alleles in Valais Red sheep. Anim Genet. 2019;50: 778–782.
doi:10.1111/age.12863

73. Cirera S, Markakis MN, Kristiansen T, Vissenberg K, Fredholm M, Christensen K, et al. A
large insertion in intron 2 of the TYRP1 gene associated with American Palomino
phenotype in American mink. Mamm Genome. 2016;27: 135–143. doi:10.1007/s00335-
016-9620-4

74. Budd P, Jackson I. Structure of the Mouse Tyrosinase-Related Protein-2 Dopachrome
Tautomerase (tyrp2/Dct) Gene and Sequence of 2 Novel Slaty Alleles. Genomics.
1995;29: 35–43. doi:10.1006/geno.1995.1212

75. Jackson I, Chambers D, Tsukamoto K, Copeland N, Gilbert D, Jenkins N, et al. A 2nd
Tyrosinase-Related Protein, Trp-2, Maps to and Is Mutated at the Mouse Slaty Locus.
Embo J. 1992;11: 527–535. doi:10.1002/j.1460-2075.1992.tb05083.x

76. Chintala S, Li W, Lamoreux ML, Ito S, Wakamatsu K, Sviderskaya EV, et al. Slc7a11
gene controls production of pheomelanin pigment and proliferation of cultured cells. Proc
Natl Acad Sci U S A. 2005;102: 10964–10969. doi:10.1073/pnas.0502856102

77. Lamason RL, Mohideen M, Mest JR, Wong AC, Norton HL, Aros MC, et al. SLC24A5, a
putative cation exchanger, affects pigmentation in zebrafish and humans. Science.
2005;310: 1782–1786. doi:10.1126/science.1116238

78. Winkler PA, Gornik KR, Ramsey DT, Dubielzig RR, Venta PJ, Petersen-Jones SM, et al.
A Partial Gene Deletion of SLC45A2 Causes Oculocutaneous Albinism in Doberman
Pinscher Dogs. PLoS One. 2014;9: e92127. doi:10.1371/journal.pone.0092127

79. Mack M, Kowalski E, Grahn R, Bras D, Penedo MCT, Bellone R. Two Variants in
SLC24A5 Are Associated with "Tiger-Eye" Iris Pigmentation in Puerto Rican Paso Fino
Horses. G3-Genes Genomes Genet. 2017;7: 2799–2806. doi:10.1534/g3.117.043786

80. Fukamachi S, Shimada A, Shima A. Mutations in the gene encoding B, a novel
transporter protein, reduce melanin content in medaka. Nature Genet. 2001;28: 381–385.
doi:10.1038/ng584

81. Gunnarsson U, Hellstrom AR, Tixier-Boichard M, Minvielle F, Bed'hom B, Ito S, et al.
Mutations in SLC45A2 cause plumage color variation in chicken and Japanese quail.
Genetics. 2007;175: 867–877. doi:10.1534/genetics.106.063107

82. Mariat D, Taourit S, Guerin G. A mutation in the MATP gene causes the cream coat
colour in the horse. Genet Sel Evol. 2003;35: 119–133. doi:10.1051/gse:2002039

83. Newton JM, Cohen-Barak O, Hagiwara N, Gardner JM, Davisson MT, King RA, et al.
Mutations in the human orthologue of the mouse underwhite gene (uw) underlie a new
form of oculocutaneous albinism, OCA4. Am J Hum Genet. 2001;69: 981–988.
doi:10.1086/324340

84. Xu X, Dong G-X, Hu X-S, Miao L, Zhang X-L, Zhang D-L, et al. The Genetic Basis of
White Tigers. Curr Biol. 2013;23: 1031–1035. doi:10.1016/j.cub.2013.04.054

85. Prado-Martinez J, Hernando-Herraez I, Lorente-Galdos B, Dabad M, Ramirez O, Baeza-
Delgado C, et al. The genome sequencing of an albino Western lowland gorilla reveals
inbreeding in the wild. BMC Genomics. 2013;14: 363. doi:10.1186/1471-2164-14-363

86. Lai X, Wichers HJ, Soler-Lopez M, Dijkstra BW. Structure and Function of Human
Tyrosinase and Tyrosinase-Related Proteins. Chem-Eur J. 2018;24: 47–55.
doi:10.1002/chem.201704410

87. Bellono NW, Escobar IE, Lefkovith AJ, Marks MS, Oancea E. An intracellular anion
channel critical for pigmentation. eLife. 2014;3: e04543. doi:10.7554/eLife.04543

88. Ginger RS, Askew SE, Ogborne RM, Wilson S, Ferdinando D, Dadd T, et al. SLC24A5
encodes a trans-golgi network protein with potassium-dependent sodium-calcium
exchange activity that regulates human epidermal melanogenesis. J Biol Chem.
2008;283: 5486–5495. doi:10.1074/jbc.M707521200

89. Vitavska O, Wieczorek H. The SLC45 gene family of putative sugar transporters. Mol Asp
Med. 2013;34: 655–660. doi:10.1016/j.mam.2012.05.014

90. Castoe TA, de Koning APJ, Hall KT, Card DC, Schield DR, Fujita MK, et al. The Burmese
python genome reveals the molecular basis for extreme adaptation in snakes. Proc Natl
Acad Sci U S A. 2013;110: 20645–20650. doi:10.1073/pnas.1314475110

91. Stephens M, Scheet P. Accounting for decay of linkage disequilibrium in haplotype
inference and missing-data imputation. Am J Hum Genet. 2005;76: 449–462.
doi:10.1086/428594

92. Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction
from population data. Am J Hum Genet. 2001;68: 978–989. doi:10.1086/319501

93. Simeonov DR, Wang X, Wang C, Sergeev Y, Dolinska M, Bower M, et al. DNA Variations
in Oculocutaneous Albinism: An Updated Mutation List and Current Outstanding Issues in
Molecular Diagnostics. Hum Mutat. 2013;34: 827–835. doi:10.1002/humu.22315

94. Gardner J, Nakatsu Y, Gondo Y, Lee S, Lyon M, King R, et al. The Mouse Pink-Eyed
Dilution Gene - Association with Human Prader-Willi and Angelman Syndromes. Science.
1992;257: 1121–1124. doi:10.1126/science.257.5073.1121

95. Ni-Komatsu L, Orlow SJ. Heterologous expression of tyrosinase recapitulates the
misprocessing and mistrafficking in oculocutaneous albinism type 2: Effects of altering
intracellular pH and pink-eyed dilution gene expression. Exp Eye Res. 2006;82: 519–528.
doi:10.1016/j.exer.2005.08.013

96. Puri N, Gardner JM, Brilliant MH. Aberrant pH of melanosomes in pink-eyed dilution (p)
mutant melanocytes. J Invest Dermatol. 2000;115: 607–613. doi:10.1046/j.1523-
1747.2000.00108.x

97. Lee S, Nicholls R, Bundey S, Laxova R, Musarella M, Spritz R. Mutations of the P-Gene
in Oculocutaneous Albinism, Ocular Albinism, and Prader-Willi-Syndrome Plus Albinism.
N Engl J Med. 1994;330: 529–534. doi:10.1056/NEJM199402243300803

98. Sato H, Tamba M, Ishii T, Bannai S. Cloning and expression of a plasma membrane
cystine/glutamate exchange transporter composed of two distinct proteins. J Biol Chem.
1999;274: 11455–11458. doi:10.1074/jbc.274.17.11455

99. Bin B-H, Bhin J, Yang SH, Shin M, Nam Y-J, Choi D-H, et al. Membrane-Associated
Transporter Protein (MATP) Regulates Melanosomal pH and Influences Tyrosinase
Activity. PLoS One. 2015;10: e0129273. doi:10.1371/journal.pone.0129273

100. Sturm R, Osullivan B, Box N, Smith A, Smit S, Puttick E, et al. Chromosomal Structure of
the Human Tyrp1 and Tyrp2 Loci and Comparison of the Tyrosinase-Related Protein
Cone Family. Genomics. 1995;29: 24–34. doi:10.1006/geno.1995.1211

101. Gershonibaruch R, Rosenmann A, Droetto S, Holmes S, Tripathi R, Spritz R. Mutations of
the Tyrosinase Gene in Patients with Oculocutaneous Albinism from Various Ethnic-
Groups in Israel. Am J Hum Genet. 1994;54: 586–594.

102. Tripathi RK, Strunk KM, Giebel LB, Weleber RG, Spritz RA. Tyrosinase gene mutations in
type I (tyrosinase-deficient) oculocutaneous albinism define two clusters of missense
substitutions. Am J Med Genet. 1992;43: 865–871. doi:10.1002/ajmg.1320430523

103. Tsai CH, Tsai FJ, Wu JY, Lin SP, Chang JG, Yang CF, et al. Insertion/deletion mutations
of type I oculocutaneous albinism in chinese patients from Taiwan. Hum Mutat. 1999;14:
542. doi:10.1002/(SICI)1098-1004(199912)14:6<542::AID-HUMU14>3.0.CO;2-3

104. Forshew T, Khaliq S, Tee L, Smith U, Johnson CA, Mehdi SQ, et al. Identification of novel
TYR and TYRP1 mutations in oculocutaneous albinism. Clin Genet. 2005;68: 182–184.
doi:10.1111/j.1399-0004.2005.00460.x

105. Lasseaux E, Plaisant C, Michaud V, Pennamen P, Trimouille A, Gaston L, et al.
Molecular characterization of a series of 990 index patients with albinism. Pigment Cell
Melanoma Res. 2018;31: 466–474. doi:10.1111/pcmr.12688

106. Rooryck C, Morice-Picard F, Elcioglu NH, Lacombe D, Taieb A, Arveiler B. Molecular
diagnosis of oculocutaneous albinism: new mutations in the OCA1-4 genes and practical
aspects. Pigment Cell Melanoma Res. 2008;21: 583–587. doi:10.1111/j.1755-
148X.2008.00496.x

107. Rooryck C, Morice F, Lacombe D, Taieb A, Arveiler B. Genetic basis of oculocutaneous
albinism. Expert Review of Dermatology. 2009;4: 611–622. doi:10.1586/edm.09.53

108. Allen WL, Baddeley R, Scott-Samuel NE, Cuthill IC. The evolution and function of pattern
diversity in snakes. Behav Ecol. 2013;24: 1237–1250. doi:10.1093/beheco/art058

109. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3-
new capabilities and interfaces. Nucleic Acids Res. 2012;40: e115.
doi:10.1093/nar/gks596

110. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation
of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.
2011;7: 539. doi:10.1038/msb.2011.75

1082

Supporting information

1083

S1 Fig. Raw gel images. (A) Raw image of the gel displayed in Fig 2. (B) Raw image of the gel

displayed in Fig 3. (C) Raw image of the gel displayed in Fig 4. Brightness and contrast settings

have not been adjusted in these images. x, experiment unrelated to the current study.

S1 Table. Variants used in the Albino association study and primers to genotype these

variants.

S2 Table. Conservation of gene structure of melanogenesis genes.

S3 Table. Primers to amplify and sequence coding regions of melanogenesis genes.

S4 Table. Primers to amplify non-coding regions of *TYR*.

S5 Table. Primers to investigate the putative deletion in *TYRP1*.

S6 Table. Genotypes.

Wildtype | Ultramel | Lavender Albino | Albino

A **B** **C** **D** Alb^{Albino} **E** Alb^{Candy} **F** Alb^{Toffee}

More melanin | Less melanin

Hypothesized causative gene: | *TYRP1, TYRP2, SLC7A11, SLC24A5,* or *SLC45A2* | *OCA2* | *TYR*

A

TYR

Association study variant

5 kb

Albinos | Non-Albinos

TYR^{Albino}
TYR^{D394G}
TYR^{P384L}

Haplotype

n = 50 | n = 56

Haplotype frequency (%)

B

TYR protein sequence

Position: 384 394

Elephant shark ... S A N D **P** I F L L H H V F V D S I F E ...
Zebrafish ... S A N D **P** I F I I H H A F **I** D S I F E ...
Tropical clawed frog ... S A N D **P** V F V L H H A F V D S I F E ...
Human ... S A N D **P** I F L L H H A F V D S M F E ...
Chicken ... S A N D **P** I F I L H H A F V D S I F E ...
Painted turtle ... S A N D **P** I F I L H H A F V D S I F E ...
Anole lizard ... S A N D **P** I F I L H H A F V D S I F E ...
Burmese python ... S A N D **P** V F I P H H A F V D S I Y E ...
Ball python ... S A N D **P** V F I P H H A F V D S I Y E ...

Ball python *D394G* ... S A N D **P** V F I P H H A F V **G** S I Y E ...
Ball python *P384L* ... S A N D **L** V F I P H H A F V D S I Y E ...

C

TYR genotype

Breeder designation	TYR^{Albino} / TYR^{Albino}	TYR^{Albino} / TYR^{D394G}	TYR^{D394G} / TYR^{D394G}	TYR^{Albino} / TYR^{P384L}	TYR^{D394G} / TYR^{P384L}	TYR^{P384L} / TYR^{P384L}	TYR^{Albino} / +	TYR^{P384L} / +	+ / +
Alb^{Albino} / Alb^{Albino}	25	8	2	1					
Alb^{Albino} / Alb^{Candy}	3			5	1				
Alb^{Albino} / Alb^{Toffee}				1					
Alb^{Candy} / Alb^{Candy}						3			
Alb^{Toffee} / Alb^{Toffee}						1			
Non-Albino							1	1	54

D

Wildtype
TYR^{Albino} / TYR^{Albino}
Non-Albino

3 kb

~400 bp

1 kb

0.5 kb

E

Albino haplotypes

TYR^{D394G} — *D394G*

TYR^{P384L} — *P384L*

TYR^{Albino} — No coding or splice-site variants

5 kb

— Tiled PCR amplifications to search for indels
* Longer allele of variable number tandem repeat (VNTR)
▶ ◀ Primers for panel D

A OCA2

Deletion in Lavender Albino

10 kb

1,514 bp   Deletion in Lavender Albino

Primers for panel B:

Primers for panel C:

B

Wildtype

Lavender Albino

3 kb — 1,789 bp

1 kb

0.5 kb — 275 bp

C

Wildtype

Lavender Albino

Het. Lavender Albino

3 kb

1 kb

0.5 kb — 429 bp

349 bp

D

OCA2 genotype

Breeder designation	*Deletion* / *Deletion*	*Deletion* / +	+ / +
Lavender Albino	14		
Het. Lavender Albino		1	
Non-Lavender Albino			76

E

Lavender Albino haplotype

OCA2^*Deletion*

Deletion

10 kb

1,514 bp

- - - - Hypothetical splicing between coding regions flanking deletion

▯ Coding regions out of frame compared to coding regions upstream of deletion

Phenotype	Color morph	Associated genetic variant(s)
Severe loss of melanin	Albino	Three alleles of *TYR*: • Missense variant *D394G* • Missense variant *P384L* • Haplotype *TYR^Albino*, which lacks coding variants and splice-site variants compared to wildtype
Moderate loss of melanin	Lavender Albino	One allele of *OCA2*: • Deletion of coding region 18
Mild loss of melanin	Ultramel	Two alleles of *TYRP1*: • Missense variant *R305H* • Putative deletion of coding regions 6 and 7