1

Mechanisms underlying microglial colonization of developing neural

retina in zebrafish

Nishtha Ranawat and Ichiro Masai*

Developmental Neurobiology Unit, Okinawa Institute of Science and Technology

Graduate University, Tancha 1919-1, Onna, Okinawa 904-0495, Japan

* Correspondence should be addressed: masai@oist.jp

Running title: Microglia colonization to retina

Key words: microglia, zebrafish, retina, blood vessel, neurogenesis

Word count of the main text (Introduction/Results/Discussion): 5,735

Abstract

Microglia are brain-resident macrophages that function as the first line of defense in

brain. Embryonic microglial precursors originate in peripheral mesoderm and migrate

into the brain during development. However, the mechanism by which they colonize the

brain is incompletely understood. The retina is one of the first brain regions to

accommodate microglia. In zebrafish, embryonic microglial precursors use intraocular

hyaloid blood vessels as a pathway to migrate into the optic cup via the choroid fissure.

Once retinal progenitor cells exit the cell cycle, microglial precursors associated with

hyaloid blood vessels start to infiltrate the retina preferentially through neurogenic

regions, suggesting that colonization of retinal tissue depends upon the neurogenic state.

Along with blood vessels and retinal neurogenesis, IL34 also participates in microglial

precursor colonization of the retina. Altogether, CSF receptor signaling, blood vessels,

and neuronal differentiation function as cues to create an essential path for microglial

migration into developing retina.

(150 words)

Introduction

Microglia are the resident macrophages of brain. These dedicated CNS phagocytes form the innate immune system of embryonic and adult brain. Microglia eliminate cellular debris to prevent neuro-inflammation and to promote neuronal protection in vertebrates (Ashwell, 1991; Caldero et al., 2009; Lawson et al., 1990; Neumann et al., 2009; Sierra et al., 2010). They also prune unnecessary synapses to establish functional, mature neural circuits during brain development, performing a variety of cellular functions (Paolicelli et al., 2011; Tremblay et al., 2010). In contrast to other CNS cells, like neurons and astrocytes, microglia do not originate from neural plate, but are derived from mesoderm (Ashwell, 1991; Boya et al., 1979) through hematopoiesis (Ginhoux et al., 2013). In developing zebrafish, embryonic hematopoiesis occurs in successive waves that are separated anatomically and temporally. The primitive or first wave of microglial precursors is generated from myeloid cells originating in the rostral blood island (RBI) at about 11 hours post-fertilization (hpf) (Stachura and Traver, 2011; Xu et al., 2012). The definitive wave is contributed by the ventral wall of the dorsal aorta (VDA), giving rise to hematopoietic stem cells (HSCs) (Xu et al., 2015). In addition, a short intermediate wave also originates from the posterior blood island (PBI) (Bertrand et al., 2007). After 2 weeks post-fertilization, VDA derived-microglia progressively replace RBI-derived microglia throughout the CNS (Ferrero et al., 2018; Xu et al., 2015). Thus, primitive and definitive hematopoiesis contribute embryonic and adult microglia, respectively, during zebrafish development.

Generation of embryonic microglial precursors and their colonization of brain areas has been extensively described in zebrafish (Herbomel et al., 2001). In zebrafish, embryonic microglial precursors are initially specified in lateral plate mesoderm and then spread on yolk. They start to migrate into the cephalic mesenchymal region after 22 hpf. At 26 – 30 hpf, a few microglia are observed in the vitreous space or choroid fissure of the optic cup, and around 30 microglia colonize the neural retina by 48 hpf. Microglial colonization of the optic tectum and other regions of zebrafish brain occurs after 48 hpf, indicating that the retina is one of the first brain regions to be colonized by microglia during development.

Previous studies have suggested various signals that promote microglial
colonization in brain. In mice, Cxcl12/CxcR4 signaling orchestrates microglial
migration into developing cerebral cortex (Arno et al., 2014; Hattori and Miyata, 2018).
In zebrafish, microglia migrate from the yolk-sac and colonize the brain in an apoptosis-
dependent manner (Casano et al., 2016; Xu et al., 2016). Microglial precursors also
migrate into the cephalic mesenchymal area in a Colony Stimulating Factor-Receptor
(CSF-R)-dependent manner (Herbomel et al., 2001; Wu et al., 2018). Zebrafish *fms*
mutants carry a genetic mutation in CSF-R and show severe delays in microglial
colonization of both brain and retina, as well as an increase in neuronal apoptosis
(Herbomel et al., 2001). Recently, it was reported that brain colonization by microglial
precursors depends primarily on one zebrafish CSF-R, CSF1ra, and one CSF-R ligand,
IL34, and that this combination of CSF ligand and receptor dominates this process (Wu
et al., 2018). Importantly, the number of microglia in the brain and retina is reduced in
zebrafish *il34* mutants that overexpress anti-apoptotic protein, Bcl2. Thus, apoptosis
and the IL34-CSF1ra signaling pathway cooperate to promote microglial colonization
of the brain and retina during zebrafish development.

In developing zebrafish retina, neurogenesis is initiated in the ventro-nasal
retina, adjacent to the optic stalk at 25 hpf and progresses to the whole region of the
neural retina, suggesting a spatio-temporal pattern of retinal neurogenesis in zebrafish
(Hu and Easter, 1999; Masai et al., 2000). Retinal progenitor cells are multipotent and
give rise to six major classes of neurons and one type of glial cells. Two types of
photoreceptors, rods, and cones, form the outer nuclear layer (ONL). Three interneurons,
amacrine cells, bipolar cells, and horizontal cells form the inner nuclear layer (INL).
Retinal ganglion cells (RGCs) form the RGC layer. Synaptic connections between
photoreceptors and bipolar/horizontal cells form the outer plexiform layer (OPL), and
synaptic connections between RGCs and bipolar/amacrine cells form the inner
plexiform layer (IPL). Cell fate determination is less dependent on the cell lineage of
retinal progenitor cells, suggesting that both extrinsic and intrinsic mechanisms
influence the status of retinal progenitor multipotency, leading to generation of diverse
retinal cell types (He et al., 2012). These developmental profiles of retinal neurogenesis
and cell differentiation may be coupled with microglial colonization. Although
apoptosis and CSF-R signaling are suggested in microglial colonization of the retina in

zebrafish (Wu et al., 2018), the mechanism underlying microglial colonization of the
retina remains to be determined.

In this study, using zebrafish, we examined the developmental profile of retinal
colonization by microglia precursors. The number of ocular microglial precursors
progressively increases from 32 to 54 hpf. Most microglial precursors do not proliferate,
suggesting that microglial colonization of the retina depends on cell migration from
outside the optic cup. We found three guidance mechanisms driving microglial
precursor colonization of the retina. First, IL34 initiates microglial precursor movement
from yolk toward the brain and the retina. Second, microglia precursors enter the optic
cup via ocular hyaloid blood vessels in the choroid fissure, suggesting that these blood
vessels guide microglia to the retina. Third, microglial precursors infiltrate the neural
retina preferentially through the neurogenic region, suggesting that the neurogenic state
of retinal tissue acts as an entry signal for microglial precursors to infiltrate the retina.
Thus, a series of guidance mechanisms promote microglial colonization from yolk to
the neural retina in zebrafish.

Results

Embryonic microglial precursors progressively colonize developing zebrafish retina

In zebrafish, early macrophages are generated from myeloid cells originating in the RBI around 11 hpf and these macrophages colonize the brain and retina by 55 hpf (Xu et al., 2015). Around 60 hpf, brain and retina-resident macrophages undergo a phenotypic transition, which indicates expression of mature microglial markers, such as apolipoprotein E (apo E) and phagocytic behavior toward dead cells (Herbomel et al., 2001). Importantly, early macrophages outside the brain never express apo E (Herbomel et al., 2001), suggesting that only brain and retina-resident macrophages give rise to microglia. Thus, early macrophages localized in the brain and retina by 60 hpf are generally accepted as microglial precursors in zebrafish. In this study, we focused on two macrophage markers, *macrophage expressing gene 1.1* (*mpeg1.1*) (Ellett et al., 2011) and *microfibrillar-associated protein 4* (*mfap4*) (Walton et al., 2015), and define *mpeg1.1*; *mfap4*-positive cells inside the optic cup as microglial precursors colonizing the zebrafish retina.

To ascertain how microglia precursors migrate from peripheral tissues into the neural retina during development, we generated a zebrafish transgenic line, *Tg[mpeg1.1:EGFP],* using the original DNA construct (Ellett et al., 2011). As previously reported (Ellett et al., 2011), our established transgenic line visualized ocular microglial precursors and enabled us to monitor their number and location in the optic cup from 24 to 54 hpf. Accordingly, we obtained 3D images using confocal laser scanning microscopy (LSM) (Figure 1A and 1B). The first microglial precursor cells appeared near the choroid fissure and lens around 30 – 32 hpf. After that, the number of ocular microglial precursors increased to 19.1±1.26 at 42 hpf and 31.0±4.44 at 54 hpf (Figure 1C), indicating a progressive increase in the number of ocular microglial precursors. We also confirmed a similar progressive increase in the number of ocular *L-plastin*-positive cells, although *L-plastin* is expressed in microglial precursors and neutrophils in zebrafish brain (Figure 1–figure supplement 1). Next, to determine more precisely the spatial distribution of microglial precursors in the optic cup, we generated another transgenic line, *Tg[mfap4:tdTomato-CAAX],* using the original DNA construct (Walton et al., 2015). As previously reported (Walton et al., 2015), our established

transgenic line efficiently labeled ocular microglial precursor membranes. We labeled this transgenic embryo using Bodipy ceramide conjugated with fluorescent Alexa-488, which visualizes retinal layer structures (Figure 1–figure supplement 2). From 32 – 36 hpf, mfap4$^+$ cells were mostly located in the vitreous space between the neural retina and lens, and possibly associated with ocular blood vessels, which develop around the lens. In 42 – 44-hpf retina, a few microglial precursor cells start to enter the neural retina and spread toward the emerging IPL, where they are associated with newly born amacrine cells (Figure 1–figure supplement 3). By 54 hpf, IPL formation is complete and microglial precursors were observed throughout all retinal tissue, except the OPL. Thus, microglial precursors enter the optic cup along the choroid fissure at 30 hpf, remain temporarily in the vitreous space between the lens and the retina, and then begin spreading into differentiating retinal tissue after 42 hpf.

Next, to evaluate the contribution of cell proliferation to the increasing number of ocular microglial precursors, we labeled ocular microglial precursors with markers of DNA replication. Here we used a zebrafish transgenic line, *Tg[EF1α: mCherry-zGem]* that specifically marks proliferative cells in S and G2 phases (Mochizuki et al., 2017; Mochizuki et al., 2014). We combined this *Tg[EF1α: mCherry-zGem]* system with *Tg[mpeg1.1:EGFP]* to calculate the fraction of proliferative microglial precursors undergoing S phase (Figure 1D and Video 1). First, we observed mCherry-zGem; mpeg1.1:EGFP double-positive cells in the peripheral tissue (Figure 1–figure supplement 4A-C) and found that more than 60% of mpeg1.1:EGFP-positive cells expressed mCherry-zGem (Figure 1–figure supplement 4D), confirming that this *Tg[EF1α: mCherry-zGem]* system works in early macrophages in zebrafish. However, in the retina, the fraction of mCherry-zGem; mpeg1.1:EGFP double-positive cells was less than 2% of all microglial precursors from 33.5 to 54 hpf (Figure 1E). Furthermore, more than 80% of mpeg1.1:EGFP-positive cells did not incorporate BrdU at 48 hpf (Figure 1F–1H), suggesting that a majority of ocular microglial precursors do not undergo S phase. Thus, microglial colonization of the retina mostly depends on cell migration from outside the optic cup.

Embryonic microglial precursor migration into the retina depends on blood vessels

The zebrafish retina receives its blood supply from two blood vessel systems, intraocular hyaloid blood vessels encapsulating the lens (Hartsock et al., 2014) and superficial choroidal blood vessels (Kaufman et al., 2015). Developing hyaloid blood vessels start to enter the space between the lens and retina through the ventral fissure at 18 – 20 hpf. Loop formation occurs around the lens at 24 – 28 hpf, and a branched hyaloid network forms after 35 hpf (Hartsock et al., 2014). Our live imaging showed that microglial precursors enter the optic cup through the choroid fissure and remain temporarily in the vitreous space between the lens and the retina before they infiltrate the neural retina (Figure 1—figure supplement 2). Furthermore, microglial precursors start to enter the optic cup after loop formation of hyaloid blood vessels is completed, suggesting a guiding role of blood vessels in microglial precursor colonization of the optic cup. To confirm whether microglial precursors entering the ocular space are associated with developing hyaloid blood vessels, we conducted time-lapse imaging of *Tg[kdrl:EGFP; mfap4-tdTomato-CAAX]* transgenic embryos, which visualizes endothelial cells of blood vessels (Jin et al., 2005) and ocular microglial precursors, respectively. The first microglial precursor was always associated with ocular hyaloid blood vessels around 30 hpf (Figure 2A) and moved along blood vessel surfaces (Video 2), so it is very likely that microglial precursors use blood vessels as a scaffold to enter the vitreous space between the lens and the neural retina. Microglial precursors move along hyaloid blood vessels in the ventral fissure, gradually leave vessel surfaces, and invade the neural retina through the basement membrane (Figure 2B, Figure 2–figure supplement 1, and Video 3).

Troponin T2A (tnnt2a; silent heart) is specifically expressed in heart and is essential for heart contraction (Sehnert et al., 2002). In zebrafish brain and mouse retina, haemodynamics drive blood vessel pruning, and loss of blood circulation causes blood vessel regression (Chen et al., 2012; Lobov et al., 2011; Yashiro et al., 2007). To examine whether the entry of microglial precursors into retina is altered upon blood vessel regression, we blocked blood circulation by injecting morpholino antisense oligos against tnnt2a (tnnt2a MO). When blood circulation is inhibited, ocular hyaloid blood vessels do not develop fully and microglial precursors are less likely to be associated with these thin blood vessels (Figure 2C). The number of ocular microglial precursors was significantly reduced at 36 hpf (Figure 2D), showing that microglial

colonization of the optic cup depends upon normal development of the blood vessel

network. This is in contrast to the case of microglial colonization of zebrafish midbrain

and optic tectum, which is independent of the blood vessel network (Xu et al., 2016).

Indeed, we confirmed that the number of microglial precursors in the optic tectum was

not significantly different between *tnnt2a* morphants and standard MO-injected

embryos at 72 hpf, although microglial precursor colonization of the optic tectum was

enhanced in *tnnt2a* morphants at 48 hpf (Figure 2–figure supplement 2).

Recent studies indicate that microglia facilitate ocular blood vessel development

(Checchin et al., 2006; Fantin et al., 2010; Rymo et al., 2011), and that macrophages

initiate a cell-death program in endothelial cells for blood vessel regression in

developing mouse retina (Lang and Bishop, 1993; Lobov et al., 2005). However, we

eliminated microglial precursors with morpholino antisense oligos against pu.1 (pu.1-

MO) or *interferon regulatory factor 8* (*irf8*) mutation (*irf8* gene knockdown causes

apoptosis of pu.1-positive myeloid cells) (Shiau et al., 2015), and confirmed that

microglial precursor elimination did not affect hyaloid blood vessel formation in

zebrafish at least by 48 hpf (Figure 2–figure supplement 3).

Microglial precursors infiltrate the neural retina preferentially through the

differentiating neurogenic area

In zebrafish, retinal neurogenesis occurs at the ventronasal retina adjacent to the optic

stalk at 25 hpf and propagates into the entire region of the neural retina at 33 hpf (Masai

et al., 2000). Microglial precursors start to migrate from the vitreous space into the

neural retina after 42 hpf, when the earliest differentiating retinal neurons, RGCs, start

to form the IPL (Mumm et al., 2006). To examine the role of retinal neurogenesis and

RGC differentiation in microglia precursor infiltration of the neural retina, we used two

transgenic lines, *Tg[EF1α: mCherry-zGem; mpeg1.1:EGFP]*, which enable us to

examine the relationship between microglial precursor migration and retinal progenitor

cells (Mochizuki et al., 2014). Live imaging of *Tg[EF1α: mCherry-zGem;*

mpeg1.1:EGFP] retinas at 42 and 48 hpf clearly showed that microglial precursors

avoid mCherry-zGem-positive proliferating regions and are preferentially positioned in

the region of mCherry-zGem-negative post-mitotic cells (Figure 3A–B, Figure 3–figure

supplement 1). The fraction of microglial precursors that infiltrated mCherry-zGem-

positive proliferating regions was 7.37% at 42 hpf and 6.13% at 48 hpf (Figure 3C), suggesting that >90% of microglial precursors infiltrate the retina through the mCherry-zGem-negative post-mitotic cell region. We also used another transgenic line *Tg[ath5:EGFP; mfap4-tdTomato-CAAX]*. In the *Tg[ath5:EGFP]* line, EGFP starts to be expressed in G2 phase of the final neurogenic cell division of retinal progenitor cells and is inherited by their daughter cells, which are negative for BrdU incorporation (Poggi et al., 2005; Yamaguchi et al., 2010), suggesting that ath5:EGFP specifically marks early differentiating retinal neurons. We conducted live imaging of *Tg[ath5:EGFP; mfap4-tdTomato-CAAX]* retinas at 36, 42, 48 hpf, and found that infiltration of mfap4-positive microglia preferentially occurs in the ath5:EGFP-positive region (Figure 3D). These data suggest that microglial precursors infiltrate the neural retina preferentially through the neurogenic area, raising the possibility that the neurogenic retinal region acts as a gateway through which microglial precursors move from the vitreous space into the neural retina.

Colonization of the optic tectum by microglial precursors depends on neuronal apoptosis in zebrafish (Casano et al., 2016; Xu et al., 2016). Therefore, it is still possible that microglial precursors preferentially infiltrate the neural retina through the neurogenic region, because of neuronal apoptosis. We inhibited retinal apoptosis by injecting morpholino antisense oligos against p53 (p53 MO) and confirmed that p53 MO effectively suppresses retinal apoptosis at 24 and 36 hpf (Figure 3–figure supplement 2). However, the number of microglial precursors did not differ between *p53* morphant retinas and standard-MO-injected retinas at 48 hpf (Figure 3–figure supplement 3AB), whereas the number of microglial precursors was significantly decreased in *p53* morphant optic tectum compared with standard-MO-injected optic tectum at 96 hpf (Figure 3–figure supplement 3CD). Thus, in contrast to microglial colonization of the optic tectum, neuronal apoptosis is not the major cue for microglial precursor colonization of the retina, at least prior to 54 hpf.

Neurogenesis acts as a gateway for microglial precursors to enter the retina

To confirm the possibility that the neurogenic retinal region functions as a gateway for microglial precursors to infiltrate the retina, we examined whether microglial precursor migration into the retina is compromised when retinal neurogenesis is affected.

Previously, we found that histone genesis slowed in zebrafish *stem loop binding protein*

1 (*slbp1*) mutants, leading to severe delays in retinal neurogenesis (Imai et al., 2014).

Our bulk RNAseq analysis confirmed that retinal neurogenesis and subsequent neuronal

differentiation were markedly delayed in zebrafish *slbp1* mutants (Figure 4—figure

supplement 1), such that ath5 expression spread into the entire *slbp1* mutant retina only

at 48 hpf, an event that occurs in wild-type retina at 33 hpf (Imai et al., 2014). We

combined *slbp1* mutants with transgenic lines *Tg[ath5:EGFP; mfap4: tdTomato-CAAX]*

and examined the number of ocular mfap4-tdTomato-CAAX-positive microglial

precursors (Figure 4A, Figure 4–figure supplement 2A and 3). In 48-hpf *slbp1* mutant

retinas, the number of ocular microglial precursors was 4.67 ± 2.42 (Figure 4A, B),

which is similar to the number in wild-type retinas at 32 hpf (Figure 1B), whereas the

number of ocular microglial precursors in wild-type siblings was 17.60 ± 5.13 at 48 hpf

(Figure 4A, B). To confirm whether the *slbp1* mutation interferes with genesis of early

macrophages, we examined peripheral mfap4$^+$ cells in the tail/trunk region of *slbp1*

mutants and wild-type sibling embryos. There was no significant difference in mfap4$^+$

cells between *slbp1* mutants and wild-type siblings in the trunk/tail region (Figure 4C,

D), indicating that the *slbp1* mutation does not influence early macrophage specification

in zebrafish embryos. Although inhibition of retinal apoptosis by p53 MO does not

influence microglial precursor colonization of the retina (Figure 3–figure supplement

3AB), we examined the level of retinal apoptosis in *slbp1* mutants. TUNEL revealed

that apoptosis was increased in *slbp1* mutant retinas compared with wild-type sibling

retinas (Figure 4–figure supplement 4). These data exclude the possibility that

decreased retinal apoptosis affects microglial precursor colonization of the retina in

slbp1 mutants, and again confirms that neuronal apoptosis is not the major cue for

microglial precursor colonization of the retina.

Mouse brain cortex colonization by microglia depends on the Cxcl12a-CxcR4

signaling axis (Arno et al., 2014). We previously reported that *cxcl12a* expression is

absent in the optic stalk of zebrafish *slbp1* mutants (Imai et al., 2014). To exclude the

possibility that the absence of *cxcl12a* expression in the optic stalk affects microglial

colonization of the retina in zebrafish *slbp1* mutants, we examined zebrafish *cxcl12a*

morphants. Injection of *cxcl12a*-MO at 500 µM, which effectively induces RGC axon

trajectory defects reported in zebrafish *odysseys* mutants carrying mutations in *cxcl12a*

receptor, *cxcr4b* (Li et al., 2005), did not affect the number of ocular microglial

precursors (Figure 4–figure supplement 5). Thus, Cxcl12a-CxcR4 signaling is not

involved in microglial colonization defects in *slbp1* mutants. We also confirmed that

elimination of microglial precursors with pu.1 MO did not affect the rate of retinal

neurogenesis or cell differentiation by 72 hpf (Figure 4–figure supplement 6).

We previously showed that overexpression of Notch1 intracellular domain

(NICD) suppresses retinal neurogenesis in zebrafish (Yamaguchi et al., 2005). We

confirmed that overexpression of NICD suppresses retinal neurogenesis in zebrafish by

injecting a DNA expression construct encoding UAS:myc-NICD (Scheer and Campos-

Ortega, 1999) into *Tg[hsp:gal4; ath5:EGFP]* double transgenic embryos (Figure 4-

figure supplement 7). Next, we examined whether microglial precursor infiltration of

the retina is compromised in retinas overexpressing NICD. We established a zebrafish

transgenic line, *Tg[rx1:gal4-VP16]*, which expresses Gal4-VP16 under control of a

retinal progenitor-specific promoter *rx1* (Chuang et al., 1999), and then injected two

DNA expression constructs encoding UAS:EGFP (Koster and Fraser, 2001) and

UAS:myc-NICD into *Tg[rx1:gal4-VP16; mfap4:tdTomato-CAAX]* double-transgenic

embryos. Embryos injected with only the DNA construct of UAS:EGFP served as a

positive control. We selected embryos in which EGFP was expressed in most retinal

cells at 24 hpf and used them for further analysis. The number of ocular microglial

precursors was significantly reduced in embryos overexpressing NICD and EGFP,

compared with control embryos overexpressing EGFP, at 44 hpf (Figure 4E, F, Figure

4–figure supplement 2B and 3). These data support the possibility that the retinal

neurogenic region functions as a gateway for microglia to infiltrate the retina.

The blockade of retinal neurogenesis delays differentiation of the first-born

retinal cell-type, RGCs. To examine whether blockade of RGC differentiation affects

microglial precursor colonization of the neural retina, we applied an antisense

morpholino against ath5 (known as atoh7) (ath5 MO). As with the zebrafish *ath5*

mutant, *lakritz* (Kay et al., 2001), RGC differentiation was specifically inhibited in *ath5*

morphant retinas (Figure 4–figure supplement 8A, B). In *ath5* morphants, the timing of

the first appearance of microglial precursors in the ocular vitreous space was not altered,

but the number of ocular microglial precursors was significantly decreased at 49 hpf

(Figure 4G, H, and Figure 4–figure supplement 2C, 3 and 8C), suggesting that RGC

differentiation or RGC-mediated IPL formation is required for microglial precursor
infiltration into the neural retina.

Microglial precursors preferentially associate with neurogenic retinal columns
To determine whether microglia precursors have greater affinity for differentiating
neurons than for retinal progenitor cells, we carried out two sets of experiments. First
we conducted cell transplant experiments using a wild-type donor line and an *slbp1*
mutant recipient line carrying *Tg[mfap4:tdTomato-CAAX]*. Wild-type donor cells were
transplanted into *slbp1* mutant recipient embryos at the blastula stage. We selected
slbp1 mutant and wild-type sibling embryos in which wild-type, donor retinal cell
columns were introduced in a mosaic manner at 48 hpf (Figure 5A). Host microglial
precursors and donor retinal cells were visualized with mfap4:tdTomato-CAAX and
Alexa-488 Dextran, respectively. In *slbp1* mutant host retinas, microglial precursors
were likely to be associated with donor wild-type retinal columns more frequently than
in wild-type host retinas (Figure 5B). To analyze these data statistically, we compared
eyes in which wild-type donors were transplanted into wild-type hosts with those in
which wild-type donors were transplanted into *slbp1* mutant hosts (Figure 5-figure
supplement 1AB). The fraction of microglial precursors associated with donor wild-type
retinal columns in total ocular microglial precursors was significantly higher in *slbp1*
mutant host retinas than in wild-type sibling host retinas at 48 hpf (Figure 5C),
suggesting that microglial precursors are more attracted by wild-type donor neurogenic
retinal columns than surrounding *slbp* mutant proliferative retinal cells. Since the
fraction of microglial precursors associated with donor retinal columns in total
microglial precursors may depend on the number of donor retinal columns incorporated
into host retinas, we next estimated trapping efficiency of microglial precursors per
donor column by dividing the fraction of microglial precursors associated with donor
columns with the transplanted donor column number in each eye (Figure 5–figure
supplement 1B). Trapping efficiency of microglial precursors per donor column was
significantly higher in *slbp1* mutant host retinas than in wild-type sibling host retinas
(Figure 5D), suggesting that microglial precursors are preferentially associated with
donor-derived wild-type retinal cells than with host-derived *slbp1* mutant retinal cells.

Second, we injected two DNA constructs encoding UAS:EGFP and

UAS:mycNICD into *Tg[hsp:gal4; mfap4:tdTomato-CAAX]* double-transgenic wild-

type embryos. Two rounds of heat-shock treatment at 18 and 30 hpf induced expression

of NICD and EGFP in a mosaic manner in the retina (Figure 5E). We examined the

fraction of *mfap4:tdTomato-CAAX*-positive microglial precursors associated with

EGFP-expressing retinal columns in the total number of *mfap4:tdTomato-CAAX*-

positive microglial precursors (Figure 5F). This fraction was significantly lower in

retinas overexpressing NICD and EGFP than in control retinas overexpressing only

EGFP (Figure 5G, Figure 5–figure supplement 1C). We also confirmed that trapping

efficiency of *mfap4:tdTomato-CAAX*-positive microglial precursors per EGFP-positive

retinal column was significantly lower in retinas overexpressing NICD and EGFP than

in retinas overexpressing only EGFP (Figure 5H, Figure 5–figure supplement 1C). Thus,

microglial precursors are less attracted by retinal columns in which neurogenesis is

arrested. Taken together, these data suggest that microglial precursors preferentially

associate with neurogenic retinal columns as opposed to proliferative retinal columns.

IL34 is involved in microglial precursor colonization of the retina

Recently, it was reported that microglial colonization of zebrafish brain, including retina,

depends on CSF-R, and that one of the CSF-R ligands, IL34, dominates this process

(Wu et al., 2018). In adult mouse retina, RGCs express IL34, which attracts one subset

of microglia and retains them around the IPL niche (O'Koren et al., 2019). First, we

confirmed that retinal cell differentiation proceeds normally until 72 hpf in zebrafish

il34 mutants, although pyknotic nuclei were stochastically observed in RGC and

amacrine cell layers (Figure 6-figure supplement 1). Next, we examined microglial

precursor colonization of the retina. The number of ocular microglial precursors was

significantly lower in *il34* homozygous mutants than in wild-type siblings at 34 hpf

(Figure 6–figure supplement 2) and 48 hpf (Figure 6A, B). Thus, consistent with the

previous report (Wu et al., 2018), IL34 is required for microglial precursor colonization

of the retina in zebrafish. However, *il34* mRNA expression is comparable in *slbp1*

mutant heads and wild-type sibling heads at 48 hpf (Figure 6–figure supplement 3),

suggesting that *il34* mRNA expression is not linked to retinal neurogenesis. Since the

number of ocular microglial precursors in *il34* homozygous mutants was zero at 34 hpf

(Figure 6–figure supplement 2) and no more than two, if any, at 48 hpf (Figure 6B), it is

likely that Csf1r-il34 signaling promotes microglial precursor movement from yolk to

the optic cup upstream of the blood vessel-mediated guidance mechanism (Figure 6C).

Discussion

In zebrafish, primitive microglia originate from the RBI, which is a hematopoietic tissue equivalent to mouse yolk sac, whereas definitive microglia are generated from hematopoietic stem cells that are specified in the VDA (Ferrero et al., 2018; Xu et al., 2015). Primitive and definitive waves of hematopoiesis generate embryonic and adult microglia, respectively. Using zebrafish as an animal model, several groups investigated microglial colonization from the periphery into developing brain, especially the optic tectum, which is part of the midbrain (Casano et al., 2016; Herbomel et al., 2001; Svahn et al., 2013; Wu et al., 2018; Xu et al., 2016). Colonization of the optic tectum by microglial precursors depends on neuronal apoptosis, probably through attraction by an apoptotic cell-secreted phospholipid, lysophosphatidylcholine (LPC) (Casano et al., 2016; Xu et al., 2016). In addition, microglial colonization of brain is CSF receptor-dependent (Herbomel et al., 2001; Wu et al., 2018). In mice, microglial colonization of brain requires functional blood circulation (Ginhoux et al., 2010). However, in zebrafish, microglial colonization of the optic tectum is independent of blood circulation (Xu et al., 2016). A series of elegant studies revealed the molecular network that promotes microglial colonization of the midbrain. However, it remains to be seen whether this mechanism fully explains colonization of other brain regions by microglial precursors. In this study, we focused on zebrafish retina and investigated the mechanism that regulates migration of embryonic microglial precursors into developing retina.

We first conducted live imaging of zebrafish microglial precursors from 24 to 54 hpf. Microglial precursors progressively increase in number during embryonic development. Interestingly, almost all microglial precursors enter the optic cup through the choroid fissure. However, peripheral macrophages located in the mesenchymal region between the eye and the brain did not enter the optic cup across the ciliary marginal zone. This may be consistent with the observation that these peripheral macrophages never enter the retina following rod cell death (White et al., 2017), suggesting a functional difference between peripheral macrophages and ocular microglia. Next, we found that the majority of ocular microglial precursors do not undergo S phase and are probably in G1 phase. Thus, the increase of ocular microglial precursors is due to migration from outside the eye. In developing mouse retina, microglial precursors appear from the vitreous area near the optic disk at E11.5,

progressively increase in number, and then infiltrate the neural retina. These retinal

microglia were also negative for a proliferative marker, Ki67 (Santos et al., 2008),

suggesting that mouse embryonic retinal microglia are also non-proliferative.

Another interesting finding is that entry of microglial precursors into the optic

cup through the choroid fissure depends on ocular blood vessels. We observed that

migrating microglial precursors are closely associated with hyaloid blood vessels after

loop formation. These microglial precursors pass along these vessels, which traverse the

choroid fissure and surround the posterior region of the lens. Furthermore, the number

of ocular microglial precursors was reduced when blood circulation was blocked. Since

inhibition of blood circulation compromises the structural integrity of blood vessels in

zebrafish, we conclude that ocular blood vessel formation is required for microglial

precursor entry into the optic cup through the choroid fissure. One possibility is that

blood vessels function as a path upon which microglial precursors enter the optic cup.

Membrane proteins or extracellular matrix proteins on blood endothelial cells may

facilitate the association of microglial precursors with blood vessel surfaces.

Alternatively, substances that attract microglial precursors may be released from

hyaloid blood endothelial cells. Previous studies on human and murine microglia

demonstrated that microglial colonization of the retina takes place prior to retinal

vascularization, and that microglia facilitate ocular blood vessel development (Checchin

et al., 2006; Fantin et al., 2010; Rymo et al., 2011). Macrophages initiate endothelial

cell death for blood vessel regression in developing mouse retina (Lang and Bishop,

1993; Lobov et al., 2005). However, in contrast to mammals, elimination of microglia

by pu.1 MO or *irf8* mutation did not affect ocular blood vessel formation, suggesting

that microglia do not regulate ocular blood vessel formation in zebrafish. Interestingly,

classic histological studies on mouse retinas showed that early emerging ocular

microglia are associated with the hyaloid artery (Hume et al., 1983; Santos et al., 2008),

which is located in the vitreous area and regresses in later stages before retinal

vasculature formation (Ito and Yoshioka, 1999). Thus, further investigation will be

necessary to determine whether the hyaloid artery guides microglial precursors into the

optic cup in vertebrate species such as mice. In zebrafish, colonization of the optic

tectum by microglia is independent of blood circulation (Xu et al., 2016). We confirmed

that the number of microglia in the optic tectum did not differ between *tnnt2a*

morphants and control embryos at 72 hpf; however, microglial colonization of the optic

tectum was enhanced and microglial shape was round rather ramified in *tnnt2a*

morphants at 48 hpf. Further study will be necessary to clarify the role of blood

circulation in microglial colonization of the optic tectum.

After 42 hpf, microglial precursors detach from hyaloid blood vessels and start

to infiltrate the neural retina. Interestingly, we found that more than 90% of microglial

precursors enter the neural retina through the neurogenic area. Indeed, the number of

microglial precursors is reduced in *slbp1* mutant retinas and NICD-overexpressing

retinas, in both of which retinal neurogenesis is severely delayed. Furthermore, we

conducted two sets of experiments: the first was cell transplantation from wild-type

donor cells into *slbp1* mutant host retinas, which introduced neurogenic wild-type

retinal cell columns in proliferative *slbp1* mutant retinas, and the second was

overexpression of NICD in wild-type retina, which introduced proliferative retinal cell

columns in neurogenic retinas. Consistently, in both cases, microglial precursors were

preferentially associated with neurogenic retinal cell columns. Thus, neurogenesis is

required for infiltration of microglial precursors into the neural retina after 42 hpf. We

observed that the number of microglial precursors is diminished in *ath5* morphant

retinas, suggesting that RGCs are required for infiltration of microglial precursors into

the neural retina. There are at least three possible mechanisms for this infiltration. First,

the basal region of retinal progenitor cells may function as a physical barrier that

inhibits microglial precursor infiltration of the neural retina. Second, microglial

precursors may be attracted to surfaces of differentiating retinal neurons or RGCs. Third,

differentiating retinal neurons or RGCs may release a specific attractant for microglia.

There are several candidate molecules that suggest the third possibility. In adult mice,

RGCs express IL34, which attracts microglia and retains them around the IPL niche

(O'Koren et al., 2019). Indeed, microglial colonization of zebrafish brain depends on

CSF-R, and one of the CSF-R ligands, IL34, dominates this process (Wu et al., 2018).

We confirmed that microglial precursor colonization of retina is severely affected in

il34 mutants. However, *il34* mRNA expression is comparable in *slbp1* mutants and their

wild-type siblings, suggesting that IL34 is not linked to neurogenesis-mediated

microglial precursor infiltration. Rather, the number of ocular microglial precursors in

il34 mutants was almost zero at 34 and 48 hpf, so it is very likely that Csf1r-il34

signaling initiates microglial precursor movement from yolk toward brain and retina,
followed by blood vessel- and neurogenesis-mediated guidance.

It was reported that apoptosis attracts microglia in zebrafish developing brain
(Casano et al., 2016; Xu et al., 2016). However, microglial precursor colonization of the
retina is normal in zebrafish *p53* morphants, suggesting that apoptosis does not promote
microglial precursor colonization of the retina. Why are microglial precursors
insensitive to retinal apoptosis? We found that apoptosis is enhanced in zebrafish *slbp*
mutant retinas, in which microglial precursor colonization is severely affected due to a
delay of retinal neurogenesis. It is likely that spontaneous apoptotic cells fail to be
eliminated because of the reduced number of microglial precursors in *slbp* mutant
retinas; however, interestingly, these increased dead cells did not promote microglial
precursor infiltration into *slbp* mutant retinas, suggesting that neurogenesis primarily
opens the gate through which microglial precursors enter the neural retina. Since retinal
neurogenesis normally occurs from 24 to 48 hpf in zebrafish, microglial precursors
could not be attracted by apoptosis without the infiltration path opened by neurogenesis
before 48 hpf. Further studies will be necessary to unveil the molecular mechanism
underlying microglial infiltration into neural retina.

In summary, there are three mechanisms for microglial colonization of
developing zebrafish retina (Figure 6C). IL34-CSF-R signaling initiates microglial
precursor movement from yolk toward brain and retina. Microglial precursors further
use ocular hyaloid blood vessels as a pathway to enter the optic cup and then infiltrate
the neural retina preferentially through the neurogenic region. In the future, it remains to
identify molecules involved in blood vessel- and neurogenesis-mediated guidance
mechanisms, and to assess whether these mechanisms are used for microglial
colonization of other brain regions in other vertebrate species.

Materials and Methods

Fish strains

Zebrafish (*Danio rerio*) were maintained using standard procedures (Westerfield, 1993). RIKEN wako (RW) was used as a wild-type strain for mutagenesis (Masai et al., 2003). *slbp1*[rw440] (Imai et al., 2014), *irf8*[st96] (Shiau et al., 2015) and *il34*[hkz11] (Wu et al., 2018) were used. Transgenic lines *Tg[ath5:EGFP]*[rw021] were used to monitor *ath5* gene expression (Masai et al., 2005). *Tg[EF1α:mCherry-zGem]*[oki011] (Mochizuki et al., 2014) was used for visualization of cell-cycle phases. *Tg[mfap4:tdTomato-CAAX]*[oki058] and *Tg[mpeg1.1:EGFP]*[oki053] were used to visualize microglial precursors. *Tg[kdrl:EGFP]*[s843Tg] was employed to visualize blood vessels (Jin et al., 2005). *Tg[hsp:gal4]*[kca4] (Scheer et al., 2002) and *Tg[rx1:gal4-VP16]*[oki065] were used for UAS-mediated expression of target genes. For confocal scanning, embryos were incubated with 0.003% phenyltiourea (PTU) (Nacalai tesque, 27429-22) to prevent melanophore pigmentation. The zebrafish pigmentation mutant, *roy orbison (roy)* (D'Agati et al., 2017) was used to remove iridophores.

Ethics statement

All zebrafish experiments were performed in accordance with the Animal Care and Use Program of Okinawa Institute of Science and Technology Graduate University (OIST), Japan, which is based on the Guide for the Care and Use of Laboratory Animals by the National Research Council of the National Academies. The OIST animal care facility has been accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC International). All experimental protocols were approved by the OIST Institutional Animal Care and Use Committee.

Establishment of *Tg[mpeg1.1:EGFP]* and *Tg[mfap4:tdTomato-CAAX]* transgenic lines

The DNA construct encoding mpeg1.1:EGFP was kindly provided by Dr. Graham Lieschke and we are indebted to Dr. David Tobin for the construct encoding mfap4-tdTomato-CAAX. These DNA constructs were injected into fertilized eggs with Tol2 transposase mRNA, to establish transgenic lines, *Tg[mpeg1.1:EGFP]* and *Tg[mfap4:tdTomato-CAAX]* in our lab.

Histology

Plastic sectioning and immunolabeling of cryosections were carried out as previously described (Masai et al., 2003). Anti-GFP (Themo Fisher Scientific, A11122), anti-myc-tag (Invitrogen, R950-25), zn5 (Oregon Monoclonal Bank) and zpr1 (Oregon Monoclonal Bank) antibodies were used at 1:200; 1:250, 1:100, and 1:100 dilutions, respectively. For detection of BrdU incorporation, BrdU (Nacalai, tesque, 05650-95) was applied to 52-hpf embryos, chased for 2 h at 28.5°C and fixed with 4% paraformaldehyde (PFA). Labeling of retinal sections with anti-BrdU antibody (BioRad, MCA2060) was carried out as previously described (Yamaguchi et al., 2005). TUNEL was performed using an In Situ Cell Death Detection Kit (Roche, 11684795910). Bodipy-ceramide (Thermo Fisher Scientific, B22650) was applied to visualize retinal layers as previously described (Masai et al., 2003). Nuclear staining was performed using 1nM TOPRO3 (Thermo Fisher Scientific, T3605).

Morpholino

Morpholino antisense oligos were designed as shown below.

tnnt2a MO: 5'-CATGTTTGCTCTGATCTGACACGCA-3' (Sehnert et al., 2002)

p53 MO: 5'-GCGCCATTGCTTTGCAAGAATTG-3' (Langheinrich et al., 2002)

cxcl12a MO: 5'-ACTTTGAGATCCATGTTTGCAGTG-3'(Li et al., 2005)

pu.1 MO: 5'-GATATACTGATACTCCATTGGTGGT-3'(Rhodes et al., 2005)

ath5 MO: 5'-TTCATGGCTCTTCAAAAAAGTCTCC-3'

Standard MO: 5'-CCTCTTACCTCAGTTACAATTTATA-3'

Morpholino antisense oligos were injected into fertilized eggs at 500 μM for tnnt2a MO and cxcl12a MO; 250 μM for ath5 MO and pu.1 MO and 100 μM for p53 MO. The same concentration was used for Standard MO in each MO experiment.

Cell transplantation

Cell transplantation was performed as previously described (Masai et al., 2003). Wild-type zygotes were injected with Alexa-488 dextran (Thermo Fisher Scientific, D22910) and used for donor embryos. *slbp1* mutant embryos carrying *Tg[mfap4-tdTomato-CAAX]* were used as host embryos. Host embryos carrying donor retinal cells were

selected by observing Alexa 488 fluorescence at 24 hpf. *slbp1* mutant and wild-type

sibling embryos were sorted based on the *slbp1* mutant morphological phenotype at 48

hpf and used for live imaging. After confocal images were obtained, the number of

ocular mfap4-positive microglial precursors associated with Alexa-488 dextran-labeled

donor transplanted retinal columns was counted. The fraction of ocular mfap4-positive

microglial precursors associated with donor transplanted retinal columns in total ocular

microglial precursors was calculated. The trapping efficiency of ocular mfap4-positive

microglial precursors per transplanted donor retinal column was calculated using the

total number of donor transplanted retinal columns in the retina. Detailed information

on each transplanted eye is shown in Figure 5-supplement 1AB.

Live Imaging and Analyses

Transgene   lines   *Tg[mpeg1.1:EGFP]*   or   *Tg[mfap4:tdTomato-CAAX]*,   and

Tg[kdrl:EGFP], were used for time-lapse imaging of microglial precursors and blood

vessels. 3D confocal images were obtained using a confocal LSM, LSM710 (Zeiss) or

an FV3000RS (Olympus), and analyzed using ImageJ (2.0.0-rc-69/1.52p) and Imaris

software (ver.9.1.2 Bitplane). The DNA construct encoding Ptf1a:EGFP was used for

visualizing amacrine cells or their progenitors (Jusuf and Harris, 2009).

RNA extraction

Heads of 48 hpf wild-type sibling and *slbp1* mutant embryos were dissected and

transferred to 100 μL Sepasol (Nacalai tesque, 09379) on ice. Heads were then

homogenized using a hand homogenizer (~20 pulses). Twenty μL $CHCl_3$ were then

added to samples and mixed gently. After centrifugation at 15,000 g for 15 min, the

aqueous phase was collected and mixed with 100 μL isopropanol. One μL of RNase-

free glycogen (Nacalai tesque 11170-11, 20 mg/mL) was added to all samples to

increase the yield. After incubating at room temperature for 10 min, samples were

centrifuged at 15,000 g at 4°C for 15 min. Supernatant was removed and the pellet was

washed 3 times with 500 μL of 75% ethanol and centrifuged at 8,000 g at 4°C. The

pellet was then resuspended in a desired amount of nuclease-free water and stored at –

80°C. RNA concentration and purity of samples were determined using a Nanodrop.

RNA sequencing and analysis

RNA samples with RIN >7 were subjected to paired-end sequencing using an Illumina HiSeq4000. First, a quality check was performed using FastQC and read trimming was done with Trimomatic (Bolger et al., 2014). PRINSEQ lite (Schmieder and Edwards, 2011) was used for PolyA trimming and quality filtering. Trimmed sequences were then mapped to the zebrafish reference genome (GRCz11) using hisat2.1.0 (Kim et al., 2019) and mapped reads are counted using featureCounts (Liao et al., 2014). With the R package, EdgeR (Robinson et al., 2010), differentially expressed genes with $Log_2FC >$ |2| and FDR values < 0.01 were extracted. EnhancedVolcano package (https://github.com/kevinblighe/EnhancedVolcano) was used to draw volcano plots. A heat map was generated with the pheatmap package (version 1.10.12) (https://cran.r-project.org/web/packages/pheatmap/index.html).

Evaluation of *il34* mRNA expression by semi-quantitative PCR

Extracted RNA from 48-hpf wild-type sibling and *slbp1* mutant heads was used to prepare cDNA, using ReverTra Ace® qPCR RT master mix with gDNA remover (Toyobo, FSQ-301). The expression level of *il34* mRNA was evaluated with quantitative PCR using the primers below. mRNA of cytoplasmic actin β2, namely *actb2* (ZFIN), was used for normalization.

Forward primer for *il34* mRNA: 5'- tggtccagtccgaatgct-3'

Reserve primer for *il34* mRNA: 5'- gctgcactactgcacactgg -3'

Forward primer for *actb2* mRNA: 5'- tgtcttcccatccatcgtg -3'

Reserve primer for *actb2* mRNA: 5'- tgtcttcccatccatcgtg-3'

Mosaic expression of NICD in retinal cells using *Tg[rx1:gal4-VP16]* and *Tg[hsp:gal4]* transgenic lines

The DNA fragment that covers a 2,892-bp genomic region upstream from the start codon of *rx1* cDNA (Chuang et al., 1999), was amplified by PCR and inserted between *XhoI* and *BamHI* sites of the Tol2 base expression vector, pT2AL200R150G (Urasaki et al., 2006). Next, DNA fragments encoding *gal4-VP16* (Koster and Fraser, 2001) were further inserted between *BamHI* and *ClaI* sites of pT2AL200R150G to fuse the *rx1*

promoter. The plasmid was injected with Tol2 transposase mRNA into fertilized eggs of

the UAS:EGFP transgenic line to establish a transgenic line, *Tg[rx1:gal4-VP16]^{oki065}*. A

mixture of plasmids of UAS:EGFP (Koster and Fraser, 2001) and UAS:myc-NICD

(Scheer and Campos-Ortega, 1999) (each 10 ng/μL) were injected into fertilized eggs of

the *Tg[mfap4:tdTomato-CAAX*; *rx1:gal4-VP16]* or *Tg[mfap4:tdTomato-CAAX*;

hsp:gal4] transgenic line. In the case of the *Tg[mfap4:tdTomato-CAAX*; *hsp:gal4]*

transgenic line, two rounds of heat shock at 37°C for 1 h were applied at 18 and 30 hpf.

Embryos expressing EGFP in the optic cup were selected at 24 hpf, fixed with PFA at

48 hpf and used to prepare serial retinal sections for imaging analysis. After confocal

images were obtained, the number of ocular mfap4-positive microglial precursors

associated with EGFP-expressing columns was counted. The fraction of ocular mfap4-

positive microglial precursors associated with EGFP-expressing columns in total

microglial precursors was calculated and the trapping efficiency of ocular mfap4-

positive microglial precursors per EGFP-expressing column was calculated using the

total number of EGFP-expressing columns in the retina. Detailed information on each

injected eye is shown in Figure 5-figure supplement 1C. To confirm that NICD inhibits

retinal neurogenesis, UAS:myc-NICD or UAS:mCherry (each 10 ng/μL) was injected

into zebrafish transgenic embryos *Tg[ath5:EGFP; hsp:gal4]*. Three rounds of heat

shock at 37°C for 1 h were applied at 18, 24 and 30 hpf. Embryos were fixed at 36 hpf

and labeled with anti-myc tag antibody to visualize myc-NICD expressing retinal cells

with Alexa-543-conjugated secondary antibody. Whole retinas were used for confocal

scanning with an FV3000 (Olympus). Controls were UAS:mCherry-injected samples

and used directly for live confocal scanning. Confocal 3D retinal images were used to

count the number of ath5:EGFP-positive and negative retinal columns in myc-NICD or

mCherry expressing retinal columns from 5 independent embryos.

Evaluation of microglial precursor colonization of the retina in *il34* mutants.

The *il34^{hkz11}* allele (Wu et al., 2018) was combined with the *Tg[mfap4:tdTomato]*

transgenic line and used for analysis. Embryos were generated by pair-wise crosses

between heterozygous mutant male and female fish, and maintained with N-phenyl

thiourea (PTU)-containing water to prevent melanophore pigmentation. Whole retinas

of 19 embryo at 34 hpf and 29 embryos at 48 hpf were scanned with confocal
microscopy, using an LSM710 (Zeiss) or an FV3000RS (Olympus). Embryos were
fixed with 4% PFA and used for genotyping. A DNA fragment containing the 4-base
deletion mutation of the *il34^{hkz11}* allele was amplified by PCR and sequenced to
determine genotypes. Primers used for PCR amplification and sequencing are below.
Forward primer for PCR: 5'-tgcaattaaacagccaatgtg-3'
Reverse primer for PCR: 5'-ctgagtcacagccctcaaatc-3'
Forward primer for sequencing: 5'-ccatttgtttttacctgaccaaa-3'
Reverse primer for sequencing: 5'-gctaattggtgtgggacgtt-3'
Using the surface rendering tool of Imaris software (Bitplane, ver.9.1.2), we eliminated
signals of iridophore-derived noise or peripheral microphages around the optic cup and
extracted only ocular microglial precursors. The number of ocular microglial precursors
was counted in each retina and compared between genotypes.
705

Statistical analysis
Statistical analyses were performed using GraphPad Prism version 8.2.1. Statistical
significance was determined using two-tailed unpaired Student's t-tests for Fig. 1G;
Figs. 2D; 4BDH; 5CDGH; Fig.1-figure supplement 2; Fig.2-figure supplement 2B;
Fig.3-figure supplement 3BD; Fig.4-figure supplement 4B; Fig.4-figure supplement 5D;
Fig.6-figure supplement 3, Tukey's multiple comparison test for Figs.4F; 6B; Fig.6-
figure supplement 2B, and Bonferroni's multiple comparison test for Fig.3-figure
supplement 2B. Chi square tests were used for Fig. 4-figure supplement 7C. Detailed
information on each dataset is provided in Excel files in Raw data.
715

716　**Data availability**
717　Raw RNA-seq dataset of *slbp1* mutant and wild-type sibling is available at Gene
718　Expression Omnibus (GSE144517).
719

Acknowledgements

We thank Graham Lieschke for DNA constructs encoding mepg1.1:EGFP, David Tobin

for DNA constructs encoding mfap4-tdTomato-CAAX, Francesco Argenton for DNA

construct encoding Ptf1a:EGFP, William Talbot for zebrafish *irf8* mutant line, Zilong

Wen for zebrafish *il34* mutant line, and José Campos-Ortega for DNA constructs

encoding UAS-myc tagged NICD. We also thank lab members, especially Yuko

Nishiwaki, Yuki Takeuchi, Yutaka Kojima, Jeff Liner, Mamoru Fujiwara, and Tetsuya

Harakuni for supporting experiments. We thank Steven D. Aird for editing the

manuscript.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by a grant from the Okinawa Institute of Science and

Technology Graduate University to IM.

References
739
Arno, B., Grassivaro, F., Rossi, C., Bergamaschi, A., Castiglioni, V., Furlan, R.,
Greter, M., Favaro, R., Comi, G., Becher, B., et al. (2014). Neural progenitor cells
orchestrate microglia migration and positioning into the developing cortex. *Nat*
Commun **5**.

Ashwell, K. (1991). The distribution of microglia and cell death in the fetal rat
forebrain. *Brain Res Dev Brain Res* **58**, 1-12.

Bertrand, J. Y., Kim, A. D., Violette, E. P., Stachura, D. L., Cisson, J. L. and
Traver, D. (2007). Definitive hematopoiesis initiates through a committed
erythromyeloid progenitor in the zebrafish embryo. *Development* **134**, 4147-4156.

Bolger, A. M., Lohse, M. and Usadel, B. (2014). Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinformatics* **30**, 2114-2120.

Boya, J., Calvo, J. and Prado, A. (1979). The origin of microglial cells. *Journal of*
anatomy **129**, 177-186.

Caldero, J., Brunet, N., Ciutat, D., Hereu, M. and Esquerda, J. E. (2009).
Development of microglia in the chick embryo spinal cord: implications in the
regulation of motoneuronal survival and death. *J Neurosci Res* **87**, 2447-2466.

Casano, A. M., Albert, M. and Peri, F. (2016). Developmental Apoptosis Mediates
Entry and Positioning of Microglia in the Zebrafish Brain. *Cell Rep* **16**, 897-906.

Checchin, D., Sennlaub, F., Levavasseur, E., Leduc, M. and Chemtob, S. (2006).
Potential role of microglia in retinal blood vessel formation. *Invest Ophthalmol Vis Sci*
47, 3595-3602.

Chen, Q., Jiang, L., Li, C., Hu, D., Bu, J. W., Cai, D. and Du, J. L. (2012).
Haemodynamics-driven developmental pruning of brain vasculature in zebrafish. *PLoS*
Biol **10**, e1001374.

Chuang, J. C., Mathers, P. H. and Raymond, P. A. (1999). Expression of three Rx
homeobox genes in embryonic and adult zebrafish. *Mech Dev* **84**, 195-198.

D'Agati, G., Beltre, R., Sessa, A., Burger, A., Zhou, Y., Mosimann, C. and White,
R. M. (2017). A defect in the mitochondrial protein Mpv17 underlies the transparent
casper zebrafish. *Dev Biol* **430**, 11-17.

Ellett, F., Pase, L., Hayman, J. W., Andrianopoulos, A. and Lieschke, G. J. (2011).
mpeg1 promoter transgenes direct macrophage-lineage expression in zebrafish. *Blood*
117, e49-56.

Fantin, A., Vieira, J. M., Gestri, G., Denti, L., Schwarz, Q., Prykhozhij, S., Peri, F.,
Wilson, S. W. and Ruhrberg, C. (2010). Tissue macrophages act as cellular

chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell
induction. *Blood* **116**, 829-840.

Ferrero, G., Mahony, C. B., Dupuis, E., Yvernogeau, L., Di Ruggiero, E.,
Miserocchi, M., Caron, M., Robin, C., Traver, D., Bertrand, J. Y., et al. (2018).
Embryonic Microglia Derive from Primitive Macrophages and Are Replaced by cmyb-
Dependent Definitive Microglia in Zebrafish. *Cell Rep* **24**, 130-141.

Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M. F.,
Conway, S. J., Ng, L. G., Stanley, E. R., et al. (2010). Fate mapping analysis reveals
that adult microglia derive from primitive macrophages. *Science* **330**, 841-845.

Ginhoux, F., Lim, S., Hoeffel, G., Low, D. and Huber, T. (2013). Origin and
differentiation of microglia. *Front Cell Neurosci* **7**, 45.

Hartsock, A., Lee, C., Arnold, V. and Gross, J. M. (2014). In vivo analysis of hyaloid
vasculature morphogenesis in zebrafish: A role for the lens in maturation and
maintenance of the hyaloid. *Dev Biol* **394**, 327-339.

Hattori, Y. and Miyata, T. (2018). Microglia extensively survey the developing cortex
via the CXCL12/CXCR4 system to help neural progenitors to acquire differentiated
properties. *Genes Cells* **23**, 915-922.

He, J., Zhang, G., Almeida, A. D., Cayouette, M., Simons, B. D. and Harris, W. A.
(2012). How variable clones build an invariant retina. *Neuron* **75**, 786-798.

Herbomel, P., Thisse, B. and Thisse, C. (2001). Zebrafish early macrophages colonize
cephalic mesenchyme and developing brain, retina, and epidermis through a M-CSF
receptor-dependent invasive process. *Dev Biol* **238**, 274-288.

Hu, M. and Easter, S. S. (1999). Retinal neurogenesis: the formation of the initial
central patch of postmitotic cells. *Dev Biol* **207**, 309-321.

Hume, D. A., Perry, V. H. and Gordon, S. (1983). Immunohistochemical localization
of a macrophage-specific antigen in developing mouse retina: phagocytosis of dying
neurons and differentiation of microglial cells to form a regular array in the plexiform
layers. *J Cell Biol* **97**, 253-257.

Imai, F., Yoshizawa, A., Matsuzaki, A., Oguri, E., Araragi, M., Nishiwaki, Y. and
Masai, I. (2014). Stem-loop binding protein is required for retinal cell proliferation,
neurogenesis, and intraretinal axon pathfinding in zebrafish. *Dev Biol* **394**, 94-109.

Ito, M. and Yoshioka, M. (1999). Regression of the hyaloid vessels and pupillary
membrane of the mouse. *Anat Embryol (Berl)* **200**, 403-411.

Jin, S. W., Beis, D., Mitchell, T., Chen, J. N. and Stainier, D. Y. (2005). Cellular and
molecular analyses of vascular tube and lumen formation in zebrafish. *Development* **132**,
5199-5209.

Jusuf, P. R. and Harris, W. A. (2009). Ptf1a is expressed transiently in all types of
amacrine cells in the embryonic zebrafish retina. *Neural Dev* **4**, 34.

Kaufman, R., Weiss, O., Sebbagh, M., Ravid, R., Gibbs-Bar, L., Yaniv, K. and
Inbal, A. (2015). Development and origins of zebrafish ocular vasculature. *BMC Dev*
Biol **15**, 18.

Kay, J. N., Finger-Baier, K. C., Roeser, T., Staub, W. and Baier, H. (2001). Retinal
ganglion cell genesis requires lakritz, a Zebrafish atonal Homolog. *Neuron* **30**, 725-736.

Kim, D., Paggi, J. M., Park, C., Bennett, C. and Salzberg, S. L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*
37, 907-915.

Koster, R. W. and Fraser, S. E. (2001). Tracing transgene expression in living
zebrafish embryos. *Dev Biol* **233**, 329-346.

Lang, R. A. and Bishop, J. M. (1993). Macrophages are required for cell death and
tissue remodeling in the developing mouse eye. *Cell* **74**, 453-462.

Langheinrich, U., Hennen, E., Stott, G. and Vacun, G. (2002). Zebrafish as a model
organism for the identification and characterization of drugs and genes affecting p53
signaling. *Curr Biol* **12**, 2023-2028.

Lawson, L. J., Perry, V. H., Dri, P. and Gordon, S. (1990). Heterogeneity in the
distribution and morphology of microglia in the normal adult mouse brain.
Neuroscience **39**, 151-170.

Li, Q., Shirabe, K., Thisse, C., Thisse, B., Okamoto, H., Masai, I. and Kuwada, J. Y.
(2005). Chemokine signaling guides axons within the retina in zebrafish. *J Neurosci* **25**,
1711-1717.

Liao, Y., Smyth, G. K. and Shi, W. (2014). featureCounts: an efficient general
purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**,
923-930.

Lobov, I. B., Cheung, E., Wudali, R., Cao, J., Halasz, G., Wei, Y., Economides, A.,
Lin, H. C., Papadopoulos, N., Yancopoulos, G. D., et al. (2011). The Dll4/Notch
pathway controls postangiogenic blood vessel remodeling and regression by modulating
vasoconstriction and blood flow. *Blood* **117**, 6728-6737.

Lobov, I. B., Rao, S., Carroll, T. J., Vallance, J. E., Ito, M., Ondr, J. K., Kurup, S.,
Glass, D. A., Patel, M. S., Shu, W., et al. (2005). WNT7b mediates macrophage-
induced programmed cell death in patterning of the vasculature. *Nature* **437**, 417-421.

Masai, I., Lele, Z., Yamaguchi, M., Komori, A., Nakata, A., Nishiwaki, Y., Wada,
H., Tanaka, H., Nojima, Y., Hammerschmidt, M., et al. (2003). N-cadherin mediates
retinal lamination, maintenance of forebrain compartments and patterning of retinal
neurites. *Development* **130**, 2479-2494.

Masai, I., Stemple, D. L., Okamoto, H. and Wilson, S. W. (2000). Midline signals
regulate retinal neurogenesis in zebrafish. *Neuron* **27**, 251-263.

Masai, I., Yamaguchi, M., Tonou-Fujimori, N., Komori, A. and Okamoto, H.
(2005). The hedgehog-PKA pathway regulates two distinct steps of the differentiation
of retinal ganglion cells: the cell-cycle exit of retinoblasts and their neuronal maturation.
Development **132**, 1539-1553.

Mochizuki, T., Luo, Y. J., Tsai, H. F., Hagiwara, A. and Masai, I. (2017). Cell
division and cadherin-mediated adhesion regulate lens epithelial cell movement in
zebrafish. *Development* **144**, 708-719.

Mochizuki, T., Suzuki, S. and Masai, I. (2014). Spatial pattern of cell geometry and
cell-division orientation in zebrafish lens epithelium. *Biol Open* **3**, 982-994.

Mumm, J. S., Williams, P. R., Godinho, L., Koerber, A., Pittman, A. J., Roeser, T.,
Chien, C. B., Baier, H. and Wong, R. O. (2006). In vivo imaging reveals dendritic
targeting of laminated afferents by zebrafish retinal ganglion cells. *Neuron* **52**, 609-621.

Neumann, H., Kotter, M. R. and Franklin, R. J. (2009). Debris clearance by
microglia: an essential link between degeneration and regeneration. *Brain* **132**, 288-295.

O'Koren, E. G., Yu, C., Klingeborn, M., Wong, A. Y. W., Prigge, C. L., Mathew,
R., Kalnitsky, J., Msallam, R. A., Silvin, A., Kay, J. N., et al. (2019). Microglial
Function Is Distinct in Different Anatomical Locations during Retinal Homeostasis and
Degeneration. *Immunity* **50**, 723-737 e727.

Paolicelli, R. C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P.,
Giustetto, M., Ferreira, T. A., Guiducci, E., Dumas, L., et al. (2011). Synaptic
pruning by microglia is necessary for normal brain development. *Science* **333**, 1456-
1458.

Poggi, L., Vitorino, M., Masai, I. and Harris, W. A. (2005). Influences on neural
lineage and mode of division in the zebrafish retina in vivo. *J Cell Biol* **171**, 991-999.

Rhodes, J., Hagen, A., Hsu, K., Deng, M., Liu, T. X., Look, A. T. and Kanki, J. P.
(2005). Interplay of pu.1 and gata1 determines myelo-erythroid progenitor cell fate in
zebrafish. *Dev Cell* **8**, 97-108.

Robinson, M. D., McCarthy, D. J. and Smyth, G. K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139-140.

Rymo, S. F., Gerhardt, H., Wolfhagen Sand, F., Lang, R., Uv, A. and Betsholtz, C.
(2011). A two-way communication between microglial cells and angiogenic sprouts
regulates angiogenesis in aortic ring cultures. *PLoS One* **6**, e15846.

Santos, A. M., Calvente, R., Tassi, M., Carrasco, M. C., Martin-Oliva, D., Marin-
Teva, J. L., Navascues, J. and Cuadros, M. A. (2008). Embryonic and postnatal
development of microglial cells in the mouse retina. *J Comp Neurol* **506**, 224-239.

Scheer, N. and Campos-Ortega, J. A. (1999). Use of the Gal4-UAS technique for
targeted gene expression in the zebrafish. *Mech Dev* **80**, 153-158.

Scheer, N., Riedl, I., Warren, J. T., Kuwada, J. Y. and Campos-Ortega, J. A.
(2002). A quantitative analysis of the kinetics of Gal4 activator and effector gene
expression in the zebrafish. *Mech Dev* **112**, 9-14.

Schmieder, R. and Edwards, R. (2011). Quality control and preprocessing of
metagenomic datasets. *Bioinformatics* **27**, 863-864.

Sehnert, A. J., Huq, A., Weinstein, B. M., Walker, C., Fishman, M. and Stainier, D.
Y. R. (2002). Cardiac troponin T is essential in sarcomere assembly and cardiac
contractility. *Nat Genet* **31**, 106-110.

Shiau, C. E., Kaufman, Z., Meireles, A. M. and Talbot, W. S. (2015). Differential
requirement for irf8 in formation of embryonic and adult macrophages in zebrafish.
PLoS One **10**, e0117513.

Sierra, A., Encinas, J. M., Deudero, J. J., Chancey, J. H., Enikolopov, G.,
Overstreet-Wadiche, L. S., Tsirka, S. E. and Maletic-Savatic, M. (2010). Microglia
shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell*
Stem Cell **7**, 483-495.

Stachura, D. L. and Traver, D. (2011). Cellular dissection of zebrafish hematopoiesis.
Methods Cell Biol **101**, 75-110.

Svahn, A. J., Graeber, M. B., Ellett, F., Lieschke, G. J., Rinkwitz, S., Bennett, M. R.
and Becker, T. S. (2013). Development of ramified microglia from early macrophages
in the zebrafish optic tectum. *Dev Neurobiol* **73**, 60-71.

Tremblay, M. E., Lowery, R. L. and Majewska, A. K. (2010). Microglial interactions
with synapses are modulated by visual experience. *PLoS Biol* **8**, e1000527.

Urasaki, A., Morvan, G. and Kawakami, K. (2006). Functional dissection of the Tol2
transposable element identified the minimal cis-sequence and a highly repetitive
sequence in the subterminal region essential for transposition. *Genetics* **174**, 639-649.

Walton, E. M., Cronan, M. R., Beerman, R. W. and Tobin, D. M. (2015). The
Macrophage-Specific Promoter mfap4 Allows Live, Long-Term Analysis of
Macrophage Behavior during Mycobacterial Infection in Zebrafish. *PLoS One* **10**,
e0138949.

Westerfield, M. (1993). *The zebrafish book : a guide for the laboratory use of*
zebrafish (Brachydanio rerio). Eugene, OR: M. Westerfield.

White, D. T., Sengupta, S., Saxena, M. T., Xu, Q., Hanes, J., Ding, D., Ji, H. and
Mumm, J. S. (2017). Immunomodulation-accelerated neuronal regeneration following
selective rod photoreceptor cell ablation in the zebrafish retina. *Proc Natl Acad Sci U S*
A **114**, E3719-E3728.

Wu, S., Xue, R., Hassan, S., Nguyen, T. M. L., Wang, T., Pan, H., Xu, J., Liu, Q.,
Zhang, W. and Wen, Z. (2018). Il34-Csf1r Pathway Regulates the Migration and
Colonization of Microglial Precursors. *Dev Cell* **46**, 552-563.e554.

Xu, J., Du, L. and Wen, Z. (2012). Myelopoiesis during zebrafish early development.
J Genet Genomics **39**, 435-442.

Xu, J., Wang, T., Wu, Y., Jin, W. and Wen, Z. (2016). Microglia Colonization of
Developing Zebrafish Midbrain Is Promoted by Apoptotic Neuron and
Lysophosphatidylcholine. *Dev Cell* **38**, 214-222.

Xu, J., Zhu, L., He, S., Wu, Y., Jin, W., Yu, T., Qu, J. Y. and Wen, Z. (2015).
Temporal-Spatial Resolution Fate Mapping Reveals Distinct Origins for Embryonic and
Adult Microglia in Zebrafish. *Dev Cell* **34**, 632-641.

Yamaguchi, M., Imai, F., Tonou-Fujimori, N. and Masai, I. (2010). Mutations in N-
cadherin and a Stardust homolog, Nagie oko, affect cell-cycle exit in zebrafish retina.
Mech Dev **127**, 247-264.

Yamaguchi, M., Tonou-Fujimori, N., Komori, A., Maeda, R., Nojima, Y., Li, H.,
Okamoto, H. and Masai, I. (2005). Histone deacetylase 1 regulates retinal
neurogenesis in zebrafish by suppressing Wnt and Notch signaling pathways.
Development **132**, 3027-3043.

Yashiro, K., Shiratori, H. and Hamada, H. (2007). Haemodynamics determined by a
genetic programme govern asymmetric development of the aortic arch. *Nature* **450**,
285-288.

Figure Legends

Figure 1. Microglial precursors progressively colonize developing zebrafish retinas

A) Lateral view of zebrafish eyes used for confocal scanning shown in panel (B).

Anterior is left and dorsal is up. The choroid fissure (cf, arrows) is formed at the

ventral retina. At 32 hpf, the interface space between the neural retina (nr) and lens

appears, in which ocular blood vessels are formed after 36 hpf. At 48 hpf, RGCL

and INL are distinct. At 54 hpf, the ONL becomes evident.

B) Three-dimensional confocal images of mpeg1.1:EGFP-positive microglial

precursors (green) in the retina from 32 to 54 hpf. Dotted circles indicate the outline

of the optic cup. The first microglial precursors appear in the choroid fissure and

near the lens at 32 hpf. Microglial precursors in the optic cup progressively increase

in number. At 42 hpf, they start to enter retinal tissue and spread into the entire

neural retina by 54 hpf.

C) Histogram of the number of intraocular microglial precursors from 32 to 54 hpf.

Horizonal and vertical bars indicate means ± SD.

D) Three-dimensional confocal images of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]*

retinas from 32 to 54 hpf. *Tg[EF1α:mCherry-zGem]* (magenta) indicates cells

undergoing S and G2 phases. mpeg1.1:EGFP-positive microglial precursors (green)

are mostly negative for mCherry-zGem, suggesting that most ocular microglial

precursors are in G1 phase.

E) Histogram of numbers of intraocular microglial precursors expressing only

mpeg1.1:EGFP , and microglial precursors expressing both mCherry-zGem and

mpeg1.1:EGFP in retinas from 32 to 54 hpf. Double-positive microglial precursors

represent proliferating microglial precursors undergoing S/G2 phase. Single

mpeg1.1:EGFP-positive microglial precursors represent microglial precursors in G1

phase. Bars and lines indicate means ± SD.

F) Sections of *Tg[mpeg1.1:EGFP]* transgenic retinas with BrdU incorporated and

labeled with anti-BrdU (magenta) and anti-EGFP (green) antibody. Nuclei were

counterstained by TOPRO3 (blue). The arrowhead indicates BrdU- and

mpeg1.1:EGFP double-positive cells. Most mpeg1.1:EGFP$^+$ cells are BrdU-negative

at 54 hpf, suggesting that they are not proliferative.

G) Histogram of numbers of mpeg1.1:EGFP$^+$ cells and BrdU-positive mpeg1.1:EGFP$^+$

cells per retinal section. Bars and lines indicate means ± SD. **p<0.01.

H) Fraction of BrdU-positive proliferative mpeg1.1:EGFP$^+$ cells in total

mpeg1.1:EGFP$^+$ cells. The average is less than 20%, indicating that more than 80%

of ocular microglial precursors are in the G1 phase. Bars and lines indicate means ±

SD.

Scale bars: 30 μm (B, C). 20 μm (F).

Source data 1. Data for Figure 1C.

Source data 2. Data for Figure 1E.

Source data 3. Data for Figure 1GH.

Source data 4. Data for Figure 1-figure supplement 1B.

Source data 5. Data for Figure 1-figure supplement 2.

Source data 6. Data for Figure 1-figure supplement 4D.

Figure supplement 1. The number of L-plastin-positive cells colonizing developing

retinas.

Figure supplement 2. Spatio-temporal profile of microglial precursor colonization of

developing retinas

Figure supplement 3. Ocular microglial precursors are associated with newly

differentiating amacrine cells

Figure supplement 4. Peripheral microphages of *Tg[EF1α:mCherry-zGem;*

mpeg1.1:EGFP] transgenic embryos

Figure 2. Microglial precursors migrate into the retina along blood vessels

A) Live confocal images of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]* retinas at 30 hpf.

Microglial precursors and blood vessels are visualized using fluorescence of

mfap4tdTomato-CAAX (magenta) and kdrl:EGFP (green), respectively. Higher

magnification image of a dotted square in the left panel is shown in the right panel.

The first microglial precursor (arrow) approaches along developing hyaloid blood

vessels near the lens through the choroid fissure. Arrowheads indicate peripheral

macrophages outside the optic cup. Scale bar: 30 μm.

B) Time-lapse 3D snapshots of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]* eyes for
around 3.5 hr after 32 hpf. Ocular microglial precursors and peripheral
macrophages outside the optic cup are indicated as yellow- and magenta-colored,
surface-rendered objects, respectively, which were prepared from the original
scanning image (Fig.2-figure supplement 1). Ocular blood vessels are visualized in
green. Microglia associated with hyaloid blood vessels around the lens (white
arrows) gradually increase and infiltrate neurogenic retinal tissue (Video 3). Scale
bar: 30 μm.
C) Live 3D images of eyes of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]* embryos
injected with standard MO and tnnt2a MO. kdrl:EGFP-positive blood vessels (green)
are thinner in *tnnt2a* morphants. Scale bar: 50 μm.
D) Histogram of the number of intraocular microglial precursors in embryos injected
with standard MO and tnnt2a MO. Bars and lines indicate means ± SD. ***p<0.001.
1020
Source data 1. Data for Figure 2D.
Source data 2. Data for Figure 2-figure supplement 2B.
Figure supplement 1. Time-lapse snapshots of *Tg[kdrl:EGFP; mfap4:tdTomato-*
CAAX] heads for around 3.5 hr after 32 hpf
Figure supplement 2. Microglial precursor colonization of the optic tectum does not
depend on blood vessel formation
Figure supplement 3. Elimination of microglial precursors does not affect ocular blood
vessel formation
1029
Figure 3. Microglial precursors infiltrate the retina through the neurogenic area
A) Schematic drawing of confocal scanning planes (superficial, middle, and deep
layers) in the optic cup shown in (B) and (D).
B) Live images of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* retinas at 42 hpf (upper
panels) and 48 hpf (lower panels). Two levels of confocal scanning planes are
indicated as superficial (a', a'') and deep positions (c', c''). mpeg1.1:EGFP positive
microglial precursors avoid mCherry-zGem positive proliferating retinal cell area.
Scale bar: 50 μm.

C) Histogram of the fraction of microglial precursors associated with the mCherry-zGem-positive area (black) and the mCherry-zGem-negative area (grey). The fraction of microglial precursors associated with the mCherry-zGem-positive area is only 7.37% at 42 hpf and 6.13% at 48 hpf. Thus, more than 90% of microglial precursors are located in the mCherry-zGem-negative retinal area.

D) Live images of *Tg[ath5:EGFP; mfap4:tdTomato-CAAX]* retinas at 36 (upper panels), 42 (middle panels) and 48 hpf (bottom panels). Three confocal scanning plane levels are indicated as superficial (a'-a'''), middle (b'-b'''), and deep (c-c'''). Dotted circles indicate the outline of the optic cup. The right-most column images indicate higher magnification images shown in the square of left panels. mfap4-positive microglia (magenta, arrows) are closely associated with ath5-positive neurogenic cells (green). Scale bar: 50 µm, except the right-most column images (Scale bar: 15 µm).

Source data 1. Data for Figure 3C.

Source data 2. Data for Figure 3-figure supplement 2B.

Source data 3. Data for Figure 3-figure supplement 3BD.

Figure supplement 1. Microglial precursors infiltrate the neural retina through the neurogenic region

Figure supplement 2. p53 MO effectively inhibits retinal apoptosis in zebrafish

Figure supplement 3. Microglial precursor colonization depends on apoptosis in the optic tectum, but not in the retina

Figure 4. Microglial precursor infiltration into the retina depends on retinal neurogenesis

A) Live 3D images of wild-type and *slbp1* mutant retinas with *Tg[mfap4:tdTomato-CAAX; ath5:EGFP]* at 49 hpf. Only mfap4:tdTomato-CAAX-positive ocular microglial precursors and peripheral macrophages are shown as surface-rendered objects. Original images are shown in Figure 4-figure supplement 2A. Scale bar: 30 µm.

B) Histogram of numbers of ocular microglial precursors in *slbp1* mutants and wild-type siblings. mfap4-positive microglial precursors are significantly fewer in *slbp1* mutants. Bars and lines indicate means ± SD. ***p<0.001.

C) Live 3D images of wild-type and *slbp1* mutant trunk with *Tg[mfap4:tdTomato-CAAX; ath5:EGFP]* at 49 hpf. Scale bar: 70 μm.

D) Histogram of numbers of trunk macrophages in *slbp1* mutants and wild-type siblings. There is no significant difference in mfap4-positive macrophage number in trunks of *slbp1* mutants. Bars and lines indicate means ± SD.

E) Live 3D images of retinas of *Tg[rx1:gal4-VP16; mfap4:tdTomato-CAAX]* embryos injected with one DNA construct encoding *UAS:EGFP* (left) or two DNA constructs encoding *UAS:EGFP; UAS:myc-tagged NICD* (right) at 44 hpf. Only mfap4:tdTomato-CAAX-positive ocular microglial precursors and peripheral macrophages are shown as surface-rendered objects. Original images are shown in Figure 4-figure supplement 2B. Scale bar: 30 μm

F) Histogram of numbers of ocular microglial precursors in *rx1:gal4-VP16; UAS:EGFP* expressed and *rx1:gal4-VP16; UAS:EGFP; UAS:myc-NICD* expressed wild-type retinas. mfap4-positive microglia are significantly decreased in *myc-NICD* expressed retinas, compared with non-injection control and *EGFP* expressed control retinas. Bars and lines indicate means ± SD. *p<0.05, ***p<0.001.

G) Live 3D images of standard MO- and ath5 MO-injected retinas of *Tg[mfap4:tdTomato-CAAX; ath5:EGFP]* embryos at 49 hpf. Only mfap4:tdTomato-CAAX-positive ocular microglial precursors and peripheral macrophages are shown as surface-rendered objects. Original images are shown in Figure 4-figure supplement 2C. Scale bar: 30 μm

H) Histogram of numbers of ocular microglial precursors in standard MO and ath5 MO-injected wild-type retinas. mfap4-positive microglial precursors are significantly less numerous in *ath5* morphant retinas. Bars and lines indicate means ± SD. *p<0.05.

Source data 1. Data for Figure 4B.

Source data 2. Data for Figure 4D.

Source data 3. Data for Figure 4F.

Source data 4. Data for Figure 4H.

Source data 5. Data for Figure 4-figure supplement 4B.

Source data 6. Data for Figure 4-figure supplement 5D.

Source data 7. Data for Figure 4-figure supplement 7C

Figure supplement 1. Retinal neurogenesis and cell differentiation are markedly
delayed in zebrafish *slbp1* mutants.

Figure supplement 2. Original scanning images of Figure 4A, E, G.

Figure supplement 3. Extraction process of ocular microglial precursors from original
3D scanning images.

Figure supplement 4. Retinal apoptosis is increased in zebrafish *slbp1* mutants.

Figure supplement 5. Microglial precursor colonization of the retina is independent of
the Cxcl12a signaling pathway.

Figure supplement 6. Elimination of microglial precursors does not affect retinal cell
differentiation.

Figure supplement 7. Overexpression of NICD suppresses retinal neurogenesis in
zebrafish.

Figure supplement 8. Microglial precursor colonization of the retina is affected in *ath5*
morphants.

1115

Figure 5. Microglial precursors are preferentially associated with neurogenic
retinal columns

A) Schematic drawing of cell transplantation experiments. Wild-type donor embryos
are labeled with Alexa-448-dextran and transplanted into *slbp1* mutant recipient
embryos at blastula stage. In *slbp1* mutant recipient embryos, transplanted wild-type
donor cells form retinal cell columns. The host *slbp1* mutant line is combined with
Tg[mfap4:tdTomato-CAAX], to investigate whether mfap4-positive microglial
precursors (magenta) infiltrate the neural retina preferentially through Alexa-448-
dextran-labeled, wild-type donor columns (green) in *slbp1* mutant recipient embryos.

B) Live images of *slbp1* mutant retinas with transplanted wild-type donor retinal cell
columns at 48 hpf. Donor wild-type retinal cell columns are labeled with Alexa-488
dextran (green). Host microglial precursors are visualized with the transgene
Tg[mfap4:tdTomato-CAAX] (magenta). Dotted circles indicate the outline of the
optic cup. Many microglial precursors are associated with wild-type donor retinal

columns in *slbp1* mutant host retinas (right panel), compared with wild-type sibling

host retinas (left panel). Scale bar: 30 μm

C) The fraction of mfap4-positive microglial precursors associated with donor

transplanted retinal cell columns versus the total number of microglial precursors in

the optic cup. The average fraction of mfap4-positive cells associated with donor

retinal cell columns is significantly higher in *slbp1* mutant host retinas than in wild-

type host retinas. Bars and lines indicate means ± SD. *$p < 0.05$.

D) The trapping efficiency of mfap4-positive microglial precursors per donor column.

The average trapping efficiency is significantly higher in *slbp1* mutant host retinas

than in wild-type host retinas, suggesting higher affinity of microglial precursors for

neurogenic retinal cells. Bars and lines indicate means ± SD. **$p < 0.01$.

E) Schematic drawing of mosaic expression of NICD in retinas. A mixture of

UAS:EGFP and UAS-myc-NICD plasmids was injected into fertilized eggs of the

Tg[hsp:gal4; mfap4:tdTomato] transgenic line, which were treated by heat shock at

18 and 30 hpf. At 48 hpf, embryos were fixed to prepare serial retinal sections for

imaging analysis.

F) Confocal scanning of retinal sections of *Tg[hsp:gal4; mfap4:tdTomato]* transgenic

embryos injected with plasmids encoding UAS:EGFP or UAS:EGFP+UAS:myc-

NICD. Scale bar: 30 μm.

G) The fraction of mfap4-positive microglial precursors associated with EGFP-

expressing retinal cell columns versus the total number of microglial precursors in

the optic cup. The average fraction of mfap4-positive cells associated with EGFP-

positive retinal columns is significantly lower in retinas injected with

UAS:EGFP+UAS:myc-NICD than with only UAS:EGFP control. Bars and lines

indicate means ± SD. ***$p < 0.005$.

H) The trapping efficiency of mfap4-positive microglial precursors per EGFP-

expressing retinal cell columns. The average trapping efficiency is significantly

lower in retinas injected with UAS:EGFP+UAS:myc-NICD than with only

UAS:EGFP control, suggesting less affinity of microglial precursors for

proliferative NICD-expressing retinal cells. Bars and lines indicate means ± SD.

*$p < 0.05$.

Source data 1. Data for Figure 5CDGH.

Figure supplement 1. Calculation of trapping efficiency of microglial precursors by

retinal columns.

1164

Figure 6. IL34 is required for colonization of the optic cup by microglial

precursors.

A) Confocal 3D scanning of 48 hpf wild-type, *il34* heterozygous and homozygous

mutant retinas carrying the *Tg[mfap4:tdTomato-CAAX]* transgene. At 48 hpf,

iridophores start to differentiate around the optic cup, which causes a noise signal

(magenta) in confocal scanning. Using the surface-rendering tool of Imaris software

(Bitplane), we eliminated iridophore-derived noise and extracted mfap4:tdTomato-

CAAX signals from ocular microglial precursors (green) (See the legend of Figure

4-figure supplement 3). Scale bar: 50 μm.

B) Histogram of numbers of ocular microglial precursors in wild-type, *il34*

heterozygous and homozygous mutant retinas at 48 hpf. The number of ocular

microglial precursors is almost zero, and very few, if any (one or two), in *il34*

homozygous mutants, indicating that ocular microglial precursors are significantly

reduced in *il34* homozygous mutants. The number of ocular microglial precursors is

mildly reduced in *il34* heterozygous mutants, but does not differ significantly from

that of wild-type siblings. Bars and lines indicate means ± SD. ***$p<0.005$.

C) A possible model of the guidance mechanism of microglial precursor into zebrafish

retina. IL34 is involved in movement of microglial precursors toward the brain.

Microglial precursors continue into the optic cup along blood vessels, and

subsequently infiltrate the neural retina through the neurogenic area.

Source data 1. Data for Figure 6B.

Source data 2. Data for Figure 6-figure supplement 2B

Source data 3. Data for Figure 6-figure supplement 3

Figure supplement 1. Retinal cell differentiation normally proceeds in *il34* mutants

Figure supplement 2. Microglial precursor colonization of the optic cup in *il34*

mutants

Figure supplement 3. *il34* mRNA expression is comparable between wild-type sibling

and *slbp1* mutant heads.

1193

 Supplementary Figure Legends

 Figure 1–figure supplement 1. The number of L-plastin-positive cells colonizing

 developing retinas.

 (A) Lateral view of the zebrafish optic cup labeled with *L-plastin* mRNA probe at 32, 36,

    42, 48, 54 hpf. *L-plastin* mRNA is expressed in microglial precursors and

    neutrophils in the developing zebrafish brain, including the retinas. Since

    neutrophils do not invade the brain area during embryonic development, *L-plastin*-

    positive ocular cells are thought to be microglial precursors. *L-plastin*-positive

    ocular cells increase progressively in number during development. Scale: 50 μm.

 (B) Histogram of the number of *L-plastin*-positive ocular cells. Horizontal and vertical

    bars indicate means ± SD.

 Figure 1–figure supplement 2. Spatio-temporal profile of microglial precursor

 colonization of developing retinas.

 Left and middle columns indicate *Tg[mfap4:TdTomato-CAAX]* transgenic zebrafish

 retinas labeled with Bodipy ceremide at 32, 36, 42, 48 and 54 hpf. The right column

 shows a histogram of numbers of mfap4$^+$ cells in the vitreous space between the retina

 and the lens and in the neural retina (means ± SD). At 32 and 36 hfp, most mfap4$^+$

 microglial precursors are associated with hyaloid blood vessels that form in the vitreous

 space between the retina and the lens. After 42 hpf, mfap4$^+$ cells start to infiltrate the

 neural retina and are often associated with emerging IPL. The number of mfap4$^+$ cells in

 the neural retina is lower than in the vitreous space at 48 hpf, but becomes higher at 54

 hpf. Scale bars: 30 μm.

 Figure 1–figure supplement 3. **Ocular microglial precursors are associated with**

 newly differentiating amacrine cells.

 Confocal image of a 44-hpf *Tg[mfap4:tdTomato-CAAX]* transgenic retina injected with

 a DNA construct encoding Ptf1a:EGFP. The lower panel indicates a higher

 magnification image of the dotted square shown in the upper panel. The dotted circle

 and dotted line in the lower panel indicate the outline of the lens and retinal pigmented

epithelium (pe), respectively. Microglial precursors (magenta) are transiently located in the vitreous space between the lens and the retina (arrowheads), and start to infiltrate the neural retina after 42 hpf. Most infiltrated microglial precursors (arrows) are associated with newly differentiating amacrine cells, which express Ptf1a:EGFP (green). Magenta signals associated with pigmentated epithelium are noise and derived from autofluorescence of iridophores. Scale bars: 30 μm.

Figure 1–figure supplement 4. Peripheral microphages of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* transgenic embryos.

(A) Schematic drawing of 36 hpf zebrafish embryos indicating the scanning area in yolk and tail, which are shown in panels (B) and (C), respectively. RBI, rostral blood island; VDA, ventral wall of dorsal aorta.

(B) Confocal scanning of peripheral macrophages in yolk of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* transgenic embryos at 36 hpf. Yellow arrows indicate mCherry-zGem and mpeg1.1:EGFP double-positive cells, whereas white arrows indicate only mpeg1.1:EGFP-positive cells. The bottom four panels are higher magnifications of the white square in the top-left panel. Scale bars: 40 μm.

(C) Confocal scanning of peripheral macrophages in tails of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* transgenic embryos at 36 hpf. Yellow arrows indicate mCherry-zGem and mpeg1.1:EGFP double-positive cells, whereas white arrows indicate only mpeg1.1:EGFP-positive cells. The bottom four panels are higher magnifications of the white square in the top-left panel. Scale bars: 20 μm.

(D) The fraction of mCherry-zGem and mpeg1.1:EGFP double-positive cells among all mpeg1.1:EGFP-positive cells. We detected a total of 7 mpeg1.1:EGFP$^+$ cells in the yolk area and 39 cells in the tail area of confocal scanning images obtained from different *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* transgenic embryos, and then calculated the fraction of mCherry-zGem and mpeg1.1:EGFP double-positive cells among all mpeg1.1:EGFP-positive cells. The average fraction of double-positive cells was 71.4% for yolk and 61.5% for tail. Thus, more than 60% of peripheral macrophages undergo S phase in zebrafish embryos, suggesting that the *Tg[EF1α:mCherry-zGem]* transgenic system works as an indicator of cell-cycle

1256 phases in peripheral macrophages.

1257

Figure 2–figure supplement 1. Time-lapse snapshots of *Tg[kdrl:EGFP;*

***mfap4:tdTomato-CAAX]* heads for around 3.5 hr after 32 hpf.**

Original confocal 3D scanning images of Figure 2B. Microglial precursors and

peripheral macrophages (magenta) and blood vessels (green) are visualized. Arrows

indicate microglial precursors associated with hyaloid blood vessels around the lens,

which are indicated as surface-rendered objects in yellow in Figure 2B. Scale bar: 30

μm.

1265

Figure 2–figure supplement 2. Microglial precursor colonization of the optic

tectum does not depend on blood vessel formation.

(A) Live images of the optic tectum of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]*

embryos injected with standard MO and tnnt2a MO at 48 and 72 hpf. The optic

tectum area is indicated by a dotted line. Blood vessels (green) are visualized with

kdrl:EGFP and were thinner in *tnnt2a* morphants. In the standard MO-injected

control embryos, the number of mfap4:tdTomato-CAAX-positive microglial

precursors was 5.2±1.3 at 48 hpf and increased to 33.5±9.1 at 72 hpf, indicating

progressive colonization of the optic tectum during development. On the other hand,

the number of microglial precursors in the optic tectum was 17.8±3.7 at 48 hpf and

higher in *tnnt2a* morphants than in control embryos. Furthermore, their shape was

round rather than ramified, suggesting activation of microglial precursors. This

observation implies that microglial colonization of the optic tectum is enhanced

when blood vessel function is compromised. However, the number of microglial

precursors was 23.3±5.6 in *tnnt2a* morphant optic tectum at 72 hpf, and microglial

precursors still showed a round shape. Since microglial precursors show a

phagocytic morphology in *tnnt2a* morphants and apoptosis normally promotes

microglial colonization of the optic tectum in zebrafish, increased apoptosis may

enhance microglial colonization of the optic tectum in *tnnt2a* morphants at 48 hpf.

Scale bar: 50 μm.

(B) Histogram of the number of optic tectum-resident microglial precursors in embryos

injected with standard MO and tnnt2a MO. The number was significantly higher in

tnnt2a morphants than in standard MO-injected embryos at 48 hpf; however, as

reported in previous studies by other groups, there was no statistical difference

between standard MO-injected embryos and *tnnt2a* morphants at 72 hpf. Bars and

lines indicate means ± SD. ***$p<0.001$.

(C) Embryonic morphology of embryos injected with standard MO and *tnnt2a*

morphants at 48 and 72 hpf.

1294

Figure 2–figure supplement 3. Elimination of microglial precursors does not affect

ocular blood vessel formation.

(A, B) Confocal 3D scanning images of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]*

transgenic retinas injected with standard control MO (A) and pu.1 MO (B) at 48 hpf.

mfap4:tdTomato-CAAX signals indicate ocular microglial precursors and peripheral

microphages outside the optic cup; however, scanning with mfap4:tdTomato-CAAX

after 48 hpf also detects noise signal caused by reflection of iridophore pigment cells,

which are positioned surrounding the optic cup.

(A', B') 3D images of *kdrl:EGFP*-positive ocular blood vessels and *mfap4:tdTomato-*

CAAX surface-rendered objects indicating ocular microglial precursors in retinas

injected with standard control MO (A) and pu.1 MO (B). To prepare surface-rendered

objects indicating ocular microglial precursors, we digitally removed noise signals

caused by reflection of iridophore pigment cells as well as peripheral macrophages

outside the optic cup, using the surface-rendering tool of Imaris software (Bitplane) (see

the legend of Figure 4-figure supplement 3).

(A'', B'') Only *mfap4:tdTomato-CAAX* surface-rendered objects indicating ocular

microglial precursors in (A', B') are shown.

(A''', B''') Only *kdrl:EGFP* positive ocular blood vessels in (A', B') are shown.

Ocular blood vessels are normally formed in *pu.1* morphants, suggesting that ocular

blood vessel formation is independent of microglial precursor differentiation at least by

48 hpf.

(C, D) Confocal 3D scanning images of *Tg[kdrl:EGFP]* transgenic retinas combined

with *irf8* mutants (D) and wild-type siblings (C) at 48 hpf. Ocular blood vessels

visualized with *Tg[kdrl:EGFP]* are normally formed in *irf8* mutants, suggesting that

ocular blood vessel formation is independent of microglial precursor differentiation.

Scale bars: 30 μm.

1321

Figure 3–figure supplement 1. Microglial precursors infiltrate the neural retina

through the neurogenic region.

(A) Confocal scanning of the neural retina along the Z-axis.

(B) Scanning images of *Tg[EF1α:mCherry-Gem; mpeg1.1:EGFP]* transgenic retinas.

Panels 1 to 8 indicate the Z axis-position of the confocal scanning plane shown in

(A). Yellow arrowheads indicate the choroid fissure. Microglial precursors (green,

arrows) infiltrate the neural retina through the mCherry-Gem (magenta)-negative

region, suggesting that microglial precursors infiltrate the neural retina through the

neurogenic region. Scale bar: 70 μm.

1331

Figure 3–figure supplement 2. p53 MO effectively inhibits retinal apoptosis in

zebrafish.

(A) TUNEL (red) of retinas injected with standard MO and p53 MO at 24, 36 and 48

hpf. Scale bars: 30 μm.

(B) The number of TUNEL-positive cells per a retinal section of embryos injected with

standard MO and p53 MO at 24, 36 and 48 hpf. The number of TUNEL-positive

cells was significantly lower in p53 morphant retinas than in standard MO injected

retinas at 24 and 36 hpf. However, the number of TUNEL-positive cells was low in

both retinas injected with standard MO and p53 MO at 48 hpf, indicating no

statistical difference between them. Bars and lines indicate means ± SD. *$p<0.05$.

1342

Figure 3–figure supplement 3. Microglial precursor colonization depends on

apoptosis in the optic tectum, but not in the retina.

(A) Confocal scanning images of *Tg[mfap4:tdTomato-CAAX]* transgenic retinas

injected with standard MO and p53 MO at 48 hpf. Scale bars: 30 μm.

(B) Histogram of the number of ocular mfap4:tdTomato-CAAX$^+$ cells at 48 hpf. There

is no significant difference between standard MO-injected retinas and *p53*

morphant retinas, suggesting that apoptosis is not a major attractant for microglial

precursors migrating to the retina. Bars and lines indicate means ± SD.

(C) Confocal scanning images of a *Tg[mfap4:tdTomato-CAAX]* transgenic optic tectum

injected with standard MO and p53 MO at 96 hpf. The optic tectum area is outlined.

Scale bars: 50 μm.

(D) Histogram of the number of optic tectum-resident mfap4:tdTomato-CAAX$^+$ cells at

96 hpf. The number of optic tectum-resident microglial precursors is significantly

lower in *p53* morphants, suggesting that microglial precursor colonization of the

optic tectum depends on apoptosis. Bars and lines indicate means ± SD. *p<0.05.

1358

Figure 4–figure supplement 1. Retinal neurogenesis and cell differentiation are

markedly delayed in zebrafish *slbp1* mutants.

(A) Plastic sections of wild-type sibling and *slbp1* mutant retinas at 72 hpf. In wild-type

retinas, retinal lamination is formed, whereas only IPL is emerging in *slbp1* mutant

retinas, which are equivalent to wild-type retinas at 48 hpf.

(B) Volcano plotting of up- (red) and down-regulated (blue) genes in *slbp1* mutants, as

determined by RNAseq analysis. Only genes of FDR<0.01 and logFC>|2| are shown.

Genes that promote neurogenesis and photoreceptor cell differentiation are

downregulated.

(C) Heat map of expression of four neurogenic genes; neurod1, neurod2, neurod6a, and

neurod6b in wild-type siblings and *slbp1* mutants. All four genes are decreased in

slbp1 mutants.

1371

Figure 4–figure supplement 2. Original scanning images of Figure 4A, E, G.

(A, B, C) Original confocal 3D scanning images of Figures 4A, 4E, and 4G,

respectively. Figures 4A, E, and G show surface-rendered objects of mfap4:tdTomato-

CAAX signals, which correspond to ocular microglial precursors and peripheral

macrophages surrounding the optic cup. Arrows indicate noise signals derived from

iridophore-mediated reflection, which are removed by the surface-rendering tool of

Imaris in Figures 4A, 4E, and 4G (See the legend of Figure 4-figure supplement 3). The

number of iridophore-derived noise signals was generally lower in eyes of *slbp1*

mutants, myc-NICD overexpressing embryos and *ath5* morphants than in their controls.

It is likely that the timing of iridophore differentiation correlates with retinal
development, so a delay of retinal neurogenesis may delay the timing of iridophore
appearance. Scale bar: 30 μm for (A, B, C).

1384

Figure 4–figure supplement 3. Extraction process of ocular microglial precursors
from original 3D scanning images.
Confocal scanning of retinas with mfap4:tdTomato-CAAX after 48 hpf detects noise
signals caused by reflection of iridophore pigment cells. In this case, we digitally
removed iridophore pigment-derived noise from 3D scanning images. Using the Imaris
surface-rendering tool (Bitplane), mfap4:tdTomato-CAAX signals are converted into
surface-rendered objects. Surface-rendered objects corresponding to iridophore-derived
signals are selected with the editing tool and are eliminated manually. Only
mfap4:tdTomato-CAAX signals derived from ocular microglial precursors and
peripheral macrophages surrounding the optic cup are retained, and merged to
reconstruct 3D images.

1396

Figure 4–figure supplement 4. Retinal apoptosis is increased in zebrafish *slbp1*
mutants.
(A) TUNEL of wild-type and *slbp1* mutant retinas at 48 hpf. The number of TUNEL-
positive signals (magenta) was high (n=14.1±7.4) in *slbp1* mutant retinas, whereas
it was normally low (n=1.8±1.3) in wild-type sibling retinas, suggesting that
apoptosis is enhanced in *slbp1* mutant retinas. Nuclei were counterstained with
TOPRO3 (green). Scale bars: 30 μm.
(B) The number of TUNEL-positive cells per retinal section. The number is
significantly higher in *slbp1* mutant retinas than in wild-type sibling retinas. Bars
and lines indicate means ± SD. *p<0.05.

1407

1408 **Figure 4–figure supplement 5. Microglial precursor colonization of the retina is**
1409 **independent of the Cxcl12a signaling pathway.**

(A) Anti-acetylated tubulin antibody labeling of 72 hpf retinas injected with standard MO and cxcl12a MO. Arrowheads indicate RGC axons. In *cxcl12a* morphants, an abnormal trajectory of RGC axons inside the retina. Scale bars: 30 μm.

(B) ath5:EGFP expression in frontally viewed heads of embryos injected with standard MO and cxcl12a MO at 72 hpf. ath5:EGFP labels RGCs and their axons, which normally exit from the optic cup and create a chiasm at the midline. *cxcl12a* morphants show an abnormal trajectory of RGC axons inside the retina (arrowheads). Scale bars: 30 μm.

(C) Confocal scanning images of *Tg[mpeg1.1:EGFP]* transgenic retinas injected with standard control MO and cxcl12a MO at 48 hpf. Scale bars: 30 μm.

(D) Histogram of the number of ocular mpeg1.1:EGFP$^+$ cells. There is no significant difference between standard MO-injected retinas and *cxcl12*a morphant retinas, suggesting that Cxcl12a signaling is not involved in microglial precursor colonization of the retina. Bars and lines indicate means ± SD.

Figure 4–figure supplement 6. Elimination of microglial precursors does not affect retinal cell differentiation.

Labeling of *pu.1* morphant retinas with zpr1 antibody, ath5:EGFP and mfap4:tdTomato-CAAX. mapf4:tdTomato-CAAX$^+$ microglia are absent in *pu.1* morphants. However, ath5:EGFP and zrp1 expression are normally observed in *pu.1* morphants, suggesting that elimination of microglia does not affect retinal cell differentiation. Scale bar: 30 μm.

Figure 4-figure supplement 7. Overexpression of NICD suppresses retinal neurogenesis in zebrafish.

(A) Experimental design of overexpression of NICD in zebrafish retinas. A DNA construct encoding UAS:myc-NICD was injected into fertilized eggs produced by pairwise crosses of *Tg[ath5:EGFP]* and *Tg[hsp:gal4]* transgenic fish. Injected embryos were maintained with PTU to prevent melanophore pigmentation and treated with heat shock at 18, 24 and 30 hpf, and fixed with 4% PFA at 36 hpf. Whole-mount embryos were labeled with anti-myc-tag antibody and anti-EGFP antibody to investigate whether myc-NICD-expressing retinal cells are ath5:EGFP-

negative. On average, 50% of embryos did not have the *Tg[hsp:gal4]* transgene and were used for negative controls.

(B) Labeling of *Tg[ath5:EGFP; hsp:gal4]* transgenic retinas injected with the UAS:myc-NICD or UAS:mCherry expression construct. We observed that retinal cell columns expressing myc-NICD (magenta) do not express ath5:EGFP (green) (white arrows), whereas retinal cells columns expressing mCherry normally express ath5:EGFP in the neurogenic area. ath5:EGFP normally spread into the whole retina of non-*Tg[hsp:gal4]* transgenic embryos, which did not express myc-NICD. Scale bars: 40 μm.

(C) Percentages of ath5:EGFP-negative and -positive retinal columns expressing myc-NICD (left panel) and mCherry (right panel). Among the 85 retinal columns expressing myc-NICD from 5 independent embryos, 97.65% were ath5:EGFP negative. On the other hand, among the 224 retinal columns expressing mCherry from 5 independent embryos, 53.13% were ath5:EGFP negative. Since ath5:EGFP is the earliest marker of retinal neurogenesis in zebrafish, overexpression of NICD effectively suppresses retinal neurogenesis. The statistical difference was significant ($p < 0.0001$, chi square).

Figure 4–figure supplement 8. Microglial precursor colonization of the retina is affected in *ath5* morphants.

zn5 antibody labeling of 49 hpf *Tg[mfap4:TdTomato-CAAX]* transgenic retinas injected with standard MO and ath5 MO. Panels (B) and (C) display only green and magenta channels, respectively. RGCs are visualized with zn5 antibody (green) and their differentiation is inhibited in *ath5* morphants. The number of ocular microglial precursors (magenta) is drastically reduced in *ath5* morphants. Nuclei were counterstained with TOPRO3 (white). Scale bar: 30 μm.

Figure 5–figure supplement 1. Calculation of trapping efficiency of microglial precursors by retinal columns.

(A) Schematic drawing of five sets of wild-type sibling retinas and three sets of *slbp1* mutant host retinas, into which wild-type donor retinal cells were transplanted, and

data of which are used for Fig. 5A-D. At 48 hpf, *slbp1* mutant host retinas consist of

proliferative retinal cells (yellow circles), whereas wild-type sibling retinas consists

of neurogenic retinal cells (light green circles). Wild-type donor-transplanted

columns contain neurogenic retinal cells (green circles). Microglial precursors

associated with transplanted donor columns are indicated by red squares. Multiple

red squares indicate multiple microglial precursors associated with one transplanted

retinal column.

(B) Table indicating the total number of host ocular microglial precursors, the number

of microglial precursors associated with transplanted donor columns, the percentage

of microglial precursors associated with transplanted donor columns relative to total

numbers of microglia per host retina, total numbers of transplanted donor columns,

and trapping efficiency of microglial precursors per neurogenic donor column. The

trapping efficiency of microglial precursors by each neurogenic column is

calculated by dividing the percentage of microglial precursors associated with donor

columns by the number of donor retinal columns per host retina. These results are

shown in Fig. 5CD.

(C) Table indicating the total number of ocular microglial precursors, the number of

microglial precursors associated with EGFP or EGFP/NICD-expressing columns,

the percentage of microglial precursors associated with EGFP or EGFP/NICD-

expressing columns relative to total numbers of microglial precursors per host retina,

total number of EGFP or EGFP/NICD-expressing columns, and trapping efficiency

of microglial precursors per EGFP or EGFP/NICD-expressing column. These

results are shown in Fig. 5GH.

1495

Figure 6–figure supplement 1. Retinal cell differentiation normally proceeds in *il34*

mutants.

Wild-type sibling and *il34* mutant retinas at 72 hpf. Retinal sections of three

independent *il34* mutants are shown. Although there were 6-10 pyknotic nuclei (arrows)

in RGC and amacrine cell layers in *il34* mutants, retinal lamination is normally formed

at 72 hpf, indicating that neurogenesis and subsequent cell differentiation normally

proceed in *il34* mutants.

1503

Figure 6–figure supplement 2. Microglial precursor colonization of the optic cup in

il34 **mutants.**

(A) Confocal 3D scanning images of 34 hpf wild-type, *il34* heterozygous, and

homozygous mutant retinas carrying the *Tg[mfap4:tdTomato-CAAX]* transgene.

Using the surface-rendering tool of Imaris software (Bitplane), we labeled ocular

microglial precursors (green). Scale bar: 40 μm.

(B) Histogram of numbers of ocular microglial precursors in wild-type, *il34*

heterozygous, and homozygous mutant retinas at 34 hpf. The number of ocular

microglial precursors is zero in *il34* homozygous mutants, and significantly lower

than that of wild-type siblings. Bars and lines indicate means ± SD. ***p<0.005.

1514

Figure 6–figure supplement 3. *il34* **mRNA expression is comparable between wild-**

type sibling and *slbp1* **mutant heads.**

il34 mRNA expression was evaluated in wild-type siblings and *slbp1* mutants at 48 hpf

using semi-quantitative PCR. *il34* mRNA expression level is normalized against *actb2*

mRNA expression. There is no significant difference between wild-type siblings and

slbp1 mutants. Bars and lines indicate means ± SD.

1521

1522 **Legends of Video files**

1523

Video 1: **3D rendering of an eye of *Tg[EF1α: mCherry-zGem;mpeg1.1:EGFP]***
zebrafish embryo at 42 hpf.

mCherry-zGem signals indicate cells in S and G2 phase (magenta), whereas
mpeg1.1:EGFP -positive cells are ocular microglial precursors (green). The fraction of
mCherry-zGem; mpeg1.1:EGFP double-positive cells in mpeg1.1:EGFP-positive cells
is very small, suggesting that almost all microglial precursors are in G1 phase.

1530

Video 2: Live imaging of *Tg[kdrl:EGFP; mfap4-tdTomato-CAAX]* embryo at 30 hpf.
mfap4-tdTomato-CAAX-positive cells indicate microglial precursors (magenta),
whereas kdrl:EGFP-positive cells indicate endothelial cells of blood vessels (green).
Microglial precursors are moving on the surface of a developing superficial ocular
blood vessel, suggesting that blood vessels act as scaffolds for migration of microglial
precursors.

1537

Video 3: Live imaging of *Tg[kdrl:EGFP; mfap4-tdTomato-CAAX]* embryos from 32
to 36 hpf.

mfap4-tdTomato-CAAX-positive cells indicate microglial precursors (magenta),
whereas kdrl:EGFP-positive cells indicate endothelial cells of blood vessels (green).
Microglial precursors use blood vessels as scaffolds to migrate into the ocular space,
and gradually invade the neural retina. Surface rendering indicates amoeboid microglial
precursors, which are attached to the hyaloid loop around the lens and infiltrate the
neural retina.

1546

1547

List of Source Data files

Figure 1-Source data 1. Data for Figure 1C.

Figure 1-Source data 2. Data for Figure 1E.

Figure 1-Source data 3. Data for Figure 1GH.

Figure 1-Source data 4. Data for Figure 1-figure supplement 1B.

Figure 1-Source data 5. Data for Figure 1-figure supplement 2.

Figure 1-Source data 6. Data for Figure 1-figure supplement 4D.

Figure 2-Source data 1. Data for Figure 2D.

Figure 2-Source data 2. Data for Figure 2-figure supplement 2B.

Figure 3-Source data 1. Data for Figure 3C.

Figure 3-Source data 2. Data for Figure 3-figure supplement 2B.

Figure 3-Source data 3. Data for Figure 3-figure supplement 3BD.

Figure 4-Source data 1. Data for Figure 4B.

Figure 4-Source data 2. Data for Figure 4D.

Figure 4-Source data 3. Data for Figure 4F.

Figure 4-Source data 4. Data for Figure 4H.

Figure 4-Source data 5. Data for Figure 4-figure supplement 4B.

Figure 4-Source data 6. Data for Figure 4-figure supplement 5D.

Figure 4-Source data 7. Data for Figure 4-figure supplement 7C

Figure 5-Source data 1. Data for Figure 5CDGH.

Figure 6-Source data 1. Data for Figure 6B.

Figure 6-Source data 2. Data for Figure 6-figure supplement 2B

Figure 6-Source data 3. Data for Figure 6-figure supplement 3

List of Supplementary figure files

Figure 1–figure supplement 1. Number of L-plastin-positive cells colonizing the developing retinas.

Figure 1–figure supplement 2. Spatio-temporal profile of microglial precursor colonization of developing retinas.

Figure 1–figure supplement 3. Ocular microglial precursors are associated with newly differentiating amacrine cells.

Figure 1–figure supplement 4. Peripheral microphages of *Tg[EF1α:mCherry-zGem; mpeg1.1:EGFP]* transgenic embryos

Figure 2–figure supplement 1. Time-lapse snapshots of *Tg[kdrl:EGFP; mfap4:tdTomato-CAAX]* heads for around 3.5 hr after 32 hpf.

Figure 2–figure supplement 2. Microglial precursor colonization of the optic tectum does not depend on blood vessel formation.

Figure 2–figure supplement 3. Elimination of microglial precursors does not affect ocular blood vessel formation.

Figure 3–figure supplement 1. Microglial precursors infiltrate the neural retina through the neurogenic region.

Figure 3–figure supplement 2. p53 MO effectively inhibits retinal apoptosis in zebrafish.

Figure 3–figure supplement 3. Microglial precursor colonization depends on apoptosis in the optic tectum, but not in the retina.

Figure 4–figure supplement 1. Retinal neurogenesis and cell differentiation are markedly delayed in zebrafish *slbp1* mutants.

Figure 4–figure supplement 2. Original scanning images of Figure 4A, E, G.

Figure 4–figure supplement 3. Extraction process of ocular microglial precursors from original 3D scanning images.

Figure 4–figure supplement 4. Retinal apoptosis is increased in zebrafish *slbp1* mutants.

Figure 4–figure supplement 5. Microglial precursor colonization of the retina is independent of the Cxcl12a signaling pathway.

Figure 4–figure supplement 6. Elimination of microglial precursors does not affect retinal cell differentiation.

Figure 4–figure supplement 7. Overexpression of NICD suppresses retinal neurogenesis in zebrafish.

Figure 4–figure supplement 8. Microglial precursor colonization of the retina is affected in *ath5* morphants.

Figure 5–figure supplement 1. Calculation of trapping efficiency of microglial precursors by retinal columns.

Figure 6–figure supplement 1. Retinal cell differentiation normally proceeds in *il34* mutants.

Figure 6–figure supplement 2. Microglial precursor colonization of the optic cup in *il34* mutants.

Figure 6–figure supplement 3. *il34* mRNA expression is comparable between wild-type sibling and *slbp1* mutant heads.

List of Video files

Video 1: 3D rendering of an eye of *Tg[EF1α: mCherry-zGem;mpeg1.1:EGFP]* zebrafish embryo at 42 hpf.

Video 2: Live imaging of *Tg[kdrl:EGFP; mfap4-tdTomato-CAAX]* embryo at 30 hpf.

Video 3: Live imaging of *Tg[kdrl:EGFP; mfap4-tdTomato-CAAX]* embryos from 32 to 36 hpf.

Appendix

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (zebrafish, Danio rerio)	Okinawa wild type	PMID: 28196805		
strain, strain background (zebrafish, Danio rerio)	RIKEN wild type	PMID: 12702661	ZFIN: ZDB-GENO-070802-4	https://shigen.nig.ac.jp/zebra/
genetic reagent (zebrafish, Danio rerio)	*slbp1^{rw440}*	PMID: 25106852	ZFIN: ZDB-ALT-150115-5	
genetic reagent (zebrafish, Danio rerio)	*irf8^{st96}*	PMID: 25615614	ZFIN: ZDB-ALT-150504-8	Dr. William Talbot (Stanford University)
genetic reagent (zebrafish, Danio rerio)	*il34^{hkz11}*	PMID: 30205037	ZFIN: ZDB-ALT-181210-3	Dr. Zilong Wen (The Hong Kong University of Science and Technology)
genetic reagent (zebrafish, Danio rerio)	*Roy orbison*	PMID: 28760346	ZFIN: ZDB-ALT-980203-444	
genetic reagent (zebrafish, Danio rerio)	*Tg[ath5:EGFP]^{rw021}*	PMID: 12702661	ZFIN: ZDB-ALT-050627-2	

genetic reagent (zebrafish, Danio rerio)	Tg[EF1a:mCherry-zGem] ^{oki011}	PMID: 25260917	ZFIN: ZDB-ALT-150128-2	
genetic reagent (zebrafish, Danio rerio)	Tg[mpeg1.1:EGFP] ^{oki053}	This paper		See "Materials and Methods"
genetic reagent (zebrafish, Danio rerio)	Tg[mfap4:tdTomato] ^{oki083}	This paper		See "Materials and Methods"
genetic reagent (zebrafish, Danio rerio)	Tg[kdrl:EGFP] ^{s843Tg}	PMID: 16251212	ZFIN: ZDB-ALT-050916-14	ZIRC
genetic reagent (zebrafish, Danio rerio)	Tg[hsp:gal4] ^{kca4}	PMID: 11850174	ZFIN: ZDB-ALT-020918-6	Reugels/Campos-Ortega lab (Köln University)
genetic reagent (zebrafish, Danio rerio)	Tg[rx1:gal4-VP16] ^{oki065}	This paper		See "Materials and Methods"
antibody	zn5 (mouse monoclonal)	ZIRC	ZFIN: ZDB-ATB-081002-19	IHC (1:100)
antibody	zpr1 (mouse monoclonal)	ZIRC	ZFIN: ZDB-ATB-081002-43	IHC (1:100)
antibody	Anti-GFP (rabbit polyclonal)	Thermo Fisher Scientific	Cat# A11122	IHC (1:200)

antibody	Anti-myc tag (mouse monoclonal)	Invitrogen	Cat# R950-25	IHC (1:250)
antibody	Anti-BrdU (rat monoclonal)	BioRad	Cat# MCA2060	Monoclonal (BU1/75(ICR1)) IHC (1:200)
recombinant DNA reagent	pT2AL200R150G (plasmid)	PMID: 16959904		Dr. Koichi Kawakami (Institute of Genetics)
recombinant DNA reagent	UAS:EGFP (plasmid)	PMID: 11336499		10 ng/μL for injection
recombinant DNA reagent	UAS:mCherry (plasmid)	This paper		10 ng/μL for injection
recombinant DNA reagent	UAS:myc-NICD (plasmid)	PMID: 10072782		Reugels/Campos-Ortega lab (Köln University) 10 ng/μL for injection
recombinant DNA reagent	Ptf1a:EGFP (plasmid)	PMID: 19732413		Dr. Francesco Argenton (University of Padova) 10 ng/μL for injection
sequence-based reagent	tnnt2a MO	PMID: 11967535	Morpholino antisense oligos	5'-CATGTTTGCTCTGATCTGACACGCA-3' Use at 500 μM

sequence-based reagent	p53 MO	PMID: 12477391	Morpholino antisense oligos	5'-GCGCCATTGCTTTGCAAGAATTG-3' Use at 100 μM
sequence-based reagent	cxcl12a MO	PMID: 15716407	Morpholino antisense oligos	5'-ACTTTGAGATCCATGTTTGCAGTG-3' Use at 500 μM
sequence-based reagent	pu.1 MO	PMID: 15621533	Morpholino antisense oligos	5'-GATATACTGATACTCCATTGGTGGT-3' Use at 250 μM
sequence-based reagent	ath5 MO	This paper	Morpholino antisense oligos	5'-TTCATGGCTCTTCAAAAAAGTCTCC-3' Use at 250 μM
sequence-based reagent	standard MO	other	Morpholino antisense oligos	5'-CCTCTTACCTCAGTTACAATTTATA-3' Use at the same concentration for each MO experiments

sequence-based reagent	il34 qPCR primer forward	This paper	PCR primers	5'-tggtccagtccgaatgct-3'
sequence-based reagent	il34 qPCR primer reverse	This paper	PCR primers	5'-gctgcactactgcacactgg -3'
sequence-based reagent	actb2 qPCR primer forward	This paper	PCR primers	5'-tgtcttcccatccatcgtg -3'
sequence-based reagent	actb2 qPCR primer reverse	This paper	PCR primers	5'-tgtcttcccatccatcgtg-3'
sequence-based reagent	il34 genotyping primer forward	This paper	PCR primers	5'-tgcaattaaacagccaatgtg-3'
sequence-based reagent	il34 genotyping primer reverse	This paper	PCR primers	5'-ctgagtcacagccctcaaatc-3'
sequence-based reagent	il34 sequencing primer forward	This paper	PCR primers	5'-ccatttgtttttacctgaccaaa-3'
sequence-based reagent	il34 g sequencing primer reverse	This paper	PCR primers	5'-gctaattggtgtgggacgtt-3'
commercial assay or kit	In Situ Cell Death Detection Kit, Fluorescein	Roche	Cat# 11684795910	
commercial assay or kit	Sepasol-RNA/Super G	Nacalai tesque	Cat# 09379	

commercial assay or kit	ReverTra Ace®☐ aPCR master mix with gDNA remover	Toyobo	Cat# FSQ-301	
chemical compound, drug	BrdU	Nacalai tesque	Cat# 05650-95	
chemical compound, drug	Bodipy ceramide	Thermo Fisher Scientific	Cat# B22650	
chemical compound, drug	TO-PRO™-3 Iodide (642/661)	Thermo Fisher Scientific	Cat# T3605	
chemical compound, drug	N-Phenyl thiourea (PTU)	Nacalai tesque	Cat# 27429-22	
chemical compound, drug	Dextran, Alexia Flour-488	Thermo Fisher Scientific	Cat# D22910	
software, algorithm	GraphPad Prism	GraphPad Software	Ver 8.2.1	https://www.graphpad.com/scientific-software/prism/
software, algorithm	IMARIS	Bitplane	Ver 9.1.2	http://www.bitplane.com/imaris; RRID: SCR_007370
software, algorithm	Image-J	NIH	2.0.0-rc-69/1.52p)	
software, algorithm	Trimomatic	PMID: 24695404	v0.39	http://www.usadellab.org/cms/?page=trimmomatic

software, algorithm	PRINSEQ lite	PMID: 21278185	v0.20.4	http://prinseq.sourceforge.net/
software, algorithm	HISAT2	PMID: 25751142	v2.1.0	https://github.com/DaehwanKimLab/hisat2
software, algorithm	featureCounts	PMID: 24227677	Packaged ub Subread v1.5.2	http://subread.sourceforge.net/
software, algorithm	EdgeR	PMID: 19910308	v3.13	https://bioconductor.org/packages/release/bioc/html/edgeR.html
software, algorithm	EnhancedVolcano package	other	v1.6.0	https://github.com/kevinblighe/EnhancedVolcano
software, algorithm	pheatmap package	other	v1.10.12	https://cran.r-project.org/web/packages/pheatmap/index.html

A Eye anatomy

32h — interface space between nr and lens — nr, lens, cf

36h — blood vessel — nr, lens, cf

42h — blood vessel — nr, lens, cf

48h — blood vessel — lens, INL, RGCL, cf

54h — blood vessel — ONL, INL, RGCL, lens, cf

B mpeg1.1:EGFP

32h   36h   42h   48h   54h

D EF1-α: mCherry-zGem / mpeg1.1:EGFP

32h   36h   42h   48h   54h

C Number of ocular mpeg1.1:EGFP⁺ cells

(n) 0–40 — 32h 36h 42h 48h 54h

E Number of mpeg1.1:EGFP⁺ cells in S/G2/M and G1 phase

S/G2/M phase (mCherry-zGem; mpeg1.1:EGFP)
G1 phase (mpeg1.1:EGFP)

(n) — 32h 36h 42h 49h 54h

G Number of BrdU⁺ mpeg1.1:EGFP⁺ cells per retina

(n) ** p=0.0082 — mpeg1.1:EGFP cells / BrdU+ mpeg1.1:EGFP cells

F BrdU / mpeg1.1:EGFP / nuclei

54 h

H Fraction of BrdU⁺mpeg1.1:EGFP⁺ cells in total mpeg1.1EGFP⁺ cells

(%) 0–100

Figure 1

Figure 2

A confocal scanning

Superficial layer Deep layer

a b c

B

Superficial layer ←→ Deep layer

EF1-α: mCherry-zGem / mpeg1.1:EGFP

42h a' c'

48h a'' c''

C Location of mpeg1.1:EGFP cells

7.37% 6.13%

42 hpf 48 hpf

■ zGem-positive area
■ zGem-negative area

D

Superficial layer ←→ Deep layer Magnified

mfap4:TdT-CAAX / ath5:EGFP

36h a' b' c' b'

42h a'' b'' c'' a''

48h a''' b''' c''' b'''

Figure 3

Figure 4

Figure 5

Figure 6