

Structural basis of PETISCO complex assembly during piRNA biogenesis in *C. elegans*

Cecilia Perez-Borrajero¹, Nadezda Podvalnaya^{2,3#}, Kay Holleis^{4#}, Raffael Lichtenberger⁴, Jérôme Basquin⁵, Janosch Hennig^{1*}, René F. Ketting^{2,6*}, Sebastian Falk^{4†}

these authors contributed equally to this work

¹Structural and Computational Biology Unit, EMBL Heidelberg, Meyerhofstraße 1, 69117, Heidelberg, Germany

²Biology of Non-coding RNA Group, Institute of Molecular Biology (IMB), Mainz, Germany

³International PhD Programme on Gene Regulation, Epigenetics & Genome Stability, Mainz, Germany

⁴Department of Structural and Computational Biology, Max Perutz Labs, University of Vienna, 1030 Vienna, Austria

⁵Max-Planck-Institute of Biochemistry, Department of Structural Cell Biology, Am Klopferspitz 18, 82152 Martinsried, Germany

⁶Institute of Developmental Biology and Neurobiology, Johannes Gutenberg University, 55099 Mainz, Germany

*Correspondence should be addressed to Sebastian Falk, René F. Ketting or Janosch Hennig

Sebastian Falk sebastian.falk@univie.ac.at

René F. Ketting r.ketting@imb-mainz.de

Janosch Hennig janosch.hennig@embl.de

†Lead contact: Sebastian Falk sebastian.falk@univie.ac.at

Running title: Structural and biochemical analysis of PETISCO

Keywords: piRNA, 21U RNA, Piwi, *C. elegans*, PETISCO, RRM domain, ERH-2, IFE-3, TOFU-6, PID-3, TOST-1, PID-1

Abstract

Piwi-interacting RNAs (piRNAs) constitute a class of small RNAs that bind PIWI proteins and are essential to repress transposable elements in the animal germline, thereby promoting genome stability and maintaining fertility. *C. elegans* piRNAs (21U RNAs) are transcribed individually from minigenes as precursors that require 5' and 3' processing. This process depends on the PETISCO complex, consisting of four proteins: IFE-3, TOFU-6, PID-3, and ERH-2. We employ biochemical and structural biology approaches to characterize the PETISCO architecture and its interaction with RNA, together with its effector proteins TOST-1 and PID-1. These two proteins define different PETISCO functions: PID-1 governs 21U processing whereas TOST-1 links PETISCO to an unknown process essential for early embryogenesis.

Here, we show that PETISCO forms an octameric assembly with each subunit present in two copies. Determination of structures of the TOFU-6/PID-3 and PID-3/ERH-2 subcomplexes, supported by *in vivo* studies of subunit interaction mutants, allows us to propose a model for the formation of the TOFU-6/PID-3/ERH-2 core complex, and its functionality in germ cells and early embryos. Using NMR spectroscopy, we demonstrate that TOST-1 and PID-1 bind to a common surface on ERH-2, located opposite its PID-3 binding site, explaining how PETISCO can mediate different cellular roles.

Introduction

RNA molecules typically require processing after transcription before becoming fully functional. In eukaryotes, messenger RNAs (mRNAs) and many non-coding RNAs are spliced, capped, and poly-adenylated by processing factors (Hocine et al. 2010). Transcripts can also be chemically modified, trimmed or cleaved by ribonucleases, or extended by the addition of non-templated nucleotides (Roundtree et al. 2017; Yu and Kim 2020). Such processing steps are crucial for activation, (de)stabilization, localization, and many other aspects relevant to RNA function.

Piwi-interacting RNAs (piRNAs) constitute one of the largest classes of non-coding RNA transcripts whose processing is only starting to be understood (Weick and Miska 2014; Ozata et al. 2019). piRNAs play a crucial role in the germline, where they act as specificity factors in genome defense pathways with transposable elements as major targets (Luteijn and Ketting 2013; Czech and Hannon 2016). The proteins guided by piRNAs, Piwi proteins, are an animal-specific subgroup of the Argonaute family. Piwi proteins are guided by piRNAs that bind to their target site, leading to either transcript cleavage or modification of chromatin, depending on the subclass of Piwi proteins involved (Luteijn and Ketting 2013). In either case, the sequence of the piRNAs bound by the Piwi proteins dictates the target specificity of the silencing process and is therefore crucial for function. Hence, the mechanism(s) that act in piRNA precursor selection and processing determine the specificity of Piwi proteins.

The precursors of piRNAs are single-stranded RNA transcripts. In *Caenorhabditis elegans* (*C. elegans*), these are produced from a multitude of miniature genes, each producing a small transcript of 27-30 nucleotides (Ruby et al. 2006; Gu et al. 2012). Mature piRNAs are bound by the Piwi protein PRG-1 and are typically 21 nucleotides long with an uracil base at the 5' end. For these reasons, the piRNAs in *C. elegans* are often named 21U RNAs. To form mature 21U

RNAs, the precursor transcripts are shortened at both ends, including the removal of the 5' cap and trimming of the 3' end (Ruby et al. 2006; Wang and Reinke 2008; Batista et al. 2008; Das et al. 2008; Gu et al. 2012; Weick and Miska 2014; Tang et al. 2016). The vast majority of 21U RNAs stem from dedicated loci characterized by a specific sequence motif in their promoter termed the Ruby motif (Ruby et al. 2006; Cecere et al. 2012; Weick et al. 2014). Within the *C. elegans* genome, these loci are strongly clustered, suggesting that they may act in concert (Ruby et al. 2006). The transcription of these loci, including the termination of their transcription, bears hallmarks of small nuclear RNA (snRNA) biogenesis, suggesting that the *C. elegans* piRNA system has evolutionary connections to these non-coding snRNAs that play essential roles in splicing (Kasper et al. 2014; Beltran et al. 2019; Weng et al. 2019; Beltran et al. 2020; Berkyurek et al. 2021).

Following the genetic identification of several 21U RNA processing genes (Goh et al. 2014; Albuquerque et al. 2014), we and others identified a four-member protein complex required for 21U RNA biogenesis, which we named PETISCO (PID-3, ERH-2, TOFU-6, IFE-3 small RNA complex) (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019). PETISCO interacts with and stabilizes 21U RNA precursors. Interestingly, PETISCO was shown to be additionally required for early embryogenesis, a function independent of 21U RNA biogenesis (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019). In this case, loss of PETISCO function leads to a so-called 'maternal effect lethal' (Mel) phenotype, in which first-generation homozygous mutant animals develop normally, but their offspring arrest in embryogenesis. At the molecular level, low levels of the splice-leader transcript SL1, a small nuclear RNA (snRNA) involved in trans-splicing, were found to be bound by PETISCO. SL1-derived 21U RNAs have also been described, albeit at very low levels (Gu et al. 2012). Whether these findings relate to the Mel phenotype of

PETISCO mutants is unclear, but they do strengthen the link between 21U RNAs and snRNAs in *C. elegans*. The two described functions of PETISCO are specified by two different effector proteins, PID-1 and TOST-1. PID-1:PETISCO mediates 21U RNA biogenesis (Albuquerque et al. 2014), while TOST-1:PETISCO is required for early embryogenesis (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019).

The PETISCO subunits contain domains often present in RNA-binding proteins (Cordeiro Rodrigues et al. 2019). PID-3 and TOFU-6 are restricted to the nematode phylum and contain two domains. PID-3 has a predicted RNA-recognition (RRM) domain and an Argonaute-related middle (MID) domain. TOFU-6 contains an RRM domain, an extended Tudor (eTudor) domain, and a C-terminal eIF4E interaction motif. IFE-3 is one of the five highly conserved *C. elegans*' eIF4E homologs (Keiper et al. 2000), and binds to the C-terminus of TOFU-6 (Cordeiro Rodrigues et al. 2019). Finally, ERH-2 is one of the two *C. elegans* paralogs of 'enhancer of rudimentary' (Erh), a factor that is conserved throughout eukaryotes (Weng and Luo 2013). Erh was shown to participate in the RNA exosome-mediated degradation of meiotic RNAs in *Schizosaccharomyces pombe* (*S. pombe*) (Sugiyama et al. 2016) and to facilitate miRNA processing in human cells (Fang and Bartel 2020; Hutter et al. 2020; Kwon et al. 2020).

An approximate architecture of PETISCO was previously derived from yeast two-hybrid (Y2H) studies (Fig. 1A) (Cordeiro Rodrigues et al. 2019). However, the structural basis of PETISCO assembly and its interaction with the effector proteins PID-1/TOST1 and RNA substrates remain poorly understood, limiting our understanding of PETISCO function.

Here, we study PETISCO assembly using a bottom-up approach with purified proteins, interaction studies, and structural analyses. We find that PETISCO forms a dimer of tetramers, in which dimerization is mediated both by PID-3 and ERH-2. Crystal structures of the PID-

3/TOFU-6 and ERH-2/PID-3 subcomplexes reveal insights into PETISCO assembly, function, and subcellular localization. Using NMR spectroscopy, we also characterize the mutually exclusive interplay of ERH-2 with the two effector proteins TOST-1 and PID-1. These results represent the first structural characterization of a piRNA biogenesis complex, and we start to reveal how PETISCO may execute its dual role *in vivo*.

Results

PETISCO forms an octameric assembly

PETISCO consists of the proteins IFE-3, TOFU-6, PID-3, and ERH-2. Previous Y2H experiments (Cordeiro Rodrigues et al. 2019) revealed a linear topology in which IFE-3 binds to TOFU-6, which in turn binds PID-3, which associates with ERH-2 (Fig. 1A, Supplemental Fig. 1A). To analyze the oligomeric state and stoichiometry of PETISCO, we recombinantly expressed and purified PETISCO components from bacterial cells and subjected the complex to size-exclusion chromatography (SEC) coupled to multiangle light scattering (MALS) (Fig. 1B). The complete PETISCO complex eluted as a single peak and showed an average molecular mass of 236 kDa as determined by SEC-MALS (Fig. 1B), twice the sum of the individual components (assuming a 1:1:1:1 complex, 116 kDa), suggesting that PETISCO forms a hetero-octameric assembly, a dimer of tetramers, with 2:2:2:2 stoichiometry (Fig. 1B). Next, we set out to determine which protein or domains mediate oligomerization. The linear topology of PETISCO suggested that the complex could be divided into IFE-3/TOFU-6 and PID-3/ERH-2 subcomplexes (Fig. 1A). We individually purified these and analyzed their molecular mass by SEC-MALS (Fig. 1C). In the case of IFE-3/TOFU-6, we measured an average mass of 69 kDa, in line with the calculated mass of a heterodimer. The PID-3/ERH-2 subcomplex, however, had an average mass of 92 kDa, consistent with that of a hetero-tetramer, suggesting that the PID-3/ERH-2 module mediates oligomerization. Since the human and fission yeast ERH orthologs have been shown to form homodimers (Wan et al. 2005; Xie et al. 2019; Hazra et al. 2020), we hypothesized that ERH-2 is responsible for dimerization. We, therefore, determined the oligomeric state of both ERH-2 and PID-3 separately. However, PID-3 full-length protein could not be prepared in a quality suitable for SEC-MALS, and therefore we analyzed the PID-3 MID and RRM domains individually, denoted here as PID-3^{MID} and PID-3^{RRM}, respectively.

We determined average masses of 19, 28, and 27 kDa for PID-3^{MID}, PID-3^{RRM}, and full-length ERH-2^{FL}, respectively (Fig. 1D,E). This suggests that PID-3^{MID} is monomeric, whereas both PID-3^{RRM} and ERH-2 are homodimers (Fig. 1E). We thus concluded that both PID-3^{RRM} and ERH-2 contribute to PID-3/ERH-2 subcomplex dimerization, and through binding of two IFE-3/TOFU-6 subcomplexes, this results in the formation of the octameric PETISCO.




Figure 1. PETISCO assembles into a hetero-octamer with 2:2:2:2 stoichiometry. (A) Top: schematic representation of the PETISCO topology and subunit interactions and binding of the effector proteins TOST-1 and PID-1. Bottom: domain organization of IFE-3, TOFU-6, PID-3 and ERH-2 from *C. elegans*. Rounded rectangles indicate predicted domains. The asterisk marks the position of the IF4E interaction motif that mediates binding of TOFU-6 to IFE-3. (B-D) SEC-MALS chromatograms showing UV absorption at 280 nm and the calculated molecular mass in kilodaltons (kDa). The UV absorption signal was normalized to the highest peak. (B) SEC-MALS profile of PETISCO. The inset shows a Coomassie-stained SDS polyacrylamide gel of PETISCO after SEC. (C) SEC-MALS profiles of the IFE-3/TOFU-6 (black line) and PID-3/ERH-2 (orange line) subcomplexes. (D) SEC-MALS profiles of ERH-2 (black line), PID-3^{MID} (purple line) and PID-3^{RRM} (orange line) domains. (E) Summary of molecular masses and stoichiometries of PETISCO and its subunits.

Crystal structure of the TOFU-6/PID-3 RRM core complex

The topological arrangement of PETISCO places TOFU-6 and PID-3 at the core, and we thus proceeded to narrow down their interacting regions. Our previous experiments indicated that the interaction between TOFU-6 and PID-3 is mediated by the RRM domains (denoted as TOFU-6^{RRM} and PID-3^{RRM}) (Fig. 1A, Supplemental Fig. S1A) (Cordeiro Rodrigues et al. 2019). To better map which domains mediate the interaction between TOFU-6 and PID-3, we used a combination of pull-down experiments and SEC. We recombinantly co-expressed maltose binding protein (MBP)-tagged TOFU-6 constructs with glutathione-S-transferase (GST)-tagged PID-3 constructs. Both MBP and GST pull-downs revealed that the RRM domains of PID-3 and TOFU-6 mediate the interaction between the two proteins and that neither the eTudor domain of TOFU-6 (TOFU-6^{eTUDOR}) nor the PID-3^{MID} are required (Supplemental Fig. S2A). This is supported by SEC using purified TOFU-6^{RRM} and PID-3^{RRM} proteins (Supplemental Fig. S2B).

To gain structural insights into the TOFU-6^{RRM}/PID-3^{RRM} complex, we determined the structures of PID-3^{RRM} and the TOFU-6^{RRM}/PID-3^{RRM} complex, at 1.8 Å and 1.7 Å resolution, respectively (Fig. 2A and Supplemental Fig. S2C, Supplemental Table S1). Both structures are very similar. The PID-3^{RRM} forms a homodimer in both cases, with slight differences in the relative orientation of the RRM domains. (Supplemental Fig. S2C). Here, we will focus on the analysis of the TOFU-6^{RRM}/PID-3^{RRM} complex structure, containing one tetrameric TOFU-6^{RRM}/PID-3^{RRM} complex in the asymmetric unit (Fig. 2A), consistent with the SEC-MALS analysis (Supplemental Fig. S2D). The PID-3^{RRM} adopts a canonical RRM fold with an antiparallel four-stranded β-sheet packing opposite two α-helices (α1 and α2) (Supplemental Fig. S2E). The TOFU-6^{RRM} has a similar architecture but contains an additional fifth β-strand β4* located between the α2 and β4 elements (Supplemental Fig. S2F). The PID-3^{RRM} dimerizes

via the $\alpha 1$ helix by a combination of hydrophobic and polar interactions (Fig. 2A and 2B) reminiscent, for instance, of the dimer interface in HuR-RRM3 (Pabis et al. 2019; Ripin et al. 2019) and RBPMS (Teplova et al. 2016). Phe217 from one PID-3^{RRM} protomer packs in a hydrophobic pocket created by Phe217, Ala220, Val228, and Ile231 from the other protomer. Moreover, Gln221 forms a hydrogen bond with the polypeptide backbone of the neighboring protomer (Fig. 2B).

Two TOFU-6 RRM domains bind on either side of the interfaces created through the PID-3^{RRM} dimerization and contact both PID-3 protomers (Fig. 2A). The TOFU-6 $\alpha 1$ helix, $\beta 2$ strand, and $\beta 3$ - $\alpha 2$ loop form a surface that interacts with the $\alpha 1$ - $\beta 2$ loop and $\alpha 2$ helix from the PID-3^{RRM} (Interface I/protomer 1) by a combination of hydrophobic and polar interactions (Fig. 2C). The TOFU-6^{RRM} interaction with the second PID-3^{RRM} protomer involves electrostatic interactions between the TOFU-6^{RRM} N-terminus and the $\alpha 2$ helix with the PID-3^{RRM} $\alpha 2$ - $\beta 5$ loop (Interface II/protomer 2), featuring a salt bridge between TOFU-6 Asp67 and PID-3 Arg263, as well as a hydrogen bond between TOFU-6 Asp12 and PID-3 Asn264 (Fig. 2D).

Our structural analysis suggests that dimerization of the PID-3^{RRM} is a prerequisite for TOFU-6^{RRM} binding and that two interfaces at the two protomers contribute to the interaction. To test this prediction, we engineered a series of substitutions both at the PID-3 dimerization interface (PID-3^{RRM} A220E) and at interface I (TOFU-6^{RRM} F30E and M61E; PID-3^{RRM} F247E/Q251R) and analyzed their impact on the interaction by bacterial co-expression pull-down experiments. We verified that the A220E mutation renders PID-3^{RRM} monomeric by SEC-MALS (Supplemental Fig. S2G). Mutations at the PID-3^{RRM} dimerization surface and interface I disrupted TOFU-6 binding, suggesting that they contribute to a stable association with TOFU-6^{RRM} (Fig. 2E).




Figure 2. The TOFU-6 and PID-3 RRM domains form the PETISCO core. (A) Crystal structure of the PID-3^{RRM}/TOFU-6^{RRM} complex shown in cartoon representation in two orientations related by the 90° rotation about the horizontal axis. The two PID-3^{RRM} protomers are shown in different shades of pink, while the two TOFU-6^{RRM} protomers are shown in different shades of orange. The N- and C-terminal residues are highlighted. (B) Zoomed-in view of the homodimerization interface of PID-3^{RRM}. Interacting residues are shown in stick representation and labeled. (C, D) Zoomed-in view of a representative set of residues at the PID-3/TOFU-6 interaction interface I (C) and interface II (D). Interacting residues are shown in stick representation as indicated. (E) Analysis of the effect of structure-guided mutations on the PID-3/TOFU-6 interaction by GST pull-down assays. Wild-type (WT) and mutant versions of MBP-tagged TOFU-6 constructed were co-expressed with WT and mutant versions of GST-tagged PID-3 constructs. Input and elution fractions were analyzed by SDS-PAGE followed by Coomassie staining. (F) Box-plot showing the percentage of hatched embryos of *pid-3::gfp(wt)* and *pid-3[a220e)::gfp* animals grown at 20 and 25°C. Progeny of 30 different mothers were analyzed for each condition, and development of at least 2600 eggs was

scored. (G,H) Single plane confocal micrographs of PID-3::GFP(WT) and PID-3[A220E]::GFP at 25°C. The boxes indicate the regions (above the spermatheca) from which three zoomed-in examples are shown below. Scale bars: 20 μm in overview, 8 μm in zoom-in.

Next, we tested the effect of the monomer-inducing A220E mutation in PID-3 *in vivo*. Using CRISPR-Cas9 mediated gene editing, we first created a strain expressing C-terminally GFP-tagged PID-3, such that *in vivo* expression could be monitored by fluorescence microscopy. We then introduced the A220E mutation and scored its effect on subcellular localization and embryonic viability. *pid-3(a220e)* animals showed a strong Mel phenotype, which was fully penetrant at 25°C (Fig. 2F), consistent with loss of PETISCO function. In addition, PID-3(A220E) did not form peri-nuclear foci (Fig. 2G, 2H), suggesting that most likely TOFU-6 mediates the previously described P granule localization of PETISCO (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019).

RNA-binding properties of the TOFU-6 and PID-3 RRM domains

To investigate the RNA-binding properties of the TOFU-6^{RRM}/PID-3^{RRM} complex, we compared the RRM domains of both TOFU-6 and PID-3 to well-characterized RRMs using structure-based sequence alignments (Pei et al. 2008). RRM domains bind single-stranded RNA molecules via the outer β -sheet surface with the contribution of two conserved motifs called RNP1 and RNP2, located in the β_3 and β_1 strand, respectively (Maris et al. 2005). The residues critical for RNA binding, located in RNP1 and RNP2, are retained in TOFU-6^{RRM}, whereas the PID-3^{RRM} shows several differences, including the absence of aromatic residues in RNP1 and RNP2 (Fig. 3A). This analysis suggested that TOFU-6^{RRM}, but not PID-3^{RRM}, might bind RNA. We tested this hypothesis using fluorescence anisotropy assays to quantitatively measure the affinity of PID-3^{RRM}, TOFU-6^{RRM}, and the PID-3^{RRM}/TOFU-6^{RRM} complex for a Cy5-labeled 16-mer oligo(U) RNA. The resulting binding isotherms were fitted with a single binding site

model. PID-3^{RRM} showed only weak interaction with RNA, whereas TOFU-6^{RRM} bound RNA with a K_d of 60 μ M. The tetrameric PID-3^{RRM}/TOFU-6^{RRM} complex bound RNA with higher affinity (K_d 10 μ M) than the isolated TOFU-6^{RRM}. The higher affinity for RNA could be a result of oligomerization of TOFU-6^{RRM} in the tetramer or due to the contribution of additional contacts from PID-3^{RRM}. We modeled RNA onto the PID-3^{RRM}/TOFU-6^{RRM} complex guided by known RRM-RNA structures (Upadhyay and Mackereth 2020; Auweter et al. 2006; Teplova et al. 2016). The RNA is bound in a specific orientation with the 5' end located near the β 1/ β 4 region of the β -sheet and aromatic residues in RNP1 and RNP2 interact with the nucleobases of the RNA by stacking interactions (Maris et al. 2005). When superposing the four RRMs of the PID-3^{RRM}/TOFU-6^{RRM} complex with an RNA-bound RRM, we saw that all RNA molecules were located on the same side of the PID-3^{RRM}/TOFU-6^{RRM} complex (Fig. 3C). Even though PID-3^{RRM} alone does not strongly bind RNA, the electrostatic surface potential revealed a positively charged region on the PID-3^{RRM} surface, in close vicinity to the TOFU-6^{RRM} RNA binding site. One might envisage the presence of an extended RNA path reaching from the TOFU-6^{RRM} to the PID-3^{RRM}.




Figure 3. RNA binding properties of the PID-3 and TOFU-6 RRM domains. (A) Structure-based multiple sequence alignment of PID-3^{RRM} and TOFU-6^{RRM} in comparison to well-characterized RNA-binding RRMs of RBM20, FOX-1

and RBPMS. The RNP1 and RNP2 motifs contributing to RNA binding are indicated and differences are highlighted with pink triangles. (B) Fluorescence anisotropy binding assay using an U16-mer RNA labeled with a Cy5-label at the 5' end. Values are presented as average and error bars correspond to the standard deviation of three ($n=3$) technical replicates. (C) Model showing the potential RNA binding sites of the PID-3^{RRM}/TOFU-6^{RRM} complex. The PID-3^{RRM}/TOFU-6^{RRM} complex is colored as in Fig. 2 and shown as an electrostatic surface potential. Blue and red indicate positively and negatively charged regions, respectively; while white represents neutral surfaces. The positioning of the four modelled RNA molecules are shown in ribbon presentation. RNAs on the TOFU-6^{RRM} and PID-3^{RRM} are shown in yellow and green, respectively. Positively charged regions on the PID-3^{RRM} are highlighted by green ovals.

ERH-2 binds to a region upstream of the PID-3 RRM domain

Next, we investigated the interaction between PID-3 and ERH-2. ERH-2 consists of an enhancer of rudimentary (ERH) domain followed by a C-terminal region of ~ 15 amino acids (Fig. 1A). Previous experiments (Zeng et al. 2019; Cordeiro Rodrigues et al. 2019) suggested a direct interaction between ERH-2 and PID-3, in particular between the PID-3^{RRM} and ERH-2 (Cordeiro Rodrigues et al. 2019). However, the PID-3 constructs used in the latter study contained additional N-terminal and C-terminal regions flanking the PID-3^{RRM} (Supplemental Fig. 1A). To better define the ERH-2 binding site of PID-3, we used PID-3 constructs covering the RRM domain (PID-3^{RRM}) or the RRM domain with an N-terminal extension (PID-3^{RRM-N}) and performed bacterial co-expression pull-down experiments. While PID-3^{RRM-N} pulled down ERH-2, PID-3^{RRM} failed to bind ERH-2 (Fig. 4A). To rule out that the C-terminal extension downstream of the PID-3 RRM domain contributed to binding, we performed SEC experiments. We incubated PID-3^{RRM-N} and PID-3^{RRM-C} with ERH-2 and analyzed the mixtures by SEC to assess binding. This confirmed that PID-3^{RRM-N} did, but PID-3^{RRM-C} did not interact with ERH-2 (Supplemental Fig. S3A,B). We then asked if the N-terminal extension of the PID-3 RRM alone was sufficient for ERH-2 binding. We purified a GST-tagged PID-3 peptide (PID-3^{pep}, residues 171-203) corresponding to this region and found that ERH-2 could indeed bind to GST-PID-3^{pep} (Fig. 4A). To obtain quantitative insights into PID-3^{pep}/ERH-2 interaction, we used isothermal titration calorimetry (ITC). We determined a dissociation constant (K_d) of 0.65

μM and a stoichiometry $N \sim 1$ (0.97) (Supplemental Fig. S3C, Table 1), consistent with the 2:2 stoichiometry observed in the case of the full-length PID-3/ERH-2 complex by SEC-MALS (Fig. 1C). Finally, we tested if PID-3^{pep} binding specificity may explain our previous observation that PETISCO specifically incorporates ERH-2, and not its close paralogue ERH-1 (36% sequence identity, 60% similarity; Supplemental Fig. S3D, (Cordeiro Rodrigues et al. 2019)). In GST-pull-down experiments, ERH-1 did not bind GST-PID-3^{pep} (Supplementary Fig. 3E), indicating that PID-3^{pep} can discriminate between the two ERH paralogs. We concluded that the N-terminal extension of the PID-3^{RRM} domain binds to ERH-2, and is likely responsible for the ERH-2 specificity of PETISCO.

Structural insights into the formation of the ERH-2/PID-3 complex

We next determined the crystal structures of a C-terminally truncated ERH-2 (residues 1-99, ERH-2^{ΔC}) in the free and PID-3^{pep} bound states, at a resolution of 1.50 Å and 2.17 Å, respectively (Fig. 4B, Supplemental Table S1). The overall structure of ERH-2^{ΔC} is similar to human ERH (Wan et al. 2005) and *S. pombe* Erh (Xie et al. 2019) (Supplemental Fig. S4A). ERH-2^{ΔC} adopts a mixed α - β -fold, with a four-stranded antiparallel β -sheet that packs against three amphipathic α -helices on the backside (Fig. 4B). The front side of the β -sheet mediates dimerization which results in the formation of a pseudo- β -barrel structure. The structure of free ERH-2^{ΔC} is very similar to PID-3^{pep}-bound ERH-2^{ΔC}, with the exception that the loop connecting helices α 1 and α 2 is not visible in the electron density, suggesting it becomes ordered upon PID-3^{pep} binding. (Supplemental Fig. S4B).




Figure 4. Structural and functional analysis of the PID-3/ERH-2 interaction. (A) GST pull-down assays. GST-tagged PID-3 constructs were co-expressed with Trx-tagged ERH-2 in bacterial cells and then subjected to co-precipitation using glutathione-coupled beads. Input and elution fraction were analyzed on SDS-PAGE gels with Coomassie brilliant blue staining. (B) Top: The crystal structure of the ERH-2 Δ C/PID-3^{pep} complex is shown in cartoon representation. The two ERH-2 protomers are shown in different shades of blue, while the two PID-3 peptides are in different shades of pink. Bottom: Zoomed-in view of the interaction between ERH-2 and PID-3^{pep}. Interacting residues are shown in stick representation. (C) Binding of PID-3^{pep} to ERH-2 monitored by NMR spectroscopy. Inset: upon incremental addition of PID-3^{pep}, the ¹H-¹⁵N amide signals of ¹⁵N-labeled ERH-2 move in slow exchange

from their free (shown in red), to their bound state (in maroon) with a 2.3-fold molar excess of PID-3^{pep}. Two representative zoomed-in regions of the overlaid spectra are shown (full view in Supplementary Fig. S5B). The arrows indicate the direction of the change in CSP from free to bound states. The CSP values determined for a saturated 1:3 ERH-2:PID-3^{pep} complex are plotted below as a function of residue number. The most highly perturbed amino acid, Thr21, is indicated. The grey dashed line corresponds to the average CSP of all residues. The secondary structure elements as found in the crystal structure are shown above. (D) Mutational analysis of the PID-3/ERH-2 interface. GST-tagged PID-3 wild-type and mutants were co-expressed with wild-type or mutant Trx-tagged wild-type or mutant ERH-2 in bacterial cells as described in (A). Pink circles highlight the bands corresponding to the bait and blue circles the prey in the input. (E) Box-plot showing the percentage of hatched embryos of *pid-3::gfp(wt)* and *pid-3[i182a;v186a]::gfp* and *pid-3[i182e;v186e]::gfp* animals grown at 20 and 25°C. Progeny of 30 different mothers were analyzed for each condition, and development of at least 2600 eggs was scored. (F) Single-plane confocal micrographs of PID-3::GFP(WT) and PID-3[I182A; V186A]::GFP at 25°C. The boxes indicate the regions (above the spermatheca) from which three zoomed-in examples are given below. Scale bars: 20 μm in overview, 8 μm in zoom-in.

The PID-3^{pep} is unfolded in solution (Supplemental Fig. S4C) but forms an amphipathic α -helix when bound to ERH-2^{AC} (Fig. 4B, Supplemental Fig. S4D). PID-3^{pep} shows well-ordered electron density for residues 177-193 (Chain C) and 179-193 (Chain D) and occupies a similar surface on ERH as DGCR8 in the human ERH/DGCR8^{peptide} complex (Kwon et al. 2020), and Mmi1 in the fission yeast Erh1/Mmi1^{peptide} complex (Xie et al. 2019) (Supplemental Fig. 4A). However, while both Mmi1 and DGCR8 bind as extended peptides lacking defined secondary structure, PID-3^{pep} binds as an α -helix (Supplemental Fig. S4A). The hydrophobic interface of the amphipathic PID-3^{pep} helix is formed by Ile182, Val186, Phe187, Val189, and Leu190 and points towards a hydrophobic groove in ERH-2 formed by strands β 1, β 2, as well as the α 1 helix (Fig. 4B and Supplemental Fig. S4E). In addition to the hydrophobic interactions, hydrogen bonds formed between PID-3^{pep} Ser185 and ERH-2 Asp41, as well as between PID-3^{pep} His191 and ERH-2 Thr21 further contribute to the affinity and specificity of the interaction. In a complementary approach, we employed NMR spectroscopy and assigned the backbone chemical shifts (¹H-¹⁵N, C ^{α} , and C ^{β}) of free full-length ERH-2. We used these chemical shifts to determine the secondary structural elements of ERH-2, which were fully consistent with the crystal structure, and also showed a disordered C-terminus (residues 104-113, Supplemental

Fig. S5A). In addition, the loop region between $\alpha 1$ and $\alpha 2$ encompassing residues 46-55 was found to contain two sets of amide peaks. This indicates the presence of two well-defined alternative conformations that exchange in the millisecond timescale, and explains the lack of order in the crystal structure of free ERH-2^{ΔC} (Supplemental Fig. 4B).

We then monitored the interaction between PID-3^{pep} and ERH-2. Unlabeled PID-3^{pep} was titrated into ¹⁵N-labeled ERH-2, and the positions of the amide chemical shifts were measured at each point by ¹H-¹⁵N HSQC experiments (Fig. 4C and Supplemental Fig. S5B). We observed binding in the slow exchange regime, whereby peaks disappear from one location and reappear at a new position corresponding to the bound complex. This indicates a strong interaction with a dissociation constant in the low micromolar to nanomolar range, in agreement with the ITC data (Supplemental Fig. S3C). We next assigned the chemical shifts of the ERH-2/PID-3^{pep} complex and calculated the chemical shift perturbations (CSPs). The ERH-2 residues most affected by the interaction with PID-3^{pep} were located in $\beta 2$, with Thr21, Trp22, and Gly23 exhibiting the largest CSPs (Fig. 4C). Mapping of the amide CSPs onto the crystal structure of ERH-2^{ΔC} was consistent with the binding interface seen in the crystal structure (Supplemental Fig. S5C). Taken together, the crystal structure and NMR data provide a clear view for the basis of the ERH-2/PID-3 interaction.

Mutational analysis of the ERH-2/PID-3 interface

To test the relevance of residues found at the complex interface, we used GST-pulldown experiments with purified proteins. Mutation of PID-3 Ile182 and Ser185 to glutamate residues (I182E; S185E), completely abrogated binding of PID-3^{pep} to ERH-2 (Fig. 4D), and even milder substitutions of PID-3 Ile182 and Val186 to alanine (I182A; V186A) were sufficient to abolish binding (Fig. 4D). Mutation of Thr13 in ERH-2, which lines the hydrophobic pocket (Fig. 4B), to glutamate (T13E), also abrogated binding to PID-3^{pep} (Fig. 4D). *In vivo*, PID-3(I182E; V186E)

worms displayed a full Mel phenotype at 20°C (Fig. 4E), while the weaker PID-3(I182A; V186A) mutations showed a Mel phenotype only at 25°C. Finally, at the subcellular level, loss of ERH-2 binding did not affect the localization of PID-3 (Fig. 4F). These results strongly support the *in vivo* relevance of the structures that we determined, but also suggest that additional interactions between ERH-2 and PID-3 may exist within the full PETISCO complex.

ERH-2 interacts with the effector proteins TOST-1 and PID-1 through a common interface

PID-1 and TOST-1 share a common sequence motif with which they interact with ERH-2, as previously assessed through Y2H and MS analysis (Fig. 5A) (Cordeiro Rodrigues et al. 2019). To understand the interplay of ERH-2 with these proteins, we purified full-length TOST-1 and PID-1, C-terminally tagged with monomeric Venus StrepII tag (mVenus-Strep), and probed their binding to ERH-2 and ERH-1. Both PID-1 and TOST-1 bound ERH-2 (Fig. 5B), but unlike PID-3, they also interacted with ERH-1 (Fig. 5B). We noted as well that TOST-1 was a more efficient bait than PID-1 (Fig. 5B), suggesting that it binds more strongly to ERH-2 and ERH-1 than PID-1.

To define the binding interface at the residue level, we analyzed the interaction of PID-1 and TOST-1 with ERH-2 using NMR spectroscopy. We titrated peptides corresponding to the conserved region of TOST-1 (residues 28-53, TOST-1^{pep}) and PID-1 (residues 47-74, PID-1^{pep}) into ¹⁵N-labeled full-length ERH-2 to monitor amide CSPs (Fig. 5C and Supplemental Fig, S6A, B). The changes occurred in slow-intermediate exchange for TOST-1^{pep}, and in intermediate exchange for PID-1^{pep} (Figure 5C). This indicates binding constants in the micromolar range for both peptides, with PID-1^{pep} binding being weaker than TOST-1^{pep}. To determine where PID-1^{pep} and TOST-1^{pep} bind ERH-2, we assigned the backbone chemical shifts of the respective complexes and calculated their amide CSPs (Fig. 5D).




Figure 5. PID-1^{pep} and TOST-1^{pep} bind to ERH-2 through a common interface. (A) Sequence alignment of *C. elegans* PID-1 and TOST-1 corresponding to the conserved motif previously determined (Cordeiro Rodrigues et al. 2019). (B) Analysis of the interaction between TOST-1/PID-1 with ERH-1/ERH-2 by Streptavidin pull-down assays. Purified, recombinant TOST-1 or PID-1 mVenus-Strep fusion proteins were used as baits and ERH-1 and ERH-2 as prey. Input and elution fractions were analyzed by SDS-PAGE followed by Coomassie staining. (C) PID-1^{pep} and TOST-1^{pep} bind to ERH-2 in a similar fashion. NMR-monitored titrations of ¹⁵N-labeled ERH-2 upon addition of unlabeled TOST-1 and PID-1 peptides, as indicated (full view in Supplemental Fig. S6). (D) The bound ERH-2/peptide complexes (1:3 molar ratios) were assigned and the extent of the CSPs quantified. The peptides produced

changes in similar directions (indicated by the arrows), and generally included the same set of perturbed amino acids. (E) Mapping of the CSP values onto the dimeric structure of ERH-2 (PDB: 7O6N) shows that the most affected residues lie in strand β 4 and adjacent helix α 3, opposite the interface bound by PID-3^{pep}. (F) Interplay of TOST-1^{pep} and PID-1^{pep} binding to ERH-2. Comparison of ¹⁵N-labeled ERH-2 moieties in its free form (red), bound to PID-1^{pep} (green, 1:3 molar ratio), TOST-1^{pep} (yellow, 1:3), in the presence of equimolar amounts of both peptides (blue, 1:3:3), or in the presence of excess TOST-1^{pep} (magenta, 1:3:7). (G) Pull-down assays with purified recombinant wild-type and mutant versions of ERH-2 as prey and TOST-1-mVenus-Strep as bait (streptavidin pull-down) or GST-PID-3^{pep} as bait (GST pull-down). Input and elution fractions were analyzed by SDS-PAGE followed by Coomassie staining.

The changes observed occurred in similar residues and mostly in the same direction, indicating that both peptides bound in analogous fashion and to the same surface of ERH-2 (Fig. 5C). Both peptides caused the largest perturbations at the interface formed by strand β 4, including the highly perturbed amino acids Ile81, Gly82, and Arg83, as well as the adjacent α 3 helix, including Trp87 (Fig. 5 D, E). Mapping of the largest CSPs on the surface of ERH-2 showed that the interface bound by TOST-1/PID-1 laid opposite the interface used for PETISCO binding through PID-3^{pep} (Fig. 5E). Of note, we observed only one set of peaks for both the ERH-2 homodimer (save for the flexible loop), as well as the protein-peptide complexes. This suggests a 2:2 stoichiometry for protein to effector peptide, as any other scenario would break the symmetry required to observe only one species in solution. We then explored the PID-1 and TOST-1 interplay on ERH-2, and compared spectra of free ERH-2 and in the presence of PID-1^{pep}, TOST-1^{pep}, as well as mixtures of both. When the peptides were present at the same concentrations, we observed chemical shifts at intermediate locations between that of ERH-2/TOST-1^{pep} and ERH-2/PID-1^{pep} complexes (Fig. 5F). However, once TOST-1^{pep} was present in excess, we observed chemical shifts consistent with that of an ERH-2/TOST-1^{pep} complex only (Fig. 5F, Supplemental Fig. S7). Since the peptides associate at the same interface, we thus concluded that PID-1 and TOST-1 compete for ERH-2 binding, and their relative concentrations will determine which of the two complexes will be favored. Despite the larger size of the ERH-2/TOST-1^{FL} complex (60 kDa), we detected very similar

amide CSPs upon addition of full-length TOST-1 (TOST-1^{FL}) to ¹⁵N-labeled ERH-2, indicating that the shorter peptide recapitulates full-length TOST-1 binding (Supplemental Fig. S8).

To validate the PID-1/TOST-1 binding interface observed by NMR, we designed two ERH-2 mutants (I81A/R83A and W87A), and performed pull-down experiments with purified proteins. Both ERH-2 mutants, in particular W87A, showed a weaker interaction with TOST-1 (Fig. 5G). These mutations in ERH-2 did not affect PID-3^{pep} binding, and conversely, the mutation in ERH-2 (T13E) that disrupted association with PID-3 did not significantly affect TOST-1 binding (Fig. 5B), consistent with their binding at opposite sides.

Analysis of the interplay between TOST-1 and PID-3 upon binding to ERH-2

Our analysis suggested that ERH-2 can simultaneously bind TOST-1 and PID-3. To test this, we performed pull-down assays with TOST-1-mVenus-Strep as bait and ERH-2 alone, as part of an ERH-2/PID-3 subcomplex, or with full PETISCO as prey. We found that TOST-1 pulled down ERH-2 in all conditions (Fig. 6A). We observed that more ERH-2 is pulled down when ERH-2 is in complex with PID-3 and as part of PETISCO, indicating that the presence of PID-3 might facilitate the binding to TOST-1. The formation of a trimeric complex was supported by SEC experiments with purified full-length proteins, as a sample containing ERH-2, TOST-1, and PID-3 elutes at smaller elution volumes than the dimeric subcomplexes (Fig. 6B).

To obtain quantitative insights, we performed ITC measurements. TOST-1^{FL} and TOST-1^{pep} bound to ERH-2 with dissociation constants of 13 and 23 μ M (Table 1, Supplemental Fig. S9). The binding of TOST-1^{FL} and TOST-1^{pep} differed modestly by a factor of two, which is consistent with the NMR experiments (Supplemental Fig. S8). Similar affinities were obtained when the experiments were performed in the presence of PID-3^{pep} (Table 1 or Supplemental Fig. S9). Taken together, the results from the pull-down, SEC, and ITC experiments allowed

us to conclude that ERH-2 forms a binding platform for simultaneous binding of PID-3^{pep} and TOST-1^{pep}.




Figure 6. PID-3 and TOST-1 associate with ERH-2 simultaneously. (A) Analysis of the interaction between TOST-1 with PETISCO components. Pull-down experiments with purified, recombinant TOST-1-mVenus-Strep fusion as bait with the indicated preys. Input and elution fractions were analyzed by SDS-PAGE followed by Coomassie staining. (B) Interactions between ERH-2, TOST-1, and PID-3 were assessed by SEC. Purified proteins were incubated alone or in the indicated mixtures in 1:1 ratio and subjected to SEC. (C) The formation of a ternary ERH-2/TOST-1^{pep}/PID-3^{pep} complex shown by NMR spectroscopy. Zoomed-in views of overlaid ^1H - ^{15}N HSQCs of ERH-2 in its free form (red), in complex with TOST-1^{pep} (blue, 1:4 molar ratio), in complex with PID-3^{pep} (orange, 1:3 molar ratio), or in the presence of both (green, 1:4:1.5 ERH-2:TOST-1^{pep}:PID-3^{pep}). Residues Thr21 (T21), Glu26 (E26), and Gly82 (G82) are indicated, with their positions color coded according to the same scheme. The black arrows highlight amide peaks present in the ternary complex that do not correspond to either the ERH-2/TOST-1^{pep} or

ERH-2/PID-3^{pep} subcomplexes. (D) Schematic model of the PETISCO core complex (TOFU-6^{RRM}/PID-3^{RRM}/ERH-2) and its interaction with the effector proteins PID-1 and TOST-1. TOFU-6^{RRM}/PID-3^{RRM} form a tetrameric core and ERH-2 binds to a region upstream of the PID-3^{RRM}. TOST-1 and PID-1 bind to a common surface on ERH-2, located opposite to its PID-3 binding site, and thereby specify PETISCO function.

We also used NMR spectroscopy to monitor the interplay of TOST-1^{pep} and PID-3^{pep} on ERH-

2. Upon addition of PID-3^{pep} to a saturated ERH-2/TOST-1^{pep} complex, we observed changes in amide positions of ¹⁵N-labeled ERH-2 to locations consistent with the formation of an ERH-2/PID-3^{pep} complex. However, signature chemical shifts corresponding to TOST-1-bound ERH-2 remained, suggesting the formation of a ternary complex (Fig. 6C, full view in Supplemental Fig. S10). Consistent with this idea, we observed the appearance of new peaks not present in either the ERH-2/TOST-1^{pep} or ERH-2/PID-3^{pep} subcomplexes (Fig. 6C).

Interestingly, some amide peaks diagnostic of the ERH-2/TOST-1^{pep} interaction became stronger in the presence of PID-3^{pep} (Fig. 6C), suggesting that PID-3^{pep} stabilizes the interaction of TOST-1^{pep} and ERH-2. Given that the two binding sites are connected through a central β -sheet, this synergy in binding could be explained by an allosteric mechanism. Taking together, these results clearly show that both PID-3^{pep} and TOST-1^{pep} can simultaneously associate with ERH-2 to form a trimeric complex and that the association of ERH-2 with PID-3^{pep} facilitates its interaction with TOST-1^{pep}.

Discussion

PETISCO - Oligomeric state

We show that PETISCO is an octameric protein complex, consisting of two copies of each IFE-3, TOFU-6, PID-3, and ERH-2 with a mass of ~236 kDa. We find that the PID-3^{RRM}/TOFU-6^{RRM} complex binds a model RNA substrate with higher affinity than the isolated TOFU-6^{RRM}, suggesting that oligomerization enhances RNA binding properties, but the functional consequence of PETISCO dimerization for piRNA processing remains to be determined. However, there are interesting parallels with other RNA processing complexes. First, it was recently demonstrated that the yeast and human THO (TREX) complexes involved in the transcription and export of RNA also associate into higher-order oligomers (Schuller et al. 2020; Pühringer et al. 2020). Whereas the yeast THO complex forms a dimer, the human THO complex forms a tetramer (Schuller et al. 2020; Pühringer et al. 2020). Although the functional consequence of oligomerization remains to be shown experimentally, it was hypothesized that dimerization of the yeast THO complex plays a crucial role in preventing R-loop formation during transcription of mRNA by RNA polymerase II (Schuller et al. 2020). In addition, dimerization is required for the function of the *D. melanogaster* SFiNX complex, which facilitates co-transcriptional gene silencing downstream of the piRNA-PIWI complex. Dimerization of SFiNX is promoted through the interaction with the dynein light chain protein 'Cup up'/LC8, which forms a homodimer and binds a short linear motif present in one of the SFiNX subunits. On the functional level, dimerization of SFiNX is required for the nucleic acid-stimulated formation of biomolecular condensates *in vitro* and heterochromatin formation at piRNA target loci *in vivo* (Schnabl et al. 2021). In the case of SFiNX, dimerization thus plays a direct role in the formation of condensates. Since the processing of piRNAs also takes place in biomolecular condensates, the P granules, dimerization might play a role in modulating the phase separation properties of PETISCO.

Finally, the human ERH homolog has recently been shown to support the microprocessor complex during microRNA (miRNA) processing (Fang and Bartel 2020; Hutter et al. 2020; Kwon et al. 2020). Microprocessor is a trimeric complex consisting of one copy of DROSHA and two copies of DGCR8 and dimerization of DGCR8 is necessary for miRNA processing (Faller et al. 2007). The dimeric ERH protein binds to a short linear motif located in the N-terminal region of DGCR8 and is thought to additionally contribute to DGCR8 dimerization (Kwon et al. 2020). At the functional level, oligomerization facilitates processing of suboptimal miRNA hairpins located in clusters (Fang and Bartel 2020; Hutter et al. 2020). In PETISCO, we find a similar architecture: PID-3 dimerizes through its RRM domain and is supported by the binding of ERH-2 to a motif upstream of the RRM (Figure 6D). Within PETISCO, ERH-2 not only reinforces dimerization, but also binds the effector proteins TOST-1 and PID-1 at a surface opposite of PID-3. Similarly, SAFB has been shown to bind human ERH (Drakouli et al. 2017) and help process suboptimal miRNAs (Hutter et al. 2020; Fang and Bartel 2020). It thus appears that ERH homologs may act as signal integrators to control RNA processing. Because human ERH stimulates processing of suboptimal miRNA precursors, we hypothesize that PETISCO dimerization may likewise stimulate 21U RNA processing, either by modulating the RNA binding or phase separation properties of PETISCO. However, dissecting whether these effects stem from dimerization or PETISCO integrity will be difficult, given the inter-dependence of PID-3 dimerization and TOFU-6 binding.

Subcellular localization of PETISCO

We find that loss of ERH-2 binding by PID-3 does not affect its subcellular distribution over the cytoplasm and P granules. However, we show that dimerization of PID-3 is a prerequisite

for TOFU-6 binding and for localization of PID-3 to P granules. Previous work from Zeng et al. (2019) showed that an allele of *tofu-6(ust95)* that results in partial deletion of the eTudor domain specifically affected TOFU-6 P granule localization and 21U RNA production, but not embryonic viability. Together, these results strongly imply that TOFU-6 plays an important role in P granule localization. Interestingly, the depletion of PID-3 by RNAi leads to loss of TOFU-6 from P granules (Zeng et al. 2019), suggesting that PID-3 can somehow enhance TOFU-6's ability to mediate this function. Whether this relates to the dimerization we describe, or to other effects, cannot currently be resolved.

Interplay between TOST-1 and PID-1

We find that TOST-1 can outcompete PID-1 for PETISCO binding when present in excess. Is this relevant for PETISCO functionality *in vivo*? A conclusive answer to this question will require an experiment in which the onset of expression of PID-1 and TOST-1 in the various stages of germ cell development can be assessed. Due to the stability of the fluorescent tags, and their large size with respect to the proteins of interest, standard localization studies under steady-state conditions (Zeng et al. 2019) are not suitable to resolve this. However, we do know that PID-1 does not have a role in the embryo, and that 21U RNAs are already expressed at early stages, as judged from PRG-1 expression patterns (Batista et al. 2008; Wang and Reinke 2008). At the same time, germ cells do not show any defects in *tost-1* mutants, while it is present in, and required for the development of embryos (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019). Interestingly, in *tost-1* mutants, some increase in 21U RNA levels have been reported (Cordeiro Rodrigues et al. 2019; Zeng et al. 2019), indicating a competition between the two PETISCO functions. Everything considered, it seems likely that PID-1-bound PETISCO assemblies are present in germ cells when TOST-1 starts to be upregulated, and that the PID-

1-TOST-1 exchange that we describe is relevant to establish a good balance between PETISCO functions *in vivo*.

Evolutionary aspects

Evolutionary analysis of PETISCO components revealed that PID-3, TOFU-6, TOST-1, and PID-1 are restricted to nematodes, while ERH (ERH-2) proteins and eIF4E proteins (IFE-3) are present throughout eukaryotes (Cordeiro Rodrigues et al. 2019). *C. elegans* contains two ERH paralogs, ERH-1 and ERH-2. Interestingly, while TOST-1 and PID-1 can interact with both ERH-paralogs, PID-3 only interacts with ERH-2. Also, TOST-1 is present in nematode species that only contain one ERH paralog (Cordeiro Rodrigues et al. 2019), suggesting that in these species the cognate TOST-1 binding partner might be ERH-1. There are also several nematode species that contain PID-3 but no ERH-2 paralog. Even though in *C. elegans* PID-3 does not bind ERH-1, it would be interesting to see whether PID-3 can interact with ERH-1 in those species, possibly representing an ancestor complex to what we now see in *C. elegans*.

Analysis of sequence conservation revealed that the most conserved feature of the PID-3^{RRM} domain is the surface that mediates homo-dimerization. In contrast, the residues involved in TOFU-6 binding are less well conserved (Supplemental Fig. S11A). On the other hand, most residues of the TOFU-6^{RRM} involved in PID-3^{RRM} binding are well conserved (Supplementary Figure S11B). This is consistent with our previous evolutionary analysis, which revealed that PID-3 is more widespread than TOFU-6 in the nematode phylum (Cordeiro Rodrigues et al. 2019). It might be that the TOFU-6 binding interface's evolutionary pressure is reduced compared to the PID-3 dimerization interface, or that TOFU-6 is replaced with another protein, possibly also an RRM domain protein, in species that have PID-3 but no TOFU-6.

These aspects clearly show that ERH is a nexus around which many RNA processing reactions concentrate, and which is amenable to significant variation in interacting proteins. The many parallels between ERH function in miRNA processing and PETISCO show that small RNA biogenesis pathways represent one domain of RNA processing that exploits ERH as an interaction platform. However, given the conservation of ERH also in species lacking small RNAs, such as *S. cerevisiae*, and the role of ERH in RNA decay in *S. pombe*, it is clear that ERH proteins fill a niche in RNA processing that is much more general (Weng and Luo 2013).

Table 1: Interplay of TOST-1 and PID-3 binding to ERH-2.

Titrand (cell)	Titrant (syringe)	K_d (μM)	N	ΔH (kcal/mol)	-TAS (kcal/mol)
ERH-2	PID-3 ^{pep}	0.65 ± 0.08	0.97	-12.3 ± 0.1	3.9 ± 0.2
ERH-2	TOST-1 ^{pep}	23 ± 10^a	1 ^b	-14 ± 5^a	8 ± 5^a
ERH-2	TOST-1 ^{full}	13 ± 1^a	1 ^b	-12 ± 1^a	5.4 ± 0.5^a
ERH-2: PID-3 ^{pep} 1:2	TOST-1 ^{pep}	4 ± 2^a	1 ^b	-13 ± 5^a	6 ± 6^a
ERH-2: PID-3 ^{pep} 1:2	TOST-1 ^{full}	15 ± 2^a	1 ^b	-16 ± 1^a	9.3 ± 0.6^a

^aThe values reported correspond to the average value and standard deviation of the mean of two measurements at 25 °C. ^bThe N value was fixed at 1 due to difficulties accurately determining the concentrations of the TOST-1^{pep} lacking aromatic amino acids. One representative binding isotherm for each is shown in Supplemental Fig. S9

Material and Methods

Protein production

The genes coding for PETISCO subunits (IFE-3, TOFU-6, PID-3, ERH-2), TOST-1, PID-1 and ERH-1 were cloned into modified pET vectors using ligation independent cloning. All proteins were produced as an N-terminal His-Tagged fusion protein with varying fusion partners. IFE-3, ERH-1, ERH-2, PID-1 and TOST-1 contained a His₆-Trx-3C tag, and TOFU-6 a His₁₀-MBP-3C tag and PID-3 a His₆-GST-3C tag. Addition of 3C protease allowed to cleave this His-fusion protein tag from the protein of interest. Proteins were produced in the *E. coli* BL21(DE3) derivatives strain in terrific broth medium. Briefly, cells were grown at 37°C, and when the culture reached an optical density (OD) at 600 nm of 2-3, the temperature was reduced to 18°C. After 2 h at 18°C, 0.2 mM IPTG was added to induce protein production for 12-16 h overnight. To reconstitute PETISCO, IFE-3 was co-expressed with TOFU-6 and PID-3 with ERH-2. Cells were lysed by sonication, and PETISCO was purified by immobilized metal affinity chromatography (IMAC) using Ni²⁺-chelating beads. The His-fusion tags were cleaved by the addition of 3C protease, and the His-fusion tag and His-tagged 3C protease were removed by a second IMAC step. PETISCO was subsequently purified using Heparin affinity chromatography and size-exclusion chromatography in a buffer containing 20 mM Tris/HCl pH 7.5, 150 mM NaCl, 10% (v/v) glycerol, 2 mM DTT.

Size-exclusion chromatography assay

Purified proteins were incubated alone or in different combinations as indicated, in concentrations between 20-40 µM (total volume of 200 µl) in SEC buffer (20 mM Tris/HCl pH 7.5, 150 mM NaCl, 2 mM DTT). Samples were incubated for 1 h on ice to allow complex formation. Complex formation was assayed by comparing the elution volumes in SEC on a

Superdex 200 Increase 10/300 column (GE Healthcare). The SEC peak fractions were analyzed by SDS-PAGE and visualized by Coomassie brilliant blue staining.

Size-exclusion chromatography coupled to multi-angle light scattering (SEC MALS)

The molecular mass and the oligomeric state of PETISCO, its subunits and other proteins in solution were determined by size exclusion chromatography (SEC) coupled to multi-angle light scattering (MALS). Individual proteins or protein complexes were analyzed at concentrations between 2 and 5 mg/mL in a buffer consisting of 20 mM Tris/HCl pH 7.5, 150 mM NaCl, 2 mM DTT. Superdex 200 Increase 10/300 GL column (GE Healthcare Life Sciences) was connected to a 1260 Infinity HPLC system (Agilent Technologies) coupled to a MiniDawn Treos detector (Wyatt Technologies) with a laser emitting at 690 nm. An RI-101 detector (Shodex) was used for refractive index measurement. Data analysis was performed using Astra 7 software package (Wyatt Technologies).

ERH-2^{ΔC} and PID-3^{pep} production and purification

ERH-2^{ΔC} containing residues 1-99 of *C. elegans* ERH-2 was cloned into a modified pET-vector with a 6xHis-Thioredoxin (Trx) tag followed by a 3C protease cleavage site. PID-3^{pep} containing residues 171-203 of *C. elegans* PID-3 was cloned into a modified pET-vector with a 6xHis-GST tag followed by a 3C protease cleavage site. Cell growth and protein production was performed as described above. ERH-2^{ΔC} and PID-3^{pep} were purified by immobilized metal affinity chromatography using a Ni²⁺ matrix. The His-Trx and His-GST tags were cleaved with His-tagged 3C protease, and both the tags and 3C protease were removed in a reverse IMAC step. ERH-2^{ΔC} and PID-3^{pep} were present in the flow-through and were concentrated using ultrafiltration and subjected to size-exclusion chromatography on a Superdex 75 (16/600)

column equilibrated with 20 mM Tris/HCl pH 7.5, 150 mM NaCl, and 5 mM 2-mercaptoethanol. After size-exclusion chromatography, ERH-2^{ΔC} was concentrated to 15 mg ml⁻¹ and PID-3^{pep} to 5 mg ml⁻¹ and stored at -80°C until further use.

PID-3^{RRM} and TOFU-6^{RRM} production and purification

TOFU-6^{RRM} (residues 1-99) and PID-3^{RRM} (residues 196-274) were cloned into a modified pET-vector containing His₁₀-MBP-3C and His₆-GST-3C tags, respectively. TOFU-6^{RRM} and PID-3^{RRM} were expressed individually as described for ERH-2^{ΔC} and PID-3^{pep}. To purify the PID-3^{RRM}/TOFU-6^{RRM} complex, the cultures expressing individual PID-3^{RRM} and TOFU-6^{RRM} proteins were mixed before cell harvest. The PID-3^{RRM}/TOFU-6^{RRM} complex was purified by immobilized metal affinity chromatography using a Ni²⁺ matrix, followed by cation-exchange chromatography and size-exclusion chromatography on a Superdex 200 (16/600) column equilibrated in 20 mM Tris/HCl pH 7.5, 150 mM NaCl, and 5 mM 2-mercaptoethanol.

Crystallization

All crystallization trials were performed using a vapor diffusion set-up by mixing the protein complex and crystallization solution in a 1:1 and 2:1 ratio. For ERH-2^{ΔC}, the best diffracting crystals grew in condition A2 from the Morpheus screen (Molecular Dimensions). For crystallization of the ERH-2^{ΔC}/PID-3^{pep} complex, these were mixed in a 1:1 ratio. The best diffracting crystals grew in condition A5 from the JCSG+ (Molecular Dimensions). The best crystals of PID-3^{RRM} were obtained in condition C5 from the Morpheus screen (Molecular Dimensions). Crystals were harvested and directly frozen in liquid nitrogen before data collection at 100 K. Crystals from ERH-2^{ΔC}, ERH-2^{ΔC}/PID-3^{pep}, and the PID-3^{RRM} grew in

conditions that did not require further cryoprotection and were therefore directly frozen in liquid nitrogen before data collection at 100 K.

Initial small crystals for the PID-3^{RRM}/TOFU-6^{RRM} complex were obtained in several commercial screens in conditions containing sodium acetate buffer at pH 4-5 and various precipitants and additives. Through several rounds of micro-seeding, we obtained larger crystals. The best crystals grew in 0.1 M Sodium acetate pH 5.0, 200 mM NaCl and 17% (v/v) PEG3350. Crystals were soaked with a mother liquor containing 300 mM NaI instead of 200 mM NaCl, and with 25% (v/v) glycerol for cryoprotection, and then frozen in liquid nitrogen.

Data processing, phase determination, refinement, and modelling

All data were processed with Xia2/Dials (Winter et al. 2013) within CCP4i2 (Potterton et al. 2018). For ERH-2^{AC} the phases were determined by molecular replacement using the human ERH structure (PDB 2nml) as model, while in the case of the ERH-2^{AC}/PID-3^{pep} complex, we used the *C. elegans* ERH-2^{AC} structure (PDB 7O6L). The phases of the PID-3^{RRM}/TOFU-6^{RRM} complex were solved by single isomorphous replacement with anomalous scattering (SIRAS) phasing from iodide with Autosol from Phenix (Terwilliger et al. 2009). The mean figure of merit over all resolution shells had a value of 0.30 and an estimated map correlation coefficient value of 40 ± 11 . The phases for the PID-3^{RRM} were determined by molecular replacement using *C. elegans* the PID-3^{RRM} domain from the PID-3^{RRM}/TOFU-6^{RRM} complex structure (PDB 7OCZ) as model. The models were automatically built using Buccaneer (Cowtan 2006) or Autobuild (Terwilliger et al. 2008), manually completed with COOT (Emsley et al. 2010), and refined with phenix.refine (Afonine et al. 2012) and refmac5 (Kovalevskiy et al. 2018). PDB redo (Joosten et al. 2012) and molprobity (Williams et al. 2018) were used for validation. Data collection,

phasing, and refinement statistics are listed in Supplemental Table S1. Molecular graphics of the structures were created using UCSF ChimeraX (Goddard et al. 2018).

Pull-down assays with purified proteins

For interaction studies with purified proteins, appropriate protein mixtures (bait 10-20 μ M, prey in 1.2-fold molar excess) were incubated in 20 mM Tris/HCl (pH 7.5), 150 mM NaCl, 10% (v/v) glycerol, 0.05% (v/v) NP40, 1 mM DTT for 30 min at 4°C. The protein mixtures were then incubated with the indicated beads: Glutathione sepharose beads (Cube Biotech), Amylose sepharose beads (New England Biolabs)), and Streptactin XT beads (IBA) for 2 h. Post incubation, the beads were washed three times with 0.2 mL incubation buffer, and the retained material was eluted with 0.05 mL incubation buffer supplemented with 20 mM reduced glutathione, 20 mM maltose, or 50 mM biotin. Input material and eluates were analyzed by SDS-PAGE and Coomassie brilliant blue staining.

Co-expression pull-down assays

For interaction studies by the co-expression co-purification strategy, the proteins of interest were co-expressed in BL21(DE3) derivative strains as described above. Cell pellets were resuspended in 2 ml lysis buffer (50 mM Sodium phosphate, 20 mM Tris/HCl, 250 mM NaCl, 10 mM Imidazole, 10% (v/v) glycerol, 0.05% (v/v) NP-40, 5 mM beta-mercaptoethanol pH 8.0) per gram of wet cell mass. Cells were lysed by ultrasonic disintegration, and insoluble material was removed by centrifugation at 21,000xg for 10 min at 4°C. For GST pull-downs, 500 μ L of supernatant was applied to 20 μ L glutathione-coupled resin (Cube Biotech); for MBP pull-downs, 500 μ L supernatant was applied to 20 μ L amylose resin (New England Biolabs) and incubated for two hours at 4°C. Subsequently, the resin was washed three times with 500 μ L

of lysis buffer. The proteins were eluted in 50 μ L of lysis buffer supplemented with 20 mM reduced glutathione or 20 mM maltose in the case of glutathione or amylose beads, respectively. Input material and eluates were analyzed by SDS-PAGE and Coomassie brilliant blue staining.

Isothermal titration calorimetry

ITC experiments were carried out using the PEAQ-ITC Isothermal titration calorimeter (Malvern). The data were processed and curves were fitted using the PEAQ-ITC software. Before the measurements, all samples were dialyzed simultaneously against 1 L of sample buffer (50 mM Tris, 250 mM NaCl, 1 mM TCEP, pH 7.50).

PID-3^{pep} - ERH-2 interaction: ERH-2 (the reactant) samples were concentrated to 23 μ M and PID-3 peptide (the injectant) to 250 μ M. Titrations were carried out at 25°C with 2 μ L of the injectant per injection added to 200 μ L of reactant cell solution. For titrations involving TOST-1, the cell contained ~ 300 μ L of either ERH-2 at 20 μ M, or the ERH-2/PID-3^{pep} complex mixed in a 1:2 ratio (20 μ M ERH-2 and 40 μ M PID-3^{pep}) pre-equilibrated at 25 °C. The TOST-1^{pep} and TOST-1^{FL} molecules were concentrated to 600 and 300 μ M, respectively, and ~60 μ L placed in the syringe as the titrant. Due to difficulties in accurately measuring the TOST-1^{pep} concentration (low extinction coefficient at 280 nm), this value was floated and the N value fixed at 1. The heats of dilution were obtained by repeating the experiments with sample buffer in the calorimeter cell, while keeping all other experimental parameters identical. These dilution heats were fit to a straight line by linear regression and subtracted from the heats of binding. The baseline-corrected binding isotherms were then fit to a 1:1 model to give the reported thermodynamic parameters. For the purposes of comparing binding of TOST-1 in the presence and absence of PID-3^{pep}, the heats of association were assumed to derive solely from

TOST-1 binding. For multiple measurements, the averages and population standard deviations of the fit results (K_d , $\Delta H^{\circ}_{\text{bind}}$, and $-\Delta S^{\circ}_{\text{bind}}$) are reported.

Fluorescence anisotropy experiments

U16 RNA carrying a 5' Cy5-label was purchased from Ella Biotech (Martinsried, Germany). 100 nM of RNA and increasing concentrations of protein were incubated in a volume of 20 μL in a buffer containing 10 mM Tris/HCl pH 7.5, 50 mM KCl, and 0.5 mM EDTA at 15°C for at least 2 h. The anisotropy was recorded on a FS5 Spectrofluorometer (Edinburgh Instruments) with the following settings: Excitation 649 nm, bandwidth 3 nm; Emission 675 nm, bandwidth 3 nm. The anisotropy was measured 5 times in 30 s, and the average anisotropy value was calculated.

Production of fractionally deuterated $^{13}\text{C}/^{15}\text{N}$ -labeled ERH-2

The gene encoding full-length *Caenorhabditis elegans* ERH-2 (NCBI gene ID: 185323) was cloned into pET28a (Addgene: 69864-3), containing an N-terminal His₆ tag followed by a TEV protease cleavage site (MGSSHHHHHHSSGENLYFQGHMAS) using NheI and BamHI restrictions sites. Due to the size of the dimeric ERH-2 protein (~ 27 kDa), fractional deuteration in combination with stable isotope labeling was used for recombinant protein expression in *Escherichia coli* BL21 (λDE3) to allow NMR triple resonance experiments for backbone assignments ($^{13}\text{C}^{\alpha}$, $^{13}\text{C}^{\beta}$, ^{15}NH). A starter 50-mL culture was grown at 37°C for 16 hours with shaking at 200 rpm, using M9 minimal medium (dissolved in H₂O supplemented with 2g/L of ^{12}C -glucose and 0.5 g/L $^{14}\text{NH}_4\text{Cl}$). The cells were collected by centrifugation and resuspended in 500 mL of M9 minimal medium dissolved in ~ 98 % D₂O supplemented with 2g/L ^{13}C -glucose and 0.5 g/L $^{15}\text{NH}_4\text{Cl}$, followed by incubation for 16 hours at 37°C. The resulting culture

was used to inoculate additional 2.5 L of minimal M9/D₂O/¹³C/¹⁵N medium, to a starting OD₆₀₀ of ~ 0.3. After growth at 37°C, protein expression was induced at an OD₆₀₀ of ~ 1 with 2 mM IPTG, and the culture was allowed to shake for 16 hours at 20 °C. Subsequently, the cells were collected by centrifugation, resuspended in lysis buffer (20 mM Na₂HPO₄, 20 mM imidazole, 1 M NaCl, 1 M urea, 10 % v/v glycerol, pH 7.4), and frozen at -20 degrees until use. The cell pellet was then thawed, and the lysis buffer supplemented with lysozyme (Genaxxon Bioscience) and protease inhibitor cocktail (Roche) according to instructions. After 15-minute incubation on ice, the mixture was homogenized (Microfluidics Corp. Microfluidizer M-110L Fluid Processor) by passing four times. Unless otherwise noted, all purification steps were performed at room temperature. The lysate was cleared by centrifugation, and applied to tandem fast-flow Ni²⁺-NTA HisTrap (GE Healthcare) columns. These were washed with 20 column volumes of lysis buffer. His-tagged ERH-2 was eluted in one step with ~35 mL of 20 mM Na₂HPO₄, 500 mM imidazole, and 0.5 M NaCl, at pH 7.4, and dialyzed against 2 L of TEV cleavage buffer (50 mM Tris, 250 mM NaCl, 1 mM DTT, pH 7.5) for 30 minutes at room temperature. At this point, ~ 1 mg of TEV protease was added, and ERH-2 was allowed to continue dialyzing in its presence at 4°C for 16 hours. The resulting cleaved product contained five non-native residues at the N-terminus and was further purified using a second HisTrap, followed by size exclusion chromatography column (Superdex 75 16/600, GE Healthcare) equilibrated with NMR buffer (50 mM Tris, 250 mM NaCl, 1 mM TCEP, pH 7.50). The appropriate pure fractions were pooled, and the protein was concentrated using Amicon 3 kDa MWCO centrifugal filters (Merck Millipore). The protein concentration was calculated using the measured absorbance at 280 nm under native conditions and assuming an extinction coefficient of 19940 M⁻¹ cm⁻¹.

NMR spectroscopy

NMR experiments were performed at 25°C using a Bruker Avance III 800 MHz spectrometer equipped with a cryoprobe. The ERH-2 sample was concentrated to 200-600 micromolar and contained 0.02% sodium azide, 10 % D₂O (as a locking agent), and 90 % NMR buffer. For backbone assignments and titrations, 2D and 3D transverse relaxation optimized spectroscopy (TROSY) pulse sequences with ²H-decoupling and apodization weighted sampling were utilized (Pervushin et al. 1997; Salzman et al. 1998; Simon and Köstler 2019). The resulting spectra were processed and analyzed using NMRPipe (Delaglio et al. 1995) and NMR-FAM Sparky (Lee et al. 2015), respectively. 90% of all amides were assigned. The resulting chemical shifts (¹H^N, ¹⁵N, ¹³C^α, and ¹³C^β) were deposited under BMRB: 50914 and used as input for secondary structure prediction using the program MICS (Shen and Bax 2012) (Supplementary Figure S5A). To assign the peptide-bound ERH-2 complexes, NMR-monitored titrations were used in combination with TROSY-HNCA experiments to track the new positions of the amide peaks (Salzman et al. 1998).

Peptides and NMR-monitored titrations

The lyophilized TOST-1 and PID-1 peptides were purchased from Proteogenix at 95 % purity. Approximately 10 mg of each peptide was resuspended in ~ 0.8 mL of NMR buffer, and further dialyzed using Pur-A-Lyzer dialysis containers (Sigma) with a MWCO of 1 kDa, against an additional 2L of NMR buffer to ensure complete buffer match. The peptide concentration was typically ~ 2 mM, calculated using the measured absorbance at 280 nm under native conditions and assuming extinction coefficients of 2980 M⁻¹ cm⁻¹ for both TOST-1 and PID-1 peptides. The peptides were added in a stepwise manner to labeled ERH-2 at a concentration of ~ 200 μM, and the changes in chemical shifts were monitored with TROSY-HSQC experiments at each

titration point. The extent of amide ^1H - ^{15}N chemical shift perturbation (CSP) in free versus saturated peptide-bound ERH-2 were calculated according to Williamson et al. (Williamson 2013) to compensate for frequency bandwidths differences between ^{15}N and ^1H dimensions. The resulting values are plotted as a function of residue number.

List of peptides used in this study.

Name	Sequence	Length	Corresponding protein residues
TOST-1	KYKTATNKRITLNERFGVLEKGYTIQ	26	28-53
PID-1	NYNTNLHRKVTLSDRFELAALGYEMKAK	28	47-74
PID-3 short	(GPDSMW) FNSEDIKDSVFKVLHAEED	25	177-195
PID-3 long	(GPDSMW) TFDKVLFNSEDIKDSVFKVLHAEEDPRGADQEN	39	171-203

Worm methods

Worms were cultivated at 20°C on the OP50 plates. For experiments at 25°C, animals were grown at 25°C for 48 hours.

List of strains

Strain	Genotype
N2	<i>C. elegans</i> wild isolate
RFK1344	<i>pid-3(xf271[pid-3::gfp]) I.</i>
RFK1346	<i>pid-3(xf272[pid-3[A220E]::gfp]) I.</i>
RFK1348	<i>pid-3(xf273[pid-3[I182A/V186A]::gfp]) I.</i>
RFK1349	<i>pid-3(xf274[pid-3[I182E/V186E]::gfp]) I.</i>

CRISPR/Cas9 genome editing

Protospacer sequences were chosen using CRISPOR (<http://crispor.tefor.net>) cloned in pRK2412 by site-directed, ligase-independent mutagenesis (SLIM).

gene	allele	Protospacer sequence (5' to 3')
<i>pid-3</i>	<i>xf271</i>	CAAATTGGGGCACATTTAAG (AGG)
	<i>xf272</i>	GATCCATTGACACGATATG (GGG)
	<i>xf273</i>	CTTCAACAGCGAAGACATCA (AGG)
	<i>xf274</i>	CTTCAACAGCGAAGACATCA (AGG)

A PCR product from plasmid pDD282 (Addgene plasmid # 66823) was used as a donor template for insertion of *gfp*. For creation of point mutations, the following Ultramer® DNA oligodeoxynucleotides from Integrated DNA Technologies™ were used as a donor template:

<i>xf271</i>	TGGGAAAGCCGATCAGACTTTCGAAAGTCTCCTCTGGAGCATCGGGAGCCTCAGGAGCATC GATGAGTAAAGGAGAAGAATTGTTCACTGGAGTTGTCCAATCCTCGTCGAGCTCGACGGA GACGTCAACGGACACAAGTTCTCCGTCTCCGGAGAGGGAGAGGGAGACGCCACCTACGGA AAGCTCACCTCAAGTTCATCTGCACCACCGAAAGCTCCCAGTCCCATGGCCAACCCTCG TCACCACCTTCTGCTACGGAGTCCAATGCTTCTCCCGTTACCCAGACCACATGAAGCGTCAC GACTTCTTCAAGTCCGCCATGCCAGAGGGATACGTCCAAGAGCGTACCATCTTCTTCAAGG TAAGTTTAAACATATATACTAATACTACTGATTATTTAAATTTTCAGGACGACGGAAACTAC AAGACCCGTGCCGAGGTCAAGTTCGAGGGAGACACCCTCGTCAACCGTATCGAGCTCAAG GTAAGTTTAAACAGTTCGGTACTAATACTAACCATAATTTAAATTTTCAGGGAATCGACTT CAAGGAGGACGGAAACATCCTCGGACACAAGCTCGAGTACAATACTCCACAACGT CTACATCATGGCCGACAAGCAAAAAGAACGGAATCAAGGTCAACTCAAGGTAAGTTTAAA CATGATTTTACTAATACTAATACTGATTTAAATTTTCAGATCCGTCACAACATCGAGGACG GATCCGTCCAACCTCGCCGACCACTACCAACAAAACACCCCAATCGGAGACGGACCAGTCC TCCTCCAGACAACCACTACCTCTCCACCCAATCCGCCCTCTCCAAGGACCCAAACGAGAA GCGTGACCACATGGTCTCTCGAGTTCGTCACCGCCCGGAATCACCCACGGAATGGAC GAGCTCTACAAGTAAATGTGCCCAATTTGCTTATAAGTTATTTTTTCT
<i>xf272</i>	GTGGATATCCGGGATGCTCAACACGTTCCGCATCGAGCAATTGCTCACTCCATATCGTGT CAATGGGATCACCATACCGGCGCCAG
<i>xf273</i>	CGACGTTGACAAGGTTCTCTTCAACAGCGAAGACGCTAAAGATTCTGCCTTCAAAGTTCT GCATGCCGAAGAAGAGCCGAGAGGTGCGGATCAGGAGAA
<i>xf274</i>	CGACGTTGACAAGGTTCTCTTCAACAGCGAAGACGAGAAAAGATTCTGAGTTCAAAGTTCT GCATGCCGAAGAAGAGCCGAGAGGTGCGGATCAGGAGAA

In all cases *dpy-10* co-conversion was used. All alleles were outcrossed twice before experiments.

Mel Phenotype

For RFK1344, 1346 and 1348 30 young adult worms were singled and left for 24 hour (20°C) or overnight (25°C). Afterwards, mothers were removed and the eggs were counted, and after two days, developed animals were scored. RFK1349 is Mel (maternal embryonic lethal) at 20°C, therefore progeny of heterozygous mothers was singled and genotyped. At least 2200 eggs from at least 23 worms were counted for each condition.

Microscopy

Young adult worms or young gravid adults grown at 25°C were washed in M9 buffer and paralyzed by 60 mM sodium azide in M9. As soon as worms stopped moving, they were imaged on TCS SP5 Leica confocal microscope at 25°C. Images were processed using Fiji and Adobe illustrator.

Accession numbers

The coordinates and the structure factors have been deposited in the Protein Data Bank with accession codes PDB ID 7O6L, 7O6N, 7OCX, and 7OCZ. NMR backbone chemical shifts of *C. elegans* ERH-2 were deposited at the BMRB under accession number 50914.

Competing Interest Statement

The authors declare no conflict of interest.

Acknowledgments

We thank all of the members of the Falk, Ketting, and Hennig laboratories for great help and discussion. Svenja Hellmann is thanked for support in genome editing, and the IMB Media Laboratory and the Microscopy Core Facilities for equipment and support. We also thank the

beamline scientists from the ESRF beamlines ID-23-1 and ID23-2 (Grenoble, France) and Swiss light source PXII (Villigen, Switzerland) for excellent support with data collection, as well as the EMBL Biophysics Facility. The Hennig lab gratefully acknowledges support via an Emmy-Noether Fellowship of the Deutsche Forschungsgemeinschaft (DFG) (HE 7291). This study was supported by funding from the DFG via the priority program SPP1935 to J.H. (EP37/3-1 and EP37/3-2) and R.F.K. (KE1888/7-1) and lab start-up funding from the University of Vienna to S.F. C.P.B. thanks the EMBL Interdisciplinary Postdoc (EIPoD) Program fellowship under Marie Skłodowska-Curie Actions COFUND (grant number 664726) for support.

Author Contributions

C.P.B., K.H., N.P., R.L., J.B., J.H., and S.F. planned and executed experiments. K.H. and S.F. generated crystals. S.F. and J.B. collected crystallographic data, S.F. solved and modeled all crystal structures. N.P. generated and analyzed *C. elegans* strains and performed protein purifications. C.B.P. and J.H. performed and analyzed NMR experiments. S.F., R.F.K. and J.H. supervised the projects and provided funds. All authors read and approved the manuscript.

References

- Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD. 2012. Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**: 352–367.
- Albuquerque BFM de, de Albuquerque BFM, Luteijn MJ, Cordeiro Rodrigues RJ, van Bergeijk P, Waaijers S, Kaaij LJT, Klein H, Boxem M, Ketting RF. 2014. PID-1 is a novel factor that operates during 21U-RNA biogenesis in *Caenorhabditis elegans*. *Genes & Development* **28**: 683–688.
- Auweter SD, Fasan R, Reymond L, Underwood JG, Black DL, Pitsch S, Allain FH-T. 2006. Molecular basis of RNA recognition by the human alternative splicing factor Fox-1. *EMBO J* **25**: 163–173.
- Batista PJ, Ruby JG, Claycomb JM, Chiang R, Fahlgren N, Kasschau KD, Chaves DA, Gu W, Vasale JJ, Duan S, et al. 2008. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in *C. elegans*. *Mol Cell* **31**: 67–78.
- Beltran T, Barroso C, Birkle TY, Stevens L, Schwartz HT, Sternberg PW, Fradin H, Gunsalus K, Piano F, Sharma G, et al. 2019. Comparative Epigenomics Reveals that RNA Polymerase II Pausing and Chromatin Domain Organization Control Nematode piRNA Biogenesis. *Dev Cell* **48**: 793-810.e6.
- Beltran T, Pahita E, Ghosh S, Lenhard B, Sarkies P. 2020. Integrator is recruited to promoter-proximally paused RNA Pol II to generate *Caenorhabditis elegans* piRNA precursors. *EMBO J* e105564.
- Berkyurek AC, Furlan G, Lampersberger L, Beltran T, Weick E-M, Nischwitz E, Cunha Navarro I, Braukmann F, Akay A, Price J, et al. 2021. The RNA polymerase II subunit RPB-9 recruits the integrator complex to terminate *Caenorhabditis elegans* piRNA transcription. *EMBO J* n/a: e105565.
- Cecere G, Zheng GXY, Mansisidor AR, Klymko KE, Grishok A. 2012. Promoters recognized by forkhead proteins exist for individual 21U-RNAs. *Mol Cell* **47**: 734–745.
- Cordeiro Rodrigues RJ, de Jesus Domingues AM, Hellmann S, Dietz S, de Albuquerque BFM, Renz C, Ulrich HD, Sarkies P, Butter F, Ketting RF. 2019. PETISCO is a novel protein complex required for 21U RNA biogenesis and embryonic viability. *Genes Dev* 1–14.
- Cowtan K. 2006. The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta Crystallogr D Biol Crystallogr* **62**: 1002–1011.
- Czech B, Hannon GJ. 2016. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. *Trends Biochem Sci* **41**: 324–337.
- Das PP, Bagijn MP, Goldstein LD, Woolford JR, Lehrbach NJ, Sapetschnig A, Buhecha HR, Gilchrist MJ, Howe KL, Stark R, et al. 2008. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the *Caenorhabditis elegans* germline. *Mol Cell* **31**: 79–90.
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. 1995. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* **6**: 277–293.
- Drakouli S, Lyberopoulou A, Papathanassiou M, Mylonis I, Georgatsou E. 2017. Enhancer of rudimentary homologue interacts with scaffold attachment factor B at the nuclear matrix to regulate SR protein phosphorylation. *FEBS J* **284**: 2482–2500.
- Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**: 486–501.

- Faller M, Matsunaga M, Yin S, Loo JA, Guo F. 2007. Heme is involved in microRNA processing. *Nat Struct Mol Biol* **14**: 23–29.
- Fang W, Bartel DP. 2020. MicroRNA Clustering Assists Processing of Suboptimal MicroRNA Hairpins through the Action of the ERH Protein. *Mol Cell* **78**: 289–302.e6.
- Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE. 2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**: 14–25.
- Goh WSS, Seah JWE, Harrison EJ, Chen C, Hammell CM, Hannon GJ. 2014. A genome-wide RNAi screen identifies factors required for distinct stages of *C. elegans* piRNA biogenesis. *Genes and Development* **28**: 797–807.
- Gu W, Lee H-C, Chaves D, Youngman EM, Pazour GJ, Conte D Jr, Mello CC. 2012. CapSeq and CIP-TAP identify Pol II start sites and reveal capped small RNAs as *C. elegans* piRNA precursors. *Cell* **151**: 1488–1500.
- Hazra D, Andrić V, Palancade B, Rougemaille M, Graille M. 2020. Formation of *S. pombe* Erh1 homodimer mediates gametogenic gene silencing and meiosis progression. *Sci Rep* **10**: 1034.
- Hocine S, Singer RH, Grünwald D. 2010. RNA processing and export. *Cold Spring Harb Perspect Biol* **2**: a000752.
- Hutter K, Lohmüller M, Jukic A, Eichin F, Avci S, Labi V, Szabo TG, Hoser SM, Hüttenhofer A, Villunger A, et al. 2020. SAFB2 Enables the Processing of Suboptimal Stem-Loop Structures in Clustered Primary miRNA Transcripts. *Mol Cell* **78**: 876–889.e6.
- Joosten RP, Joosten K, Murshudov GN, Perrakis A. 2012. PDB_REDO: constructive validation, more than just looking for errors. *Acta Crystallogr D Biol Crystallogr* **68**: 484–496.
- Kasper DM, Wang G, Gardner KE, Johnstone TG, Reinke V. 2014. The *C. elegans* SNAPc component SNPC-4 coats piRNA domains and is globally required for piRNA abundance. *Dev Cell* **31**: 145–158.
- Keiper BD, Lamphear BJ, Deshpande AM, Jankowska-Anyszka M, Aamodt EJ, Blumenthal T, Rhoads RE. 2000. Functional Characterization of Five eIF4E Isoforms in *Caenorhabditis elegans* *. *J Biol Chem* **275**: 10590–10596.
- Kovalevskiy O, Nicholls RA, Long F, Carlon A, Murshudov GN. 2018. Overview of refinement procedures within REFMAC5: utilizing data from different sources. *Acta Crystallogr D Struct Biol* **74**: 215–227.
- Kwon SC, Jang H, Shen S, Baek SC, Kim K, Yang J, Kim J, Kim J-S, Wang S, Shi Y, et al. 2020. ERH facilitates microRNA maturation through the interaction with the N-terminus of DGCR8. *Nucleic Acids Res* **48**: 11097–11112.
- Lee W, Tonelli M, Markley JL. 2015. NMRFAM-SPARKY: enhanced software for biomolecular NMR spectroscopy. *Bioinformatics* **31**: 1325–1327.
- Luteijn MJ, Ketting RF. 2013. PIWI-interacting RNAs: from generation to transgenerational epigenetics. *Nat Rev Genet* **14**: 523–534.
- Maris C, Dominguez C, Allain FH-T. 2005. The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *FEBS J* **272**: 2118–2131.
- Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. 2019. PIWI-interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20**: 89–108.
- Pabis M, Popowicz GM, Stehle R, Fernández-Ramos D, Asami S, Warner L, García-Mauriño SM, Schlundt A, Martínez-Chantar ML, Díaz-Moreno I, et al. 2019. HuR biological function involves RRM3-mediated dimerization and RNA binding by all three RRMs. *Nucleic Acids Res* **47**: 1011–1029.

- Pei J, Kim B-H, Grishin NV. 2008. PROMALS3D: a tool for multiple protein sequence and structure alignments. *Nucleic Acids Res* **36**: 2295–2300.
- Pervushin K, Riek R, Wider G, Wüthrich K. 1997. Attenuated T2 relaxation by mutual cancellation of dipole-dipole coupling and chemical shift anisotropy indicates an avenue to NMR structures of very large biological macromolecules in solution. *Proc Natl Acad Sci U S A* **94**: 12366–12371.
- Potterton L, Agirre J, Ballard C, Cowtan K, Dodson E, Evans PR, Jenkins HT, Keegan R, Krissinel E, Stevenson K, et al. 2018. CCP4i2: the new graphical user interface to the CCP4 program suite. *Acta Crystallogr D Struct Biol* **74**: 68–84.
- Pühringer T, Hohmann U, Fin L, Pacheco-Fiallos B, Schellhaas U, Brennecke J, Plaschka C. 2020. Structure of the human core transcription-export complex reveals a hub for multivalent interactions. *Elife* **9**. <http://dx.doi.org/10.7554/eLife.61503>.
- Ripin N, Boudet J, Duszczyc MM, Hinniger A, Faller M, Krepl M, Gadi A, Schneider RJ, Šponer J, Meisner-Kober NC, et al. 2019. Molecular basis for AU-rich element recognition and dimerization by the HuR C-terminal RRM. *Proc Natl Acad Sci U S A* **116**: 2935–2944.
- Roundtree IA, Evans ME, Pan T, He C. 2017. Dynamic RNA Modifications in Gene Expression Regulation. *Cell* **169**: 1187–1200.
- Ruby JG, Jan C, Player C, Axtell MJ, Lee W, Nusbaum C, Ge H, Bartel DP. 2006. Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell* **127**: 1193–1207.
- Salzmann M, Pervushin K, Wider G, Senn H, Wüthrich K. 1998. TROSY in triple-resonance experiments: new perspectives for sequential NMR assignment of large proteins. *Proc Natl Acad Sci U S A* **95**: 13585–13590.
- Schnabl J, Wang J, Hohmann U, Gehre M, Batki J, Andreev VI, Purkhauser K, Fasching N, Duchek P, Novatchkova M, et al. 2021. Molecular principles of Piwi-mediated cotranscriptional silencing through the dimeric SFiNX complex. *Genes Dev* **35**: 392–409.
- Schuller SK, Schuller JM, Prabu JR, Baumgärtner M, Bonneau F, Basquin J, Conti E. 2020. Structural insights into the nucleic acid remodeling mechanisms of the yeast THO-Sub2 complex. *Elife* **9**. <http://dx.doi.org/10.7554/eLife.61467>.
- Shen Y, Bax A. 2012. Identification of helix capping and b-turn motifs from NMR chemical shifts. *J Biomol NMR* **52**: 211–232.
- Simon B, Köstler H. 2019. Improving the sensitivity of FT-NMR spectroscopy by apodization weighted sampling. *J Biomol NMR* **73**: 155–165.
- Sugiyama T, Thillainadesan G, Chalamcharla VR, Meng Z, Balachandran V, Dhakshnamoorthy J, Zhou M, Grewal SIS. 2016. Enhancer of Rudimentary Cooperates with Conserved RNA-Processing Factors to Promote Meiotic mRNA Decay and Facultative Heterochromatin Assembly. *Mol Cell* **61**: 747–759.
- Tang W, Tu S, Lee HC, Weng Z, Mello CC. 2016. The RNase PARN-1 Trims piRNA 3' Ends to Promote Transcriptome Surveillance in *C. elegans*. *Cell* **164**: 974–984.
- Teplova M, Farazi TA, Tuschl T, Patel DJ. 2016. Structural basis underlying CAC RNA recognition by the RRM domain of dimeric RNA-binding protein RBPMS. *Q Rev Biophys* **49**: e1.
- Terwilliger TC, Adams PD, Read RJ, McCoy AJ, Moriarty NW, Grosse-Kunstleve RW, Afonine PV, Zwart PH, Hung LW. 2009. Decision-making in structure solution using Bayesian estimates of map quality: the PHENIX AutoSol wizard. *Acta Crystallogr D Biol Crystallogr* **65**: 582–601.
- Terwilliger TC, Grosse-Kunstleve RW, Afonine PV, Moriarty NW, Zwart PH, Hung LW, Read RJ, Adams PD. 2008. Iterative model building, structure refinement and density

- modification with the PHENIX AutoBuild wizard. *Acta Crystallogr D Biol Crystallogr* **64**: 61–69.
- Upadhyay SK, Mackereth CD. 2020. Structural basis of UCUU RNA motif recognition by splicing factor RBM20. *Nucleic Acids Res* **48**: 4538–4550.
- Wan C, Tempel W, Liu Z-J, Wang B-C, Rose RB. 2005. Structure of the conserved transcriptional repressor enhancer of rudimentary homolog. *Biochemistry* **44**: 5017–5023.
- Wang G, Reinke V. 2008. A *C. elegans* Piwi, PRG-1, regulates 21U-RNAs during spermatogenesis. *Curr Biol* **18**: 861–867.
- Weick E-M, Miska EA. 2014. piRNAs: from biogenesis to function. *Development* **141**: 3458–3471.
- Weick E-M, Sarkies P, Silva N, Chen RA, Moss SMM, Cording AC, Ahringer J, Martinez-Perez E, Miska EA. 2014. PRDE-1 is a nuclear factor essential for the biogenesis of Ruby motif-dependent piRNAs in *C. elegans*. *Genes Dev* **28**: 783–796.
- Weng C, Kosalka J, Berkyurek AC, Stempor P, Feng X, Mao H, Zeng C, Li W-J, Yan Y-H, Dong M-Q, et al. 2019. The USTC co-opts an ancient machinery to drive piRNA transcription in *C. elegans*. *Genes Dev* **33**: 90–102.
- Weng MT, Luo J. 2013. The enigmatic ERH protein: Its role in cell cycle, RNA splicing and cancer. *Protein and Cell* **4**: 807–812.
- Williams CJ, Headd JJ, Moriarty NW, Prisant MG, Videau LL, Deis LN, Verma V, Keedy DA, Hintze BJ, Chen VB, et al. 2018. MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci* **27**: 293–315.
- Williamson MP. 2013. Using chemical shift perturbation to characterise ligand binding. *Prog Nucl Magn Reson Spectrosc* **73**: 1–16.
- Winter G, Lobley CMC, Prince SM. 2013. Decision making in xia2. *Acta Crystallogr D Biol Crystallogr* **69**: 1260–1273.
- Xie G, Vo TV, Thillainadesan G, Holla S, Zhang B, Jiang Y, Lv M, Xu Z, Wang C, Balachandran V, et al. 2019. A conserved dimer interface connects ERH and YTH family proteins to promote gene silencing. *Nat Commun* **10**. <http://dx.doi.org/10.1038/s41467-018-08273-9>.
- Yu S, Kim VN. 2020. A tale of non-canonical tails: gene regulation by post-transcriptional RNA tailing. *Nat Rev Mol Cell Biol* **21**: 542–556.
- Zeng C, Weng C, Wang X, Yan YH, Li WJ, Xu D, Hong M, Liao S, Dong MQ, Feng X, et al. 2019. Functional Proteomics Identifies a PICS Complex Required for piRNA Maturation and Chromosome Segregation. *Cell Rep* **27**: 3561-3572.e3.