1 **Title:**
2 **Cochlear nucleus small cells use olivocochlear collaterals to encode sounds in noise**
3
4 **Running title:**
5 **Small cells encode sounds in noise**
6

7 Adam Hockley[1], Calvin Wu[1], Susan E Shore[1,2,3]
8
9 [1] Kresge Hearing Research Institute, Department of Otolaryngology, [2] Department of Molecular
10 and Integrative Physiology, and [3] Department of Biomedical Engineering
11 University of Michigan, Ann Arbor, MI, 48109, U.S.A.
12
Correspondence:
Dr. Susan E. Shore,
Department of Otolaryngology, University of Michigan, 1150 West Medical Center Drive, Ann
Arbor, MI 48109
sushore@umich.edu

Author contributions:
A.H., C.W. and S.E.S. designed research; A.H. and C.W. performed research; A.H. and CW
analysed data; A.H., C.W. and S.E.S. wrote the paper.
22
23 **Conflict of Interest Statement:** The authors declare that the research was conducted in the
24 absence of any commercial or financial relationships that could be construed as a potential
25 conflict of interest.
26
27 **Acknowledgements:**
28 The authors would like to thank Mr. Dileepkumar for engineering support throughout this
29 project. This study was supported by National Institutes of Health grant R01-DC017119 (S.E.S.).
30
31 **Keywords:**
32 Cochlear nucleus, intensity coding, small cell cap

ABSTRACT
Understanding communication signals, especially in noisy environments, is crucial to social interactions. Yet, as we age, acoustic signals can be disrupted by cochlear damage and the subsequent auditory nerve fiber degeneration. The most vulnerable—medium and high-threshold—auditory nerve fibers innervate various cell types in the cochlear nucleus, among which, the small cells are unique in receiving this input exclusively. Furthermore, small cells project to medial olivocochlear (MOC) neurons, which in turn send branched collaterals back into the SCC. Here, we use single-unit recordings to characterise small cell firing characteristics and demonstrate superior intensity coding in this cell class. We show converse effects when activating/blocking the MOC system, demonstrating that small-cell unique coding properties are facilitated by direct cholinergic input from the MOC system. Small cells also maintain tone-level coding in the presence of background noise. Finally, small cells precisely code low-frequency modulation more accurately than other ventral CN (VCN) cell types, demonstrating accurate envelope coding that may be important for vocalisation processing. These results highlight the small cell–olivocochlear circuit as a key player in signal processing in noisy environments, which may be selectively degraded in aging or after noise insult.

INTRODUCTION
Auditory perception in ever-changing soundscapes is key for communication. However, this important task is degraded with age (Pichora-Fuller, 1997). Crucial to acoustic signal processing are the temporal fine structure and envelope information present in complex signals (Shannon et al., 1995), and difficulties in resolving these signals by the brain are prevalent in age-related hearing deficits (Bharadwaj et al., 2014). In the auditory periphery, the high threshold, low spontaneous rate (SR) auditory nerve fibers (ANFs) may be particularly important as their best dynamic range overlaps with normal levels of communicative signals. These specialised ANFs project along with low threshold ANFs throughout the cochlear nucleus (CN). However, the small-cell cap region receives exclusive input from the low (and medium) SR ANFs (Liberman, 1991; Ryugo, 2008).

Located between the granule-cell and magnocellular domains of the VCN (Cant, 1993), the small-cell cap (SCC), as the name suggests, is a region containing small cells (<20 μm). In humans, the SCC occupies a larger proportion of the CN than in rodents, and is therefore poised to play a major role in central mechanisms of speech perception (Moore & Osen, 1979). Small cells project either to medial olivocochlear (MOC) neurons in the ventral nucleus of the trapezoid body (VNTB; Darrow et al., 2012; De Venecia et al., 2005; Thompson & Thompson, 1991; Ye et al., 2000) or directly to the medial geniculate body, the latter serving as short-latency relay to the auditory cortex (Schofield, Mellott, et al., 2014; Schofield, Motts, et al., 2014). Small cells, in turn, receive branched collaterals from MOC neurons (Benson et al., 1996; Benson & Brown, 1990; Ryan et al., 1990), whose major function is suppression of cochlear outer hair cell electromotility to alter cochlear gain. MOC activation releases the cochlea from background masking by shifting its dynamic range (Dolan & Nuttall, 1988; Kawase & Liberman, 1993). However, small cells remain largely unexplored. Here, we hypothesise that MOC collateral projections to the VCN, and especially onto small cells, may counteract the effect of cochlear suppression, allowing small cell firing to accurately represent stimulus intensity.

78
In this study, we characterised small cell response properties that are consistent with its unique
circuit arrangement. We showed that small cells can more accurately encode intensity as well
as temporal information compared to other VCN cell types. Further, we found both MOC
activation and inhibition of cholinergic input to the VCN bidirectionally modulated small cell
firing properties. In the presence of background noise, small cells maintained superior intensity
coding. Taken together, these findings reveal a special role for small cells in processing
communicative signals and that its degradation after selective auditory neuropathy may
underlie hearing-in-noise difficulties.

METHODS

All animal experimental procedures were performed in accordance with the protocols established by the National Institutes of Health (Publication 80-23) and approved by the University Committee on Use and Care of Animals at the University of Michigan. Male and female (n = 11 and 15, respectively) pigmented guinea pigs weighing 280-800g obtained from Elm Hill Labs were used. No differences in unit responses for animal's sex or age was found. Guinea pigs were dual-housed on a 12/12 h light-dark cycle, with food and water readily available. Auditory brainstem responses (ABRs) were used to confirm normal auditory thresholds of ≤20 dB SPL at 4-20kHz in all animals. ABRs (0–90 dB SPL tone bursts; 5 ms duration, 1 ms rise/fall times, 21 Hz presentation rate, 512 repetitions in 10 dB steps; Tucker-Davis Technologies RZ6) were recorded at 4, 8, 12, 16 and 20 kHz.

Surgery

Guinea pigs were initially anaesthetised with ketamine (50 mg kg^{-1}; Hospira Inc., Lake Forrest, IL, USA) and xylazine (5 mg kg^{-1}; Akorn Inc.; Lake Forrest, IL, USA). Anaesthetic depth was maintained using 10 mg kg^{-1} ketamine and 1 mg kg^{-1} xylazine supplements. Atropine (0.05 mg kg^{-1}) was administered during the initial surgery to reduce bronchial secretions. Animals were placed in a stereotaxic frame (Kopf; Tuijunga, U.S.A.) within a sound-attenuating and electrically shielded double-walled chamber. The skull was exposed, and a midline incision made, temporalis muscle removed, and a craniotomy performed to allow access to the left CN. The dura mater was removed and the exposed brain surface was kept moist by regular applications of saline.

Sound presentation

Auditory stimuli were delivered monaurally via a closed-field, calibrated system (modified DT770 drivers; Beyerdynamic Heilbronn, Germany) coupled to hollow ear bars. The speakers were driven by a Tucker-Davis Technologies (TDT; Alachua, FL, USA) System 3 (RZ6, PA5 & HB7), controlled by TDT Synapse and custom MATLAB 2021a (Mathworks; Natick, U.S.A) software.

Neural recordings

Multi-channel recording probes (NeuroNexus; Ann Arbor, MI, USA) were advanced stereotaxically through the cerebellum towards the left CN using an MP-285 microdrive (Sutter Instruments; Novato, CA, USA). Signals were amplified by a TDT PZ5 preamp connected to a TDT RZ2 processor for filtering (0.3 − 5 kHz), and data was collected using Synapse software. Spikes were detected on-line with threshold set at 4 standard deviations from mean background noise, then spike-sorted post-hoc using a PCA approach. Electrodes were positioned so that shanks passed through the dorsal cochlear nucleus (DCN) to reach the SCC and VCN. In 2 animals, electrode tracts were marked to confirm SCC positioning. Units were characterised by responses to tone-burst and broadband noise (BBN) signals, also used to create receptive fields, peri-stimulus time histograms (PSTHs) and rate-level functions (RLFs) (Ghoshal & Kim, 1997; Palmer, 1987; Stabler et al., 1996; Winter & Palmer, 1995), and manually characterised divided into 5 categories; PL (bushy cells), Ch (T-stellates), SC (small cells), B (buildups) and On (onsets). A machine learning model was then used to confirm manual typing had produced discrete classifications of neurons (Fig 1). Poorly driven units were discarded.

131
Intensity coding analyses
Receptive field stimuli consisted of randomised tonebursts (50 ms duration, 200 ms ISI, 0.1
octave spacing, 5 dB steps, 15 repeats, randomised). From these, threshold and BF were
obtained from the lowest sound level to produce firing >2 standard deviations above
spontaneous firing rates. RLFs at BF were then used for further analyses. Total inhibitory area of
the receptive field was calculated as the number of squares of the receptive field where firing
rate was <2 standard deviations below spontaneous firing rates.
139
For further intensity coding analysis, a PSTH-based stimulus classification was performed using
methods described in Foffani & Moxon (2004). Briefly, spike responses (serialised PSTH; 1 ms
bins concatenated across units) of the neural population generated in each trial (50 trials total)
were compared to the "template," or average PSTH, of a given stimulus intensity. Euclidean
distances between each trial and the templates (19 total: 0 to 90 dB with 5 dB steps) were
computed and the template with the shortest distance was then assigned as "classified
intensity." Classification accuracy was defined as the percentage of trials that were classified
within <10 dB of the actual intensity, or "hits." Coefficient of variation for classification result
across all 20–50 trials was calculated as $std(x^2)/mean(x^2)$, where x counts the discretised step
away from the "hit."
150
Drug delivery
Local drug delivery of the muscarinic acetylcholine receptor antagonist, atropine was achieved
using a NeuroNexus D16 puffer-probe. The probe was connected to a 100 µl syringe containing
an 80 µM solution. The syringe was fixed to a digital syringe driver (UMP3 microsyringe injector
with Micro4 controller; World Precision Instruments; Sarasota, FL). This system allows
simultaneous drug delivery and neural recordings as described and validated in previous studies
(Rohatgi et al., 2009; Stefanescu & Shore, 2015, 2017). After assessing baseline CN neuronal
activity, 2 µl of solution was delivered at a rate of 100 nl/min, as in previous studies (Rohatgi et
al., 2009; Stefanescu & Shore, 2015, 2017).
160
Round window recordings
Cochlear compound action potential (CAP) recordings were performed using an insulated silver-
wire ball inserted through the bulla to rest on the cochlear round window. The trailing wire was
sealed to the skull, with a small hole left open for fluid wicking. CAPs were recorded in response
to 10 ms tones with rise/fall times of 2 ms. Amplitudes were calculated as P_1-N_1 of the mean
CAP waveform (200 repeats).
167
Electrical stimulation
MOC neurons in the VNTB were electrically stimulated with biphasic current pulses (100 ms
duration, 0.1 ms pulse width, 100 Hz, 500-1000 µA; 5/sec) applied through concentric bipolar
electrodes (CBCEF75; FHC; Bowdoin, ME). CAPs were recorded to verify MOC neuron location
(16 kHz, 70 dB SPL tone). CAP amplitudes were compared at baseline and 5 ms after the
cessation of electrical stimulation. A CAP-amplitude reduction of >30% confirmed accurate

MOC location in the VNTB for stimulation. Current levels were titrated to ensure maximal MOC
activation with no elicited muscular contractions.
176
Imaging
In a separate guinea pig, a tract tracing experiment was performed to confirm VNTB projections
to and from SCC. FluroEmerald (10%; 0.5 µl, Thermo Fisher Scientific, Waltham, MA, USA) was
pressure-injected (0.1 µl min-1) into the contralateral VNTB (1 mm anterior to the interaural
line, 1.5 mm lateral to midline, 12 mm from the dural surface). The animal was anesthetised
with ketamine-xylazine mixture in the same manner as described for electrophysiology
experiments but allowed to recover and maintained in the vivarium for 5 days before
transcardial perfusion (100 mL 1× PBS, 400 mL 4% paraformaldehyde) after euthanasia. The
brain was extracted, post-fixed for 2 h, immersed in 30% sucrose solution for 2 days, frozen and
cryosectioned at 50 µm in the coronal plane (Leica, CM3050S). Coverslipped brain sections
were examined under confocal epifluorescence (PMT; Leica, SP5-x).
188
189 **Experimental design and statistical analysis**
190 Small cell sound evoked responses were compared to other VCN cell types. Changes in these
191 responses during MOC modulation were also analysed. Statistical tests, including Kruskal-Wallis
192 test, ANCOVA, and two-way ANOVA were used to compare between cell-types or conditions (α
193 = 0.05). Post hoc analyses were performed using Tukey's multiple comparison test.
194
195 **RESULTS**
196 Using multichannel electrodes, we isolated 1523 single units from 83 VCN electrode locations in
197 26 guinea pigs. Unit distributions are shown in Table 1. We excluded onset and buildup units in
198 subsequent analyses due to their low numbers.
199

	PL	Ch	SC	On	B	Total
Control	722	247	310	57	187	1523
VNTB stimulation	260	96	139	15	75	585
Atropine infusion	20	36	34	4	8	102

200 *Table 1: Number of single CN units recorded. Cell types are Primary-like (PL), Chopper (Ch), Small*
201 *cell (SC), Onset (On) and Buildup (B).*
202
Single-shank 32-channel electrodes with 50 µm spacing of electrode sites were positioned to
simultaneously record units from the SCC and VCN (Fig 1A). Units were classified by their
temporal and frequency response patterns (Fig 1B)(Stabler et al., 1996; Winter & Palmer, 1995;
Young et al., 1988). Small cells, located dorsoventrally between DCN and VCN, displayed several
unusual PSTH shapes, the most common being a 'flat' or bimodal PSTH shape (Fig 1B). Small
cells were therefore identified by a combination of their PSTH shapes and unique locations
within the cochlear nucleus and by their distinct best frequency (BF) progressions with depth. A
machine learning model (RUSBoosted Trees) was used to confirm that manual typing had
produced discrete types of neurons (Fig 1E). This model used 23 inputs, each of which was a

metric obtained from either the PSTH, inter-spike interval histogram (ISIH), receptive field or
RLF.
214

215
Fig 1: Validating recordings from CN small cells. A) An example 3D plot of unit types within the
left CN of one guinea pig. Cells exhibiting buildup PSTH shapes were found more medial and
caudal, in the DCN. The VCN is more rostral and lateral, with the SCC units located between
these two structures. B) Example PSTHs from 4 different CN cell types: small cell (SC) with flat,
bimodal PSTH, primary-like (PL), buildup (B) and chopper (Ch). C) Receptive field of a CN small
cell. D) Coronal section of the cochlear nucleus, containing a Fluorogold-marked electrode track
in the SCC. (Bar = 0.5 mm) E) Confusion matrix of the machine learning model used to confirm
recordings from discrete types of neurons.
224
225   To confirm reciprocal projections from small cells to MOC neurons in the VNTB, a tracer
226   (FluoroEmerald) was pressure-injected into the VNTB. One week later, retrograde and
227   anterograde staining was observed in the SCC (Fig 2), demonstrating, along with the
228   physiological data, that SCC neurons form a circuit with MOC neurons, confirming previous
229   findings (Benson et al., 1996; Benson & Brown, 1990; Ye et al., 2000).
230

231
232

233
234
235 **Fig. 2: VNTB neurons project to and receive projections from the SCC.**
236 A) Schematic showing the locations of photomicrographs in B (VNTB) and C (SCC). B)
237 FluoroEmerald injection in the contralateral VNTB (cVNTB). C) Anterograde labeling of puncta
238 (arrowsheads) in SCC originating in VNTB and retrograde labelling of SCC somata. Bar = 100 μm.

239
Small cell firing patterns resemble those of low/medium SR ANFs
Afferent input to small cells is exclusively from low/med SR, high threshold ANFs, while small
cells project to, and receive input from MOC neurons. Due to this unique circuit configuration,
we hypothesised that small cells would have different patterns of intensity coding (rate-level
functions) compared to VCN cell types that also receive input from high SR, low threshold ANFs.
small cell response characteristics were like their low/medium SR, high threshold ANF inputs,
with low SRs, high thresholds, and long first-spike latencies compared to other VCN cell types
(Fig S1). Small-cell receptive fields contained inhibitory areas (Figs 1C S1D), signifying a
currently unknown source of inhibition.

249
To examine intensity coding at speech-relevant levels, we used a PSTH-based classifier to
compare the trial to trial reliability across cell types in response to different intensity stimuli
(Foffani & Moxon, 2004). Intensity coding fidelity is represented by accurate representations of
stimuli for each single-trial spiking response (Fig 3C). Primary-like units displayed better
intensity discrimination between 20-45 dB as well as very low variability, while small cells were
better at intensity coding above 60 dB (Fig 3D–E; ANCOVA F=9.1, P=1.1e-4, effect size (partial
eta squared) = 0.006). The high-fidelity intensity coding and absence of RLF plateaus in small
cells is likely due to their combined input from low SR ANFs and excitatory input from MOCs at
high sound levels. The dual action of peripheral and central excitation would allow small cells to
better encode tone intensities at levels relevant for speech processing.
260
261

Fig 3: CN small cells show superior intensity coding at high sound levels. A) Example PSTH and RLFs for a CN small cell and a primary-like unit. B) Mean (± SEM) RLFs of 4 major CN cell types. C) Confusion matrix for sound intensity classification across the population (all cell types). Greyscale — white pixels: 0%, black pixels: 100%. D) Probability of correct classification across unit-types. E) Trial to trial variability (coefficient of variation) of classifier performance across unit-types.

MOC neurons modulate small-cell excitability

Cholinergic MOC neurons send projections into the SCC (Benson et al., 1996; Benson & Brown, 1990; Ryan et al., 1990). Acetylcholine is known to act on CN cells through muscarinic receptors, resulting in a cell-type specific excitation or modulation of neural plasticity (Fujino & Oertel, 2001; Stefanescu & Shore, 2017). To investigate whether cholinergic MOC inputs mediate the greater accuracy of intensity coding in small cells, first, we applied the muscarinic acetylcholine receptor (mAChR) antagonist, atropine using drug delivery probes. Atropine reduced small cell medium to high intensity tone-evoked activity, (e.g. for 75 dB SPL, α = 0.00263, p < 0.0001), confirming a role of cholinergic input in the more accurate intensity encoding of small cells at higher tone intensities (Fig 4). Chopper cells, which also receive cholinergic input (Fujino & Oertel, 2001), likewise showed reduced sound-evoked activity following atropine infusion, (75 dB SPL, α = 0.00263, p = 0.00098) but over a narrow intensity range. Atropine had no significant effect on the RLF of primary-like cells (75 dB SPL, α = 0.00263, p = 0.7419). These findings demonstrate that the primary action of cholinergic input on small cells and choppers is to increase sound-driven responses above 60 dB, resulting in a widened dynamic range that is essential for intensity coding.

Next, to determine whether the MOCs were the key cholinergic modulators of small cells, we
electrically stimulated MOC neuron somata in the VNTB while performing CN recordings.
Successful targeting of MOC cells was confirmed by the reduction of CAP amplitudes during
electrical stimulation. MOC stimulation significantly increased small cell firing rates in response
to tones above 70 dB SPL but not to lower intensity or sub threshold tones (Fig 4, 75 dB SPL, α =
0.00263, p < 0.0001), suggesting that cholinergic MOC input to CN small cells enhances ANF-
evoked activity. As expected, the overall effect of MOC activation on CN small cells was
opposite to that achieved by cholinergic blocking. In contrast, VNTB stimulation had no effect
on chopper firing (75 dB SPL, α = 0.00263, p = 0.461), while decreasing primary-like firing across
the entire range of tone intensities (75 dB SPL, α = 0.00263, p < 0.0001). The reciprocal findings
of increased firing rates with MOC stimulation and decreased firing rates with atropine in small
cells confirm the role of MOC input to small cells in enhancing sound-driven responses.
Remarkably, small cell excitation by MOC neurons can overcome the MOC suppression of
cochlear output. The failure of MOC stimulation to alter firing rates in choppers suggests that
the cholinergic effect observed originates rather from VNTB cells that are not MOCs.
303

304
Fig 4: CN small cells receive excitatory cholinergic input from MOCs. A) Blocking muscarinic
cholinergic receptors with atropine supresses CN small-cell and chopper responses to tones.
Top row shows mean RLFs ± SEM for primary-like, chopper cells and small cells before and after

local application of atropine via a drug-delivery probe. B) Electrically stimulating MOC neurons
increases tone-evoked responses in small cells but not choppers. Mean RLFs ± SEM for primary-
like, chopper cells and small cells are shown before and during MOC electrical stimulation. All
statistical comparisons made using paired t-test with Bonferroni's multiple comparison
correction (α = 0.00263, stars signify p<0.05).
313
Small cells preserve higher-intensity tone coding in background noise
MOC projections to the cochlea suppress cochlear outer hair cell electromotility, thereby
reducing cochlear amplification. This aids in unmasking of sounds from background noise,
which facilitates the detection of tones and speech in noise (Dolan & Nuttall, 1988; Kawase &
Liberman, 1993). Since small cells receive projections from MOC collaterals, we hypothesised
that small cells are involved in encoding sounds in the presence of background noise. Signal-in-
noise discrimination was tested in VCN neurons first by comparing RLFs to tones in silence with
tones embedded in broadband noise.
322
As seen in individual and mean RLFs, small cells maintained their firing rates to medium to high
intensity tones even in the presence of background noise (Fig 5, 75 dB p = 0.9677). In fact,
small-cell spike rates in response to BF tones in the presence of 40- and 60 dB background noise
were equivalent to their responses to tones in silence (Fig 6). Chopper units maintained their
firing rates to BF tones in background noise at 40, but not 60 dB SPL (75 dB, p = 0.9677). In
contrast, background noise degraded the responses to BF tones in primary-like cells, which
showed increased rates and altered RLF patterns in the presence of 40- and 60 dB SPL
broadband noise. These differences were not due to thresholds as they were apparent for cells
of similar thresholds.
332
To further examine intensity coding, we applied the PSTH-based classifier analysis on these
same trials in both 40- and 60 dB broadband noise. Single-trial spike trains were classified based
on templates generated in silence (Fig. 3D) in the same population of units. In primary-like cells
during 40-dB background noise, we found that single-trial spike trains could not encode levels
above 55 dB (Fig. 6A). Chopper and small cells retained coding above 65 dB with increased
variability. The summary data showed that small cells are the most accurate coder with the
lowest variability at the highest intensity tested (Fig. 6B–C; ANCOVA: $F_{(1,2)}=0.34$, P=0.00085
across cell type). In 60-dB background noise, tone coding was nearly nonexistent for primary-
like, and sparse above 60 dB for choppers (Fig. 6D). Only in small cells were high intensity
coding above 75 dB preserved. At the highest intensity tested small cell retained 50% of
accuracy with lowest variability among cell types (Fig. 6E–F; ANCOVA: $F_{(1,2)}=26.9$, P=0.064).
Taken together, these results demonstrate that, vis-à-vis VCN cell types, small cells reliably
encode medium to high stimulus intensity in the presence of background noise.
346
347
348

Fig 5: CN small cells maintain tone-level coding in the presence of background broadband noise. A) Single unit examples of RLFs in the presence of 0, 40- or 60 dB SPL background noise, for the 3 major cell-type classifications (columns). Insets show PSTH shapes. B) Mean (± SEM) RLFs of 3 major CN cell types in the presence of background noise.

Fig 6: Small cells retain superior medium to high-level intensity coding in background noise. A) Representative PSTH-based classification of tone intensity for primary-like (PL), chopper (Ch),

361 and small cells (SC) populations during constant 40 dB BBN. Colour bar = normalised kernel
362 density across trials. B) Proportion of trials that show classified correctly (within ±5 dB of actual
363 level) as a function of intensity compared among the three cell types. C) The variability of
364 classification, or deviation from 'correct', measured by coefficient of variation (CV). D–F) Same
365 format as (A–C) but for tone coding in 60 dB BBN. Gray boxes denote background noise levels.
366
367

Small cells display precise envelope coding
Tracking sound envelopes is important for decoding human speech or animal vocalisations
(Shannon et al., 1995). Here, we used amplitude modulated broadband noise (75 dB SPL) to
evaluate envelope coding in CN neurons. Figure 7 (A-C) shows an example vector strength
calculation for a single small cell at a modulation frequency of 32 Hz and depth of 12%,
(Goldberg & Brown, 1969). This unit had a vector strength value of 0.43191, demonstrating
accurate spike timing to the amplitude modulation. Figure 7 D shows that mean vector strength
of the different VCN neurons peaked at a modulation frequency of 256 Hz, similar to that in
previous studies (Sayles et al., 2013). At low modulation frequencies and depths, small cells had
significantly greater vector-strength values than other VCN types (Fig 7) demonstrating superior
envelope coding. For example, at 6% modulation depth and a modulation frequency of 16 Hz,
Tukey's multiple comparison test found that the mean value of small cells was significantly
greater than choppers (p < 0.0001, 95% C.I. = 0.0437, 0.1345) and primary-like cells (p = 0.0014,
95% C.I. = 0.0172, 0.0877). The enhanced coding of low-frequency amplitude modulation in CN
cells may be important for species-specific vocalisation processing, as guinea pig vocalisations
contain predominantly low frequency amplitude modulation bursts (Berryman, 1976).
384

385
Fig 7: Small cells show more accurate amplitude-modulation coding than other VCN cell
types. Modulation transfer functions show vector strength of spiking to sinusoidally modulated
BBN at 5 different modulation depths (100-6%). CN small cells have significantly greater vector
strength of firing at modulation frequencies below 128 Hz and at lower modulation depths.
Stars indicate significance between CN small cells and other cell types (One-way ANOVA with
Tukey's multiple comparison test, p<0.05).

DISCUSSION

Small cells receive exclusive afferent input from high-threshold, low/med SR ANFs, positioning them to encode sounds at levels well above threshold. Conversational speech (above 60 dB SPL) is encoded by temporal cues within amplitude modulated signals. Here, we showed that small cells encode both intensity and amplitude modulation with higher precision than other VCN cell types. For tones in background noise, small cells were the only VCN cell type that maintained coding above 60 dB SPL. The unique properties of small cells are likely mediated by cholinergic MOC input, which provide excitation to small cells capable of overcoming the suppressive effect of MOCs on the cochlea, thereby increasing their suprathreshold firing rates.

Small cells exhibited low SRs, high thresholds and steep, unsaturating (up to 90 dB SPL) rate-intensity slopes. These characteristics resemble the firing properties of the low/medium SR ANFs that innervate small cells and are similar to those identified in cells in the marginal region of the cat CN (Ghoshal & Kim, 1996, 1997). Those early studies also showed flat, bimodal or Unusual PSTH shapes, signifying a heterogenous population, similar to that shown here for guinea-pig CN small cells. However, in the present study, the difference between small-cell thresholds and other VCN cell types (~5 dB) was less than the threshold difference between low/medium and high SR ANFs (~20 dB; Liberman, 1978; Winter and Palmer, 1991), putatively due to the excitatory MOC cholinergic input to small cells demonstrated here. Blocking muscarinic, cholinergic receptors with atropine reduced small-cell spike rates while MOC stimulation, confirmed by CAP reductions, increased their firing rates. Both effects are consistent with the presence of excitatory cholinergic MOC input to the SCC. The greatest effect of blocking muscarinic cholinergic receptors was observed during suprathreshold sound stimulation, which would be expected to also activate the sound-driven, cholinergic projections from MOC collaterals to the SCC (Lee et al., 2006).

MOC collaterals to the CN can also influence other cell types (Baashar et al., 2019; Mulders et al., 2003, 2007) including T-stellate cells (planar multipolar cells with chopper responses) and D-stellate cells (radiate multipolar cells with onset chopper responses). Previous studies showed an increased firing rate in 94% of T-stellate cells following application of carbachol (AChR agonist) in vitro (Fujino & Oertel, 2001). However, Baashar et al. (2019) used tract tracing and double labelling to view contacts from MOC collaterals onto T- and D-stellate cells and demonstrated that MOC input to these CN cells was weaker than would be expected based on the early electrophysiological studies. Some studies have suggested that there is inhibitory MOC projection to the CN (Mulders et al., 2002), however the present study found no evidence for inhibitory MOC or cholinergic effect on CN small cells or choppers. One possible reason for this discrepancy is the more targeted stimulation of contralateral MOC neurons in the present study, compared to midline stimulation, (Mulders et al., 2002), which would activate all crossed olivocochlear fibers including the lateral olivocochlear system.

The present data reveal strong, excitatory MOC-collateral innervation of small cells in the SCC but weak input to T-stellate cells. Small-cell receptive fields in the present study had inhibitory sidebands, revealing a currently unknown source of inhibition. This could be the glycinergic, D-

stellate cells, which provide wideband inhibition to several cell types in the CN (Arnott et al.,
2004; Oertel et al., 1990). However, further studies are required to confirm this pathway. An
alternative source of inhibition is the L-stellate cell, which also responds to carbachol and thus
potentially of olivocochlear origin (Ngodup et al., 2020). However, prediction of sources based
on bath infusion of cholinergic modulators must be made with caution, due to multiple sources
of cholinergic modulation to the CN (Mellott et al., 2011).

We demonstrated that small cells more accurately encode tones-in-noise compared to other
VCN cell types, putatively due to their restricted ANF inputs and unique input from MOC
collaterals. Activating the MOC pathway to the cochlea suppresses the cochlear amplifier,
benefitting speech-in-noise processing (de Boer et al., 2012; Winslow & Sachs, 1987). The
function of MOC collaterals to the SCC may be to further aid speech coding. Small cells have the
characteristics necessary for superior suprathreshold sound encoding and accurately encode
stimulus intensity (Figs 3&6). Pre-emptive MOC-pathway enhancement protects the cochlea
from synaptopathic acoustic trauma (Boero et al., 2018), but whether increased MOC activation
following synaptopathic noise exposure could reverse speech-in-noise deficits is still unknown.
Conversely, in humans with tinnitus or hyperacusis, an increased contralateral suppression of
otoacoustic emissions (Knudson et al., 2014) suggests an over-responsive MOC system in that
pathology. MOC neurons are modulated by descending input from the inferior colliculus
(Malmierca et al., 1996; Robertson & Mulders, 2000; Schofield & Cant, 1999), auditory cortex
(Coomes & Schofield, 2004; Mulders & Robertson, 2000), and local inhibitory interneurons of
the medial nucleus of the trapezoid body (Torres Cadenas et al., 2020). Descending modulation
may be a route for auditory attention to affect peripheral processing, which may explain
attentional facilitation of speech-in-noise understanding (de Boer et al., 2012). Descending
modulation of MOC activity would also affect MOC collateral input to CN small cells, which
accurately encode intensity, providing an additional action pathway for auditory attention.

Hearing damage in the absence of threshold shifts preferentially affects the higher threshold
ANFs (Furman et al., 2013) that provide the sole afferent input to small cells. Cochlear
synaptopathy is a potential major health issue in humans, as human temporal bones
demonstrate widespread synaptic damage (Viana et al., 2015). One downstream effect of
cochlear synaptopathy may be impaired speech perception, especially in the presence of
background noise. This widespread complaint has been attributed to cochlear synaptopathy,
but this relationship is still unclear (Plack et al., 2014). Human studies suggest that such
perceptual deficits reflect difficulties in coding temporal envelopes (Bharadwaj et al., 2014),
expected to impair speech recognition (Shannon et al., 1995). Temporal-coding deficits after
synaptopathy are not apparent in ANFs (Heeringa et al., 2020), suggesting their origin may be
central. The enhanced intensity coding and precise envelope coding shown here for small cells
primes the SCC as an important subcortical area for speech processing is expected to be
preferentially targeted by cochlear synaptopathy. In humans, the SCC occupies a large
proportion of the CN and is therefore poised to play a major role in central mechanisms of
auditory neuropathy (Moore & Osen, 1979). Speech coding in human small cells may be
impaired following cochlear synaptopathy, and modulation of CN activity via the MOC pathway
may be a potential treatment for the associated speech-in-noise deficits.

REFERENCES

Arnott, R. H., Wallace, M. N., Shackleton, T. M., & Palmer, A. R. (2004). Onset Neurones in the
Anteroventral Cochlear Nucleus Project to the Dorsal Cochlear Nucleus. *Journal of the*
Association for Research in Otolaryngology, 5(2), 153–170.
Baashar, A., Robertson, D., Yates, N. J., & Mulders, W. H. A. M. (2019). Targets of Olivocochlear
Collaterals in Cochlear Nucleus of Rat and Guinea Pig. *Journal of Comparative Neurology,*
527(March), 2273–2290.
Benson, T. E., Berglund, A. M., & Brown, M. C. (1996). Synaptic input to cochlear nucleus
dendrites that receive medial olivocochlear synapses. *Journal of Comparative Neurology,*
365(1), 27–41.
Benson, T. E., & Brown, M. C. (1990). Synapses formed by olivocochlear axon branches in the
mouse cochlear nucleus. *The Journal of Comparative Neurology, 295*(1), 52–70.
Berryman, J. C. (1976). Guinea-pig Vocalizations: Their Structure, Causation and Function.
Zeitschrift Für Tierpsychologie, 41(1), 80–106. https://doi.org/10.1111/j.1439-
0310.1976.tb00471.x
Bharadwaj, H. M., Verhulst, S., Shaheen, L., Liberman, M. C., Shinn-Cunningham, B. G., Peelle, J.
E., Léger, A. C., & Elisabeth Gockel, H. (2014). Cochlear neuropathy and the coding of
supra-threshold sound. *Frontiers in Systems Neuroscience, 8*(26).
Boero, L. E., Castagna, V. C., Di Guilmi, M. N., Goutman, J. D., Elgoyhen, A. B., & Gómez-Casati,
M. E. (2018). Enhancement of the Medial Olivocochlear System Prevents Hidden Hearing
Loss. *Journal of Neuroscience, 38*(34), 7440–7451.
Cant, N. B. (1993). The Synaptic Organization of the Ventral Cochlear Nucleus of the Cat: The
Peripheral Cap of Small Cells. In *The Mammalian Cochlear Nuclei* (pp. 91–105). Springer
US.
Coomes, D. L., & Schofield, B. R. (2004). Projections from the auditory cortex to the superior
olivary complex in guinea pigs. *European Journal of Neuroscience, 19*(8), 2188–2200.
https://doi.org/10.1111/j.0953-816X.2004.03317.x
Darrow, K. N., Benson, T. E., & Brown, M. C. (2012). Planar multipolar cells in the cochlear
nucleus project to medial olivocochlear neurons in mouse. *The Journal of Comparative*
Neurology, 520(7), 1365–1375.
de Boer, J., Thornton, A. R. D., & Krumbholz, K. (2012). What is the role of the medial
olivocochlear system in speech-in-noise processing? *Journal of Neurophysiology, 107*(5),
1301–1312.
De Venecia, R. K., Liberman, M. C., Guinan, J. J., & Brown, M. C. (2005). Medial olivocochlear
reflex interneurons are located in the posteroventral cochlear nucleus: A kainic acid lesion
study in guinea pigs. *Journal of Comparative Neurology, 487*(4), 345–360.
Dolan, D. F., & Nuttall, A. L. (1988). Masked cochlear whole-nerve response intensity functions
altered by electrical stimulation of the crossed olivocochlear bundle. *J. Acoust. Soc. Am,*
83(3), 1081–1086.
Foffani, G., & Moxon, K. A. (2004). PSTH-based classification of sensory stimuli using ensembles
of single neurons. *Journal of Neuroscience Methods, 135*(1–2), 107–120.
https://doi.org/10.1016/j.jneumeth.2003.12.011
Fujino, K., & Oertel, D. (2001). Cholinergic modulation of stellate cells in the mammalian ventral

cochlear nucleus. *Journal of Neuroscience*, *21*(18), 7372–7383.

Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2013). Noise-induced cochlear neuropathy is
selective for fibers with low spontaneous rates. *Journal of Neurophysiology*, *110*(3), 577–
586.

Ghoshal, S., & Kim, D. O. (1996). Marginal shell of the anteroventral cochlear nucleus: intensity
coding in single units of the unanesthetized, decerebrate cat. *Neuroscience Letters*, *205*,
71–74.

Ghoshal, S., & Kim, D. O. (1997). Marginal Shell of the Anteroventral Cochlear Nucleus: Single-
Unit Response Properties in the Unanesthetized Decerebrate Cat. *J Neurophysiol*, *77*(4),
2083–2097.

Goldberg, J. M., & Brown, P. B. (1969). Response of Binaural Neurons of Dog Superior Olivary
Complex to Dichotic Tonal Stimuli: Some Physiological Mechanisms of Sound Localization.
American Physiological Society, *32*(4), 613–636.

Heeringa, A. N., Zhang, L., Ashida, G., Beutelmann, R., Steenken, F., & Köppl, C. (2020).
Temporal Coding of Single Auditory Nerve Fibers Is Not Degraded in Aging Gerbils. *The*
Journal of Neuroscience⊡: The Official Journal of the Society for Neuroscience, *40*(2), 343–
354.

Kawase, T., & Liberman, M. C. (1993). Antimasking Effects of the Olivocochlear Reflex. I.
Enhancement of Compound Action Potentials to Masked Tones. *Journal of Physiology*,
70(6).

Knudson, I. M., Shera, C. A., & Melcher, J. R. (2014). Increased contralateral suppression of
otoacoustic emissions indicates a hyperresponsive medial olivocochlear system in humans
with tinnitus and hyperacusis. *Journal of Neurophysiology*, *112*(12), 3197–3208.

Kujawa, S. G., Glattke, T. J., Fallon, M., & Bobbin, R. P. (1993). Contralateral sound suppresses
distortion product otoacoustic emissions through cholinergic mechanisms. *Hearing*
Research, *68*(1), 97–106. https://doi.org/10.1016/0378-5955(93)90068-C

Lee, D. J., De Venecia, R. K., Guinan, J. J., & Brown, M. C. (2006). Central auditory pathways
mediating the rat middle ear muscle reflexes. *Anatomical Record - Part A Discoveries in*
Molecular, Cellular, and Evolutionary Biology, *288*(4), 358–369.
https://doi.org/10.1002/ar.a.20296

Liberman, M. C. (1978). Auditory-nerve response from cats raised in a low-noise chamber. *The*
Journal of the Acoustical Society of America, *63*, 442.

Liberman, M. C. (1991). Central projections of auditory-nerve fibers of differing spontaneous
rate. I. Anteroventral cochlear nucleus. *Journal of Comparative Neurology*, *313*(2), 240–
258.

Malmierca, M. S., Le Beau, F. E. N., & Rees, A. (1996). The topographical organization of
descending projections from the central nucleus of the inferior colliculus in guinea pig.
Hearing Research, *93*(1–2), 167–180.

Mellott, J. G., Motts, S. D., & Schofield, B. R. (2011). Multiple origins of cholinergic innervation
of the cochlear nucleus. *Neuroscience*, *180*, 138–147.

Moore, J. K., & Osen, K. K. (1979). The cochlear nuclei in man. *American Journal of Anatomy*,
154(3), 393–417.

Mulders, W. H. A. M., Harvey, A. R., & Robertson, D. (2007). Electrically Evoked Responses in
Onset Chopper Neurons in Guinea Pig Cochlear Nucleus. *Journal of Neurophysiology*, *97*(5),

3288–3297.

Mulders, W. H. A. M., Paolini, A. G., Needham, K., & Robertson, D. (2003). Olivocochlear

collaterals evoke excitatory effects in onset neurones of the rat cochlear nucleus. *Hearing*

Research, 176(1–2), 113–121.

Mulders, W. H. A. M., & Robertson, D. (2000). Evidence for direct cortical innervation of medial

olivocochlear neurones in rats. *Hearing Research, 144*(1–2), 65–72.

https://doi.org/10.1016/S0378-5955(00)00046-0

Mulders, W. H. A. M., Winter, I. M., & Robertson, D. (2002). Dual action of olivocochlear

collaterals in the guinea pig cochlear nucleus. *Hearing Research, 174*(1–2), 264–280.

https://doi.org/10.1016/S0378-5955(02)00701-3

Ngodup, T., Romero, G. E., & Trussell, L. O. (2020). Identification of an inhibitory neuron

subtype, the l-stellate cell of the cochlear nucleus. *ELife, 9*, 1–40.

https://doi.org/10.7554/eLife.54350

Oertel, D., Wu, S. H., Garb, M. W., & Dizack, C. (1990). Morphology and physiology of cells in

slice preparations of the posteroventral cochlear nucleus of mice. *Journal of Comparative*

Neurology, 295(1), 136–154.

Palmer, A. R. (1987). Physiology of the cochlear nerve and cochlear nucleus. *Br Med Bull, 43*(4),

838–855.

Pichora-Fuller, M. K. (1997). Language comprehension in older listeners. *Journal of Speech-*

Language Pathology and Audiology, 21(2), 125–142. https://psycnet.apa.org/record/1997-

05664-003

Plack, C. J., Barker, D., & Prendergast, G. (2014). Perceptual Consequences of "Hidden" Hearing

Loss. *Trends in Hearing, 18*, 1–11.

Robertson, D., & Mulders, W. H. A. M. (2000). Distribution and possible functional roles of some

neuroactive peptides in the mammalian superior olivary complex. *Microscopy Research*

and Technique, 51(4), 307–317.

Rohatgi, P., Langhals, N. B., Kipke, D. R., & Patil, P. G. (2009). In vivo performance of a

microelectrode neural probe with integrated drug delivery. *Neurosurg Focus, 27*(1), 8.

Ryan, A. F., Keithley, E. M., Wang, Z. -X, & Schwartz, I. R. (1990). Collaterals from lateral and

medial olivocochlear efferent neurons innervate different regions of the cochlear nucleus

and adjacent brainstem. *Journal of Comparative Neurology, 300*(4), 572–582.

Ryugo, D. K. (2008). Projections of low spontaneous rate, high threshold auditory nerve fibers

to the small cell cap of the cochlear nucleus in cats. *Neuroscience, 154*(1), 114–126.

Sayles, M., Füllgrabe, C., & Winter, I. M. (2013). Neurometric amplitude-modulation detection

threshold in the guinea-pig ventral cochlear nucleus. *The Journal of Physiology*

Neuroscience C, 591(13), 3401–3419.

Schofield, B. R., & Cant, N. B. (1999). Descending auditory pathways: Projections from the

inferior colliculus contact superior olivary cells that project bilaterally to the cochlear

nuclei. *Journal of Comparative Neurology, 409*(2), 210–223.

Schofield, B. R., Mellott, J. G., & Motts, S. D. (2014). Subcollicular projections to the auditory

thalamus and collateral projections to the inferior colliculus. *Frontiers in Neuroanatomy,*

8(70), 1–16.

Schofield, B. R., Motts, S. D., Mellott, J. G., & Foster, N. L. (2014). Projections from the dorsal

and ventral cochlear nuclei to the medial geniculate body. *Frontiers in Neuroanatomy,*

8(10), 1–12.

Shannon, R. V, Zeng, F., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech Recognition with
Primarily Temporal Cues. *Science*, *270*(5234), 303–304.

Stabler, S. E., Palmer, A. R., & Winter, I. M. (1996). Temporal and mean rate discharge patterns
of single units in the dorsal cochlear nucleus of the anesthetized guinea pig. *Journal of*
Neurophysiology, *76*(3), 1667–1688.

Stefanescu, R. A., & Shore, S. E. (2015). NMDA receptors mediate stimulus-timing-dependent
plasticity and neural synchrony in the dorsal cochlear nucleus. *Frontiers in Neural Circuits*,
9(November), 1–14.

Stefanescu, R. A., & Shore, S. E. (2017). Muscarinic acetylcholine receptors control baseline
activity and hebbian stimulus timing-dependent plasticity in fusiform cells of the dorsal
cochlear nucleus. *Journal of Neurophysiology*, *117*(3), 1229–1238.

Thompson, A. M., & Thompson, G. C. (1991). Posteroventral cochlear nucleus projections to
olivocochlear neurons. *The Journal of Comparative Neurology*, *303*(2), 267–285.

Torres Cadenas, L., Matthew, X., Fischl, J., & Weisz, C. J. C. (2020). Synaptic Inhibition of Medial
Olivocochlear Efferent Neurons by Neurons of the Medial Nucleus of the Trapezoid Body.
The Journal of Neuroscience, *40*(3), 509–525.

Viana, L. M., O, J. T., Burgess, B. J., Jones, D. D., ACP Oliveira, C., Santos, F., Merchant, S. N.,
Liberman, L. D., & Charles Liberman, M. (2015). Cochlear neuropathy in human
presbycusis: confocal analysis of hidden hearing loss in post-mortem tissue HHS Public
Access. *Hear Res*, *327*, 78–88. https://doi.org/10.1016/j.heares.2015.04.014

Winslow, R. L., & Sachs, M. B. (1987). Effect of Electrical Stimulation of the Crossed
Olivocochlear Bundle on Auditory Nerve Response to Tones in Noise. *Journal of*
Neurophysiology, *57*(4), 1002–1021.

Winter, I. M., & Palmer, A. R. (1991). Intensity coding in low-frequency auditory-nerve fibers of
the guinea pig. *The Journal of the Acoustical Society of America*, *90*(4), 1958–1967.

Winter, I. M., & Palmer, A. R. (1995). Level dependence of cochlear nucleus onset unit
responses and facilitation by second tones or broadband noise. *Journal of*
Neurophysiology, *73*(1), 141–159.

Ye, Y., Machado, D. G., & Kim, D. O. (2000). Projection of the marginal shell of the anteroventral
cochlear nucleus to olivocochlear neurons in the cat. *The Journal of Comparative*
Neurology, *420*(1), 127–138.

Young, E. D., Robert, J. M., & Shofner, W. P. (1988). Regularity and latency of units in ventral
cochlear nucleus: implications for unit classification and generation of response properties.
Journal of Neurophysiology, *60*(1), 1–29.

649
650 Fig S1: Small cells show lower SFRs and longer FSLs than other VCN units. A) Bar plot (mean ±
651 SEM) of SFR for CN cell types (primary-like (PL), chopper (Ch), small cell (SC) and buildup. B) Bar
652 plot (mean ± SEM) for FSL. C) Bar plot (mean ± SEM) of thresholds for each CN cell type. D) Bar
653 plot (mean ± SEM) of total inhibitory (TI) area of the receptive field. Comparisons between
654 groups were made using a Kruskal-Wallis test with Tukey's multiple comparison test *$p<0.05$.
655

Model input	Description
SFR	Spontaneous firing rate (Hz)
T	Threshold at BF (dB SPL)
Q10	Tuning width at BF and 10 dB above threshold.
Q40	Tuning width at BF and 40 dB above threshold.
DR	Dynamic range
ISIHcv	Coefficient of variation of the ISIH (0.5ms bin width, 0-25ms).
FSL	First spike latency
Plat	RLF plateau
PSTHfr	Mean driven firing rate throughout PSTH (0-50ms)
PSTHon	Onset firing rate of PSTH (0-15ms)
PSTHsus	Sustained firing rate of PSTH (35-50ms)
PSTHonsusrat	Ratio of sustained rate to onset rate (PSTHon/PSTHsus)
PSTHsl	PSTHsus / PSTH rate 15-35ms
PSTHsl2	Slope of linear fit to PSTH between 15-50ms
PSTHonpks	Number of PSTH peaks between 0-20ms, with a min prominence of 25 Hz, and min firing rate of 100Hz.
PSTHnotch	Using a window from the first onset peak as 0, to 3ms. Minimum firing rate within this window / maximum firing rate.
PSTHcv	Coefficient of variation (CV) of the first 20ms of the PSTH.
PSTHcvsl1	The slope of a linear fit on a plot of how PSTH CV changes from 5-25ms.
PSTHcvsl2	The slope of a linear fit on a plot of how PSTH CV changes from 31-50ms.
RLFin	Increase in firing rate from 60 dB to 90 dB at BF.
RLFsl	RLF slope between 80 and 90 dB SPL.
RLFsl6090	RLF slope between 60 and 90 dB SPL.
TI	Total inhibitory squares on the receptive field.

656 Table S1: Inputs for machine learning CN cell typing model.
657