Decoding the effects of synonymous variants

Zishuo Zeng[1*], Ariel A. Aptekmann[1], and Yana Bromberg[1,2*]

1 Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, NJ 08873, USA
2 Department of Genetics, Rutgers University, Piscataway, New Jersey 08854, USA
* Corresponding author: zzeng@bromberglab.org, yana@bromberglab.org
Tel: +1-732-932-9763 Ext. 218; Fax +1-732-932-8965

ABSTRACT

Synonymous single nucleotide variants (sSNVs) are common in the human genome but are often overlooked. However, sSNVs can have significant biological impact and may lead to disease. Existing computational methods for evaluating the effect of sSNVs suffer from the lack of gold-standard training/evaluation data and exhibit over-reliance on sequence conservation signals. We developed synVep (synonymous Variant effect predictor), a machine learning-based method that overcomes both of these limitations. Our training data was a combination of variants reported by gnomAD (observed) and those unreported, but possible in the human genome (generated). We used positive-unlabeled learning to purify the generated variant set of any likely unobservable variants. We then trained two sequential extreme gradient boosting models to identify subsets of the remaining variants putatively enriched and depleted in effect. Our method attained 90% precision/recall on a previously unseen set of variants. Furthermore, although synVep does not explicitly use conservation, its scores correlated with evolutionary distances between orthologs in cross-species variation analysis. synVep was also able to differentiate pathogenic vs. benign variants, as well as splice-site disrupting variants (SDV) vs. non-SDVs. Thus, synVep provides an important improvement in annotation of sSNVs, allowing users to focus on variants that most likely harbor effects.

INTRODUCTION

The recent increase in accessibility of sequencing has facilitated a rise in precision medicine efforts focused on the interpretation of the effects of individual-specific genome variation (1). Genome-wide association studies (GWAS) have identified multiple variants marking specific phenotypes (2). However, the evaluation of variants in terms of their functional contributions to molecular pathogenicity mechanisms holds promise for both a better understanding of disease and drug discovery/optimization (3). SNVs (single nucleotide variants) are the most common variants in the human genome (4). Three types of SNVs are of particular interest – regulatory (*i.e.* changing the quantity/production of the gene product, *e.g.* transcription or splice site variants), non-synonymous (*i.e.* altering product protein sequence), and synonymous (*i.e.* variants in protein-coding regions that, due to the degeneracy of the genetic code, do not alter the protein sequence). Many computational tools have been developed to evaluate the functional effects of regulatory and non-synonymous variants (5,6). However, while an individual genome carries as

many synonymous as non-synonymous SNVs (7), the former are often disregarded as functionally irrelevant. Still, sSNVs can cause disease (8) and affect gene function via multiple mechanisms, including binding of transcription factors (9), splicing (10), mRNA stability (11-13), co-translational folding (14-16), *etc.*, as reviewed in our earlier work (17).

Existing methods for predicting sSNV effects are either (1) sSNV-specific tools, including SilVA (18), reg-SNP-splicing (19), DDIG-SN (20), TraP (21), and IDSV (22), or (2) general-purpose ones, including CADD (23,24), DANN (25), FATHMM-MKL (26), and MutationTaster2 (27). The number of computational sSNV effect predictors is limited in comparison to that of nsSNV (non-synonymous single nucleotide variant) effect predictors, as reviewed in (6,17). Partially, this paucity is due to the limited available experimental data evaluating variant effects, which could be used for training or testing of such methods. In fact, all existing predictors, except CADD and DANN, are trained using "pathogenic" variants from databases such as Human Gene Mutation Database (HGMD) (28) and ClinVar (29). Here we note that "pathogenicity" is not equivalent to "functional effect" (30,31) and inferring variant-disease causality is complicated by this inequality. The experimental disease variant annotations are also often unreliable (17), as it is difficult to distinguish causative variants from simply associated ones. Moreover, the pathogenic label is inconsistent across databases, and possibly over time/database releases. Finally, even these labeled effect variants are few; even fewer are experimentally labeled neutral polymorphisms. Thus, predictors trained on these variants are likely insufficient to predict the effects of tens of millions of possible sSNVs in human genome.

Using positive-unlabeled learning (32-34), we inferred a subset of human sSNVs that could be used for training a predictor of sSNV molecular effect. We then developed, synVep (synonymous Variant Effect Predictor), a machine learning-based method for scoring putative effect for each possible human sSNV. synVep discriminated experimentally validated pathogenic sSNVs from randomly sampled common variants. Its predictions also displayed the expected trends (35) in evolutionary distances between orthologs, where the sSNVs corresponding to evolutionarily-close human relatives' (*e.g.* chimp) reference nucleotide, have lower effect scores than those corresponding to the nucleotides of further away organisms (e.g. fruitfly). Furthermore, nucleotides that are not identified in any of the species evaluated here are deemed to have most effect when substituted into the human reference. However, many of the sSNVs that are not observed in the human population, tend to be scored very high (most effect), regardless of their appearance in other species.

In line with our earlier observations (17), we find that the variant frequency in the population is poorly correlated with the effect score; *i.e* rare variants are about equally likely to have no effect on gene function as common variants (65% common vs 69% rare). synVep does not rely on conservation and is developed without an experimental or explicitly evolutionarily estimated gold standard training/development set. Its success thus suggests the feasibility of a similar approach for the development of a training set for other variant types, *e.g.* nsSNVs or indels. We expect that synVep sSNV

effect predictions will greatly contribute to our understanding of pathogenicity pathways and to the
prioritization of synonymous variants in disease.

MATERIALS AND METHODS

Data Collection. We extracted all 93,437 human protein-coding transcripts from the Ensembl BioMart
(36) GRCh37 p.13 assembly (37) and discarded the ones containing unknown nucleotides, lacking a
start/stop codon, or having patched (https://grch37.ensembl.org/Homo_sapiens/Info/Annotation)
chromosome IDs. We then generated all possible sSNVs for the remaining 72,400 transcripts. We further
used ANNOVAR (38) (installed Aug 5, 2019) to extract sSNVs in these transcripts, and their allele-count
based frequencies, from the Genome Aggregation Database exome subset (gnomAD exome) (39). An
sSNV present in gnomAD was labeled a *singleton* if it was seen in only one individual and otherwise
labeled *observed*. *Generated* sSNVs were those in the set of all possible variants in the 72,400
transcripts that were not *singleton* or *observed*. Thus, we collected 4,160,063 *observed*, 3,438,470
singleton, and 57,208,450 *generated* sSNVs (https://zenodo.org/record/4763256). Note that these
correspond to 1,520,334 *observed*, 1,233,878 *singleton*, and 21,314,668 *generated* sSNVs with unique
genomic coordinates and reference/alternative alleles, *i.e.* in one transcript per gene.

To evaluate and compare the performance of our predictor to other predictors, we manually curated a
dataset of 42 *curated-effect* sSNVs with known biological effects, including the 33 pathogenic variants
from the Buske et al. study (18). We required that all sSNVs in this set were strongly associated with
disease and that there was experimental evidence of their molecular effects. These 42 sSNVs
(Supplementary Table S1) mapped to 170 transcript-based sSNVs and were excluded from model
training throughout this manuscript.

Variant Features. We collected 35 variant and sequence features (Supplementary Table S2), grouped
into six categories: codon bias and autocorrelation (10 features), protein structure (3), mRNA stability (8),
distance to regulatory factors (4), expression profile (3), and miscellaneous (7). The reasons for selecting
these features are described in our earlier paper (17). We further calculated the correlation of feature
values across all sSNVs using the dython package (v0.6.7, https://github.com/shakedzy/dython), where
correlations between continuous-continuous, continuous-categorical, and categorical-categorical features
were computed using Pearson correlation (40), Cramer's V (41), and correlation ratio (42), respectively.
Feature importance was obtained by calculating the average performance gain across all splits where the
feature was present.

1. Transcript expression profiles. We downloaded the GTEx (43) "Transcript TPMs" dataset (dbGaP
Accession phs000424.v7.p2) and standardized the transcript expression across tissue samples. We then
used the average expression of each transcript over all samples from the same tissue as the
representative transcript expression for that tissue.

Calculations of some of the codon bias metrics described below require a reference set of coding
sequences, which are typically a set of highest expressed transcripts (44). To identify these references,

we collected the maximum expression values for all transcripts across the 53 tissues. We then selected

the transcripts within the highest 1% expression per tissue. We also used $\log_{10}$ (minimum expression per

tissue), $\log_{10}$ (median expression per tissue), and $\log_{10}$ (maximum expression per tissue) for each

transcript as features.

<u>2. Codon bias and autocorrelation</u>. A variety of measures and/or their "Δ" form (difference in measure

value after mutation vs. value before mutation) are adopted as features to characterize the codon bias of

transcripts (see Supplementary text for more details), including the Codon Adaptation Index (CAI,

Supplementary text Eqn. 1) (44), Fraction of Optimal Codons (fracOpt, Supplementary text Eqn. 2) (45),

Codon Usage Bias (CUB, Supplementary text Eqn. 3) (46), Intrinsic Codon Deviation Index (ICDI,

Supplementary text Eqn. 4) (47), Synonymous Codon Usage Order (SCUO, Supplementary text Eqn. 5)

(48), and tRNA Adaptation Index (tAI, Supplementary Eqn. 6) (49). The calculation of these values was

performed in R (50) and is available as an R package in

<u>https://bitbucket.org/bromberglab/codonbiasmetrics/src/master/</u>.

These measures describe codon bias from different perspectives. CAI, fracOpt, and CUB rely on a

reference set of optimal codons, found in highly expressed genes (51). CAI computes the geometric

mean of relative usage of a codon compared to the most frequently used codon for the same amino acid

(44). fracOpt is the fraction of optimal codons in a sequence of a certain length. CUB weighs the

frequency of amino acids in calculating codon bias. ICDI is independent of a reference set of genes (47).

SCUO borrows the idea of entropy from Shannon information theory to describe codon usage bias of

sequences (48). tAI focuses on translational efficiency by taking tRNA levels into account (49).

We also considered codon autocorrelation – a feature that has not yet been used by any sSNV

predictors. In autocorrelated sequences same codons cluster together, whereas they are separated in

anticorrelated sequences (*e.g.* XXXYYY is more autocorrelated than XYXYXY, where X and Y are two

different codons) (52). Cannarozzi et al. noted the association between codon autocorrelation and

translation dynamics and proposed the tRNA pairing index (TPI) to describe a sequence's codon

autocorrelation. Autocorrelated sequences benefit from rapid translation due to the recycling of

isoaccepting tRNAs (52). However, we note that the significance of recycling is likely weaker if the interval

between two issoaccepting codons is larger – a feature that is not accounted for in TPI. Therefore, we

proposed a new measure, Codon Autocorrelation Measure (CAM, Supplementary Eqn. 7), to describe the

variant-specific codon autocorrelation impact penalized by the distance between the synonymous codons.

Finally, we also introduced the change of frequency measure (CF, Supplementary text Eqn. 8), to

describe the amount of impact on codon's frequency in a sequence due to the introduction of the variant.

<u>3. Distance to regulatory and splicing sites</u>. We used as features the distances to the nearest splice sites,

transcription factor binding site (TFBS), RNA-binding protein (RBP) motif, and exonic splicing regulator

(ESR). Their genomic coordinates were obtained from different sources as described below. We then

computed the distance of a variant (in nucleotides) to all regulatory sites and selected the minimum value

as the feature distance. We categorized these distances (d) into six categories as feature inputs: d=0,
0<d<=3, 3<d<=5, 5<d<=10, 10<d<=20, and d>20.

Genomic coordinates of regulatory regions were inferred as follows: (1) *Splice sites* were inferred from the
"Genomic coding start" and "Genomic coding end" of all human protein-coding transcripts annotated in
Ensembl BioMart GRCh37 p.13 assembly. (2) We downloaded the Gene Transcription Regulation
Database (GTRD, version 18.06) (53) and identified the genomic coordinates of *TFBS,* using hg38 to
hg19 conversion via CrossMap (54) for correspondence with our transcript coordinates. (3) We
downloaded the ATtRACT database of RNA binding proteins and AssoCiated moTifs (55) and mapped
the human *RPB* motifs to our set of transcript sequences. (4) We also downloaded the supplementary
data of Cáceres et al (56) gold standard *ESR* motif set and mapped these to our transcripts.

4. Protein structure. We ran PredictProtein (57), a collection of tools for protein structure predictions, on
all of the translated transcript sequences. We were particularly interested in protein secondary structure
(PSS), residue solvent accessibility (SS), and disorder (PD) predictions; in PredictProtein, PROFphd (58)
predicts PSS and SS, while Meta-disorder (MD) (59) predicts PD.

5. mRNA stability, structure, and structural changes. We ran RNAfold (60) to predict (with calculation of
partition function and base pairing probability matrix) the secondary structure and stability of all
transcripts. We extracted the frequency of the structure with minimum free energy (MFE) in the structure
ensemble, the free energy of the centroid structure, and its distance to the structure ensemble, as well as
the local mRNA structure (strongly paired, strongly up/down -stream paired, weakly paired, weakly
up/down -stream paired, or unpaired bases).

We also used RNAsnp (61) to predict the variant-induced local secondary structure changes for all
sSNVs. The "mode" and "winsizeFold" parameters should be assigned according to the length in
nucleotides (L) of the input sequence. We assigned the parameters as follows: (1) for L <= 200, mod=1
and winsizeFold=100; (2) for 200 < L <= 500, mod=1 and winsizeFold=200; (3) for L > 500, mod=2 and
winsizeFold=500. We recorded the local structure dissimilarity, global structural dissimilarity and their
statistical significance (p-values).

Model construction *Classifier setup.* We standardized all continuous features and label-encoded
categorical features. We compared two classifiers for differentiating *observed* and *generated* variants:
deep neural network (DNN) (62) and XGBoost (63); we selected XGBoost as the classification algorithm
for its higher accuracy and speed (preliminary experiment described in Supplementary Text, Page 2).
XGBoost is implemented in Python (v3.6.4) *xgboost* package (v0.8.2) integrated with sci-kit learn (0.20.3)
(64) (https://xgboost.readthedocs.io/en/latest/python/python_api.html).

Balancing variant data by transcript. The *generated* set of sSNVs is much larger than the *observed* set,
but the number of *observed* sSNVs per transcript varies greatly. Moreover, some classifier input features
are transcript-specific. Thus, a predictor may "memorize" transcripts that have more *observed* sSNVs,
and preferentially assign its variants *observed* status, instead of finding variant-specific differences

between *observed* and *generated*. To avoid this, we assigned sampling likelihood weights for the
generated set, *i.e.* the sampling likelihood weight of a *generated* variant is the number of *observed* sSNVs
in the corresponding transcript. In all further balancing of data sets, *generated* sSNVs were
probabilistically added to the set on the basis of their weights. Thus, the number of *generated* sSNVs on a
transcript that were selected for a particular training set was correlated with the number *observed* sSNVs
on this transcript.

Positive unlabeled learning (PUL) to identify unobservable sSNVs. PUL is a semi- supervised approach
applicable to scenarios where only positive data points are labeled and the rest can be positive or
negative (32-34). We employed the modified version of PUL (34) to separate the *generated* sSNVs into
unobservable and *not-seen* sets. To prevent overfitting, we adopted relatively conservative
hyperparameters of XGBoost (100 trees [n_estimor], 5 maximum depth [max_depth], 30% of the features
per tree [colsample_bytree], 30% subsamples per tree [subsample]). We left out from PUL a fraction of
observed as a test set, aiming to reach <5% incorrect predictions for this set at the end of the PUL.

In one epoch of PUL, a classifier was trained to differentiate the *observed* sSNVs from the same number
of unlabeled ones (*generated;* selected via transcript-based set balancing as described above). All
unlabeled sSNVs, including the ones not used in training, were evaluated with the resulting model and the
generated variants classified as *observed* (scoring below 0.5) were added to the *not-seen* pool. The PUL
process was repeated until convergence (Supplementary text). sSNVs scoring >0.5 in prediction from the
last PUL model were further excluded from our data set. One pitfall of this PUL strategy is that a fraction
of the unlabeled samples may become positive (*observed*) with more sequencing in the future.

Differentiating the observable from not-seen using an intermediate model. We trained a model to
differentiate the *observable* sSNVs from the *not-seen* sSNVs (termed intermediate model from here on).
We excluded 10% (9,274) of the common sSNVs (MAF > 0.01; *excluded* set) and all *curated-effect*
sSNVs (170) from the construction of the intermediate model for testing and final model parameter
optimization. We split the *observed* sSNVs into subsets of 9: 0.5: 0.5 size ratio for training (3,631,441
variants), validation and testing (201,746 variants each). We then randomly sampled the *not-seen*
variants to match the *observed* validation and test set sizes; this left 47,923,258 *not-seen* sSNVs for
training. We then up-sampled the 3.6M *observed* variants in the training set to create a balanced set of
47,923,258 *observed* and *not-seen* variants, each). We tuned the model hyperparameters by optimizing
the F-score (Eqn. 3) of performance on the validation set and evaluated the resulting model on the test
set.

$$Precision = \frac{TP}{TP+FP}$$ **(Eqn. 1)**

$$Recall = \frac{TP}{TP+FN}$$ **(Eqn. 2)**

$$F\_core = \frac{2 \times Precision \times Recall}{(Precision+Recall)}$$ **(Eqn. 3)**

where TP, TN, FP, FN are respectively, true positive, *i.e. observed* sSNVs predicted to be *observed*; true
negative, *not-seen* sSNVs predicted to be *not-seen*; false positive, *not-seen* sSNVs predicted to be
observed; false negative, *observed* sSNVs predicted to be *not-seen*.

Final model (synVep) training. We used the intermediate model to score the excluded common and
curated-effect sSNVs, as well as all *observed* and *not-seen* sSNVs. Here we assumed that common
variants should be enriched in no-effect/neutral variation. Based on the scores of excluded sSNVs, we
defined *effect* and *no-effect* synVep development sets, where sSNVs (both *observed* and *not-seen*)
scoring above the median of the *curated-effect* predictions were deemed *effect*; while sSNVs (both
observed and *not-seen*) scoring below the median of the excluded common sSNV predictions were
labeled *no-effect*. We thus collected 7,385,137 *no-effect* and 32,117,625 *effect* sSNVs.

We split the *no-effect* and *effect* sSNVs into subsets of 9: 0.5: 0.5 size ratio (in the same way as for the
intermediate model) for training, validation, and test sets (62,758,222: 735,194: 735,194 variants per set).
We sampled equal numbers of *effect* sSNVs to match the *no-effect* sSNVs in validation and test sets. We
trained the final model on the training set using the hyperparameters optimized (F-score; Eqn. 3) on the
validation set. We finally evaluated the model on the test set. Note that none of the *curated-effect*, the
excluded common sSNVs, or the ClinVar (described below) dataset variants were included in our model
training.

Performance comparison with other predictors. For all comparisons with other predictors, we
calculated the area under the receiver operating characteristic curve (auROC) using the pROC package
(65) (v1.17.0.1), and the area under the precision recall curve (auRPC) using PRROC (66) (v1.3.1).
Statistical significance of the differences between the auROC/auPRC of synVep and those of other
predictors were tested using the pROC package (bootstrap method with the default settings, n=2,000;
source code was modified to accommodate testing for auPRC).

Common/curated-effect dataset comparison. To evaluate synVep in comparison with other predictors, we
used the 170 (transcript-based*;* 42 genomic coordinate-based) *curated-effect* sSNVs and the 9,274
(transcript-based; 7,957 genomic coordinate-based) *excluded* common sSNVs. Here we again assumed
that common variants should be enriched in no-effect/neutral variation.

Other predictors in this comparison included: CADD (phred-like scaled scores) (23), DANN (25),
FATHMM-MKL (26), DDIG-SN (20), and EIGEN (67). EIGEN scores were collected using ANNOVAR (38)
annotations; for other predictors, the scores were collected with default parameters as described in our
earlier work (17). We did not include SilVA (18) or TraP (21) in this comparison because 33 of 42 of the
curated-effect sSNVs are in their training sets.

Note that synVep scores are produced per variant per transcript, while other predictors use the genomic
coordinates, *i.e.* one reference sequence per variant. For the purposes of our comparison, we randomly
re-sampled each tool's predictions of the *effect* set (42 variant scores) to produce 170 scores.
Furthermore, as the common sSNVs (putatively no-effect) outnumbered the *effect* set, we randomly

sampled 170 common variant scores in 100 comparison iterations. For each re-sampling, we performed a

one-sided permutation test (null hypothesis: mean of common variant scores is equal to mean of *effect*

scores; alternative hypothesis: mean of common variant sores is lower than mean of *effect* scores) and

recorded the p-value and the difference of the common and *effect* variant score distribution medians

(∆medians). To assure that all predictor scores fall into the same [0,1] range, we standardized CADD and

EIGEN ∆medians to their maximum non-outlier score (20 for CADD and 3 for EIGEN).

We also computed the Spearman correlation across predictor scores and the Fraction of Consensus

Binary Predictions (FCBP; *i.e.* the number of binarized predictions agreed upon by all predictors, divided

by total number of predictions) (17). An *effect/no-effect* scoring threshold for the FCBP computation is

required; we used the default value of score=0.5 for DANN, FATHMM-MKL, and DDIG-SN. For CADD,

we used score=15 as the threshold recommended by its online documentation

(https://cadd.gs.washington.edu/info). As there was no recommended cutoff in the EIGEN publication

(67), we selected the cutoff (score=1.35) at the 75-percentile of EIGEN scores of 1,000 randomly

sampled *observed* sSNVs.

ClinVar dataset comparison. We downloaded all ClinVar (68) submissions from the FTP site

(https://ftp.ncbi.nlm.nih.gov/pub/clinvar/) and identified the sSNVs among these. We only considered the

sSNVs with the "reviewed by expert panel" *review status.* From these we selected the (1) pathogenic and

pathogenic/likely pathogenic variants as the *pathogenic* set and (2) benign and benign/likely benign as

the *benign* set. There were 51 *benign* (genomic coordinate-based; 254 transcript-based) and 17

pathogenic (genomic coordinate-based; n=68 transcript-based) sSNVs (Supplementary Table S3). We

also annotated these ClinVar sSNVs with the precomputed GERP++ scores

(http://mendel.stanford.edu/SidowLab/downloads/gerp/) (69).

Splicing dataset comparisons. We downloaded and analyzed a dataset of SNV splicing effects (70)

(https://github.com/KosuriLab/MFASS), referenced by genomic coordinates and Ensembl transcript IDs.

For comparison with synVep, we downloaded and ran spliceAI (71) (https://github.com/Illumina/SpliceAI)

and retrieved CADD-splice (72) annotation from https://cadd.gs.washington.edu/score. spliceAI

predictions are composed of probabilities of splice acceptor and donor's gain and loss. Since these

outputs are predominantly zero, we took the maximal value for evaluation purpose, as in (72).

Cross-species sequence variation (CSV) analysis. Cross-species variation (CSV) describes the

nucleotide difference between the human reference sequence and the ortholog reference sequence of

another species. In this study, we selected 20 species to generate CSVs: yeast (*Saccharomyces*

cerevisiae), worm (*Caenorhabdiis elegans*), fruitfly (*Drosophila melanogaster*), zebrafish (*Danio rerio*),

xenopus (*Xenopus laevis*), anole lizard (*Anolis carolinensis*), chicken (*Gallus gallus*), platypus

(*Ornithorhynchus anatinus*), opossum (*Monodelphis domestica*), dog (*Canis familiaris*), pig (*Sus scrofa*),

dolphin (*Tursiops truncatus*), mouse (*Mus musculus*), rabbit (*Oryctolagus cuniculus*), tree shrew (*Tupaia*

belangeri), tarsier (*Carlito syrichta*), gibbon (*Nomascus leucogenys*), gorilla (*Gorilla gorilla*), bonobo (*Pan*
paniscus), and chimpanzee (*Pan troglodytes*).

To represent the evolutionary distance of the CSV species to human, we obtained the value in million
years since divergence from the TimeTree database (73). Given a human transcript *T* and its
corresponding human gene *G*, we queried Ensembl BioMart for *G*'s orthologs in the 20 species, $\mathbf{G_{orthologs}}$
= [G_{yeast}, G_{worm}, $G_{fruitfly}$, …, $G_{chimpanzee}$]. We downloaded all coding DNA sequences (CDS) for these
orthologs from Ensembl (release-94) (74). For each gene in $G_{orthologs}$, we identified its longest transcript
per organism, $\boldsymbol{T_{orthologs}}$ = [T_{yeast}, T_{worm}, $T_{fruitfly}$, …, $T_{chimpanzee}$]. We then used PRANK (75) to generate a
multiple sequence alignment (MSA) for each *T*. PRANK aligns CDSs by first translating them into protein
sequences so that gaps tend to be placed between codons, instead of within codons. For each codon in
each human transcript, we could identify if other organisms carried the same codon or another, even if
the amino acid remained the same. If the codon was different, the corresponding human sSNV was
termed a CSV.

Evaluation of synVep predictions according to constraint on coding regions. Constrained regions
(76), referenced by genomic coordinates, were downloaded from https://s3.us-east-
2.amazonaws.com/ccrs/ccr.html. The constraint of human coding region is measured by percentile (of
residuals from a linear regression for distance-to-mutation prediction as computed in (76)), where a high
percentile indicates a more constrained region. We annotated the sparsity of sSNVs, *i.e.* the fraction of
observed sSNVs among all possible sSNVs in a region of a certain constraint level, and the median
synVep prediction of variants in these regions.

Analysis of sSNVs identified in Qatari Genome . We downloaded all VCF files containing variants
identified from the Qatari Genome project (QTRG) (77) from NCBI Sequence Read Archive (78)
(https://trace.ncbi.nlm.nih.gov/Traces/sra/?study=SRP061943). We then parsed these VCF files,
extracted the variants, and mapped the sSNVs to our *observed*, *singleton*, *not-seen*, and *unobservable*
sets.

312
Figure 1. Pipeline of predictor construction. Starting with 4,160,063 *observed* and 57,208,450 generated sSNVs,
63 epochs of positive unlabeled learning (PUL) was conducted to separate the *generated* set into *not-seen*
(observable) and *unobservable* set (Supplementary text). An intermediate model was trained using the *observed* and
not-seen sets (*observed* set was up-sampled to equal amount of *not-seen* variants). The intermediate model's
predictions for common and pathogenic sSNVs were used as guideline to set cutoffs assigning *no-effect* and *effect*
set. The final predictor was trained using the *no-effect* and *effect* sets (*no-effect* set was up-sampled to equal amount
of *effect* sets). After the final synVep model was trained, it was evaluated on independent datasets as shown. Here,
singletons are sSNVs found in only one individual in gnomAD; *observed* are any other sSNVs found in gnomAD;
generated are all possible sSNVs, except *singletons* or *observed*; *unobservable* are sSNVs PUL-labeled to be unlike
the *observed*; *not-seen* are any other *generated* sSNVs; *effect/no-effect* are sSNVs that affect/do not affect the
function or quantity of a gene product.
324
325

Figure 2. Extraction of cross-species sequence variants (CSV). For each human protein coding transcript *T*, codon-oriented multiple sequence alignment was performed with 20 species' longest coding sequencing of the same ortholog as *T*. The CSV are represented as "codon>codon" format for specific transcript positions and may coincide with human sSNVs.

RESULTS AND DISCUSSION

Generated sSNVs may be observable in the future. In the absence of a gold-standard experimentally validated data set describing sSNV functional effects, we sought an alternative for the development of our method. We had previously proposed to use sSNVs that have been observed in major sequencing projects *vs.* all other possible human genome sSNVs (the *generated* set) for method evaluation (17). We collected 72,400 human transcripts with 4,160,063 (n=1,520,334 genomic coordinate-based) *observed* sSNVs and 3,438,470 (n=1,233,878 genomic coordinate-based) *singletons* (observed in only one individual) from the exome sequencing data of the Genome Aggregation Database (gnomAD exome) (39). We then created a *generated* set of 57,208,450 (n=21,314,668 genomic coordinate-based) all possible sSNVs in these transcripts that were not found in gnomAD data. Note that only ~12% of all sSNVs in our set were ever reported by gnomAD. We annotated these sSNVs with 35 transcript- and variant-specific features, including codon bias, codon autocorrelation, transcript stability, expression level, distance to regulatory sites, predicted protein secondary structures, *etc.* (Methods; Supplementary Table S2).

While the *observed* sSNVs are not necessarily functionally neutral, they are at least compatible with life. The *generated* sSNVs, on the other hand, likely comprise two subtypes: the *not-seen* sSNVs, which may or may not become *observed* with more sequencing, and the *unobservable* ones, which cannot be

observed given the contemporary variant-discovery capability. Note that the *unobservable* character of
sSNVs may be due to a broad range of technical and biological reasons such as sequencing (79,80),
molecular functional constraints (81), and analytical biases or extreme deleteriousness resulting in early
embryonic incompatibility with life (82,83). We also note that in our modelling, the *unobservable* set may
simply be poorly described by our selection of variant features.

We used *observed* sSNVs as positives in positive-unlabeled learning (PUL) (32-34) to differentiate the
not-seen sSNVs (similar to *observed*) from *unobservable* ones in the *generated* (unlabeled) set (Figure
1). At convergence (epoch 63, Methods; Supplementary Figure 1), PUL partitioned all *generated* sSNVs
into *unobservable* (n=6,278,254 transcript-based and 2,764,229 genomic coordinate-based; 11%) and
not-seen (n=50,930,196 transcript-based and 19,730,623 genomic coordinate-based; 89%). Additionally,
8% (n=266,192) of *singletons* were deemed *unobservable* by the PUL model, as were 2% of the
observed sSNVs (n=79,639). The latter result highlights the possible insufficiency of our variant
descriptors for capturing the complete observable variant diversity, while the former may also indicate
sequencing errors. The difference in percentages of variants misidentified by the model (11% of
generated vs 2% of *observed*), however, suggests that deleteriousness of variants also plays a role in
defining *unobservable* variants.

Observed and not-seen variant sets contain both no-effect and effect sSNVs. We trained the *intermediate*
model (Figure 1) to recognize *observed* vs. *not-seen* sSNVs. The model accurately (F-score=0.71; Eqn.
3) recognized the two classes in a previously unseen variant test set (Supplementary Figure 2). It also
predicted 9% (9,282,542) of the *not-seen* sSNVs to be *observed* (scoring < 0.5), implying that these may
be sequenced in the future.

Although large effect sSNVs may be enriched in the *not-seen* group, the intermediate model cannot be
directly used to evaluate effect, because it is only meant to predict whether an sSNV has been observed
or not. To build a model for effect evaluation, we leveraged the intermediate model's predictions on
common variants *excluded* from training and the experimentally validated *effect* sSNVs (*curated-effect*;
Methods; 170 transcript-based sSNVs). While these *curated-effect* sSNVs are, in fact, observed, their
prediction scores were higher than those of the *excluded* common set (Supplementary Figure 2, Mann-
Whitney U test p-value<2.2e-16). This observation is likely due to the fact that the *not-seen* set is
enriched, while the common variant set is depleted, in large effect sSNVs. We assume this for common
variants because large-effect deleterious variants would not become common and large-effect
advantageous variants would tend to become wild-type.

We excluded 10% (7,957) of the common sSNVs from training of the intermediate model for selecting the
cutoff of *effect/no-effect* variants as next described. For training of the final model, we selected as *no-*
effect those sSNVs (both *observed* and *not-seen*) scoring below the intermediate model prediction
median (0.38) of the excluded common sSNVs; variants scoring above the intermediate model median of
the *curated-effect* sSNVs (score = 0.63) were labeled *effect* (Supplementary Figure 2). We thus obtained

7,385,137 (2,580,540 *observed* and 4,804,597 not-seen) *no-effect* and 32,117,625 (405,170 *observed*
and 31,712,455 *not-seen*) *effect* sSNVs. We trained the final model (synVep, Figure 1) to differentiate the
no-effect and *effect* sSNVs (in balanced class training), using a 9: 0.5: 0.5 split of data for training,
validation, and testing purposes (Methods). synVep was accurate (F-score=0.90; binary score cutoff=0.5)
in evaluating the hold-out test set (369,257 *no-effect* and 369,257 *effect*). Note that synVep prediction
scores did not correlate with allele frequency (Pearson correlation=0.02).

Feature importance in discriminating effect. We collected 35 features (Supplementary Table S2; Methods)
highlighting the different ways how sSNVs can impact gene function (17). We examined the correlation of
feature scores across all sSNVs (Supplementary Figure 3; Methods) and computed feature importance
for the final model (Supplementary Figure 4; Methods). Feature scores correlated within the same
feature category for some categories (*e.g.* codon context, codon bias, and expression profile), but not
across different categories. The most important feature for our model was codon_mutation (*i.e.* the wild
type/mutant codon pair), which is consistent with our earlier observation that some codons are
preferentially mutated in observed sSNVs (17) and with the Karczewski et al (39) observation that CpG-
transitions in the population are closer to saturation than other mutation types. Another codon context
feature – next_codon, which is highly correlated with the last_codon and codon_mutation features – was
the third most important feature. This reflects the biological importance of codon pairs in modulating
translational efficiency (84-86). Codon bias measures (and their changes due to mutation) were also of
high importance (starting at second highest rank), in line with the abundant evidence of the relationship
between codon composition and a variety of biologically-relevant factors, including gene expression
(49,87,88), translational efficiency (14,89,90), and mRNA stability (91-94). Since codon selection
modulates translational speed and thus cotranslational folding (95), sSNVs can also affect protein
structure (96) without altering protein sequence. We incorporated protein annotations (predicted
secondary structure, solvent accessibility, and disorder) as features; curiously, solvent accessibility
ranked 8[th] in importance for the synVep model. Surprisingly, most other features had low importance;
including features related to mRNA structure and stability, which are known to be directly influenced by
sSNVs (8). This is perhaps due to the fact it is difficult to accurately predict RNA structure/stability for
sequences longer than 500 nucleotides (97), and over 74% of transcripts in our data are longer than that.

Note that we did not use conservation as a synVep feature as it is usually the overarching signal of effect
for most predictors (6) and we were hoping to capture additional, more subtle, signals orthogonal to those
already reported. However, we also evaluated synVep's potential performance loss due to this choice by
re-training the final model with an additional conservation feature (we used GERP++ scores for evaluation
purposes (69)). This model was not significantly better in discriminating *no-effect*/*effect* variants (0.9 vs
0.9 F-score with and without conservation, respectively, in evaluating the test set); we also note that the
difference in distribution of conservation scores across the *effect* and *no-effect* data sets was minimal
(Supplementary Figure 5). Driven by this somewhat unexpected lack of conservation difference between
the *effect/no-effect* sets, we further aimed to validate our set selection at the intermediate model level. We

trained an intermediate model using conservation as one of the features. This model identified a new,
conservation-included set of *effect* (n=6,263,638) and *no-effect* (n=32,010,049) sSNVs. This new partition
overlapped significantly with the original one (75% of the original sSNVs were present and had identical
effect/no-effect labels in both data sets). Moreover, synVep (without the conservation feature) predicted
both data sets equally well. Using a balanced test set from the original data, it achieved 89%/90% *no-*
effect precision/recall (Eqn. 1 and Eqn. 2) and 90%/89% *effect* precision/recall; it achieved a similar
performance for a balanced subset of the conservation-included *effect/no-effect* set (87%/88% *no-effect*
precision/recall and 87%/88% *effect* precision/recall). Given these results, we chose to further continue to
exclude conservation from synVep features. This choice makes synVep scores orthogonal to those of
other predictors and allows for the further described cross-species variant analysis to be performed.

433
Figure 3. Predictor performance on common *vs. curated-effect* sSNVs. Panels **A-F** show the differential
predictions on sets of *curated-effect* (n=170) and common sSNVs (randomly selected n=170) for CADD (phred-like
scaled scores), DANN, DDIG-SN, FATHMM-MKL, EIGEN, and synVep, respectively. Gray line indicates scoring
cutoff suggested by tool authors. Neither the common set nor the *curated-effect* set were included in synVep training.
In panel **(G)**, this comparison was repeated 100 times to compute the medians (i.e., median$_{curated-effect}$ −
median$_{common}$). Permutation tests show that all predictors give significantly different scores between the two sets in

every iteration, except for DANN where 11 of 100 comparisons were not significant (p-value>0.05 after Bonferroni
correction). Panel H reports the performance (auROC and *effect* auPRC) of each predictor on the left-out common
sSNVs (negative; n=9,274) and *curated-effect* sSNVs (positive; n=170). All other predictors' (except CADD_phred
and DANN) auROC and auPRC are significantly different from synVep's (p-value<0.05; Methods). Note that the
utility of auROC and auPRC is limited in single variant effect prediction; instead, a cutoff is needed.

Predictors identify sSNV effect. To evaluate the performance of synVep, we needed a gold standard set

of designated effect and no effect variants. However, since there is no experimentally validated "neutral"

sSNVs, we used the common sSNVs excluded in training as neutrals (*no-effect*), *i.e.* as described above

we assumed that the majority of common sSNVs have little effect. On the level of molecular function, the

effect of a variant is not directional (neither advantageous nor deleterious). On the level of individual

fitness (evolutionary), however, variants could be advantageous, neutral, or deleterious. Note that

evolutionarily *effect* variants must have a molecular effect, but not vice versa. The neutral theory of

molecular evolution, as well as some experimental work, suggest that advantageous mutations are rare,

and most effect mutations are deleterious (98,99). Variants with large deleterious effects on evolutionary

fitness would have been purified out and will not be seen in the population. Neutral or weak *effect*

mutations could, depending on the population size, become common or even fixed due to genetic drift

(98). Thus, in the absence of experimentally neutral variants, common variation may be considered a

reference point for neutral or weak *effect* variants.

We used the set of *curated-effect* sSNVs (n=170; excluded from synVep training) as the *effect* group and

the subset of common sSNVs left-out from synVep training (n=9,274) as the *no-effect* group to compare

the predictor performances (Figure 3). Of the considered methods, synVep had the highest auPRC but

the lowest auROC (Figure 3H), suggesting that other predictors' decision cutoffs may not be optimized for

sSNV effect classification. Note that the utility of auROC and auPRC is, however, questionable in

application to individual variants; here, a default cutoff is required to prioritize a variant. We note that most

predictors failed to differentiate the two groups of variants at the default binary cutoff, placing both sets of

variants below (CADD, DDIG-SN) or above the default threshold (DANN, FATHMM-MKL). Although

possibly helped by the intermediate model choices, the majority of synVep's predictions for the two

groups fall on two sides of the default cutoff.

We further designed an additional testing scheme to evaluate predictor performance. Since the test set of

no effect common sSNVs (n=9,274) greatly outnumbered the *curated-effect* set (n=170), we randomly

sampled 170 sSNVs from the common set 100 times; each time, we computed the difference between

the *effect*-median and *no-effect*-median scores (Methods). synVep outperformed other predictors (Figure

3F; CADD and EIGEN scores normalized to 0-1 scale), although all predictors identified most *curated-*

effect sSNVs as having more *effect* than common sSNVs. Note that the "general-purpose" predictor

(CADD, DANN, and EIGEN) cutoffs are not optimized to specifically evaluate synonymous variants and

may thus be expecting a larger signal of effect.

We also examined the correlation of the predicted scores (Figure 4A) and the Fraction of Consensus
Binary Prediction (17) (FCBP; Figure 4B) on all 4,160,063 *observed* sSNVs for all predictors (synVep,
CADD, DANN, FATHMM-MKL, DDIG-SN, and EIGEN). synVep's scores were poorly correlated with other
predictor scores (Pearson correlation ranging from -0.1 [DANN] to 0.23 [FATHMM-MKL]), while binary
classification was more similar (FCBP ranging from 0.37 [DANN] to 0.69 [CADD and DDIG-SN]).

482
Figure 4. Comparing predictions made by synVep and other tools. Panel **(A)** Spearman correlation and **(B)**
Fraction of Consensus Binary Prediction (FCBP) highlight similarity and lack thereof among the predictors for all
observed sSNVs (n=4,160,063).

Singletons are more likely than observed to have an effect. We re-predicted scores of all variants in our
data (excluding *unobservable*) with the final synVep model. As expected, of *observed* sSNVs, only 31.3%
were effect (median score 0.15), while 72.0% of *not-seen* sSNVs were effect (median score 0.88);
singletons were scored/distributed bimodally (Supplementary Figure 6) into the two classes (48.2% effect;
median score 0.47). Note that synVep predictions did not appear to be driven by site mutability (putative
mutation rate). synVep scores of *not-seen* sSNVs that share a genomic position with none, 1, or
2 observed variants do not significantly differ from each other (median synVep scores 0.86, 0.94, and
0.88 respectively; Supplementary Figure 7).

Singletons were not included in our training because it is difficult to estimate how many of them are
artifacts due to the 0.1-0.6% error rates of next-generation sequencing (100). If the *singletons* are not
artifacts, then they are likely to be individual or ultrarare variants. These are more likely to be *effect* than
higher frequency variants (101,102). An excess burden of ultrarare variants (although not necessarily

synonymous) is also often seen in diseases, such as schizophrenia (103-105), Parkinson disease (106),

and bipolar disorder (107). In line with these expectations, we found that *singletons* were, on average,

scored higher than *observed* sSNVs (Supplementary Figure 6), suggesting that *singletons* are more likely

to have an *effect* than the *observed*.

Variant effect predictors differentiate benign and pathogenic variants. Among the predictors considered in

this work, only two (FATHMM-MKL and DDIG-SN) are explicitly aimed to assess variant pathogenicity. To

investigate whether predictors for variant functional *effect* (*i.e.* not pathogenicity) can identify pathogenic

sSNVs, we obtained from ClinVar 17 *pathogenic* (genomic coordinate-based, 68 transcript-based) sSNVs

and 51 *benign* sSNVs (genomic coordinate-based, 254 transcript-based) variants reviewed by an expert

panel. Of these 68 variants, one benign and one pathogenic (genomic coordinate-based, 13 transcript-

based) were deemed *unobservable* by our model and were removed from consideration. Note that for

fairness of evaluation, these ClinVar sSNVs were also excluded from training of synVep. All variant-effect

predictors, including synVep, assigned higher scores to *Pathogenic* than *Benign* variants (Figure 5A-C

and E). However, at the default/recommended cutoff, only synVep placed the majority of *Benign* vs.

Pathogenic variants on opposite sides of the cutoff (*Pathogenic* recall=0.58; *Benign* recall=0.66).

synVep also attained the highest *effect* auPRC (Figure 5F), suggesting that it can identify disease-

causing sSNVs well even though it was not explicitly trained to do so. However, synVep attained the

lowest auROC, which may be due to the fact that benign ClinVar variants are actually functionally

significant (*effect*) and are thus predicted by synVep as such but classified as wrong by ClinVar

annotation (*FP*). In our definition, a variant of some *effect* is not necessarily pathogenic, but pathogenic

variants are expected to have *effect*. Thus, experimentally validated pathogenic variants predicted to be

no-effect by synVep are likely errors, but benign variants predicted be *effect* are quite possibly correctly

identified as having functional impact, which does not necessarily correspond to disease.

Note that because synVep's predictions are transcript-based, they can differ for the same variant

across multiple transcripts. Aggregating these predictions to score a variant is not trivial: one can

use the mean, maximum, or median scores. A more sophisticated approach would be to weigh the

scores from different transcripts by their expression level in multiple tissues. Specifically, if the

question is about a disease and if the disease is primarily associated with one tissue, only the

transcript most expressed in that tissue cab ne considered. However, given the complicated

regulation and genetic interactions, this idea needs further validation. An evaluation of predictions for

the same variant, however, highlights an interesting observation: only 5.1% of *not-seen*, 5.5% of

observed, and 7.4% of *singletons* had at least one transcript, whose effect prediction differed from

others.

Figure 5. Evaluating variant effect predictors using ClinVar data. *Benign* (negative; n=254) are variants labeled "Benign" and "Benign/Likely benign" in ClinVar, with "Review by expert panel" as review status. *Pathogenic* (positive; n=68) are those labeled "Pathogenic" and "Pathogenic/Likely Pathogenic" in ClinVar, with "Review by expert panel" as review status or with "research" method and at least one publication experimentally validating the *effect*. Panel A-E show the predictions from for CADD (phred-like scaled scores), DANN, EIGEN, GERP++ score, and synVep, respectively. Grey dashed lines show the method author-recommended cutoffs, where available. Differences between scores of *Benign* and *Pathogenic* variants in panels A-E are all statistically significant (one-sided permutation test p-value = 0). Panel F reports auROC and auPRC for each predictor; all predictor auROCs and auPRCs are significantly different from that of synVep (p-value<0.05; Methods). Note that the utility of auROC and auPRC is limited in single variant effect prediction; instead, a cutoff is needed.

All variant-effect predictors, including synVep, assigned higher scores to *pathogenic* than *benign* variants (Figure 5A-C and E, all statistically significant, one-sided permutation test p-value = 0). Notably, conservation (GERP++ (69)) carried sufficient signal to recognize pathogenic variants as well, suggesting that these are often found in conserved positions, which may not be the case for variants of less severe effect. Note that all other predictors (CADD, DANN, and EIGEN) incorporate GERP++ as a feature, but their auROC and auPRC are not substantially higher (or even lower) than those of GERP++. Highly conserved genomic positions often have experienced extensive purifying selection (108). Therefore, conservation is understandably a commonly used feature for disease variant prioritization (109). However, in the scenario of disease variant prioritization synVep, offers discriminative power independent of conservation, so it may be used in combination with a conservation score or other predictors.

One major challenge in disease variant prioritization is that for complex diseases, causality can rarely be

explained by a single variant (110). The utility of variant pathogenicity score is thus questionable: does a

high score suggest a high likelihood of an individual developing a disease or a high likelihood of this

variant contributing to a disease? Also, would an individual with many predicted-pathogenic variants carry

many diseases or be very certain to carry at least one disease? A potential way to solve this puzzle is to

establish the variant-disease relationship using the collective *effect* from the whole variome instead of a

single or a few variants. A modification of polygenic risk scoring methods (111,112) to only account for

effect variants may represent one approach, although it would be limited by the location of most GWAS

SNPs in non-coding regions. Another approach is to unite only the coding variant effects by aggregating

all variants per gene to predict disease predisposition (*e.g.*, (113,114)) synVep predictions (as well as

those of other predictors) may be plugged in these pipelines to explore the contribution of sSNVs to

complex diseases.

synVep highlights correlation between conservation and effect. We annotated all sSNVs as CSVs (cross-

species variation) or not (Figure 2; Methods). CSVs are codon differences between the human reference

sequence and another species' ortholog. For example, if the proline-coding codon in a human transcript *T*

is CCC, while the aligned proline codon on *T*'s chimp ortholog is CCT, then the human sSNV CCC->CCT

is considered a chimp-CSV. We thus annotated 15,618,155 unique (only exists in one species) and

35,102,565 non-unique (overlapping across species) CSVs (Supplementary Figure 8). Since less than

10% (7,026 of 72,400) of the human transcripts can be mapped to orthologs in all 20 species, we

analyzed separately the CSVs in (1) *all transcripts* (n=32,264,860) and (2) only the transcripts that have

orthologs in all 20 species (n=3,321,574) and that are likely ancient (*ancient genes*) (115) .

Figure 6. Variant effect prediction from the perspective of cross-species variation (CSV). Panels **A-C** show synVep-predicted scores for variants grouped by the number of species carrying the mutant nucleotide; separately for *observed*, *singletons*, and *not-seen* sets. The red dashed line is synVep's default cutoff for *effect* and *no-effect*. The number in each box indicates the number of variants of that group (in thousands). Panels **D-F** show the median score (y-axis) across species at log$_2$ million years since divergence from common ancestor with human (x-axis) and linear regression trendline (red line) between the two. The Spearman correlations between median synVep score and log2(million years) for panel D-F are 0.68, 0.64, and 0.66, respectively.

The distribution of synVep prediction scores for CSVs in the *ancient genes* and for those in *all transcripts* were similar (mean=0.05, Mann-Whitney U test p-value<2.2e-16), suggesting that synVep's evaluation of variants does not discriminate by gene age. For *all transcripts*, *observed* sSNVs had more CSVs (67%, n=2,823,142) than did the *not-seen* variants (53%, n=26,976,016; Supplementary Figure 9). CSVs overall were predicted less likely to be *effect* [reviewer3] than non-CSV for both *ancient* and *all transcripts* (Figure 6 A-C; Supplementary Figure 10 A-C). While this is in line with the scoring trends of the *observed* and *not-seen* variants overall, it also mirrors earlier findings of few CSV nsSNVs corresponding to a known human disease (116-119). synVep scores also trended lower for CSVs whose substituting nucleotide was found in more species, for both *ancient* (Figure 6 A-C) and *all transcripts* (Supplementary Figure 10 A-C). Since the number of CSV species is somewhat indicative of codon conservation, this trend suggests that, although synVep was trained without using conservation features, its predictions still identify conserved codons that are often functionally relevant (120).

To further elucidate the effect of sequence conservation across species, we calculated codon mutation
fraction (CMF, Supplementary Eqn. 9) to describe how common a human's alternative codon is,
compared to the reference codon, among the 20 species included for CSV analysis. For example, if in a
multiple sequence alignment of the 20 species orthologs, the human CCC codon is aligned to 10 CCC, 5
CCT, and 5 other codons, then the CMF of the corresponding synonymous variant, CCC>CCT, is
5/15=0.33. We observed that predicted scores generally decrease with higher CMF (Supplementary
Figure 11 A-C), indicating that sSNVs with alternative codons commonly present as reference codons
among other species have less *effect*.

We additionally investigated the relationship between the evolutionary distance of CSV species from
human and the *effects* of the corresponding sSNVs. Since one sSNV can correspond to multiple species
CSVs, we only considered CSVs that are uniquely found in one species for this evaluation. The medians
of synVep scores of these species-exclusive CSVs in both *ancient genes* (Figure 6 D-F) and *all*
transcripts (Supplementary Figure 10 D-F) correlated with the evolutionary distance of the corresponding
species to human. However, for ancient genes, the median scores of *observed* variants unique to further
related (*i.e.* beyond Tarsier) species were in the *effect* range (synVep>0.5). Arguably, this means that
human sSNVs that introduce nucleotides likely present in recent ancestors tend to be *no-effect*, while
similarity to further removed relatives carries no such benefit (Figure 6 D). These findings agree with our
recent work on nsSNV CSV analysis (35). We note that species relationship had much less impact on
binary *effect* classification for singleton variants and none for not-seen variants (Figure 6E-F). The same
observations could not be made for the *all transcript* set of variants, where *observed* and *singleton* CSVs
were predicted to be *no-effect* for a large portion of species (Supplementary Figure 10 D-F). This
observation suggests that ancient genes are functionally crucial and have been sufficiently optimized over
time to only permit minor levels of variation without impact on functionality.

synVep differentiates splice-disrupting variants. Cheung et al. (70) measured the splice-disrupting effects
of genomic variants (3,297 transcript-specific sSNVs) and defined a group of large-effect splice-disrupting
variants (140 SDV sSNVs). As expected, synVep scores of SDVs were on average higher than those of
non-SDVs (Figure 7 A-C). Curiously, 140 SDVs comprised only six *observed* (4.6%) and 18 *singletons*
(13.7%) variants; nine were deemed *unobservable* (6.4%) and 107 were *not-seen* (76.4%). The fact that
most of SDVs are *not-seen* reinforces our assumption that *not-seen* sSNVs are enriched for large-effect
deleterious sSNVs that may have been purified.

Figure 7. Evaluation of synVep, CADD-splice, and spliceAI using large-effect splice-disrupting variants
(SDVs) and non-SDVs. SynVep predictions are higher scoring for a set of experimentally determined SDV
(positive; n=140) than non-SDV (negative; n=3,157) variants across *observed* (**A**), *singleton* (**B**), and *not-seen* (**C**)
data sets. Note that non-SDVs may still carry other functional effects. The *observed* SDVs mispredicted as *no-effect*
highlight the limitations of our training data, although two of the five (40%) *observed* SDVs are correctly annotated
as *effect*. Panels **D-F** show the distribution of scores on complete non-SDV and SDV collections predicted by
CADD-splice, spliceAI, and synVep, respectively. Panel G reports tha auROC and *effect* auPRC measures. CADD-
splice auROC and auPRC are significantly different from synVep (p-value<0.05; Methods); while spliceAI's are not.

We evaluated two state-of-the-art predictors for splicing effect evaluation: CADD-splice (72) and spliceAI

(71) on this set of experimentally determined variants (Figure 7D-G). spliceAI is a 32-layer deep learning

model that predicts splicing donor/acceptor gain/loss probabilities. CADD-splice was developed based on

CADD (23) with the addition of two splicing-specific predictor (MMsplice (121) and spliceAI) outputs as

features. MMsplice and spliceAI were selected to be incorporated into CADD-splice because they

performed best among several other splicing-specific predictors on the same non-SDV/SDV dataset (not

limited to synonymous variants). CADD-splice has higher auROC (0.75) than both spliceAI (auROC=0.67)

and synVep (auROC=0.67); meanwhile, the auRPC of all the three predictors are similar (Figure 7 G).

Splicing disruption is a well-known and well-studied mechanism of sSNV *effect* (122). In fact, most of the

experimental validations of our *curated-effect* and ClinVar pathogenic variants refer to elucidating splicing

effects (Supplementary Table S1 and S3). Moreover, many cancer driver mutations are found to be

splice-disrupting synonymous variants (123). Aside from splicing, experimental validation of variant *effect*

is rare, arguably due to technical challenges (124). Perhaps, since the experimental evidence for splicing

disruption is more abundant than non-splicing effects', the former is considered a major factor in clinical

consideration for sSNVs. For example, according to the guidelines from American College of Medical Genetics and Genomics (125), an sSNV is clinically benign if it is not in a conserved position and is predicted to be non-impacting to a splice site (*e.g.* via GeneSplicer (126), NNsplice (127)). Thus, synVep's ability to identify *effect* and score sSNVs regardless of their splice effects or conservation makes it an ideal tool for prioritization of all possible variants, regardless of their mechanism or evolutionary evidence of *effect*.

sSNV effects reflect genomic constraints. Havrilla *et al.* developed the concept of "coding constrained regions" (CCR) to describe the regional scarcity of protein-changing (missense or loss-of-function) variants in the human genome (76). Here, a region with fewer of these variants observed in the human population has a higher CCR percentile score. For our set of variants, the fraction of *observed* (number of *observed* sSNVs divided by all possible sSNVs in this region) negatively correlated (Pearson ρ=-0.61) with CCR percentile (Figure 8 A); *i.e.* higher constraint indicates fewer sSNVs. Furthermore, synVep predictions positively correlated with CCR percentiles for *observed* (ρ=0.58, Figure 8 B), *i.e.* lower CCR percentile (less constrained regions) indicated lower (*no-effect*) synVep scores.

Figure 8. sSNV effect measured by region constraint. Coding constrained regions (CCR) describe the regional scarcity of nsSNVs; higher percentile regions represent have fewer observed nsSNVs. *Observed* sSNVs are relatively scarce in constrained regions **(A)**, while their median synVep scores are higher **(B)**. Pearson correlation are indicated in blue.

The negative correlation between the fraction of sSNVs and CCR indicates a positive correlation between synonymous mutation rate and missense or loss-of-function mutation rate. This observation is in line with

earlier studies (128,129), but raises a question of the utility of Ka/Ks ratio (non-synonymous divided by
synonymous mutation rate), which is widely used to measure the strength of evolutionary selection at
certain genomic sites (130). The application of the Ka/Ks ratio is based on the assumption that
synonymous mutations are neutral and thus Ks can serve as a baseline for Ka. However, it has been
demonstrated that a high Ka/Ks can also result from a low Ks due to strong negative selection at the
synonymous sites (10,11,131-133). Efforts have been made to improve the utility of Ka/Ks by
incorporating codon preference (134-136), but the question remains: how often is the selection at
synonymous sites sufficiently underestimated so that Ka/Ks is no longer accurate? Lawrie et al. found
that 22% of the fourfold synonymous sites (where the amino acid can be encoded by four codons) in the
fruitfly genome are under strong selection (137). Lu and Wu estimated that 90% synonymous mutations
in human and chimp are deleterious (138). Hellmann et al. estimated that 39% mutations at the human-
chimp-diverged non-CpG fourfold synonymous sites have been purified (139). Zhou et al. showed that 9%
of all yeast genes and 5% all worm genes undergo purifying selection on synonymous sites (136). In turn,
our results show that, excluding *unobservable* (9.6%), ~67% of all possible human sSNVs are *effect*
(synVep score>0.5), but we cannot estimate the strength of selection acting upon these. These findings
suggest that Ka/Ks measures of genomic site constraints may be underpowered.

synVep sheds light on future variant discovery and interpretation. Whenever a human genome variant is
sequenced, it will automatically be reassigned a class in our collection. Thus, a newly sequenced variant
will first become a *singleton* and may, eventually, be a member of the *observed* group. An enrichment in
observed variants will likely come from large-scale sequencing. The ethnic diversity of gnomAD
represents the ethnic diversity in the United States, but not the global ethnicity diversity; although only
16% of global population are of European descent (140), 53% of the samples from gnomAD exomes
database are (141) are; i.e. there is a significant underrepresentation of sSNVs from other ethnicities.
When more diverse genomes are sequenced, will there be a significant addition to the *observed* set (*i.e.*
significant reduction of the *not-seen* set)?

To answer this question, we obtained all variants from the Qatar Genome (QTRG) project (77) and
mapped them to our set of sSNVs. QTRG comprises 1,376 individuals and may serve as a representative
pool of genomic variants in Middle East and north Africa (MENA) area (77); thus, this set is
complementary to gnomAD. We identified 526,616 transcript-based sSNVs (n=192,246 genomic
coordinate-based) from QTRG sequencing. Importantly, only 0.6% of the Qatari sSNVs mapped to our
unobservable set – a fraction that is lower than the misprediction rate (5%) that we allowed during PUL.
Moreover, two thirds of these variants were singletons in QTRG. This observation suggests that our
unobservable variants are indeed unlikely to be ever observed in future sequenced human populations.
The majority of QTRG sSNVs (81.9%) mapped to our *observed* set; 4.6% and 12.8% were *singleton* and
not-seen, respectively (Figure 9A). Interestingly, 63.5% and 64.6% QTRG sSNVs mapping to our
singleton and *not-seen* sets, respectively, were singletons in the Qatari cohort. We also found that
gnomAD *singletons* that were present in QTRG, on average, scored higher than QTRG variants

overlapping with *observed* sSNVs (34.7% vs 26.6% *effect* variants, respectively). Thus finding further

confirms that *singletons* are more likely to have an effect than *observed* variants.

Figure 9. Distributions of the Qatar Genome sSNVs. In both panels, our gnomAD-based *observed* (orange),
singleton (blue) and *not-seen* (dark orange) sets are highlighted. **(A)** represents the fraction of the QTRG sSNVs
mapped our *observed*, *singleton*, *not-seen*, and *unobservable* (gray) sets. **(B)** synVep scores for our (gnomAD-based)
variant sets, as well as the scores for QTRG sSNVs (white) mapping to the corresponding gnomAD-classes.
Importantly, the synVep scores of QTRG variants that were previously classified as singletons or not-seen score
much lower than other variants in the corresponding groups.

How many of the previously *not-seen* sSNVs are *effect*? New sSNVs are likely to come from clinical

sequencing and could thus could often be deemed disease-associated. We expect, however, that these

variants will carry little or no *effect*. In other words, currently *not-seen no-effect* observable sSNVs

(n=14,259,180 transcript-based and n=5,975,076 genomic coordinate-based) are more likely to be

discovered in the future than an *effect* ones – even if a sample is taken from a sick individual. Recall our

assumption that the *not-seen* set is composed of those sSNVs that carry a large effect and have been

purified, as well as those that are putatively neutral and will be seen in the future if more sequencing is

performed. The synVep scores of the QTRG sSNVs mapping to our *not-seen* set were, on average, much

lower than those of the entire *not-seen* set (Figure 9B, average synVep score 0.49 vs. 0.88, Mann-

Whitney U test p-value<2.2e-16). Similarly, the median synVep score of the 2,469,205 sSNVs *not-seen*

according to gnomAD, but present in dbSNP (142) was lower than for the entire *not-seen* set (0.47 vs

0.88; p-value<2.2e-16). These results confirm our assumption, as these newly identified sSNVs are

actually observable (not purified) and thus they are generally less likely to have large effect (and thus

lower synVep scores). It may also be that the newly identified predicted *effect* variants (from QTRG, and

other sequencing efforts in the future) are the ethnicity-differentiating, *i.e.* not necessarily affecting overall

fitness, but contributing to individual differences (as in *e.g.* (143)).

CONCLUSION

We developed synVep—a machine learning-based model for evaluating the *effect* of human sSNVs. Our

model does not use disease/deleteriousness-labeled training data. Instead, we used the signals derived

from *observed* (and corresponding *generated*) sSNVs from large sequencing projects. Our model

successfully distinguishes sSNVs with experimentally validated *effect*, e.g. splice-site disrupters, as well

as pathogenic sSNVs. Moreover, our model's predictions of cross-species variants (CSVs) correlate with

the evolutionary distance between human and CSV-species. While further experimental validations of

effect prediction are necessary, synVep's evaluation on sSNV *effect* will greatly contribute to our

understanding of biological molecular pathways in general, and of pathogenicity pathways in particular.

DATA AVAILABILITY

synVep webserver for online query: https://services.bromberglab.org/synvep; For local run, Python script

(https://bitbucket.org/bromberglab/synvep_local) and prediction database

(https://zenodo.org/record/4763256) are also available.

AUTHOR CONTRIBUTION

ZZ and YB designed the study, evaluated the results, and wrote the manuscript; ZZ conducted the study; ZZ and AA

built the webserver.

ACKNOWLEDGEMENT

We thank our current and former lab members, Dr. Yannick Mahlich, Dr. Chengsheng Zhu, Dr. Maximillian Miller,

and Dr. Yanran Wang (all Rutgers), for all discussions and constructive suggestions. We also thank Kyle Flannery

(Rutgers) for the idea of testing our predictions with Qatari Genome variant data. We are also grateful to the

Rutgers Office of Advanced Research Computing (OARC) for making high-performance compute resources

available to this project, Thomas Pawlowski (Rutgers Office of Information Technology) for setting up the host of

synVep webserver, and to the Ensembl team for their help and feedback. Last but not least, we want to thank all

researchers and human subjects who made the data and tools used in this study available.

FUNDING

ZZ and YB were supported by the NIH/NIGMS grant R01 [GM115486]; AA is supported by the Astrobiology Institute

grant [80NSSC18M0093]; YB was also supported by NIH grant R01 [MH115958].

CONFLICT OF INTEREST

None declared.

REFERENCES

1.   Ashley, E.A. (2016) Towards precision medicine. *Nature Reviews Genetics*, **17**, 507.

2.   Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A. and Yang, J. (2017)
10 years of GWAS discovery: biology, function, and translation. *The American Journal of Human*
Genetics, **101**, 5-22.

3.   Pan, B., Kusko, R., Xiao, W., Zheng, Y., Liu, Z., Xiao, C., Sakkiah, S., Guo, W., Gong, P., Zhang,
C. *et al.* (2019) Similarities and differences between variants called with human reference
genome HG19 or HG38. *BMC Bioinformatics*, **20**, 101.

4.   Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Abecasis, G.R., Bentley, D.R.,
Chakravarti, A., Clark, A.G., Donnelly, P., Eichler, E.E. *et al.* (2015) A global reference for human
genetic variation. *Nature*, **526**, 68-74.

5.   Ward, L.D. and Kellis, M. (2012) Interpreting noncoding genetic variation in complex traits and
human disease. *Nature biotechnology*, **30**, 1095-1106.

6.   Zhu, C., Miller, M., Zeng, Z., Wang, Y., Mahlich, Y., Aptekmann, A. and Bromberg, Y. (2020)
Computational Approaches for Unraveling the Effects of Variation in the Human Genome and
Microbiome. *Annual Review of Biomedical Data Science*, **3**.

7.   Shen, H., Li, J., Zhang, J., Xu, C., Jiang, Y., Wu, Z., Zhao, F., Liao, L., Chen, J., Lin, Y. *et al.*
(2013) Comprehensive Characterization of Human Genome Variation by High Coverage Whole-
Genome Sequencing of Forty Four Caucasians. **8**, e59494.

8.   Sauna, Z.E. and Kimchi-Sarfaty, C. (2011) Understanding the contribution of synonymous
mutations to human disease. *Nature reviews. Genetics*, **12**, 683.

9.   Stergachis, A.B., Haugen, E., Shafer, A., Fu, W., Vernot, B., Reynolds, A., Raubitschek, A.,
Ziegler, S., LeProust, E.M. and Akey, J.M. (2013) Exonic transcription factor binding directs
codon choice and affects protein evolution. *Science*, **342**, 1367-1372.

10.   Pagani, F., Raponi, M. and Baralle, F.E. (2005) Synonymous mutations in CFTR exon 12 affect
splicing and are not neutral in evolution. *Proceedings of the National Academy of Sciences*, **102**,
6368-6372.

11.   Chamary, J.V., Parmley, J.L. and Hurst, L.D. (2006) Hearing silence: non-neutral evolution at
synonymous sites in mammals. *Nature Reviews Genetics*, **7**, 98-108.

12.   Meyer, I.M. (2005) Statistical evidence for conserved, local secondary structure in the coding
regions of eukaryotic mRNAs and pre-mRNAs. **33**, 6338-6348.

13.   Duan, J., Shi, J., Ge, X., Dölken, L., Moy, W., He, D., Shi, S., Sanders, A.R., Ross, J. and
Gejman, P.V. (2013) Genome-wide survey of interindividual differences of RNA stability in human
lymphoblastoid cell lines. *Scientific reports*, **3**, 1318.

14.   Shah, P., Ding, Y., Niemczyk, M., Kudla, G. and Joshua. (2013) Rate-Limiting Steps in Yeast
Protein Translation. *Cell*, **153**, 1589-1601.

15.   Pechmann, S. and Frydman, J. (2013) Evolutionary conservation of codon optimality reveals
hidden signatures of cotranslational folding. *Nature structural & molecular biology*, **20**, 237.

16.   Stoletzki, N. and Eyre-Walker, A. (2006) Synonymous codon usage in Escherichia coli: selection
for translational accuracy. *Molecular biology and evolution*, **24**, 374-381.

17.   Zeng, Z. and Bromberg, Y. (2019) Predicting Functional Effects of Synonymous Variants: A
Systematic Review and Perspectives. *Frontiers in genetics*, **10**.

18.   Buske, O.J., Manickaraj, A., Mital, S., Ray, P.N. and Brudno, M. (2013) Identification of
deleterious synonymous variants in human genomes. *Bioinformatics*, **29**, 1843-1850.

19.   Zhang, X., Li, M., Lin, H., Rao, X., Feng, W., Yang, Y., Mort, M., Cooper, D.N., Wang, Y. and
Wang, Y. (2017) regSNPs-splicing: a tool for prioritizing synonymous single-nucleotide
substitution. *Human genetics*, **136**, 1279-1289.

20.   Livingstone, M., Folkman, L., Yang, Y., Zhang, P., Mort, M., Cooper, D.N., Liu, Y., Stantic, B. and
Zhou, Y. (2017) Investigating DNA-, RNA-, and protein-based features as a means to
discriminate pathogenic synonymous variants. *Human mutation*, **38**, 1336-1347.

21.   Gelfman, S., Wang, Q., McSweeney, K.M., Ren, Z., La Carpia, F., Halvorsen, M., Schoch, K.,
Ratzon, F., Heinzen, E.L. and Boland, M.J. (2017) Annotating pathogenic non-coding variants in
genic regions. *Nature communications*, **8**, 236.

22.   Shi, F., Yao, Y., Bin, Y., Zheng, C.-H. and Xia, J. (2019) Computational identification of
deleterious synonymous variants in human genomes using a feature-based approach. *BMC*
medical genomics, **12**, 12.

23.   Kircher, M., Witten, D.M., Jain, P., O'roak, B.J., Cooper, G.M. and Shendure, J. (2014) A general
framework for estimating the relative pathogenicity of human genetic variants. *Nature genetics*,
46, 310.

24.   Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J. and Kircher, M. (2019) CADD: predicting
the deleteriousness of variants throughout the human genome. *Nucleic acids research*, **47**, D886-
D894.

25.   Quang, D., Chen, Y. and Xie, X. (2014) DANN: a deep learning approach for annotating the
pathogenicity of genetic variants. *Bioinformatics*, **31**, 761-763.

26.   Shihab, H.A., Rogers, M.F., Gough, J., Mort, M., Cooper, D.N., Day, I.N., Gaunt, T.R. and
Campbell, C. (2015) An integrative approach to predicting the functional effects of non-coding
and coding sequence variation. *Bioinformatics*, **31**, 1536-1543.

27.   Schwarz, J.M., Cooper, D.N., Schuelke, M. and Seelow, D. (2014) MutationTaster2: mutation
prediction for the deep-sequencing age. *Nature methods*, **11**, 361.

28.   Stenson, P.D., Mort, M., Ball, E.V., Evans, K., Hayden, M., Heywood, S., Hussain, M., Phillips,
A.D. and Cooper, D.N. (2017) The Human Gene Mutation Database: towards a comprehensive
repository of inherited mutation data for medical research, genetic diagnosis and next-generation
sequencing studies. *Human genetics*, **136**, 665-677.

29.   Landrum, M.J., Lee, J.M., Riley, G.R., Jang, W., Rubinstein, W.S., Church, D.M. and Maglott, D.R.
(2014) ClinVar: public archive of relationships among sequence variation and human phenotype.
Nucleic Acids Research, **42**, D980-D985.

30.   Bromberg, Y., Kahn, P.C. and Rost, B. (2013) Neutral and weakly nonneutral sequence variants
may define individuality. **110**, 14255-14260.

31.   Pejaver, V., Babbi, G., Casadio, R., Folkman, L., Katsonis, P., Kundu, K., Lichtarge, O., Martelli,
P.L., Miller, M. and Moult, J. (2019) Assessment of methods for predicting the effects of PTEN
and TPMT protein variants. *Human mutation*, **40**, 1495-1506.

32.   Liu, B., Lee, W.S., Yu, P.S. and Li, X. (2002), *ICML*. Citeseer, Vol. 2, pp. 387-394.

33.   Liu, B., Dai, Y., Li, X., Lee, W.S. and Yu, P.S. (2003), *Third IEEE International Conference on*
Data Mining. IEEE, pp. 179-186.

34.   Fusilier, D.H., Montes-y-Gómez, M., Rosso, P. and Cabrera, R.G. (2015) Detecting positive and
negative deceptive opinions using PU-learning. *Information processing & management*, **51**, 433-
443.

35.   Mahlich, Y., Miller, M., Zeng, Z. and Bromberg, Y. (2021) Low diversity of human variation despite
mostly mild functional impact of de novo variation. *Frontiers in Molecular Biosciences*, **8**, 74.

36.   Kinsella, R.J., Kahari, A., Haider, S., Zamora, J., Proctor, G., Spudich, G., Almeida-King, J.,
Staines, D., Derwent, P., Kerhornou, A. *et al.* (2011) Ensembl BioMarts: a hub for data retrieval
across taxonomic space. **2011**, bar030-bar030.

37.   Church, D.M., Schneider, V.A., Graves, T., Auger, K., Cunningham, F., Bouk, N., Chen, H.-C.,
Agarwala, R., McLaren, W.M., Ritchie, G.R.S. *et al.* (2011) Modernizing Reference Genome
Assemblies. **9**, e1001091.

38.   Wang, K., Li, M. and Hakonarson, H. (2010) ANNOVAR: functional annotation of genetic variants
from high-throughput sequencing data. *Nucleic Acids Research*, **38**, e164-e164.

39.   Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L.,
Laricchia, K.M., Ganna, A., Birnbaum, D.P. *et al.* (2019) Variation across 141,456 human exomes

and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding
genes. *bioRxiv*, 531210.

40.   Freedman, D., Pisani, R., Purves, R. and Adhikari, A. (2007). WW Norton & Company New York.

41.   Acock, A.C. and Stavig, G.R. (1979) A measure of association for nonparametric statistics. *Social*
Forces, **57**, 1381-1386.

42.   Fisher, R.A. (1992), *Breakthroughs in statistics*. Springer, pp. 66-70.

43.   Lonsdale, J., Thomas, J., Salvatore, M., Phillips, R., Lo, E., Shad, S., Hasz, R., Walters, G.,
Garcia, F. and Young, N. (2013) The genotype-tissue expression (GTEx) project. *Nature genetics*,
45, 580.

44.   Sharp, P.M. and Li, W.-H. (1987) The codon adaptation index-a measure of directional
synonymous codon usage bias, and its potential applications. *Nucleic acids research*, **15**, 1281-
1295.

45.   Ikemura, T. (1981) Correlation between the abundance of Escherichia coli transfer RNAs and the
occurrence of the respective codons in its protein genes: a proposal for a synonymous codon
choice that is optimal for the E. coli translational system. *Journal of molecular biology*, **151**, 389-
409.

46.   Karlin, S. and Mrázek, J. (1996) What drives codon choices in human genes? *Journal of*
molecular biology, **262**, 459-472.

47.   Freire-Picos, M.A., Gonzalez-Siso, M.I., Rodríguez-Belmonte, E., Rodriguez-Torres, A.M., Ramil,
E. and Cerdan, M.E. (1994) Codon usage in Kluyveromyces lactis and in yeast cytochrome c-
encoding genes. *Gene*, **139**, 43-49.

48.   Wan, X., Xu, D. and Zhou, J. (2003) A new informatics method for measuring synonymous codon
usage bias. *Intelligent engineering systems through artificial neural networks Volume*, **13**.

49.   Dos Reis, M., Wernisch, L. and Savva, R. (2003) Unexpected correlations between gene
expression and codon usage bias from microarray data for the whole Escherichia coli K-12
genome. *Nucleic acids research*, **31**, 6976-6985.

50.   Team, R.C. (2013) R: A language and environment for statistical computing.

51.   Supek, F. and Vlahoviček, K. (2005) Comparison of codon usage measures and their applicability
in prediction of microbial gene expressivity. *BMC bioinformatics*, **6**, 182.

52.   Cannarozzi, G., Schraudolph, N.N., Faty, M., Von Rohr, P., Friberg, M.T., Roth, A.C., Gonnet, P.,
Gonnet, G. and Barral, Y. (2010) A Role for Codon Order in Translation Dynamics. *Cell*, **141**,
355-367.

53.   Yevshin, I., Sharipov, R., Valeev, T., Kel, A. and Kolpakov, F. (2016) GTRD: a database of
transcription factor binding sites identified by ChIP-seq experiments. *Nucleic acids research*,
gkw951.

54.   Zhao, H., Sun, Z., Wang, J., Huang, H., Kocher, J.-P. and Wang, L. (2013) CrossMap: a versatile
tool for coordinate conversion between genome assemblies. *Bioinformatics*, **30**, 1006-1007.

55.   Giudice, G., Sánchez-Cabo, F., Torroja, C. and Lara-Pezzi, E. (2016) ATtRACT—a database of
RNA-binding proteins and associated motifs. *Database*, **2016**.

56.   Cáceres, E.F. and Hurst, L.D. (2013) The evolution, impact and properties of exonic splice
enhancers. *Genome biology*, **14**, R143.

57.   Yachdav, G., Kloppmann, E., Kajan, L., Hecht, M., Goldberg, T., Hamp, T., Hönigschmid, P.,
Schafferhans, A., Roos, M. and Bernhofer, M. (2014) PredictProtein—an open resource for online
prediction of protein structural and functional features. *Nucleic acids research*, **42**, W337-W343.

58.   Rost, B. (1996), *Methods in enzymology*. Elsevier, Vol. 266, pp. 525-539.

59.   Schlessinger, A., Punta, M., Yachdav, G., Kajan, L. and Rost, B. (2009) Improved disorder
prediction by combination of orthogonal approaches. *PloS one*, **4**, e4433.

60.   Lorenz, R., Bernhart, S.H., Zu Siederdissen, C.H., Tafer, H., Flamm, C., Stadler, P.F. and
Hofacker, I.L. (2011) ViennaRNA Package 2.0. *Algorithms for Molecular Biology*, **6**, 26.

61.   Sabarinathan, R., Tafer, H., Seemann, S.E., Hofacker, I.L., Stadler, P.F. and Gorodkin, J. (2013)
RNA snp: efficient detection of local RNA secondary structure changes induced by SNP s.
Human mutation, **34**, 546-556.

62.   LeCun, Y., Bengio, Y. and Hinton, G. (2015) Deep learning. *nature*, **521**, 436-444.

63.   Chen, T. and Guestrin, C. (2016), *Proceedings of the 22nd acm sigkdd international conference*
on knowledge discovery and data mining. ACM, pp. 785-794.

64.   Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R. and Dubourg, V. (2011) Scikit-learn: Machine learning in Python.
Journal of machine learning research, **12**, 2825-2830.
65.   Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C. and Müller, M. (2011)
pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC*
bioinformatics, **12**, 1-8.
66.   Grau, J., Grosse, I. and Keilwagen, J. (2015) PRROC: computing and visualizing precision-recall
and receiver operating characteristic curves in R. *Bioinformatics*, **31**, 2595-2597.
67.   Ionita-Laza, I., McCallum, K., Xu, B. and Buxbaum, J.D. (2016) A spectral approach integrating
functional genomic annotations for coding and noncoding variants. *Nature genetics*, **48**, 214.
68.   Landrum, M.J. and Kattman, B.L. (2018) ClinVar at five years: Delivering on the promise. *Human*
Mutation, **39**, 1623-1630.
69.   Davydov, E.V., Goode, D.L., Sirota, M., Cooper, G.M., Sidow, A. and Batzoglou, S. (2010)
Identifying a high fraction of the human genome to be under selective constraint using GERP++.
PLoS computational biology, **6**, e1001025.
70.   Cheung, R., Insigne, K.D., Yao, D., Burghard, C.P., Wang, J., Hsiao, Y.-H.E., Jones, E.M.,
Goodman, D.B., Xiao, X. and Kosuri, S. (2019) A multiplexed assay for exon recognition reveals
that an unappreciated fraction of rare genetic variants cause large-effect splicing disruptions.
Molecular cell, **73**, 183-194. e188.
71.   Jaganathan, K., Panagiotopoulou, S.K., McRae, J.F., Darbandi, S.F., Knowles, D., Li, Y.I.,
Kosmicki, J.A., Arbelaez, J., Cui, W. and Schwartz, G.B. (2019) Predicting splicing from primary
sequence with deep learning. *Cell*, **176**, 535-548. e524.
72.   Rentzsch, P., Schubach, M., Shendure, J. and Kircher, M. (2021) CADD-Splice—improving
genome-wide variant effect prediction using deep learning-derived splice scores. *Genome*
medicine, **13**, 1-12.
73.   Kumar, S., Stecher, G., Suleski, M. and Hedges, S.B. (2017) TimeTree: a resource for timelines,
timetrees, and divergence times. *Molecular biology and evolution*, **34**, 1812-1819.
74.   Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M.R., Barrell, D., Bhai, J., Billis, K., Cummins,
C., Gall, A. and Girón, C.G. (2017) Ensembl 2018. *Nucleic acids research*, **46**, D754-D761.
75.   Löytynoja, A. and Goldman, N. (2005) An algorithm for progressive multiple alignment of
sequences with insertions. *Proceedings of the National Academy of Sciences*, **102**, 10557-10562.
76.   Havrilla, J.M., Pedersen, B.S., Layer, R.M. and Quinlan, A.R. (2019) A map of constrained coding
regions in the human genome. *Nature genetics*, **51**, 88-95.
77.   Fakhro, K.A., Staudt, M.R., Ramstetter, M.D., Robay, A., Malek, J.A., Badii, R., Al-Marri, A.A.-N.,
Abi Khalil, C., Al-Shakaki, A. and Chidiac, O. (2016) The Qatar genome: a population-specific tool
for precision medicine in the Middle East. *Human genome variation*, **3**, 1-7.
78.   Leinonen, R., Sugawara, H., Shumway, M. and Collaboration, I.N.S.D. (2010) The sequence read
archive. *Nucleic acids research*, **39**, D19-D21.
79.   Dohm, J.C., Lottaz, C., Borodina, T. and Himmelbauer, H. (2008) Substantial biases in ultra-short
read data sets from high-throughput DNA sequencing. *Nucleic acids research*, **36**, e105.
80.   Rieber, N., Zapatka, M., Lasitschka, B., Jones, D., Northcott, P., Hutter, B., Jäger, N., Kool, M.,
Taylor, M. and Lichter, P. (2013) Coverage bias and sensitivity of variant calling for four whole-
genome sequencing technologies. *PloS one*, **8**, e66621.
81.   Forsdyke, D.R. (2001) Functional constraint and molecular evolution. *e LS*.
82.   Alazami, A.M., Awad, S.M., Coskun, S., Al-Hassan, S., Hijazi, H., Abdulwahab, F.M., Poizat, C.
and Alkuraya, F.S. (2015) TLE6 mutation causes the earliest known human embryonic lethality.
Genome biology, **16**, 1-8.
83.   Shamseldin, H.E., Tulbah, M., Kurdi, W., Nemer, M., Alsahan, N., Al Mardawi, E., Khalifa, O.,
Hashem, A., Kurdi, A. and Babay, Z. (2015) Identification of embryonic lethal genes in humans by
autozygosity mapping and exome sequencing in consanguineous families. *Genome biology*, **16**,
116.
84.   Shao, Z.-Q., Zhang, Y.-M., Feng, X.-Y., Wang, B. and Chen, J.-Q. (2012) Synonymous codon
ordering: a subtle but prevalent strategy of bacteria to improve translational efficiency. *PLoS One*,
7, e33547.

85. Tesina, P., Lessen, L.N., Buschauer, R., Cheng, J., Wu, C.C.C., Berninghausen, O., Buskirk,
A.R., Becker, T., Beckmann, R. and Green, R. (2020) Molecular mechanism of translational
stalling by inhibitory codon combinations and poly (A) tracts. *The EMBO journal*, **39**, e103365.
86. Gamble, C.E., Brule, C.E., Dean, K.M., Fields, S. and Grayhack, E.J. (2016) Adjacent codons act
in concert to modulate translation efficiency in yeast. *Cell*, **166**, 679-690.
87. Boël, G., Letso, R., Neely, H., Price, W.N., Wong, K.-H., Su, M., Luff, J.D., Valecha, M., Everett,
J.K. and Acton, T.B. (2016) Codon influence on protein expression in E. coli correlates with
mRNA levels. *Nature*, **529**, 358.
88. Plotkin, J.B., Robins, H. and Levine, A.J. (2004) Tissue-specific codon usage and the expression
of human genes. *Proceedings of the National Academy of Sciences of the United States of*
America, **101**, 12588-12591.
89. Pop, C., Rouskin, S., Ingolia, N.T., Han, L., Phizicky, E.M., Weissman, J.S. and Koller, D. (2014)
Causal signals between codon bias, mRNA structure, and the efficiency of translation and
elongation. *Molecular systems biology*, **10**, 770.
90. Qian, W., Yang, J.-R., Pearson, N.M., Maclean, C. and Zhang, J. (2012) Balanced Codon Usage
Optimizes Eukaryotic Translational Efficiency. **8**, e1002603.
91. Hia, F., Yang, S.F., Shichino, Y., Yoshinaga, M., Murakawa, Y., Vandenbon, A., Fukao, A.,
Fujiwara, T., Landthaler, M. and Natsume, T. (2019) Codon bias confers stability to human mRNA
s. *EMBO reports*, **20**, e48220.
92. Presnyak, V., Alhusaini, N., Chen, Y.-H., Martin, S., Morris, N., Kline, N., Olson, S., Weinberg, D.,
Baker, K.E. and Graveley, B.R. (2015) Codon optimality is a major determinant of mRNA stability.
Cell, **160**, 1111-1124.
93. Chamary, J.-V. and Hurst, L.D. (2005) Evidence for selection on synonymous mutations affecting
stability of mRNA secondary structure in mammals. *Genome biology*, **6**, 1-12.
94. Duan, J., Wainwright, M.S., Comeron, J.M., Saitou, N., Sanders, A.R., Gelernter, J. and Gejman,
P.V. (2003) Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA
stability and synthesis of the receptor. *Human molecular genetics*, **12**, 205-216.
95. Jacobs, W.M. and Shakhnovich, E.I. (2017) Evidence of evolutionary selection for cotranslational
folding. *Proceedings of the National Academy of Sciences*, **114**, 11434-11439.
96. Kimchi-Sarfaty, C., Oh, J.M., Kim, I.-W., Sauna, Z.E., Calcagno, A.M., Ambudkar, S.V. and
Gottesman, M.M. (2007) A" silent" polymorphism in the MDR1 gene changes substrate specificity.
Science, **315**, 525-528.
97. Lorenz, R., Wolfinger, M.T., Tanzer, A. and Hofacker, I.L. (2016) Predicting RNA secondary
structures from sequence and probing data. *Methods*, **103**, 86-98.
98. Hughes, A.L. (2008) Near-neutrality: the leading edge of the neutral theory of molecular evolution.
Annals of the New York Academy of Sciences, **1133**, 162.
99. Eyre-Walker, A. and Keightley, P.D. (2007) The distribution of fitness effects of new mutations.
Nature Reviews Genetics, **8**, 610-618.
100. Wall, J.D., Tang, L.F., Zerbe, B., Kvale, M.N., Kwok, P.-Y., Schaefer, C. and Risch, N. (2014)
Estimating genotype error rates from high-coverage next-generation sequence data. *Genome*
research, **24**, 1734-1739.
101. Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-Luria,
A.H., Ware, J.S., Hill, A.J., Cummings, B.B. *et al.* (2016) Analysis of protein-coding genetic
variation in 60,706 humans. *Nature*, **536**, 285-291.
102. Bloom, J.S., Boocock, J., Treusch, S., Sadhu, M.J., Day, L., Oates-Barker, H. and Kruglyak, L.
(2019) Rare variants contribute disproportionately to quantitative trait variation in yeast. *Elife*, **8**,
e49212.
103. Purcell, S.M., Moran, J.L., Fromer, M., Ruderfer, D., Solovieff, N., Roussos, P., O'dushlaine, C.,
Chambert, K., Bergen, S.E. and Kähler, A. (2014) A polygenic burden of rare disruptive mutations
in schizophrenia. *Nature*, **506**, 185-190.
104. Teng, S., Michonova-Alexova, E. and Alexov, E. (2008) Approaches and resources for prediction
of the effects of non-synonymous single nucleotide polymorphism on protein function and
interactions. *Current pharmaceutical biotechnology*, **9**, 123-133.
105. Genovese, G., Fromer, M., Stahl, E.A., Ruderfer, D.M., Chambert, K., Landén, M., Moran, J.L.,
Purcell, S.M., Sklar, P. and Sullivan, P.F. (2016) Increased burden of ultra-rare protein-altering
variants among 4,877 individuals with schizophrenia. *Nature neuroscience*, **19**, 1433-1441.

106.   Bobbili, D.R., Banda, P., Krüger, R. and May, P. (2020) Excess of singleton loss-of-function
1039   variants in Parkinson's disease contributes to genetic risk. *Journal of Medical Genetics*.
1040   107.   Zhang, D., Cheng, L., Qian, Y., Alliey-Rodriguez, N., Kelsoe, J.R., Greenwood, T., Nievergelt, C.,
1041   Barrett, T.B., McKinney, R. and Schork, N. (2009) Singleton deletions throughout the genome
1042   increase risk of bipolar disorder. *Molecular psychiatry*, **14**, 376-380.
1043   108.   Cooper, G.M. and Shendure, J. (2011) Needles in stacks of needles: finding disease-causal
1044   variants in a wealth of genomic data. *Nature Reviews Genetics*, **12**, 628-640.
1045   109.   Eilbeck, K., Quinlan, A. and Yandell, M. (2017) Settling the score: variant prioritization and
1046   Mendelian disease. *Nature Reviews Genetics*, **18**, 599-612.
1047   110.   MacArthur, D., Manolio, T., Dimmock, D., Rehm, H., Shendure, J., Abecasis, G., Adams, D.,
1048   Altman, R., Antonarakis, S. and Ashley, E. (2014) Guidelines for investigating causality of
1049   sequence variants in human disease. *Nature*, **508**, 469-476.
1050   111.   Torkamani, A., Wineinger, N.E. and Topol, E.J. (2018) The personal and clinical utility of
1051   polygenic risk scores. *Nature Reviews Genetics*, **19**, 581-590.
1052   112.   Khera, A.V., Chaffin, M., Aragam, K.G., Haas, M.E., Roselli, C., Choi, S.H., Natarajan, P., Lander,
1053   E.S., Lubitz, S.A. and Ellinor, P.T. (2018) Genome-wide polygenic scores for common diseases
1054   identify individuals with risk equivalent to monogenic mutations. *Nature genetics*, **50**, 1219-1224.
1055   113.   Wang, Y. and Bromberg, Y. (2019) Identifying mutation-driven changes in gene functionality that
1056   lead to venous thromboembolism. *Human mutation*, **40**, 1321-1329.
1057   114.   Raimondi, D., Simm, J., Arany, A., Fariselli, P., Cleynen, I. and Moreau, Y. (2020) An
1058   interpretable low-complexity machine learning framework for robust exome-based in-silico
1059   diagnosis of Crohn's disease patients. *NAR Genomics and Bioinformatics*, **2**, lqaa011.
1060   115.   Capra, J.A., Stolzer, M., Durand, D. and Pollard, K.S. (2013) How old is my gene? *Trends in
1061   Genetics*, **29**, 659-668.
1062   116.   Briscoe, A.D., Gaur, C. and Kumar, S. (2004) The spectrum of human rhodopsin disease
1063   mutations through the lens of interspecific variation. *Gene*, **332**, 107-118.
1064   117.   Waterston, R.H. and Pachter, L. (2002) Initial sequencing and comparative analysis of the mouse
1065   genome. *Nature*, **420**, 520-562.
1066   118.   Kondrashov, A.S., Sunyaev, S. and Kondrashov, F.A. (2002) Dobzhansky–Muller
1067   incompatibilities in protein evolution. *Proceedings of the National Academy of Sciences*, **99**,
1068   14878-14883.
1069   119.   Subramanian, S. and Kumar, S. (2006) Evolutionary anatomies of positions and types of disease-
1070   associated and neutral amino acid mutations in the human genome. *BMC genomics*, **7**, 306.
1071   120.   Li, W., Manktelow, E., von Kirchbach, J.C., Gog, J.R., Desselberger, U. and Lever, A.M. (2010)
1072   Genomic analysis of codon, sequence and structural conservation with selective biochemical-
1073   structure mapping reveals highly conserved and dynamic structures in rotavirus RNAs with
1074   potential cis-acting functions. *Nucleic acids research*, **38**, 7718-7735.
1075   121.   Cheng, J., Nguyen, T.Y.D., Cygan, K.J., Çelik, M.H., Fairbrother, W.G. and Gagneur, J. (2019)
1076   MMSplice: modular modeling improves the predictions of genetic variant effects on splicing.
1077   *Genome biology*, **20**, 1-15.
1078   122.   Wang, G.-S. and Cooper, T.A. (2007) Splicing in disease: disruption of the splicing code and the
1079   decoding machinery. *Nature Reviews Genetics*, **8**, 749-761.
1080   123.   Supek, F., Miñana, B., Valcárcel, J., Gabaldón, T. and Lehner, B. (2014) Synonymous mutations
1081   frequently act as driver mutations in human cancers. *Cell*, **156**, 1324-1335.
1082   124.   Weile, J. and Roth, F.P. (2018) Multiplexed assays of variant effects contribute to a growing
1083   genotype–phenotype atlas. *Human genetics*, **137**, 665-678.
1084   125.   Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., Grody, W.W., Hegde, M.,
1085   Lyon, E. and Spector, E. (2015) Standards and guidelines for the interpretation of sequence
1086   variants: a joint consensus recommendation of the American College of Medical Genetics and
1087   Genomics and the Association for Molecular Pathology. *Genetics in medicine*, **17**, 405-423.
1088   126.   Pertea, M., Lin, X. and Salzberg, S.L. (2001) GeneSplicer: a new computational method for splice
1089   site prediction. *Nucleic acids research*, **29**, 1185-1190.
1090   127.   Reese, M.G., Eeckman, F.H., Kulp, D. and Haussler, D. (1997) Improved splice site detection in
1091   Genie. *Journal of computational biology*, **4**, 311-323.

1092   128.   Comeron, J.M. and Kreitman, M. (1998) The correlation between synonymous and
1093   nonsynonymous substitutions in Drosophila: mutation, selection or relaxed constraints? *Genetics*,
1094   **150**, 767-775.

1095   129.   Wyckoff, G.J., Malcom, C.M., Vallender, E.J. and Lahn, B.T. (2005) A highly unexpected strong
1096   correlation between fixation probability of nonsynonymous mutations and mutation rate. *Trends in*
1097   *Genetics*, **21**, 381-385.

1098   130.   Li, J., Zhang, Z., Vang, S., Yu, J., Wong, G.K.-S. and Wang, J. (2009) Correlation between Ka/Ks
1099   and Ks is related to substitution model and evolutionary lineage. *Journal of molecular evolution*,
1100   **68**, 414-423.

1101   131.   Hurst, L.D. and Pál, C. (2001) Evidence for purifying selection acting on silent sites in BRCA1.
1102   *TRENDS in Genetics*, **17**, 62-65.

1103   132.   Orban, T.I. and Olah, E. (2001) Purifying selection on silent sites–a constraint from splicing
1104   regulation? *TRENDS in Genetics*, **17**, 252-253.

1105   133.   Parmley, J.L. and Hurst, L.D. (2007) How Common Are Intragene Windows with K A> K S Owing
1106   to Purifying Selection on Synonymous Mutations? *Journal of molecular evolution*, **64**, 646-655.

1107   134.   McVean, G.A. and Vieira, J. (2001) Inferring parameters of mutation, selection and demography
1108   from patterns of synonymous site evolution in Drosophila. *Genetics*, **157**, 245-257.

1109   135.   Nielsen, R., Bauer DuMont, V.L., Hubisz, M.J. and Aquadro, C.F. (2007) Maximum likelihood
1110   estimation of ancestral codon usage bias parameters in Drosophila. *Molecular Biology and*
1111   *Evolution*, **24**, 228-235.

1112   136.   Zhou, T., Gu, W. and Wilke, C.O. (2010) Detecting positive and purifying selection at
1113   synonymous sites in yeast and worm. *Molecular biology and evolution*, **27**, 1912-1922.

1114   137.   Lawrie, D.S., Messer, P.W., Hershberg, R. and Petrov, D.A. (2013) Strong purifying selection at
1115   synonymous sites in D. melanogaster. *PLoS Genet*, **9**, e1003527.

1116   138.   Lu, J. and Wu, C.-I. (2005) Weak selection revealed by the whole-genome comparison of the X
1117   chromosome and autosomes of human and chimpanzee. *Proceedings of the National Academy*
1118   *of Sciences*, **102**, 4063-4067.

1119   139.   Hellmann, I., Zöllner, S., Enard, W., Ebersberger, I., Nickel, B. and Pääbo, S. (2003) Selection on
1120   human genes as revealed by comparisons to chimpanzee cDNA. *Genome research*, **13**, 831-837.

1121   140.   Martin, A.R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B.M. and Daly, M.J. (2019) Clinical use
1122   of current polygenic risk scores may exacerbate health disparities. *Nature genetics*, **51**, 584-591.

1123   141.   Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L.,
1124   Laricchia, K.M., Ganna, A. and Birnbaum, D.P. (2020) The mutational constraint spectrum
1125   quantified from variation in 141,456 humans. *Nature*, **581**, 434-443.

1126   142.   Sherry, S.T., Ward, M.-H., Kholodov, M., Baker, J., Phan, L., Smigielski, E.M. and Sirotkin, K.
1127   (2001) dbSNP: the NCBI database of genetic variation. *Nucleic acids research*, **29**, 308-311.

1128   143.   Bromberg, Y., Kahn, P.C. and Rost, B. (2013) Neutral and weakly nonneutral sequence variants
1129   may define individuality. *Proceedings of the National Academy of Sciences*, **110**, 14255.

1130