3   6',6'-Difluoro-aristeromycin is a potent inhibitor of MERS-coronavirus replication

Natacha S. Ogando[1], Jessika C. Zevenhoven-Dobbe[1], Dnyandev B. Jarhad[2], Sushil Kumar

Tripathi[2], Hyuk Woo Lee[3], Lak Shin Jeong[2], Eric J. Snijder[1#], Clara C. Posthuma[1]*

[1] Molecular Virology Laboratory, Department of Medical Microbiology, Leiden University Medical

Center, Leiden, The Netherlands

[2] Research Institute of Pharmaceutical Sciences, College of Pharmacy, Seoul National University,

Seoul 08826, Korea

[3] Future Medicine Company Ltd., Seongnam, Gyeonggi-do 13449, Korea

#Address correspondence to Eric J. Snijder; E.J.Snijder@lumc.nl

* Present address: The Netherlands Commission on Genetic Modification, Bilthoven, The

Netherlands

Running title: Antiviral activity of DFA against coronaviruses

Abstract: 247 words

Figures: 5

Tables: 2

Abstract

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has highlighted the

lack of treatments to combat infections with human or (potentially) zoonotic CoVs. Thus, it is

critical to develop and evaluate antiviral compounds that either directly target CoV functions or

modulate host functions involved in viral replication. Here, we demonstrate that low-micromolar

concentrations of 6',6'-difluoro-aristeromycin (DFA), an adenosine nucleoside analogue, strongly

inhibit the replication of Middle East respiratory syndrome coronavirus (MERS-CoV) in a cell-

based infection assay. DFA was designed to target S-adenosylhomocysteine (SAH) hydrolase and,

consequently, may affect intracellular levels of the methyl donor S-adenosylmethionine, which is

used by two CoV methyltransferases involved in the capping of the 5' end of the viral mRNAs.

Passaging of wild-type MERS-CoV in the presence of DFA selected a virus population with a ~100-

fold decreased DFA sensitivity, which carried various amino acid substitutions in viral

nonstructural proteins (nsps). Specifically, mutations were present in the RNA polymerase

subunit (nsp12) and in nsp13, the helicase subunit containing a nucleoside triphosphate

hydrolase activity that has been implicated in CoV capping. We hypothesize that DFA directly or

indirectly affects viral cap methylation, either by inhibiting the viral enzymes involved or by

binding to SAH hydrolase. We also evaluated the antiviral activity of DFA against other

betacoronaviruses, but found it to have limited impact on their replication, while being quite

cytotoxic to the Calu-3 cells used for this comparison. Nevertheless, our results justify the further

characterization of DFA derivatives as an inhibitor of MERS-CoV replication.

Importance

Currently, there is a lack of antiviral drugs with proven efficacy against human CoV infections

including the MERS-CoV that is endemic in the Middle East, the pandemic SARS-CoV-2 and

potential future zoonotic CoV. This highlights the importance to investigate new drug targets and

identify compounds that can be used to inhibit CoV replication. In this study, we characterize the

inhibitory effect of DFA on MERS-CoV replication by phenotypic studies, time-of-addition studies,

and the generation and genotyping of a DFA-resistant virus population. Our results revealed that

DFA needs further improvement to reduce its cytotoxic side-effects and potentially enhance its

broad-spectrum activity. Despite this observation, we think that DFA can be used to understand

the function and metabolic interactions of the CoV RNA-synthesizing machinery, or as a starting

point for the design of new compounds of the same class.

Introduction

Previously, the emergence of severe acute respiratory syndrome coronavirus (SARS-CoV; in 2003 in China) and Middle East respiratory syndrome coronavirus (MERS-CoV; in 2012 in Saudi Arabia) highlighted the potential pandemic threat posed by this type of zoonotic pathogens and the need to develop rapid response options to contain them (1-4). Due to the severity of the diseases caused by SARS-CoV and MERS-CoV, and their potential for zoonotic transmission and global spread, both these agents received a priority status from the World Health Organization and other government agencies for the development of prophylactic and therapeutic treatment strategies (5, 6). The current SARS-CoV-2 pandemic (7, 8) and its burden on public health worldwide further emphasize the critical nature of the quest for anti-CoV drugs with high clinical efficacy. Many drug classes currently are under evaluation as inhibitors of CoV replication, including compounds directly targeting viral functions, like viral proteases and the RNA polymerase, and host factor-targeting inhibitors (reviewed in (9-12)).

Coronaviruses are positive-stranded RNA (+RNA) viruses with a single genomic RNA of approximately 30 kb that is replicated in the cytoplasm of infected cells. Following entry, the 5'-capped viral genome is recognized and translated by host ribosomes to yield the replicase polyproteins pp1a and pp1ab (13). Subsequently, these large precursors are processed into 16 individual nonstructural proteins (nsp 1 to 16), which are released following polyprotein cleavage by two or three internal proteases. Together, the nsps form a multi-enzyme complex that ensures the replication of the viral genome and the transcription of a set of subgenomic mRNAs (reviewed in (14, 15)). The enzymatic core of this complex is formed by the nsp12 RNA-dependent RNA polymerase (RdRp) that synthesizes RNA with the help of the auxiliary factors nsp7 and nsp8 (16, 17), the nsp13 helicase that unwinds RNA duplexes (18-20), and several other RNA-processing enzymes residing in nsp12-nsp16 (reviewed in (15, 21, 22)). These also include a 3'-to-5' exoribonuclease (nsp14-ExoN) that is thought to increase replication fidelity by correcting mismatches sustained during RNA synthesis (reviewed in (23-26)). The viral structural and accessory proteins, encoded by smaller open reading frames located in the 3'-proximal part of the genome, are expressed from a set of 5'-capped and 3'-polyadenylated subgenomic mRNAs (reviewed in (15, 22)). Apart from ensuring mRNA recognition during formation of the ribosomal

preinitiation complex, the 5'-terminal cap structure protects the viral mRNAs from degradation

by cellular ribonucleases and prevents detection by the host's intracellular pathogen recognition

receptors, which would trigger innate immune responses (reviewed in (27)).

The CoV capping mechanism is thought to consist of four sequential reactions: (i) an RNA

triphosphatase activity residing in nsp13 removes the γ-phosphate group from the 5'-

triphosphorylated RNA (28, 29); (ii) a guanosine monophosphate (GMP) is transferred to the 5'-

diphosphate terminus by a yet to be confirmed guanylyltransferase (GTase)(30), which was

recently proposed to reside in the N-terminal nucleotidyl transferase (NiRAN) domain of nsp12

(31); (iii) the nsp14 methyltransferase (MTase) methylates the cap's 5'-terminal guanine at the

N7-position, producing the so-called cap-0 structure, ^{7me}GpppN (32); (iv) finally, a cap-1 structure

is formed when nsp16, in complex with its nsp10 co-factor, methylates the ribose 2'-O-position

of the first transcribed nucleotide of each viral RNA, converting ^{7me}GpppN into ^{7me}GpppN$_{2'me}$ (33).

Given the central position of the RNA-synthesizing and capping machinery in the CoV replication

cycle, each single component constitutes a potential target for direct-acting antiviral drug

development.

As in cellular methylation reactions, S-adenosyl-L-methionine (SAM) is the most common methyl

donor used by viral MTases, such as those present in CoV nsp14 and nsp16 (34, 35). Thus, the

identification of compounds that can interfere with viral mRNA capping, by either directly

targeting viral MTases or indirectly affecting the concentrations of essential cellular metabolites,

constitutes a viable strategy to develop broad-spectrum CoV inhibitors. S-Adenosyl-

homocysteine (SAH) is released upon the transfer of the methyl group of SAM to a nucleic acid

substrate by a SAM-dependent MTase. Consequently, accumulation of SAH can interfere with

SAM-dependent MTase function due to product inhibition (36). Inhibitors targeting S-adenosyl-

homocysteine (SAH) hydrolase have been reported as potential broad-spectrum antiviral drugs

in different studies (37-39). This hydrolase catalyzes the reversible conversion of SAH into

adenosine and L-homocysteine, which both are then further metabolized for use in different

cellular pathways (40, 41).

Recently, using cell-based assays for MERS-CoV, SARS-CoV, chikungunya and Zika virus

replication, we described the inhibitory potential of a set of adenosine and selenoadenosine

analogues (38). These compounds were derived from aristeromycin, a well-known carbocyclic

nucleoside compound that inhibits SAH hydrolase and exhibits anti-viral, anti-cancer and anti-

toxoplasma activities (reviewed in (42)). These aristeromycin derivatives are nucleoside

analogues designed to directly target viral RdRp activity and/or indirectly target the methylation

of viral RNA by inhibiting the host SAH hydrolase (38). From this library, we identified 6',6'-

difluoro-aristeromycin (DFA) as the aristeromycin derivative that inhibited MERS-CoV replication

most efficiently in cell-based assays (38). In different cell lines, DFA inhibited MERS-CoV

replication at low-micromolar concentrations and could potently reduce the progeny titers

produced by MERS-CoV. Evaluation of the potential of DFA as a broad-spectrum antiviral

compound revealed limited inhibition of the replication of different betacoronaviruses at non-

cytotoxic concentrations. This suggests that DFA-based derivatives need to be developed to

improve the antiviral activity of this compound class and reduce the cytotoxic side- effects.

Results

<u>DFA inhibits MERS-CoV replication at low-micromolar concentrations in different cell lines</u>

DFA was part of a library of more than 80 adenosine and selenoadenosine analogues that was previously evaluated for its antiviral activity against MERS-CoV, SARS-CoV and mouse hepatitis virus (MHV) using cell-based cytopathic effect (CPE) reduction assays. From this analysis, DFA was identified as the most potent inhibitor of MERS-CoV and SARS-CoV replication, with EC_{50} values (half-maximum effective concentration) of 0.2 µM and 0.5 µM, respectively. The compound was found to be more effective in reducing the progeny titers of MERS-CoV than those of SARS-CoV, yielding reductions of more than 3 $\log_{10}$ and 1 $\log_{10}$, respectively, when treating Vero cells with 1.2 µM of DFA (38). Here, we evaluated the antiviral activity of DFA against MERS-CoV in more detail using two independent cell-based assays: a CPE-reduction assay and a dose response assay, using previously described protocols (43, 44). Both assays were performed in different cell lines of human (Huh7 and MRC-5) and non-human origin (Vero).

Remdesivir (RDV) and chloroquine (CHO) were included as positive controls for inhibition of viral replication. The mean EC_{50} values in Vero cells for RDV and CHO were 0.4 µM and 25 µM, respectively, similar to what was described previously (45, 46). Using CPE reduction assays, EC_{50} values in the low-micromolar range were measured for DFA in each of the three cell lines: 0.2 µM in Vero cells, 5.2 µM in Huh7 cells, and 2.3 µM in MRC-5 cells (Fig. 1A-C). In cytotoxicity control studies, the corresponding CC_{50} values (the compound concentration resulting in 50% cytotoxicity) were calculated to be 3.6 µM in Vero, 64 µM in Huh7, and >100 µM in MRC-5 cells (Fig 1A-C). Differences between cell lines in sensitivity (cytotoxicity) to DFA treatment, as observed here, may reflect variation in SAH hydrolase expression (a target of DFA) or uptake and metabolization of the compound.

In order to analyze the inhibitory effect of DFA on MERS-CoV progeny production in more detail, a multiple-cycle dose response assay was performed. A dose-dependent reduction of viral progeny was observed, with a 4 to 5 $\log_{10}$ decrease following treatment with >1.2 µM of DFA in Vero cells, >2.4 µM in Huh7, and >12.5 µM in MRC-5, respectively (Fig. 1D-F). Similar or lower EC_{50} values than in the CPE reduction assay were calculated from these studies: 0.2 µM in Vero

cells, ~0.8 µM in Huh7 cells, and 1.4 µM in MRC-5 cells. These results indicated that DFA exhibits

a similar antiviral activity across multiple cell lines resulting in a consistent ~3.5 to 4-$\log_{10}$

reduction of MERS-CoV progeny titers.

Having established the strong inhibition of MERS-CoV replication by DFA, we also tested its

monophosphoramidate pro-drug (pDFA; Fig. 2A) in a CPE-reduction assay. This compound was

synthesized in order to circumvent the rate-limiting first phosphorylation step that presumably

restricts the efficient metabolization of nucleoside analogues like DFA following their uptake by

the cell (reviewed in (47)). Unfortunately, in this case the pro-drug was less active than DFA itself,

independent of the cell line used (Fig. 2B). Although the chemical and structural modifications of

the prodrug decreased its cytotoxicity, the calculated EC_{50} values, 9 µM in Vero cells and 36 µM

in MRC-5 cells, were more than 10 times higher than the ones measured for DFA (Fig. 1).

Therefore, pDFA was not included in subsequent experiments.

<u>DFA inhibits the early stage of MERS-CoV replication</u>

To characterize the mechanism of action of DFA in more detail, a time-of-addition assay was

performed to determine which stage of the viral replication cycle was inhibited by the compound.

For this purpose, Vero or MRC-5 cells were infected with MERS-CoV at high MOI (3 PFU/cell) and

treated with DFA at different time points pre and post infection at a concentration equaling 4

times the EC_{50}. We observed inhibition of replication when the compound was administered

before infection and at time points up to 4 h p.i. in Vero (Fig. 3A) and 8 h p.i. in MRC-5 (Fig. 3B).

In Vero cells, a 2 $\log_{10}$ reduction of progeny virus titers was observed when the compound was

administered between 24 h before infection and 1 h p.i. In MRC-5 cells, DFA treatment led to a

larger decrease of viral progeny production, >3 $\log_{10}$, when treatment was started up to 4 h p.i.

Using this DFA dose, no cytotoxicity was detected in either cell line (data not shown). Replication

kinetics of MERS-CoV is similar in Vero and MRC-5 cells (48, 49). Thus, the different levels of

progeny titer reduction observed between the two cell lines may be explained by variation in

uptake or metabolic conversion of the compound (50). In any case, these results demonstrated

that DFA inhibits an early stage of MERS-CoV replication.

Selection of MERS-CoV mutants with 100-fold increased DFA resistance

In order to explore the mode of action of DFA, we selected for compound-resistant MERS-CoV
mutants. For this purpose, wild-type MERS-CoV (wtP0) was serially passaged 10 times in MRC-5
cells in the presence of increasing DFA concentrations (from 2.5 µM up to 45 µM). Development
of CPE and drug resistance were monitored microscopically, and plaque phenotype and viral
progeny production were evaluated by plaque assay after each passage. From passage 8 (P8)
onwards, two of the three independently generated lineages showed no increased CPE compared
to uninfected, DFA-treated control cells, meaning that these virus populations could not replicate
in the presence of DFA concentrations above 35 µM, which were used in these later passages.
When P8 virus from lineages 1 or 2 was tested in a CPE-reduction assay, no increased DFA
resistance was noticed compared to an untreated wt virus control (data not shown). In contrast,
lineage 3 (L3) virus did show clear signs of developing DFA resistance. After 10 passages, infection
of cells with L3P10 virus in the presence of 45 µM DFA led to full CPE, which developed equally
fast as for the untreated wt control. When L3P10 virus was tested in a dose response assay, only
a small (<0.5 $\log_{10}$) effect of DFA treatment on viral progeny production was observed in the
presence of up to 100 µM of DFA (Fig. 4A). When compared to untreated wt (wtP10) or parental
virus (wtP0), L3P10 virus displayed a more than 100-fold increased drug resistance, with an EC_{50}
value >100 µM against 0.8 µM for wtP10 and 0.4 µM for wtP0.

In order to assess if the increased DFA resistance of L3P10 affected its replication kinetics in
comparison to the wt control, multi-cycle infections of MRC-5 cells were performed. The two
viruses showed similar growth kinetics (Fig. 4B) with peak titers of 6.1×10^6 PFU/ml (wt) and 7.5
$\times 10^6$ PFU/ml (L3P10) at 48 h p.i. Taken together, the replication kinetics and strongly increased
DFA resistance suggested that, during serial passaging in the presence of DFA, the L3P10 virus
population had acquired mutations that account for a strongly increased resistance to the
compound.

Mutations in the L3P10 virus population implicate DFA in the inhibition of viral capping

In order to identify mutations that contribute to DFA resistance, we sequenced the wtP10 and

L3P10 virus populations by Illumina next-generation sequencing. Subsequently, sequencing reads

were mapped to the reference sequence of MERS-CoV strain EMC/2012 (NC_019843.3; (3)).

Sequence variants constituting less than 10% of the total population of viral reads were excluded

from further analysis. Compared to the original viral sequence, a total of 14 changes were

identified: five synonymous and nine non-synonymous mutations distributed across genes

encoding nine different viral proteins. Two of the identified non-synonymous mutations were

present both in wtP10 and L3P10, suggesting they were cell culture adaptations acquired during

repeated passaging. These mutations, G12033-to-A and C21068-to-U, resulted in D73N and L56F

substitutions in nsp7 and nsp16, respectively. Likewise, the five translationally silent mutations

in the L3P10 population were considered unlikely to be relevant for its phenotypic profile. Of the

remaining seven (Table 1) non-synonymous mutations, one mapped to the accessory protein

encoded by ORF5, which is not essential for viral replication in cell culture (51)-(52), and one to

the Spike protein (53), which also is an unlikely target for inhibition by nucleoside analogues. This

left five L3P10-specific mutations leading to amino acid substitutions in the viral replicase

subunits nsp1, 3, 12, and 13 that may be associated with DFA resistance. As shown in Table 1, all

of these were present in only part of the viral population (in 37% to 55% of the total reads),

suggesting a complex pattern of virus evolution with DFA resistance possibly relying on (different)

combinations of mutations (see Discussion). The short NGS read-length (150 nucleotides) did not

allow us to determine which mutations were combined in the same genome. Since further

optimization of this compound class is needed to improve its selectivity index (Table 2), we did

not perform follow-up experiments to elucidate its mode of action at this stage.

<u>Evaluation of DFA potential as a pan-coronaviral inhibitor</u>

To explore the potential of DFA as a broad-spectrum antiviral, Calu-3 cells, human lung cells that

supports MERS-CoV, SARS-CoV and SARS-CoV-2 replication (54, 55), were treated with increasing

concentrations of DFA and infected with each of these viruses in a dose response assay. By using

the same cell line for all three CoVs, differences in DFA up-take or metabolic conversion to its

triphosphate form were eliminated. The results showed a dose-dependent decrease in the

production of viral progeny for MERS-CoV (Fig. 5A) and SARS-CoV-2 (Fig. 5C) that followed the

cytotoxicity of the compound. At a DFA concentration of 3.2 μM, only a small reduction of MERS-

CoV and SARS-CoV-2 progeny was observed, 0.5 to 1log$_{10}$. Surprisingly, the antiviral activity of

DFA against MERS-CoV in Calu-3 cells was severely reduced when compared to results obtained

in other cell lines, including another human lung cell line MRC-5 (Fig. 1D-F). In the case of SARS-

CoV infection, a minor inhibitory effect was observed at concentrations that appeared to be

somewhat cytotoxic (Fig. 5B and 5D), contrary to what was demonstrated in Vero E6 cells ((38)

and Table 2). Unfortunately, in Calu-3 cells cytotoxicity was detected at low compound

concentrations (>6.2 μM) and the inhibitory effects observed could thus be associated with an

overall decrease in relative cell viability. This indicates that the design of improved DFA

derivatives is needed to decrease cytotoxicity and improve inhibitory potency.

Discussion

This study describes that treatment with low-micromolar DFA concentrations exhibits a strong antiviral effect on MERS-CoV replication in cell culture-based infection models (Fig. 1). Time-of-addition assays indicated that DFA reduced MERS-CoV progeny production when cells were treated prior to, at the time of, or within 4 h after infection (Fig. 3), suggesting that DFA interferes with the early stage of replication. Propagation of MERS-CoV in the presence of DFA led to the selection of a virus population with strongly enhanced resistance to this compound (Fig. 4). Subsequent sequence analysis revealed a potentially complex pattern of resistance evolution, exhibiting multiple mutations that are present in only part of the virus population, including several that map to enzymes involved in viral RNA synthesis and mRNA capping (Table 1).

DFA was originally designed to target the host SAH hydrolase directly and was demonstrated to inhibit this enzyme *in vitro* with an IC_{50} (50% inhibitory concentration) of 1.06 μM (38). The compound is a carbocyclic adenosine analogue based on the parental inhibitor aristeromycin (56, 57), which was further modified by incorporation of a difluorine group at the 6' (top) position of its sugar ring ((38) and Fig. 2A). This modification improved the binding affinity of the compound for human SAH hydrolase and, consequently, the inhibition of its enzymatic activity. Previous studies demonstrated that treatment of cells with high-affinity SAH hydrolase inhibitors, such as neplanocin A and aristeromycin, increases the intracellular SAH concentration, preventing the metabolic conversion of SAH to adenosine and L-homocysteine (reviewed in (58)). Therefore, SAH hydrolase inhibitors reduce or deplete the intracellular pools of homocysteine and adenosine, the latter being produced exclusively by SAH hydrolysis. As the SAM methyl donor is formed via homocysteine trans-sulfuration or the adenosine kinase pathway, SAH hydrolase regulates the intracellular SAM levels and consequently the cell's SAM-dependent methylation reactions. Moreover, SAH accumulation can also reduce the activity of SAM-dependent methyltransferases by feed-back inhibition, as SAH can bind to their active site with higher affinity than SAM itself (36).

A correlation between the antiviral effect of adenosine analogues and their ability to interfere with viral capping has been demonstrated in previous studies with chikungunya virus, dengue virus, West Nile virus and vaccinia virus (59-62). Both CoV methyltransferases use SAM as a

methyl donor for their enzymatic activity (63-65). Thus, SAH hydrolase inhibition and reduced

SAM concentrations may impact, if not block, their activity. Previous studies with 5'-β-

fluoroadenosine and derivatives of aristeromycin demonstrated that inhibition of SAH hydrolase

affects viral replication by reducing RNA methylation (reviewed in (58)). Biochemical analysis of

the MERS-CoV nsp16/nsp10 complex showed the capacity of nsp16 to bind SAH with greater

affinity than SAM (66), whereas superimposition of the SARS-CoV nsp16/nsp10 in complex with

SAH demonstrated that the same binding site is used by both substrates (67). In addition,

increased SAH concentrations reduced the 2'-O-methylation of N7-methylated substrates (65,

66).

In light of the above, direct or indirect inhibition of the CoV capping pathway is a possible mode

of action (MoA) of DFA, although the genotypic profile of the L3P10 virus population (Table 1)

suggests that the compound may inhibit CoV replication using multiple mechanisms. As DFA is a

nucleoside analog, mutations in viral enzymes involved in RNA synthesis and capping may

contribute to the observed DFA-resistance of the L3P10 virus population. Consequently,

mutations Y218F in the nsp12-NiRAN domain and R22K and R161H in the nsp13 ZBD-helicase

subunit attract more attention than nsp1-D172H and nsp3-R1314C. Recently, the nsp12 NiRAN

domain was proposed to function as the capping GTase (31), while also nsp13 has been

implicated in the CoV capping pathway (see Introduction; reviewed in (30)). As the identified

mutations have not been characterized in structural or biochemical studies, one can only

speculate about their potential role in viral replication and DFA resistance

Further phenotypic and mechanistic studies will be needed to better understand the mode of

action of DFA. Additionally, cloning of L3P10 viruses by plaque picking could help to define the

combination(s) of mutations that are the basis for DFA resistance, by evaluating their frequency

of occurrence and associated replication and plaque phenotype.

As a nucleoside analogue, DFA was also considered to be a potential RdRp inhibitor. This would

require uptake by the cell's nucleoside transporters, and subsequent phosphorylation into a

triphosphorylated product that could be incorporated into the RNA chain during viral RNA

synthesis (reviewed on (47)). In order to improve absorption of the compound by the cells and

metabolism into its active form, a prodrug of DFA was synthesized and its antiviral activity was

evaluated. In theory, the monophosphoramidate mask would promote the second
phosphorylation to occur once the compound enters the cytoplasm by circumventing the rate-
limiting step of the first phosphorylation. However, when compared to DFA, the EC_{50} of the
prodrug was more than 10 times higher (Fig. 2), in contrast to results obtained with prodrugs of
other nucleoside analogues (46, 68). In previous work, structure-activity studies and tests of
several purine and pyrimidine analogues of DFA suggested that DFA is most likely not targeting
the RdRp (38, 61, 69). This notion is also supported by the fact that the genotypic profile obtained
for L3P10 did not reveal mutations in the RdRp domain of nsp12.

In this study, we demonstrate that DFA can inhibit the replication of MERS-CoV, but that the
design and development of DFA-based derivatives will be required to reduce cytotoxic side
effects. Combining our results in this study with our previous report (38), showing that DFA can
inhibit chikungunya and Zika virus, DFA appears to be an interesting compound for further
development as a broad-spectrum antiviral agent.

Materials and Methods

<u>Cell culture and viruses</u>

Vero cells were a kind gift from the Department of Viroscience, Erasmus Medical Center, Rotterdam, the Netherlands, and Huh7 cells were provided by Dr. Ralf Bartenschlager, Heidelberg University, Germany. Vero, Vero E6, Huh7, MRC-5 and Calu-3 were cultured as described before (48, 49, 70-72). All cells were incubated at 37°C with 5% CO_2. Infections were carried out in Eagle's minimum essential medium (EMEM; Lonza) containing 25 mM HEPES (Lonza), 2% fetal calf serum (FCS; Bodinco), 100 units/ml penicillin (Lonza), 100 units/ml streptomycin (Lonza) and 2 mM L-glutamine (PAA) (abbreviated from now on as EMEM-2%FCS). MERS-CoV (strain EMC/2012; (3, 4)), SARS-CoV (Frankfurt-1 strain,(73)) and SARS-CoV-2/Leiden-0002 (GenBank accession nr. MT510999; (72)) were used for infections with wild type virus. CoV infections were performed inside biosafety cabinets in a certified biosafety level 3 (BSL3) facilities at Leiden University Medical Center.

<u>Compounds</u>

6',6'-Difluoro-aristeromycin (DFA) and its adenine phosphoramidate pro-drug (pDFA) were designed and synthesized, designated as 2c and 3a, respectively, as described in a previous report (38). Different batches of powder were dissolved in DMSO to a final concentration of 20 mM and single use aliquots were stored at 4°C. Remdesivir (RDV; HY-104077) was purchased from MedChemexpress and chloroquine (C6628) from Sigma. Both compounds were dissolved in adequate solvents (DMSO or PBS, respectively) and single use aliquots were stored at -20°C.

<u>Cytopathic effect (CPE) reduction assay</u>

Cells were seeded in 96-well flat bottom plates in 100 µl at a density of 10000 cells/well of Huh7, 15000 cells/well of MRC-5 or 20000 cells/well of Vero cells. After overnight culture at 37°C, cells were pre-incubated for 30 min with 50 µl of two-fold serial dilutions of compounds prepared in EMEM-2%FCS. Subsequently, half of the wells were infected with MERS-CoV at low MOI in a total

volume of 150 µl of medium with increasing concentrations between 0.05 to 100 uM of

compound, to evaluate the inhibitory effect of compound. The other half of the wells were

"mock"-infected with medium to monitor the (potential) cytotoxicity of the compound. Plates

were incubated for three days (or as mentioned) at 37°C, after which cell viability was measured

using the colorimetric CellTiter 96® Aqueous Non-Radioactive Cell Proliferation kit (Promega).

The absorption at 495 nm was measured using a monochromatic filter in a multimode plate

reader (Envision; Perkin Elmer). Data were normalized to the "mock"-infected control, after

which EC_{50} and CC_{50} values were calculated using non-linear regression with Graph-Pad Prism

V8.0. Each experiment was performed at least in quadruplicate and repeated at least twice.

<u>Dose response assay</u>

To evaluate the effect of compound treatment on viral progeny titers, confluent monolayers of

Vero, Huh7 or MRC-5 were seeded in 24-well plates. Cells were incubated for 30 min at 37°C with

solvent or a range of DFA concentrations (from 0.1 to 100 uM). Then, cells were infected with

MERS-CoV at an MOI of 0.01 for 1 h. After infection, cells were washed three times with PBS and

1 ml of medium with compound at corresponding concentration was added. Supernatants were

collected at 48 h p.i. and viral progeny titers were determined by plaque assay in Vero cells as

described before (74).

In 96-well clusters, Calu-3 cells were seeded at a density of 3×10^4 cells per well in 100 µl culture

medium. Two days later, cells were pre-incubated for 30 min with 2-fold serial dilutions of

compound, starting at a concentration of 25 µM. Subsequently, cells were infected with MERS-

CoV, SARS-CoV or SARS-CoV-2 (MOI of 1 based on titer determined on Vero cells) in the presence

of compound for 1 h. Next, cells were washed three times with PBS and 100 µl of compound

solution in EMEM-2%FCS was added. Supernatants were collected at 24 h p.i. and progeny virus

titers were determined by plaque assay.

<u>Time of addition assay</u>

Confluent monolayers of MRC-5 or Vero cells were seeded in 12-well plates in 1 ml/well of the

appropriate medium (see above), and were grown overnight at 37°C. Treatment of cells (before,

during or after infection) was performed using 0.6 µM (for Vero) and 12.5 µM (for MRC-5) of

compound solution freshly prepared in EMEM-2%FCS medium. Cells were infected with MERS-

CoV inoculum (MOI of 5) for 1h and washed three times with PBS. Subsequently, EMEM-2%FCS

medium was added to the cells and supplemented with compound solution in 2-h intervals to a

final concentration as mentioned above. Supernatants were collected 16 h p.i. and viral titers

were determined by plaque assay.

<u>Resistance culturing and next-generation sequencing (NGS)</u>

Recombinant wt MERS-CoV strain EMC/2012(rMERS-CoV) was passaged in triplicate in presence

of increasing concentrations of DFA ranging from 3.2 µM to 45 µM. Infections were performed

at an MOI of 0.05 in every passage in MRC-5 monolayers. In parallel, rMERS-CoV wt was passaged

in the same conditions in the absence of compound, to identify possible mutations associated

with cell culture adaptation. Additionally, a "mock"-infected well treated with the same

concentration of compound in each passage was evaluated for cytotoxicity by light microcopy.

Supernatants were harvested when 80% to full CPE was observed (usually at 3 d p.i.). Three

lineages were generated by serial passaging, but only lineage 3 was used for next-generation

sequencing. To this end, RNA was isolated from 200 µl of virus-containing cell culture

supernatants using TriPure isolation reagent (Roche Applied Science) and purified according to

manufacturer's instructions. The RNA concentration was measured using a Qubit fluorometer

and RNA High Sensitivity kit (Thermo Fisher Scientific). NGS sample preparation and analysis were

performed as described previously (75). After filtration and trimming of data, the remaining reads

were mapped to the MERS-CoV GenBank reference sequence (NC_019843;(3, 4)). Raw NGS data

sets for wtP10 and L3P10 samples analysed in this study were deposited in NCBI Bioproject and

are available under the following link: http://www.ncbi.nlm.nih.gov/bioproject/730836. Only

MERS-CoV-specific reads were included in these data files.

Acknowledgments

N.S.O. was supported by the Marie Skłodowska-Curie ETN European Training Network 'ANTIVIRALS' (EU Grant Agreement No. 642434). We thank Adriaan de Wilde and Martijn van Hemert for their support and for helpful discussions. We thank Igor Sidorov for his technical support in the analysis of NGS data.

409   **References**

410   1.   Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, Tong S, Urbani C,
411   Comer JA, Lim W, Rollin PE, Dowell SF, Ling AE, Humphrey CD, Shieh WJ, Guarner
412   J, Paddock CD, Rota P, Fields B, DeRisi J, Yang JY, Cox N, Hughes JM, LeDuc JW,
413   Bellini WJ, Anderson LJ, Group SW. 2003. A novel coronavirus associated with severe
414   acute respiratory syndrome. N Engl J Med 348:1953-66.
415   2.   Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, Nicholls J, Yee WK, Yan WW,
416   Cheung MT, Cheng VC, Chan KH, Tsang DN, Yung RW, Ng TK, Yuen KY, group Ss.
417   2003. Coronavirus as a possible cause of severe acute respiratory syndrome. Lancet
418   361:1319-25.
419   3.   Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. 2012. Isolation
420   of a novel coronavirus from a man with pneumonia in Saudi Arabia. N Engl J Med
421   367:1814-1820.
422   4.   van Boheemen S, de Graaf M, Lauber C, Bestebroer TM, Raj VS, Zaki AM, Osterhaus
423   AD, Haagmans BL, Gorbalenya AE, Snijder EJ, Fouchier RA. 2012. Genomic
424   characterization of a newly discovered coronavirus associated with acute respiratory
425   distress syndrome in humans. MBio 3:e00473-12.
426   5.   Brende B, Farrar J, Gashumba D, Moedas C, Mundel T, Shiozaki Y, Vardhan H, Wanka
427   J, Rottingen JA. 2017. CEPI-a new global R&D organisation for epidemic preparedness
428   and response. Lancet 389:233-235.
429   6.   Mehand MS, Al-Shorbaji F, Millett P, Murgue B. 2018. The WHO R&D Blueprint: 2018
430   review of emerging infectious diseases requiring urgent research and development
431   efforts. Antiviral Res 159:63-67.
432   7.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL,
433   Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng
434   XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL.
435   2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin.
436   Nature 579:270-273.
437   8.   Munster VJ, Koopmans M, van Doremalen N, van Riel D, de Wit E. 2020. A novel
438   coronavirus emerging in China - key questions for impact assessment. N Engl J Med
439   382:692-694.
440   9.   Zumla A, Chan JF, Azhar EI, Hui DS, Yuen KY. 2016. Coronaviruses - drug discovery
441   and therapeutic options. Nat Rev Drug Discov 15:327-47.
442   10.   Garcia-Serradilla M, Risco C, Pacheco B. 2019. Drug repurposing for new, efficient,
443   broad spectrum antivirals. Virus Res 264:22-31.
444   11.   Dittmar M, Lee JS, Whig K, Segrist E, Li M, Jurado K, Samby K, Ramage H, Schultz D,
445   Cherry S. 2020. Drug repurposing screens reveal FDA approved drugs active against
446   SARS-Cov-2. Cell Rep doi:10.1016/j.celrep.2021.108959:108959.
447   12.   Valle C, Martin B, Touret F, Shannon A, Canard B, Guillemot JC, Coutard B, Decroly E.
448   2020. Drugs against SARS-CoV-2: What do we know about their mode of action? Rev
449   Med Virol 30:1-10.
450   13.   Snijder EJ, Bredenbeek PJ, Dobbe JC, Thiel V, Ziebuhr J, Poon LL, Guan Y, Rozanov
451   M, Spaan WJ, Gorbalenya AE. 2003. Unique and conserved features of genome and
452   proteome of SARS-coronavirus, an early split-off from the coronavirus group 2 lineage. J
453   Mol Biol 331:991-1004.

14. Fehr AR, Perlman S. 2015. Coronaviruses: an overview of their replication and
pathogenesis. Methods Mol Biol 1282:1-23.
15. V'Kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. 2020. Coronavirus biology and
replication: implications for SARS-CoV-2. Nat Rev Microbiol doi:10.1038/s41579-020-
00468-6.
16. Subissi L, Posthuma CC, Collet A, Zevenhoven-Dobbe JC, Gorbalenya AE, Decroly E,
Snijder EJ, Canard B, Imbert I. 2014. One severe acute respiratory syndrome coronavirus
protein complex integrates processive RNA polymerase and exonuclease activities. Proc
Natl Acad Sci U S A 111:E3900-E3909.
17. Kirchdoerfer RN, Ward AB. 2019. Structure of the SARS-CoV nsp12 polymerase bound
to nsp7 and nsp8 co-factors. Nat Commun 10:2342.
18. Seybert A, Hegyi A, Siddell SG, Ziebuhr J. 2000. The human coronavirus 229E
superfamily 1 helicase has RNA and DNA duplex-unwinding activities with 5'-to-3'
polarity. RNA 6:1056-68.
19. Adedeji AO, Singh K, Calcaterra NE, DeDiego ML, Enjuanes L, Weiss S, Sarafianos SG.
2012. Severe acute respiratory syndrome coronavirus replication inhibitor that interferes
with the nucleic acid unwinding of the viral helicase. Antimicrob Agents Chemother
56:4718-28.
20. Chen J, Malone B, Llewellyn E, Grasso M, Shelton PMM, Olinares PDB, Maruthi K,
Eng ET, Vatandaslar H, Chait BT, Kapoor TM, Darst SA, Campbell EA. 2020. Structural
Basis for Helicase-Polymerase Coupling in the SARS-CoV-2 Replication-Transcription
Complex. Cell 182:1560-1573 e13.
21. Posthuma CC, Te Velthuis AJW, Snijder EJ. 2017. Nidovirus RNA polymerases:
Complex enzymes handling exceptional RNA genomes. Virus Res 234:58-73.
22. Snijder EJ, Decroly E, Ziebuhr J. 2016. The Nonstructural Proteins Directing
Coronavirus RNA Synthesis and Processing. Adv Virus Res 96:59-126.
23. Ulferts R, Ziebuhr J. 2011. Nidovirus ribonucleases: Structures and functions in viral
replication. RNA Biol 8:295-304.
24. Denison MR, Graham RL, Donaldson EF, Eckerle LD, Baric RS. 2011. Coronaviruses:
an RNA proofreading machine regulates replication fidelity and diversity. RNA Biol
8:270-9.
25. Ogando NS, Ferron F, Decroly E, Canard B, Posthuma CC, Snijder EJ. 2019. The curious
case of the nidovirus exoribonuclease: its role in RNA synthesis and replication fidelity.
Front Microbiol 10:1813.
26. Robson F, Khan KS, Le TK, Paris C, Demirbag S, Barfuss P, Rocchi P, Ng WL. 2020.
Coronavirus RNA Proofreading: Molecular Basis and Therapeutic Targeting. Mol Cell
79:710-727.
27. Kindler E, Thiel V. 2014. To sense or not to sense viral RNA--essentials of coronavirus
innate immune evasion. Curr Opin Microbiol 20:69-75.
28. Ivanov KA, Thiel V, Dobbe JC, van der Meer Y, Snijder EJ, Ziebuhr J. 2004. Multiple
enzymatic activities associated with severe acute respiratory syndrome coronavirus
helicase. J Virol 78:5619-32.
29. Ivanov KA, Ziebuhr J. 2004. Human coronavirus 229E nonstructural protein 13:
characterization of duplex-unwinding, nucleoside triphosphatase, and RNA 5'-
triphosphatase activities. J Virol 78:7833-8.
30. Decroly E, Ferron F, Lescar J, Canard B. 2011. Conventional and unconventional
mechanisms for capping viral mRNA. Nat Rev Microbiol 10:51-65.

31. Yan L, Ge J, Zheng L, Zhang Y, Gao Y, Wang T, Huang Y, Yang Y, Gao S, Li M, Liu Z,
Wang H, Li Y, Chen Y, Guddat LW, Wang Q, Rao Z, Lou Z. 2021. Cryo-EM Structure
of an Extended SARS-CoV-2 Replication and Transcription Complex Reveals an
Intermediate State in Cap Synthesis. Cell 184:184-193 e10.
32. Chen Y, Cai H, Pan J, Xiang N, Tien P, Ahola T, Guo D. 2009. Functional screen reveals
SARS coronavirus nonstructural protein nsp14 as a novel cap N7 methyltransferase. Proc
Natl Acad Sci U S A 106:3484-3489.
33. Decroly E, Imbert I, Coutard B, Bouvet M, Selisko B, Alvarez K, Gorbalenya AE,
Snijder EJ, Canard B. 2008. Coronavirus nonstructural protein 16 is a cap-0 binding
enzyme possessing (nucleoside-2'O)-methyltransferase activity. J Virol 82:8071-84.
34. Lu SC. 2000. S-Adenosylmethionine. Int J Biochem Cell Biol 32:391-5.
35. Struck AW, Thompson ML, Wong LS, Micklefield J. 2012. S-adenosyl-methionine-
dependent methyltransferases: highly versatile enzymes in biocatalysis, biosynthesis and
other biotechnological applications. Chembiochem 13:2642-55.
36. Kloor D, Osswald H. 2004. S-Adenosylhomocysteine hydrolase as a target for
intracellular adenosine action. Trends Pharmacol Sci 25:294-7.
37. De Clercq E. 2002. Strategies in the design of antiviral drugs. Nat Rev Drug Discov 1:13-
25.
38. Yoon JS, Kim G, Jarhad DB, Kim HR, Shin YS, Qu S, Sahu PK, Kim HO, Lee HW,
Wang SB, Kong YJ, Chang TS, Ogando NS, Kovacikova K, Snijder EJ, Posthuma CC,
van Hemert MJ, Jeong LS. 2019. Design, Synthesis, and Anti-RNA Virus Activity of 6'-
Fluorinated-Aristeromycin Analogues. J Med Chem doi:10.1021/acs.jmedchem.9b00781.
39. Chen Q, Schneller SW, Liu C, Jones KL, Singer T. 2020. 5'-Nor-3-Deaza-1',6'-
Isoneplanocin, the Synthesis and Antiviral Study. Molecules 25.
40. Cantoni GL. 1975. Biological methylation: selected aspects. Annu Rev Biochem 44:435-
51.
41. Palmer JL, Abeles RH. 1979. The mechanism of action of S-adenosylhomocysteinase. J
Biol Chem 254:1217-26.
42. Rawal RK, Bariwal J, Singh V. 2016. Chemistry and Bioactivities of Aristeromycins: An
Overview. Curr Top Med Chem 16:3258-3273.
43. de Wilde AH, Raj VS, Oudshoorn D, Bestebroer TM, van Nieuwkoop S, Limpens RW,
Posthuma CC, van der Meer Y, Barcena M, Haagmans BL, Snijder EJ, van den Hoogen
BG. 2013. MERS-coronavirus replication induces severe in vitro cytopathology and is
strongly inhibited by cyclosporin A or interferon-alpha treatment. J Gen Virol 94:1749-
60.
44. de Wilde AH, Falzarano D, Zevenhoven-Dobbe JC, Beugeling C, Fett C, Martellaro C,
Posthuma CC, Feldmann H, Perlman S, Snijder EJ. 2017. Alisporivir inhibits MERS- and
SARS-coronavirus replication in cell culture, but not SARS-coronavirus infection in a
mouse model. Virus Res 228:7-13.
45. de Wilde AH, Jochmans D, Posthuma CC, Zevenhoven-Dobbe JC, van Nieuwkoop S,
Bestebroer TM, van den Hoogen BG, Neyts J, Snijder EJ. 2014. Screening of an FDA-
approved compound library identifies four small-molecule inhibitors of Middle East
respiratory syndrome coronavirus replication in cell culture. Antimicrob Agents
Chemother 58:4875-84.
46. Agostini ML, Andres EL, Sims AC, Graham RL, Sheahan TP, Lu X, Smith EC, Case JB,
Feng JY, Jordan R, Ray AS, Cihlar T, Siegel D, Mackman RL, Clarke MO, Baric RS,

Denison MR. 2018. Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is
Mediated by the Viral Polymerase and the Proofreading Exoribonuclease. MBio 9.

47.   Jordheim LP, Durantel D, Zoulim F, Dumontet C. 2013. Advances in the development of
nucleoside and nucleotide analogues for cancer and viral diseases. Nat Rev Drug Discov
12:447-64.

48.   de Wilde AH, Raj VS, Oudshoorn D, Bestebroer TM, van Nieuwkoop S, Limpens R,
Posthuma CC, van der Meer Y, Barcena M, Haagmans BL, Snijder EJ, van den Hoogen
BG. 2013. MERS-coronavirus replication induces severe in vitro cytopathology and is
strongly inhibited by cyclosporin A or interferon-alpha treatment. J Gen Virol 94:1749-
1760.

49.   Knaap RCM, Fernández-Delgado R, Dalebout TJ, Oreshkova N, Bredenbeek PJ,
Enjuanes L, Sola I, Snijder EJ, Kikkert M. 2019. The deubiquitinating activity of Middle
East respiratory syndrome coronavirus papain-like protease delays the innate immune
response and enhances virulence in a mouse model. bioRxiv doi:10.1101/751578:751578.

50.   El-Awady R, Saleh E, Hashim A, Soliman N, Dallah A, Elrasheed A, Elakraa G. 2016.
The Role of Eukaryotic and Prokaryotic ABC Transporter Family in Failure of
Chemotherapy. Front Pharmacol 7:535.

51.   Menachery VD, Mitchell HD, Cockrell AS, Gralinski LE, Yount BL, Jr., Graham RL,
McAnarney ET, Douglas MG, Scobey T, Beall A, Dinnon K, 3rd, Kocher JF, Hale AE,
Stratton KG, Waters KM, Baric RS. 2017. MERS-CoV Accessory ORFs Play Key Role
for Infection and Pathogenesis. mBio 8.

52.   Liu DX, Fung TS, Chong KK, Shukla A, Hilgenfeld R. 2014. Accessory proteins of
SARS-CoV and other coronaviruses. Antiviral Res 109:97-109.

53.   Mou H, Raj VS, van Kuppeveld FJ, Rottier PJ, Haagmans BL, Bosch BJ. 2013. The
receptor binding domain of the new Middle East respiratory syndrome coronavirus maps
to a 231-residue region in the spike protein that efficiently elicits neutralizing antibodies.
J Virol 87:9379-83.

54.   Hoffmann M, Mosbauer K, Hofmann-Winkler H, Kaul A, Kleine-Weber H, Kruger N,
Gassen NC, Muller MA, Drosten C, Pohlmann S. 2020. Chloroquine does not inhibit
infection of human lung cells with SARS-CoV-2. Nature 585:588-590.

55.   Lau SK, Lau CC, Chan KH, Li CP, Chen H, Jin DY, Chan JF, Woo PC, Yuen KY. 2013.
Delayed induction of proinflammatory cytokines and suppression of innate antiviral
response by the novel Middle East respiratory syndrome coronavirus: implications for
pathogenesis and treatment. J Gen Virol 94:2679-90.

56.   Kusaka T, Yamamoto H, Shibata M, Muroi M, Kishi T. 1968. Streptomyces citricolor
nov. sp. and a new antibiotic, aristeromycin. J Antibiot (Tokyo) 21:255-63.

57.   Wolfe MS, Lee Y, Bartlett WJ, Borcherding DR, Borchardt RT. 1992. 4'-modified
analogues of aristeromycin and neplanocin A: synthesis and inhibitory activity toward S-
adenosyl-L-homocysteine hydrolase. J Med Chem 35:1782-91.

58.   WANG JR, R. K.; CHU, C. K. 2011. Recent Advances in Carbocyclic Nucleosides:
Synthesis and Biological Activity. Med Chem Nucleic Acids
doi:10.1002/chin.201204251:1-100.

59.   Hasobe M, McKee JG, Borchardt RT. 1989. Relationship between intracellular
concentration of S-adenosylhomocysteine and inhibition of vaccinia virus replication and
inhibition of murine L-929 cell growth. Antimicrob Agents Chemother 33:828-34.

60. Ransohoff RM, Narayan P, Ayers DF, Rottman FM, Nilsen TW. 1987. Priming of
influenza mRNA transcription is inhibited in CHO cells treated with the methylation
inhibitor, neplanocin A. Antiviral Res 7:317-27.
61. Kovacikova K, Morren BM, Tas A, Albulescu IC, van Rijswijk R, Jarhad DB, Shin YS,
Jang MH, Kim G, Lee HW, Jeong LS, Snijder EJ, van Hemert MJ. 2020. 6'-beta-Fluoro-
Homoaristeromycin and 6'-Fluoro-Homoneplanocin A Are Potent Inhibitors of
Chikungunya Virus Replication through Their Direct Effect on Viral Nonstructural
Protein 1. Antimicrob Agents Chemother 64.
62. Chen H, Zhou B, Brecher M, Banavali N, Jones SA, Li Z, Zhang J, Nag D, Kramer LD,
Ghosh AK, Li H. 2013. S-adenosyl-homocysteine is a weakly bound inhibitor for a
flaviviral methyltransferase. PLoS One 8:e76900.
63. Jin X, Chen Y, Sun Y, Zeng C, Wang Y, Tao J, Wu A, Yu X, Zhang Z, Tian J, Guo D.
2013. Characterization of the guanine-N7 methyltransferase activity of coronavirus nsp14
on nucleotide GTP. Virus Res 176:45-52.
64. Bouvet M, Debarnot C, Imbert I, Selisko B, Snijder EJ, Canard B, Decroly E. 2010. In
vitro reconstitution of SARS-coronavirus mRNA cap methylation. PLoS Pathog
6:e1000863.
65. Aouadi W, Eydoux C, Coutard B, Martin B, Debart F, Vasseur JJ, Contreras JM, Morice
C, Querat G, Jung ML, Canard B, Guillemot JC, Decroly E. 2017. Toward the
identification of viral cap-methyltransferase inhibitors by fluorescence screening assay.
Antiviral Res 144:330-339.
66. Aouadi W, Blanjoie A, Vasseur JJ, Debart F, Canard B, Decroly E. 2017. Binding of the
Methyl Donor S-Adenosyl-l-Methionine to Middle East Respiratory Syndrome
Coronavirus 2'-O-Methyltransferase nsp16 Promotes Recruitment of the Allosteric
Activator nsp10. J Virol 91.
67. Decroly E, Debarnot C, Ferron F, Bouvet M, Coutard B, Imbert I, Gluais L,
Papageorgiou N, Sharff A, Bricogne G, Ortiz-Lombardia M, Lescar J, Canard B. 2011.
Crystal structure and functional analysis of the SARS-coronavirus RNA cap 2'-O-
methyltransferase nsp10/nsp16 complex. PLoS Pathog 7:e1002059.
68. Meier C, Balzarini J. 2006. Application of the cycloSal-prodrug approach for improving
the biological potential of phosphorylated biomolecules. Antiviral Res 71:282-92.
69. Shin YS, Jarhad DB, Jang MH, Kovacikova K, Kim G, Yoon JS, Kim HR, Hyun YE,
Tipnis AS, Chang TS, van Hemert MJ, Jeong LS. 2020. Identification of 6'-beta-fluoro-
homoaristeromycin as a potent inhibitor of chikungunya virus replication. Eur J Med
Chem 187:111956.
70. Ogando NS, Zevenhoven-Dobbe JC, van der Meer Y, Bredenbeek PJ, Posthuma CC,
Snijder EJ. 2020. The enzymatic activity of the nsp14 exoribonuclease is critical for
replication of MERS-CoV and SARS-CoV-2. J Virol doi:10.1128/JVI.01246-20.
71. Nedialkova DD, Gorbalenya AE, Snijder EJ. 2010. Arterivirus Nsp1 modulates the
accumulation of minus-strand templates to control the relative abundance of viral
mRNAs. PLoS Pathog 6:e1000772.
72. Salgado-Benvindo C, Thaler M, Tas A, Ogando NS, Bredenbeek PJ, Ninaber DK, Wang
Y, Hiemstra PS, Snijder EJ, van Hemert MJ. 2020. Suramin Inhibits SARS-CoV-2
Infection in Cell Culture by Interfering with Early Steps of the Replication Cycle.
Antimicrob Agents Chemother 64.
73. Pfefferle S, Krahling V, Ditt V, Grywna K, Muhlberger E, Drosten C. 2009. Reverse
genetic characterization of the natural genomic deletion in SARS-Coronavirus strain

Frankfurt-1 open reading frame 7b reveals an attenuating function of the 7b protein in-
vitro and in-vivo. Virol J 6:131.
74. van den Worm SH, Eriksson KK, Zevenhoven JC, Weber F, Zust R, Kuri T, Dijkman R,
Chang G, Siddell SG, Snijder EJ, Thiel V, Davidson AD. 2012. Reverse genetics of
SARS-related coronavirus using vaccinia virus-based recombination. PLoS One
7:e32857.
75. Ogando NS, Dalebout TJ, Zevenhoven-Dobbe JC, Limpens R, van der Meer Y, Caly L,
Druce J, de Vries JJC, Kikkert M, Barcena M, Sidorov I, Snijder EJ. 2020. SARS-
coronavirus-2 replication in Vero E6 cells: replication kinetics, rapid adaptation and
cytopathology. J Gen Virol doi:10.1099/jgv.0.001453.

Figures

Figure 1- DFA inhibits MERS-CoV replication in different cell lines. Vero (A and D), Huh7 (B and E) and MRC-5 (C and F) were treated with a two-fold dilution series of DFA in the low-micromolar range and infected with MERS-CoV. Inhibitory effect was evaluated by a CPE-reduction assay (A-C) or dose response assay (D-F). For the CPE-reduction assay, cell viability was assayed using the CellTiter 96 Aqueous One Solution cell proliferation assay (MTS assay) 3 d p.i.. The graphs show the results of at least two independent experiments (mean ± sd are shown). A non-linear regression analysis was applied. In the dose response assay, cell supernatants were collected after 2 d.p.i and viral progeny was titrated by plaque assay on Vero cells. Error bars represent standard deviation. Statistical significance was determined by one-way ANOVA. *, $p<0.1$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$.

Figure 2- DFA prodrug inhibits MERS-CoV replication. (A) DFA and pDFA schematic structure. pDFA antiviral activity was evaluated by a CPE-reduction assay. (B) Vero or (C) MRC-5 cells were treated with two-fold serial dilution of pDFA and infected with MERS-CoV. After 3 d p.i., cell viability was measured using the CellTiter 96 Aqueous One Solution cell proliferation assay (MTS assay). The graphs show the results of two independent experiments (mean ± sd are shown). A non-linear regression analysis was applied.

653
654

Figure 3- DFA inhibits early steps of MERS-CoV replication. Vero (A) and MRC-5 (B) cells were treated with 0.6 and 12.5 µM, respectively, at the indicated time points pre- and post-infection. Viral progeny in supernatant harvested at 16 h p.i. was determined by plaque assay in Vero cells. The data represent the results from duplicates of 2 independent experiments. Error bars represent standard deviations. Statistical significance was determined by one-way ANOVA.; *, $p<0.1$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$.

Figure 4- Resistant MERS-CoV mutants selected by passaging in the presence of DFA. Replication in MRC-5 cells of a DFA-resistant virus population (L3P10) in the presence of increasing concentrations of DFA, compared to the parental virus (wtP0) and untreated wt virus (wtP10). Cells were infected with MOI 0.01 and virus progeny in supernatant harvested at 48 h p.i. The data represent the results from four replicates obtained in 2 independent experiments. (B) Characterization of growth kinetics of selected resistant mutant (L3P10). MRC-5 cells were infected with MOI 0.01 and supernatants were harvested at indicated time points from triplicate wells. Viral progeny titers were determined by plaque assay in Vero cells (mean ± sd is presented). Statistical significance was determined by one-way ANOVA. *, $p < 0.1$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

657
658

Figure 5- DFA antiviral activity is reduced in Calu-3 cells. Calu-3 cells were infected with MERS-CoV (A), SARS-CoV (B) and SARS-CoV-2 (C) in the presence of various concentrations of DFA. An MOI of 1 was used, based on titrations of virus stocks on Vero cells. Progeny virus titers in supernatants harvested at 24 h p.i. were determined by plaque assay in Vero cells. (D) Cytotoxicity of DFA was measured in mock-infected cells, and was determined at 24 h p.i. in a CPE-reduction assay by use of the CellTiter 96 Aqueous One Solution cell proliferation assay (MTS assay). The data represent triplicates of 2 independent experiments and error bars show standard deviations. Statistical significance was determined by one-way ANOVA. No*, no significance; *, $p < 0.1$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Table 1- Summary of non-synonymous mutations specifically identified in MERS-CoV L3P10 by NGS

Coding region	nt change	aa change	Domain	Presence in L3P10 (NGS)
nsp1	G795C	D172H	CTD	55%
nsp3	C6777T	R1314C	PLnc	37%
nsp12	A14061T	Y218F	NiRAN	48%
nsp13	G16272A	R22K	ZBD	49%
	G16689A	R161H	1B	40%
spike	T24085_24086*ins*ACTCAACAGGTG	P876_V877ins TQQV		37%
ORF5	G26927_26928*ins*T	Not in frame		48%

nt, nucleotide; aa, amino acid; CTD, C-terminal domain; PLnc, papain-like noncanonical domain; NiRAN, Nidovirus RdRp associated nucleotidyl transferase domain; ZBD, Zinc binding domain; 1B, 1B domain of helicase; * also present in wtP10 control

Table 2 – The antiviral effect of DFA on the replication of different betacoronaviruses.

Virus	Cell line	EC_{50} (µM)	CC_{50} (µM)	SI
MERS-CoV	Vero	0.2	>3.2	>16
	HuH7	0.6	>50	>80
	MRC-5	0.8	>50	>60
	Calu-3	~2	>25	>12
SARS-CoV	Calu-3	~5	>25	>5
SARS-CoV-2	Calu-3	~2	>25	>12

EC_{50}s values were calculated based on results obtained in dose response assay, while CC_{50}s values were determined in a cell viability assay as described in materials and methods. SI, selectivity index was calculated by comparing CC_{50} with EC_{50} values.