Significance of Oct-4 transcription factor as a pivotal therapeutic target for CD44$^+$/24$^-$ mammary tumour initiating cells; aiming at the root of the recurrence

Utsav Sen[1], Apoorva Gowda[1], Hari Krishnareddy Rachamalla[2,3], Rajkumar Banerjee[2,3],

Sudheer Shenoy P*[1], Bipasha Bose*[1]

[1]*Stem cells and regenerative medicine centre, Yenepoya Research Centre, Yenepoya (Deemed to be university), Deralakatte, Mangalore-575018, Karnataka.*

[2]*Applied Biology Division, CSIR-Indian Institute of Chemical Technology, Uppal Road, Hyderabad, Telangana-500007, India.*

[3]*Academy of Scientific & Innovative Research (AcSIR), CSIR-HRDC Campus, Ghaziabad, Uttar Pradesh-201002, India.*

***Correspondence**: Bipasha Bose (Bipasha.bose@yenepoya.edu.in, Bipasha.bose@gmail.com) +91 9730889609. Sudheer Shenoy P. (shenoy@yenepoya.edu.in, shenoy2000@yahoo.com)
+91 9108585282.

Address: Stem Cells and Regenerative Medicine Centre, Yenepoya Research Centre, Yenepoya Deemed to be University, University Road, Deralakatte, Mangalore, 575018, Karnataka, India.

Abstract:

Breast cancer (BC) remains one of the deadliest and most frequently diagnosed metastatic cancers worldwide. Cancer stem cells (CSCs) are the cell population within the tumour niche, having an epithelial to mesenchymal (EMT) transition phenotype, high self-renewal, and vigorous metastatic capacity, drug resistance, and ultimate cause of tumour relapse in patients. Identification of targets for apoptosis induction is essential to provide novel therapeutic approaches in breast cancer. Our earlier studies showed that Vitamin C induces apoptotic cell death by losing redox balance in TNBC CSCs. In this current study, we have made an attempt to identify previously unrecognized CSC survival factor/factors that can be used as druggable targets for bCSCs apoptosis regulators isolated from the triple negative breast cancer cell line: MDA MB 468. A thorough literature review was carried out to identify candidates, and Oct-4 was identified as the most promising marker for its unique abundance in cancer and not in normal cells and the contribution of Oct4 to the sustenance of cancer cells. We then validated a very high expression of Oct-4 in the MDA MB 468 bCSCs population using flowcytometry. Loss of function of Oct-4 was carried out using siRNA mediated knockdown in the bCSCs followed by assessing for cellular apoptosis. Our results indicated that Oct-4 knockdown, induced cell death in bCSCs as evident by a change in cellular morphology and positive staining for Annexin-V expression thereby indicating the role of Oct4 in bCSC survival. Moreover, our findings also suggest that Oct-4 directly interacts with Vitamin C as revealed by in silico docking and Oct-4 downregulation inhibited mammosphere formation in vitro. This data, hence, contributes towards novel information about Oct-4 highlighting this molecule as a novel survival factor in bCSCs.

Keywords: *Breast cancer stem cells, POUF1, Oct-4 transcription factor, Ascorbic acid, Cancer stem cells therapeutics, siRNA*

Introduction:

Triple-negative breast cancer (TNBC) is a notable clinical problem to which 30% of all patients often turn up with a recurrence of the disease within 2 to 5 years after completion of treatment (1). Despite extensive progress in BC's diagnosis and treatment, several clinical and scientific problems remain unsolved (2). Procedures for advanced stages of this disease are still relatively limited and inefficient (3). The limitation of these therapies is due to not yet efficiently targeting two essential events that are happening to the breast cancer cells, i.e., epithelial to mesenchymal transition (EMT) and cancer stem cells (CSCs) turnover (4).

Based on pathology and gene expression profiling, triple-negative breast cancer cells (ER$^-$, PR$^-$, HER2$^-$) (TNBCs) are heterogeneous and enriched with CD44$^+$/24$^-$ tumour-initiating cells or cancer stem cells (TICs/CSCs) (5, 6). These pathobiological modifications make TNBC cells aggressive, metastatic, and less sensitive to standard chemotherapy and tumour relapse. These cancer stem cells are interlinked with each other and play crucial roles in BC progression and relapse (7-9). For metastasis, cancer cells modify their cytoskeleton structure, thus initiating invasion and migration of the CSCs (10). The initiation of the migration invasion and metastasis is associated with EMT (11, 12). Where the cells lose the epithelial markers like E-cadherin, EpCAM, and gain mesenchymal markers like N-cadherin, Vimentin, Snail, and Twist (13).

Homeodomain transcription factor of the Pit-Oct-Unc (POU) family, named Octamer-binding transcription factor 4 (Oct-4) is considered to play an essential role in the self-renewal epithelial-mesenchymal transition (EMT) (14) and drug resistance development of CSCs and breast cancer metastasis (7). Oct-4 is well established as one of the most important transcription factors that control the self-renewal of pluripotent stem cells and cancer cells (15). Most interestingly, Oct-4 expression is predominantly observed in embryonic stem cells (ESCs) or some cancer cells (16). Hence, this uniqueness of abundance and the crucial developmental and sustenance role could make Oct-4 one of the critical factors worth targeting in stem cell-specific cancer therapy (17). In our previous study, we have used Vitamin C, and demonstrated the apoptotic effect of Vitamin C on bCSCs (18).

In this study, we have first isolated the breast CSCs from the TNBC cells MDA MB 468,

using flow-sorting based on surface marker CD44$^+$/CD24$^-$. Then the abundance of the Oct-4

was confirmed by using flow cytometric analysis in bCSCs. Loss of function of Oct-4 on

bCSCs survival was established upon Oct 4 knockdown using siRNA. In summary, we have

deciphered the importance of the transcription factor Oct-4 in bCSC survival and sustenance.

Also, we sought to determinize the efficiency of Oct-4 as a potential target for the bCSCs

toward cancer/cancer stem cell therapeutics.

2.0 Material and Methods:

2.1 Cell culture:

The TNBC line MDA MB 468 was procured from the National Centre for Cell Science,

Pune, India. The cancer cells were cultured in media containing DMEM supplemented with

10% fetal bovine serum (Cat. No. RM9955) (all from Hi-Media Laboratories, India), 1%

Penicillin and Streptomycin, GlutaMAX, non-essential amino acids, sodium pyruvate, and

0.1% of 2-mercaptoethanol (all from Gibco Thermo Fisher Scientific, USA). The cells were

grown in incubator with 5% CO_2 with humidified atmosphere. The cells were trypsinized

upon reaching 75–80% confluency using 0.25% trypsin-EDTA (Thermo Fisher Scientific,

USA).

2.2 Immunophenotyping by flow cytometry:

For analytical flow cytometry (immunophenotyping), the bCSCs were fixed with 4% PFA

and permeabilized with 0.3% Triton X-100. Cells were then incubated with the fluorescently

tagged antibody (CD44 and CD24) for 1 hour. Then the cells were given PBS wash and

acquired on guava EasyCyte flow cytometer (10,000 events). Obtained data were analyzed

using DE Novo FCS Express 5 software, USA (antibody details are available in

Supplementary Table-1).

2.3 Migration assay:

Sorted and MDA MB 468 CSC, and the WT cells were seeded in a 6-well plate. After

reaching 80% confluency, the cell monolayer was scraped/scratch using a 100μl pipette tip.

The cells were washed with PBS to remove all single cells. Phase-contrast images (10X

magnifications) were captured after 24h of the scratch for two cell types. Percentage cell

migration was calculated by measuring the distance between the scratches comparing the

final gap width to the initial gap width at Time T0 (zero) using Image-J software. Covered

area was calculated in 5 random fields and represented as percentage of area covered. Experiments were performed in triplicate.

2.4 Mammosphere assay:

Sorted and MDA MB 468 CSC and the WT cells were harvested by trypsinization. Cells were counted and around 800 cells per 20μl drops were kept as hanging drops on the lid of 100mm dish at 37°C under humidified conditions. 3D mammosphere of 200-250μm (average diameter) size were formed at the end of 48h for all the four cell types. At the end of four days (96h), mammospheres were collected and observed under the microscope (Primovert inverted microscope from CARL ZEISS) and images were captured. Relative area, roundness score and solidity of 3D mammosphere were calculated using Image-J software.

2.5 Immunocytochemistry assay:

The cells were cultured on 8 well chamber slides in 70% confluency followed by fixing using 4% paraformaldehyde (PFA) and permeabilization with 0.3% Triton X-100 (Sigma) in PBS. The non-specific binding sites were blocked with 5% fetal bovine serum in PBS (Himedia Laboratories, India) for 1 hour. Cells were then incubated with primary antibodies overnight at 4°C, followed by the appropriate secondary antibody for 1 hour, counterstained with DAPI for 10 minutes. Finally, the slides were mounted with Prolonged Glass Antifade (Thermo Fisher, USA). The slides were observed, and images were captured under a ZOE Fluorescent cell imager (Bio-Rad, USA). The details of the antibodies are provided in (Supplementary table-1).

2.6 Gene expression study by qRT-PCR:

One million (1×10^6) cells were used to isolate RNA using TRIzol reagent (15596026, Thermo Scientific, USA) as per the manufacturer's instructions. Quantification of RNA was done by using a Colibri Micro volume Spectrometer (Titertek-Berthold, Germany). 1μg of RNA was reverse transcribed to cDNA using iSCRIPT™ cDNA synthesis kit (Bio-Rad Catalog Number-1708891), and qRT-PCR reactions were performed using SSO-Fast™ Eva Green Supermix (Bio-Rad, USA) as per the manufacturer's instructions. The individual mRNA expressions of the tested genes of all the cell types (Ct values) were first normalized with their respective GAPDH values for getting the δCt values. Finally, the respective mRNA expressions were represented as fold change. The primer sequences were designed by the authors (Sigma, United States). The details of primers are provided in (Supplementary table-2)

157   **2.7 siRNA transfection:**

158   For transfection experiments, 0.5×10^6 cells were plated in a 6-well plate and allowed to reach

159   75% confluency. Cells were transfected with 10nM of anti-Oct-4 siRNA and

160   scrambled/control (10 nM) siRNA. D1X cationic lipid-based formulation (developed in

161   CSIR-IICT, Hyderabad) was used as a potent siRNA transfection reagent instead of the

162   commercially available transfection reagents such as lipofectamine or fugene, in the presence

163   of serum containing media as described previously (29). After 16h incubation at 37ºC 5%

164 CO_2, the media was changed, and transfection was confirmed by fluorescence microscopy.

165   Other experiments for validation were done simultaneously. We had initially transfected the

166   Cy5 labelled scrambled siRNA using the lipid-based gene delivery system (D1X) to

167   determine the transfection efficiency which was 100% as assessed using the confocal

168   microscopy.

169   **2.8 Annexin-V-FITC assay:**

170   Cancer stem cells (0.5 million) control and transfected with siRNA were washed with ice

171   cold PBS and resuspended in annexin-V binding buffer. Cells were incubated with annexin-V

172   conjugated to FITC antibody (Thermo Fisher Scientific, USA) and propidium iodide (PI) for

173   15 min in dark. Further, the cells were washed and resuspended in annexin-V binding buffer.

174   Cells (10,000 events) were acquired using Guava EasyCyte Flow Cytometer (Millipore

175   Sigma, USA) data analysis was carried out by using DE Novo FCS Express 5 software,

176   USA.The cells were analyzed on the basis of four groups in cells appearing in quadrants in

177   the flow cytometry Annexin-V-FITC dot plot: healthy (Q1), early apoptotic (Q2), late

178   apoptotic (Q3) and necrotic cells (Q4).

179   **2.9 Molecular modelling to study Oct4 and Vitamin C interactions: Development of**

180   **protein structure:**   Protein target Oct-4 structure was downloaded from database Protein

181   Data Bank (PDB) (OCT4. PDB). Vitamin C structure was downloaded from Pub Chem. (Pub

182   Chem ID54670067) in .sdf format. Conversion of Ligand i.e., Vitamin C from .sdf format to

183   .pdb format was done by using Open Babel (version 2.4.1). Hydrogen atoms and Kollman

184   charges (7.0) were added to the protein by AutoDock MGL-Tool version 1.5.6 and saved as

185   PDBQT file. Hydrogen atoms, Gasteiger charges were added to ligand and were saved in

186   PDBQT format.

187   **Protein ligand Interaction:** Vitamin C was docked with Oct4 using Auto Dock 4.2.6. Grid

188   box was obtained by taking inside the entire protein structure (blind docking method). The

grid box size maintained in configuration was 36Å × 30Å × 48Å and centre was 57.908,

0.508, and 3.182. All the information like exhaustiveness, coordinates size of grid box was

saved in .txt file and assigning the docking report in an out. pdbqt file. Ligand was aligned

with protein molecule and analysed for different binding interactions and interacting residues

using PyMol. The main purpose of this experiment is to study the mode of interactions and

binding site of Vitamin C with Oct-4.

2.10 Statistical analysis:

All the experiments were carried as biological and technical triplicates and results have been

represented as mean ± SD. The differences between more than two groups were analysed by

one way analysis of variance (ANOVA) followed by Bonferroni post-hoc test and between

two groups student t-test was used. The P values of $P \leq 0.05$ (*) and $P \leq 0.01$ (**) $P \leq 0.001$

(***) were considered as significant. Error bar represents the ± standard error of mean.

3. Results:

3.1 CD44$^+$/24$^-$ bCSCs expressed EMT genes, exhibited mammosphere forming abilities:

CD44 is considered a potential CSC marker in most cancers, and CD24 is another vital

marker whose prognostic value and significance are investigated in combination with CD44

in various cancers, including breast cancer. By using fluorescent activated cells sorting

CD44$^+$/24$^-$ CSCs were isolated from the heterogeneous TNBC cell line MDA MB 468.

Almost, 61.70% of the pure CSC population were isolated and utilized in all the further

experiments (Fig. 1A). Gene expression of the CD44$^+$/24$^-$ CSCs was performed by qRT-PCR,

the results indicated significant expression (2.5-fold) of stem cell marker CD44 (up-

regulation) and CD24 (down-regulation). Less expression of EMT marker such as EpCAM

was observed in MDA MB 468 CSCs. The qRT-PCR analysis results also showed up-

regulation of metastatic MMP-2, angiogenesis marker SP-1, proliferative marker AKT,

perhaps indicating EMT in concordance with increased stemness metastatic nature of CSCs

as compared to WT or heterogeneous population (Fig. 1F).

In order to test the intrinsic mammosphere forming potential of CD44$^+$/24$^-$ bCSCs, the

mammosphere assay was performed. The mammosphere formed by bCSC were bigger in size

and numbers compared to the WT populations (Fig. 1B). The 2D images of the

mammospheres upon analysis, exhibited relatively large area (0.570 mm^2) from CSC as

compared that of the mammospheres from WT cells (0.481 mm^2) (Fig.1C). Furthermore,

higher percentage roundness score from CSCs (80%) mammosphere was observed when

compared to mammospheres derived from the WT cells (40%) (Fig.1C). These results

validated the CSC subpopulation of MDA MB 468, for their self-renewal capacity. In other

words, the sorted CSC subpopulation demonstrated more stemness, as compared to, the WT/

parent population.

3.2 CD44$^+$/24$^-$ bCSC population exhibited migratory properties and expression of
pluripotent stem cell markers

To validate an in vitro relevance of the migration or metastasis, wound healing assay was

performed on CSC populations MDA MB 468 cell line. After 24 hours of the scratch, the

distance covered by the cells was highest in the CD44$^+$/24$^-$ CSCs and less in the WT

population (Fig. 1D) indicating a faster migration potential of bCSCs. The percentage of CSC

invading the scratch space was more than 80% in MDA MB 468 when compared to WT

control (60%) after 24h (Fig.1E).

The stem cell properties of CSCs were also reflected in immunocytochemistry (ICC)

analysis. ICC showed that the pluripotent stem cell markers such as Oct-4, Sox-2, Nanog,

expression was significantly higher in the CD44$^+$/24$^-$ bCSCs than the WT/unsorted cells (Fig.

2A and B). The sorted cell types also expressed vimentin suggesting the fact that CSC-like

property co-exists with EMT phenotype in bCSC (Wang et al., 2014) (Fig. 2A and B). These

results confirmed that the CD44$^+$/24$^-$ population exhibits CSC characteristics in conjunction

with EMT phenotype.

3.3 Oct-4 plays crucial role in CSCs sustenance and mammosphere formation:

Oct-4 is an anti-apoptotic factor and presents itself abundantly in the CSCs to save the cells

from programmed cell death (30). To assess the importance of the Oct-4 in bCSCs, the Oct-4

was knocked-down using Oct-4 siRNA. When the bCSCs were transfected with the Oct-4

siRNA, as shown in figure-3, the expression level decreased after 16h of transfection (Fig.

3C). Further, the expression level was verified using analytical flow cytometry. In the control

sample, 65.23% (Fig.3C) of the cells expressed Oct-4 protein when compared with cells

transfected with the Oct-4 siRNA which exhibited only 6-13% (Fig. 3C) of Oct-4 expression.

This assay validated the transfection efficiency and downregulation of the Oct-4 expression

in bCSCs. Moreover, morphological changes were observed in bCSCs of MDA MB 468 cells

after 16 hours post-transfection (Fig.3B). No morphological changes were observed in the

cells transfected with Cy5 labelled scrambled siRNA. As Oct-4 has been reported as an anti-

apoptotic agent, we hypothesized that siRNA mediated Oct-4 knockdown might induce CSCs

apoptosis that was further established by annexin-V expression in the Oct4 knock-down
bCSCs (Fig. 3G). In summary, knock-down of Oct-4 caused morphological changes along
with apoptotic cell-death in bCSCs.

We next sought to explore the role of Oct-4 in mammosphere formation. As shown in figure
3D, the appearance of the tumour/mammosphere was disrupted in the transfected samples
compared to the control (Fig. 3D). Both the sphere roundness and the compactness were
significantly reduced in the mammospheres formed from si-RNA-Oct4 transfected bCSCs, as
compared to, the control non-transfected bCSCs (Fig. 3D and 3F). Here, 1.5 µM Doxorubicin
was used to obtain a positive inhibition of the mammosphere formation. Taken together, the
knock-down of Oct-4 caused inhibition of mammosphere formation in bCSCs (Fig. 3D).

3.4 Vitamin C directly interacts with bCSCs by down regulation of the pluripotency
factor Oct-4

As shown in figure 4A, the bCSCs/CD44$^+$24$^-$ population when subjected to treated with
Vitamin C (10 mM) for 2h, exhibited down-regulation of Oct-4 (Fig. 4A). In the untreated
control, the Oct-4 expression was 91.99% in CSCs (Fig. 4A). Interestingly, after Vitamin C
treatment, the expression reduced drastically to 3.65% in bCSCs (Figure 4A). As we have
mentioned earlier (Sen et al., 2017, Sen et al., 2020) (18, 26), Vitamin C is responsible for
cellular damage, apoptosis, and inhibition of cell proliferation Here we have, observed the
down-regulation of the pluripotency-regulating octamer-binding transcription factor (Oct-4)
expression upon treatment with Vitamin C for 2h (Fig.4B). Based on the above-mentioned
findings, we hypothesized a co-relation/possible interaction between Oct-4 and Vitamin C
worth validation.

To further understand the possible interaction between the Oct-4 and Vitamin C, Auto dock
vina was used to investigate the binding sites and type of interactions occurring between
Vitamin C and Oct4 protein. Vitamin C – Oct-4 complexes were analysed using the
molecular (blind) docking approach. Out of nine different predicted conformations for
Vitamin C–Oct-4, top three ligand conformations having minimum binding energies were
considered. The top minimum energies were as mentioned **-5.1**, **-5.0** and **-4.9 Kcal** mol^{-1}
respectively. All the structural or binding parameters are summarized in Table 1. These
results predicted that Vitamin C binds with Oct-4 mainly in beta position of K9 to K11
sequences. The major type of interaction between Vitamin C and Oct-4 is hydrogen bond

between hydroxyl groups of the ligand to the oxygen or nitrogen atoms of amino acid main
chain of the protein. The number of hydrogen bonds found for all the three different positions
were seen in PyMol as shown in Fig. -4C-G.

Discussion:

Octamer-binding transcription factor 4 (Oct-4), a homeodomain transcription factor of the
Pit-Oct-Unc (POU) family, is considered to play an essential role in the self-renewal
epithelial-mesenchymal transition (EMT) (14) and drug resistance development of CSCs and
breast cancer metastasis (7). Oct-4 is abundantly proved as one of the most significant
transcription factors that control self-renewal and pluripotency of pluripotent stem cells and
malignant tumour succession and differentiation in CSCs (17). Hence, Oct-4 could be one of
the critical factors that target stem cell-specific cancer therapeutics.

Breast cancer, which is now a frequently detected type of cancer, is still one of the most
lethal malignancies despite recent advances in early detection and therapy (19). Conventional
therapies such as immune, chemo, and radiotherapy, can target breast cancer cells. However,
conventional therapies are often unaffordable and toxic to the patients (20, 21). Hence,
certain alternatives/additives to conventional therapies such as oral supplements,
nutraceuticals, and antioxidants might provide a better quality of life and the mitigation of the
disease burden in breast cancer patients (22). CSCs play critical roles in regulating tumour
initiation, relapse, and chemoresistance, and hence such processes must be disrupted as
therapeutic strategies (23). In the current study, we have used the MDA MB 468 cell line as
an in vitro TNBC model and isolated the CSCs based upon the surface markers $CD44^+/24^-$.
The $CD44^+/24^-$ CSCs were further characterized for the validation of the CSC phenotype. We
obtained a high expression of Oct-4 in the cancer CSC population by IF and flow-cytometric
analysis (94%). Moreover, detection of Oct-4 in metastatic cancer cells and tissues indicates
its enrichment CSCs (15). Considerable uncertainties and controversies persist, and only a
few studies have been reported attempting to target Oct-4 directly (16). According to the
available reports, inhibition of Oct-4 effectively suppressed the propagation of human
embryonal carcinoma cells and triggered their apoptotic death (15). This current study
additionally talks about the dependence of bCSC on Oct-4 for their survival/ sustenance.

Nevertheless, our results indicate a negative co-relation between Oct 4 expression and
apoptosis. Oct-4 is also known as an anti-apoptotic protein; hence repression of Oct-4 can

cause apoptosis in breast cancer (24). This current study added to the growing evidence on
the previous study's conclusion wherein we demonstrated the cellular apoptosis of bCSCs
upon Oct-4 knock-down. Furthermore, in this work, we have demonstrated the morphological
changes and disruption of mammosphere formation of bCSCs upon si-RNA mediated Oct-4
knock-down in vitro. Thus, depending on the cellular contexts, restraining Oct-4 by siRNA
may activate or inactivate its downstream counterparts, resulting in significant apoptosis in
bCSCs, resulting in therapeutic outcomes. (27). Previous reports attributed the inhibition of
Oct-4 to effectively suppress the propagation of human embryonal carcinoma cells and
triggered their apoptotic death (24, 27). This study established that Vitamin C in bCSC
downregulates the Oct-4 expression, further leading to apoptosis as it is known that the
Vitamin C increases cellular ROS levels. The interaction between Vitamin C and Oct-4 was
finally confirmed by in Silico docking, which added novel information in the literature.

Moreover, we anticipate that Oct-4 knockout by CRISPR-Cas9 technology may attain a
higher degree of inhibition on cell propagation than si-Oct4 in bCSCs (24). We established
an essential proof-of-concept that inhibiting cancer-specific Oct-4 can effectively target CSCs
and the entire bulk of cancer cells (25). Such a strategy may help tackle the significant
challenges in treating human cancers, which is worth exploring in next step. We, hence,
propose down-regulation of Oct4 first at a pre-clinical level in tumour xenograft animal
models followed by possible clinical trials using either RNAi or CRISPR-CAS9 approaches.

Conclusion:

This study depicts the role of Oct-4 in the sustenance of the bCSCs. Oct-4 is, hence, one of
the essential cancer-specific transcription factors; which gets downregulated upon treatment
with Vitamin C leading to cellular apoptosis. in bCSCs. Further, we established the
interaction between the Oct-4 and Vitamin C by the in-silico chemical docking technique. In
addition, Oct-4 could be the best candidate target for future breast cancer therapeutics for its
unique abundance in metastatic cancer and its contribution to its sustenance. Finally, the
molecule Oct-4 can be a promising diagnostic marker and therapeutic target to improvise the
lives of numerous breast cancer patients.

Acknowledgements:

The authors would like to thank Indian Council of Medical Research (ICMR) for providing the senior research fellowship to Mr Utsav Sen (Ph.D. scholar) (ICMR-2017-3769/CMB/BMS). HKR thanks Council of Scientific & Industrial Research, Govt. of India for his doctoral fellowship. Authors also thank the Yenepoya Research Centre, Yenepoya (Deemed to be University) Mangalore for its infrastructural and administrative support for conducting this research. This is IICT communication number IICT/Pubs./2021/125

Author contribution:

US designed and performed the experiments, analysed, interpreted data, and wrote the initial manuscript, and prepared the figures; AG carried out in silico docking and its data interpretation; HKR developed the lipid transfection reagent and helped in siRNA experiment. RB provided his valuable inputs in the entire manuscript, provided cationic lipid delivery system, reviewed and approved the manuscript; SS and BB directed the project, involved in designing experiments and analysis of data, wrote and approved the manuscript.

7. Supporting information:

Additional tables and information can be found in the Supporting information section.

8. Conflict of interest:

There is no conflict of interest among the authors.

9. References:

1. Yin L, Duan JJ, Bian XW, Yu SC. Triple-negative breast cancer molecular subtyping and treatment progress. Breast Cancer Res. 2020 Jun 9;22(1):61. doi: 10.1186/s13058-020-01296-5.
2. Harbeck N, Penault-Llorca F, Cortes J, Gnant M, Houssami N, Poortmans P, Ruddy K, Tsang J, Cardoso F. Breast cancer. Nat Rev Dis Primers. 2019 Sep 23;5(1):66. doi: 10.1038/s41572-019-0111-2.
3. Mattina J, MacKinnon N, Henderson VC, Fergusson D, Kimmelman J. Design and Reporting of Targeted Anticancer Preclinical Studies: A Meta-Analysis of Animal Studies Investigating Sorafenib Antitumor Efficacy. Cancer Res. 2016 Aug 15;76(16):4627-36. doi: 10.1158/0008-5472.CAN-15-3455. Epub 2016 Jun 3.
4. Saadin K, White IM. Breast cancer stem cell enrichment and isolation by mammosphere culture and its potential diagnostic applications. Expert Rev Mol Diagn. 2013 Jan;13(1):49-60. doi: 10.1586/erm.12.117.
5. Collina F, Di Bonito M, Li Bergolis V, De Laurentiis M, Vitagliano C, Cerrone M, Nuzzo F, Cantile M, Botti G. Prognostic Value of Cancer Stem Cells Markers in

Triple-Negative Breast Cancer. Biomed Res Int. 2015; 2015:158682. doi: 10.1155/2015/158682. Epub 2015 Oct 4.

6. Das A, Dhar K, Maity G, Sarkar S, Ghosh A, Haque I, Dhar G, Banerjee S, Banerjee SK. Deficiency of CCN5/WISP-2-Driven Program in breast cancer Promotes Cancer Epithelial cells to mesenchymal stem cells and Breast Cancer growth. Sci Rep. 2017 Apr 27;7(1):1220. doi: 10.1038/s41598-017-00916-z.

7. Wang D, Lu P, Zhang H, Luo M, Zhang X, Wei X, Gao J, Zhao Z, Liu C. Oct-4 and Nanog promote the epithelial-mesenchymal transition of breast cancer stem cells and are associated with poor prognosis in breast cancer patients. Oncotarget. 2014 Nov 15; 5(21):10803-15. doi: 10.18632/oncotarget.2506.

8. Fabregat I, Malfettone A, Soukupova J. New Insights into the Crossroads between EMT and Stemness in the Context of Cancer. J Clin Med. 2016 Mar 12; 5(3):37. doi: 10.3390/jcm5030037.

9. Radosa JC, Eaton A, Stempel M, Khander A, Liedtke C, Solomayer EF, Karsten M, Pilewskie M, Morrow M, King TA. Evaluation of Local and Distant Recurrence Patterns in Patients with Triple-Negative Breast Cancer According to Age. Ann Surg Oncol. 2017 Mar; 24(3):698-704. doi: 10.1245/s10434-016-5631-3. Epub 2016 Oct 25.

10. Fares J, Fares MY, Khachfe HH, Salhab HA, Fares Y. Molecular principles of metastasis: a hallmark of cancer revisited. Signal Transduct Target Ther. 2020 Mar 12;5(1):28. doi: 10.1038/s41392-020-0134-x.

11. Lu W, Kang Y. Epithelial-Mesenchymal Plasticity in Cancer Progression and Metastasis. Dev Cell. 2019 May 6;49(3):361-374. doi: 10.1016/j.devcel.2019.04.010.

12. Denisenko TV, Gorbunova AS, Zhivotovsky B. Mitochondrial Involvement in Migration, Invasion and Metastasis. Front Cell Dev Biol. 2019 Dec 20; 7:355. doi: 10.3389/fcell.2019.00355.

13. Loh CY, Chai JY, Tang TF, Wong WF, Sethi G, Shanmugam MK, Chong PP, Looi CY. The E-Cadherin and N-Cadherin Switch in Epithelial-to-Mesenchymal Transition: Signaling, Therapeutic Implications, and Challenges. Cells. 2019 Sep 20;8(10):1118. doi: 10.3390/cells8101118.

14. Zeineddine D, Hammoud AA, Mortada M, Boeuf H. The Oct4 protein: more than a magic stemness marker. Am J Stem Cells. 2014 Sep 5;3(2):74-82.

15. Kim, R. J., & Nam, J. S. (2011). OCT4 expression enhances features of cancer stem cells in a mouse model of breast cancer. Laboratory animal research, 27(2), 147.

16. Wang YJ, Herlyn M. The emerging roles of Oct4 in tumor-initiating cells. Am J Physiol Cell Physiol. 2015 Dec 1;309(11):C709-18. doi: 10.1152/ajpcell.00212.2015. Epub 2015 Oct 7.

17. Wu G, Schöler HR. Role of Oct4 in the early embryo development. Cell Regen. 2014 Apr 29;3(1):7. doi: 10.1186/2045-9769-3-7.

18. Liu A, Yu X, Liu S. Pluripotency transcription factors and cancer stem cells: small genes make a big difference. Chin J Cancer. 2013 Sep;32(9):483-7. doi: 10.5732/cjc.012.10282. Epub 2013 Feb 19.

19. Sen U, Chaudhury D, Shenoy P S, Bose B. Differential sensitivities of triple-negative breast cancer stem cell towards various doses of Vitamin C: An insight into the

internal antioxidant systems. J Cell Biochem. 2021 Apr;122(3-4):349-366. doi: 10.1002/jcb.29863. Epub 2020 Nov 2.

20. Keegan TH, DeRouen MC, Press DJ, Kurian AW, Clarke CA. Occurrence of breast cancer subtypes in adolescent and young adult women. Breast Cancer Res. 2012 Mar 27;14(2):R55. doi: 10.1186/bcr3156.

21. Arruebo M, Vilaboa N, Sáez-Gutierrez B, Lambea J, Tres A, Valladares M, González-Fernández A. Assessment of the evolution of cancer treatment therapies. Cancers (Basel). 2011 Aug 12;3(3):3279-330. doi: 10.3390/cancers3033279.

22. Schirrmacher V. From chemotherapy to biological therapy: A review of novel concepts to reduce the side effects of systemic cancer treatment (Review). Int J Oncol. 2019 Feb;54(2):407-419. doi: 10.3892/ijo.2018.4661. Epub 2018 Dec 10.

23. Lopes CM, Dourado A, Oliveira R. Phytotherapy and Nutritional Supplements on Breast Cancer. Biomed Res Int. 2017;2017:7207983. doi: 10.1155/2017/7207983. Epub 2017 Aug 6.

24. Phi LTH, Sari IN, Yang YG, Lee SH, Jun N, Kim KS, Lee YK, Kwon HY. Cancer Stem Cells (CSCs) in Drug Resistance and their Therapeutic Implications in Cancer Treatment. Stem Cells Int. 2018 Feb 28;2018:5416923. doi: 10.1155/2018/5416923.

25. Li W, Zhou Y, Zhang X, Yang Y, Dan S, Su T, She S, Dong W, Zhao Q, Jia J, Yao H, Zheng M, Kang B, Wang YJ. Dual inhibiting OCT4 and AKT potently suppresses the propagation of human cancer cells. Sci Rep. 2017 Apr 6;7:46246. doi: 10.1038/srep46246.

26. Yang L, Shi P, Zhao G, Xu J, Peng W, Zhang J, Zhang G, Wang X, Dong Z, Chen F, Cui H. Targeting cancer stem cell pathways for cancer therapy. Signal Transduct Target Ther. 2020 Feb 7;5(1):8. doi: 10.1038/s41392-020-0110-5.

27. Sen U, Chaudhury D, Shenoy P S, Bose B. Differential sensitivities of triple□negative breast cancer stem cell towards various doses of Vitamin C: An insight into the internal antioxidant systems. Journal of Cellular Biochemistry. 2021 Apr;122(3-4):349-66. doi.org/10.1002/jcb.29863

28. Kristensen, D. M., Nielsen, J. E., Kalisz, M., Dalgaard, M. D., Audouze, K., Larsen, M. E.,& Leffers, H. (2010). OCT4 and downstream factors are expressed in human somatic urogenital epithelia and in culture of epididymal spheres. Molecular human reproduction, 16(11), 835-845.

29. Rachamalla, H. K., Mondal, S. K., Deshpande, S. S., Sridharan, K., Javaji, K., Jaggarapu, M. M. C. S., Jinka, S., Bollu, V., Misra, S., & Banerjee, R. Efficient anti-tumor nano-lipoplexes with unsaturated or saturated lipid induce differential genotoxic effects in mice. Nanotoxicology. 2019 13(9):1161-1175.

30. Wang, Y. D., Cai, N., Wu, X. L., Cao, H. Z., Xie, L. L., & Zheng, P. S. (2013). OCT4 promotes tumorigenesis and inhibits apoptosis of cervical cancer cells by miR-125b/BAK1 pathway. Cell death & disease, 4(8), e760-e760.

10. Figure Legends:

Figure-1: CD44$^+$/24$^-$ bCSCs from MDA MB 468 TNBC line exhibited CSC phenotype

A) Work flow for the entire work B) FACS sorting of CSCs as CD44$^+$ and CD24$^-$cells from

MDA MB 468. C) and E) Comparative gene expression analysis of both CSCs and WT

population from MDA MB468 cells. D) Formation of mammosphere by both Wt. and CSC

population F) Quantitative readout of Mammosphere formation by the TNBCs. G) and H)

Scratch assay / migration assay results and graphical representation of the data. Scale bar =

100μm. The differences between the two groups were analysed by student t-test. The P

values of >0.12 (ns), 0.033(*), 0.002(**), <0.0002(***) were considered as significant. Error

bar represents the ± standard error of mean.

Figure-2:CD44$^+$/24$^-$ bCSCs from MDA MB 468 TNBC line expressed pluripotency

related protein. A) and D) Immunocytochemistry of WT and CSC population from MDAMB

468 line. B), C), E), F) are graphical representation of the expression of the protein expression

based up on the Mean fluorescent intensity (MFI). Scale bar = 100μm. The differences

between the two groups were analysed by student t-test. The P values of >0.12 (ns), 0.033(*),

0.002(**), <0.0002(***) were considered as significant. Error bar represents the ± standard

error of mean.

Figure 3: Inhibition of Oct-4 expression prevaricates bCSC survival

A) Confocal images of the MDA MB 468 CSCs after transfected with the Cy5 labelled

scrambled siRNA. The Images were captured 16h post-transfection. B) Morphological

changes in bCSCs after Oct-4 siRNA transfection C) Analytical flowcytometry result showing

the Oct-4 expression before and after transfection. D) Are the representative images obtained

from the mammosphere assay before and after Oct-4 siRNA transfection (Doxorubicin was

used as positive control). E) and F) are the direct read out and representation of

mammosphere assay. G) Flow cytometry analysis of the Annexin V expression before and

after Oct-4 siRNA expression in bCSCs of MDA MB 468 cells, also after treatment with the

Vitamin C. H) graphical representation of the percentage of apoptotic cells. Scale bar = 10, 25

and 50μm. The differences between the two groups were analysed by student t-test and more

than two groups were analysed by one way ANOVA, followed by Bonferroni's post-hoc test.

The P values of >0.12 (ns), 0.033(*), 0.002(**), <0.0002(***) were considered as significant.

Error bar represents the ± standard error of mean.

Figure 4: Vitamin C could be a potential target for the Oct-4

Flow cytometry data representation of the Oct-4 expression before and after Vitamin C

treatment (10mM). B) graphical representation of the flow data. C)-G) Possible interaction of

Vitamin C and Oct-4 protein by *in sillico* docking technique.

Figure 5: Schematic representation of the work and the possible mechanism of action

after Oct-4 knockdown in bCSCs.

Oct-4 is very crucial for cell proliferation and sustenance and even for the aggressiveness of

cancer. And this unique pluripotent transcription factor is only expressed in ESCs and CSCs.

When this transcription factor was knocked down with siRNA, the cellular process altered

drastically. The bCSCs morphology was changed, the cells were undergoing apoptosis; it was

also found the expression of Oct-4 was downregulated. Most interestingly, Oct-4 knockdown

not only influenced the cellular sustenance but also affected tumour formation ability. It was

previously reported the Vitamin C causes apoptosis in bCSCs, and here we have shown that

Vitamin C also interacts with Oct 4 by flow cytometry and sillico docking. In summary, Oct-4

can be a potential target for the CSC-based therapeutic maneuvering for breast cancer

treatment.

Figure-1

Figure-2

Figure-3

(A) Unstained | Stained with Oct-4/ Untretaed | Stained with Oct-4/Vit C 10mM

(D)

Serial No.	Binding Energy (Kcal. Mol^{-1})	Atoms of Vitamin C involved in H-bonding	Oct4 atoms involved in H-bonding	Bond length (Å)
1.	-5.1	8OH	NH LYS 11	2.3
		10OH	CO LYS 11	2.6
		1OH	NH ARG 10	1.8
		1OH	NH LYS 9	2.6
2.	-5.0	6O	NH ARG 8	2.5
		6O	NH ARG 10	2.3
		10OH	CO LYS 11	2.3
		12OH	CO LYS 11	2.5
		12OH	CO LEU 15	2.1
3.	-4.9	6O	NH LYS 9	2.1
		6O	NH ARG 10	1.9
		10OH	CO ARN17	2.2

Figure-4

Figure-5