Title: Rapamycin induced hyperglycemia is associated with exacerbated age-related osteoarthritis

Authors: Dennis M. Minton[1,2], Christian J. Elliehausen[1,2], Martin A. Javors[3], Kelly S. Santangello[4], Adam R. Konopka[1,2,5]

[1]Department of Medicine, University of Wisconsin-Madison; [2]Department of Kinesiology, University of Illinois at Urbana-Champaign; [3]Departments of Psychiatry and Pharmacology at University of Texas Health, San Antonio; [4]Department of Microbiology, Immunology, Pathology, Colorado State University; [5]Geriatric Research, Education, and Clinical Center, William S. Middleton Memorial Veterans Hospital.

Running Title: Off-target effects of rapamycin worsen age-related OA

Corresponding Author:
Adam R. Konopka, PhD
Department of Medicine, Division of Geriatrics and Gerontology
University of Wisconsin-Madison
Geriatric Research, Education, and Clinical Center
William S. Middleton Memorial Veterans Hospital
akonopka@medicine.wisc.edu

Abstract

Background: The objective of this study was to determine if mechanistic target of rapamycin (mTOR) inhibition with or without AMP-activated protein kinase (AMPK) activation can protect against primary, age-related OA.

Design: Dunkin-Hartley guinea pigs develop mild primary OA pathology by 5-months of age that progresses to moderate OA by 8-months of age. At 5-months, guinea pigs sacrificed as young control (n=3) or were fed either a control diet (n=8), a diet enriched with the mTOR-inhibitor rapamycin (Rap, 14ppm, n=8), or Rap with the AMPK-activator metformin (Rap+Met, 1000ppm, n=8) for 12 weeks. Knee joints were evaluated by OARSI scoring, micro-computed tomography, and immunohistochemistry. Glenohumeral articular cartilage was collected for western blotting.

Results: Rap and Rap+Met treated guinea pigs displayed lower body weight than control. Rap and Rap+Met inhibited articular cartilage mTORC1 but not mTORC2 signaling. Rap+Met, but not Rap alone, stimulated AMPK. Despite lower body weight and articular cartilage mTORC1 inhibition, Rap and Rap+Met treated guinea pigs had greater OA severity in the medial tibial plateau due to articular cartilage structural damage and/or proteoglycan loss. Rap and Rap+Met increased plasma glucose compared to control. Plasma glucose concentration was positively correlated with proteoglycan loss, suggesting hyperglycemic stress may have contributed to worsened OA.

Conclusions: This is the first study to show that Rap induced increase in plasma glucose was associated with greater OA severity. Further, articular cartilage mTORC1 inhibition and bodyweight reduction by dietary Rap and Rap+Met did not protect against primary OA during the prevailing hyperglycemia.

Key Words: Aging, mTOR, AMPK, Dunkin Hartley Guinea Pig, Primary Osteoarthritis

Background

Primary, age-related osteoarthritis (OA) is estimated to account for as many as 90% of all knee OA cases in humans (1). However, preclinical research commonly relies on experimental models of secondary OA. Although primary and secondary OA share similar pathological outcomes, there is a growing body of evidence to suggest they are driven by distinct mechanisms. Retrospective analysis of differentially expressed genes from separate cohorts of primary and secondary OA patients relative to their healthy controls found that only 10% of differentially upregulated and 35% of differentially downregulated genes in OA vs non-OA samples are conserved between primary and secondary OA (2,3). Therefore, 65-90% of differentially expressed genes may be unique to primary versus secondary OA. Additionally, transgenic animal models have revealed that several genes are differentially involved in the progression of primary and secondary OA (4–9). For example, deletion of *Panx3* protects against secondary OA yet dramatically worsens primary OA (4), and deletion of *JNK1/2* accelerates the development of primary OA while having no effect on secondary OA progression (9). Together, these studies reinforce that unique mechanisms underpin these two forms of OA.

Age is one of the greatest risk factors for nearly every chronic disease, including primary OA. Two evolutionarily conserved kinases, mechanistic target of rapamycin (mTOR) and AMP-activated protein kinase (AMPK), are energy sensing pathways similarly dysregulated during aging and OA (10–13). The mTOR inhibitor rapamycin (Rap) can extend lifespan in mice and delay the onset of several age-related morbidities (12,14). The anti-diabetic drug metformin (Met) can activate AMPK and, when added to Rap, extends lifespan to a greater extent than historical cohorts of mice treated with Met or

Rap alone (15). Additionally, Met is the first drug being tested to slow age-related multi-

morbidity in humans (16). While the prospect of lifespan extension is tantalizing,

extending lifespan without delaying the onset or slowing the progression of the most

debilitating age-associated conditions could be viewed as detrimental. Therefore, it is

imperative to understand if purported lifespan-extending therapies that target the

fundamental biology of aging are also capable of delaying the onset of chronic diseases,

such as primary OA.

mTOR exists as complex I (mTORC1) and complex II (mTORC2). mTORC1 regulates

cellular proliferation, protein synthesis, senescence, and survival while mTORC2

functions downstream of insulin signaling on substrates such as PI3K-Akt (12). In articular

cartilage, mTORC1 activity increases with age and is sufficient to induce OA in young

male mice (10). In non-articular tissues, acute or intermittent Rap selectively inhibits

mTORC1 while chronic Rap administration for durations greater than 14 days also inhibits

mTORC2 activity (17). Cartilage-specific deletion of mTOR and systemic or intra-articular

injections of Rap and the mTORC1/2 inhibitor Torin 1 lower secondary OA in young-male

mice and rabbits (18–21). While these findings support mTOR-based therapeutics for OA,

the completed studies were exclusively in injury-induced models of OA and have not been

investigated in primary, age-related OA.

Recently, it has been proposed that the positive effects of mTOR inhibition on OA

pathology may be diminished by feedback activation of PI3K and has raised questions

about the need for a dual treatment strategy that inhibits both mTOR and upstream PI3K

signaling (22,23). In addition to activating AMPK, Met has pleotropic effects including

inhibition of PI3K signaling in rheumatoid arthritis fibroblast-like synoviocytes (24).

Moreover, Met and other AMPK-activators have chondroprotective effects against

inflammatory-induced protease expression *in vitro* (25,26) and protect against injury-

induced OA in young male mice and rhesus monkeys (27). Treatment with Met is also is

associated with a lower rate of medial tibiofemoral cartilage volume loss and risk of total

knee replacement in obese patients (28). However, Met as an adjuvant therapy to Rap

has not been investigated in primary OA.

The Dunkin-Hartley guinea pig is a well-characterized outbred model of primary OA.

The progression of OA in guinea pigs is related to bodyweight (29) and shares a similar

age-related and spatial progression to humans (30). Mild OA pathology develops by 5

months in guinea pigs that progresses to moderate OA by 8-9 months of age (30–32).

Therefore, at 5 months of age we treated guinea pigs with lifespan-extending doses of

Rap or a combination of Rap+Met for 12 weeks to slow the progression from mild to

moderate OA. This study is the first to evaluate if lifespan extending treatments can

modify primary OA, the most prevalent form of OA observed in older adults.

Methods

<u>Animal Use</u>

All tissues were collected at the University of Illinois Urbana-Champaign and approved by the Institutional Animal Care and Use Committee. Data collection and analysis were completed at University of Wisconsin-Madison and William S. Middleton Memorial Veterans Hospital. Because male Dunkin-Hartley guinea pigs develop more severe OA pathology than female (33), we used male animals to maximize the potential for the interventions to slow the progression of OA. Therefore, similar to previous work (34), male Dunkin-Hartley guinea pigs (Charles River) were singly housed in clear plastic, flat bottomed cages (Thoren, Model #6) with bedding. Guinea pigs were single housed to measure food consumption. 12-hour light/dark cycles were used beginning at 0600. Guinea pigs acclimated for 2-3 weeks and were provided standard chow diet (Evigo 2040) fortified with vitamin C (1050 ppm) and Vitamin D (1.5 IU/kg) and water ad libitum until 5 months of age. Guinea pigs were then sacrificed to serve as young control (n=3), randomized to continue the standard diet (n=8), or receive standard diets enriched with encapsulated rapamycin (14 ppm, n=8) or the combination of encapsulated rapamycin and metformin (14 ppm, 1000 ppm, n=8) for 12 weeks. Guinea pigs were randomized to match bodyweight between groups prior to beginning treatment. Diets were enriched with microencapsulated rapamycin (Rapamycin holdings) and/or metformin (AK Scientific, I506) at concentrations previously shown to extend lifespan in mice (14,15,35). Food consumption was recorded on Monday, Wednesday, and Friday between 8 and 9 AM, and body weight was recorded before feeding on Monday. Guinea pigs treated with Rap or Rap+Met diet had ad libitum access to food. Dietary Rap treatment has been shown to

significantly reduce bodyweight in mice (36,37). Therefore, we matched food

consumption in the control group to the Rap diets to minimize the influence of food intake

on dependent variables. One guinea pig in the Rap+Met group was euthanized early due

to a wound on the gums which led to suppressed appetite and infection. Tissues from this

animal were not collected for analysis. It could not be determined if this was due to a

laceration or an oral ulcer, the latter of which is a known side effect of mTOR inhibitors

(38).

Tissue Collection

Two animals were sacrificed daily between 7 and 10 AM. Food and water were

removed from the cages 2-4 hours before euthanasia. Animals were anesthetized in a

chamber containing 5% isoflurane gas in oxygen and maintained using a face mask with

1.5-3% isoflurane. Blood was collected by cardiac venipuncture followed by excision of

the heart. The right hind limb was removed at the coxofemoral joint, fixed in 10% neutral

buffered formalin (NBF) for 48 hours, and transferred to 70% ethanol until processed for

histology. Glenohumeral cartilage was collected, snap frozen in liquid nitrogen, and stored

at -80C for further analysis. Because testicular atrophy has been observed following Rap

treatment (39), the left testicle was preserved in 10% NBF and weighed. Although tissues

are commonly weighed before fixation, previous work demonstrates that fixation

negligibly effects testicle weight in similarly sized rodents (40).

Analysis of Experimental Diets and Blood

Samples of diets enriched with Rap, Met or the combination of Rap+Met, and

aliquots of whole blood (n=4 per group) were sent to the Bioanalytical Pharmacology Core

at the San Antonio Nathan Shock Center to confirm drug concentrations in the diet and

in circulation. Analysis was performed using tandem HPLC-MS as described previously

(14,41,42). Frozen aliquots of plasma were thawed to measure glucose and lactate

concentrations using the YSI Biochemistry Analyzer (YSI 2900).

Micro Computed Tomography (µCT)

Right hind limbs from half of each treatment group (n=4 per group) were scanned

using a Rigaku CT Lab GX130 at 120 μA and 110 kV for 14 minutes, achieving a pixel

size of 49 μm. Scans were first processed in Amira 6.7 (ThermoFisher) where epicondylar

width was measured and a series of dilation, erosion, filling, and image subtraction

functions were used to isolate trabecular and cortical bone as described previously (43).

Scans were then resliced 4 times along axes perpendicular to medial and lateral tibial

and femoral articular surfaces and binarized using identical thresholds. NIH ImageJ

software and BoneJ plugin were used to quantify thickness, spacing, and volume fraction

measurements. Cortical thickness was measured by placing polygonal regions of interest

(ROI) in resliced scans to encompass the articular surfaces in each joint compartment.

Trabecular thickness, spacing, and bone volume fraction were measured by placing

transverse ROIs (2.4x2.4x1mm) in the trabecular bone of each joint compartment.

<u>Histology</u>

Knee joints were decalcified in a 5% ethylenediaminetetraacetic acid, changed every 2-

3 days for 6 weeks. Joints were then cut in a coronal plane along the medial collateral

ligament, paraffin embedded and sectioned at 5um increments for Toluidine Blue

staining and immunohistochemistry (IHC). Slides were scanned using the Hamamatsu

NanoZoomer Digital Pathology System, providing 460nm resolution. Scan focus points

were set manually along the articular cartilage. Imaged slides were then scored by two

blinded reviewers for OA severity following OARSI Modified Mankin guidelines as

described (32). Briefly, toluidine blue stained histology slides were assigned scores for

severity of articular cartilage structural damage (0-8), proteoglycan loss as assessed by

absence of toluidine blue staining (0-6), disruption of chondrocyte cellularity (0-3), and

tidemark integrity (0-1), with a total possible score of 18 per joint compartment (Total

OARSI Score). One guinea pig each from the Rap and Met groups were unable to be

analyzed due to off-axis transection before embedding. One control animal was a

statistical outlier as detected by Grubb's test and was excluded from the study.

Therefore, n=7 per group were used for histopathological analysis.

<u>Immunohistochemistry</u>

Antigen retrieval was performed in 10mM sodium citrate for 7 hours at 60C.

Endogenous peroxidase activity was quenched using 3% H_2O_2 for 15min before blocking

in 5% normal goat serum diluted in TBST for 1 hour at RT. Slides were incubated

overnight in 200-300 uL of either p-RPS6 (1:200 dilution; Cell Signaling, 4858) or a rabbit

IgG isotype control (Cell Signaling, 3900) diluted to match primary antibody concentration.

Primary antibodies against p-Akt Ser473 (1:100 dilution; 4060) and p-AMPK Thr172

(1:200 dilution; 50081) from Cell Signaling were attempted, but reactivity was not seen in

guinea pig articular cartilage. 150-200uL of goat anti-rabbit secondary antibody (Cell

Signaling, 8114) was added for 1 hour at room temperature followed by exposure in 3,3'-

diaminobenzadine (DAB; Cell Signaling, 8059) for 10 minutes. Slides were then

counterstained using hematoxylin, dehydrated, and cleared through graded ethanol and

xylene, coverslipped using Permount (Electron Microscopy Sciences), and viewed and

imaged under a brightfield microscope. No DAB staining was seen following incubation

with the IgG control or secondary antibody alone, confirming specificity of the primary

antibody. For quantification, ROIs were placed to encompass areas of staining in the

medial tibial articular cartilage, and cells were counted to determine the percent-positive

cells. For intensity-based quantification, a color deconvolution for DAB staining was

applied in ImageJ, and mean integrated intensity was quantified by averaging two p-RPS6

replicates and subtracting background staining of IgG controls.

<u>Western Blot</u>

Cartilage was removed from the glenohumeral joint using a scalpel and placed in

reinforced Eppendorf tubes containing 500 mg of ceramic beads (Fisher, 15-340-160)

and 200 μL of RIPA buffer with protease and phosphatase inhibitors (Sigma,

5892970001), and homogenized by 2, 30-second cycles at 6 m/s in the Omni

BeadRuptor. Homogenate was transferred to microcentrifuge tubes and spun at 10,000g

for 10 min at 4C. Supernatants were diluted to equal concentration following a BCA assay.

Samples were prepared in reducing conditions with β-mercaptoethanol in 4x Laemmli

Sample Buffer (BioRad, 1610747) and heated at 95C for 5 minutes. 10 μg of protein was

separated on 4-15% TGX precast gels (BioRad, 4561083) and transferred to PVDF

membranes (BioRad, 1620177). Membranes were blocked in TBST with 5% bovine

serum albumin (Sigma, A9647) for 1 hour at RT and incubated overnight at 4C in primary

antibodies against p-RPS6 Ser235/236 (4858), RPS6 (2217) p-Akt Ser473 (4060), Akt

(4685), P-AMPK Thr172 (50081), AMPK (2532), and LC3B (3868) from Cell Signaling

and ADAMTS5 (ab41037), MMP-13 (ab39012), and b-Actin (ab8226) from Abcam. HRP-

conjugated anti-Rabbit (Cell Signaling) or anti-Mouse (Abcam) secondary antibodies

were diluted 1:5,000 for all proteins except b-Actin (1:10,000 dilution). All membranes

were imaged using a UVP BioSpectrum 500 (UVP) following 5-minute incubation in a 2:1

combination of SuperSignal Pico (Fisher, 34577) and Femto (Fisher, PI34095)

chemiluminescent substrates except b-Actin which received Pico alone. Densitometric

analysis was performed using VisionWorks (Analytikjena). Phosphorylated proteins are

expressed relative to their total protein and other targets are expressed relative to b-Actin.

Statistical Analysis

Previous work demonstrated that a sample size of n=6 is adequately powered to

detect changes between groups in guinea pigs (34). Therefore, we a priori determined

our sample size (n=7-8 per group) to be appropriate to detect differences between

treatment groups. All data were subjected to normality testing via the Shapiro-Wilk test.

Comparisons of normally distributed data were performed using two-way unpaired t-tests

or one-way ANOVA followed by Holm-Sidak's multiple comparison test. Data with non-

Gaussian distribution were compared using non-parametric Mann-Whitney tests or the

Kruskal-Wallis test followed by Dunn's multiple comparisons test. A two-way repeated

measures ANOVA (time x treatment) was performed to determine differences in food

consumption and body weight. Upon a significant interaction, Holm-Sidak's multiple

comparisons test was used. Because we were interested in determining if treatments

impacted the trajectory of OA pathogenesis compared to aged controls, differences in all

other variables besides plasma glucose were made using one-way ANOVA comparing

treatment groups to 8-month controls. Due to previous reports that Met can rescue the

hyperglycemic effects of Rap (37), comparisons were made between all groups for

plasma glucose. Pearson's R was used to determine correlation between variables. P-

values <0.05 were considered statistically significant. Data are presented as scatter plots

with mean or mean ± standard deviation (SD).

Results

Influence of rapamycin and rapamycin+metformin on guinea pig physical and metabolic characteristics

Figure 1A shows the average daily food consumption per week of standard diet or standard diet enriched with Rap or Rap+Met. The average daily intakes of Rap and Met based on food consumption and dietary concentration are reported in Table 1. Compared to control, there was decreased food consumption in guinea pigs receiving Rap+Met during week 2 (P=0.04). There were no significant differences between treatments. Despite largely matching food intake, there was a significant effect for treatment (P=0.004) and an interaction between time and treatment (P<0.0001) on bodyweight. Rap+Met (P=0.01) and Rap-treated guinea pigs (P=0.02) were smaller than control starting at week 3 and week 4, respectively, until the end of the study (Figure 1B). At sacrifice, Rap (P=0.002) and Rap+Met-treated guinea pigs (P=0.001) were 15% and 22% smaller than control.

Treatment with Rap (396±61 mg/dL; P<0.0001) and Rap+Met (334±53 mg/dL; P=0.007) increased plasma glucose compared to control (234±55 mg/dL), and the addition of Met to Rap decreased plasma glucose compared to Rap alone (P=0.05; Figure 1C). Lactate concentration trended to be elevated by 66% in Rap+Met-treated guinea pigs, only (P=0.07; Figure 1D). Testicle weight in guinea pigs receiving Rap (P=0.006) and Rap+Met (P=0.0003) were 27% and 44% lower than control, respectively, suggesting gonadal atrophy (Figure 1E). We analyzed blood for the circulating Rap and Met concentrations ~3-hours after food had been removed from the cage (Table 2). This timing aligns with a measurement of peak circulating Rap and Met. We saw that

experimental diets were sufficient to increase Rap and Met concentrations in the blood,

and that Rap values were not different when providing diets individually or in combination.

There was no Rap or Met detected in circulation in control animals.

Rapamycin and rapamycin+metformin treatment exacerbated the age-related

progression of OA

Consistent with the age-related progression of mild to moderate OA in guinea pigs,

we observed an increase in medial tibial total OARSI score from 5 to 8 months (P=0.03;

Figure S1A-B). Surprisingly, Rap and Rap+Met treatment resulted in a ~2-fold increase

in total OARSI score in the medial tibial plateau compared to 8 month old, age-matched

control (P=0.02 for both Rap and Rap+Met; Figure 2B). This was driven by increased

scores for articular cartilage structure (P=0.02 for Rap, P=0.11 for Rap+Met; Figure 2C)

and proteoglycan loss (P=0.02 for Rap and Rap+Met; Figure 2D). There was no

significant effect of Rap or Rap+Met on the OARSI score for the lateral tibia or medial or

lateral femur (Figure S1C).

OA pathology was correlated to plasma glucose, bodyweight, and testicle weight

Because Rap and Rap+Met treated guinea pigs displayed several side effects of

Rap, including increased plasma glucose, testicular atrophy, decreased bodyweight, and

worsened OA pathology, we evaluated the relationship between these variables and

measures of OA severity across all guinea pigs. Plasma glucose was positively correlated

to proteoglycan loss (R^2=0.19; P=0.04; Figure 3A), and total OARSI score was negatively

correlated with both bodyweight (R^2=0.19; P=0.04; Figure 3B) and testicle weight

(R^2=0.20; P=0.04; Figure 3C). However, because testicle weight and bodyweight were

also related (data not shown), the individual contribution of these variables cannot be

resolved.

Effects of rapamycin and rapamycin+metformin on mTOR, AMPK, and protease

expression

To evaluate mTORC1 signaling in articular cartilage, we measured the

phosphorylation of ribosomal protein S6 (P-RPS6) at Ser235/236 using IHC and western

blotting. Representative images of P-RPS6 IHC are shown in Figure 4A. P-RPS6 was

decreased by 90-95% in the medial tibial articular cartilage of Rap and Rap+Met treated

guinea pigs as assessed by percentage of P-RPS6-positive cells (P=0.001 for Rap,

P=0.01 for Rap+Met; Figure 4B), and by staining intensity (P=0.02 for both; Figure 4C).

mTORC1 inhibition was further supported by an 81% lower ratio of phosphorylated to

total RPS6 in glenohumeral cartilage from Rap (P=0.005; Figure 4E). Rap+Met trended

to decrease RPS6 phosphorylation by 48% (P=0.06). There were no signficant effect on

the phosphorylation of the mTORC2 substrate Akt at Ser473 in Rap or Rap+Met

compared to control (Figure 4F; P=0.11). AMPK activity was measured using western blot

to assess phosphorylation of AMPK at Thr172 (P-AMPK). P-AMPK was not changed by

Rap alone (P=0.83; Figure 4G) but was elevated 77% by Rap+Met (P=0.05). Rap or

Rap+Met did not significantly change the conversion of LC3B I to II (P>0.99 for both;

Figure 4H) nor a disintegrin and metalloproteinase with thrombospondin motifs 5

(ADAMTS5; Figure 4I; P=0.97 for Rap, P=0.35 for Rap+Met). Matrix metalloproteinase

13 (MMP13) was unchanged by Rap (P>0.99) but trended higher in Rap+Met (P=0.09;

Figure 4J).

Rapamycin and rapamycin+metformin decreased subchondral and diaphyseal bone

thickness

Representative microCT images shown in Figure 5A were used to quantify the

effect of experimental diets on subchondral bone parameters. Mean subchondral cortical

thickness was decreased by Rap and Rap+Met in the medial (29%, P=0.003 for Rap;

23%, P=0.007 for Rap+Met) and lateral (21% for Rap; 20% for Rap+Met; P=0.01 for both)

tibia (Figure 5B). Rap and Rap+Met decreased trabecular spacing by 15% and 16%,

respectively, in the lateral tibia only (P=0.006 for both; Figure S2B). Trabecular thickness,

trabecular spacing in other compartments, and bone volume fraction were not affected by

any experimental diet (Figures S2A-C). Further investigation revealed that cortical

thickness at the femoral diaphysis was decreased by Rap (P=0.001) and Rap+Met

(P=0.01; Fig 5C), and this change was proportionate to the decrease observed in the

medial tibial subchondral bone (Figure 5D). Further, medial tibial cortical thickness was

correlated to bodyweight (R^2=0.47, P=0.01; Figure 5E), suggesting the smaller body

mass of Rap and Rap+Met treated guinea pigs may have contributed to decreased

cortical thickness. Femoral epicondylar width (Figure 5F) was not statistically different

between groups (Rap, P=0.42; Rap+Met, P=0.45), suggesting our treatments did not

affect skeletal development.

Discussion

The purpose of this study was to test if dietary Rap or Rap+Met could delay the

onset of age-related OA in the outbred Dunkin-Hartley guinea pig. We found that at

concentrations shown to extend lifespan, dietary Rap and Rap+Met inhibited mTORC1

but not mTORC2 signaling in articular cartilage, and Rap+Met increased AMPK

phosphorylation. Surprisingly, guinea pigs treated with Rap, with or without Met,

developed greater age-related OA compared to control. Guinea pigs receiving Rap and

Rap+Met also displayed increased plasma glucose, which correlated with proteoglycan

loss. These findings indicate that off-target side effects of Rap are associated with greater

OA pathology. Further, in the face of these Rap-induced side effects, mTORC1 inhibition

may not slow the progression of age-related OA in Dunkin Harltey guinea pigs.

Despite inhibiting mTORC1 in articular cartilage, our findings indicate that guinea

pigs treated with Rap, with or without Met, had exacerbated age-related OA in the medial

tibial plateau. Further, Rap and Rap+Met treated guinea pigs had greater total OARSI

scores even though they weighed less, which is contrary to previous work where lower

body weight was accompanied by lower OA scores in guinea pigs (29). Although there is

precedent that mTORC1 inhibition by intra-articular injection of Rap is associated with

exacerbated temporomandibular joint (TMJ) OA (44), our findings were opposite of our

original hypothesis and previous results using Rap in secondary models of knee OA

(18,19). The guinea pigs in the current study received a dose of Rap that achieved similar

circulating Rap concentrations shown to extend lifespan in mice (14). Additionally, the

dose of Rap in guinea pigs was similar to the dose shown to protect against secondary

OA in mice (0.7 vs 1 mg/kg/day in guinea pigs vs. mice) (18). These findings suggest that

dose of Rap was not a likely factor contributing to differences between studies. In our

study, Rap and Rap+Met treatment inhibited mTORC1 but not mTORC2 in articular

cartilage. Previous work has shown that deleting articular cartilage mTOR (21) or treating

with Rap (18,19) or Torin-1 (20) can attenuate secondary OA in mice and rabbits. These

non-selective genetic and pharmacological methods likely disrupt the entire mTOR kinase

and therefore could inhibit both mTORC1 and mTORC2 signaling. However, this remains

speculative as mTORC2 signaling was not evaluated in these previous studies, and it

continues to be unknown if mTORC2 inhibition is necessary for protection against either

primary or secondary OA. In support of the notion that targeting mTORC2 modifies OA,

inhibition of the mTORC2 substrate Akt protects against PTEN-deletion-induced OA by

decreasing cellular senescence and oxidative stress (45). Further investigation is needed

to resolve the role of each mTOR complex in the initiation, progression, and treatment of

both primary and secondary OA.

Despite its lifespan-extending effects, chronic Rap treatment is commonly

associated with several metabolic and immunological side effects including glucose

intolerance, insulin resistance, hypertriglyceridemia, immunosuppression, testicular

atrophy, lower body weight, and cataracts (17,39,46). Consistent with this, we showed

that 12-weeks of dietary Rap and Rap+Met was accompanied by increased plasma

glucose, testicular atrophy, and lower body weight. Despite increasing AMPK activity in

articular cartilage and partially restoring glucose levels compared to Rap alone, the

addition of Met to Rap did not offer protection against the detrimental effects of dietary

Rap on OA pathology. The glucose lowering effects of Met are in line with previous studies

where Met alleviated Rap-induced glucose intolerance only in female mice (37).

However, our OA pathology findings are in contrast to previous studies that showed Met

attenuated hyperglycemia-induced OA in mice (55). In our study, medial tibial

proteoglycan loss was correlated with plasma glucose, and we propose that Rap-induced

hyperglycemia may have contributed to worsened OA following dietary Rap treatment. In

support of this hypothesis, diabetic mice show accelerated OA after injury, and

chondrocytes cultured in high glucose media display decreased expression of Collagen

II and increased MMP13 and inflammatory mediators IL-6 and NFkB (47,48). However,

intermittent intraperitoneal injections of Rap lowered glucose and mitigated diabetes

accelerated secondary OA (49). It is possible that Rap did offer partial protection against

hyperglycemic stress but still resulted in greater OA pathology than control, as was

observed by Ribeiro et al. (50). However, this remains speculative as we did not have a

group exposed to hyperglycemic stress alone. Previous work suggests Rap can have

divergent effects where it is beneficial in some diabetic models but causes adverse side

effects in metabolically healthy models (17,51). Collectively, these data indicate that the

adverse metabolic side-effects of dietary Rap treatment are associated with a deleterious

impact on primary OA pathology and could limit the utility of systemic Rap as a healthspan

extending treatment.

Rap has been implicated in attenuating secondary OA by increasing autophagy

and decreasing protease expression (18,19). While autophagy is a highly dynamic

process, the static marker of autophagy, LC3B, is commonly used as a surrogate for

autophagic flux. In our study, we saw no effect by any treatment on LC3B or ADAMTS5,

while Rap+Met trended to increase MMP13 in glenohumeral cartilage. Therefore, the

inability to increase markers of autophagy and decrease proteases may be one

contributing factor to why our lifespan-extending treatments did not protect and even

worsened OA during aging and hyperglycemia. However, because proteoglycan loss was

observed independent of increased protease expression in Rap-treated guinea pigs,

decreased extracellular matrix (ECM) protein synthesis may have contributed to

proteoglycan loss. More work is needed to determine the molecular and cellular

mechanisms responsible for the deleterious effects of Rap and Rap+Met.

Treatment with Rap and Rap+Met also decreased subchondral cortical bone

thickness in the medial and lateral tibia and the femoral diaphysis. As bone growth in

guinea pigs ceases by 4 months (52), and epicondylar width was not different between

groups, the differences in bone thickness were likely not the result of disrupted

development. Decreased subchondral thickness was only observed in the tibia. Intra-

articular injection of Rap into the TMJ caused subchondral bone loss by inhibiting pre-

osteoblast proliferation (44), and Rap treatment also decreased osteoblast differentiation

and bone matrix synthesis (53), which supports the idea that Rap can act directly on the

bone to decrease thickness. However, we also found that subchondral thickness was

highly correlated to bodyweight. This is in line with Wolff's law and agrees with previous

findings where bodyweight restriction decreased cortical bone thickness in the femoral

diaphysis (54). Therefore, both local and systemic effects of Rap likely contributed to

reduced cortical bone thickness.

Although we provide new insight into the role of mTOR during primary OA

progression, we recognize some study limitations. While the guinea pig is an excellent

model of primary OA, it is not a widespread model for biomedical research and molecular

probes are seldom designed for reactivity with guinea pig tissue. Due to reactivity issues

with IHC in guinea pig cartilage (Figure S3), some of our analyses relied on western blot

from glenohumeral cartilage. Although guinea pigs also develop mild glenohumeral

OA(30), this is not the site at which we measured OA pathology. Our study could not

conclusively determine if the deleterious effects of Rap stemmed from its direct effects on

the joint or off-target effects on other tissues. However, our data suggest hyperglycemia

induced by off-target actions of Rap was associated with worsened age-related OA. The

Dunkin Hartley guinea pig is an outbred model of primary OA which leads to inherent

variability. While this could be perceived as a limitation, we contend that the variability

and the choice of animal model adds translational value since this more closely

recapitulates the genetic diversity and OA heterogeneity in humans. We acknowledge

that although the sample size used in our study was in line with previous studies using

guinea pigs, the varability could have possibly limited our ability to detect more subtle

differences between groups. However, this does not detract from the findings that guinea

pigs treated with both Rap and Rap+Met had worse OA. Further, the presence of largely

overlapping and consistent deleterious outcomes in both groups receiving Rap increases

our confidence that the side effects accompanying Rap contribute to worsened primary

OA.

Conclusion

In summary, we have shown that at doses previously shown to extend lifespan, dietary

Rap and Rap+Met caused hyperglycemia and was associated with aggravated OA in

Dunkin Hartley guinea pigs despite inhibiting mTORC1 in articular cartilage. Treatments

that extend lifespan without a proportional delay in age-related chronic diseases and

disabilities is counter to the concept of healthspan extension. Our findings that guinea

pigs treated with Rap had worse OA pathology raises concerns regarding the efficacy of

dietary Rap as a life- and healthspan-extending agent. Additional work is needed to

investigate the role of alternative routes of administration or Rap anaologs that may

capture the positive benefits of Rap while minimizing off-target effects. Our data also

reveal that the contribution of mTOR in articular cartilage and chondrocyte metabolism is

incompletely understood and additional research is needed to clarify the individual and

combined role of mTORC1 and mTORC2 signaling in both primary and secondary OA.

Abbreviations

OA: osteoarthritis; mTOR: mechanistic target of rapamycin; AMPK: AMP-activated

protein kinase; Rap: rapamycin; Met: metformin; mTORC1: mTOR complex I; mTORC2:

mTOR complex II; NBF: neutral buffered formalin; µCT: micro computed tomography;

ROI: region of interest; IHC: immunohistochemistry; OARSI: osteoarthritis research

society international; SD: standard deviation; RPS6: ribosomal protein S6; ADAMTS5: a

disintegrin and metalloproteinase with thrombospondin motifs 5; MMP13: matrix

metalloproteinase 13; TMJ: temporomandibular joint; IL-6: interleukin 6; NFkB: nuclear

factor kappa-light-chain-enhancer of activated B-cells; ECM: extracellular matrix.

Acknowledgements

The authors would like to thank Greg Friesenhahn at the Analytical Pharmacology and

Drug Evaluation Core of the San Antonio Nathan Shock Center. We would also like to

acknowledge the technical assistance from William Fairfield, Oscar Safairad, Alex Nichol,

Nathan Carper, and Morgan Berland. We also acknowledge the assistance of the

University of Wisconsin Translational Research Initiatives in Pathology (TRIP) laboratory

supported by the UW Department of Pathology and Laboratory Medicine, UWCCC (P30

CA014520).

Author Contributions

Study design: AK. Data collection: DM, CE, AK, KS, MJ. Data analysis and

interpretation: DM, AK. Manuscript preparation: DM, AK. All authors approved of the

final manuscript.

Funding

The Konopka Laboratory was supported by the Campus Research Board at the University

of Illinois and startup funds from the University of Wisconsin-Madison School of Medicine

and Public Health, Department of Medicine, and National Institutes of Health grant R21

AG067464. This work was supported using facilities and resources from the William S.

Middleton Memorial Veterans Hospital. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the NIH, the Department

of Veterans Affairs, or the United States Government.

Availabililty of Data and Materials

Data from this study are available from the corresponding author upon reasonable

request.

Ethics Approval

Animal use was approved by the University of Illinois at Urbana-Champaign IRB and

IACUC.

Consent for Publication

Not applicable.

Competing Interests

The authors have no competing interests to disclose.

References

1.	Brown TD, Johnston RC, Saltzman CL, Marsh JL, Buckwalter JA. Posttraumatic
	Osteoarthritis: A First Estimate of Incidence, Prevalence, and Burden of Disease.
	J Orthop Trauma. 2006 Nov;20(10):739–44. Available from:
	https://pubmed.ncbi.nlm.nih.gov/17106388/
2.	Aki T, Hashimoto K, Ogasawara M, Itoi E. A whole-genome transcriptome
	analysis of articular chondrocytes in secondary osteoarthritis of the hip. Agarwal
	S, editor. PLoS One. 2018 Jun 26;13(6):e0199734. Available from:
	https://pubmed.ncbi.nlm.nih.gov/29944724/
3.	Xu Y, Barter MJ, Swan DC, Rankin KS, Rowan AD, Santibanez-Koref M, et al.
	Identification of the pathogenic pathways in osteoarthritic hip cartilage:
	commonality and discord between hip and knee OA. Osteoarthr Cartil. 2012
	Sep;20(9):1029–38. Available from: https://pubmed.ncbi.nlm.nih.gov/22659600/
4.	Moon PM, Shao ZY, Wambiekele G, Appleton CTG, Laird DW, Penuela S, et al.
	Global Deletion of Pannexin 3 Resulting in Accelerated Development of Aging-
	Induced Osteoarthritis in Mice. Arthritis Rheumatol. 2021 May 25; Available from:
	https://pubmed.ncbi.nlm.nih.gov/33426805/
5.	Yu D, Hu J, Sheng Z, Fu G, Wang Y, Chen Y, et al. Dual roles of misshapen/NIK-
	related kinase (MINK1) in osteoarthritis subtypes through the activation of TGFβ
	signaling. Osteoarthr Cartil. 2020 Jan;28(1):112–21. Available from:
	https://pubmed.ncbi.nlm.nih.gov/31647983/
6.	Bouderlique T, Vuppalapati KK, Newton PT, Li L, Barenius B, Chagin AS.
	Targeted deletion of Atg5 in chondrocytes promotes age-related osteoarthritis.
	Ann Rheum Dis. 2016 Mar;75(3):627–31. Available from:
	https://pubmed.ncbi.nlm.nih.gov/26438374/

7.   O'Conor CJ, Ramalingam S, Zelenski NA, Benefield HC, Rigo I, Little D, et al.
Cartilage-specific knockout of the mechanosensory ion channel TRPV4
decreases age-related osteoarthritis. Sci Rep. 2016;6(July):1–10. Available from:
http://dx.doi.org/10.1038/srep29053
8.   Usmani SE, Ulici V, Pest MA, Hill TL, Welch ID, Beier F. Context-specific
protection of TGFα null mice from osteoarthritis. Sci Rep. 2016 Sep 26;6(1):1–11.
Available from: https://pubmed.ncbi.nlm.nih.gov/27457421/
9.   Loeser RF, Kelley KL, Armstrong A, Collins JA, Diekman BO, Carlson CS.
Deletion of JNK Enhances Senescence in Joint Tissues and Increases the
Severity of Age-Related Osteoarthritis in Mice. Arthritis Rheumatol. 2020 Oct
26;72(10):1679–88. Available from: https://pubmed.ncbi.nlm.nih.gov/32418287/
10. Zhang H, Wang H, Zeng C, Yan B, Ouyang J, Liu X, et al. mTORC1 activation
downregulates FGFR3 and PTH/PTHrP receptor in articular chondrocytes to
initiate osteoarthritis. Osteoarthr Cartil. 2017 Jun;25(6):952–63. Available from:
https://pubmed.ncbi.nlm.nih.gov/28043938/
11. Zhou S, Lu W, Chen L, Ge Q, Chen D, Xu Z, et al. AMPK deficiency in
chondrocytes accelerated the progression of instability-induced and ageing-
associated osteoarthritis in adult mice. Sci Rep. 2017 Apr 22;7(1):43245.
Available from: https://pubmed.ncbi.nlm.nih.gov/28225087/
12. Johnson SC, Rabinovich PS, Kaeberlin M. mTOR is a key modulator of ageing
and age-related disease. Nature. 2013;493(7432):338–45. Available from:
https://pubmed.ncbi.nlm.nih.gov/23325216/
13. Salminen A, Kaarniranta K, Kauppinen A. Age-related changes in AMPK
activation: Role for AMPK phosphatases and inhibitory phosphorylation by
upstream signaling pathways. Ageing Res Rev. 2016;28:15–26. Available from:
https://pubmed.ncbi.nlm.nih.gov/27060201/
14. Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, et al.
Rapamycin fed late in life extends lifespan in genetically heterogeneous mice.
Nature. 2009;460(7253):392–5. Available from:
https://pubmed.ncbi.nlm.nih.gov/19587680/
15. Strong R, Miller RA, Antebi A, Astle CM, Bogue M, Denzel MS, et al. Longer
lifespan in male mice treated with a weakly estrogenic agonist, an antioxidant, an
α-glucosidase inhibitor or a Nrf2-inducer. Aging Cell. 2016 Oct 16;15(5):872–84.
Available from: https://pubmed.ncbi.nlm.nih.gov/27312235/
16. Barzilai N, Crandall JP, Kritchevsky SB, Espeland MA. Metformin as a Tool to
Target Aging. Cell Metab. 2016;23(6):1060–5. Available from:
http://dx.doi.org/10.1016/j.cmet.2016.05.011
17. Lamming DW, Ye L, Katajisto P, Goncalves MD, Saitoh M, Stevens DM, et al.
Rapamycin-Induced Insulin Resistance Is Mediated by mTORC2 Loss and
Uncoupled from Longevity. Science (80-). 2012 Mar 30;335(6076):1638–43.
Available from: https://pubmed.ncbi.nlm.nih.gov/22461615/
18. Caramés B, Hasegawa A, Taniguchi N, Miyaki S, Blanco FJ, Lotz M. Autophagy
activation by rapamycin reduces severity of experimental osteoarthritis. Ann
Rheum Dis. 2012 Apr 4;71(4):575–81. Available from:
https://pubmed.ncbi.nlm.nih.gov/22084394/
19. Takayama K, Kawakami Y, Kobayashi M, Greco N, Cummins JH, Matsushita T, et

al. Local intra-articular injection of rapamycin delays articular cartilage
degeneration in a murine model of osteoarthritis. Arthritis Res Ther. 2014 Dec
17;16(6):482. Available from: https://pubmed.ncbi.nlm.nih.gov/25403236/

20.   Cheng N-T, Guo A, Cui Y-P. Intra-articular injection of Torin 1 reduces
degeneration of articular cartilage in a rabbit osteoarthritis model. Bone Joint Res.
2016 Jun;5(6):218–24. Available from:
https://pubmed.ncbi.nlm.nih.gov/27301478/

21.   Zhang Y, Vasheghani F, Li Y, Blati M, Simeone K, Fahmi H, et al. Cartilage-
specific deletion of mTOR upregulates autophagy and protects mice from
osteoarthritis. Ann Rheum Dis. 2015 Jul;74(7):1432–40. Available from:
https://pubmed.ncbi.nlm.nih.gov/24651621/

22.   Chen J, Crawford R, Xiao Y. Vertical inhibition of the PI3K/Akt/mTOR pathway for
the treatment of osteoarthritis. J Cell Biochem. 2013;114(2):245–9.

23.   Pal B, Endisha H, Zhang Y, Kapoor M. mTOR: A potential therapeutic target in
osteoarthritis? Drugs R D. 2015;15(1):27–36. Available from:
https://pubmed.ncbi.nlm.nih.gov/25688060/

24.   Chen K, Lin ZW, He S mao, Wang C qiang, Yang J cheng, Lu Y, et al. Metformin
inhibits the proliferation of rheumatoid arthritis fibroblast-like synoviocytes through
IGF-IR/PI3K/AKT/m-TOR pathway. Biomed Pharmacother. 2019;115(April
2018):1–8. Available from: https://pubmed.ncbi.nlm.nih.gov/31028998/

25.   Wang C, Yang Y, Zhang Y, Liu J, Yao Z, Zhang C. Protective effects of metformin
against osteoarthritis through upregulation of SIRT3-mediated PINK1/Parkin-
dependent mitophagy in primary chondrocytes. Biosci Trends. 2018 Dec
31;12(6):605–12. Available from: https://pubmed.ncbi.nlm.nih.gov/30584213/

26.   Petursson F, Husa M, June R, Lotz M, Terkeltaub R, Liu-Bryan R. Linked
decreases in liver kinase B1 and AMP-activated protein kinase activity modulate
matrix catabolic responses to biomechanical injury in chondrocytes. Arthritis Res
Ther. 2013;15(4):R77. Available from: https://pubmed.ncbi.nlm.nih.gov/23883619/

27.   Li J, Zhang B, Liu W-X, Lu K, Pan H, Wang T, et al. Metformin limits osteoarthritis
development and progression through activation of AMPK signalling. Ann Rheum
Dis. 2020 May;79(5):635–45. Available from:
https://pubmed.ncbi.nlm.nih.gov/32156705/

28.   Wang Y, Hussain SM, Wluka AE, Lim YZ, Abram F, Pelletier J-P, et al.
Association between metformin use and disease progression in obese people
with knee osteoarthritis: data from the Osteoarthritis Initiative—a prospective
cohort study. Arthritis Res Ther. 2019 Dec 24;21(1):127. Available from:
https://pubmed.ncbi.nlm.nih.gov/31126352/

29.   Bendele AM, Hulman JF. Effects of Body Weight Restriction on the Development
and Progression of Spontaneous Osteoarthritis in Guinea Pigs. Arthritis Rheum.
1991 Oct 7;34(9):1180–4. Available from:
https://pubmed.ncbi.nlm.nih.gov/1930336/

30.   Bendele AM, White SL, Hulman JF. Osteoarthrosis in guinea pigs: histopathologic
and scanning electron microscopic features. Lab Anim Sci. 1989;39(2):115–21.
Available from: https://pubmed.ncbi.nlm.nih.gov/2709799/

31.   Radakovich LB, Marolf AJ, Shannon JP, Pannone SC, Sherk VD, Santangelo KS.
Development of a microcomputed tomography scoring system to characterize

disease progression in the Hartley guinea pig model of spontaneous osteoarthritis. Connect Tissue Res. 2018 Nov 2;59(6):523–33. Available from: https://pubmed.ncbi.nlm.nih.gov/29226725/

32. Kraus VB, Huebner JL, DeGroot J, Bendele A. The OARSI histopathology initiative - recommendations for histological assessments of osteoarthritis in the guinea pig. Osteoarthr Cartil. 2010;18(SUPPL. 3):S35–52. Available from: http://dx.doi.org/10.1016/j.joca.2010.04.015

33. Bendele AM. Animal models of osteoarthritis. In: Journal of Musculokeletal Neuron Interaction. 2001. p. 363–76. Available from: https://pubmed.ncbi.nlm.nih.gov/15758487/

34. Radakovich LB, Marolf AJ, Culver LA, Santangelo KS. Calorie restriction with regular chow, but not a high-fat diet, delays onset of spontaneous osteoarthritis in the Hartley guinea pig model. Arthritis Res Ther. 2019 Dec 13;21(1):145. Available from: https://pubmed.ncbi.nlm.nih.gov/31196172/

35. Martin-Montalvo A, Mercken EM, Mitchell SJ, Palacios HH, Mote PL, Scheibye-Knudsen M, et al. Metformin improves healthspan and lifespan in mice. Nat Commun. 2013 Oct 16;4(1):2192. Available from: https://pubmed.ncbi.nlm.nih.gov/23900241/

36. Miller RA, Harrison DE, Astle CM, Baur JA, Boyd AR, de Cabo R, et al. Rapamycin, But Not Resveratrol or Simvastatin, Extends Life Span of Genetically Heterogeneous Mice. Journals Gerontol Ser A. 2011 Feb;66A(2):191–201. Available from: https://pubmed.ncbi.nlm.nih.gov/20974732/

37. Weiss R, Fernandez E, Liu Y, Strong R, Salmon AB. Metformin reduces glucose intolerance caused by rapamycin treatment in genetically heterogeneous female mice. Aging (Albany NY). 2018 Mar 22;10(3):386–401. Available from: https://pubmed.ncbi.nlm.nih.gov/29579736/

38. De Oliveira MA, Martins E Martins F, Wang Q, Sonis S, Demetri G, George S, et al. Clinical presentation and management of mTOR inhibitor-associated stomatitis. Oral Oncol. 2011;47(10):998–1003. Available from: https://pubmed.ncbi.nlm.nih.gov/21890398/

39. Wilkinson JE, Burmeister L, Brooks S V., Chan C-C, Friedline S, Harrison DE, et al. Rapamycin slows aging in mice. Aging Cell. 2012 Aug;11(4):675–82. Available from: http://doi.wiley.com/10.1111/j.1474-9726.2012.00832.x

40. Fraser KW. Effect of storage in formalin on organ weights of rabbits. New Zeal J Zool. 1985;12(2):169–74.

41. Tardif S, Ross C, Bergman P, Fernandez E, Javors M, Salmon A, et al. Testing Efficacy of Administration of the Antiaging Drug Rapamycin in a Nonhuman Primate, the Common Marmoset. Journals Gerontol Ser A Biol Sci Med Sci. 2015 May;70(5):577–88. Available from: https://pubmed.ncbi.nlm.nih.gov/25038772/

42. Fernandez E, Ross C, Liang H, Javors M, Tardif S, Salmon AB. Evaluation of the pharmacokinetics of metformin and acarbose in the common marmoset. Pathobiol Aging Age-related Dis. 2019;9(1):1657756. Available from: https://pubmed.ncbi.nlm.nih.gov/31497263/

43. Buie HR, Campbell GM, Klinck RJ, MacNeil JA, Boyd SK. Automatic segmentation of cortical and trabecular compartments based on a dual threshold technique for in vivo micro-CT bone analysis. Bone. 2007 Oct;41(4):505–15.

Available from: https://pubmed.ncbi.nlm.nih.gov/17693147/

44. Li Y, Yang J, Liu Y, Yan X, Zhang Q, Chen J, et al. Inhibition of mTORC1 in the
rat condyle subchondral bone aggravates osteoarthritis induced by the overly
forward extension of the mandible. Am J Transl Res. 2021;13(1):270–85.
Available from: https://pubmed.ncbi.nlm.nih.gov/33527023/

45. Xie J, Lin J, Wei M, Teng Y, He Q, Yang G, et al. Sustained Akt signaling in
articular chondrocytes causes osteoarthritis via oxidative stress-induced
senescence in mice. Bone Res. 2019 Dec 5;7(1):23. Available from:
https://pubmed.ncbi.nlm.nih.gov/31646013/

46. Aggarwal D, Fernandez ML, Soliman GA. Rapamycin, an mTOR inhibitor,
disrupts triglyceride metabolism in guinea pigs. Metabolism. 2006 Jun;55(6):794–
802. Available from: https://pubmed.ncbi.nlm.nih.gov/16713440/

47. Chen YJ, Chan DC, Lan KC, Wang CC, Chen CM, Chao SC, et al. PPARγ is
involved in the hyperglycemia-induced inflammatory responses and collagen
degradation in human chondrocytes and diabetic mouse cartilages. J Orthop Res.
2015;33(3):373–81. Available from: https://pubmed.ncbi.nlm.nih.gov/25410618/

48. Liang H, Wang H, Luo L, Fan S, Zhou L, Liu Z, et al. Toll-like receptor 4 promotes
high glucose-induced catabolic and inflammatory responses in chondrocytes in an
NF-κB-dependent manner. Life Sci. 2019;228(April):258–65. Available from:
https://pubmed.ncbi.nlm.nih.gov/30953645/

49. Ribeiro M, López de Figueroa P, Nogueira-Recalde U, Centeno A, Mendes AF,
Blanco FJ, et al. Diabetes-accelerated experimental osteoarthritis is prevented by
autophagy activation. Osteoarthr Cartil. 2016;24(12):2116–25.

50. Ribeiro M, López de Figueroa P, Nogueira-Recalde U, Centeno A, Mendes AF,
Blanco FJ, et al. Diabetes-accelerated experimental osteoarthritis is prevented by
autophagy activation. Osteoarthr Cartil. 2016;24(12):2116–25. Available from:
https://pubmed.ncbi.nlm.nih.gov/27390029/

51. Reifsnyder PC, Flurkey K, Te A, Harrison DE. Rapamycin treatment benefits
glucose metabolism in mouse models of type 2 diabetes. Aging (Albany NY).
2016;8(11):3120–30. Available from: https://pubmed.ncbi.nlm.nih.gov/27922820/

52. Watson PJ, Hall LD, Malcolm A, Tyler JA. Degenerative joint disease in the
guinea pig: Use of magnetic resonance imaging to monitor progression of bone
pathology. Arthritis Rheum. 1996;39(8):1327–37. Available from:
https://pubmed.ncbi.nlm.nih.gov/8702441/

53. Xian L, Wu X, Pang L, Lou M, Rosen C, Qui T, et al. Matrix IGF-1 regulates bone
mass by activation of mTOR in mesenchymal stem cells. Nat Med.
2012;18(7):1095–101. Available from: https://pubmed.ncbi.nlm.nih.gov/22729283/

54. Hamrick MW, Ding KH, Ponnala S, Ferrari SL, Isales CM. Caloric restriction
decreases cortical bone mass but spares trabecular bone in the mouse skeleton:
Implications for the regulation of bone mass by body weight. J Bone Miner Res.
2008;23(6):870–8. Available from: https://pubmed.ncbi.nlm.nih.gov/18435579/

55. Dawood AF, Alzamil N, Ebrahim HA, Abdel Kader DH, Kamar SS, Haidara MA, et
al. Metformin pretreatment suppresses alterations to the articular cartilage
ultrastructure and knee joint tissue damage secondary to type 2 diabetes mellitus
in rats. Ultrastruct Pathol. 2020;44(3):273–82. Available from:
https://pubmed.ncbi.nlm.nih.gov/32404018/

56. Choi YH, Sang KG, Lee MG. Dose-independent pharmacokinetics of metformin in
rats: Hepatic and gastrointestinal first-pass effects. J Pharm Sci.
2006;95(11):2543–52. Available from: https://pubmed.ncbi.nlm.nih.gov/16937336/
57. Graham GG, Punt J, Arora M, Day RO, Doogue MP, Duong JK, et al. Clinical
pharmacokinetics of metformin. Clin Pharmacokinet. 2011;50(2):81–98. Available
from: https://pubmed.ncbi.nlm.nih.gov/21241070/

757
758

Figure Legends

Figure 1: Characterization of animals on experimental diets. Food consumption (A) and bodyweight (B) of guinea pigs were recorded for the duration of the study (data presented as mean with shaded bands representing SD). Plasma glucose (C), lactate (D), and testicle weight (E) are shown. **$P<0.01$ vs Con, ***$P<0.001$ vs Con, ****$P<0.0001$ vs Con.

Figure 2: Rapamycin and rapamycin plus metformin worsened primary OA. Representative images of histology from the medial tibia are shown for each group (A; scale bars are 0.5mm and 0.25mm in 5x and 10x images, respectively). Histological images were graded for total OARSI score (B; n=7 per group). The individual scores for articular cartilage structure (C), proteoglycan loss (D) and cellularity (E) are also shown. *$P<0.05$ vs Con.

Figure 3: Proteoglycan loss correlated with hyperglycemia. Correlations between proteoglycan loss and plasma glucose (A), bodyweight and total OARSI score (B), and testicle weight and total OARSI score (C) are shown. Shaded bands represent 95% CI.

Figure 4: Rapamycin and rapamycin plus metformin inhibited mTORC1 but had no effect on mTORC2 or autophagy. IHC was performed on the medial tibia for P-RPS6 (A; n=7 per group) and quantified as percent positive cells (B) and mean integrated intensity (C). Red arrowheads indicate cells staining positive for P-RPS6. Western blot was performed on glenohumeral cartilage (D) for P-RPS6 (E), P-Akt (F), P-AMPK (G), LC3B (H), ADAMTS5 (I), and MMP-13 (J). n=8 per group for Rap and n=7 per group for Con and Rap+Met. Images are outlined in black to show that, while each band is from the same blot, bands were selected for presentation to best represent the mean change. *$P<0.05$ vs Con, **$P<0.01$ vs Con.

Figure 5: Decreased subchondral bone thickness in rapamycin and rapamycin plus metformin treated guinea pigs. Representative microCT sagittal cross sections from the medial aspect of the joint are shown (A). Subchondral cortical thickness was measured in the medial and lateral tibial plateaus and femoral condyles (B), and cortical thickness was measured in the femoral diaphysis (C). Medial tibial cortical thickness relative to femoral diaphyseal cortical thickness was found to be similar between groups (D). Medial tibial cortical thickness was highly correlated to bodyweight (E). Femoral epicondylar width was found to be similar between groups (F). N=4 per group. Shaded bands represent 95% CI. *$P<0.05$ vs Con, **$P<0.01$ vs Con.

804 **Individual Tables and Figures**
805

806 **Table 1: Average consumption of rapamycin and metformin.** Using the concentration

807 of rapamycin and metformin from the diet analysis, the average doses were calculated

808 for each group. N=7-8 per group. Data are presented as mean $\pm$ SD.

	Experimental Diet	
	Rapamycin	Rapamycin+Metformin
Rapamycin consumed (mg/kg/day)	0.72 ± 0.09	0.68 ± 0.08
Metformin consumed (mg/kg/day)	-	45 ± 5.6

831 **Table 2: Concentrations of rapamycin and metformin in circulation.** Whole blood

832 was collected ~3 hours after food had been removed from the cages of guinea pigs and

833 was analyzed for rapamycin and metformin concentration by tandem HPLC/MS. N=4 per

834 group. Data are presented as mean with $\pm$ SD.

	Experimental Diet		
	Control	Rapamycin	Rapamycin+Metformin
Circulating rapamycin (ng/mL)	0.4 ± 0	72 ± 8	78 ± 10
Circulating metformin (ng/mL)	2 ± 0	-	282 ± 54

 Figure 1

Figure 2

893 **Figure 3**

Figure 4

 Figure 5

Supplementary Material

Figure S1: OA pathology increased from 5- to 8-months of age. Total OARSI scores

are shown from the lateral tibia, medial femur, and lateral femur (A). Histological images

of knee joints from 5- and 8-month-old guinea pigs (B; scale bars are 0.5mm and 0.25mm

in 5x and 10x images, respectively) were graded for total OARSI score and individual

OARSI criteria (C). N=3 for 5-month and N=7 for 8-month. *P<0.05 vs Con.

Figure S2: Trabecular bone changes in response to experimental diets. Trabecular

thickness (A), spacing (B), and bone volume fraction (C) were measured using microCT.

N=4 per group. *P<0.05 vs Con.

Figure S3: Antibody reactivity with guinea pig articular cartilage was limited.

Immunohistochemical staining was performed, and no reactivity was observed using

primary antibodies against P-Akt Ser473 or P-AMPK Thr172.

1002
1003
1004 <u>Supplementary Figures</u>
1005 **Figure S1**

1012
1013
1014   **Figure S2**

1029

1030

1031 **Figure S3**

1032