Transcriptional and Functional Activity of Canine Hemangiosarcoma to Support Hematopoiesis Demonstrate Bone Marrow Nurse Cell Ontogeny

Jong Hyuk Kim[1-3,*], Ashley J. Schulte[1-3], Aaron L. Sarver[1,3,4], Mathew G. Angelos[5,6,7,8], Aric M. Frantz[1-3], Colleen L. Forster[10], Timothy D. O'Brien[1,3,5,9], Ingrid Cornax[1,3,9,11], M. Gerard O'Sullivan[1,3,9], Nuojin Cheng[12,13], Mitzi Lewellen[1-3], LeAnn Oseth[3], Sunil Kumar[9], Susan Bullman[14,15], Chandra Sekhar Pedamallu[14,16], Sagar M. Goyal[9], Matthew Meyerson[14,16], Troy C. Lund[17], Jessica Alfoldi[14], Kerstin Lindblad-Toh[14,18], Matthew Breen[19,20], Erin B. Dickerson[1-3], Dan S. Kaufman[3,5,6,21,22], Jaime F. Modiano[1-3,5,19, 23]

[1]Animal Cancer Care and Research Program, University of Minnesota, St Paul, MN; [2]Department of Veterinary Clinical Sciences, College of Veterinary Medicine, University of Minnesota, St Paul, MN; [3]Masonic Cancer Center, University of Minnesota, Minneapolis, MN; [4]Institute for Health Informatics, University of Minnesota, Minneapolis, MN; [5]Stem Cell Institute, University of Minnesota, Minneapolis, MN; [6]Department of Medicine (Division of Hematology, Oncology, and Transplantation), Medical School, University of Minnesota, Minneapolis, MN; [7]Microbiology, Immunology and Cancer Biology (MICaB) Graduate Program, University of Minnesota, Minneapolis, MN; [8]Internal Medicine, Hospital of the University of Pennsylvania, Philadelphia, PA; [9]Department of Veterinary Population Medicine, College of Veterinary Medicine, University of Minnesota, St Paul, MN; [10]The University of Minnesota Biological Materials Procurement Network (BioNet), University of Minnesota, Minneapolis, MN; [11]Janssen Research and Development, LLC; [12]School of Mathematics, College of Science and Engineering at the University of Minnesota, Minneapolis; [13]Applied Mathematics, University of Colorado Boulder, Boulder, CO; [14]Broad Institute of MIT and

Harvard, Cambridge, MA; [15]Human Biology Division, Fred Hutchinson Cancer Research Center,

Seattle, WA; [16]Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard

Medical School, Boston, MA; [17]Department of Pediatrics, Medical School, University of

Minnesota, Minneapolis, MN; [18]Science of Life Laboratory, Department of Medical

Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden; [19]Department of

Molecular Biomedical Sciences, College of Veterinary Medicine, & Comparative Medicine

Institute, North Carolina State University, Raleigh, NC; [20]Cancer Genetics Program, University

of North Carolina Lineberger Comprehensive Cancer Center, Raleigh, NC; [21]Center for

Immunology, University of Minnesota, Minneapolis, MN; [22]Division of Regenerative Medicine,

Department of Medicine, , La Jolla, CA; [23]Department of Laboratory Medicine and Pathology,

Medical School, University of Minnesota, Minneapolis, MN

*Corresponding author: Jong Hyuk Kim, Masonic Cancer Center, University of Minnesota,

Minneapolis, MN 55455, USA.

Tel.:+1 612-624-3612; E-mail: jhkim@umn.edu

Running Title: Bone marrow nurse cell ontogeny of hemangiosarcoma

Key Points:

• Canine hemangiosarcoma supports expansion and differentiation of human hematopoietic

progenitors, enabling development of hematopoietic tumors.

• Molecular programs of canine hemangiosarcoma that support hematopoiesis and create a

tumor immune niche resemble those of bone marrow nurse cells.

Abstract

Hemangiosarcoma and angiosarcoma are soft tissue sarcomas of malignant blood vessel-forming cells in dogs and humans, respectively. These vasoformative sarcomas are aggressive and highly metastatic, with disorganized, irregular blood-filled vascular spaces. Our objective was to define molecular programs that support the niche, enabling progression of canine hemangiosarcoma and human angiosarcoma. Here, we show that the transcriptional landscape of canine hemangiosarcoma and human angiosarcoma included comparable angiogenic and inflammatory programs. Dog-in-mouse hemangiosarcoma xenografts recapitulated the vasoformative and highly angiogenic morphology and molecular characteristics of primary tumors. Blood vessels in the tumors were complex and disorganized, and they were lined by both donor and host cells, a trait that was not observed in xenografts from canine osteosarcoma and lymphoma. In some cases, the xenografted hemangiosarcoma cells created exuberant myeloid hyperplasia and gave rise to lymphoproliferative tumors of mouse origin. We did not uncover a definitive transmissible etiology, but our data indicate that transcriptional programs of hemangiosarcoma cells resemble those of hematopoietic nurse cells, and these malignant cells support expansion and differentiation of human hematopoietic progenitors. We conclude that canine hemangiosarcoma, and possibly human angiosarcoma, originate from nurse cells that make up the stromal bone marrow niche, and that these cells may also support the growth of hematopoietic tumors.

Keywords: angiosarcoma, bone marrow, hemangiosarcoma, hematopoiesis, tumor immunity, tumor niche, tumor ontogeny.

Introduction

Canine hemangiosarcoma is a vasoformative tumor originating from bone marrow (BM)-derived progenitor cells.[1-4] Unlike human angiosarcoma, which is a rare disease, spontaneous hemangiosarcoma occurs commonly in dogs. The histology and natural history of these tumors is similar in both species: both are comprised of disorganized, tortuous, dilated blood vessels with moderate to high proliferative activity and extremely high metastatic potential.[5,6] Furthermore, canine hemangiosarcoma and human angiosarcoma appear to establish convergent molecular programs despite their genomic complexity.[7]

Angiogenesis and inflammation are key molecular features of canine hemangiosarcoma,[8] and convergent transcriptional programs regulating these processes are also observed in human angiosarcoma.[7] These molecular programs are the foundation for a novel subclassification of canine hemangiosarcoma.[4] However, the contribution of the tumor microenvironment to these programs remains incompletely understood. The microenvironment creates the tumor niche, a complex of cellular and non-cellular components that are essential for tumor cell survival, disease progression, and metastasis.[9] Emerging data suggest that cancer cells re-educate niche cells; and conversely, niche cells modulate the function of cancer cells.[9-11] This effect is especially important for sustaining stemness and self-renewal of cancer stem cells in both solid and hematopoietic tumors.[12-14]

The hematopoietic niche consists of osteoblasts, and various stromal nurse cells, including endosteal endothelial cells, fibroblasts, sinusoidal nestin-positive mesenchymal stromal cells, sinusoidal leptin receptor-positive stromal cells, and CXCL12-abundant reticular (CAR) cells.

Together, these cells support hematopoietic stem and progenitor cells (HSPCs) to proliferate and

differentiate into lineage-committed cells, promoting hematopoiesis.[15] Impairment of the

regulation of HSPCs can cause numerous blood disorders,[16,17] as well as hematopoietic

malignancies such as leukemia and lymphoma.[18-20]

Our goal was to define the molecular origin of canine hemangiosarcoma and human

angiosarcoma. Our data indicate that components of the transcriptional programs in these tumors

resemble those of bone marrow nurse cells, and that in addition to their capability to form

vasoformative tumors, canine hemangiosarcoma cells create a niche for expansion and

differentiation of blood cells.

Methods

Canine samples and cell lines

HSA cell lines (SB, COSB, Emma, DD1, JHE, and JLU) were previously established.[1,2,21-24]

Canine tissue samples were obtained from surgical removal or biopsy of tumor at the University

of Minnesota or private veterinary clinics, and additional HSA cell lines (DHSA-1401, DHSA-

1420, and DHSA-1426) were generated and cultured as described previously.[1,2,22] All protocols

and procedures for sample procurement were reviewed by the Institutional Animal Care and Use

Committee (IACUC; protocols 0802A27363, 1101A94713, 1312-31131A) of the University of

Minnesota.

Human and mouse cell lines

Human BM-derived mesenchymal stromal cells (hBM-MSCs) were isolated from whole bone

marrow purchased from AllCells (Emeryville, CA, USA) as previously described.[25-28] M2-10B4

murine bone marrow stromal cells were purchased from the ATCC (Manassas, VA, USA) and

maintained as previously described.[26] Human umbilical cord blood (hUCB) samples were

obtained from the University of Colorado Cord Blood Bank, ClinImmune Stem Cell

Laboratory.[29]

Mice and xenotransplantation

Female NOD-Scid-Gamma (NSG) and Beige-Nude-Xid (BNX) mice were used for adult

xenografts. Both male and female NSG mice were used for neonatal xenografts. All mice were

housed and handled according to the Research Animal Resources (RAR) husbandry and care

protocols. Procedures for breeding animals and for tumor implantation were reviewed by the

IACUC of the University of Minnesota (protocols 1006A84813, 1106A00649, 1306-30712A,

and 1311-31104A). We performed xenotransplantations in a total of 132 mice as detailed in

Table S1.

First, we injected cultured-tumor cells from three HSA cell lines (SB, Emma, and JHE) in 150

µL of PBS into irradiated (200gy) NSG mice (N=20): 5×10^6 SB cells were injected into the

subcutaneous (SQ) space of four mice and 5 or 10×10^6 SB cells were injected intraperitoneally

(IP) into four mice; 2×10^6 Emma cells were injected into four mice each by the SQ and IP

routes; and 3×10^5 JHE cells were injected into four mice by the SQ route. Second, we injected

tumor cells from five HSA cell lines (Emma, DD1, JLU, DHSA-1401, and COSB) into NSG

neonates (1 or 2 days after birth; N=52): 5×10^5 Emma, DD1, JLU, or DHSA-1401 cells, or

6.25×10^5 COSB cells were injected IP in 50 µL of PBS. Third, we injected 5×10^6 cells from

three HSA cell lines (JLU, DHSA-1420, and DHSA-1426) in a mixture of 100 µL of PBS and

100 µL of BD Matrigel[TM] Basement Membrane Matrix (Growth Factor Reduced; BD

Biosciences, Bedford, MA) into the SQ space of BNX mice (N=12). For DHSA-1426, we

injected tumor cells from passage-5 and passage-14, independently. We used BNX mice for

patient-derived tumor xenograft. Sections of viable tumor from four dogs affected with HSA

(DHSA-1413, DHSA-1416, DHSA-1420, and DHSA-1426) were each dissected and implanted

into subcutaneous pockets of four mice (N=16). Sections of non-HSA splenic tissues from seven

dogs were similarly implanted into 18 mice as controls. The tissues included sections from

splenic hematomas (DHSA-1412, DHSA-1417, DHSA-1419, DHSA-1421, and DHSA-1430).

Finally, after visible tumors developed in mice, we serially transplanted the tumors by

inoculation of cultured tumor cells (N=3 mice); or by direct implantation of single cell

suspensions of the tumor (N=8 mice). Mice were sacrificed when they reached a tumor endpoint,

including a mass measuring 1.5 cm in the longest diameter, or at the end of a 16-week period

after xenotransplantation.

Statistical analysis

Mann-Whitney U test or Welch's (Heteroscedastic) t-test were performed to determine

differences of continuous values between two groups. Pearson's correlation coefficient was

calculated for correlation between two variables. Statistical analysis was performed using

GraphPad Prism 6 (GraphPad Software, Inc., San Diego, CA, USA) or Microsoft Excel. Kaplan-

Meier survival analysis and log rank test were performed for survival difference using R

programming. P-values are presented without inference of significance, consistent with the

American Statistical Association's Statement on P-Values.[30]

Detailed and additional materials and methods are provided in the supplemental Methods.

Results

Canine hemangiosarcoma and human angiosarcoma show parallel transcriptional

programs and patterns of immune cell infiltration

Canine hemangiosarcomas can be classified into angiogenic, inflammatory, and adipogenic

subtypes based on their transcriptional programs (**Figure S1A**).[4,7] This separation is

prognostically significant: of the 43 dogs in this cohort for which we had treatment and outcome

data, those with inflammatory hemangiosarcomas accounted for nearly all that showed prolonged

survival (>150 days, **Figure S1B**). Intriguingly, the subset of dogs whose hemangiosarcomas had

a predominant angiogenic phenotype could be further subdivided into a group with a "pure

angiogenic" molecular signature and a group with a "mixed angiogenic and inflammatory"

molecular signature. Of these, the dogs whose tumors were purely angiogenic (*i.e.*, had deficient

inflammatory signatures), showed the shortest overall survival times (**Figure S1B**).

Consistent with previously reports,[31,32] human angiosarcomas showed highly enriched

angiogenic signatures.[7] However, the presence of inflammatory signatures in this tumor has not

been thoroughly examined. Using a defined methodology to compare transcriptional signatures

in tumors from different species,[33] we identified 588 differentially expressed genes in canine

angiogenic and inflammatory hemangiosarcomas and evaluated the expression of their homologs

in human angiosarcomas. **Figure S1C** shows that comparable angiogenic, inflammatory, and

mixed angiogenic/inflammatory gene expression signatures were present in canine

hemangiosarcomas and in human angiosarcomas.

We then used bioinformatics approaches and immunostaining to establish whether the

inflammatory gene signatures arose from the malignant cells themselves[4] or from inflammatory

cell infiltrates within the tumors. ESTIMATE was used to assign tumor purity, and xCell was

used to assign immune scores to each tumor: both tools resulted in consistent scores (**Figure**

S2A and B). Predictably, the immune score for canine samples was correlated with the

predominant transcriptional phenotype for the tumor; *i.e.*, angiogenic tumors had low immune

scores and inflammatory tumors had high immune scores (**Figure S2C and D**). There was a

similar correlation between transcriptional signatures and immune scores in human

angiosarcomas (**Figure S2E and F**). To further verify that these were the result of immune and

inflammatory cells present in the tumors, we stained sections from 11 canine hemangiosarcomas

and 10 human angiosarcomas with antibodies against T cells (CD3), B cells (Pax5), myeloid

cells (Mac387) and macrophages (Iba1 for canine; CD163 for human) (**Figure 1A and B**). CD3+

T cells, PAX5+ B cells, MAC387+ myeloid cells and Iba1+ or CD163+ macrophages were

detectable in both canine and human tumors. Myeloid cells were found most abundantly, while T

cells ranged from rare to abundant and B cells were infrequent, and when they were present, the

inflammatory cells were diffusely distributed throughout the tumor tissue. There was a direct

correlation between xCell immune scores and immunohistochemistry scores for both the canine

(Spearman R=0.38; $P = 0.255$) and human (Spearman R=0.78; $P = 0.011$) samples examined

(**Figure 1C and D**).

We next segregated human angiosarcomas into tumors with high and low immune scores

("immune-high" vs "immune-low") and identified 461 up-regulated genes (FDR $P < 0.05$) in the

immune-high group compared to immune-low group (**Figure 1E**). Fifty-eight of these genes

were also found in canine inflammatory hemangiosarcomas, and they were associated with T-

and B-cell activation (**Figure 1F and G**). The similarity in tissue organization and infiltrating

patterns of immune cells between the canine and human samples was striking, yet we were

unable to confirm the prognostic value of the immune scores in human angiosarcoma because

this sample set was not annotated with outcome data.

Canine hemangiosarcoma cells can recapitulate the disease *in vivo*

We used *in vivo* xenografts to examine the significance of the angiogenic and inflammatory

phenotypes in tumor biological behavior.[21,24,34-36] We inoculated mice with canine

hemangiosarcoma cell lines or primary tissues as described in Table S1. The engraftment

efficiency of canine hemangiosarcoma xenografts was low: in these experiments, only cultured

DHSA-1426 cells injected subcutaneously were able to reproducibly generate vasoformative

tumors in immunodeficient BNX mice (**Figure 2**), and the tumorigenic potential of this cell line

was maintained over multiple passages. These tumors could be serially passaged from cells

cultured out of the tumor xenografts and injected into new recipient mice (**Figure S3**).

Xenografts enable quantification of the stromal contribution to the hemangiosarcoma

microenvironment

To establish the contribution of stromal elements to the formation of canine hemangiosarcoma,

we used fluorescence *in situ* hybridization (FISH) to enumerate canine and mouse cells in the

tumors. We selected probes for canine *CXCL8*, since this gene is absent from the mouse genome,

and for a unique region of the mouse X chromosome, to identify canine and mouse cells,

respectively. The data show that the hemangiosarcoma xenografts achieved a complex

topological organization, with blood vessels lined concomitantly by donor and host cells (**Figure**

3A). The pattern of organization seen in canine hemangiosarcoma xenografts was distinct from

that of canine osteosarcoma and canine lymphoma xenografts (**Figure 3B**). The

hemangiosarcoma tumors were comprised of a mixture of 50-70% malignant canine cells and

30-50% mouse stromal cells. The cellular composition of orthotopic osteosarcoma xenografts

was similar, but none of the malignant canine cells were seen lining blood vessels. The

composition of lymphoma xenografts was remarkably different, including fewer than 5% mouse

stromal cells. Malignant canine cells were also absent from the blood vessels in these tumors

(**Figure 3C**).

To establish the contribution of stromal elements to the gene expression signatures of canine

hemangiosarcoma, we used a bioinformatics method that aggregates and then segregates

transcripts by species[37] and identified 459 differentially expressed mouse (stromal) genes in the

tumor xenografts (**Table S2**). We name-mapped these genes to the canine genome and examined

their expression across all of the 76 samples of primary and metastatic canine hemangiosarcoma

tumors. We then selected overlapping genes between the 1,477 subtype-specific genes that were

differentially expressed across the three canine hemangiosarcoma subtypes and the 371 mouse-

defined stromal genes (**Figure S4A**). This generated a list of 56 genes, whose expression was

also examined across the primary and metastatic canine hemangiosarcoma samples **(Figure**

S4B). Strikingly, the results from both analyses showed that the separation of tumors according

to their molecular phenotype was retained in its entirety. It suggests that the transcriptional

programs of tumor stroma are distinct among different types of hemangiosarcomas, and that

stromal elements are necessary and sufficient to define these different tumor types. However, our

experiments do not exclude the possibility that "tumor-education" is responsible for the distinct

patterns of gene expression in stromal elements.

Canine hemangiosarcoma cells can induce expansion of hematopoietic cells as well as

lymphoma cells *in vivo*

An unexpected series of results helped to inform the relationship between hemangiosarcoma and

the presence of inflammation in the microenvironment. We and others showed previously that

canine hemangiosarcoma cells could form hemangiosarcomas in NSG mice.[7,21,24,35,36] We also

observed that some of the NSG mice inoculated with SB-HSA cells or with another cell line

called Emma-brain (EFB), and some BNX mice inoculated with DHSA-1426 canine patient-

derived tumor fragments developed what appeared to be round cell tumors. Four mice that

received 5 x 10^6 SB-HSA cells intraperitoneally and one of four mice that received 2 x 10^6 EFB

cells subcutaneously died acutely two weeks after inoculation. The mice showed evidence of

anemia and splenomegaly. The spleens in all five mice were expanded by monomorphic

populations of hematopoietic cells (**Figure S5A and B**). Upon further analysis, the cells were

determined to be of mouse origin, representing erythroid progenitors (Ter-119$^+$), with few canine

hemangiosarcoma cells admixed in the population (**Figure S5C - F**). We were unable to

definitively establish that these cells had undergone malignant transformation.

Three of four mice that received DHSA-1426 tumor fragments (1st-generation canine patient -

derived xenograft or CPDX) formed apparent xenograft tumors in multiple organs including

spleen, lymph nodes, meninges, cerebrum, and mesentery 12 weeks after implantation. However,

the cellular morphology of these tumors was consistent with that of round cell tumors (**Figure**

S6A and B). We determined that the tumor cells expressed CD45 of mouse origin, B220, and

Pax5, without expression of CD3, Ter-119, and MPO, indicating a mouse B-cell origin (**Figure**

S6C - H). We verified the mouse origin using flow cytometry based on the tumor cells staining

with an antibody directed against mouse CD45, but not with antibodies directed against canine

CD45 and human $\alpha V\beta_3$-integrin, which cross reacts with canine, but not with mouse $\alpha V\beta_3$-

integrin (**Figure S7**). These B-cells could be serially passaged, forming B-cell lymphomas in

recipient BNX mice without the need for supporting canine hemangiosarcoma cells (**Figure**

S8A). Intriguingly, we observed a similar result when we inoculated single cell suspensions

derived from fresh tumor fragments of canine hemangiosarcoma xenografts into BNX recipients

(2^{nd} generation CPDX from a 1^{st} generation cell line tumor). The resulting tumors had similar

morphology and were also of mouse B-cell origin and could be serially passaged in BNX mice

(**Figure S8B**).

Canine hemangiosarcoma supports expansion and differentiation of CD34+ human

umbilical cord blood cells

The potential etiology of these tumors was perplexing. We first examined whether the tumors

might be caused by horizontal gene transfer. We used the method described above (in **Figure 3**

and **Figure S4**) to identify species-specific transcripts.[37] Canine DHSA-1426 cells expressed

only canine genes, whereas canine hemangiosarcoma xenograft tumors expressed a mixture of

canine and mouse genes (**Figure S9**). In contrast, the genes expressed by the B-cell lymphomas

arising from the hemangiosarcoma xenograft were almost exclusively of mouse origin.

Next, we sequenced viral RNA from the xenografts and used PathSeq platform to evaluate the

potential for a transmissible, infectious agent as the cause of these tumors. Notably, no bacterial

or viral sequences with tumorigenic potential were identified in the xenografts as well as in the

primary or metastatic canine hemangiosarcoma tumor samples or cell lines. A recent study

reported an association between *Bartonella spp* and canine hemangiosarcoma.[38] Ten of 24 dogs

tested in our study had detectable *Bartonella spp* sequences, although these were present in low

abundance (**Table S3**). Furthermore, four of the samples contained only sequences from *B.*

bacilliformis, B. grahamii, or *B. tribocorum*, which infect humans and rats, respectively, and

none of which is known to infect dogs as a primary or accidental host.[39,40] Three of the remaining

six dogs had sequences for *B. clarridgeiae* and three had sequences for multiple *Bartonella spp*.,

including the human and rodent-specific types. Together, the low abundance and the presence of

sequences from organisms that do not infect dogs suggest that the *Bartonella spp*. sequences

might have been contaminants.

On the other hand, murine leukemia virus (MuLV) reads were reproducibly identified in the

mouse B-cell lymphomas arising from the hemangiosarcoma xenografts (**Table S4**). The

presence of MuLV sequences was further confirmed in one mouse B-cell lymphoma arising from

the hemangiosarcoma xenografts and in two subcutaneous canine hemangiosarcoma xenograft

tumors through independent viral RNA sequencing experiments. Moreover, MuLV sequences

could be amplified by PCR from normal mouse tissues (liver and spleen), from each of three

mouse B-cell lymphomas arising from the hemangiosarcoma xenografts, and from two

subcutaneous canine hemangiosarcoma xenograft tumors. MuLV appears to be a promiscuous

virus, and we also found it in SB-HSA cells that had been previously passaged through mice as

hemangiosarcomas.

These results suggested that MuLV might be the transforming agent giving rise to the mouse

lymphomas, and pointed to a third possibility, that canine hemangiosarcoma cells were able to

provide a suitable environment for expansion of these MuLV-transformed B-cells. We thus

examined the potential for DHSA-1426 and EFB canine hemangiosarcoma cells to promote and

maintain hematopoiesis in long-term culture initiating cell (LTC-IC) assays designed to promote

expansion and differentiation of $CD34^+$ human umbilical cord blood hematopoietic progenitor

cells (HPCs). Mouse M2-10B4 and human bone marrow derived mesenchymal stromal cells

(MSCs) were used as positive controls, and HPCs cultured in the absence of feeder cells were

used as a negative control. We found that DHSA-1426 cells promoted expansion of human

$CD34^+$ HPCs with at least equal, if not better efficiency than conventional mouse or human

feeder cells, and they achieved comparable proportions of hematopoietic cell differentiation *in*

vitro across all cell lineages (**Figure 4 and Figure S10**). Similar results were obtained for EFB

cells, although the expansion and differentiation were somewhat more limited.

Canine hemangiosarcoma and human angiosarcoma establish transcriptional programs

that support hematopoietic expansion and differentiation to create a tumor immune niche

Finally, we evaluated whether hemangiosarcoma cells expressed genes that characterize distinct

stromal cells in the bone marrow niche, including sinusoidal nestin$^+$ mesenchymal stromal cells,

CXCL12-abundant reticular (CAR) cells, and sinusoidal leptin receptor (LeptinR)$^+$ stromal cells.

Figure 5 and Table S5 show that canine hemangiosarcoma tissues showed enrichment of genes

expressed by CAR cells, sinusoidal stromal cells, endosteal niche cells, endothelial progenitors,

and hematopoietic progenitors. The tumors showed higher expression of cytokines such as *CSF3*,

IL6, *IL11*, and *LIF*, associated with hematopoiesis and expansion of hematopoietic progenitors,

whereas expression of *FLT3* and *IL7*, which are also involved in hematopoiesis and lymphoid

homeostasis, were lower in hemangiosarcomas than the hematoma tissues. The tumors also

showed down-regulation of genes associated with myeloid cells and macrophages. Next, we

sought to determine if these gene signatures were also enriched in cultured hemangiosarcoma

cells (cell lines) where the tumor microenvironmental factors were depleted. **Figure 6 and Table**

S6 show that hemangiosarcoma cells up-regulate some genes associated with endothelial

progenitors and hematopoietic cytokines such as *PECAM1*, *TIE1*, *KDR*, *CD34*, *CSF3*, *IL6*, *IL11*,

and *LIF in vitro*, with significant down-regulation of *NES*. Unlike hemangiosarcoma tissues, the

hemangiosarcoma cell lines showed a higher level of genes expressed by myeloid cells and

macrophage such as *CSF2RB*, *CSF1R* and *CD68*. DHSA-1426 and EFB cells in particular,

showed remarkable enrichment of those genes, consistent with their documented capability to

support hematopoietic expansion and differentiation (**Figure 4**).

Discussion

Vasoformative sarcomas are uniformly aggressive tumors with uncertain cellular origin. In

humans, angiosarcomas are rare tumors that can arise idiopathically, or they can be secondary to

genotoxic exposures from monomeric vinyl chloride, thorium, or radiation therapy.[41,42] In dogs,

hemangiosarcomas are common and mostly idiopathic, although heritable risk factors might play

a role.[6,43] These tumors are also seen in mice after exposure to certain genotoxic agents, as well

as to non-genotoxic agents that cause severe tissue hypoxia,[44,45] and they are seen rarely in other

domestic and non-domestic species. The anatomical organization of these tumors across species

is remarkably consistent. Angiosarcomas of humans and hemangiosarcomas of dogs have been

proposed to arise from a multipotent bone marrow cell, possibly a hemangioblastic progenitor,

while hemangiosarcomas of mice may arise from lineage committed endothelial progenitor

cells.[1-4,7]

Angiosarcomas of humans and hemangiosarcomas of dogs are chemoresponsive tumors, but for

most human patients and for dogs, the benefits are transient and the prognosis for long-term

survival is poor.[46-48] A better understanding of the cell of origin and the mutational spectrum for

these tumors would help to inform the development of effective treatments for this group of

diseases.

We and others have reported on the mutational spectrum of human angiosarcomas and canine

hemangiosarcomas.[7,48-53] Human angiosarcomas and canine hemangiosarcomas have a limited

shared mutational spectrum, primarily in the canine visceral forms of the disease and in human

breast angiosarcomas.[50] But the tumors in both species, as well as in zebrafish, seem to activate

convergent transcriptional programs characterized by deregulation of phosphoinositide 3-kinase

pathways.[7,54]

Here, we document additional shared properties of human angiosarcoma and canine

hemangiosarcoma. The transcriptional landscape of these vasoformative sarcomas is strongly

pro-angiogenic; however, a subset of tumors from both species is characterized by the presence

of robust transcriptional immune and inflammatory signatures, which are proportional to the

number of detectable T cells and macrophages. Our results also suggest that at least in dogs, the

tumors with such signatures accounts for virtually all of the long-term survivors. Additional

work will be necessary to establish if the different outcomes we observed in this cohort is

generally representative of canine splenic hemangiosarcoma, where approximately 15% of dogs

with localized or regional disease are expected to survive a year or longer when treated with the

standard of care.[55,56] If this were the case, our work would provide the first potential set of

biomarkers to inform personalized treatment strategies for this disease, with surgery and

adjuvant cytotoxic chemotherapy providing adequate tumor control for some dogs with

inflammatory hemangiosarcomas, but not for dogs with angiogenic hemangiosarcomas.

The data from our canine hemangiosarcoma xenograft experiments confirm the notion that the

chaotic vascular organization is driven by the tumor cells.[1,21,24] The experiments show that the

tumor vessels are formed by a combination of malignant tumor cells and non-malignant host

endothelial cells. This observation was unique to hemangiosarcoma among the three types of

xenografts we examined, and it supports the capacity of hemangiosarcoma cells to adopt

endothelial functions. In parallel, it suggests that normal (non-malignant) cells are necessary for

the formation of malignant blood vessels in the vasculogenic tumors. Intriguingly, these

experiments also show that the stromal contribution to the angiogenic and inflammatory

transcriptional signatures is highly conserved in the xenografts, suggesting that the stromal cells

in these tumors are heavily conditioned or reprogrammed by the malignant cells. Together, the

results suggest that the interactions between the tumor and its microenvironment are rigorously

orchestrated in the formation of the hemangiosarcoma niche, highlighting a potential point of

vulnerability.

The development of exuberant myeloid and erythroid hyperplasia, as well as of bona fide

lymphomas arising from mouse cells in animals with primary or secondary hemangiosarcoma

xenografts was initially perplexing. Our results indicate that a transmissible etiology (from dog

to mouse) is unlikely to be the cause of these expanded hematopoietic cell populations. Rather,

the data indicate that at least a subset of canine hemangiosarcomas is capable of supporting

robust expansion and differentiation of hematopoietic progenitor cells *in vitro*, and we interpret

that this property is responsible for the expansion of myeloid cells *in vivo*, both in mice and in

primary canine tumors. In the case of lymphomas, which occurred repeatedly and independently

in animals receiving different tumor preparations, MuLV might have provided the driver events

for transformation of residual lymphoid elements within a hyperproliferative environment

created by the hemangiosarcoma cells. Our findings are also consistent with a report showing

angiosarcoma in the bone marrow of a human patient with tumor-associated myeloid

proliferation and extramedullary hematopoiesis.[57] The presence of transcriptional programs seen

in hemangiosarcoma cells that resemble those of CAR cells, sinusoidal stromal cells, and

endosteal niche cells, is consistent with this interpretation, and it increases the probability that

canine hemangiosarcomas – and possibly human angiosarcomas – arise from one or more of

these bone marrow nurse cells.

Other unexpected tumors have been reported in xenograft experiments and in pre-clinical models

of stem cell transplantation. For instance, transplantation of murine MSCs has been reported to

induce tumor formation and tissue malformation, potentially as a result of their possible genetic

instability and/or cellular transformation.[58-60] It has also been reported that patient-derived

xenografts of human solid cancers including breast, colon, pancreatic cancer, and

rhabdomyosarcoma induce lymphomagenesis or lymphocytic tumors in immunodeficient mice,

but in these cases, the tumors were derived from human tumor-infiltrating lymphocytes

transformed by Epstein-Barr virus.[61-64] These previously reported tumors were all of donor origin,

while the tumors in our study originated from the mouse recipients and were distinct from the

donor hemangiosarcomas. We were unable to identify and reports of hematopoietic tumors of

recipient origin arising from xenotransplantation experiments using other types of canine cancers

in the published literature, and we have not observed such an event in our own studies.[37,65,66]

Thus, this finding appears to be unique to canine hemangiosarcoma, and may be due to the

ability of hemangiosarcoma cells to support hematopoietic expansion. Our findings also suggest

that the normal counterparts of canine hemangiosarcoma cells might contribute to the

development of hematopoietic malignancies through the creation of a permissive niche.

Additionally, transdifferentiation between lymphoma and sarcoma may occur through cellular

reprogramming, potentially initiated by hematopoietic disruption.[67] In this light, it is especially

interesting that a shared region of the canine genome was found to be significantly associated

with B-cell lymphomas and hemangiosarcomas of golden retrievers.[43]

The capacity to create space for bone marrow transplants and adoptive cell therapies has taught

us that the hematopoietic niche is resilient, and that bone marrow stromal cells are highly

resistant to chemotherapy and radiation. These intrinsic properties could explain the relatively

poor long-term responses of human patients with angiosarcoma and of dogs with

hemangiosarcoma to cytotoxic therapies and may open the door to develop more effective

treatments. Nevertheless, we must recognize that therapies targeting the hematopoietic niche

might themselves carry the potential for high toxicity.

Our data provide a new model to understand the etiology and cell of origin of canine

hemangiosarcomas and possibly human angiosarcomas. We propose that the malignant cells

originate from a bone marrow nurse cell, which has the potential to create a niche that favors

angiogenic proliferation or hematopoietic expansion, as illustrated by the models in **Figure 7**.

The robust inflammation observed in some of these tumors, then, may be intrinsic to the tumor,

and not simply due to extrinsic factors associated with tissue disruption. These paths of

differentiation may also control the biological behavior of the tumors, with those showing strong

angiogenic propensity also having the most aggressive behaviors.[7]

Finally, our data do not support a transmissible etiology for hemangiosarcoma, but they do

suggest that the permissive niche created by these cells can lead to the development of

hematopoietic tumors driven by leukemia viruses in mice, raising the possibility that the bone

marrow niche plays a similar role in viral lymphomas and leukemias of humans.

Acknowledgements

The authors acknowledge Keumsoon Im for assistance with experiments and data acquisition.

The authors would also like to acknowledge Milcah Scott for processing the next generation

sequencing data and assistance with data analysis. Artistic design of Figures 2A and 7 was

created with Biorender (biorender.com). This work was partially supported by grants

1R03CA191713- 01 (to J.F. Modiano, A.L. Sarver, and J.H. Kim) from the NCI of the NIH,

grants #02759 (to J.H. Kim), #422 (to J.F. Modiano), and 1889-G (to J.F. Modiano, M.Breen,

and K. Lindblad-Toh) from the AKC Canine Health Foundation, grant JHK15MN-004 (to J.H.

Kim) from the National Canine Cancer Foundation, grant D10-501 (to J.F. Modiano and K.

Lindblad-Toh) from Morris Animal Foundation, and a grant from Swedish Cancerfonden (to K.

Lindblad-Toh). This work was also supported by an NIH NCI R50 grant, CA211249 (to A.L.

Sarver). The NIH Comprehensive Cancer Center Support Grant to the Masonic Cancer Center,

University of Minnesota (P30 CA077598) provided support for the cytogenetic analyses

performed in the Cytogenomics Shared Resource. M. Breen is supported in part by the Oscar J.

Fletcher Distinguished Professorship in Comparative Oncology Genetics at North Carolina State

University. K. Lindblad-Toh is supported by a Distinguished Professor award from the Swedish

Research Council. J.F. Modiano is supported by the Alvin and June Perlman Chair in Animal

Oncology. The authors gratefully acknowledge donations to the Animal Cancer Care and

Research Program of the University of Minnesota that helped support this project. The content of

this manuscript is solely the responsibility of the authors and does not necessarily represent the

official views of any of the funding agencies listed above.

Authorship Contributions

J.H.K., and J.F.M conceptualized the study, designed experiments, and supervised the project.

J.H.K., A.L.S., N.C., J.A., M.B., and K.L.T. analyzed RNA-seq data and provided computational

methodology. S.B., C.S.P., and M.M ran PathSeq algorithms and analyzed data. S.K., and

S.M.G. performed PCR experiment from viral RNA and analyzed MiSeq data. M.M., and M.B.

contributed to analysis and interpretation of PathSeq and MiSeq data. J.H.K, A.J.S., and A.M.F.

performed mouse experiments. M.G.A., T.C.L., D.S.K. performed LTC-IC and CFU assays and

analyzed data. A.J.S. performed flow cytometrical analysis. C.L.F. generated histological

samples and performed IHC. A.J.S., and M.L. provided administrative, technical, and material

support. J.H.K., I.C., T.D.O., and M.G.O. performed histopathological review and scoring

analysis. L.O. conducted FISH experiments. J.H.K., A.J.S., A.L.S., E.B.D., and J.F.M. wrote,

reviewed, and revised the manuscript with help from all authors. All authors read the manuscript

and approved the final draft.

Disclosure of Conflicts of Interest

No potential conflicts of interest were disclosed.

References

1. Fosmire SP, Dickerson EB, Scott AM, et al. Canine malignant hemangiosarcoma as a model

of primitive angiogenic endothelium. *Lab Invest*. 2004;84(5):562-572.

2. Lamerato-Kozicki AR, Helm KM, Jubala CM, Cutter GC, Modiano JF. Canine

hemangiosarcoma originates from hematopoietic precursors with potential for endothelial

differentiation. *Exp Hematol*. 2006;34(7):870-878.

3. Liu L, Kakiuchi-Kiyota S, Arnold LL, Johansson SL, Wert D, Cohen SM. Pathogenesis of

human hemangiosarcomas and hemangiomas. *Hum Pathol*. 2013;44(10):2302-2311.

4. Gorden BH, Kim JH, Sarver AL, et al. Identification of three molecular and functional

subtypes in canine hemangiosarcoma through gene expression profiling and progenitor cell

characterization. *Am J Pathol*. 2014;184(4):985-995.

5. Italiano A, Thomas R, Breen M, et al. The miR-17-92 cluster and its target THBS1 are

differentially expressed in angiosarcomas dependent on MYC amplification. *Genes

Chromosomes Cancer*. 2012;51(6):569-578.

6. Kim J-H, Graef A, Dickerson E, Modiano J. Pathobiology of Hemangiosarcoma in Dogs:

Research Advances and Future Perspectives. *Vet Sci*. 2015;2(4):388.

7. Kim JH, Megquier K, Thomas R, et al. Genomically Complex Human Angiosarcoma and

Canine Hemangiosarcoma Establish Convergent Angiogenic Transcriptional Programs

Driven by Novel Gene Fusions. *Mol Cancer Res*. 2021;19(5):847-861.

8. Tamburini BA, Phang TL, Fosmire SP, et al. Gene expression profiling identifies

inflammation and angiogenesis as distinguishing features of canine hemangiosarcoma. *BMC*

Cancer. 2010;10:619.

9. Quail DF, Joyce JA. Microenvironmental regulation of tumor progression and metastasis.

Nat Med. 2013;19(11):1423-1437.

10. Colmone A, Amorim M, Pontier AL, Wang S, Jablonski E, Sipkins DA. Leukemic cells

create bone marrow niches that disrupt the behavior of normal hematopoietic progenitor

cells. *Science*. 2008;322(5909):1861-1865.

11. Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in

cancer suppression and promotion. *Science*. 2011;331(6024):1565-1570.

12. Lane SW, Scadden DT, Gilliland DG. The leukemic stem cell niche: current concepts and

therapeutic opportunities. *Blood*. 2009;114(6):1150-1157.

13. Charles N, Ozawa T, Squatrito M, et al. Perivascular nitric oxide activates notch signaling

and promotes stem-like character in PDGF-induced glioma cells. *Cell Stem Cell*.

2010;6(2):141-152.

14. Vermeulen L, De Sousa EMF, van der Heijden M, et al. Wnt activity defines colon cancer

stem cells and is regulated by the microenvironment. *Nat Cell Biol*. 2010;12(5):468-476.

15. Adams GB, Scadden DT. The hematopoietic stem cell in its place. *Nat Immunol*.

2006;7(4):333-337.

16. de Haan G, Lazare SS. Aging of hematopoietic stem cells. *Blood*. 2018;131(5):479-487.

17. Filippi MD, Ghaffari S. Mitochondria in the maintenance of hematopoietic stem cells: new

perspectives and opportunities. *Blood*. 2019;133(18):1943-1952.

18. Wu JY, Purton LE, Rodda SJ, et al. Osteoblastic regulation of B lymphopoiesis is mediated

by Gs{alpha}-dependent signaling pathways. *Proc Natl Acad Sci U S A*.

2008;105(44):16976-16981.

19. Raaijmakers MH, Mukherjee S, Guo S, et al. Bone progenitor dysfunction induces

myelodysplasia and secondary leukaemia. *Nature*. 2010;464(7290):852-857.

20. Kode A, Manavalan JS, Mosialou I, et al. Leukaemogenesis induced by an activating beta-

catenin mutation in osteoblasts. *Nature*. 2014;506(7487):240-244.

21. Akhtar N, Padilla ML, Dickerson EB, et al. Interleukin-12 inhibits tumor growth in a novel

angiogenesis canine hemangiosarcoma xenograft model. *Neoplasia*. 2004;6(2):106-116.

22. Tamburini BA, Trapp S, Phang TL, Schappa JT, Hunter LE, Modiano JF. Gene expression

profiles of sporadic canine hemangiosarcoma are uniquely associated with breed. *PLoS One*.

2009;4(5):e5549.

23. Gorden BH, Kim JH, Sarver AL, et al. Identification of Three Molecular and Functional

Subtypes in Canine Hemangiosarcoma through Gene Expression Profiling and Progenitor

Cell Characterization. *Am J Pathol*. 2014;184(4):985-995.

24. Kim JH, Frantz AM, Anderson KL, et al. Interleukin-8 promotes canine hemangiosarcoma

growth by regulating the tumor microenvironment. *Exp Cell Res*. 2014;323(1):155-164.

25. Hicok KC, Thomas T, Gori F, Rickard DJ, Spelsberg TC, Riggs BL. Development and

characterization of conditionally immortalized osteoblast precursor cell lines from human

bone marrow stroma. *J Bone Miner Res*. 1998;13(2):205-217.

26. Kopher RA, Penchev VR, Islam MS, Hill KL, Khosla S, Kaufman DS. Human embryonic

stem cell-derived CD34+ cells function as MSC progenitor cells. *Bone*. 2010;47(4):718-728.

27. Zou L, Kidwai FK, Kopher RA, et al. Use of RUNX2 expression to identify osteogenic

progenitor cells derived from human embryonic stem cells. *Stem Cell Reports*.

2015;4(2):190-198.

28. Zou L, Chen Q, Quanbeck Z, Bechtold JE, Kaufman DS. Angiogenic activity mediates bone

repair from human pluripotent stem cell-derived osteogenic cells. *Sci Rep*. 2016;6:22868.

29. Ferrell PI, Hexum MK, Kopher RA, Lepley MA, Gussiaas A, Kaufman DS. Functional

assessment of hematopoietic niche cells derived from human embryonic stem cells. *Stem*

Cells Dev. 2014;23(12):1355-1363.

30. Ronald L. Wasserstein, Lazar NA. The ASA's Statement on p-Values: Context, Process, and

Purpose. *The American Statistician*. 2016;70(2):129-133.

31. Antonescu CR, Yoshida A, Guo T, et al. KDR activating mutations in human angiosarcomas

are sensitive to specific kinase inhibitors. *Cancer Res*. 2009;69(18):7175-7179.

32. Antonescu C. Malignant vascular tumors--an update. *Mod Pathol*. 2014;27 Suppl 1:S30-38.

33. Scott MC, Sarver AL, Gavin KJ, et al. Molecular subtypes of osteosarcoma identified by

reducing tumor heterogeneity through an interspecies comparative approach. *Bone*.

2011;49(3):356-367.

34. Kodama A, Sakai H, Matsuura S, et al. Establishment of canine hemangiosarcoma xenograft

models expressing endothelial growth factors, their receptors, and angiogenesis-associated

homeobox genes. *BMC Cancer*. 2009;9:363.

35. Murai A, Asa SA, Kodama A, Hirata A, Yanai T, Sakai H. Constitutive phosphorylation of

the mTORC2/Akt/4E-BP1 pathway in newly derived canine hemangiosarcoma cell lines.

BMC Vet Res. 2012;8:128.

36. Andersen NJ, Nickoloff BJ, Dykema KJ, et al. Pharmacologic inhibition of MEK signaling

prevents growth of canine hemangiosarcoma. *Mol Cancer Ther*. 2013;12(9):1701-1714.

37. Scott MC, Tomiyasu H, Garbe JR, et al. Heterotypic mouse models of canine osteosarcoma

recapitulate tumor heterogeneity and biological behavior. *Dis Model Mech*. 2016;9(12):1435-

1444.

38. Lashnits E, Neupane P, Bradley JM, et al. Molecular prevalence of Bartonella, Babesia, and

hemotropic Mycoplasma species in dogs with hemangiosarcoma from across the United

States. *PLoS One*. 2020;15(1):e0227234.

39. Chomel BB, Boulouis HJ, Maruyama S, Breitschwerdt EB. Bartonella spp. in pets and effect

on human health. *Emerg Infect Dis*. 2006;12(3):389-394.

40. Engel P, Dehio C. Genomics of host-restricted pathogens of the genus bartonella. *Genome*

Dyn. 2009;6:158-169.

41. Fayette J, Martin E, Piperno-Neumann S, et al. Angiosarcomas, a heterogeneous group of

sarcomas with specific behavior depending on primary site: a retrospective study of 161

cases. *Ann Oncol*. 2007;18(12):2030-2036.

42. Young RJ, Brown NJ, Reed MW, Hughes D, Woll PJ. Angiosarcoma. *Lancet Oncol*.

2010;11(10):983-991.

43. Tonomura N, Elvers I, Thomas R, et al. Genome-wide association study identifies shared risk

loci common to two malignancies in golden retrievers. *PLoS Genet*. 2015;11(2):e1004922.

44. Cohen SM, Storer RD, Criswell KA, et al. Hemangiosarcoma in rodents: mode-of-action

evaluation and human relevance. *Toxicol Sci*. 2009;111(1):4-18.

45. Kakiuchi-Kiyota S, Vetro JA, Suzuki S, et al. Effects of the PPARgamma agonist

troglitazone on endothelial cells in vivo and in vitro: differences between human and mouse.

Toxicol Appl Pharmacol. 2009;237(1):83-90.

46. Clifford CA, Mackin AJ, Henry CJ. Treatment of canine hemangiosarcoma: 2000 and

beyond. *J Vet Intern Med*. 2000;14(5):479-485.

47. Mullin C, Clifford CA. Histiocytic Sarcoma and Hemangiosarcoma Update. *Vet Clin North*

Am Small Anim Pract. 2019;49(5):855-879.

48. Painter CA, Jain E, Tomson BN, et al. The Angiosarcoma Project: enabling genomic and

clinical discoveries in a rare cancer through patient-partnered research. *Nat Med*.

2020;26(2):181-187.

49. Wang G, Wu M, Maloneyhuss MA, et al. Actionable mutations in canine hemangiosarcoma.

PLoS One. 2017;12(11):e0188667.

50. Megquier K, Turner-Maier J, Swofford R, et al. Comparative Genomics Reveals Shared

Mutational Landscape in Canine Hemangiosarcoma and Human Angiosarcoma. *Mol Cancer*

Res. 2019;17(12):2410-2421.

51. Wang G, Wu M, Durham AC, et al. Molecular subtypes in canine hemangiosarcoma reveal

similarities with human angiosarcoma. *PLoS One*. 2020;15(3):e0229728.

52. Beca F, Krings G, Chen YY, et al. Primary mammary angiosarcomas harbor frequent

mutations in KDR and PIK3CA and show evidence of distinct pathogenesis. *Mod Pathol*.

2020;33(8):1518-1526.

53. Chan JY, Lim JQ, Yeong J, et al. Multiomic analysis and immunoprofiling reveal distinct

subtypes of human angiosarcoma. *J Clin Invest*. 2020;130(11):5833-5846.

54. Choorapoikayil S, Weijts B, Kers R, de Bruin A, den Hertog J. Loss of Pten promotes

angiogenesis and enhanced vegfaa expression in zebrafish. *Dis Model Mech*. 2013;6(5):1159-

1166.

55. Borgatti A, Koopmeiners JS, Sarver AL, et al. Safe and Effective Sarcoma Therapy through

Bispecific Targeting of EGFR and uPAR. *Mol Cancer Ther*. 2017;16(5):956-965.

56. Borgatti A, Fieberg A, Winter AL, et al. Impact of repeated cycles of EGF bispecific

angiotoxin (eBAT) administered at a reduced interval from doxorubicin chemotherapy in

dogs with splenic haemangiosarcoma. *Vet Comp Oncol*. 2020;18(4):664-674.

57. Xie W, Lin P, Konoplev S. An unexpected diagnosis: angiosarcoma with bone marrow

involvement mimicking a myeloproliferative neoplasm. *Br J Haematol*. 2019;184(4):495.

58. Miura M, Miura Y, Padilla-Nash HM, et al. Accumulated chromosomal instability in murine

bone marrow mesenchymal stem cells leads to malignant transformation. *Stem Cells*.

2006;24(4):1095-1103.

59. Foudah D, Redaelli S, Donzelli E, et al. Monitoring the genomic stability of in vitro cultured

rat bone-marrow-derived mesenchymal stem cells. *Chromosome Res*. 2009;17(8):1025-1039.

60. Jeong JO, Han JW, Kim JM, et al. Malignant tumor formation after transplantation of short-

term cultured bone marrow mesenchymal stem cells in experimental myocardial infarction

and diabetic neuropathy. *Circ Res*. 2011;108(11):1340-1347.

61. Chen K, Ahmed S, Adeyi O, Dick JE, Ghanekar A. Human solid tumor xenografts in

immunodeficient mice are vulnerable to lymphomagenesis associated with Epstein-Barr

virus. *PLoS One*. 2012;7(6):e39294.

62. Fujii E, Kato A, Chen YJ, Matsubara K, Ohnishi Y, Suzuki M. Characterization of EBV-

related lymphoproliferative lesions arising in donor lymphocytes of transplanted human

tumor tissues in the NOG mouse. *Exp Anim*. 2014;63(3):289-296.

63. Bondarenko G, Ugolkov A, Rohan S, et al. Patient-Derived Tumor Xenografts Are

Susceptible to Formation of Human Lymphocytic Tumors. *Neoplasia*. 2015;17(9):735-741.

64. Tillman H, Vogel P, Rogers T, Akers W, Rehg JE. Spectrum of Posttransplant

Lymphoproliferations in NSG Mice and Their Association With EBV Infection After

Engraftment of Pediatric Solid Tumors. *Vet Pathol*. 2020;57(3):445-456.

65. Weiskopf K, Anderson KL, Ito D, et al. Eradication of Canine Diffuse Large B-Cell

Lymphoma in a Murine Xenograft Model with CD47 Blockade and Anti-CD20. *Cancer

Immunol Res*. 2016;4(12):1072-1087.

66. Ito D, Endicott MM, Jubala CM, et al. A tumor-related lymphoid progenitor population

supports hierarchical tumor organization in canine B-cell lymphoma. *J Vet Intern Med*.

2011;25(4):890-896.

67. Zhang Q, Orlando EJ, Wang HY, et al. Transdifferentiation of lymphoma into sarcoma

associated with profound reprogramming of the epigenome. *Blood*. 2020;136(17):1980-1983.

Figure legends

Figure 1. Immune cell infiltration and comparative immune signatures between canine hemangiosarcoma and human angiosarcoma. (A and **B)** Representative photomicrographs of H&E and immunohistochemical staining showing histological morphology and immune cell infiltration in canine hemangiosarcoma (**A**) and human angiosarcoma tissues (**B**) using anti-CD3, anti-PAX5, anti-MAC387, and anti-Iba1 (for canine) or anti-CD163 (for human) antibodies for detecting T cell, B cell, and macrophages. H&E = hematoxylin and eosin. Horseradish peroxidase or alkaline phosphate (for Iba1) conjugates were used. Counterstain = hematoxylin. Bar = 100 um. (**C** and **D**) Scatter plots display correlation between transcriptional and immunohistochemical immune score in canine hemangiosarcoma (**C**) and human angiosarcoma (**D**). Spearman's correlation coefficient (R) was calculated. (**E**) 461 upregulated genes were identified in immune-high (N = 8) compared to immune-low (N = 5) groups in human angiosarcomas (FDR P value < 0.05). (**F**) 567 immune gene signatures were identified among three molecular subtypes of canine hemangiosarcomas (N = 76; FDR P value < 0.001; fold change > 3). The heatmaps show up-regulated (red) and down-regulated (green) genes by unsupervised hierarchical clustering (average linkage; mean-centered; $\log^2$ transformed). (**G**) Venn diagram shows 58 common genes associated with signaling pathways of immune cell functions between human and canine tumors.

Figure 2. Establishment of xenografts derived from canine hemangiosarcoma in

immunodeficient mice. (**A**) Schematic illustration depicts process of tumor xenografts in beige-

nude-xid (BNX) mice. (**B**) DHSA-1426 hemangiosarcoma cells were established from canine

patient-derived tumor fragments. DHSA-1426 cells formed tumors histologically classified as

hemangiosarcoma. Left panel: H&E stain was done in xenograft tumor tissue. Right panel:

Immunohistochemistry was performed for detection of CD31 protein (alkaline phosphatase

conjugates; counterstain = hematoxylin). Top panel: 100X magnification (Bar = 100 um);

Bottom panel: 200X magnification (Bar = 50 um).

Figure 3. Organization of tumor and stromal cells in mouse xenografts of canine

hemangiosarcoma, osteosarcoma, and lymphoma. (**A**) Fluorescence *in situ* hybridization

images using canine-specific (*IL8*, red) and mouse-specific (X chromosome, green) probes in a

canine hemangiosarcoma (HSA) xenograft, a canine osteosarcoma (OS) xenograft, and a canine

lymphoma xenograft transplanted into receptive immunodeficient female mouse hosts. Red and

green arrows point to representative xenograft canine tumor cells and mouse stromal cells,

respectively, to aid in identification. (**B**) Schematic representation of A, illustrating the

organization of HSA, OS, and lymphoma xenografts. (**C**) Individual points on graph represent

relative quantity of donor (dog) and host (mouse) cells in each tumor type. 10-12 fields of

pictures at high magnification (400X) per slide were acquired. A total of approximately 1,000

cells in individual xenograft tumor was counted, and the percentages for each species-cells are

presented.

Figure 4. Canine hemangiosarcoma cells support expansion and differentiation of CD34+

human umbilical cord blood (hUCB) cells. (**A**) Flow cytometric data show populations of cells

expressing CD43 and CD45 differentiated from CD34+ hUCB cells. CD34+ hUCB cells were

pooled from two patients. M2-10B4, hBM-MSCs, and canine HSA cells (DHSA-1426 and EFB)

were seeded on gelatin-coated 24-well plates at a density of $1x10^5$ cells/well. Gelatin-coated

wells without stroma served as a negative control. Surface antigens of CD34, CD43, and CD45

were analyzed at week 5. (**B** and **C**) Bar graphs show number of different colonies formed by

hUCB CD34+ cells co-cultured with feeder cells. Both DHSA-1426 and EFB canine

hemangiosarcoma cell lines expanded hUCB CD34+ cells similar to the M2-10B4 and hBM-

MSC positive control lines, while gelatin-coated wells alone failed to support expansion. Burst-

forming unit-erythroid (BFU-E), CFU-Erythroid (CFU-E), CFU-Granulocyte/Macrophage

(CFU-GM), CFU-Macrophage (CFU-M), and CFU-

Granulocyte/Erythroid/Macrophage/Megakaryocyte (CFU-GEMM) were determined for colony-

forming unit assay.

Figure 5. Gene expression signature of hematopoietic and immune cell function enriched in

hemangiosarcoma tissues. Bar graphs show representative gene signature enrichment

associated with bone marrow niche cells, endothelial and hematopoietic progenitor, myeloid and

macrophage, signaling network, and hematopoietic cytokines between hemangiosarcoma (N =

76) and hematoma tissues (N = 10). Expression values represent count per million reads

calculated by RNA-seq transcriptomic data. CAR = CXC chemokine ligand 12 (CXCL12)-

abundant reticular; MSC = mesenchymal stromal cell; LepR = Leptin Receptor. P values and

statistical significance were determined by Mann-Whitney U test. *, P <0.05; **, P < 0.01; ***,

P < 0.001; ****, P < 0.0001.

Figure 6. Gene expression signature of hematopoietic and immune cell function enriched in

hemangiosarcoma cells. Bar graphs representative gene expression associated with bone

marrow niche cells, endothelial and hematopoietic progenitor, myeloid and macrophage,

signaling network, and hematopoietic cytokines between hemangiosarcoma cell lines (N = 11)

and hematoma cells (N = 4). Expression value represents count per million reads calculated by

RNA-seq transcriptomic data. CAR = CXC chemokine ligand 12 (CXCL12)-abundant reticular;

MSC = mesenchymal stromal cell; LepR = Leptin Receptor. P values and statistical significance

were determined by Mann-Whitney U test. *, P <0.05; **, P < 0.01; ***, P < 0.001; ****, P <

0.0001.

Figure 7. Hypothetical models for establishment of distinct molecular phenotypes of canine

hemangiosarcoma. Non-mutually exclusive models illustrate discreet cells of origin for distinct

molecular subtypes of hemangiosarcoma. Hemangiosarcoma may progress from bone marrow

nurse cells that create a niche for hematopoietic expansion and inflammation, to a transitional

pro-angiogenic state and full progression to a pure angiogenic state, or to malignant

transformation with an inflammatory phenotype.

Figure 1

Figure 2

H&E   **CD31**

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Angiogenic hemangiosarcoma

Inflammatory hemangiosarcoma

Hematopoietic progenitor

Lymphoid cell

Hemangiosarcoma cell

Macrophage

Cytokines

Neutrophil