Plant pathogenic bacterium can rapidly evolve tolerance to an antimicrobial plant allelochemical

Carrie Louise Alderley[1]#, Samuel Terrence Edwards Greenrod[1] and Ville-Petri Friman[1]

[1]University of York, Department of Biology, York, UK

Running Head: ITC tolerance in *Ralstonia solanacearum*

#Address correspondence to Carrie Louise Alderley, carrie.alderley@york.ac.uk

Abstract word-count: 266

Abstract

Crop losses to plant pathogens are a growing threat to global food security and more effective control strategies are urgently required. Biofumigation, an agricultural technique where *Brassica* plant tissues are mulched into soils to release antimicrobial plant allelochemicals called isothiocyanates (ITCs), has been proposed as an environmentally friendly alternative to agrochemicals. While biofumigation has been shown to suppress a range of plant pathogens, its effects on plant pathogenic bacteria remain largely unexplored. Here we used a laboratory model system to compare the efficacy of different types of ITCs against *Ralstonia solanacearum* plant bacterial pathogen. Additionally, we evaluated the potential for ITC-tolerance evolution under high, intermediate and low transfer frequency ITC exposure treatments. We found that allyl-ITC was the most efficient compound at suppressing *R. solanacearum* growth, and its efficacy was not improved when combined with other types of ITCs. Despite consistent pathogen growth suppression, ITC tolerance evolution was observed in the low transfer frequency exposure treatment, leading to cross-tolerance to ampicillin beta-lactam antibiotic. Mechanistically, tolerance was linked to insertion sequence movement at four positions in genes that were potentially associated with stress responses (H-NS histone like protein), cell growth and competitiveness (acyltransferase), iron storage ((2-Fe-2S)-binding protein) and calcium ion sequestration (calcium-binding protein). Interestingly, pathogen adaptation to the growth media also indirectly selected for increased ITC tolerance through potential adaptations linked with metabolism and antibiotic resistance (dehydrogenase-like protein) and transmembrane protein movement (Tat pathway signal protein). Together, our results suggest that *R. solanacearum* can rapidly evolve tolerance to allyl-ITC plant allelochemical which could

31   constrain the long-term efficiency of biofumigation biocontrol and potentially shape

32   pathogen evolution with plants.

33   **Keywords:** Antimicrobial tolerance, Plant allelochemicals, Plant pathogenic bacteria,

34   Biofumigation, Isothiocyanates (ITCs), *Ralstonia solanacearum*

Introduction

Plant pathogens are a growing threat to global food security, accounting for up to 40% of crop losses annually (Savary et al., 2012). The phasing out of environmentally toxic chemical fumigants, such as methyl bromide, has directed attention towards alternative biocontrol strategies (Qin et al., 2004). Plant-derived antimicrobial allelochemicals, such as phenolic acids, terpenes and volatile isothiocyanates (ITCs), are naturally exuded by the roots of legumes (Mondal et al., 2015; Wink, 2013), cereals (Larkin & Halloran, 2015; Mazzola & Gu, 2002) and other crops such as *Brassica* (Kirkegaard et al., 1996; Sarwar et al., 1998). These compounds could potentially be used to control pathogens by biofumigation, which involves mulching plant tissues into soils to release biocidal allelochemicals. While biofumigation has previously been shown to suppress the growth of soil-borne fungal (Angus et al., 1994; Rumberger & Marschner, 2003; Sarwar et al., 1998), nematode (Lord et al., 2011; Ngala et al., 2015) and bacterial pathogens (Hu et al., 2015; Ji et al., 2007), outcomes are still varied, ranging from clear pathogen suppression (Larkin & Griffin, 2007; Matthiessen & Kirkegaard, 2006) to having no effect (Hartz et al., 2005; Kirkegaard et al., 2000; Stirling & Stirling, 2003). A better understanding of the antimicrobial and biocidal effects of plant allelochemicals on pathogens is thus required.

The success of biofumigation is influenced by various factors including soil conditions, the biofumigant plant species, timing of application and the half-life of biocidal compounds (Matthiessen & Kirkegaard, 2006). The biocidal effects of *Brassica*-based biofumigation are believed to result primarily from the release of toxic ITCs from their glucosinolate (GSL) pre-cursors (Gimsing & Kirkegaard, 2009; Lord *et al.,* 2011; Matthiessen & Kirkegaard, 2006). Moreover, other allelochemicals such as dimethyl sulfide and methyl

iodide might contribute to the biocidal activity of biofumigant plants (Vervoort et al., 2014;

Wang et al., 2009). Even though ITC-liberating GSL levels can potentially reach as high as

45.3 mM/m^2 following initial mulching of plant material into the soil (Kirkegaard & Sarwar,

1998), their concentrations often decline rapidly due to high volatility, sorption to organic

matter, leaching from the soil and microbial degradation (Frick et al., 1998; Gimsing et al.,

2007; Hanschen et al., 2015; Matthiessen & Kirkegaard, 2006; Warton et al., 2001). As ITCs

often have short half-lives of up to sixty hours (Borek et al., 1995; Gimsing & Kirkegaard,

2006), it is important to identify ITCs that are highly effective against pathogens even during

short-term exposure.

The antimicrobial activity of different types of ITCs can vary depending on their

mode of action and the species and genotype of the target pathogen. In the case of

bacterial pathogens, several antimicrobial mechanisms have been suggested. For instance,

ITCs could damage the outer cell membrane of Gram-negative bacteria leading to changes

in cell membrane potential (Sofrata et al., 2011) and leakage of cell metabolites (Lin et al.,

2000). Further, it has been suggested that ITCs could bind to bacterial enzymes, such as

thioredoxin reductases and acetate kinases and disrupt their tertiary structure and

functioning (Luciano & Holley, 2009). It is also possible that some ITCs, such as allyl-ITC,

could have multiple targets, making them relatively more toxic to pathogenic bacteria

(Luciano & Holley, 2009). However, antimicrobial activity and potential tolerance evolution

to ITCs are still poorly understood in plant pathogenic bacteria.

Antibiosis is an important mechanism underlying bacterial competition in soils and

soil bacteria often produce and are resistant to several antimicrobials, enabling them to

outcompete surrounding bacteria for space and nutrients (Hibbing et al., 2010).

Antimicrobial tolerance is also important for plant-bacteria interactions, as it can help

bacteria to tolerate antimicrobials secreted by plants, such as coumarins, giving them a

selective advantage in the plant rhizosphere microbiome (Stringlis et al., 2018). Such

tolerance has recently been shown to evolve *de novo* in *Pseudomonas protegens* CHA0

bacterium against the antimicrobial scopoletin secreted by *Arabidopsis thaliana* (Li et al.,

2020). Prolonged exposure to plant allelochemicals could thus select for more tolerant plant

pathogen genotypes also during biofumigation and will likely be affected by the strength

and duration of ITC exposure, which is important in determining whether potential

tolerance or resistance mutations have enough time to sweep through pathogen

populations. If the mutations enabling ITC tolerance are costly, their selective benefit could

be further reduced by competition or growth trade-offs, leading to loss of tolerance

mutations in the absence of ITCs. While ITC concentrations are known to reach antimicrobial

levels during biofumigation in the field (Sarwar et al., 1998), no direct experimental

evidence for ITC tolerance evolution in plant pathogenic bacteria exists.

To study these questions, we developed a model laboratory system where we tested

the growth-inhibiting effects of ITCs produced by Indian mustard (*Brassica juncea*) on

Ralstonia solanacearum plant pathogenic bacterium, which is the causative agent of

bacterial wilt and potato brown rot diseases and a globally important pathogen, affecting

over 200 different plant species including various important crops (Elphinstone, 2005;

Yabuuchi et al., 1995). The *R. solanacearum* genome is bipartite, consisting of a

chromosome and megaplasmid (Salanoubat et al., 2002). Disease control techniques such as

crop rotation, the use of clean and certified seeds or resistant plant cultivars, have shown

only limited success in controlling *R. solanacearum* (Chellemi et al., 1997; Ciampi-Panno et

al., 1989; Ramesh et al., 2009). Indian mustard was chosen as a model biofumigant plant

due to its well-established allelochemical properties (Bending & Lincoln, 1999; Kirkegaard &

Matthiessen, 2005; Mazzola et al., 2015; Sarwar et al., 1998), which are predominantly

caused by the release of allyl, sec-butyl and 2-phenylethyl ITCs (Bangarwa et al., 2011;

Olivier et al., 1999; Yim et al., 2016). As these ITCs might vary in their biocidal activity, we

first tested to what extent they suppress *R. solanacearum* growth when applied either alone

or in combination at concentrations relevant to field biofumigation (Gimsing et al., 2007;

Hanschen et al., 2012; Kirkegaard & Sarwar, 1998; Matthiessen & Kirkegaard, 2006; Rudolph

et al., 2015). Subsequently, we explored whether long-term exposure to the most effective

ITC type could select for resistant or more ITC-tolerant pathogens in the lab, and if ITC

tolerance is associated with competitive costs or cross-tolerance to other antimicrobials. It

was found that allyl-ITC was the most suppressive allelochemical. However, long-term

exposure selected for ITC-tolerant pathogen mutants that also had increased cross-

tolerance to the beta-lactam antibiotic ampicillin. At the molecular level, adaptations were

associated with a few parallel mutations and loss of insertion sequences mainly in the

megaplasmid. Together these results suggest that while Indian mustard could be used as a

biofumigant plant against *R. solanacearum* due to the high antimicrobial activity of allyl-ITC,

its long-term efficacy could be constrained by rapid ITC tolerance evolution.

Materials and Methods

(a) Pathogen strain and culture media

We used a *Ralstonia solanacearum* strain (21415687) which was originally isolated from the

river Loddon (phylotype II sequevar 1) in the UK as our ancestral pathogen strain (Source:

John Elphinstone, Fera Science, 2014). This strain was chosen as river water is the most

common environmental source of potato brown rot outbreaks in the UK (Elphinstone *et al.*,

1998), and hence highly relevant for UK *R. solanacearum* epidemics. The strain was cultured

in CPG broth (1 g casamino acids, 10 g peptone and 5 g glucose per litre of ddH$_2$O) for 48

hours at 28 °C to create cryostocks (20% w/v glycerol) that were preserved at -80 °C. CPG

was also used as the main growth media in all experiments except for fitness assays, where

lysogeny broth (LB: 10 g tryptone, 5 g yeast, 10 g NaCl per litre of ddH$_2$O) was also used as a

'naïve' growth media to control the effects of *R. solanacearum* adaptation to CPG media

during the selection experiment.

(b) Comparing the effects of different types of ITCs for pathogen suppression

To determine antimicrobial activity of ITCs, we first identified concentrations that caused a

significant reduction in *R. solanacearum* growth relative to the no-ITC control treatments.

To this end, we conducted short-term growth assays where *R. solanacearum* was exposed

to allyl, sec-butyl and 2-phenylethyl ITCs at 63, 125, 250, 500, 1000, 2500 and 5000 µM

concentrations in CPG media (Suppl. Fig. 2). For this experiment, *R. solanacearum* was

revived from cryostocks by growing with shaking (250 rpm) for 48 hours at 28 °C before

normalising bacterial density to an optical density (OD) reading of 0.1 (600 nm; Tecan,

Sunrise), equalling ~10^7 cells per ml. This method was consistently used to revive and adjust

bacterial densities in all growth experiments. *R. solanacearum* was grown in 200 µl CPG

media in different ITC concentrations for 148 hours and bacterial densities were measured

every 24 hours (OD600 nm). We found that allyl-ITC concentrations as low as 125 µM

inhibited *R. solanacearum* growth, while relatively higher concentrations of 500 µM of sec-

butyl and 2-phenylethyl ITC were required to inhibit pathogen growth (Suppl. Fig. 2). Based

on this data, 500 µM and 1000 µM ITC concentrations were selected because they showed

pathogen growth suppression in the case of all measured ITCs (Suppl. Table 1). Furthermore,

these concentrations are known to be achievable at least transiently during biofumigation in

the field (Gimsing et al., 2007; Hanschen et al., 2012; Kirkegaard & Sarwar, 1998;

Matthiessen & Kirkegaard, 2006; Rudolph et al., 2015). To explore the effects of ITCs on

pathogen growth alone and in combination, different ITCs were mixed in all possible two-

way and three-way combinations using equal concentrations of each ITC within

combinations (two-way 50:50%; three-way 33:33:33%) to achieve final low (500 μM) and

high (1000 μM) ITC concentrations in 200 μl of CPG media in 96-well microplates.

Microplates were cultured at 28 °C (N= 8 for all treatments) and the experiment was run for

three days (72 hours), with population density measurements recorded every 24 hours as

optical density at 600 nm.

(c) Determining pathogen ITC and beta-lactam tolerance evolution in response

to repeated allyl-ITC exposure

To investigate the potential for ITC tolerance evolution, we set up a 16-day selection

experiment where we exposed *R. solanacearum* to 500 μM of allyl-ITC, which has the

strongest effect on pathogen growth suppression of all tested ITCs (Fig. 1A; Suppl. Fig. 2).

We also manipulated the frequency of ITC exposure using high (1-day), intermediate (2-day)

and low (3-day) serial transfer frequency treatments. At each serial transfer, a subset of

evolved bacteria (5% of the homogenised bacterial population) was serially transferred to

fresh CPG media in the absence (control) and presence of allyl-ITC. ITC treatments thus

manipulated both resource renewal and exposure to fresh ITC. The selection experiment

was set-up following the same protocols described earlier and following this, separate

fitness assays were conducted to directly compare the growth of ancestral and evolved

populations (and individual colonies) in the absence and presence of 500 μM allyl-ITC. In

addition to testing potential ITC tolerance evolution, we quantified changes in the growth of

evolved bacteria in the absence of ITCs to reveal potential adaptations to the CPG growth

media. All fitness assays were also repeated in 'naïve' LB media to control the potential

effects of pathogen adaptation to the CPG growth media during the selection experiment. In

all assays, bacteria were revived and prepared as described earlier, and grown in 96-well

microplates in different media (CPG or LB) in the absence or presence of 500 μM allyl-ITC for

72 hours. Changes in ITC tolerance were quantified as bacterial growth relative to the

ancestral and control treatments based on optical density at 600 nm (48-hour time point).

Fitness assays were also conducted for individual bacterial colonies at the final time point

where a single ancestral colony and one colony from each replicate selection line per

treatment were selected resulting in a total of 49 clones.

To explore potential ITC-tolerance mechanisms, we tested if ITC tolerance correlated

with tolerance to ampicillin beta-lactam antibiotic (growth assays), which is commonly

produced by various soil bacteria (Ranjan et al., 2021). Moreover, we specifically tested for

ampicillin tolerance as we identified potential antibiotic-linked insertion sequence

movement in our evolved clones, which has previously been shown to confer beta-lactam

antibiotic tolerance in clinical settings (Boutoille et al., 2004; Poirel et al., 2003). Ampicillin

tolerance was tested using the sequenced isolated clones from the final time point of the

selection experiment (intermediate transfer frequency no-ITC, low transfer frequency no-

ITC and low transfer frequency ITC exposure treatments) and the ancestral strain (total of 24

evolved clones and 8 replicate ancestral clones per treatment). Clones were prepared as

described earlier and grown in 96-well microplates in CPG media in the absence or presence

of 15 or 30 μg/ml ampicillin. Ampicillin tolerance was quantified as bacterial growth relative

to the ancestral clones based on optical density at 600nm (48-hour time point).

(d) Genome sequencing of evolved bacterial clones

A subset of evolved clones was whole genome sequenced to identify potential single

nucleotide polymorphisms (SNPs), genomic rearrangements (small insertions and deletions)

and potential changes in prophage and insertion sequence movement linked with *R.*

solanacearum adaptation. Based on phenotypic data, we chose eight clones (1 per replicate

selection line) from the low transfer frequency treatments that had evolved in the absence

or presence of ITC (16 clones). Moreover, we sequenced the ancestral strain (1 clone) and

eight clones from the intermediate transfer frequency no-ITC treatment (8 clones), that

showed no evidence of ITC tolerance adaptation (a total of 25 clones), as controls. Genomic

DNA was extracted using the Qiagen DNeasy UltraClean Microbial Kit according to the

manufacturer's protocol. DNA was quantified using the NanoDrop microvolume

spectrophotometer and quality checked by gel electrophoresis imaging. DNA yields of all

samples were diluted with EB buffer to 30 ng/μl concentrations and DNA samples were sent

to MicrobesNG for sequencing (Illumina 30x coverage; http://www.microbesng.uk).

MicrobesNG conducted library preparation using Nextera XT Library Prep Kit (Illumina, San

Diego, USA) following the manufacturer's protocol with the following modifications: 2 ng of

DNA were used as input, and PCR elongation lasted 1 min. Hamilton Microlab STAR

automated liquid handling system was used for DNA quantification and library preparation.

Pooled libraries were quantified using the Kapa Biosystems Library Quantification Kit for

Illumina on a Roche light cycler 96 qPCR machine. Libraries were sequenced on the Illumina

HiSeq 2500 using a 250 bp paired end protocol. Reads were adapter trimmed using

Trimmomatic 0.30 with a sliding window quality cut-off of Q15 (Bolger et al., 2014).

Assembly was performed on samples using SPAdes v.3.7 (Bankevich et al., 2012) and contigs

were annotated using Prokka v.1.11 (Seemann, 2014). Genomes were analysed using a

standard analysis pipeline (Guarischi-Sousa et al., 2016), where reads were first mapped to a

high quality and well annotated UY031 reference genome (NCBI accession: NZ_CP012687)

which showed 99.95% similarity with our ancestral *R. solanacearum* strain at the

chromosome level and 97.87% similarity at the mega-plasmid level. Variant calling was

performed using Snippy v.3.2, a rapid haploid variant calling pipeline (Seemann, 2015).

When comparing the sequenced genomes, the SNPs identified in both the ancestral strain

and the evolved clones were first filtered out as these likely represent pre-existing

phylogenetic differences between the reference genome and our ancestral *R. solanacearum*

strain. We also compared the control treatment clones isolated from low and intermediate

transfer frequency treatments (no ITC exposure) to identify potential mutations linked with

CPG media adaptation. The software IMSindel v.1.0.2 (Shigemizu et al., 2018) was used to

identify potential intermediate indels with options "—indelsize 10000" and using UY031 as a

reference. After running IMSindel, putative indels in all isolates were combined. Putative

short indels that were < 50 bp in length were removed. To investigate potential insertion

sequences underlying ITC tolerance and media adaptation, insertion sequences were

detected in the UY031 with ISEScan v.1.7.2.3; (Xie & Tang, 2017) using default parameters.

Potential false positives were determined by blasting insertion sequences against the

ISFinder database (https://isfinder.biotoul.fr/) and removing hits with an E-value > e-04.

Experimental isolates were then screened for the insertion sequences identified with

ISEScan using ISMapper v.2.0; (Hawkey et al., 2015) with default settings. In line with a

previous study (Hawkey et al., 2020), ISMapper was run using an IS-removed UY031

assembly to improve insertion site precision. The genes flanking putative IS sites were

determined by annotating the UY031 assembly using the stand-alone NCBI prokaryotic

genome annotation pipeline 2021-07-01.build5508 (Tatusova et al., 2016). Additionally, we

determined isolate prophage content and positions to identify potential phenotypic changes

via mobile genetic elements. Isolate draft assemblies were generated using Unicycler

Illumina-only assembly v.0.4.7 (Wick et al., 2017). Prophages were then identified in draft

assemblies using the PHASTER (PHAge Search Tool Enhanced Release) web server (Arndt et

al., 2016). Prophage movement was detected by parsing out the 5kb (or to end of contig)

flanking regions either side of the prophages in the draft assemblies and mapping them to a

closely related complete UY031 genome sequence. Prophage movement was detected if the

flanking regions map to different parts of the UY031 genome between isolates. Prophage

movement analyses were conducted using custom R and Python scripts available at

(https://github.com/SamuelGreenrod/Prophage_movement). All genomes including the

ancestral strain have been deposited in the European Nucleotide Archive database under

the following accession number: PRJEB42551.

(e) Statistical analysis

Repeated measures ANOVA was performed to analyse all the data with temporal sampling

structure and pairwise differences were determined using *post-hoc* t-test with Bonferroni

correction. All other statistical analyses (ITC tolerance and cost of tolerance in CPG and LB

media and cross-tolerance in ampicillin) were conducted focusing on the 48-hour

measurement time point (where ITC was still actively suppressive to *R. solanacearum,* Suppl.

Fig. 1) and two-way ANOVA was used to explain variation in bacterial growth between

different treatments. Tukey *post-hoc* tests were used to compare differences between

subgroups (p< 0.05). Where data did not meet the assumptions of a parametric test, non-

parametric Kruskal-Wallis test and *post-hoc* Dunn test were used. All statistical analyses and

graphs were produced using R (R Foundation for Statistical Computing, R Studio v.3. 5. 1)

using ggplot2, tidyverse, ggpubr, lme4, rcompanion and reshape2 packages.

Results

(a) Only allyl-ITC suppressed pathogen growth irrespective of the presence of other ITCs

We first determined the effects of different ITCs on *R. solanacearum* growth alone and in combination. Overall, there was a significant reduction in *R. solanacearum* densities in the presence of ITCs (ITC presence: $F_{1, 120}$= 6.33, p< 0.01; Tukey: p< 0.05; Fig. 1B). However, this effect was mainly driven by the allyl-ITC, which significantly reduced bacterial densities compared to the no-ITC control treatment (ITC type: $F_{7, 114}$= 49.45, p< 0.001; Tukey: p< 0.05), while other ITCs had no significant effect on the pathogen (p> 0.05; Fig. 1B). Increasing the ITC concentration from low to high (500 to 1000 μM) had no effect on inhibitory activity in either single or combination ITC treatments (ITC concentration in single ITC treatment: $F_{1, 43}$= 2.0, p= 0.17; combination ITC treatment: $F_{1, 59}$= 0.68, p= 0.41; Fig. 1B). However, a significant interaction between ITC type and ITC concentration in both single and combination treatments was found (ITC concentration × ITC type in single ITC treatment: $F_{2, 39}$= 4.67, p< 0.05; in combination ITC treatment: $F_{3, 53}$= 4.94, p< 0.01; Fig. 1B), which was driven by the increased inhibitory activity of allyl-ITC at high concentration (Tukey: p< 0.05). As a result, ITC combinations were less inhibitory than single ITC treatments (Number of ITCs: $F_{2, 103}$= 3.82, p<0.05; Fig. 1B), which was due to reduced allyl-ITC concentration in combination treatments (total ITC concentrations were kept the same between treatments). Similarly, ITC combinations that included allyl-ITC significantly reduced bacterial densities relative to the control treatment (Allyl-ITC presence: $F_{1, 57}$= 36.21, p< 0.001; Fig. 1B), and the presence of allyl-ITC had a clearer effect at the high ITC concentration (Allyl-ITC presence × ITC concentration: $F_{1, 57}$= 7.51, p< 0.01; Fig. 1B). Together

these results suggest that allyl-ITC was the most inhibitory compound and its antimicrobial

activity was not enhanced by the presence of other ITCs.

(b) Pathogen growth was more clearly suppressed in high and intermediate

ITC exposure treatments during an experimental evolution experiment

To study the evolutionary effects of ITCs, we exposed the ancestral *R. solanacearum* strain

to allyl-ITC at the low concentration (500 µM) and manipulated the frequency of exposure

to ITC by transferring a subset of evolved bacterial population to fresh ITC-media mixture

everyday (high), every second day (intermediate) and every third day (low) for a total of 16

days. As a result, this manipulation also affected the resource renewal rate. Overall, bacteria

reached the highest population densities in the low transfer frequency treatments and the

second highest in the intermediate transfer frequency treatments (Transfer frequency: $F_{2,}$

$_{45}$= 4.66, p< 0.001; p< 0.05 for pairwise comparison; Fig. 2). While allyl-ITC exposure

significantly reduced bacterial densities in all ITC-containing treatments (ITC presence: $F_{1,46}$=

30.68, p< 0.001; Fig. 2), bacterial growth was least affected in the low transfer frequency

treatment (ITC presence × Transfer frequency: $F_{2,42}$= 4.36, p< 0.05; p< 0.001 for all pairwise

comparisons; Fig. 2). The inhibitory activity of allyl-ITC also varied over time: while relatively

more constant suppression was observed in the high and intermediate transfer frequency

treatments, pathogen growth suppression became clear in the low transfer frequency

treatment only towards the end of the selection experiment potentially due to media

growth adaptation in the no-ITC control treatment (Time × Transfer frequency × ITC

presence: $F_{2,673}$= 7.33, p< 0.001; Fig. 2). Together these results suggest that the long-term

ITC activity varied temporally and depended on the ITC exposure and serial transfer

frequency.

(c) ITC tolerance evolution was observed only in the low transfer frequency ITC exposure treatment

Fitness assays were conducted at the end of the selection experiment to compare the growth of the ancestral strain and evolved populations from different treatments in the presence and absence of allyl-ITC (experimental concentration: 500 μM). The ancestral strain reached lower densities in the presence of ITC compared to evolved populations regardless of the ITC treatment they had evolved in during the selection experiment (Evolutionary history: $F_{2, 45}$= 5.39, p< 0.01; Tukey: p< 0.05; Fig. 3A). However, ITC tolerance was mainly observed in the low transfer frequency ITC exposure treatment, while populations that had evolved in the high or intermediate transfer frequency treatments did not significantly differ from the ancestral strain (Transfer frequency within ITC-exposed populations: $F_{2, 19}$= 24.72, p< 0.001; Tukey: p< 0.05; Fig. 3A). Surprisingly, even the control populations that had evolved in the absence of ITCs in the low transfer frequency treatment showed an increase in ITC tolerance (p< 0.05; Fig. 3A). One potential explanation for this is that these populations adapted to grow better in CPG media, which could have helped to compensate for the mortality imposed by allyl-ITC during the fitness assays. To test this, we compared the growth of ancestral and evolved populations in the absence of allyl-ITC in the CPG media (Fig. 3B). We found that all control populations showed improved growth in the CPG media compared to ITC-exposed populations regardless of the transfer frequency treatment (Evolutionary history: $F_{1, 40}$= 20.00, p< 0.001; Transfer frequency: $F_{2, 40}$= 2.66, p= 0.08, in all pairwise comparisons, Tukey: p< 0.05; Fig. 3B). In contrast, none of the ITC-exposed populations showed improved growth in CPG media relative to the ancestral strain (Tukey: p< 0.05; Fig. 3B), which suggests that ITC exposure constrained *R. solanacearum* adaptation to the growth media.

To disentangle the effects due to adaptation to the media and allyl-ITC, we repeated

fitness assays in 'naïve' LB growth media which the bacteria had not adapted to. ITC

tolerance was observed only when bacterial populations had previously been exposed to

allyl-ITC (Evolutionary history: $F_{2, 49}$= 18.82, p< 0.001; Tukey: p< 0.05; Fig. 3C), and this effect

was driven by adaptation in the low transfer frequency ITC exposure treatment (no ITC

tolerance was observed in the high and intermediate transfer frequency treatment; Transfer

frequency: $F_{2, 49}$= 4.37, p< 0.01; Tukey: p< 0.05; Fig. 3C). Crucially, CPG-adapted control

populations showed no signs of ITC tolerance, but instead, suffered reduced growth in LB

media relative to the ancestral strain and ITC-exposed populations (Evolutionary history: $F_{2,}$

$_{49}$= 94.89, p< 0.001; Fig. 3D), which was clearest in the low transfer frequency exposure

treatment (Evolutionary history × Transfer frequency: $F_{2, 49}$= 23.17, p< 0.001; Fig. 3D).

We further validated our population level fitness results using individual clones (one

randomly chosen clone per replicate population per treatment). In line with previous

findings, ITC-exposed clones showed increased ITC tolerance compared to the control and

ancestral bacterium in the LB media (Evolutionary history: $F_{2, 49}$= 14.20, p< 0.001; Fig. 4A),

and tolerance evolution was the greatest in the low transfer frequency ITC exposure

treatment (Transfer frequency: $F_{2, 49}$= 11.15, p< 0.001; Tukey: p< 0.05; Evolutionary history ×

Transfer frequency: $F_{2, 49}$= 3.04, p< 0.05; Fig. 4A). Together, our results suggest that ITC

tolerance, which evolved in the low transfer frequency ITC exposure treatment was robust

and independent of the growth media it was quantified in. Moreover, while all control

populations adapted to grow better in the CPG media, this adaptation had a positive effect

on ITC tolerance only when quantified in CPG media and when the clones had evolved in the

low transfer frequency treatment.

(d) Evolution of ITC-tolerance confers cross-tolerance to ampicillin beta-

lactam antibiotic

We also tested if exposure to allyl-ITC could have led to cross-tolerance to other

antimicrobials such as the beta-lactam antibiotic ampicillin. Overall, both low (15 µg/ml) and

high (30 µg/ml) ampicillin concentrations had negative effects on *R. solanacearum* growth

relative to the no-ampicillin control treatment (Ampicillin concentration: $F_{2, 93}$= 50.12, p<

0.001; Tukey: p< 0.05; high concentration was relatively more inhibitory, Suppl. Fig. 3).

However, the evolved clones from the low transfer frequency ITC exposure treatment

reached significantly higher bacterial densities than the ancestral strain (Evolutionary

history: $F_{3, 92}$= 3.51, p< 0.05; Tukey: p< 0.05; Suppl. Fig. 3), while evolved clones derived from

low and intermediate transfer frequency control treatments (no prior ITC exposure) did not

differ from the ancestral strain (Tukey: p> 0.05; Suppl. Fig. 3). Ampicillin tolerance was only

observed in the high ampicillin concentration (High ampicillin concentration: $F_{3, 28}$= 8.22, p<

0.001; Suppl. Fig. 3C; Low ampicillin concentration: $F_{3, 28}$= 1.551, p= 0.223; Suppl. Fig. 3B).

Together these results suggest that ITC tolerance conferred cross-tolerance to ampicillin for

clones that had evolved in the low transfer frequency ITC exposure treatment.

(e) Media adaptation and ITC tolerance are linked to a few mutations and loss

of insertion sequences

A subset of clones which were phenotyped regarding ITC and ampicillin tolerance were

selected for genome sequencing (N=25). All isolated colonies showed ancestral, fluid colony

morphotype with no evidence for spontaneous evolution of small colony types as observed

previously (Khokhani et al., 2017; Perrier et al., 2019). Specifically, we focused on comparing

parallel small mutations and indels, intermediate indels (>50 bp) and prophage and

insertion sequence (IS) movement between populations that had evolved in the absence

and presence of ITC in the low transfer frequency treatments (evidence of ITC tolerance

evolution) with ancestral and control populations from the intermediate transfer frequency

treatment (no ITC tolerance evolution observed). Potential genetic changes were

investigated in both the chromosome and megaplasmid of the bipartite genome.

Only a few mutations were observed in 1 to 6 different genes, which was expected

considering the relatively short duration of the selection experiment (16 days). Of these

mutations, 8 were non-synonymous and 4 synonymous (Table 1). Some mutations were

observed across all treatments, indicative of adaptation to the culture media or other

experimental conditions. For example, parallel non-synonymous mutations in *hisH1* gene

controlling imidazole glycerol phosphate synthase were observed in 6/8 to 8/8 replicate

clones in all treatments (Table 1; Fig. 5). Similarly, non-synonymous mutations in

serine/threonine protein kinase genes (between 5/8 to 8/8 replicate clones) and

synonymous mutations in putative deoxyribonuclease *RhsC* gene (between 1/8 to 5/8

replicate clones) were found across all treatments (Table 1; Fig. 5). A single clone that had

evolved in the absence of allyl-ITC in the intermediate transfer frequency treatment had a

unique non-synonymous mutation in the gene encoding the putative HTH-type

transcriptional regulator *DmlR* and another clone originating from this treatment had a

mutation in the IS5/IS1182 family transposase encoding gene (Table 1; Fig. 5). Additionally,

we observed mutations exclusively in the low transfer control clones in genes encoding the

dehydrogenase-like uncharacterised protein (3/8 replicate clones) and Tat pathway signal

protein (2/8 replicate clones; Table 1; Fig. 5), which may explain ITC tolerance via media

adaptation. However, no clear parallel mutations exclusive to the low frequency ITC-

exposed populations were found.

In terms of putative intermediate indels (>50 bp), we identified 122 and 116 indel

sites in the chromosome (Chr) and megaplasmid (MP), respectively. Almost all of these were

insertions (Chr, 119/122; MP, 113/116) and the majority were singletons (Chr, 101/122; MP,

95/116) or doubletons (Chr, 14/122; MP, 13/116). The number of intermediate indels did

not differ between evolutionary treatments either in the chromosome (Kruskal-Wallis: x^2=

3.65; df= 2; p= 0.161) or megaplasmid (Kruskal-Wallis: x^2= 3.46; df= 2; p= 0.178). As a result,

this genetic variation was likely non-adaptive and driven by random drift.

To identify other potential molecular mechanisms, variation in prophages and

insertion sequences (ISs) was investigated. Two prophages were found in all sequenced

isolates: Inoviridae prophage φRS551 and a novel, unclassified prophage (Table S2).

Prophage genome positions were almost identical between all sequenced isolates (Table

S2). Therefore, no evidence for systematic prophage movement was observed in the

evolved isolates relative to the ancestral strain. In contrast, ISs appeared to be highly mobile

regarding 15 variable positions in the chromosome and 15 variable positions in the

megaplasmid (Suppl. Fig. 5). In most variable positions (7 in the chromosome and 9 in the

megaplasmid), the gain or loss of ISs was infrequent, occurring in up to three clones per

treatment (Suppl. Fig. 5), which is indicative of non-adaptive, random IS movement.

However, the remaining IS positions showed higher frequency of gain or loss, indicating of

potentially adaptive IS movement which was also in some cases treatment-specific. For

example, an IS element in position 2302900 on the chromosome absent in the ancestral

strain was observed in 2 clones in the intermediate transfer frequency control and 2 low

transfer ITC treatment clones, while it was gained by 5 clones in low transfer control

treatment. The IS element in this position was found to be close to the start codon (~50 bp)

of an acyltransferase. In two of the low transfer control clones, the IS was found to disrupt

the gene (Fig. 5), potentially knocking out acyltransferase gene expression after inserting

into this position. Moreover, three IS elements in the megaplasmid were almost exclusively

lost in the low transfer frequency treatment (Fig. 5). In one of the positions (209500), the IS

disrupted a putative calcium-binding protein in the intermediate transfer control clones but

was absent in 4/8 low transfer control and 4/8 low transfer ITC treatment clones. In the

other two positions (243500 and 253900), the ISs were intergenic (positioned 450 bp and

104 bp (243500) and 301 bp and 46 bp (253900) from their left- and right-flanking genes;

Fig. 5). The right-flanking genes closest to the ISs included a (2Fe-2S)-binding protein

(243500) and an H-NS histone family protein (253900), whilst the left-flanking genes

included the type III effector *HopG1* (243500) and an unknown hypothetical protein

(253900). The frequency of IS absence in these positions (243500 and 253900) differed

between low transfer treatments. Specifically, in position 243500, the IS was absent in 7/8

low transfer control and 5/8 low transfer ITC treatment clones. Meanwhile, in position

253900, the IS was absent in 4/8 low transfer control and 6/8 low transfer ITC treatment

clones. However, despite these patterns, the extent of IS loss did not differ statistically

between low transfer control and ITC-exposed clones when analysed individually (Mann-

Whitney: 209500: w= 32, n1= 8, n2= 8, p= 1; 243500: w= 40, n1= 8, n2= 8, p= 0.29; 253900:

w= 24, n1= 8, n2= 8, p= 0.35) or in combination (Mann-Whitney: w= 32, n1= 8, n2= 8, p= 1).

Together, these results suggest that media adaptation and ITC tolerance was potentially

driven by parallel mutations in a few genes and more frequent loss of IS elements in the low

transfer frequency treatments.

Discussion

Here we studied the effects of *Brassica*-derived ITC allelochemicals for the suppression and tolerance evolution of plant pathogenic *R. solanacearum* bacterium in a model biofumigation experiment. We found that only allyl-ITC suppressed *R. solanacearum* growth, while no reduction in pathogen densities were observed when sec-butyl and 2-phenylethyl ITCs were applied alone or in combination. By using experimental evolution, we further showed that long-term allyl-ITC exposure selected for ITC tolerance in the low transfer frequency ITC exposure treatment and was associated with cross-tolerance to ampicillin. At the genetic level, tolerance evolution was associated with the loss of IS elements. Together, our results suggest that allyl-ITC derived from Indian mustard is effective at suppressing the growth of the *R. solanacearum* pathogen *in vitro*. However, prolonged exposure could select for increased ITC tolerance, potentially reducing the efficiency of ITC-based biocontrol.

Only allyl-ITC suppressed pathogen growth and its effects were not enhanced by the presence of other ITCs. This contradicts previous studies which demonstrated *R. solanacearum* sensitivity to 2-phenylethyl ITC at concentrations as low as 330 µM (Smith & Kirkegaard, 2002). However, in the previous experiment *R. solanacearum* was exposed to 2-phenylethyl ITC in agar instead of liquid media, which has been shown to increase the toxicity of ITCs (Sarwar et al., 1998). Moreover, it is possible that different *R. solanacearum* strains respond differently to ITCs, which could also explain discrepancy between ours and other studies. While the suppressive effects of sec-butyl ITC have previously been demonstrated against dust mites (Yun et al., 2012) and fungi (Bainard et al., 2009), no antimicrobial activity has been observed in bacteria. Variation in the antimicrobial activity of

ITCs could be explained by differences in chemical side-chain structure and molecular

weight which govern ITC volatility and hydrophobicity (Sarwar et al., 1998). Previous studies

have shown greater pathogen suppression by ITCs with aliphatic compared to aromatic

sidechains in fungal pathogens (Kurt et al., 2011; Sarwar et al., 1998), insect pests

(Matthiessen & Shackleton, 2005), and weeds (Vaughn et al., n.d.). With bacteria, the

toxicity of allyl-ITC could be attributed to its high volatility, very short R-side chains and high

reactivity (Kirkegaard & Sarwar, 1998; Manici et al., 1997; Neubauer et al., 2014). These

properties could enable rapid diffusion through the liquid media before ITC is lost in the

gaseous phase (Wang et al., 2009). This is supported by a study by Sarwar *et al.* (Sarwar et

al., 1998)*,* where a droplet of aliphatic allyl-ITC was shown to volatilise at room temperature

in 5 minutes, whilst aromatic 2-phenylethyl ITC remained in the liquid for over 72 hours.

Together, our result suggests that high volatility and reactivity could be important

properties determining the antibacterial effects of ITCs.

The evolution of ITC tolerance was mainly observed in the low transfer frequency ITC

exposure treatment. However, we also found that low transfer frequency control

populations showed improved tolerance measured in CPG media even though they had not

been exposed to allyl-ITC during the experiment. As all treatments were kept separate from

each other using tightly sealed bags, this effect is unlikely explained by 'cross selection' due

to ITC volatilisation. Alternatively, ITC tolerance evolution could have been linked to certain

metabolic adaptations in this transfer frequency treatment. In support of this, we found that

evolved control bacterial populations showed improved growth in the CPG media relative to

ancestral and ITC-exposed populations, indicative of media adaptation. While similar media

adaptations were observed in all control treatment populations, it is not clear why ITC

tolerance did not evolve under one- and two-day transfer frequency treatments. One

potential explanation for this could be growth-dependent effects on mutation rates. For

example, prior studies have shown that bacterial mutation rates can be elevated at

stationary phase (Loewe et al., 2003; Navarro Llorens et al., 2010), which could have

promoted ITC tolerance and media adaptation in the low transfer frequency treatment

where bacteria had spent the relatively longest time at stationary phase (Suppl. Fig. 1).

Alternatively, stationary phase growth conditions could have triggered expression of stress

tolerance genes, enabling selection for mutants with relatively higher ITC tolerance (Navarro

Llorens et al., 2010). For example, expression of *RpoS* sigma factor in *P. aeruginosa* has

previously been linked to elevated antibiotic resistance and biofilm formation at stationary

phase (Murakami et al., 2005; Olsen, 2015). While more work is needed to elucidate these

mechanisms, it is likely that the periodic 3-day growth cycle was important for driving ITC

tolerance evolution in our experimental conditions. Interestingly, the ITC tolerance that

evolved in the absence of allyl-ITC exposure was specific to CPG media and disappeared

when measured in 'naïve' LB media. This result suggests that ITC tolerance observed in

control populations was likely driven by adaptation to CPG growth media. Such adaptation

may have helped to offset the suppressive effects of allyl-ITC by boosting pathogen growth

to compensate increased mortality. Alternatively, it is possible that the glucose availability

in the CPG media indirectly favoured the evolution of ITC tolerance via metabolic

adaptations, which has previously been shown to occur both in the absence (Knöppel et al.,

2017) and presence of clinical antibiotics (Zampieri et al., 2017). Together, our results

suggest that prior exposure to allyl-ITC was required for the evolution of robust ITC

tolerance, which was independent of the growth media.

At the genetic level, ITC tolerance was not associated with any clear parallel mutations

or indels in the low transfer frequency treatments. Three clones from the low transfer

frequency control treatment had unique mutations in a gene encoding a dehydrogenase-like

uncharacterised protein. Dehydrogenase genes have previously been associated with both

metabolism and antibiotic resistance (Marshall et al., 1999). For instance, in *Escherichia coli*,

a mutation in a glucose dehydrogenase gene has been shown to function in

lipopolysaccharide modification and calanic acid biosynthesis, which enabled resistance to

polymyxin and other antimicrobial peptides (Lacour et al., 2008; Rodionova et al., 2020),

and may have contributed to ITC tolerance in these clones. Additionally, two clones from

the low transfer control treatment had mutations in a gene encoding a Tat pathway signal

protein which is involved in protein translocation across membranes (Palmer et al., 2005),

and may have enabled improved growth in the CPG media. Three clones from the

intermediate transfer frequency treatment had unique mutations in a gene encoding a

probable transcription regulator protein. While there is little information available regarding

this gene, it is located beside the IS2 transposase *TnpB* gene, potentially affecting its

regulation in DNA replication, recombination and repair activity (Pasternak et al., 2013).

Instead of treatment-specific parallel mutations, certain mutations were found across all

treatments. For example, mutations in genes encoding putative serine/threonine protein

kinases, amino acid biosynthesis (*hisH1* gene) and DNA replication, recombination and

repair (putative *RhsC* gene) were common for clones isolated from all treatments.

Mutations observed in serine/threonine protein kinase genes could have potentially

affected ITC tolerance if these enzymes were targeted by the ITCs as has been shown before

in the fungus *Alternaria brassicicola* (Calmes et al., 2015), and bacterial pathogen *E. coli*

(Luciano & Holley, 2009). However, as these mutations were not specific to ITC-treatment

clones, they were probably associated with bacterial growth and metabolism.

In *R. solanacearum*, insertion sequences (ISs) have been shown to affect host

virulence and phenotypic plasticity by inserting into and disrupting type III effectors and

global virulence regulators (Gonçalves et al., 2020; Jeong & Timmis, 2000). Therefore, we

investigated whether IS movement may be the cause of *R. solanacearum* ITC tolerance

adaptation. We identified one IS position in the chromosome and three positions in the

megaplasmid which showed treatment specific patterns. The gain of IS at position 2302900

was primarily observed with low transfer control isolates and was situated either ~50 bp

from the start codon or inside of a putative acyltransferase. Acyltransferases have a broad

range of functions including lipid storage (Ohlrogge & Browse, 1995), phospholipid

biosynthesis (Li et al., 2017), and the production of toxins (Greene et al., 2015) and

antibiotics (Kozakai et al., 2020). Whilst many of these functions are critical to cell growth,

some such as the production of toxins would be redundant when grown in media.

Therefore, gene disruption by ISs in the low transfer control may increase fitness by allowing

energy and nutrients to be re-directed towards promoting cell growth and competitivity,

potentially at the expense of reduced virulence *in planta*. We also found loss of two ISs in

the intergenic region of the megaplasmid in the low transfer control and ITC treatments.

While these were intergenic, they were close (~50-100 bp) to the start codons of their right

flanking genes and could have affected gene expression. In position 243500, the IS was

situated close to a (2Fe-2S)-binding protein gene. Iron-sulfur clusters have been implicated

in cellular metabolism, protein structural stabilisation, iron storage, and the regulation of

gene expression (Johnson et al., 2005). In the other position (253900), the IS was situated

close to an H-NS histone like protein gene and while non-significant, was lost more

frequently across low transfer ITC treatment clones (6/8) than low transfer control isolates

(4/8). H-NS histone like proteins are transcriptional repressors generally involved in

adaptation to environmental challenges like temperature stress and osmolarity gradients

(Atlung & Ingmer, 1997). Further, H-NS histone like proteins have been shown to stabilise

the sigma factor *RpoS* (Hommais et al., 2001) which acts as a master regulator of the

bacterial stress response. Whilst the H-NS histone-like protein could affect ITC tolerance by

mediating the bacterial stress response, the impact of the (2Fe-2S)-binding protein is less

clear. Notably, in *Campylobacter jejuni*, genes containing iron-sulfur clusters have been

found to be upregulated in response to ITCs, potentially due to their susceptibility to

oxidative stress caused by ITC exposure (Dufour et al., 2013). Therefore, by altering the

expression of the (2Fe-2S)-binding protein, IS loss could increase the pool of cellular iron-

sulfur cluster proteins and compensate for losses caused by ITC oxidative stress. In the final

megaplasmid IS position (209500), we identified a loss of IS from a calcium-binding protein

gene, which had likely disrupted gene expression or protein function in this gene with the

ancestral strain. In human breast cancer cells, ITCs, including phenethyl- (Tusskorn et al.,

2013) and allyl-ITC (Bo et al., 2016) have been found to induce mitochondrial calcium ion

mobilisation resulting in cytotoxicity through a reduction in mitochondrial membrane

potential. Whilst further work is required to determine the causes of ITC cytotoxicity in *R.*

solanacearum, upregulation of calcium-binding protein gene expression could have

increased ITC tolerance by facilitating the sequestration of free calcium ions. However, like

other genetic changes, loss of this IS did not occur statistically more often in the presence of

ITC selection. As a result, specific genetic mechanisms underlying ITC tolerance remain

elusive.

In conclusion, our findings demonstrate that allyl-ITC could potentially be used to

suppress the growth of *R. solanacearum* plant pathogen. However, repeated ITC exposure

could select for mutants with increased ITC tolerance, potentially weakening the long-term

efficiency of ITCs and biofumigation. Future work should focus on validating these findings

in more complex natural environments. For example, it is currently not clear if *R.*

solanacearum ITC tolerance evolves in the plant rhizosphere in the presence of other

microbes that could constrain mutation supply rate via resource and direct competition.

Moreover, different resistance mechanisms could be selected depending on soil

physiochemical properties and nutrient and plant root exudate availability, while it is not

clear if the ITC concentrations used in this experiment are achievable through biofumigation

and whether they might have negative effects on beneficial soil microbes. More efficient ITC

application could be attained by drilling the biofumigant plants into fields at the time of

flowering when GSL levels are highest using finely chopped plant material, which maximises

cell disruption and ITC release to the soil (Back et al., 2019). In addition, the efficacy of

Brassica-based biofumigation could potentially be improved by using plant cultivars with

elevated levels of sinigrin, the GSL precursor to allyl-ITC. Comprehensive *in vivo* work is thus

required to validate the potential of allyl-ITC for *R. solanacearum* biocontrol in the field. It

would also be interesting to study if ITC tolerance leads to life-history traits in *R.*

solanacearum, potentially affecting its virulence or competitiveness in the rhizosphere.

Acknowledgements

614 This work was funded by the NERC ACCE DTP (CLA), the Royal Society (RSG\R1\180213 and

615 CHL\R1\180031; V-P.F) and jointly by a grant from UKRI, Defra, and the Scottish

616 Government, under the Strategic Priorities Fund Plant Bacterial Diseases programme

617 (BB/T010606/1; V-P.F) at the University of York. The funders had no role in study design,

618 data collection and interpretation, or the decision to submit the work for publication.

References

Angus, J. F., Gardner, P. A., Kirkegaard, J. A., & Desmarchelier, J. M. (1994). Biofumigation: Isothiocyanates released frombrassica roots inhibit growth of the take-all fungus. *Plant and Soil*, *162*(1), 107–112. https://doi.org/10.1007/BF01416095

Arndt, D., Grant, J. R., Marcu, A., Sajed, T., Pon, A., Liang, Y., & Wishart, D. S. (2016). PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Research*, *44*(W1), W16–W21. https://doi.org/10.1093/nar/gkw387

Atlung, T., & Ingmer, H. (1997). H-NS: a modulator of environmentally regulated gene expression. *Molecular Microbiology*, *24*(1), 7–17. https://doi.org/10.1046/J.1365-2958.1997.3151679.X

Back, M., Barker, A., & Evans, K. (2019). *Effectiveness of biofumigant crops for the management of PCN in GB*. Agricultural And Horticultural Development Board (AHDB). https://pure.sruc.ac.uk/en/publications/effectiveness-of-biofumigant-crops-for-the-management-of-pcn-in-g

Bainard, L. D., Brown, P. D., & Upadhyaya, M. K. (2009). Inhibitory Effect of Tall Hedge Mustard (*Sisymbrium loeselii*) Allelochemicals on Rangeland Plants and Arbuscular Mycorrhizal Fungi. *Weed Science*, *57*(4), 386–393. https://doi.org/10.1614/WS-08-151.1

Bangarwa, S. K., Norsworthy, J. K., Mattice, J. D., & Gbur, E. E. (2011). Glucosinolate and Isothiocyanate Production from Brassicaceae Cover Crops in a Plasticulture Production System. *Weed Science*, *59*(2), 247–254. https://doi.org/10.1614/WS-D-10-00137.1

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477. https://doi.org/10.1089/cmb.2012.0021

Bending, G. D., & Lincoln, S. D. (1999). Characterisation of volatile sulphur-containing compounds produced during decomposition of Brassica juncea tissues in soil. *Soil Biology and Biochemistry*, *31*(5), 695–703. https://doi.org/10.1016/S0038-0717(98)00163-1

Bo, P., Lien, J.-C., Chen, Y.-Y., Yu, F.-S., Lu, H.-F., Yu, C.-S., Chou, Y.-C., Yu, C.-C., & Chung, J.-G. (2016). Allyl Isothiocyanate Induces Cell Toxicity by Multiple Pathways in Human Breast Cancer Cells. *Http://Dx.Doi.Org/10.1142/S0192415X16500245*, *44*(2), 415–437. https://doi.org/10.1142/S0192415X16500245

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/BIOINFORMATICS/BTU170

Borek, V., Morra, M. J., Brown, P. D., & McCaffrey, J. P. (1995). Transformation of the Glucosinolate-Derived Allelochemicals Allyl Isothiocyanate and Allylnitrile in Soil. *Journal of Agricultural and Food Chemistry*, *43*(7), 1935–1940. https://doi.org/10.1021/jf00055a033

Boutoille, D., Corvec, S., Caroff, N., Giraudeau, C., Espaze, E., Caillon, J., Plésiat, P., & Reynaud, A. (2004). Detection of an IS21 insertion sequence in the mexR gene of Pseudomonas aeruginosa increasing β-lactam resistance. *FEMS Microbiology Letters*, *230*(1), 143–146. https://doi.org/10.1016/S0378-1097(03)00882-6

Calmes, B., N'Guyen, G., Dumur, J., Brisach, C. A., Campion, C., Lacomi, B., PignÉ, S., Dias, E., Macherel, D., Guillemette, T., & Simoneau, P. (2015). Glucosinolate-derived isothiocyanates impact mitochondrial function in fungal cells and elicit an oxidative stress response necessary for growth recovery. *Frontiers in Plant Science*, *6*(June), 414. https://doi.org/10.3389/fpls.2015.00414

Chellemi, D. O., Olson, S. M., Mitchell, D. J., Secker, I., & McSorley, R. (1997). Adaptation of soil solarization to the integrated management of soilborne pests of tomato under humid conditions. *Phytopathology*, *87*(3), 250–258. https://doi.org/10.1094/PHYTO.1997.87.3.250

Ciampi-Panno, L., Fernandez, C., Bustamante, P., Andrade, N., Ojeda, S., & Contreras, A. (1989). Biological control of bacterial wilt of potatoes caused byPseudomonas solanacearum. *American Potato Journal*, *66*(5), 315–332. https://doi.org/10.1007/BF02854019

Dufour, V., Stahl, M., Rosenfeld, E., Stintzi, A., & Baysse, C. (2013). Insights into the mode of action of benzyl isothiocyanate on Campylobacter jejuni. *Applied and Environmental Microbiology*, *79*(22), 6958–6968. https://doi.org/10.1128/AEM.01967-13

Elphinstone, J. G. (2005). The current bacterial wilt situation: a global overview. *Bacterial Wilt Disease and the Ralstonia Solanacearum Species Complex*, 9–28.

Frick, A., Zebarth, B. J., & Szeto, S. Y. (1998). Behavior of the Soil Fumigant Methyl Isothiocyanate in Repacked Soil Columns. *Journal of Environmental Quality*, *27*(5), 1158–1169. https://doi.org/10.2134/jeq1998.00472425002700050022x

Gimsing, A L, & Kirkegaard, J. A. (2006). Glucosinolate and isothiocyanate concentration in soil following incorporation of Brassica biofumigants. *Soil Biology & Biochemistry*, *38*, 2255–2264. https://doi.org/10.1016/j.soilbio.2006.01.024

Gimsing, Anne Louise, & Kirkegaard, J. A. (2009). Glucosinolates and biofumigation: fate of glucosinolates and their hydrolysis products in soil. *Phytochemistry Reviews*, *8*(1), 299–310. https://doi.org/10.1007/s11101-008-9105-5

Gimsing, Anne Louise, Poulsen, J. L., Pedersen, H. L., & Hansen, H. C. B. (2007). Formation and degradation kinetics of the biofumigant benzyl isothiocyanate in soil. *Environmental Science and Technology*, *41*(12), 4271–4276. https://doi.org/10.1021/es061987t

Gonçalves, O. S., Campos, K. F., Assis, J. C. S. de, Fernandes, A. S., Souza, T. S., Rodrigues, L. G. do C., Queiroz, M. V. de, & Santana, M. F. (2020). Transposable elements contribute to the genome plasticity of Ralstonia solanacearum species complex. *Microbial Genomics*, *6*(5), 1–12. https://doi.org/10.1099/MGEN.0.000374

Greene, N. P., Crow, A., Hughes, C., & Koronakis, V. (2015). Structure of a bacterial toxin-activating acyltransferase. *Proceedings of the National Academy of Sciences*, *112*(23), E3058–E3066. https://doi.org/10.1073/PNAS.1503832112

Guarischi-Sousa, R., Puigvert, M., Coll, N. S., Siri, M. I., Pianzzola, M. J., Valls, M., & Setubal, J. C. (2016). Complete genome sequence of the potato pathogen Ralstonia solanacearum UY031. *Standards in Genomic Sciences*, *11*(1), 7. https://doi.org/10.1186/s40793-016-0131-4

Hanschen, F. S., Bru□, N., Brodehl, A., Mewis, I., Schreiner, M., Rohn, S., & Kroh, L. W. (2012). *Characterization of Products from the Reaction of Glucosinolate-Derived Isothiocyanates with Cysteine and Lysine Derivatives Formed in Either Model Systems or Broccoli Sprouts.* https://doi.org/10.1021/jf301718g

Hanschen, F. S., Brüggemann, N., Brodehl, A., Mewis, I., Schreiner, M., Rohn, S., & Kroh, L. W. (2012). Characterization of products from the reaction of glucosinolate-derived isothiocyanates with cysteine and lysine derivatives formed in either model systems or broccoli sprouts. *Journal of Agricultural and Food Chemistry*, *60*(31), 7735–7745. https://doi.org/10.1021/jf301718g

Hanschen, F. S., Yim, B., Winkelmann, T., Smalla, K., & Schreiner, M. (2015). Degradation of Biofumigant Isothiocyanates and Allyl Glucosinolate in Soil and Their Effects on the Microbial Community Composition. *PloS One*, *10*(7), e0132931. https://doi.org/10.1371/journal.pone.0132931

Hartz, T. K., Johnstone, P. R., Miyao, E. M., & Davis, R. M. (2005). Mustard cover crops are ineffective in suppressing soilborne disease or improving processing tomato yield. *HortScience*, *40*(7), 2016–2019. https://doi.org/10.21273/hortsci.40.7.2016

Hawkey, J., Hamidian, M., Wick, R. R., Edwards, D. J., Billman-Jacobe, H., Hall, R. M., & Holt, K. E. (2015). ISMapper: identifying transposase insertion sites in bacterial genomes from short read sequence data. *BMC Genomics 2015 16:1*, *16*(1), 1–11. https://doi.org/10.1186/S12864-015-1860-2

Hawkey, J., Monk, J. M., Billman-Jacobe, H., Palsson, B., & Holt, K. E. (2020). Impact of insertion sequences on convergent evolution of Shigella species. *PLOS Genetics*, *16*(7), e1008931. https://doi.org/10.1371/JOURNAL.PGEN.1008931

Hibbing, M. E., Fuqua, C., Parsek, M. R., & Peterson, S. B. (2010). Bacterial competition: Surviving and thriving in the microbial jungle. In *Nature Reviews Microbiology* (Vol. 8, Issue 1, pp. 15–25). Nature Publishing Group. https://doi.org/10.1038/nrmicro2259

Hommais, F., Krin, E., Laurent-Winter, C., Soutourina, O., Malpertuy, A., Caer, J.-P. Le, Danchin, A., & Bertin, P. (2001). Large-scale monitoring of pleiotropic regulation of gene expression by the prokaryotic nucleoid-associated protein, H-NS. *Molecular Microbiology*, *40*(1), 20–36. https://doi.org/10.1046/J.1365-2958.2001.02358.X

Hu, P., Hollister, E. B., Somenahally, A. C., Hons, F. M., & Gentry, T. J. (2015). Soil bacterial and fungal communities respond differently to various isothiocyanates added for biofumigation. *Frontiers in Microbiology*, *5*, 729. https://doi.org/10.3389/fmicb.2014.00729

Jeong, E. L., & Timmis, J. N. (2000). Novel insertion sequence elements associated with genetic heterogeneity and phenotype conversion in Ralstonia solanacearum. *Journal of Bacteriology*, *182*(16), 4673–4676. https://doi.org/10.1128/JB.182.16.4673-4676.2000

Ji, P., Momol, M. T., Rich, J. R., Olson, S. M., & Jones, J. B. (2007). Development of an Integrated Approach for Managing Bacterial Wilt and Root-Knot on Tomato Under Field Conditions. *Plant Disease*, *91*(10), 1321–1326. https://doi.org/10.1094/PDIS-91-10-1321

Johnson, D. C., Dean, D. R., Smith, A. D., & Johnson, M. K. (2005). STRUCTURE, FUNCTION, AND FORMATION OF BIOLOGICAL IRON-SULFUR CLUSTERS. *Http://Dx.Doi.Org/10.1146/Annurev.Biochem.74.082803.133518*, *74*, 247–281. https://doi.org/10.1146/ANNUREV.BIOCHEM.74.082803.133518

Khokhani, D., Lowe-Power, T. M., Tran, T. M., & Allen, C. (2017). A single regulator mediates strategic switching between attachment/spread and growth/virulence in the plant pathogen Ralstonia solanacearum. *MBio*, *8*(5). https://doi.org/10.1128/MBIO.00895-17

Kirkegaard, J. A., & Matthiessen, J. N. (2005). *Developing and refining the biofumigation concept.*

Kirkegaard, J. A., & Sarwar, M. (1998). Biofumigation potential of brassicas: I. Variation in glucosinolate profiles of diverse field-grown brassicas. *Plant and Soil*, *201*(1), 71–89. https://doi.org/10.1023/A:1004364713152

Kirkegaard, J. A., Sarwar, M., Wong, P. T. W., Mead, A., Howe, G., & Newell, M. (2000). Field studies on the biofumigation of take-all by Brassica break crops. *Australian Journal of Agricultural Research*, *51*(4), 445–456. https://doi.org/10.1071/AR99106

Kirkegaard, J. A., Wong, P. T. W., & Desmarchelier, J. M. (1996). In vitro suppression of fungal root pathogens of cereals by Brassica tissues. *Plant Pathology*, *45*(3), 593–603. https://doi.org/10.1046/j.1365-3059.1996.d01-143.x

Knöppel, A., Näsvall, J., & Andersson, D. I. (2017). Evolution of antibiotic resistance without antibiotic exposure. *Antimicrobial Agents and Chemotherapy*, *61*(11), 1495–1512. https://doi.org/10.1128/AAC.01495-17

Kozakai, R., Ono, T., Hoshino, S., Takahashi, H., Katsuyama, Y., Sugai, Y., Ozaki, T., Teramoto, K., Teramoto, K., Tanaka, K., Abe, I., Asamizu, S., & Onaka, H. (2020). Acyltransferase that catalyses the condensation of polyketide and peptide moieties of goadvionin hybrid lipopeptides. *Nature Chemistry 2020 12:9*, *12*(9), 869–877. https://doi.org/10.1038/s41557-020-0508-2

Kurt, Ş., Güneş, U., & Soylu, E. M. (2011). *In vitro* and *in vivo* antifungal activity of synthetic pure isothiocyanates against *Sclerotinia sclerotiorum*. *Pest Management Science*, *67*(7), 869–875. https://doi.org/10.1002/ps.2126

Lacour, S., Bechet, E., Cozzone, A. J., Mijakovic, I., & Grangeasse, C. (2008). Tyrosine Phosphorylation of the UDP-Glucose Dehydrogenase of Escherichia coli Is at the Crossroads of Colanic Acid Synthesis and Polymyxin Resistance. *PLoS ONE*, *3*(8), e3053. https://doi.org/10.1371/journal.pone.0003053

Larkin, R. P., & Griffin, T. S. (2007). Control of soilborne potato diseases using Brassica green manures. *Crop Protection*, *26*(7), 1067–1077. https://doi.org/10.1016/j.cropro.2006.10.004

Larkin, R. P., & Halloran, J. M. (2015). Management effects of disease-suppressive rotation crops on potato yield and soilborne disease and their economic implications in potato production. *American Journal of Potato Research*, *91*(5), 429–439. https://doi.org/10.1007/s12230-014-9366-z

Li, E., de Jonge, R., Liu, C., Jiang, H., Friman, V. P., Pieterse, C. M. J., Bakker, P. A. H. M., & Jousset, A. (2020). Rapid evolution of bacterial mutualism in the plant rhizosphere. In *bioRxiv* (p. 2020.12.07.414607). bioRxiv. https://doi.org/10.1101/2020.12.07.414607

Li, Z., Tang, Y., Wu, Y., Zhao, S., Bao, J., Luo, Y., & Li, D. (2017). Structural insights into the committed step of bacterial phospholipid biosynthesis. *Nature Communications 2017 8:1*, *8*(1), 1–14. https://doi.org/10.1038/s41467-017-01821-9

Lin, C.-M., Preston Iii, J. F., & Wei, C.-I. (2000). Antibacterial Mechanism of Allyl Isothiocyanate. *Journal of Food Protection*, *63*(6), 727–734. http://jfoodprotection.org/doi/pdf/10.4315/0362-028X-63.6.727

Loewe, L., Textor, V., & Scherer, S. (2003). High Deleterious Genomic Mutation Rate in Stationary Phase of Escherichia coli. *Science*, *302*(5650), 1558–1560. https://doi.org/10.1126/science.1087911

Lord, J. S., Lazzeri, L., Atkinson, H. J., & Urwin, P. E. (2011). Biofumigation for Control of Pale Potato Cyst Nematodes: Activity of Brassica Leaf Extracts and Green Manures on Globodera pallida in Vitro and in Soil. *J. Agric. Food Chem*, *59*, 7882–7890.

https://doi.org/10.1021/jf200925k

Luciano, F. B., & Holley, R. A. (2009). Enzymatic inhibition by allyl isothiocyanate and factors
affecting its antimicrobial action against Escherichia coli O157:H7. *International Journal*
of Food Microbiology, *131*(2–3), 240–245.
https://doi.org/10.1016/j.ijfoodmicro.2009.03.005

Manici, L. M., Lazzeri, L., & Palmieri, S. (1997). In Vitro Fungitoxic Activity of Some
Glucosinolates and Their Enzyme-Derived Products toward Plant Pathogenic Fungi.
Journal of Agricultural and Food Chemistry, *45*(7), 2768–2773.
https://doi.org/10.1021/jf9608635

Marshall, C. G., Zolli, M., & Wright, G. D. (1999). Molecular mechanism of VanHst, an α-
ketoacid dehydrogenase required for glycopeptide antibiotic resistance from a
glycopeptide producing organism. *Biochemistry*, *38*(26), 8485–8491.
https://doi.org/10.1021/bi982843x

Matthiessen, J. N., & Kirkegaard, J. A. (2006). Biofumigation and Enhanced Biodegradation:
Opportunity and Challenge in Soilborne Pest and Disease Management. *Critical*
Reviews in Plant Sciences, *25*(3), 235–265.
https://doi.org/10.1080/07352680600611543

Matthiessen, J. N., & Shackleton, M. A. (2005). Biofumigation: environmental impacts on the
biological activity of diverse pure and plant-derived isothiocyanates. *Pest Management*
Science, *61*(11), 1043–1051. https://doi.org/10.1002/ps.1086

Mazzola, M., & Gu, Y. H. (2002). Wheat genotype-specific induction of soil microbial
communities suppressive to disease incited by Rhizoctonia solani Anastomosis Group
(AG)-5 and AG-8. *Phytopathology*, *92*(12), 1300–1307.
https://doi.org/10.1094/PHYTO.2002.92.12.1300

Mazzola, M., Hewavitharana, S. S., & Strauss, S. L. (2015). *Brassica* Seed Meal Soil
Amendments Transform the Rhizosphere Microbiome and Improve Apple Production
Through Resistance to Pathogen Reinfestation. *Phytopathology*, *105*(4), 460–469.
https://doi.org/10.1094/PHYTO-09-14-0247-R

Mondal, M. F., Asaduzzaman, M., & Asao, T. (2015). Adverse Effects of Allelopathy from
Legume Crops and Its Possible Avoidance. *American Journal of Plant Sciences*,
06(06), 804–810. https://doi.org/10.4236/ajps.2015.66086

Murakami, K., Ono, T., Viducic, D., Kayama, S., Mori, M., Hirota, K., Nemoto, K., & Miyake,
Y. (2005). Role for rpoS gene of Pseudomonas aeruginosa in antibiotic tolerance.
FEMS Microbiology Letters, *242*(1), 161–167.
https://doi.org/10.1016/j.femsle.2004.11.005

Navarro Llorens, J. M., Tormo, A., & Martínez-García, E. (2010). Stationary phase in gram-
negative bacteria. In *FEMS Microbiology Reviews* (Vol. 34, Issue 4, pp. 476–495).
Blackwell Publishing Ltd. https://doi.org/10.1111/j.1574-6976.2010.00213.x

Neubauer, C., Heitmann, B., & Müller, C. (2014). Biofumigation potential of Brassicaceae
cultivars to Verticillium dahliae. *European Journal of Plant Pathology*, *140*(2), 341–352.
https://doi.org/10.1007/s10658-014-0467-9

Ngala, B. M., Woods, S. R., & Back, M. A. (2015). Sinigrin degradation and G. pallida
suppression in soil cultivated with brassicas under controlled environmental conditions.
Applied Soil Ecology, *95*, 9–14. https://doi.org/10.1016/j.apsoil.2015.05.009

Ohlrogge, J., & Browse, J. (1995). Lipid biosynthesis. *The Plant Cell*, *7*(7), 957.
https://doi.org/10.1105/TPC.7.7.957

Olivier, C., Vaughn, S. F., Mizubuti, E. S. G., & Loria, R. (1999). Variation in allyl
isothiocyanate production within Brassica species and correlation with fungicidal
activity. *Journal of Chemical Ecology*, *25*(12), 2687–2701.
https://doi.org/10.1023/A:1020895306588

Olsen, I. (2015). Biofilm-specific antibiotic tolerance and resistance. In *European Journal of*
Clinical Microbiology and Infectious Diseases (Vol. 34, Issue 5, pp. 877–886). Springer
Verlag. https://doi.org/10.1007/s10096-015-2323-z

Palmer, T., Sargent, F., & Berks, B. C. (2005). Export of complex cofactor-containing
proteins by the bacterial Tat pathway. *Trends in Microbiology*, *13*(4), 175–180.
https://doi.org/10.1016/J.TIM.2005.02.002

Pasternak, C., Dulermo, R., Ton-Hoang, B., Debuchy, R., Siguier, P., Coste, G., Chandler,
M., & Sommer, S. (2013). IS *Dra 2* transposition in *D einococcus radiodurans* is
downregulated by TnpB. *Molecular Microbiology*, *88*(2), 443–455.
https://doi.org/10.1111/mmi.12194

Perrier, A., Barlet, X., Rengel, D., Prior, P., Poussier, S., Genin, S., & Guidot, A. (2019).
Spontaneous mutations in a regulatory gene induce phenotypic heterogeneity and
adaptation of Ralstonia solanacearum to changing environments. *Environmental*
Microbiology, *21*(8), 3140–3152. https://doi.org/10.1111/1462-2920.14717

Poirel, L., Decousser, J. W., & Nordmann, P. (2003). Insertion sequence ISEcp1B is
involved in expression and mobilization of a blaCTX-M β-lactamase gene. *Antimicrobial*
Agents and Chemotherapy, *47*(9), 2938–2945. https://doi.org/10.1128/AAC.47.9.2938-
2945.2003

Qin, S., Gan, J., Liu, W., & Becker, J. O. (2004). *Degradation and Adsorption of Fosthiazate*
in Soil. https://doi.org/10.1021/jf049094c

Ramesh, R., Joshi, A. A., & Ghanekar, M. P. (2009). Pseudomonads: major antagonistic
endophytic bacteria to suppress bacterial wilt pathogen, Ralstonia solanacearum in the
eggplant (Solanum melongena L.). *World Journal of Microbiology and Biotechnology*,
25(1), 47–55. https://doi.org/10.1007/s11274-008-9859-3

Ranjan, V. K., Mukherjee, S., Thakur, S., Gupta, K., & Chakraborty, R. (2021). Pandrug-
resistant Pseudomonas sp. expresses New Delhi metallo-β-lactamase-1 and consumes
ampicillin as sole carbon source. *Clinical Microbiology and Infection*, *27*(3), 472.e1-
472.e5. https://doi.org/10.1016/j.cmi.2020.10.032

Rodionova, I. A., Zhang, Z., Aboulwafa, M., & Saier, M. H. (2020). UDP-glucose
dehydrogenase Ugd in E. coli is activated by Gmd and RffD, is inhibited by CheY, and
regulates swarming. In *bioRxiv* (p. 2020.01.08.899336). bioRxiv.
https://doi.org/10.1101/2020.01.08.899336

Rudolph, R. E., Sams, C., Steiner, R., Thomas, S. H., Walker, S., & Uchanski, M. E. (2015).
Biofumigation performance of four brassica crops in a green chile pepper (Capsicum
annuum) rotation system in southern New Mexico. *HortScience*, *50*(2), 247–253.
https://doi.org/10.21273/hortsci.50.2.247

Rumberger, A., & Marschner, P. (2003). 2-Phenylethylisothiocyanate concentration and
microbial community composition in the rhizosphere of canola. *Soil Biology and*
Biochemistry, *35*(3), 445–452. https://doi.org/10.1016/S0038-0717(02)00296-1

Salanoubat, M., Genin, S., Artiguenave, F., Gouzy, J., Mangenot, S., Arlat, M., Billault, A.,
Brottier, P., Camus, J. C., Cattolico, L., Chandler, M., Choisne, N., Claudel-Renard, C.,
Cunnac, S., Demange, N., Gaspin, C., Lavie, M., Moisan, A., Robert, C., … Boucher, C.
A. (2002). Genome sequence of the plant pathogen Ralstonia solanacearum. *Nature*,

896   *415*(6871), 497–502. https://doi.org/10.1038/415497a

897 Sarwar, M., Kirkegaard, J. A., Wong, P. T. W., & Desmarchelier, J. M. (1998). Biofumigation
898   potential of brassicas. *Plant and Soil*, *201*(1), 103–112.
899   https://doi.org/10.1023/A:1004381129991

900 Savary, S., Ficke, A., Aubertot, J. N., & Hollier, C. (2012). Crop losses due to diseases and
901   their implications for global food production losses and food security. In *Food Security*
902   (Vol. 4, Issue 4, pp. 519–537). Springer. https://doi.org/10.1007/s12571-012-0200-5

903 Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, *30*(14),
904   2068–2069. https://doi.org/10.1093/bioinformatics/btu153

905 Seemann, T. (2015). *Snippy: fast bacterial variant calling from NGS reads.*
906   *https://github.com/tseemann/snippy.*

907 Shigemizu, D., Miya, F., Akiyama, S., Okuda, S., Boroevich, K. A., Fujimoto, A., Nakagawa,
908   H., Ozaki, K., Niida, S., Kanemura, Y., Okamoto, N., Saitoh, S., Kato, M., Yamasaki,
909   M., Matsunaga, T., Mutai, H., Kosaki, K., & Tsunoda, T. (2018). IMSindel: An accurate
910   intermediate-size indel detection tool incorporating de novo assembly and gapped
911   global-local alignment with split read analysis. *Scientific Reports 2018 8:1*, *8*(1), 1–9.
912   https://doi.org/10.1038/s41598-018-23978-z

913 Smith, B. J., & Kirkegaard, J. A. (2002). In vitro inhibition of soil microorganisms by 2-
914   phenylethyl isothiocyanate. *Plant Pathology*, *51*(5), 585–593.
915   https://doi.org/10.1046/j.1365-3059.2002.00744.x

916 Sofrata, A., Santangelo, E. M., Azeem, M., Borg-Karlson, A.-K., Gustafsson, A., & Pütsep, K.
917   (2011). Benzyl Isothiocyanate, a Major Component from the Roots of Salvadora
918   Persica Is Highly Active against Gram-Negative Bacteria. *PLoS ONE*, *6*(8), e23045.
919   https://doi.org/10.1371/journal.pone.0023045

920 Stirling, G. R., & Stirling, A. M. (2003). The potential of Brassica green manure crops for
921   controlling root-knot nematode (Meloidogyne javanica) on horticultural crops in a
922   subtropical environment. *Australian Journal of Experimental Agriculture*, *43*(6), 623–
923   630. https://doi.org/10.1071/EA02175

924 Stringlis, I. A., Yu, K., Feussner, K., De Jonge, R., Van Bentum, S., Van Verk, M. C.,
925   Berendsen, R. L., Bakker, P. A. H. M., Feussner, I., & Pieterse, C. M. J. (2018).
926   MYB72-dependent coumarin exudation shapes root microbiome assembly to promote
927   plant health. *Proceedings of the National Academy of Sciences of the United States of
928   America*, *115*(22), E5213–E5222. https://doi.org/10.1073/pnas.1722335115

929 Tatusova, T., DiCuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E. P., Zaslavsky, L.,
930   Lomsadze, A., Pruitt, K. D., Borodovsky, M., & Ostell, J. (2016). NCBI prokaryotic
931   genome annotation pipeline. *Nucleic Acids Research*, *44*(14), 6614–6624.
932   https://doi.org/10.1093/NAR/GKW569

933 Tusskorn, O., Senggunprai, L., Prawan, A., Kukongviriyapan, U., & Kukongviriyapan, V.
934   (2013). Phenethyl isothiocyanate induces calcium mobilization and mitochondrial cell
935   death pathway in cholangiocarcinoma KKU-M214 cells. *BMC Cancer 2013 13:1*, *13*(1),
936   1–12. https://doi.org/10.1186/1471-2407-13-571

937 Vaughn, S. F., Palmquist, D. E., Duval, S. M., & Berhow, M. A. (n.d.). *Herbicidal Activity of
938   Glucosinolate-Containing Seedmeals* (Vol. 54, Issue 4).

939 Vervoort, M. T. W., Vonk, J. A., Brolsma, K. M., Schütze, W., Quist, C. W., De Goede, R. G.
940   M., Hoffland, E., Bakker, J., Mulder, C., Hallmann, J., & Helder, J. (2014). Release of
941   isothiocyanates does not explain the effects of biofumigation with Indian mustard

942 cultivars on nematode assemblages. *Soil Biology and Biochemistry*, *68*, 200–207.
943 https://doi.org/10.1016/j.soilbio.2013.10.008

944 Wang, D., Rosen, C., Kinkel, L., Cao, A., Tharayil, N., & Gerik, J. (2009). Production of
945 methyl sulfide and dimethyl disulfide from soil-incorporated plant materials and
946 implications for controlling soilborne pathogens. *Plant and Soil*, *324*(1–2), 185–197.
947 https://doi.org/10.1007/s11104-009-9943-y

948 Warton, B., Matthiessen, J. N., & Shackleton, M. A. (2001). Glucosinolate content and
949 isothiocyanate evolution - Two measures of the biofumigation potential of plants.
950 *Journal of Agricultural and Food Chemistry, 49*(11), 5244–5250.
951 https://doi.org/10.1021/jf010545s

952 Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Unicycler: Resolving bacterial
953 genome assemblies from short and long sequencing reads. *PLOS Computational*
954 *Biology*, *13*(6), e1005595. https://doi.org/10.1371/JOURNAL.PCBI.1005595

955 Wink, M. (2013). Evolution of secondary metabolites in legumes (Fabaceae). *South African*
956 *Journal of Botany*, *89*, 164–175. https://doi.org/10.1016/j.sajb.2013.06.006

957 Xie, Z., & Tang, H. (2017). ISEScan: automated identification of insertion sequence
958 elements in prokaryotic genomes. *Bioinformatics*, *33*(21), 3340–3347.
959 https://doi.org/10.1093/BIOINFORMATICS/BTX433

960 Yabuuchi, E., Kosako, Y., Yano, I., Hotta, H., & Nishiuchi, Y. (1995). Transfer of Two
961 *Burkholderia* and An *Alcaligenes* Species to *Ralstonia* Gen. Nov. *Microbiology and*
962 *Immunology, 39*(11), 897–904. https://doi.org/10.1111/j.1348-0421.1995.tb03275.x

963 Yim, B., Hanschen, F. S., Wrede, A., Nitt, H., Schreiner, M., Smalla, K., & Winkelmann, T.
964 (2016). Effects of biofumigation using Brassica juncea and Raphanus sativus in
965 comparison to disinfection using Basamid on apple plant growth and soil microbial
966 communities at three field sites with replant disease. *Plant and Soil*, *406*(1–2), 389–
967 408. https://doi.org/10.1007/s11104-016-2876-3

968 Yun, Y.-K., Kim, H.-K., Kim, J.-R., Hwang, K., & Ahn, Y.-J. (2012). Contact and fumigant
969 toxicity of *Armoracia rusticana* essential oil, allyl isothiocyanate and related compounds
970 to *Dermatophagoides farinae*. *Pest Management Science*, *68*(5), 788–794.
971 https://doi.org/10.1002/ps.2327

972 Zampieri, M., Enke, T., Chubukov, V., Ricci, V., Piddock, L., & Sauer, U. (2017). Metabolic
973 constraints on the evolution of antibiotic resistance. *Molecular Systems Biology*, *13*(3),
974 917. https://doi.org/10.15252/msb.20167028

Tables

Location	Chromosome	Chromosome	Chromosome	Chromosome	Chromosome	Chromosome	Chromosome	Chromosome	Chromosome	Plasmid	Plasmid	Plasmid
Position	293900	294413	585646	1257034	1258240	2302809	2830656	2874048	3123064	44512	105610	124931
Type	snp	snp	ins	snp	snp	snp	ins	ins	ins	del (27bp)	del (1bp)	snp
Ref	G	C	T	G	C	C	A	C	A	TCGTGAGCGGCAAGCCGGCACATCGCAA	TG	G
Alt	A	T	TCGTGCTG	C	G	T	ACAGCAACGG	CGGGCACT	AC	T	T	A
Effect	Synonymous variant	Synonymous variant	Frameshift variant	Missense variant	Missense variant	Synonymous variant	Conservative inframe insertion	Frameshift variant	Conservative inframe insertion	Frameshift variant	Synonymous variant	Missense variant
Locus tag	RSUY_02640	RSUY_02640	RSUY_05230	RSUY_11710	RS_RS11675	RSUY_21390	RSUY_26530	ATK36_5281	BSE24067_05643	NA8A_21102	RSIPO_03141	RSUY_33140
Product	Putative deoxyribonuclease RhsC	Putative deoxyribonuclease RhsC	Imidazole glycerol phosphate synthase subunit HisH1	HTH-type transcriptional regulator DmlR	Probable transcription regulator protein (86.1% similarity)	F-box domain-containing protein (83.3% similarity)	Putative serine/threonine protein kinase (84.6% similarity)	Dehydrogenase-like uncharacterised protein (90.9% similarity)	Tat pathway signal protein (90.9% similarity)	DNA-3-methyladenine glycosidase II (100% similarity)	Hypothetical transmembrane protein (100% similarity)	IS5/IS1182 family transposase (100% similarity)
Int No ITC 1	■		■		■		■					■
Int No ITC 2	■		■		■		■					
Int No ITC 3			■				■					
Int No ITC 4	■		■									
Int No ITC 5	■		■									
Int No ITC 6			■									
Int No ITC 7			■									
Int No ITC 8	■		■				■					
Low No ITC 1			■									
Low No ITC 2	■		■			■	■		■			
Low No ITC 3	■		■									
Low No ITC 4			■									
Low No ITC 5	■		■							■		
Low No ITC 6			■				■					
Low No ITC 7			■									
Low No ITC 8			■									
Low ITC 1			■				■				■	
Low ITC 2			■				■					■
Low ITC 3			■				■					
Low ITC 4			■									
Low ITC 5							■					
Low ITC 6	■		■				■					
Low ITC 7	■		■									
Low ITC 8							■					

Table 1. Mutated *Ralstonia solanacearum* genes and annotated gene functions observed in intermediate and low transfer frequency control (no-ITC), and ITC-exposed low transfer frequency treatments. Gene function predictions were derived based on BLAST using UNIPROT and percentage (%) sequence similarity is included for putative (hypothetical) proteins. Filled cells denote for the presence of mutations in given clones and white cells denote for the absence of given mutations. Replicates are named by treatments, IntNoITC= Intermediate transfer frequency, no ITC; LowNoITC= Low transfer frequency, no ITC; LowITC= Low transfer frequency, ITC.

Figures and figure legends

Figure 1. The antimicrobial activity of different ITCs against *Ralstonia solanacearum*

pathogen when applied alone and in combination. The chemical properties of the three

different ITCs predominantly released from Indian mustard biofumigant plant (*Brassica*

juncea) (A), and their effects on *R. solanacearum* growth after 48h exposure when applied

alone and in combination in liquid microcosms at low (500 µM) and high (1000 µM)

concentrations (B). In (B) boxplot colours represent different ITC treatments that are

labelled on X-axes as follows: (0): no-ITC (control); (1) allyl-ITC; (2) sec-butyl ITC and (3) 2-

phenylethyl ITC. Individual data points show bacterial densities for each technical replicate

(N=8). The boxplots show the minimum, maximum, interquartile range and the median

(black line).

Figure 2. *Ralstonia solanacearum* **density dynamics (OD600$_{nm}$) during the evolution**

experiment in the absence and presence of allyl-ITC in high, intermediate and low transfer

frequency treatments. In all panels, black and red lines correspond to *R. solanacearum*

densities in the absence and presence of 500 µM allyl-ITC, respectively. Panels A-C

correspond to high (1-day), intermediate (2-day) and low (3-day) transfer frequency

treatments, respectively. Grey shaded areas indicate the time point of serial transfers, while

optical density reads were taken at 24-hour intervals in all treatments. Each time point

shows the mean of eight biological replicates and bars show ±1 error of mean.

1008

Figure 3. Comparison of *Ralstonia solanacearum* ITC tolerance between the ancestral clone and evolved populations from high, intermediate and low transfer frequency treatments at the end of the evolution experiment in CPG and LB media. ITC tolerance was determined as *R. solanacearum* growth (OD600$_{nm}$) after 48h exposure to 500 µM allyl-ITC in CPG (A) and LB (C) media. Growth was also measured in the absence of allyl-ITC in both CPG (B) and LB (D) media. High (1-day), intermediate (2-day) and low (3-day) transfer frequency

1015   treatments are shown in grey, blue and yellow boxplots, respectively, and boxplots show

1016   the minimum, maximum, interquartile range and the median (black line). Individual data

1017   points show bacterial densities for each biological replicate population (N=8).

1018

1019

Figure 4. Comparison of *Ralstonia solanacearum* ITC tolerance between the ancestral and evolved clones from high, intermediate and low transfer frequency treatments at the end of the evolution experiment in LB media. ITC tolerance was determined as *R. solanacearum* growth ($OD600_{nm}$) after 48h exposure to 500 µM allyl-ITC in LB media (A). Growth was also measured in the absence of allyl-ITC (B). High (1-day), intermediate (2-day) and low (3-day) frequency treatments are shown in grey, blue and yellow, respectively, and boxplots show the minimum, maximum, interquartile range and the median (black line). Individual data points show bacterial densities for each biological replicate population (N=8).

1020

1021

1022

1023

1024

1025

1026

1027

1028

A

1029

1030

B

Figure 5. Mutations (A) and insertion sequences (IS; B) associated with evolved *Ralstonia*

solanacearum clones. Each ring in panel A represents the *R. solanacearum* genome

(Chromosome on the left and Megaplasmid on the right). Rings are grouped by the

sequenced treatments) in different colours (see key) and dots represent mutations at

different loci. Dots are sized and coloured by the number of replicates that had the same

mutations (N=8) in the indicated locus. Labels show the gene name, when named, or the

numbered locus tag. Distance marker is shown as Mb within each ring. In panel B, tile plot

shows presence (filled tiles) and absence (unfilled tiles) of insertion sequences (ISs) in each

isolate. The X-axis of the tile plot shows the IS position rounded to the nearest 100 bp. Gene

schematics on the right show insertion sequence at each position and nearby genes. Gene

annotation and distance between insertion sequence and genes are shown, with gene size

and distance proportional to the scale bar (bottom right).

 # Supplementary Materials

1044 **Supplementary Table 1. The mean density reduction (%) of *Ralstonia solanacearum***

1045 **bacterium when exposed to 500 or 1000 µM allyl, sec-butyl and 2-phenylethyl ITCs in CPG**

1046 **growth media after 24, 48 or 72 hours relative to when grown in the absence of ITCs.** This

1047 table is based on the same data presented in Supplementary Fig. 2.

ITC Type and Concentration (µM)	Time (h)	Bacterial density reduction (%) compared to control
Allyl-ITC, 500	24	66
	48	54
	72	27
Allyl-ITC, 1000	24	66
	48	47
	72	41
Sec-Butyl ITC, 500	24	33
	48	26
	72	9
Sec-Butyl ITC, 1000	24	30
	48	27
	72	8
2-Phenylethyl ITC, 500	24	39
	48	13
	72	10
2-Phenylethyl ITC, 1000	24	38
	48	18
	72	13

1048

1049 **Supplementary Table 2. Prophage information of ancestral and experimental isolate assemblies as determined using flanking regions**

1050 **mapped to UY031.** Replicates are named by treatments, IntNoITC= Intermediate transfer frequency, no ITC; LowNoITC= Low transfer

1051 frequency, no ITC; LowITC= Low transfer frequency, ITC.

1052

Clone	Prophage	Left flank UY031 position	Right flank UY031 position	Length (kb)	GC content (%)	Total proteins #
UY031	Unclassified A	-	-	37	62.76	44
	RS551	-	-	13.4	61.24	16
	PHAGE_Vibrio_VHML_NC_004456	-	-	18.5	64.64	29
Ancestor	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	17
IntNoITC1	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17
IntNoITC2	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17

IntNoITC3	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17
IntNoITC4	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
IntNoITC5	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
IntNoITC6	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
IntNoITC7	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1218233-1223232	13.1	58.58	17
IntNoITC8	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17
LowNoITC1	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42

	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17
LowNoITC2	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
LowNoITC3	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
LowNoITC4	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218233-1223232	NZ_CP012687.1:1236385-1237640	13.1	58.58	17
LowNoITC5	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236385-1237640	13.1	58.58	18
LowNoITC6	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	18
LowNoITC7	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	18

LowNoITC8	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	17
LowITC1	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	17
LowITC2	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	43
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	17
LowITC3	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237640	13.1	58.58	18
LowITC4	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	17
LowITC5	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	18
LowITC6	Unclassified A	NZ_CP012687.1:1121824-1126823	NZ_CP012687.1:1162291-1162775	35.4	62.85	42

	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	17
LowITC7	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	18
LowITC8	Unclassified A	NZ_CP012687.1:1121823-1126822	NZ_CP012687.1:1162290-1162775	35.4	62.85	41
	RS551	NZ_CP012687.1:1218220-1223219	NZ_CP012687.1:1236384-1237639	13.1	58.58	18

1053

1054

1055

1056 **Supplementary Figure 1. The effect of allyl-ITC pre-volatilisation for antibacterial activity**

1057 **against _Ralstonia solanacearum_.** _R. solanacearum_ bacterial growth was measured in CPG

1058 media supplemented with 0 (No allyl-ITC) or 500 µM of allyl-ITC that had been allowed to

1059 volatilise for 2, 24, 48 or 72 hours (see key). All data points show the mean of eight technical

1060 replicates and bars show ±1 standard error of the mean (SEM).

1061

Supplementary Figure 2. Effects of allyl, sec-butyl and 2-phenylethyl ITCs on *Ralstonia*

***solanacearum* growth at different ITC concentrations.** In all panels, *R. solanacearum*

bacterial densities are shown on the Y-axis as optical density ($OD600_{nm}$), measured at 24-

hour intervals (X-axis). In all panels, different line colours refer to different ITC

concentrations (see key in A). All data points show the mean of eight technical replicates

and bars show ±1 standard error of the mean (SEM).

1068

1069

Supplementary Figure 3. *Ralstonia solanacearum* **tolerance to ampicillin beta-lactam**

antibiotic. Ampicillin tolerance was measured as the growth of ancestral and evolved *R.*

solanacearum clones isolated from intermediate (Int) and low transfer frequency (Low)

control treatments (no-ITC) and ITC-exposed low transfer frequency treatment in the

absence (A) and presence (B-C) of ampicillin (15 and 30 µg/ml concentrations). Boxplots

show the minimum, maximum, interquartile range and the median (black line) after 48

hours. Individual data points show bacterial densities for each biological replicate clone (N=

8). Different small case letters above boxplots indicate significant pairwise differences

(Tukey: p< 0.05) between treatments within each panel.

1079

1080 **Supplementary Figure 4. Presence and absence of intermediate indels found in more than**

1081 **two isolates in the chromosome and megaplasmid.** The X-axis shows the indel position

1082 rounded to the nearest 100bp. The Y-axis shows isolates grouped as shown in Figure 5.

1083

1084

1085

Supplementary Figure 5. Presence and absence of insertion sequences in the chromosome and megaplasmid. The X and Y-axes show the insertion sequence position and experimental isolate, respectively, as outlined in Figure 5.

1089