Co-option of local and systemic immune responses by the hormonal signalling system triggering metamorphosis

Catarina Nunes[1†], Takashi Koyama[2], Élio Sucena[1,3†]

[1] Evolution and Development Laboratory, Instituto Gulbenkian de Ciência, Oeiras, Portugal; [2] Section for Cell and Neurobiology, Department of Biology, University of Copenhagen, Copenhagen, Denmark; [3] Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal

[†] Corresponding authors: esucena@igc.gulbenkian.pt; cdnunes@igc.gulbenkian.pt

ABSTRACT

Insect metamorphosis is regulated by the production, secretion and degradation of two peripheral hormones: 20-hydroxyecdysone (ecdysone) and juvenile hormone (JH). In addition to their roles in developmental regulation, increasing evidence suggests that these hormones are involved in innate immunity processes, such as phagocytosis and the induction of antimicrobial peptide (AMP) production. AMP regulation includes systemic responses and local responses, at surface epithelia that contact with the external environments.

At pupariation, *Drosophila melanogaster* increases dramatically the expression of three AMP genes: *drosomycin (drs)*, *drosomycin-like 2 (drsl2)* and *drosomycin-like 5 (drsl5)*. Using *D. melanogaster*, we show that the expression of *drs* at pupariation is dependent on ecdysone signalling in the fat body. This systemic immune response involving *drs* expression in the fat body operates via the ecdysone downstream target, Broad-Z4. In parallel, ecdysone also regulates local responses, specifically through the activation of *drsl2* expression in the gut. Finally, we establish the relevance of this ecdysone dependent AMP expression for the control of bacterial persistence by showing that flies lacking *drs* expression in the fat body have higher bacterial persistence over metamorphosis.

Together, our data establishes a new role for ecdysone during pupariation. We propose that the co-option of immune mechanisms by the hormonal cascade responsible for controlling metamorphosis constitutes a pre-emptive mechanism to control bacterial numbers in the pupa and increase developmental success.

INTRODUCTION

Insect development and physiology are mainly regulated by two peripheral hormones: the steroid hormone, 20-hydroxyecdsyone (hereafter, ecdysone) and the sesquiterpenoid hormone, juvenile hormone (JH) [1, 2]. In vertebrates, steroid hormones and their nuclear receptors are key regulators of systemic immune responses, namely through the enhancement of inflammation [3]. Similarly, in insects, steroid hormones are also known to be involved in innate immunity [4–9]. In insects with complete metamorphosis (holometabolans), AMP genes are usually silent in the absence of an immune challenge and their expression is induced upon injury and/or infection [10]. However, indirect evidence suggests that AMPs may be up-regulated at specific developmental stages irrespective of pathogen presence. Studies in the moth *Manduca sexta* show that expression of lysozyme and of the AMPs *cecropin A*, *cecropin B* and *hemolin* increases dramatically during metamorphosis, although it is not clear what regulates this increase [11–13]. This transcriptional AMP peak occurs before the end of histolysis of the larval midgut, that is, before cells are potentially exposed to any pathogen or bacterium. In this context, the accumulation of AMPs at the end of the feeding stage may constitute a response against the possible harmful effects inside the pupal body of bacteria carried-over from the gut remodelling process [14–17]. Indeed, studies in dipterans show that some species of bacteria can persist inside the host throughout metamorphosis from the larval guts [16–19]. In *Drosophila melanogaster,* up-regulation of several *drosomycin* (*drs*) paralogues has also been reported at pupariation [20]. Furthermore, five different AMPs (*diptericin, drs, attacin-A, metchnikowin* and *cecropin A1*) were shown to contain putative *cis*-regulatory elements for binding of the functional ecdysone receptor complex, suggesting an ecdysone-dependent control of their expression [21]. Importantly, this association between ecdysone signalling and immunity at this particular stage of development is not restricted to the humoral arm of the immune response. An ecdysone-dependent increase in phagocytic capacity of plasmatocytes at pupariation has been described and hypothesized to be particularly important given the higher predisposition for infection at the pupal stage [6].

In *D. melanogaster*, RNAseq analysis revealed a peak in the expression of AMPs at pupariation, of which around 95% consists of three AMPs: *drs, drosomycin-like 2 (drsl2)* and *drosomycin-like 5 (drsl5)* (**Fig. S1,** [20]). Considering the temporal coincidence between this phenomenon and the ecdysone peak at pupariation to regulate metamorphosis, we hypothesized that ecdysone signalling controlling development was co-opted to regulate the expression of AMPs at pupariation as a pre-emptive mechanism to control bacterial proliferation and prevent possible infections during metamorphosis associated with the gut reconstruction. We demonstrate that ecdysone regulates the expression of AMPs at pupariation, both systemically (by regulating the expression of *drs*) and locally in the midgut (by regulating the expression of *drsl2*). Importantly, the systemic expression of *drs* has an impact on the number of bacteria that persists and proliferates during metamorphosis, corroborating the hypothesis that the AMP peak has a function in controlling bacteria during metamorphosis. Furthermore, we show that this association between the expression of *drs* and ecdysone occurs solely at pupariation and through the early ecdysone-response gene Broad-Z4, which is only expressed at the end of the feeding stage [22].

RESULTS

drs, drsl2 and drsl5 are expressed at pupariation in a bacteria-independent manner

Around 95% of the AMP peak at pupariation consists of three AMPs: *drs, drsl2* and *drsl5*. Due to the gut remodelling at metamorphosis, an inefficient gut purge could lead to the activation of the canonical immunity pathways, leading to an AMP peak. To test this, we determined the expression of *drs, drsl2* and *drsl5* in mid-L3 larvae (L3) and newly pupariated (P0) pupae of germ-free (GF) or standardly-raised individuals by qPCR. We

confirmed that expression increases at pupariation for all AMPs tested (**Figure 1(A), Fig.S2(A)**), including *drs*

(lsmean, pairwise~{w-};+;+ | Stage*drs<0.0001), drsl2 (lsmeans(pairwise~{w-};+;+|Stage*drsl2<0.0001)) and

drsl5 (lsmeans(pairwise~{w-};+;+|Stage*drsl5<0.000)). This increment is not lost even under GF conditions with

the putative exceptions of *cecropinA1* and *defensin*. Importantly, the three main AMPs under scrutiny, *drs*, *drsl2* and

drsl5, show a marked independence of bacteria presence for their expression at this transition

(lsmeans(pairwise~{w-};+;+|Stage*drs<0.0001); lsmeans(pairwise~{w-}; +;+|Stage*drsl2<0.0001);

lsmeans(pairwise~{w-};+;+|Stage*drsl5<0.0001)) (**Figure 1(A),** Table S1).

Together, these results reject the hypothesis that the AMPs expression at pupariation is dependent on

bacterial induction and suggest that it may be hard-wired to the developmental programme.

<u>Ecdysone regulates the systemic expression of *drs*</u>

We focused primarily on two main sources of systemic AMP expression, the fat body and haemocytes. We

manipulated ecdysone signalling activity in these tissues, simultaneously and specifically, to ensure no perturbation

of normal development and pupariation timing. For that we used *Cg-Gal4* combined with a *tub-Gal80[ts]*, a thermo-

sensitive allele of *Gal80* under the control of the tubulin promoter (hereafter *Cg[ts]*), to drive the expression of a

dominant negative ecdysone receptor under the control of the UAS sequence [23] *(EcR-A DN W650A TP3,* hereafter

EcRDN). The reduction of ecdysone sensitivity was triggered from the onset of L3 or at wandering stage, resulting

in elimination of *drs* peak at P0, compared to control conditions (**Figure 1(B)**,

(*lsmeans(pairwise~Stage|Cg[ts]>mcD8GFP)=0.0001*, lsmeans(pairwise~Stage|Cg[ts]>EcRDN)=0.0017,

lsmeans(pairwise~Genotype|P0)<0.0001)). However, the expression of *drsl2* and *drsl5* (**Figure 1(B)**, Table S2) and

of the other AMPs (**Fig. S2(B)**, Table S2) were not eliminated. Furthermore, the expression of *dif*, the NF-κB factor

that triggers *drs* expression upon activation of the Toll pathway, was not altered when ecdysone sensitivity was

decreased (**Figure S2(C),** lsmeans(pairwise~Stage|*Cg[ts]>mcD8GFP)*=0.0162,

lsmeans(pairwise~Stage|Cg[ts]>EcRDN)=0.0015, lsmeans(pairwise~Genotype|P0)=0.3978)**,** suggesting that the

expression of *drs* at pupariation is independent of the action of the NF-κB factor.

This was further confirmed by targeting haemocytes and fat body independently, using *Hemolectin-Gal4*

(*Hml-ΔGal4*) for the haemocytes and *Adipokinetic hormone receptor-Gal4* (*AkhR-Gal4*) for the fat body. When

ecdysone sensitivity was reduced in the haemocytes but not in the fat body, expression of *drs*, *drsl2* and *drsl5*, as

well as of other 4 AMPs, was unaffected (*lsmeans(pairwise~Stage|Hml>mcD8GFP)<0.0001*,

lsmeans(pairwise~Stage|Hml>EcR DN)<0.0001), for the three genes), (**Fig. S3(A)** and **S3(B),** Table S3)**.**

However, although reduced ecdysone sensitivity in the fat body (*AkhR>EcRDN*), did not affect the expression of

drsl2 and *drsl5* (**Fig. S3(C)**, (*lsmeans(pairwise~Stage|AkhR>mcD8GFP)<0.0001*,

lsmeans(pairwise~Stage|AkhR>EcRDN)<0.0001), for both genes)) or of other AMPs (**Figure S3(D)**, Table S4), it

resulted in decreased expression of *drs* at pupariation. (**Fig. S3(C),** *lsmeans(pairwise~Stage|AkhR>mcD8GFP)*

<0.0001, lsmeans(pairwise~Stage|AkhR>EcR DN) =0.6709)).

Altogether, these results indicate that *drs* expression - but not *drsl2* or *drsl5* - is systemically regulated by

ecdysone signalling specifically in the fat body at pupariation.

Figure 1 – Expression of AMPs at pupariation is developmentally regulated by ecdysone, both systemic and locally. (A) The expression levels of *drs*, *drsl2* and *drsl5* increase at pupariation and are not lost in GF conditions. (B) Foldchange increase in *drs* expression (but not of *drsl2* or *drsl5*) at pupariation was eliminated when ecdysone sensitivity is decreased specifically in the fat body and haemocytes simultaneously. (C) Manipulation of ecdysone sensitivity specifically in the midgut reduced the expression of *drsl2*, but not of *drs* or *drsl5*. (D) Knock-down of *drsl2* specifically in the midgut resulted in an overall decrease in the expression of this AMP. Each dot represents a sample of five pooled individuals; the lines connect the median of the samples at L3 and P0; different letters represent statistically significant differences in foldchange. Foldchange was determined using the ΔΔCT method.

Local expression of *drsl2* in the midgut is regulated by ecdysone

Having determined that the expression of these other AMPs, particularly *drsl2* and *drsl5*, are developmentally regulated, but not through the fat body, we turned our attention to two of the most important tissues for local immune response: the epithelial layers of the gut and trachea. First, we overexpressed *EcRDN* specifically in the most abundant cells of the midgut, the enterocytes (*Mex>EcRDN*). Unlike the fat body, the gut is not responding to the pupariation ecdysone pulse by increasing *drs* expression (**Figure 1(C)**). In contrast, decreased ecdysone sensitivity specifically in the midgut resulted in a drastic reduction (*circa* 600-fold) of *drsl2* expression,

131 very close to the L3 basal levels (**Figure 1(C)**, *lsmeans(pairwise~Genotype|P0)<0.0001;*

132 *lsmeans(pairwise~Stage|Mex>EcRDN) <0.0001)).* Moreover, knock-down of *drsl2* specifically in the midgut

133 (*MexGal4>ds-drsl2*) resulted in an overall decrease in the expression of this AMP at pupariation (**Figure 1(D)**,

134 *lsmeans(pairwise~Genotype|P0)<0.0237; lsmeans(pairwise~Stage|Mex>ds-drsl2) <0.0001)),* which indicates that

135 the midgut is the main source for the production of *drsl2* at this stage. Other tested AMPs also appear to be, at least

136 partially, regulated by ecdysone signalling in the midgut. Under ecdysone-signalling down-regulation conditions,

137 significant decrease in mRNA levels were observed for: *cecropin A1 (lsmeans(pairwise~Genotype|P0)<0.0001;*

138 *lsmeans(pairwise~Stage|Mex>EcRDN)=0.0038), defensin (lsmeans (pairwise~Genotype|P0)<0.0001;*

139 *lsmeans(pairwise~Stage|Mex>EcR DN)=0.3866), metchnikowin (lsmeans(pairwise~Genotype|P0)=0.0060;*

140 *lsmeans (pairwise~Stage|Mex>EcRDN<0.0001), and drsl3 ((lsmeans(pairwise~Genotype|P0)=0.0008;*

141 *lsmeans(pairwise~Stage|Mex>EcRDN)=0.2084)*, **Fig. S3(E)**)). Interestingly, no difference was detected in the

142 expression of *drsl5*, one of the three main players in the AMP peak (**Figure 1(C)**, *lsmeans*

143 *(pairwise~Genotype|P0)<0.4479; lsmeans(pairwise~Stage|Mex>EcRDN)<0.0001)).*

144   On the other hand, decreased ecdysone sensitivity in the trachea (*btl>EcRDN*) did not have an impact on

145 the expression of any of the three major AMPs under study (**Fig. S3(F),** Table S6).

146   Together, these results show that the local sensing of the ecdysone peak at the onset of pupariation in, at

147 least the gut epithelium, controls the production of AMPs. The local and systemic mechanisms that share ecdysone

148 control account for the expression of the genes that constitute about 70% of the pupariation AMP peak.

149 Furthermore, the data also point towards an alternative mechanism for regulating the expression of *drsl5* at

150 pupariation, which is either independent of ecdysone or is regulated by this hormone in an alternative tissue or cell

151 type that were not tested.

153 <u>Systemic expression of *drs* is necessary to reduce bacteria derived from imperfect larval gut purge</u>

154   Before pupariation, larvae purge their gut contents to undergo metamorphosis. Any imperfection in this

155 process creates the conditions in the pupa for a putatively rampant infection. Thus, we examined whether or not gut

156 purge eliminates all bacteria from the gut. Under our standard rearing conditions, the mortality between P0 and

157 pharate adult is negligible, which allows us to infer and correlate the bacterial numbers detected in each of these

158 developmental stages (**Fig. S4(A),** Table S7).

159   After gut purge, approximately 5% of control P0 individuals contained bacteria in the order of the

160 thousands and another 12% in the order of the hundreds (**Figure 2(A)**). However, at the pharate adult stage, these

161 two classes decreased in number alongside an increase in the number of individuals without detectable bacteria

162 (**Figure 2(B)**). In contrast, *drs* null mutants showed an increase rather than a decrease in the number of pupae with

163 more bacteria between these two stages (**Figures 2(C)** and **2(D)**), supporting that Drs is necessary to eliminate gut

164 derived bacteria during metamorphosis.

165   Using a resampling strategy, we determined that most coefficients associated with the control line were

166 negative (**Figure 2(G)** (*Median: -2.279 (IQR: -2.682 -1.803); One-way Wilcoxon test for median<0: <2.2e-16, r=*

167 *0.864)* and **Figure 2(I)** (*Median: -2.317 (IQR: -2.907 -1.858); One-way Wilcoxon test for median<0: <2.2e-16,*

168 *r=0.866))*. In contrast, the coefficients associated with *drs* null mutant were mostly positive for both bacteria types

169 (**Figure 2(H)** (*(Median: 0.596 (IQR: 0.359; 0.757); One-way Wilcoxon test for median> 0: <2.2e-16, r= 0.845)* and

170 **Figure 2(J)** (*Median: 2.055 (IQR: 1.897; 2.192); One-way Wilcoxon test for median> 0: <2.2e-16, r=0.866)).*

171 These results suggest that pupae have the bacteria eliminated in a process that is dependent on an increase of AMPs

expression at pupariation. This process is necessary for the control the persistence of bacteria originating from gut
purge imperfections, throughout metamorphosis.

When ecdysone signalling activity is decreased in the fat body and haemocytes, a similar pattern can be
observed: there is a higher proportion of individuals with larger loads of persisting bacteria (**Figure 2(E-F)**). The
resampling strategy revealed that *Cg[ts]>EcR DN* samples have positive slope values for both *Lactobacillus*
(**Figure 2(K)**, *Median: 2.746 (IQR: 2.562; 2.930); One-way Wilcoxon test for median>0: <2.2e-16, r=0.866)* and
Acetobacter (**Figure 2(L)**, *Median: 1.981 (IQR: 1.795; 2.184); One-way Wilcoxon test for median>0: <2.2e-16,*
r=0.866), which is in accordance with increased bacterial loads during metamorphosis. Importantly, *Mex>EcRDN*
individuals do not have increased bacterial loads during metamorphosis, suggesting that the local immune response
does not have a measurable antibacterial effect relatively to the systemic response that is still operating in these
conditions (**Fig. S3(B-E)**).

184
185
Figure 2 – Expression of AMPs at pupariation is developmentally regulated by ecdysone, both systemic and locally. (A-F) Quantification
in P0 pupae and pharate adults of *Lactobacillus* (L), growing in selective MRS medium, and *Acetobacter* (A), growing on selective D-Mannitol.
(A-B) In control flies (*{w-};+;+*)**,** the number of pupae with more bacteria decreases between P0 and pharate adult stages. **(C-D)** In the *drs*
mutant line (*{w-};+; Drs r{w+}*), there are more individuals with higher loads of bacteria persisting from P0 to pharate adult. **(E-F)** When
ecdysone sensitivity is decreased in fat body and haemocytes, the number of both bacteria types increases in P0 and pharate adults. Y-axis
represents the proportion of individuals in each bacterial count category. **(G-H)** Slopes for the number of *Lactobacillus* in control and the mutant
lines. **(G)** Control samples have negative slopes, representative of a decrease in the number of bacteria between stages. **(H)** *drs* mutant samples
have positive slope values, representative of an increase in the number of bacteria between the stages. **(I-J)** Slopes for the number of *Acetobacter*
in the control and the mutant lines. **(I)** Control samples distribution show negative slopes, i.e., a decrease in the number of bacteria between
stages. **(J)** *drs* null mutant showed a positive slope values. **(K-L)** Coefficient (slopes) distribution in *Cg[ts]>EcRDN* samples of *Lactobacillus*
(K) and *Acetobacter***.** Doted lines represent the 0.05 and 0.95 quantiles; full line represents the mean slope**.**
197

<u>Br is involved in systemic *drs* induction through the fat body at pupariation</u>

Our data suggests that the systemic expression of Drs is more important than the local production of AMPs

for the control of bacterial persistence during metamorphosis. Hence, we focused our efforts in trying to uncover the

molecular regulation of temporal *drs* expression.

Previous RNAseq data showed that the expression of *drs* markedly increases at pupariation (**Figure S1,**

[20]). We further confirmed that the expression of *dhr3* (a gene directly regulated by the ecdysone-ecdysone

receptor complex) and that of *drs* are tightly correlated and peak exclusively at pupariation (**Figure S5(A)**).

Importantly, this robust expression of *drs* at the onset of metamorphosis seems to be a general feature regardless of

their genetic background (**Figure S5(B)**).

br is a direct target of ecdysone receptor, characterized as a holometabolan pupal specifier with increased

expression before pupariation [22, 24, 25]. We examined the association between *br* and the ecdysone-sensitive

systemic induction of *drs* at metamorphosis by knocking-down *br* specifically in the fat body and haemocytes. This

led to a significant decrease in the expression of *drs* at pupariation, when compared to the control (**Figure 3(A),**

lsmeans (pairwise~Genotype|P0)=0.0510)), suggesting a role for *br* in the regulation of AMP expression. To further

explore this possibility, we simultaneously reduced ecdysone signalling and restored *br* in the haemocytes and fat

body. This was sufficient to rescue the expression of *drs* to control levels at P0 (**Figure 3(B),**

lsmeans(pairwise~Genotype|P0)= 0.3068). Furthermore, we generated two different conditional CRISPR-mediated

knock-out constructs [26, 27], one targeting a common region to all *br* isoforms and, the other, a region specific to

br-Z4, previously associated with AMP expression in a different species [28]. Knock-out of all *br* isoforms or of *br-*

Z4 alone in haemocytes and fat body, led to the elimination of the *drs* peak at pupariation (**Figure 3(C),** *lsmeans*

(pairwise~Stage|Cg>+;Cas9;+)<0.0001;lsmeans(pairwise~Stage| Cg>Cas9;br KO)=0.0001,

lsmeans(pairwise~Stage|Cg>Cas9;br-Z4 KO)=0.4691), supporting that Br is involved in the regulation of *drs*

expression at pupariation downstream of the ecdysone receptor complex.

To further explore this regulation of *drs* expression by Br-Z4, we used the JASPAR

(http://jaspar.genereg.net) matrix MA0013.1 to run a comprehensive search for Br-Z4 binding sites over 2 kb of

putative promoter region that is upstream of the coding sequence of *drs*, obtained from Flybase (http://flybase.net/),

using FIMO software [29] (**Fig. S5(C)**). We identified three putative Br-Z4 binding sites (p-value<0.0001) in the

predicted 5' promoter region of *drs* (**Figure S5(D)**).

Together, these results support the idea that Br, specifically isoform Br-Z4, is involved in the regulation of

the expression of *drs* at pupariation.

235
236

DISCUSSION

Generally, in Holometabola, the expression of AMPs is induced by sensing of specific molecular patterns that trigger the activation of the Toll or Imd pathways [10, 30–33]. However, there is increasing evidence of alternative regulators of innate immunity, including the moulting hormone, ecdysone [4–9]. In vertebrates, it has been known for many years that the immune system influences the secretion of hormones and that these, in turn, can influence immunity [3].

In this study we explored the association between ecdysone and the increased expression of *drs* and its paralogues at pupariation. We unravelled a novel association between the endocrine and immune systems in *D. melanogaster*, which act both systemically and locally during metamorphosis. Importantly, we could determine that the systemic expression of *drs* has an effect on the number of bacteria during metamorphosis, as both mutant and ecdysone-reduced sensitivity individuals showed an increase rather than a decrease in the number of bacteria between P0 and pharate adult stages. Interestingly the effect was detected for both Gram-positive (*Lactobacillus*) and Gram-negative (*Acetobacter*) bacteria. Although Drs is generally characterized as an antifungal AMP, previous studies already demonstrated that it can also be induced by both Gram-positive and Gram-negative bacteria [33–39]. As such, the unexpected higher impact of the lack of *drs* on *Acetobacter* could actually be explained by its possible direct impact on Gram-negative bacteria. Importantly, the humoral arm of defence is not the only one acting at this stage of development. Regan and colleagues demonstrated that at pupariation, ecdysone is required to activate

haemocytes and to orchestrate the active migration of these cells towards wounding sites [6]. Therefore, at this stage

both ecdysone-regulated cellular and humoral mechanisms act in synergy to decrease the probability of infection.

Because the lack of local expression of *drsl2* did not affect bacterial persistence *per se* in our conditions,

we focused our efforts on trying to disentangle the molecular regulation of *drs* expression in the fat body. We

showed that the expression of *drs* at this stage depends upon the expression in the fat body of the ecdysone

signalling target, *br-Z4*. This is in accordance to previous indications that Br is essential for the PGRP-LC-

independent hormonal regulation of *drs* [4]. Furthermore, similar ecdysone-mediated immune activation through Br

has been previously observed. Studies in the silkworm *B. mori* showed that during metamorphosis, ecdysone

activates the transcription of lysozyme via Br-Z2 [40] and of the AMP *lebocin* via Br-Z4 [28]. This regulatory

mechanism through Br may cause the stage specificity. Although ecdysone controls all the moults throughout

development, *drs* expression increases in response to this hormone specifically at pupariation, when Br is expressed

in Holometabola [22, 41]. The regulation of AMPs by Br is an interesting feature, as it can also have implications on

the interpretation of AMP expression throughout development between holo- and hemimetabolan insects. Johnston

[42] demonstrated that these two insect groups have different AMP expression patterns, with holometabolans

showing a strong upregulation of immunity that coincides with metamorphosis timing and Br expression. In

contrast, hemimetabolans do not undergo a dramatic increase in immunity, possibly correlating with much less

drastic changes in gut morphology during progressive metamorphosis. In accordance, there is indication that insects

with complete metamorphosis suffer a reduction in density, diversity and altered composition of bacterial

populations during metamorphosis [43–45], whereas those with incomplete metamorphosis show a constant increase

in microbial density and diversity throughout development [46]. These data, as ours, are consistent with the

previously proposed hypothesis that complete metamorphosis elicits a prophylactic immune response [12, 13].

Alterations in immunity at life stages with higher risk of infection seem to be a general feature of

development, finding some parallels in both vertebrates and invertebrates. In *D. triauraria*, genes of the *drs* family

are induced during diapause, which is also considered to be a more susceptible period [47]. Most interestingly, data

from *Xenopus* also point towards an association between the timing of metamorphosis and a shift in the immune

response [48]. For example, Major Histocompatibility Complex I (MHC class I) molecules, which bind to peptide

fragments derived from pathogens, are first detected at the beginning of metamorphosis [49] and the expression of

MHC class II molecules is not triggered if metamorphosis is chemically blocked [50].

Together, our data suggests a regulatory mechanism through which ecdysone co-opted the immune system

to enhance the success of metamorphic processes. This study reveals a novel association between the endocrine and

immune systems in *D. melanogaster*, which act both systemically and locally during metamorphosis. This

recruitment of immunity mechanisms by the holometabolan metamorphosis programme presents other interesting

features. Three AMPs explain around 95% of the peak observed at the L3 to pupal transition. Interestingly, these

three AMPs have been co-opted differently and complementarily to produce, what we interpret as a full-fledged

immune prophylactic response. One of them, *drsl2*, was recruited to one of the main exposed surface epithelia of

larva, the midgut, that undergoes extensive remodelling at metamorphosis. Another of the AMPs with increased

expression, *drs*, is produced in the fat body at metamorphosis initiation, executing a putatively systemic action. The

local and systemic mechanisms that share ecdysone control and that were explored in this work account for the

expression of the genes that constitute close 70% of the pupariation AMP peak.

This specificity and multi-layered developmentally-regulated immune response underscores the sophistication of

the co-option process and hints as a possible contribution of this response to the success of metamorphosis as a whole.

REFERENCES

1. Nijhout HF (1994) Insect hormones. Princeton University Press, Princept

2. Gilbert, L. I., Rybczynski, R., Tobe SS (1996) Endocrine cascade in insect metamorphosis. In: Gilbert, L.I., Tata, J.R., Atkinson BG (ed) Metamorphosis, postembryonic reprogramming of gene expression in amphibian and insect cells. Academic Press Inc, pp 59–107

3. Webster JI, Tonelli L, Sternberg EM (2002) Neuroendocrine regulation of immunity. Annu Rev Immunol 20:125–163. https://doi.org/10.1146/annurev.immunol.20.082401.104914

4. Rus F, Flatt T, Tong M, et al (2013) Ecdysone triggered PGRP-LC expression controls Drosophila innate immunity. EMBO J 32:1626–1638. https://doi.org/10.1038/emboj.2013.100

5. Flatt T, Heyland A, Rus F, et al (2008) Hormonal regulation of the humoral innate immune response in Drosophila melanogaster. J Exp Biol 211:2712–2724. https://doi.org/10.1242/jeb.014878

6. Regan JC, Brandão AS, Leitão AB, et al (2013) Steroid hormone signaling is essential to regulate innate immune cells and fight bacterial infection in Drosophila. PLoS Pathog 9:. https://doi.org/10.1371/journal.ppat.1003720

7. Dimarcq JL, Imler JL, Lanot R, et al (1997) Treatment of l(2)mbn Drosophila tumorous blood cells with the steroid hormone ecdysone amplifies the inducibility of antimicrobial peptide gene expression. Insect Biochem Mol Biol 27:877–886. https://doi.org/10.1016/S0965-1748(97)00072-6

8. Wang JL, Chen L, Tang L, et al (2014) 20-hydroxyecdysone transcriptionally regulates humoral immunity in the fat body of Helicoverpa armigera. Insect Mol Biol 23:842–856. https://doi.org/10.1111/imb.12131

9. Nunes C, Sucena É, Koyama T (2020) Endocrine regulation of immunity in insects. FEBS J 1–20. https://doi.org/10.1111/febs.15581

10. Cociancich S, Bulet P, Hetru C, Hoffmann JA (1994) The inducible antibacterial peptides of insects. Parasitol Today 10:132–139. https://doi.org/10.1016/0169-4758(94)90260-7

11. Yu XQ, Kanost MR (1999) Developmental expression of Manduca sexta hemolin. Arch Insect Biochem Physiol 42:198–212. https://doi.org/10.1002/(SICI)1520-6327(199911)42:3<198::AID-ARCH4>3.0.CO;2-G

12. Russell VW, Dunn PE (1991) Lysozyme in the midgut of Manduca sexta during metamorphosis. 80:67–80

13. Russell V, Dunn PE (1996) Antibacterial proteins in the midgut of Manduca sexta during metamorphosis. J Insect Physiol 42:65–71. https://doi.org/10.1016/0022-1910(95)00083-6

14. Hakim RS, Baldwin K, Smagghe G (2010) Regulation of midgut growth, development, and metamorphosis. Annu Rev Entomol 55:593–608. https://doi.org/10.1146/annurev-ento-112408-085450

15. Judy KJ, Gilbert LI (1970) Histology of the alimentary canal during the metamorphosis of Hyalophora cecropia (L.). J Morphol 131:277–299. https://doi.org/10.1002/jmor.1051310304

16. Bacot A (1911) On the persistence of Bacilli in the gut of an Insect during metamorphosis. Parasitology 4:497–500

17. Leach JG (1934) The method of survival of bacteria in the puparia of the seed-corn maggot (Hylemyia Cilicrura Rond.). Zeitschrift für Angew Entomol 20:150–161. https://doi.org/10.1111/j.1439-0418.1934.tb00358.x

18. Bakula M (1969) The persistence of a microbial flora during postembryogenesis of Drosophila melanogaster. J Invertebr Pathol 14:365–374. https://doi.org/10.1016/0022-2011(69)90163-3

19.   Duneau DF, Lazzaro BP (2018) Persistence of an extracellular systemic infection across metamorphosis in a
holometabolous insect. Biol Lett 14:10–13. https://doi.org/10.1098/rsbl.2017.0771

20.   Thurmond J, Goodman JL, Strelets VB, et al (2019) FlyBase 2.0: The next generation. Nucleic Acids Res
47:D759–D765. https://doi.org/10.1093/nar/gky1003

21.   Zhang Z, Palli SR (2009) Identification of a cis-regulatory element required for 20-hydroxyecdysone
enhancement of antimicrobial peptide gene expression in Drosophila melanogaster. Insect Mol Biol 18:595–
605. https://doi.org/10.1111/j.1365-2583.2009.00901.x

22.   Zhou X, Riddiford LM (2002) Broad specifies pupal development and mediates the "status quo" action of
juvenile hormone on the pupal-adult transformation in Drosophila and Manduca. Development 129:2259–
2269

23.   Cherbas L, Hu X, Zhimulev I, et al (2003) EcR isoforms in Drosophila: Testing tissue-specific requirements
by targeted blockade and rescue. Development 130:271–284. https://doi.org/10.1242/dev.00205

24.   Truman JW, Riddiford LM (1999) The origins of insect metamorphosis. Nature 401:447–452

25.   Belles X (2011) Origin and evolution of insect metamorphosis. eLS 1–11.
https://doi.org/10.1002/9780470015902.a0022854

26.   Port F, Chen HM, Lee T, Bullock SL (2014) Optimized CRISPR/Cas tools for efficient germline and
somatic genome engineering in Drosophila. Proc Natl Acad Sci U S A 111:.
https://doi.org/10.1073/pnas.1405500111

27.   Xue Z, Wu M, Wen K, et al (2014) CRISPR/Cas9 mediates efficient conditional mutagenesis in Drosophila.
G3 Genes, Genomes, Genet 4:2167–2173. https://doi.org/10.1534/g3.114.014159

28.   Mai T, Chen S, Lin X, et al (2017) 20-hydroxyecdysone positively regulates the transcription of the
antimicrobial peptide, lebocin, via BmEts and BmBR-C Z4 in the midgut of Bombyx mori during
metamorphosis. Dev Comp Immunol 74:10–18. https://doi.org/10.1016/j.dci.2017.04.002

29.   Grant CE, Bailey TL, Noble WS (2011) FIMO: Scanning for occurrences of a given motif. Bioinformatics
27:1017–1018. https://doi.org/10.1093/bioinformatics/btr064

30.   Lemaitre B, Hoffmann J (2007) The host defense of Drosophila melanogaster. Annu Rev Immunol 25:697–
743. https://doi.org/10.1146/annurev.immunol.25.022106.141615

31.   Dunn PE (1990) Humoral immunity in insects. Bioscience 40:738–744. https://doi.org/10.2307/1311506

32.   Hultmark D (1993) Immune reactions in Drosophila and other insects: a model for innate immunity. Trends
Genet 9:178–183. https://doi.org/10.1016/0168-9525(93)90165-E

33.   Tzou P, Ohresser S, Ferrandon D, et al (2000) Tissue-specific inducible expression of antimicrobial peptide
genes in Drosophila surface epithelia. 13:737–748

34.   Lemaitre B, Reichhart JM, Hoffmann JA (1997) Drosophila host defense: Differential induction of
antimicrobial peptide genes after infection by various classes of microorganisms. Proc Natl Acad Sci U S A
94:14614–14619. https://doi.org/10.1073/pnas.94.26.14614

35.   Leulier F, Rodriguez A, Khush RS, et al (2000) The Drosophila caspase Dredd is required to resist Gram-
negative bacterial infection. EMBO Rep 1:353–358. https://doi.org/10.1093/embo-reports/kvd073

36.   Foley E, O'Farrell PH (2003) Nitric oxide contributes to induction of innate immune responses to gram-
negative bacteria in Drosophila. Genes Dev 17:115–125. https://doi.org/10.1101/gad.1018503

37.   Tanji T, Hu X, Weber ANR, Ip YT (2007) Toll and IMD pathways synergistically activate an innate
Immune Response in Drosophila melanogaster. Mol Cell Biol 27:4578–4588.

377    https://doi.org/10.1128/mcb.01814-06

378 38. Li Y, Li S, Li R, et al (2017) Genome-wide miRNA screening reveals miR-310 family members negatively

379    regulate the immune response in Drosophila melanogaster via co-targeting Drosomycin. Dev Comp

380    Immunol 68:34–45. https://doi.org/10.1016/j.dci.2016.11.014

381 39. Zhan MY, Yang PJ, Rao XJ (2018) Molecular cloning and analysis of PGRP-L1 and IMD from silkworm

382    Bombyx mori. Elsevier Inc

383 40. Ma H, Abbas MN, Zhang K, et al (2019) 20-Hydroxyecdysone regulates the transcription of the lysozyme

384    via Broad-Complex Z2 gene in silkworm, Bombyx mori. Dev Comp Immunol 94:66–72.

385    https://doi.org/10.1016/j.dci.2019.01.014

386 41. Erezyilmaz DF, Riddiford LM, Truman JW (2006) The pupal specifier broad directs progressive

387    morphogenesis in a direct-developing insect. Proc Natl Acad Sci U S A 103:6925–6930.

388    https://doi.org/10.1073/pnas.0509983103

389 42. Johnston PR, Paris V ronique, Rolff J (2019) Immune gene regulation in the gut during metamorphosis in a

390    holo- versus a hemimetabolous insect. Philos Trans R Soc B Biol Sci 374:1–6.

391    https://doi.org/10.1098/rstb.2019.0073

392 43. Johnston PR, Rolff J (2015) Host and Symbiont Jointly Control Gut Microbiota during Complete

393    Metamorphosis. PLoS Pathog 11:1–11. https://doi.org/10.1371/journal.ppat.1005246

394 44. Hammer TJ, McMillan WO, Fierer N (2014) Metamorphosis of a butterfly-associated bacterial community.

395    PLoS One 9:. https://doi.org/10.1371/journal.pone.0086995

396 45. Kim JK, Han SH, Kim CH, et al (2014) Molting-associated suppression of symbiont population and up-

397    regulation of antimicrobial activity in the midgut symbiotic organ of the Riptortus-Burkholderia symbiosis.

398    Dev Comp Immunol 43:10–14. https://doi.org/10.1016/j.dci.2013.10.010

399 46. Sudakaran S, Salem H, Kost C, Kaltenpoth M (2012) Geographical and ecological stability of the symbiotic

400    mid-gut microbiota in European firebugs, Pyrrhocoris apterus (Hemiptera, Pyrrhocoridae). Mol Ecol

401    21:6134–6151. https://doi.org/10.1111/mec.12027

402 47. Daibo S, Kimura MT, Goto SG (2001) Upregulation of genes belonging to the drosomycin family in

403    diapausing adults of Drosophila triauraria. Gene 278:177–184. https://doi.org/10.1016/S0378-

404    1119(01)00713-2

405 48. Robert J, Ohta Y (2009) Comparative and developmental study of the immune system in Xenopus. Dev Dyn

406    238:1249–1270. https://doi.org/10.1002/dvdy.21891

407 49. Flajnik MF, Du Pasquier L (1988) MHC class I antigens as surface markers of adult erythrocytes during the

408    metamorphosis of Xenopus. Dev Biol 128:198–206. https://doi.org/10.1016/0012-1606(88)90282-5

409 50. Rollins-Smith LA, Blair P (1990) Expression of class II major histocompatibility complex antigens on adult

410    T cells in Xenopus is metamorphosis-dependent. Dev Immunol 1:97–104

MATERIALS AND METHODS

Drosophila Stocks and Husbandry

If not further noticed, *w[1118]* flies were obtained from Dr. Luís Teixeira (Instituto Gulbenkian de Ciência, Portugal) and used as a wild-type strain. The origin of other fly strains is mentioned in SI. All fly lines we kept at 25ºC in *Drosophila* cornmeal food (Supplementary Information), in a 12L:12D photoperiod. For the ecdysone decreased sensitivity experiment and tissue specific CRISPR, animals were maintained at 17ºC until the moult to the L3 and then transferred to 29ºC to activate Gal4 activity.

CRISPR Knock-out Flies Generation

The tissue-specific UAS *br* CRISPR knockout constructs were cloned into pCFD6 vector (www.crisprflydesign.org, Addgene #73915) according to manufacturer's instructions. Four independent gRNA constructs were cloned to target introns (see primers listed in **Table S9**) of the *br* core region genomic DNA sequence for *br* common construct and Z4 isoform specific sequence for *br-Z4* construct to target only native genes but not transgenes. After sequenced, the constructs were inserted on either the second chromosome in a recipient line carrying phiC31 integrase and attP40 landing cite with an in-house modified line (*y,w, P{y+ nos-int.NLS}; P{CaryP y+}attP40*) or the third chromosome in a recipient line carrying phiC31 integrase and attP2 landing site, *y,w, P(y[+].nos-int. NLS); P(CaryP)attP2* (gift from Dr. Diogo Manoel, Sidra Medical and Research Center, Qatar). Transgenesis was carried out in-house at the Instituto Gulbenkian de Ciência, Portugal.

Larval and pupal staging

Larvae were resynchronized every 2h and kept in normal food for the appropriate time before collection in Trizol. White pre-pupae (P0) were collected directly in Trizol every 20 minutes. All samples were kept at -80ºC until use. More details are available as SI.

P0 and adult pharate collection and platting

P0 samples were collected as aforementioned. Pharate adults were also collected in 20 minutes intervals 48h to 72h after P0 collection and were identified as fully developed adults still to eclose. All samples were surface sterilised before platting onto MRS or D-Mannitol plates. Plates were incubated at 25ºC for 48h. More details as SI.

Statistical analysis for bacterial growth across stages

Growth across stages was assessed with the generation of simulated data from real data. The initial sample was resampled in subsets of 50 entries for 2000 cycles. In each cycle, a negative binomial regression was performed to model the change of bacteria according to the stage, materialized in a regression coefficient, or slope. These slopes were then analysed to see the central tendency of the population. In parallel, the effect size was also calculated. More details on statistical analysis on SI.

Germ-free fly generation

Newly laid eggs were collected into a sterile embryo basket and washed with autoclaved MilliQ water. The eggs were dechorionated, sterilised and transferred to axenic media supplemented with antibiotics. Plates and/or bottles with embryos were kept in a 25ºC inside sterilized containers at 25ºC. After the treatment, the germ-free status was

456 validated by plating surface-sterilized adult flies (from both germ free and control conditions) onto MRS and D-
457 Mannitol media.
458
RNA extraction, cDNA synthesis and real-quantitative PCR
RNA was isolated using the Direct-zol™ RNA MiniPrep Kit (Zymo®), following manufacturer's instructions.
DNase I treatment and cDNA synthesis were preformed using the RQ1 RNASE-FREE DNASE 1* (Promega®) and
the RevertAid H Minus First Strand cDNA Synthesis Kit (Thermo Scientific®), respectively.
qPCR was performed using the SYBR® Green PCR Master Mix (Applied Biosystems®) and ABI 7900HT (Applied
Biosystems). More details on PCR conditions and data analysis on Supplementary Information.
465
466 Br binding sites prediction
467 The DNA binding motif matrix for Br-Z4 was obtained from the JASPAR database (http://jaspar.genereg.net) and
468 the 5' 2kb region of the *drosomycin* gene was acquired in Flybase (http://flybase.org). Both sequences were
469 uploaded in FIMO software (http://meme-suite.org/tools/fimo) [29], using a p-value < 0.0001 as a filter.
470
471

Acknowledgements
We thank Drs. Lynn Riddiford, Marc Dionne, Luís Teixeira, Bruno Lemaitre, François Leulier, Kim Rewitz and
Pedro Domingos for stock sharing. We thank Liliana Vieira and Sandra Crisóstomo for technical assistance, Gastón
Guilgur for transgenesis services and André Barros for help with statistical analysis. Stocks obtained from the
Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study. This work was supported by:
Instituto Gulbenkian de Ciência/Fundação Calouste Gulbenkian; Fundação para a Ciência e a Tecnologia (FCT,
Portugal) to CN (PD/BD/114344/2016); CONGENTO, project LISBOA-01-0145-FEDER-022170, co-financed by
Lisboa Regional Operational Programme (Lisboa 2020), under the Portugal 2020 Partnership Agreement, through
the European Regional Development Fund (ERDF), and Fundação para a Ciência e a Tecnologia (FCT, Portugal).

481
482

Contributions
CN, Conceptualization, Formal analysis, Investigation, Visualization, Methodology, Writing—original draft, review
and editing
TK, Conceptualization, Investigation, Methodology, Writing—review and editing
ES, Conceptualization, Investigation, Funding acquisition, Methodology, Writing— original draft, review and
editing

SUPPLEMENTARY INFORMATION

491   **Supplementary Figure S1** – High-throughput temporal expression data using whole body, with expression levels

492   represented as linear and scaled to maximum expression (publicly available on Flybase http://flybase.org/ [20]).

Supplementary Figure S2 - **(A)** The expression levels of *cecropin A1*, *defensin*, *metchnikowin* and *drsl3* increase at

pupariation and are not lost in GF conditions, albeit to different degrees. **(B)** Reduced expression of ecdysone sensitivity in

the fa body and haemocytes does not affect the expression of *cecropin A1*, *defensin*, *metchnikowin* and *drsl3*. **(C)** *dif*

expression was not affected by the decreased ecdysone sensitivity in the haemocytes and fat body

(*lsmeans(pairwise~Genotype|P0)=0.3878*). Each dot represents a sample of five pooled individuals; the lines connect the

median of the samples at L3 and P0; different letters represent statistically significant differences in foldchange. Foldchange

was determined using the ΔΔCT method.

Supplementary Figure S3 – (A) Reduced ecdysone sensitivity in the haemocytes does not affect the expression of the AMPs under study.

(B) Expression of other 4 AMPs was also not reduced when ecdysone signalling was reduced in the haemocytes. **(C)** Reduced ecdysone

signalling in the fat body resulted in decreased *drs* expression at pupariation (*lsmeans(pairwise~Stage|AkhR>mcD8GFP)<0.0001*,

lsmeans(pairwise~Stage|AkhR>EcR DN) =0.6709), without affecting the expression of *drsl2* and *drsl5*

(*lsmeans(pairwise~Stage|AkhR>mcD8GFP)<0.0001, lsmeans(pairwise~Stage|AkhR>EcRDN)<0.0001*), for both genes). **(D)** The expression

of *cecropin A1*, *defensin*, *metchnikowin* and *drsl3* was unaffected by decreased ecdysone sensitivity specifically in the fat body. **(E)**

Decreased ecdysone signalling in the midgut resulted in significant decrease in mRNA levels for: *cecropin A1*

(*lsmeans(pairwise~Genotype|P0)<0.0001; lsmeans(pairwise~Stage|Mex>EcRDN)=0.0038), defensin (lsmeans*

(*pairwise~Genotype|P0)<0.0001; lsmeans(pairwise~Stage|Mex>EcR DN)=0.3866), metchnikowin*

(*lsmeans(pairwise~Genotype|P0)=0.0060; lsmeans (pairwise~Stage|Mex>EcRDN<0.0001*), and *drsl3*

(*(lsmeans(pairwise~Genotype|P0)=0.0008; lsmeans(pairwise~Stage|Mex>EcRDN)=0.2084*). **(F)** Ecdysone signalling in the trachea does not

affect the expression of any AMP under study.

530
531
532