Positive Selection and Heat-Response Transcriptomes Reveal Adaptive Features of the *Arabidopsis* Desert Relative, *Anastatica hierochuntica*

Gil Eshel[a,1,], Nick Duppen[a,1], Guannan Wang[b], Dong-Ha Oh[b], Yana Kazachkova[a,2], Pawel Herzyk[c], Anna Amtmann[c], Michal Gordon[d,], Vered Chalifa-Caspi[d], Michelle Arland Oscar[e], Shirli Bar-David[f], Amy Marshall-Colon[g], Maheshi Dassanayake[b] and Simon Barak[h]

[a]Albert Katz International School for Desert Studies, Ben-Gurion University of the Negev, Sde Boqer Campus, 8499000, Israel

[b]Department of Biological Sciences, Louisiana State University, Baton Rouge, LA, 70803, USA

[c]Institute of Molecular, Cell and Systems Biology, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8QQ, UK

[d]Bioinformatics Core Facility, The National Institute for Biotechnology in the Negev, Ben-Gurion University of the Negev, Beer-Sheva 8410501, Israel

[e]Blaustein Center for Scientific Cooperation, Ben-Gurion University of the Negev, Sde Boqer Campus, 8499000, Israel

[f]Mitrani Department of Desert Ecology, Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Midreshet Ben-Gurion, 8499000, Israel

[g]Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[h]French Associates' Institute for Agriculture and Biotechnology of Drylands, Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Sde Boqer Campus, 8499000, Israel

[1]These authors contributed equally to the article.

[2]Current address: Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot 7610001, Israel

Authors for correspondence: Simon Barak, Tel: +972-8-6596750. Email: simon@bgu.ac.il; Maheshi Dassanayake, Tel: +1- 225-578-0960, Email: maheshid@lsu.edu

Summary: 199 words
Main body: 6,270 words;
Introduction: 687 words; Materials and Methods: 1,041 words; Results: 2,633 words; Discussion: 1,909 words
No. figures: 7, all color
No. Tables: 1
Supporting information: 6 figures; 24 Datasets; 1 Methods.

Summary

- Plant adaptation to a desert environment and its endemic heat stress is poorly understood at the molecular level. The naturally heat-tolerant Brassicaceae species *Anastatica hierochuntica* is an ideal extremophyte model to identify genetic adaptations that have evolved to allow plants to tolerate heat stress and thrive in deserts.

- We generated an *A. hierochuntica* reference transcriptome and pinpointed extremophyte adaptations by comparing *Arabidopsis thaliana* and *A. hierochuntica* transcriptome responses to heat and identifying positively selected genes in *A. hierochuntica*.

- The two species exhibit similar transcriptome adjustment in response to heat and the *A. hierochuntica* transcriptome does not exist in a constitutive heat "stress-ready" state. Furthermore, the *A. hierochuntica* global transcriptome as well as heat-responsive orthologs, display a lower basal and higher heat-induced expression than in *A. thaliana*. Genes positively selected in multiple extremophytes are associated with stomatal opening, nutrient acquisition, and UV-B induced DNA repair while those unique to *A. hierochuntica* are consistent with its photoperiod-insensitive, early-flowering phenotype.

- We suggest that evolution of a flexible transcriptome confers the ability to quickly react to extreme diurnal temperature fluctuations characteristic of a desert environment while positive selection of genes involved in stress tolerance and early flowering could facilitate an opportunistic desert lifestyle.

Key words: extremophyte, Brassicaceae, desert species, *Anastatica hierochuntica*, heat stress, adaptation, transcriptome, positive selection

Introduction

Plant species inhabiting extreme environments - so-called extremophytes - are able to thrive in some of the most inhospitable environments on Earth that are characterized by severe abiotic stresses. These stresses include drought and temperature extremes in deserts, intense cold in the Antarctic, and habitats both on land and in the sea that are typified by acute salinity (Amtmann, 2009; John et al., 2009; Dassanayake et al., 2010; Oh et al., 2012; Lawson et al., 2014; Farrant et al., 2015; Fan et al., 2018; Kazachkova et al., 2018; Oscar et al., 2018). Understanding the molecular mechanisms by which extremophytes adapt to their stressful environments could aid in identifying targets for molecular breeding efforts to improve crop stress tolerance, as well as facilitate the development of extremophyte-based agriculture (Bressan et al., 2013; Shabala, 2013; Cheeseman et al., 2015; Ventura et al., 2015). While tolerance to salt stress has been extensively investigated in halophytes that are adapted to highly saline environments (Flowers et al., 2015; Kazachkova et al., 2018; Wang et al., 2021a), our understanding of plant molecular adaptations to stresses characteristic of desert habitats is still in its infancy (e.g. Granot et al., 2009; Yates et al., 2014; Oh et al., 2015; Obaid et al., 2016; Eshel et al., 2021; Wan et al., 2021). Yet, desert species could represent a treasure trove of molecular determinants that confer tolerance to the multiple stresses such as drought, salinity, low soil nutrient levels, and heat stress. With global temperatures projected to continue rising (IPCC, 2021), plant adaptation to heat stress is of particular importance due to its negative effects on plant physiology, particularly at the reproductive stage, leading to severe consequences upon yield (Mittler and Blumwald, 2010; Lesk et al., 2016; Chaturvedi et al., 2021; Wang et al., 2021b). Indeed, tolerance to heat stress in wheat at the reproductive stage has been identified as a key trait to increase yield potential under projected climate change (Stratonovitch and Semenov, 2015).

To gain insight into genetic adaptations that facilitate an extremophyte lifestyle, comparative physiological and molecular analyses of stress-sensitive *Arabidopsis thaliana* and its Brassicaceae extremophyte relatives have proven to be a powerful approach (Kraemer, 2010; Koenig and Weigel, 2015; Kazachkova et al., 2018). Indeed, these extremophyte relatives are becoming premier models for understanding plant adaptation to extreme environments with the development of a number of genetic resources including chromosome-level genome assemblies, natural accession collections, transformation protocols, and web resources (http://extremeplants.org/) (Zhu et al., 2015; Kazachkova et al., 2018; Wang et al., 2019). Yet, an extremophyte Brassicaceae model that represents desert species has not hitherto been developed. Such a model could leverage the functional genomics knowledge that exists for *A. thaliana* thereby facilitating comparative analyses to understand plant adaptations to the extreme desert

environment. We have therefore been studying the *A. thaliana* relative, *Anastatica hierochuntica* L., also

known as the 'True Rose of Jericho', a Saharo-Arabian desert species (Fig. 1A) which also occupies the

uppermost, driest zones of wadies or runnels of the Israeli Negev desert (Friedman and Stein, 1980;

Friedman et al., 1981; Fig. 1B). This arid region has temperatures varying between -3.6 and 46 °C, an

annual rainfall between 25 and 200 mm, and soil nitrate levels ranging from 0.4 to 4 mM (Gutterman,

2002; Ward, 2009; Eshel et al., 2017). We have demonstrated that *A. hierochuntica* is highly tolerant to

heat, low soil nitrogen, and moderately tolerant to salt stress (Eshel et al., 2017). Moreover, the plant

exhibits salt-resilient photochemistry and displays constitutively higher levels of metabolites that have a

role in scavenging reactive oxygen species, than *A. thaliana* (Eppel et al., 2014; Eshel et al., 2017).

In the current study, we assembled an *A. hierochuntica* reference transcriptome and used this

resource for two approaches to identify adaptations to an arid environment in a desert annual

extremophyte. In the first approach, comparative analysis of heat-response transcriptomes revealed an *A.*

hierochuntica transcriptome that is more reactive to heat than that of *A. thaliana*. In the second approach,

positive selection analysis identified genes that could contribute to adaptation to extreme conditions in

general, and those that could facilitate an opportunistic desert lifestyle.

Materials and Methods

119   For all analyses, detailed methods are provided in Supporting Information Methods S1.

Plant material and growth conditions

Plants for *de novo* reference transcriptome sequencing were grown on MS (Murashige and Skoog,

1962) plates for 5 d in the growth room (16 h light [150 µmol photons m^{-2} s^{-1}]/8 h dark; 22 °C). For

Illumina sequencing, plate-grown seedlings were directly used. For Roche 454 sequencing, plate-grown

seedlings were transferred to pots containing *A. thaliana* soil growth medium (Weizmann Institute of

Science), and kept in the growth room until plants developed four true fully-expanded leaves. These

plants were then treated as follows: (a) Control (field-capacity, 22 °C); (b) Drought stress (25% field

capacity for 1 week); (c) Salt shock (200 mM NaCl in the fertilizer solution), harvested after 1, 3 and 6

h; (d) Heat shock (45 °C), harvested after 0.5, 1, and 2 h. Roots, shoots and flowers (where available)

from these soil-grown plants, were harvested separately and snap-frozen in liquid nitrogen. In addition,

mature seeds, were imbibed in H_2O for 8.5 h.

For RNA-seq heat stress experiments, *A. thaliana* and *A. hierochuntica* were grown on plates

until cotyledons were fully expanded before transfer to pots containing *Arabidopsis* nitrogen-less soil

(Weizmann Institute of Science) and irrigation to field capacity with a custom-made fertilizer solution.

After 6 d in the growth room, uniform plants were transferred to two growth chambers (KBWF 720,

BINDER GmbH, Tuttlingen, Germany) (16 h light/8 h dark; 23 °C; 60% relative humidity; sunrise, 0.5

h at 100 μmol photons m^{-2} s^{-1}; daytime, 250 μmol m^{-2} s^{-1}; sunset, 0.5 h at 150 μmol photons m^{-2} s^{-1}). Ten

days after transfer to soil, heat treatment was initiated in one chamber comprising 3 d at 40/25 °C,

day/night temperatures, followed by 2 d recovery at control conditions (Fig. 2A). The other chamber was

kept as the control (23 °C). For each condition, three biological replicates comprising 6 pooled plants per

replicate (54 samples in total) were used for downstream analyses.

Reference transcriptome and RNA-seq

For sequencing the *A. hierochuntica* reference transcriptome, equal amounts of total RNA from

all samples (i.e. control, various stresses, tissues, time points, see "Plant material and growth conditions")

were pooled and sent to the GenePool genomics facility at the University of Edinburgh, UK for 454

sequencing of a normalized cDNA library. In addition, total RNA from plate-grown seedlings was sent

to the Glasgow Polyomics Facility at the University of Glasgow, UK for Illumina sequencing. The

reference transcriptome was assembled using a hybrid assembly approach that utilized both Illumina and

454 reads and was annotated based on public databases (Methods S1).

For RNA-seq, control and heat-treated total RNA samples (Fig. 2A) were delivered to the Roy J.

Carver Biotechnology Center, University of Illinois, Urbana-Champaign, USA. Libraries were prepared

with the Illumina TruSeq Stranded mRNA Sample Prep Kit (Illumina) and 100 bp HiSeq2500 Illumina

single-end reads were uniquely mapped to *A. thaliana* TAIR 10 or the *A. hierochuntica* reference

transcriptome using the Trinity align_and_estimate_abundance.pl script, which applies the RSEM

program (Grabherr et al., 2011; Li and Dewey, 2011) with the Bowtie aligner.

Differentially expressed genes (DEGs) were identified using DESeq2 (Love et al., 2014; Methods

S1). For raw read counts and DEGs identified in each species and for various functional groups, see

Dataset S2..

Ortholog pairs (17,962; Methods S1) were assigned to the five idealized modes of expression in

response to heat stress (Fig. 3A), using Weighted Gene Co-expression Network Analysis (WGCNA;

Langfelder and Horvath, 2008) to cluster normalized and quantified expression data into modules

containing genes with similar expression profiles (Fig. S3; Dataset S3; Methods S1). For direct

comparison of absolute orthologous transcript levels, TPM values of minimum or maximum expression

were analyzed for statistically significant difference ($p \leq 0.05$) with a Student *t*-test.

Phylogenomics and positive selection analysis

To identify orthologous genes among species and generate a maximum likelihood phylogenomic tree, we used coding sequences of the *A. hierochuntica* reference transcriptome and 16 sequenced Brassicaceae species (Dataset S12) with the Agalma phylogenomics pipeline (Dunn et al., 2013).

To detect positive selection, we used the Branch-Site model in the PAML v4.8, CODEML program (Yang, 1997; Yang, 2007). Ortholog groups with sequence representation in at least two of the five extremophytes, were selected to ensure sufficient statistical power (Anisimova et al., 2001). The tested branch(s) were labeled (foreground), and the log likelihood of two models (M1a and M2a), were calculated for each ortholog group. A Likelihood Ratio Test was performed (with X^2 distribution), to identify genes with log likelihood values significantly different between the two models, indicative of deviation from neutral selection. Ortholog groups with a portion of sites in the foreground branches, that had an estimated dN/dS ratio greater than 1, were considered under positive selection. To account for multiplicity, a Benjamini–Yekutieli false discovery rate (FDR) correction (Benjamini and Yekutieli, 2001) was applied using the "qvalue" R package, with a *q*-value < 0.05 cutoff for a gene to be considered as positively selected. Sites under positive selection were identified using the empirical Bayes approach with a posterior probability *p* > 0.95.

For each analysis, different branches on the tree were tested (labeled as foreground) compared with all other branches (background): (i) labeling the external branches of all five extremophyte species as the foreground (4,723 ortholog groups); (ii) labeling the *A. hierochuntica* external branch as the foreground (3,093 ortholog groups); (iii) labeling the *E. salsugineum* external branch as the foreground (4,457 ortholog groups); (iv) labeling the *S. parvula* external branch as the foreground (4,369 ortholog groups); and (v) labeling the *A. thaliana* external branch as the foreground (5,513 ortholog groups). *A. thaliana* was considered as a control/comparator species sensitive to abiotic stresses (Kazachkova et al., 2018). The Venn diagram comparing positive selected genes (Fig. 6B) was generated using an online tool: http://bioinformatics.psb.ugent.be/webtools/Venn/).

Significant positively selected genes as well as all DEGs were tested for enriched GO terms (Fisher's exact test, with a *q*-value < 0.05 cutoff) using AgriGO (Du et al., 2010), where the *A. thaliana* genome served as background. GOMCL (Wang et al., 2020) was used to summarize non-redundant functional groups).

Results

De novo assembly and annotation of the *Anastatica hierochuntica* reference transcriptome

To generate a high-quality *A. hierochuntica* reference transcriptome that maximizes coverage of genes, we sequenced and assembled transcripts using RNA pooled from multiple plant organs (root, shoot, flower, seeds), at different developmental stages (early seedling stage, and mature plants before and after flower initiation), and under control, heat, drought and salinity stress conditions (Fig. S1; Methods S1). We identified 30,670 putative protein-coding genes out of the high-confidence 36,871 assembled transcripts (Fig. 1C; Methods S1), and the distribution of transcript lengths was similar to that of *A. thaliana* cDNAs (Fig. 1C). High completeness of the reference transcriptome was evidenced by the detection of 93.6% BUSCOs (Fig. 1D; Simão et al., 2015), comparable to other *de novo* assembled Brassicaceae transcriptomes (Lopez et al., 2017). Furthermore, we obtained 88% sequenced reads that mapped back to the assembled reference transcriptome. These data indicate a high-quality reference transcriptome appropriate for downstream analyses. We annotated the reference transcriptome using sequence similarity to protein databases including NCBI, InterPro, and KEGG (Dataset S1) resulting in annotation of 96% of our assembled transcripts to a known sequence in the reference databases (Methods S1).

A. thaliana and *A. hierochuntica* global transcriptomes exhibit similar adjustment in response to heat stress

Certain stress-associated metabolites are constitutively maintained at high levels in *A. hierochuntica* even under stress-neutral control conditions suggesting that *A. hierochuntica* exists in a "stress-ready" state (Eshel et al., 2017). To test this hypothesis at the transcriptome level, we performed a comparative analysis of the *A. thaliana* and *A. hierochuntica* transcriptome response to heat stress in young plants at similar developmental stages. To simulate a realistic scenario, temperature data near *A. hierochuntica* populations during their growing season were obtained from the Israel Meteorological Service and used as a guide for diurnal heat treatments (Fig. S2). Furthermore, heat treatments were chosen that *A. thaliana* plants could also survive (Hayes et al., 2021). Thus, plants were exposed to either control (23 °C) or three consecutive daily heat waves covering the early heat response and acquired heat tolerance phases (Lindquist, 1986; Hong and Vierling, 2000), with day/night temperatures of 40 °C/25 °C followed by 2 days recovery at 23 °C (Fig. 2A). Plants were harvested either in the morning (1.5 hours after the onset of the light/heat period, red circles in Fig. 2A) or in the afternoon (7 hours after the onset

of the light/heat period, blues circles in Fig. 2A). Plants were well-watered throughout the entire

experiment to avoid any dehydration effects that could arise due to the heat treatment.

Heat stress had no significant effect on *A. hierochuntica* leaf area in contrast to *A. thaliana* where

growth in leaf area was significantly retarded by heat stress although it had almost recovered to control

levels 2 days after the end of the heat treatment (Fig. 2B). *A. thaliana* shoot fresh weight was also

significantly reduced by heat stress but did not recuperate after 2 days recovery under control conditions

while *A. hierochuntica* fresh weight was not affected by heat stress (Fig. 2C). These results illustrate that

A. hierochuntica is highly tolerant to heat stress and confirm our previous *in vitro* experiments (Eshel et

al., 2017).

Transcriptomes of both species under elevated temperature were clearly distinct from those in

control conditions (Fig. 2D). The control and heat-stressed samples harvested in the morning were

positioned separately from the samples harvested in the afternoon, which could be due to differences in

early vs. late heat-mediated gene expression or/and diurnal changes in gene expression. Transcriptomes

of plants recovering from heat stress clustered near control noon samples suggesting that, overall, the

transcriptomes return to pre-stress conditions. Because both species underwent transcriptional

adjustment in response to heat stress, we examined the median expression level across the whole

transcriptome for each condition. Compared to their respective controls (CM, CN, CR2N), the median

transcript abundance (and total abundance as depicted by the distribution) of both species decreased

under heat stress in the morning samples, increased in response to heat treatments in the noon samples,

and decreased during recovery (Fig. 2E). Furthermore, the percentage of differentially expressed genes

(out of the total number of protein-coding genes) was similar for both species under all heat conditions

(Fig. 2F; Dataset S2). These data show that the *A. thaliana* and *A. hierochuntica* global transcriptomes

adjust to heat stress with a similar magnitude and does not support a globally "stress-ready" *A.*

hierochuntica transcriptome.

254   The *A. hierochuntica* heat-response transcriptome does not exist in a "stress-ready" state

To test our contention that *A. hierochuntica* transcriptome is not "stress-ready", we used

Weighted Gene Co-expression Network Analysis (WGCNA) to identify five types of idealized

transcriptional response modes among the expression patterns of 17,962 orthologous pairs from each

species (Wang et al., 2021a; Fig. S3; Dataset S3): (a) "Stress-ready" where transcript level under control

conditions in one species is equal to the ortholog transcript level under heat in the other species; (b)

"Shared response" where expression of both orthologs exhibit a similar response to heat (i.e. both

upregulated or downregulated by heat); (c) "Unique response" where expression of an ortholog exhibits a heat response specifically in one species but not in the other; (d) "Opposite response" where expression of the ortholog in one species shows the opposite response in the other species; (e) "No response" where expression of both orthologs does not respond to heat. Of the orthologs categorized within the five response modes, only 4.4% of the orthologs belonged to the "No response" mode (Fig. 3A). The vast majority (82%) of orthologs displayed a shared response mode while about 5% exhibited a unique response and 2.1% showed an opposite response. Importantly, while 535 genes did exhibit a "stress-ready" mode in *A. hierochuntica*, we also detected 221 *A. thaliana* genes displaying a "stress-ready" mode. Similar results were obtained when we examined only orthologs associated with GO terms for abiotic stress responses (Fig. 3A; Methods S1). Overall, our data support the notion that *A. hierochuntica* does not possess a "stress-ready" transcriptome.

A. hierochuntica heat-regulated genes display a higher fold-change and/or lower basal expression compared to *A. thaliana*

Under control conditions, we observed that median basal expression of the *A. hierochuntica* transcriptome was significantly lower than in *A. thaliana* (Fig. 3B). Moreover, differentially expressed genes (DEGs) from the extremophyte displayed a greater heat-mediated fold-change in expression than *A. thaliana* DEGs (Figs. 3C, 3D) suggesting a more reactive heat-response transcriptome. To support these findings, we compared orthologous expression of specific functional groups that exhibited either a shared or unique response mode to heat stress (Dataset S2). Orthologs associated with GO-terms for abiotic stress whose expression displayed shared upregulation by heat exhibited an average lower basal expression in *A. hierochuntica* compared to *A. thaliana* and no significant difference in average % induction of expression (Fig. 4A). Abiotic stress-associated orthologs showing shared heat-mediated downregulated expression displayed both a lower basal and higher % reduction in expression in *A. hierochuntica* compared to *A. thaliana* (Fig. 4B). Similarly, both heat-mediated upregulated and downregulated abiotic stress-associated, unique-expressed orthologs showed lower basal and higher % induction/reduction in expression in the extremophyte (Figs. 4A, 4B).

Plants actively and early on reduce their growth in response to stress independent of photosynthesis, and this is apparent in a reduction in both cell size and cell elongation that can be linked to downregulation of cell cycle-associated genes (Aguirrezabal et al., 2006; Skirycz et al., 2010; Skirycz et al., 2011; Kazachkova et al., 2013). Subsequently, expression of genes involved in photosynthesis is downregulated under stress (Rizhsky et al., 2002; Zhang et al., 2018a; Huang et al., 2019). We observed

that the majority of shared- and unique-expressed orthologs associated with photosynthesis or the cell

cycle were downregulated by heat stress in both species (Dataset 2). However, for both shared- and

unique-expressed orthologs associated with photosynthesis, *A. hierochuntica* exhibited a similar basal,

but greater % reduction in expression than *A. thaliana* (Fig. 4C). Orthologs encoding proteins involved

in the cell cycle that possessed shared heat-mediated downregulated expression showed a lower basal

and higher % reduction in expression in the extremophyte while unique-expressed cell-cycle orthologs

exhibited lower basal expression in *A. hierochuntica* (Fig. 4D).

We next used WGCNA to cluster genes from all conditions into modules with similar expression

profiles and detected 22 *A. thaliana* and 21 *A. hierochuntica* co-expression modules (Fig. S3). In both

species, two modules clearly covered early heat-induced genes (1.5 h [morning] and 7 h [afternoon] after

onset of heat stress) (Fig. 5A; Datasets S4-S8). The morning heat-response modules of both species were

enriched in GO biological terms such as "response to heat", "response to high light intensity" and

"response to reactive oxygen species", (Dataset S9; Methods S1) while the *A. thaliana* and *A.*

hierochuntica afternoon heat-response modules were not enriched in any GO-terms. Importantly, both

shared- and unique-expressed *A. hierochuntica* genes associated with GO terms for abiotic stress in the

early heat-response modules exhibited the same or lower basal expression, and higher heat-mediated %

induction of expression than their *A. thaliana* orthologs (Fig. 5B).

To provide functional support for a more reactive *A. hierochuntica* heat-response transcriptome,

we identified 10,653 *A. hierochuntica* genes that exhibited significantly higher heat-mediated % change

in expression than their *A. thaliana* orthologs (Dataset S10). These genes were enriched in biological

processes related to abiotic stress including oxidative, water, and salt stresses, response to radiation

(including genes involved in defense against U/V light), and the response to DNA damage (Fig. 5C,

Dataset S11). Additionally, the "protein folding" gene list contained heat shock protein-encoding genes.

To validate our gene expression comparisons, we showed (Fig. S4; Methods S1): (i) no significant

difference between the species in the proportion of the top 10 most highly expressed genes out of the

total transcripts sequenced across all treatments; (ii) similar comparative basal expression results as

observed with DEseq2, when we used a new between-species Scale-Based Normalization method (Zhou

et al., 2019); (iii) relative and quantitative QPCR analysis confirmation of the RNA-seq fold-change and

basal gene expression patterns of selected genes.

Furthermore, we examined the basal expression of 15 orthologous housekeeping genes from both

species and found that the average ratio of basal expression of *A. thaliana* to *A. hierochuntica*

housekeeping genes was 1.0 ± 0.34 (Fig. S5). Thus, the average lower basal gene expression observed in

A. hierochuntica compared to *A. thaliana* was not due to lower metabolic activity in the extremophyte.

Brassicaceae extremophytes possess positively selected genes associated with surviving harsh environments

As a second approach to identifying adaptations to an extremophyte lifestyle, in general, and to

desert conditions in particular, we pinpointed positively selected genes that might be indicative of

adaptive evolution of stress tolerance. We first used phylogenomics to infer evolutionary relationships

between 16 Brassicaceae species including *A. hierochuntica* and representing all major lineages in this

family (Dataset S12). *Tarenaya hassleriana* (Cleomaceae) was used as an outgroup. This led to a

selection of 13,806 ortholog groups found in 17 taxa. The phylogenomic tree partitioned the species in

concordance with their previously assigned lineages (I, II, and III), where *Aethionema arabicum* is

considered to belong to a basal clade within the Brassicaceae (Fig. 6A; Franzke et al., 2011; Kiefer et al.,

2014). *A. hierochuntica* (*Anastaticeae*) was assigned to LIII (Franzke et al., 2011). It is important to note

that *A. hierochuntica* is the single representative species used for LIII due to this lineage being sparsely

represented in publicly available genomic databases unlike transcriptomes available for LI and LII

species. Thus, to the best of our knowledge, we provide the first substantial genetic resource that enables

exploration into adaptive traits that have evolved in a representative lineage III species.

The tree contains five extremophyte species (Fig. 6A, red asterisks): the halophytes *Eutrema*

salsugineum and *Schrenkiella parvula* (tolerant to high salinity and multiple other stresses; Kazachkova

et al., 2018; Wang et al., 2021a), *Thlaspi arvense* (freezing-tolerant; Sharma et al., 2007; Zhou et al.,

2007), *A. hierochuntica* (heat-, salt-, low N-tolerant; Eshel et al., 2017) and *Arabidopsis halleri* (heavy

metal hyperaccumulator, semi-alpine conditions; Hanikenne et al., 2008; Honjo and Kudoh, 2019).

Therefore, to identify genes under common positive selective pressure in the extremophytes, we used the

branch-site model (Yang, 1997; Yang, 2007) to test the external branches (foreground) of the five

extremophyte species against all the other branches (background). We then repeated this procedure to

test for positively selected genes in three specific extremophytes - the well-studied halophyte models, *E.*

salsugineum and *S. parvula*, and *A hierochuntica* - by labelling each species' external branch as the

foreground. Overall, we identified 194, 120, 131 and 99 genes with a positive selection signal in the "all

extremophyte species", *A. hierochuntica*, *E. salsugineum*, and *S. parvula* runs, respectively (Datasets

S13-S16). We also tested *A. thaliana*, as an abiotic stress-sensitive control, and identified 112 positively

selected genes (Dataset S17).

We next examined whether any common genes are under positive selection in the extremophytes
or between the extremophytes and stress-sensitive *A. thaliana*. While we could not detect a clear
convergence in the use of common positively selected orthologs in the extremophytes (Fig. 6B), the
functional attributes shared by those positively selected orthologs in each extremophyte exhibited
convergence (Fig. 6C; Fig. S6; Datasets S18 to S22). Notably, orthologs associated with the GO-term
"response to stress" [GO:0006950] were highly enriched in the extremophytes suggesting major selective
pressure for stress tolerance imposed by their extreme environments.

Positively selected genes from the "all extremophyte species", supported association with
adaptations to harsh environments. For instance, *AKS2*, *MYB52*, *WRKY75*, *ASF1B* and *PHR1/UVR2* that
have known functions in ABA responses, phosphate starvation, heat stress, and UV-B radiation stress
(Table 1 and refs. therein), were among the positively selected group of genes in the extremophytes.
Interestingly, *bZIP1* (salt/drought tolerance, and nitrogen signaling) and *APX6* (ROS-scavenging)
showed signatures of positive selection unique to *A. hierochuntica* (Table 1), which is highly tolerant to
low N and oxidative stresses, and moderately tolerant to salt stress (Eshel et al., 2017). Positively selected
genes unique to *S. parvula* included *CAX11/CCX5* and *RAB28* that are involved in high-affinity K^+
uptake and Na^+ transport, and lithium toxicity, respectively (Table 1; Borrell et al., 2002; Zhang et al.,
2011). The pinpointing of these two genes added validity to our positive selection analysis because the
native soils of *S. parvula* contain highly toxic levels of Li^+ and K^+ (Helvaci et al., 2004; Ozfidan-Konakci
et al., 2016), and this species displays extreme tolerance to both Li^+ and K^+ toxicity (Oh et al., 2014;
Pantha et al., 2021). In contrast to the extremophyte species, positively selected genes in *A. thaliana* were
related to biotic stress responses (Table 1). Of the positively selected genes in the "all extremophyte
species" or *A. hierochuntica*-specific sets, *AKS2*, *bZIP1* and *PHR1/UVR1* expression displayed clear and
significantly higher transcript levels in *A. hierochuntica* compared to *A. thaliana* whereas the expression
of *APX6* exhibited lower transcript levels in *A. hierochuntica* (Fig. 7A).

Exclusively in *A. hierochuntica,* we identified, *CYP71*, *FAS1*, *FBH2*, *SBI1/LCMT1*, and *VIP5* as
positively selected genes that are involved in photoperiodic flowering, regulation of meristems, and
control of morphology including shoot branching (Table 1). Furthermore, *AhFAS1* expression was highly
upregulated by heat stress while *AtFAS1* expression was downregulated (Fig. 7B). *AhSBI1/LCMT1*
expression was unaffected by heat stress whereas *AtSB1/LCMT1* expression was highly upregulated by
heat. Moreover, *AhSBI1/LCMT1* transcript levels were lower than *AtSBI1/LCMT1* over all time points.
Notably, genes involved in organ development and flowering time were more reactive to heat in *A.*
hierochuntica than in *A. thaliana* (Fig. 5C). Considering that *A. hierochuntica* ontogeny is very different

from *A. thaliana*, *E. salsugineum* and *S. parvula* – it exhibits a multi-branched sympodial shoot structure
supporting multiple axillary inflorescences that flower independent of day length (Fig. 1B; Gutterman,
1998; Eshel et al., 2017) - positive selection of these genes could indicate an important adaptation to the
desert environment.

Discussion

The *A. hierochuntica* transcriptome does not exist in a heat "stress-ready" state and is more
reactive to heat stress than *A. thaliana*

Our finding that *A. thaliana* and *A. hierochuntica* exhibit similar transcriptome adjustment in
response to heat stress and during recovery (Figs. 2E, 2F) distinguishes *A. hierochuntica* from other
extremophyte relatives. The extent of transcriptomic, proteomic and metabolic adjustment in response to
ionic stress in the halophytes *E. salsugineum* and *S. parvula*, is much lower than in *A. thaliana* (Taji et
al., 2004; Gong et al., 2005; Lugan et al., 2010; Pang et al., 2010; Oh et al., 2014; Vera-Estrella et al.,
2014; Wang et al., 2021a). This lower adjustment reflects their "stress-ready" state whereby transcript
and metabolite accumulation that is induced or repressed in *A. thaliana* in response to ionic stress, is
constitutively high or low, respectively, in the halophytes (Gong et al., 2005; Lugan et al., 2010;
Kazachkova et al., 2013; Oh et al., 2014; Wang et al., 2021a). A "stress-ready" transcriptome is
exemplified in *S. parvula* where basal expression of over 1,000 "stress-ready" orthologs matches the
post-boron stress expression levels observed in *A. thaliana* (Wang et al., 2021a). In contrast, the great
majority of *A. hierochuntica* and *A. thaliana* orthologs exhibit a shared response mode (Fig. 3A).
Furthermore, many stress-related *A. hierochuntica* genes show lower basal and/or higher fold-change
gene expression compared to *A. thaliana* (Figs. 3, 4, 5). Indeed, almost one third of *A. hierochuntica*
genes display a higher heat-mediated fold-change in expression compared to *A. thaliana* and are enriched
in abiotic stress-associated functions (Fig. 5; Datasets 10, 11). Taken together, our findings support a
paradigm whereby the *A. hierochuntica* transcriptome is more reactive to heat stress than *A. thaliana*.

One reason for the difference in the transcriptome responses of *A. hierochuntica* vs. its halophytic
relatives relates to the type of stress each species faces. *E. salsugineum* and *S. parvula* habitats possess
levels of ions such as Na^+ and BO_3^{3-} that are toxic to the majority of plant species and the two halophytes
are constantly exposed to ionic stress throughout their life cycle. This situation might have led to the
evolution of a stress-associated transcriptome that is continuously "switched-on". Conversely, *A.*
hierochuntica is generally exposed to heat stress later in its life cycle and on a diurnal basis thus favoring
a more reactive transcriptome.

A second reason is that *A. hierochuntica* thrives in an environment with seasonal temperatures

ranging from -3.6 to 46.8 °C (Eshel et al., 2017), with diurnal maximum variations exceeding 18 °C

(Israeli Meteorological Services). On the other hand, *E. salsugineum* can be found in locations such as

China's Shandong peninsula where temperatures range from -5 °C in the winter to 32 °C in the summer

and with diurnal temperature differences rarely exceeding 10 °C (Guedes et al., 2015). Furthermore,

diurnal temperature differences in the cold, spring-growing period of *E. salsugineum* are likely more

moderate than those experienced by *A. hierochuntica* during the warm Negev desert spring. Thus,

evolution of a flexible transcriptome that confers a strong reaction to extreme diurnal temperature

fluctuations could be advantageous for adaptation to a desert environment. Moreover, a transcriptome

with globally lower basal expression levels would require less energy to be expended in the low nutrient

desert environment.

Brassicaceae extremophytes possess common positively selected genes that are indicative of

adaptation to harsh environments

Extremophytes are present in all three Brassicaceae lineages (Fig. 6A; Franzke et al., 2011)

illustrating that adaptation to stressful habitats has occurred independently, multiple times within the

Brassicaceae and is indicative of convergent evolution (Kazachkova et al., 2018; Birkeland et al., 2020).

Consistent with this notion, we identified a set of 194 genes under positive selection across the five

extremophyte species that could commonly contribute to plant adaptation to extreme environments.

Other studies with extremophyte Brassicaceae have also detected signatures of positive selection in genes

that function in stress tolerance (Zhou et al., 2009; Jarvis et al., 2014; Birkeland et al., 2020). For instance,

positive selection of stress-associated genes was detected in three Arctic Brassicaceae species (Birkeland

et al., 2020). Similar to our findings (Figs. 6B, 6C, S6) there was little overlap of positively selected

genes between the Arctic extremophytes but considerable overlap in functional pathways. Taken

together, these data do not support adaptive molecular convergence but rather indicate evolution of

similar adaptations via distinct evolutionary pathways.

Among the genes under positive selection across the five extremophyte species in the current

study, we identified two genes encoding ABA-responsive transcription factors (TFs), AKS2 and MYB52

(Table 1), illustrating the importance of the ABA response networks in adaptive evolution of stress

tolerance (Xia et al., 2010; Fischer et al, 2011; Bondel et al., 2018). In particular, the bHLH TF, ABA-

RESPONSIVE KINASE SUBSTRATE (AKS) 2 activates transcription of K^+ channels in stomata guard

cells in an ABA-dependent manner thereby enhancing stomatal opening (Takahashi et al., 2013). It is

intriguing that a regulator of stomatal aperture has undergone positive selection across the extremophytes because alterations in stomatal aperture is a crucial early response to multiple abiotic stresses (Brugnoli and Lauteri, 1991; Chaves et al., 2009; Stepien and Johnson, 2009; Devireddy et al., 2020). Thus, positive selection of non-synonymous amino acid changes in the coding region of *AKS2*, (plus differences in heat-mediated regulation of *A. thaliana* and *A. hierochuntica AKS2* expression (Fig. 7A)) suggest that this gene may have been naturally selected for survival in extreme environments.

WRKY75 was also positively selected across the extremophytes (Table 1). In *A. thaliana*, this TF regulates the expression of several key phosphate starvation-induced genes (Devaiah et al., 2007). Extremophytes often exist on soils with low Pi availability (Thompson et al., 2006; Holzapfel, 2008; Guevara et al., 2012). For instance, the *E. salsugineum* Yukon ecotype grows in the low Pi soils of the Yukon region in Canada (Guevara et al., 2012) and is highly tolerant to Pi deficiency compared to *A. thaliana*. This tolerance is associated with higher basal expression of several Pi starvation genes including *WRKY75* (Velasco et al., 2016). Both *E. salsugineum* and *A. hierochuntica* exhibit significantly higher basal levels of Pi than *A. thaliana* (Gong et al., 2005; Kazachkova et al., 2013; Velasco et al., 2016; Eshel et al., 2017). Thus, positive selection of *WRKY75* across the five extremophyte plants, and differential expression of *A. thaliana* and *E. salsugineum WRKY75* suggests that selection for more efficient extraction of soil Pi is a common evolutionary adaptation to extreme environments.

Extremophytes are often exposed to UV-B radiation that can cause direct damage to DNA (Kimura and Sakaguchi, 2006). *PHOTOLYASE1/UV-RESISTANCE2* (*PHR1/UVR2*) and *ANTI-SILENCING FUNCTION 1b* (*ASF1B*) that are crucial for repairing UV-B-induced DNA damage were positively selected across the five extremophytes (Table 1; Ahmad et al., 1997; Landry et al., 1997; Jiang et al., 1997; Lario et al., 2013; Nie et al., 2014). *PHR1/UVR2* is also the major mechanism maintaining transgenerational genome stability in *A. thaliana* continuously exposed to UV-B (Willing et al., 2016) while *ASF1B* is also involved in the regulation of basal and acquired thermotolerance (Weng et al., 2014). Additionally, *PHR1/UVR2* displays higher basal expression in *A. hierochuntica* compared to *A. thaliana*, and while heat leads to downregulation of the gene in both species, expression is reduced to a lesser extent in the extremophyte. (Fig. 7A).

Collectively then, our data suggest common selective pressures in extremophyte plants that target key components in stomatal opening, nutrient acquisition, and UV-B-induced DNA repair. On the other hand, we found that *A. thaliana* positively selected genes were principally involved in defense against pathogens (Table 1). This supports the hypothesis that because *A. thaliana* evolved in temperate regions

where pathogen density is relatively high compared to extremophyte habitats, it encountered greater evolutionary pressures for adaptation to biotic stresses (Oh et al., 2014).

Positively selected *A. hierochuntica* genes suggest adaptive evolution for an opportunistic desert lifestyle

We pinpointed a number of positively selected genes specifically in *A. hierochuntica* indicating adaptation to the desert environment (Table 1; Fig. 7B). Intriguingly, several of these genes function in *A. thaliana* in the transition from vegetative to reproductive growth and meristem development: (i) *VERNALIZATION INDEPENDENCE VIP5* enhances transcription of the floral repressor *FLOWERING LOCUS C* (*FLC*) gene and other *MADS AFFECTING FLOWERING* (*MAF*) gene family members (Oh et al., 2004; Yu and Michaels, 2010; Crevillen and Dean, 2011; Lu et al., 2017) and *A. thaliana vip5* mutants exhibit early flowering; (ii) FLOWERING BHLH 2 (FBH2) activates transcription of the *CONSTANS* gene, a central regulator of photoperiodic flowering. *FBH2* overexpression causes photoperiod-independent early flowering (Ito et al., 2012); (iii) *FASCIATA1* (*FAS1*) appears to function in the organization of shoot and root apical meristems, and in cellular differentiation (Kaya et al., 2001; Exner et al., 2006). Mutations in *fas1* cause stem fasciation, abnormal leaf and flower morphology, and defects in the organization of apical meristems (Leyser and Furner, 1992; Kaya et al., 2001); (iv) *CYP71* plays a critical role in regulating meristem development, including the floral meristem (Li et al., 2007). Furthermore, CYP71 physically interacts with FAS1 thereby targeting FAS1 to the *KNAT1* locus (Li and Luan, 2011). KNAT1 is essential for maintenance of apical meristems (Hake et al., 2004). In addition, CYP71 interacts with LIKE HETEROCHROMATIN PROTEIN 1 (LHP1), which is involved in repressing expression of flowering time and floral identity genes (Gaudin et al., 2001; Kotake et al., 2003). Thus, *lhp1* mutations cause strong early flowering; (v) *SUPPRESSOR OF BRI1 (SBI1)/LEUCINE CARBOXYLMETHYLTRANSFERASE (LCMT1)* regulates components of the brassinosteroid signaling pathway (Di Rubbo et al., 2011; Wu et al., 2011) and the *sbi1/lcmt* mutant is early flowering in both long and short days consistent with the role of brassinosteroids in flowering (Li and He, 2010; Nolan et al., 2020).

The discovery of positively selected flowering and meristem development genes specifically in *A. hierochuntica* is consistent with its very different developmental program compared to many other Brassicaceae including the additional four extremophyte plants included in our analysis. *A. hierochuntica* does not display the distinctive transition from the vegetative rosette leaf stage to the reproductive bolting stage, which is accelerated in long-day conditions (Pouteau and Albertini, 2009; Song et al., 2013).

Instead, regardless of photoperiod, the shoot repeatedly bifurcates from the four true-leaf stage onwards,

developing an axillary inflorescence at each branch point thereby leading to a multi-branched shoot

morphology (Fig. 1B, panel (i); Eshel et al., 2017). Most interestingly, mutation in the *A. thaliana FAS1*

gene (whose *A. hierochuntica* ortholog is under positive selection) can induce stem bifurcation and

enlargement (Leyser and Furner, 1992). The shoot bifurcation, multi-branch, photoperiod-insensitive,

early flowering traits could maximize fitness in the unpredictable desert environment where plants need

to ensure development of seeds but might not survive until a critical day length induces flowering. This

idea is supported by our observations of *A. hierochuntica* populations in the Dead Sea valley of Israel,

where tiny dead plants that have still managed to produce a few seeds can be seen alongside much larger

plants presumably from a year with higher rainfall (Fig. 1B).

In conclusion, we have shown that *A. hierochuntica* possesses a more reactive heat-response

transcriptome, and stress-related genes that have undergone positive selection. Genes that could be

associated with its multi-branch, early flowering phenotype also exhibit signatures of positive selection.

Together, these evolutionary adaptations could allow survival in a hot desert environment with

unpredictable precipitation. Our study furthermore provides rich gene sets that will facilitate comparative

and functional genomics studies to reveal additional molecular mechanisms for plant tolerance to heat

stress in a desert habitat.

Acknowledgements

We would like to dedicate this paper to Dr. Dirk Hincha and Prof. Hillel Fromm, who passed

away in 2020. Great plant scientists, colleagues and friends. They will be sorely missed. We thank Ruth

Shaked and Beery Yaakov for their dedicated technical support. We are also grateful to Patrick

Barko from the University of Illinois Urbana-Champaign for excellent aid with WGCNA. Our

appreciation to Alvaro Hernandez and all the team at the Roy J. Carver Biotechnology Center, University

of Illinois Urbana-Champaign for superb sequencing services. We are grateful to Julie Galbraith

(Glasgow Polyomics) for preparing libraries and carrying out the Illumina RNA-sequencing for the

reference transcriptome. This work was supported by the Goldinger Trust Jewish Fund for the Future,

the Koshland Foundation for Support of Interdisciplinary Research In Combatting Desertification, and

the I-CORE Program of the Planning and Budgeting Committee to S.B., National Science Foundation

award MCB 1616827, Next-Generation BioGreen21 Program of Republic of Korea (PJ01317301).to

M.D., NSF-BSF-IOS-EDGE 1923589/2019610 to M.D. and S.B., and the Biotechnology and Biological

Sciences Research Council grant (BBSRC; BB/R019894/1) to AA and PH. G.E. was supported by an

Israel President Fellowship for Excellence and Scientific Innovation award, N.D was supported by a
Ben-Gurion University, Kreitman School for Advanced Research Studies High-tech, Bio-tech and
Chemo-tech award, and G.W. was supported by an Economic Development Assistantship award from
Louisiana State University, The authors also acknowledge the LSU High Performance Computing
Services, and the BGU Bioinformatics Core Facility for providing computational resources needed for
data analyses.

Author contributions

S.B. conceptualized and supervised the overall project; GE and ND performed the main analyses;
GW contributed to transcriptome response assessment and performed the GOMCL analysis; GE, D-HO,
MG, MD, and VC-C contributed to assembly of the *A. hierochuntica* reference transcriptome. MD
assisted with, and GE performed the positive selection analysis; YK and MAO contributed to plant
growth, preparation of samples and transcriptome response validation; AA and PH designed, supervised
and analysed the Illumina RNA-sequencing experiment for the reference transcriptome at Glasgow
Polyomics; AM-C contributed to the WGCNA; GE, ND, GW, SB and MD contributed to data
interpretation, GE, ND, and SB wrote the article. MD and AA critically revised and approved the final
manuscript.

Data availability

566   Reference transcriptome and RNA-seq reads as well as the fully assembled transcriptome are
567   openly available via the NCBI SRA and TSA databases under BioProject PRJNA731383. Other main
568   data that supports the findings of this study are available in the main text and Supporting Information of
569   this article.

References

Aguirrezabal L, Bouchier-Combaud S, Radziejwoski A, Dauzat M, Cookson SJ, Granier C (2006)
Plasticity to soil water deficit in *Arabidopsis thaliana*: dissection of leaf development into
underlying growth dynamic and cellular variables reveals invisible phenotypes. Plant Cell Environ
29: 2216-2227
Ahmad M, Jarillo JA, Klimczak LJ, Landry LG, Peng T, Last RL, Cashmore AR (1997) An enzyme
similar to animal type II photolyases mediates photoreactivation in *Arabidopsis*. Plant Cell 9: 199-
207

Amtmann A (2009) Learning from evolution: *Thellungiella* generates new knowledge on essential and critical components of abiotic stress tolerance in plants. Mol Plant 2: 3-12

Anisimova M, Bielawski JP, Yang Z (2001) Accuracy and power of the likelihood ratio test in detecting adaptive molecular evolution. Mol Biol Evol 18: 1585-1592

Bao F, Azhakanandam S Franks RG (2010) SEUSS and SEUSS-LIKE transcriptional adaptors regulate floral and embryonic development in *Arabidopsis*. Plant Physiol 152; 821-836

Benjamini Y, Yekutieli D (2001) The control of the false discovery rate in multiple testing under depencency. Ann of Stat 29: 1165-1188

Birkeland S, Gustafsson ALS, Brysting AK, Brochmann C, Nowak MD (2020) Multiple genetic trajectories to extreme abiotic stress adaptation in Arctic Brassicaceae. Mol Biol Evol 37: 2052-2068

Bondel KB, Nosenko T, Stephan W (2018) Signatures of natural selection in abiotic stress-responsive genes of *Solanum chilense*. R. Soc open sci 5: 171198

Borrell A, Cutanda MC, Lumbreras V, Pujal J, Goday A, Culiáñez-Macià FA, Pagès M (2002) *Arabidopsis thaliana* Atrab28: A nuclear targeted protein related to germination and toxic cation tolerance. Plant Mol Biol 50: 249-259

Bressan R, Park H, Orsini F, Oh DH, Dassanayake M, Inan G, Yun J-Y, Bohnert H.J, Maggio A (2013) Biotechnology for mechanisms that counteract salt stress in extremophile species: a genome-based view. Plant Biotechnol Rep. 7: 27-37

Brugnoli E, Lauteri M (1991) Effects of salinity on stomatal conductance, photosynthetic capacity, and carbon isotope discrimination of salt-tolerant (*Gossypium hirsutum* L.) and salt-sensitive (*Phaseolus vulgaris* L.) C_3 Non-Halophytes. Plant Physiol 95: 628-635

Cassan-Wang H, Goué N, Saidi MN, Legay S, Sivadon P, Goffner D, Grima-Pettenati J (2013) Identification of novel transcription factors regulating secondary cell wall formation in *Arabidopsis*. Front Plant Sci 4: 189

Chaturvedi P, Wiese AJ, Ghatak A, Drabkova LZ, Weckwerth W, Honys D (2021) Heat stress response mechanisms in pollen development. New Phytol 231: 571-585

Chaves MM, Flexas J, Pinheiro C (2009) Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. Ann Bot 103: 551-560

Cheeseman JM (2015) The evolution of halophytes, glycophytes and crops, and its implications for food security under saline conditions. New Phytol 206: 557-570

Chen X, Goodwin SM, Liu X, Chen X, Bressan RA, Jenks MA (2005) Mutation of the
RESURRECTION1 locus of *Arabidopsis* reveals an association of cuticular wax with embryo
development. Plant Physiol 139: 909-919

Chen C, Letnik I, Hacham Y, Dobrev P, Ben-Daniel B-H, Vanková R, Amir R, Miller G (2014)
ASCORBATE PEROXIDASE6 protects *Arabidopsis* desiccating and germinating seeds from
stress and mediates cross talk between reactive oxygen species, abscisic acid, and auxin. Plant
Physiol 166: 370-383

Chen H, Ruan J, Chu P, Fu W, Liang Z, Li Y, Tong J, Xiao L, Liu J, Li C (2020) AtPER1 enhances
primary seed dormancy and reduces seed germination by suppressing the ABA catabolism and GA
biosynthesis in *Arabidopsis* seeds. Plant J 101: 310-323

Crevillen P, Dean C (2011) Regulation of the floral repressor gene FLC: the complexity of transcription
in a chromatin context. Curr Opin Plant Biol 14: 38-44

Dassanayake M, Haas JS, Bohnert HJ, Cheeseman JM (2010) Comparative transcriptomics for mangrove
species: an expanding resource. Funct Integr Genomics 10: 523-532

Devaiah BN, Karthikeyan AS, Raghothama KG (2007) WRKY75 transcription factor is a modulator of
phosphate acquisition and root development in *Arabidopsis*. Plant Physiol 143: 1789-1801

Devireddy AR, Arbogast J, Mittler R (2020) Coordinated and rapid whole plant systemic stomatal
response. New Phytol 225: 21-25

Di Rubbo S, Irani NG, Russinova E (2011) PP2A Phosphatases: The "on-off" regulatory switches of
brassinosteroid signaling. Sci Signal 4: pe25

Du Z, Zhou X, Ling Y, Zhang Z, Su Z (2010) agriGO: A GO analysis toolkit for the agricultural
community. Nucleic Acids Res 38: 64-70

Dunn CW, Howison M, Zapata F (2013) Agalma: an automated phylogenomics workflow. BMC
Bioinformatics 14: 330

Eppel A, Shaked R, Eshel G, Barak S, Rachmilevitch S (2014) Low induction of non-photochemical
quenching and high photochemical efficiency in the annual desert plant *Anastatica hierochuntica*.
Physiol Plant 151: 544-558

Eshel G, Shaked R, Kazachkova Y, Khan A, Eppel A, Cisneros A, Acuna T, Gutterman Y, Tel-Zur N,
Rachmilevitch S et al. (2017) *Anastatica hierochuntica*, an *Arabidopsis* desert relative, is tolerant
to multiple abiotic stresses and exhibits species-specific and common stress tolerance strategies
with its halophytic relative, *Eutrema* (*Thellungiella*) *salsugineum*. Front Plant Sci 7: 1992

Eshel G, Araus V, Undurraga S, Soto DC, Moraga C, Montecinos A, Moyano T, Maldonado J, Dıaz FP, Varala K (2021) Plant ecological genomics at the limits of life in the Atacama Desert. Proc Natl Acad Sci USA 118: e2101177118

Exner V, Taranto P, Schonrock N, Gruissem W, Hennig L (2006) Chromatin assembly factor CAF-1 is required for cellular differentiation during plant development. Development 133: 4163-4172

Fan Y, van den Dool H (2008) A global monthly land surface air temperature analysis for 1948-present. J Geophys Res 113: D01103

Fan L, Wang G, Hu W, Pantha P, Tran K-N, Zhang H, An L, Dassanayake M, Qiu Q-S (2018) Transcriptomic view of survival during early seedling growth of the extremophyte *Haloxylon ammodendron*. Plant Physiol Biochem 132: 475-489

Farrant JM, Cooper K, Hilgart A, Abdalla KO, Bentley J, Thomson JA, Dace HJW, Peton N, Mundree SG, Rafudeen MS (2015) A molecular physiological review of vegetative desiccation tolerance in the resurrection plant *Xerophyta viscosa* (Baker). Planta 242: 407-426

Fischer I, Camus-Kulandaivelu L, Allal F, Stephan W (2011) Adaptation to drought in two wild tomato species: the evolution of the *Asr* gene family. New Phytol 190: 1032-1044

Flowers TJ, Munns R, Colmer TD (2015) Sodium chloride toxicity and the cellular basis of salt tolerance in halophytes. Ann Bot 115: 419-431

Franzke A, Lysak MA, Al-Shehbaz IA, Koch MA, Mummenhoff K (2011) Cabbage family affairs: The evolutionary history of Brassicaceae. Trends Plant Sci 16: 108-116

Friedman J, Stein Z (1980) The influence of seed dispersal mechanisms on the distribution and survival of *Anastatica hierochuntica* (Cruciferae) in the Negev Desert. J Ecol 68: 43-50

Friedman J, Stein Z, Rushkin E (1981) Drought tolerance of germinating seeds and young seedlings of *Anastatica hierochuntica* L. Oecologia 51: 400-403

Gaudin V, Libault M, Pouteau S, Juul T, Zhao G, Lefebvre D, Grandjean O (2001) Mutations in LIKE HETEROCHROMATIN PROTEIN 1 affect flowering time and plant architecture in *Arabidopsis*. Development 128: 4847-4858

Gong Q, Li P, Ma S, Indu Rupassara S, Bohnert HJ (2005) Salinity stress adaptation competence in the extremophile *Thellungiella halophila* in comparison with its relative *Arabidopsis thaliana*. Plant J 44: 826-839

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q et al (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol 29: 644-52

Granot G, Sikron-Persi N, Gaspan O, Florentin A, Talwara S, Paul LJ, Morgenstern Y, Granot Y, Grafi
G (2009) Histone modifications associated with drought tolerance in the desert plant *Zygophyllum*
dumosum Boiss. Planta 231: 27-34.

Guedes JDA, Jin G, Bocinsky RK (2015) The impact of climate on the spread of rice to north-eastern
China: A new look at the data from Shandong Province. PLoS One 10: 1-19

Guevara DR, Champigny MJ, Tattersall A, Dedrick J, Wong CE, Li Y, Labbe A, Ping CL, Wang Y,
Nuin P, et al. (2012) Transcriptomic and metabolomics analysis of Yukon *Thellungiella* plants
grown in cabinets and their natural habitat show phenotypic plasticity. BMC Plant Biol 12: 175

Guo P, Li Z, Huang P, Li B, Fang S, Chu J, Guo H (2017) A tripartite amplification loop involving the
transcription factor WRKY75, salicylic acid, and reactive oxygen species accelerates leaf
senescence. Plant Cell 29: 2854-2870

Gutterman Y. (1998) Ecophysiological genotypic and phenotypic strategies affecting seed ''readiness to
germinate'' in plants occurring in deserts. In: Taylor, AG., Huang, X.-L. (Eds.), Progress in Seed
Research: Proceedings of Second International Conference on Seed Science and Technology.
Communication services of the New York State Agricultural Experiment Station, Geneva, NY, pp.
348

Gutterman Y (2002) Survival Strategies of Annual Desert Plants (Adaptations of Desert Organisms).
Heidelberg: Springer

Hake S, Smith HM, Holtan H, Magnani E, Mele G, Ramirez J (2004) The role of *knox* genes in plant
development. Annu Rev Cell Dev Biol 20: 125-151

Hanikenne, M, Talke IN, Haydon M., Lanz C, Nolte A, Motte P et al. (2008) Evolution of metal
hyperaccumulation required cis-regulatory changes and triplication of HMA4. Nature 453: 391-
396

Hayes S, Schachtschabel J, Mishkind M, Munnik T, Arisz SA (2021) Hot topic: Thermosensing in plants.
Plant Cell Environ 44: 2018-2033

Helvaci C, Mordogan H, Çolak M, Gündogan I. (2004) Presence and distribution of lithium in borate
deposits and some recent lake waters of West-Central Turkey. Int Geol Rev 46: 177-190.

Hisanaga T, Ferjani A, Horiguchi G, Ishikawa N, Fujikura U, Kubo M, Demura T, Fukuda H, Ishida T,
et al. (2013) The ATM-dependent DNA damage response acts as an upstream trigger for
compensation in the *fas1* mutation during *Arabidopsis* leaf development. Plant Physiol 162: 831-
841

Holzapfel, C. (2008) Deserts. In S.E. Jørgensen & B.D. Fath (eds.), Vol.2 of Encyclopedia of Ecology,
pp. 879-898. Oxford: Elsevier.

Hong SW, Vierling E (2000) Mutants of *Arabidopsis thaliana* defective in the acquisition of tolerance
to high temperature stress. Proc Natl Acad Sci USA 97: 4392-4397

Honjo MN, Kudoh H (2019) *Arabidopsis halleri*: a perennial model system for studying population
differentiation and local adaptation. AoB Plants 11: plz076

Hu Y, Wu Q, Sprague SA, Park J, Oh M, Rajashekar CB, Koiwa H, Nakata PA, Cheng N, Hirschi KD,
et al. (2015) Tomato expressing *Arabidopsis* glutaredoxin gene *AtGRXS17* confers tolerance to
chilling stress via modulating cold responsive components. Hortic Res 2: 15051

Huang J, Zhao X, Chory J (2019) The *Arabidopsis* transcriptome responds specifically and dynamically
to high light stress. Cell Rep 29: 4186-4199

IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte,
V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I.
Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O.
Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press.

Ito S, Song YH, Josephson-Day AR, Miller RJ, Breton G, Richard G. Olmstead RG, Imaizumi T (2012)
FLOWERING BHLH transcriptional activators control expression of the CONSTANS in
Arabidopsis. Proc Natl Acad Sci USA 109: 3583-3587

Jarvis DE, Ryu C-H, Beilstein MA, Schumaker KS (2014) Distinct roles for SOS1 in the convergent
evolution of salt tolerance in *Eutrema salsugineum* and *Schrenkiella parvula.* Mol Biol Evol 31:
2094-2107

Jiang C-Z, Yee J, Mitchell DL, Britt AB (1997) Photorepair mutants of *Arabidopsis*. Proc Natl Acad Sci
USA 94: 7441-7445

John UP, Polotnianka RM, Sivakumaran KA, Chew O, Mackin L, Kuiper MJ, Talbot JP, Nugent GD,
Mautord J, Schrauf GE et al. (2009) Ice recrystallization inhibition proteins (IRIPs) and freeze
tolerance in the cryophilic Antarctic hair grass *Deschampsia antarctica* E. Desv. Plant Cell Environ
32: 336-348

Kaya H, Shibahara K, Taoka K, Iwabuchi M, Stillman B, Araki T (2001) *FASCIATA* genes for Chromatin
Assembly Factor-1 in *Arabidopsis* maintain the cellular organization of apical meristems. Cell 104:
131-142

Kazachkova Y, Batushansky A, Cisneros A, Tel-Zur N, Fait A, Barak S (2013) Growth platform-
dependent and -independent phenotypic and metabolic responses of *Arabidopsis* and its halophytic
relative, *Eutrema salsugineum*, to salt stress. Plant Physiol 162: 1583-1598

Kazachkova Y, Eshel G, Pantha P, Cheeseman JM, Dassanayake M, Barak S (2018) Halophytism: What
have we learnt from *Arabidopsis thaliana* relative model systems? Plant Physiol 178: 972-988

Kiefer M, Schmickl R, German DA, Mandakova T, Lysak MA, Al-Shehbaz IA, Franzke A,
Mummenhoff K, Stamatakis A, Koch MA (2014) BrassiBase: Introduction to a novel knowledge
database on brassicaceae evolution. Plant Cell Physiol 55: 1-9

Kimura S, Sakaguchi K (2006) DNA Repair in Plants. Chem Rev 106: 753-766

Koenig D, Weigel D (2015) Beyond the thale: comparative genomics and genetics
of *Arabidopsis* relatives. *Nat Rev Genet* 16**,** 285–298.

Kotake T, Takada S, Nakahigashi K, Ohto M, Goto K (2003) *Arabidopsis* TERMINAL FLOWER 2
gene encodes a Heterochromatin Protein 1 homolog and represses both FLOWERING LOCUS T
to regulate flowering time and several floral homeotic genes.

Kramer, U (2010) Metal hyperaccumulation in plants. Annu Rev Plant Biol 61: 517-534

Landry G, Stapleton AE, Lim J, Hoffman P, Hays JB, Walbot V, Last RL (1997) An *Arabidopsis*
photolyase mutant is hypersensitive to ultraviolet-B radiation. Proc Natl Acad Sci USA 94: 328-
332

Lange H, Ndecky SYA, Gomez-Diaz C. Pflieger D, Butel N, Zumsteg J, Kuhn L, Piermaria C, Chicher
J, Christie M, et al. (2019) RST1 and RIPR connect the cytosolic RNA exosome to the Ski complex
in *Arabidopsis*. Nat Commun **10:** 3871

Langfelder P, Horvath S (2008) WGCNA: An R package for weighted correlation network analysis.
BMC Bioinformatics 9: 559

Laporte D, Olate E, Salinas P, Salazar M, Jordana X, Holuigue L (2012) Glutaredoxin GRXS13 plays a
key role in protection against photooxidative stress in *Arabidopsis*. J Exp Bot 63: 503-515

Lario LD, Ramirez-Parra E, Gutierrez C, Spampinato CP, Casati P (2013) ANTI-SILENCING
FUNCTION1 proteins are involved in ultraviolet-induced DNA damage repair and are cell cycle
regulated by E2F transcription factors in *Arabidopsis*. Plant Physiol 162: 1164-1177

Lawson T, Davey PA, Yates SA, Bechtold U, Baeshen M, Baeshen N, Mutwakil MZ, Sabir J, Baker NR,
Mullineaux PM (2014) C_3 photosynthesis in the desert plant *Rhazya stricta* is fully functional at
high temperatures and light intensities. New Phytol 201: 862-873

Lee JE, Lampugnani ER, Bacic A, Golz JF (2014) SEUSS and SEUSS-LIKE 2 coordinate auxin
distribution and KNOXI activity during embryogenesis. Plant J 80: 122-135

Lesk C, Rowhani P, Ramankutty N (2016) Influence of extreme weather disasters on global crop
production. Nature 529: 84-87

Leyser HMO, Furner IJ (1992) Characterisation of three shoot apical meristem mutants of *Arabidopsis*
thaliana. Development 116: 397-403

Li H, He Z, Lu G, Lee SC, Alonso J, Ecker JR, Sheng Luan (2007) A WD40 domain cyclophilin interacts
with histone H3 and functions in gene repression and organogenesis in *Arabidopsis*. Plant Cell 19:
2403-2416

Li Z, He Y (2010) Roles of brassinosteroids in plant reproduction. Int J Mol Sci 21: 872

Li B, Dewey C (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. BMC Bioinform 12: 323

Li H, Luan S (2011) The cyclophilin AtCYP71 interacts with CAF-1 and LHP1 and functions in multiple
chromatin remodeling processes. Mol Plant 4: 748-758

Li J. Liu J, Wang G, Cha J-Y, Li G, Chen S., Li Z, Guo J, Zhang C, Yang Y et al. (2015). A chaperone
function of NO CATALASE ACTIVITY1 is required to maintain catalase activity and for multiple
stress responses in *Arabidopsis*. Plant Cell 27: 908-925

Liang X, Ding P, Lian K, Wang J, Ma M, Li L, Li L, Li M, Zhang X, She Chen S, et al. (2016)
Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to the FLS2 receptor.
eLife 5: e13568

Lindquist S (1986) The heat shock response. Annu Rev Biochem 55: 1151-1191

Lopez L, Wolf EM, Pires JC, Edger PP, Koch MA (2017) Resources from transcriptomes in the
Brassicaceae family. Front Plant Sci 8: 1488

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biol 15: 550

Lu C, Tian Y, Wang S, Su Y, Mao T, Huang T, Chen Q, Xu Z, Ding Y (2017) Phosphorylation of SPT5
by CDKD;2 is required for VIP5 recruitment and normal flowering in *Arabidopsis thaliana*. Plant
Cell 29: 277-291

Lugan R, Niogret MF, Leport L, Guegan J-P, Larher FR, Savouré A, Kopka J, Bouchereau A (2010)
Metabolome and water homeostasis analysis of *Thellungiella salsuginea* suggests that dehydration
tolerance is a key response to osmotic stress in this halophyte. Plant J 64: 215-229

Mang HG, Laluk KA, Parsons EP, Kosma DK, Cooper BR, Park HC, AbuQamar S, Boccongelli C,
Miyazaki S, Consiglio F, et al. (2009) The *Arabidopsis* RESURRECTION1 gene regulates a novel
antagonistic interaction in plant defense to biotrophs and necrotrophs. Plant Physiology, 151: 290-
305.

Mittler R, Blumwald E (2010) Genetic engineering for modern agriculture: challenges and perspectives.
Annu Rev Plant Biol 61: 443-462

Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue
cultures. Physiol Plant 15: 473-497

Nekrasov V, Li J, Batoux M, Roux M, Chu ZH, Lacombe S, Rougon A, Bittel P, Kiss-Papp M, Chinchilla
D, et al. (2009) Control of the pattern-recognition receptor EFR by an ER protein complex in plant
immunity. EMBO J 28: 3428-3438

Nie X, Wang H, Li J, Holec S, Berger F (2014) The HIRA complex that deposits the histone H3.3 is
conserved in *Arabidopsis* and facilitates transcriptional dynamics. Biology Open 3: 794-802

Nolan TM, Vukasinovic N, Liu D, Russinova E, Yina Y (2020) Brassinosteroids: Multidimensional
regulators of plant growth, development, and stress responses. Plant Cell 32: 295-318

Obaid AY, Sabir JSM, Atef A, Liu X, Edris S, El-Domyati FM, Mutwakil MZ, Gadalla NO, Hajrah NH,
Magdy A. Al-Kordy MA, et al. (2016) Analysis of transcriptional response to heat stress in *Rhazya*
stricta. BMC Plant Biol 16: 252

Obertello M, Krouk G, Katari MS, Runko SJ, Coruzzi GM (2010) Modeling the global effect of the
basic-leucine zipper transcription factor 1 (bZIP1) on nitrogen and light regulation in *Arabidopsis*.
BMC Systems Biol 4: 111

Oh S, Zhang H, Ludwig P, van Nocker S (2004) A mechanism related to the yeast transcriptional
regulator Paf1c is required for expression of the *Arabidopsis* FLC/MAF MADS box gene family.
Plant Cell 16: 2940-2953

Oh DH, Dassanayake M, Bohnert HJ, Cheeseman JM (2012) Life at the extreme: lessons from the
genome. Genome Biol 13: 241

Oh DH, Hong H, Lee SY, Yun DJ, Bohnert HJ, Dassanayake M (2014) Genome structures and
transcriptomes signify niche adaptation for the multiple-ion-tolerant extremophyte *Schrenkiella*
parvula. Plant Physiol 164: 2123-2138

Oh D-H, Barkla BJ, Vera-Estrella R, Pantoja O, Lee S-Y, Bohnert HJ, Dassanayake M (2015) Cell type-
specific responses to salinity – the epidermal bladder cell transcriptome of Mesembryanthemum
crystallinum. New Phytol 207: 627-644

Oscar MA, Barak S, Winters G (2018) The tropical invasive seagrass, *Halophila stipulacea*, has a
superior ability to tolerate dynamic changes in salinity levels compared to its freshwater relative,
Vallisneria americana. Front Plant Sci 9: 950

Ozfidan-Konakci C, Uzilday B, Ozgur R, Yildiztugay E, Sekmen AH, Turkan I (2016) Halophytes as a
source of salt tolerance genes and mechanisms: a case study for the Salt Lake area, Turkey. Funct
Plant Biol 43: 575-589

Pang Q, Chen S, Dai S, Chen Y, Wang Y, Yan X (2010) Comparative proteomics of salt tolerance in
Arabidopsis thaliana and *Thellungiella halophila*. J Proteome Res 9: 2584-2599

Pantha P, Oh DH, Longstreth D, Dassanayake M (2021) Living with high potassium: how plants balance
nutrient acquisition during K-induced salt stress. bioRxiv doi: 10.1101/2021.07.01.450778

Para A, Li Y, Marshall-Colon A, Varala K, Francoeur NJ, Moran TM, Edwards MB, Hackley
C, Bargmann BOR, Birnbaum KD (2014) Hit-and-run transcriptional control by bZIP1 mediates
rapid nutrient signaling in *Arabidopsis*. Proc Natl Acad Sci 111: 10371-10376

Park MY, Kang JY, Kim SY (2011). Overexpression of *AtMYB52* confers ABA hypersensitivity and
drought tolerance. Molecules and Cells 31: 447-454

Patel S, Dinesh-Kumar SP (2008) *Arabidopsis* ATG6 is required to limit the pathogen-associated cell
death response. Autophagy 4: 20-27

Pouteau S, Albertini C (2009) The significance of bolting and floral transitions as indicators of
reproductive phase change in *Arabidopsis*. J Exp Bot 60: 3367-3377

Rizhsky L, Liang H, Mittler R (2002) The combined effect of drought stress and heat shock on gene
expression in tobacco. Plant Physiol 130: 1143-1151

Serrano M, Guzmán P (2004) Isolation and gene expression analysis of *Arabidopsis thaliana* mutants
with constitutive expression of ATL2, an early elicitor-response RING-H2 zinc-finger gene.
Genetics, 167: 919-929

Shabala S (2013) Learning from halophytes: physiological basis and strategies to improve abiotic stress
tolerance in crops. Ann Bot 112: 1209-1221

Sharma N, Cram D, Huebert T, Zhou N Parkin IAP (2007) Exploiting the wild crucifer *Thlaspi arvense*
to identify conserved and novel genes expressed during a plant's response to cold stress. Plant Mol
Biol 63: 171-184

Shrestha B, Guragain B, Sridhar VV (2014) Involvement of co-repressor LUH and the adapter proteins
SLK1 and SLK2 in the regulation of abiotic stress response genes in *Arabidopsis*. BMC Plant Biol
14: 54

Shi D, Ren A, Tang X, Qi G, Xu Z, Chai G, Hu R, Zhou G, Kong Y (2018) MYB52 negatively regulates
pectin demethylesterification in seed coat mucilage. Plant Physiol 176: 2737-2749

Simão FA Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:
3210-3212

Skirycz A, De Bodt S, Obata T, De Clercq I, Claeys H, De Rycke R, Andriankaja M, Van Aken O, Van
Breusegem F, Fernie AR, et al (2010) Developmental stage specificity and the role of
mitochondrial metabolism in the response of *Arabidopsis* leaves to prolonged mild osmotic stress.
Plant Physiol 152: 226-244

Skirycz A, Claeys H, De Bodt S, Oikawa A, Shinoda S, Andriankaja M, Maleux K, Eloy NB, Coppens
F, Yoo S-D, et al (2011) Pause-and-stop: the effects of osmotic stress on cell proliferation during
early leaf development in *Arabidopsis* and a role for ethylene signaling in cell cycle arrest. Plant
Cell 23: 1876-1888

Song YH, Kubota A, Kwon MS, Covington MF, Lee N, Taagen ER, Cintron DL, Hwang DY, Akiyama
R, Hodge SK, et al. (2013) Molecular basis of flowering under natural long-day conditions in
Arabidopsis. Nat Plants 4: 824-835

Stepien P, Johnson GN (2009) Contrasting responses of photosynthesis to salt stress in the glycophyte
Arabidopsis and the halophyte *Thellungiella*: role of the plastid terminal oxidase as an alternative
electron sink. Plant Physiol 149: 1154-1165

Stratonovitch P, Semenov MA (2015) Heat tolerance around flowering in wheat identified as a key trait
for increased yield potential in Europe under climate change. J Exp Bot 66: 3599-3609

Sun X, Li Y, Cai H, Bai X, Ji W, Ding X, Zhu Y.(2012) The *Arabidopsis* AtbZIP1 transcription factor
is a positive regulator of plant tolerance to salt, osmotic and drought stresses. J Plant Res: 125, 429-
438

Taji T, Seki M, Satou M, Sakurai T, Kobayashi M, Ishiyama K, Narusaka Y, Narusaka M, Zhu JK,
Shinozaki K (2004) Comparative genomics in salt tolerance between *Arabidopsis* and *Arabidopsis*-
related halophyte salt cress using *Arabidopsis* microarray. Plant Physiol 135: 1697-1709

Taji T, Komatsu K, Katori T, Kawasaki Y, Sakata Y, Tanaka S, Kobayashi M, Toyoda A, Seki M,
Shinozaki K (2010) Comparative genomic analysis of 1047 completely sequenced cDNAs from an
Arabidopsis-related model halophyte, *Thellungiella halophila*. BMC Plant Biol 10: 261

Takahashi Y, Ebisu Y, Kinoshita T, Doi M, Okuma E, Murata Y, Shimazaki K (2013) bHLH
transcription factors that facilitate K^+ uptake during stomatal opening are repressed by abscisic acid
through phosphorylation. Sci Signal 6: ra48

Thompson TL, Zaady E, Huancheng P, Wilson TB, Martens DA (2006) Soil C and N pools in patchy
shrublands of the Negev and Chihuahuan Deserts. Soil Biol Biochem 38: 1943-1955

Velasco VME, Mansbridge J, Bremner S, Carruthers K, Summers PS, Sung WWL, Champigny MJ,
Weretilnyk EA (2016) Acclimation of the crucifer *Eutrema salsugineum* to phosphate limitation is
associated with constitutively high expression of phosphate-starvation genes. Plant Cell Environ
39: 1818-1834

Ventura Y, Eshel A, Pasternak D, Sagi M (2015) The development of halophyte-based agriculture: past
and present. Ann Bot 115: 529-540

Vera-Estrella R, Barkla BJ, Pantoja O (2014) Comparative 2D-DIGE analysis of salinity responsive
microsomal proteins from leaves of salt-sensitive *Arabidopsis thaliana* and salt-tolerant
Thellungiella salsuginea. J Proteomics 111: 113-127

Wan T, Liu Z, Leitch IJ, Xin H, Maggs-Kölling G, Gong Y, Li Z, Marais E, Liao Y, Dai C (2021) The
Welwitschia genome reveals a unique biology underpinning extreme longevity in deserts. Nat
Commun 12: 4247

Wang G, Pantha P, Tran KN, Oh DH, Dassanayake M (2019) Plant growth and Agrobacterium-mediated
floral-dip transformation of the extremophyte *Schrenkiella parvula*. J Vis Exp 2019: 1-8

Wang G, Oh DH Dassanayake M (2020) GOMCL: a toolkit to cluster, evaluate, and extract non-
redundant associations of Gene Ontology-based functions. BMC Bioinformatics 21: 139.

Wang G, DiTusa SF, Oh DH, Herrmann AD, Mendoza-Cozatl DG, O'Neill MA, Smith AP, Dassanayake
M (2021a) Cross species multi-omics reveals cell wall sequestration and elevated global transcript
abundance as mechanisms of boron tolerance in plants. New Phytol 230: 1985-2000

Wang Y, Impa SM, Jagadish SVK (2021b) The neglected other half - role of the pistil in plant heat stress
responses. Plant Cell Environ 44: 2200-2210

Ward D (2009) The biology of deserts. Oxford: Oxford University Press

Weng M, Yang Y, Feng H, Pan Z, Shen W-H, Zhu Y, Dong A (2014) Histone chaperone ASF1 is
involved in gene transcription activation in response to heat stress in *Arabidopsis thaliana*. Plant
Cell Environ 37: 2128-2138

Widiez T, Kafafia ESE, Girina T, Berr A, Ruffel S, Krouk G, Vayssièresa A, Shen W-H, Coruzzi GM,
Gojon A et al. (2011) HIGH NITROGEN INSENSITIVE 9 (HNI9)-mediated systemic repression

of root NO_3^- uptake is associated with changes in histone methylation. Proc Natl Acad Sci USA
108: 13329-13334

Willing E-M, Piofczyk T, Albert A, Winkler JB, Schneeberger K, Pecinka A (2016) UVR2 ensures
transgenerational genome stability under simulated natural UV-B in *Arabidopsis thaliana*. Nat
Commun | 7: 13522

Wu G, Wang X, Li X, Kamiya Y, Otegui MS, Chory J (2011) Methylation of a phosphatase specifies
dephosphorylation and degradation of activated brassinosteroid receptors. Sci Signal 4: ra29

Wu JX, Li J, Liu Z, Yin J, Chang Z-Y, Rong C, Wu J-L, Bi F-C, Yao N (2015) The *Arabidopsis*
ceramidase AtACER functions in disease resistance and salt tolerance. Plant J 81: 767-780

Xia H, Camus-Kulandaivelu L, Stephan W, Tellier A, Zhang Z (2010) Nucleotide diversity patterns of
local adaptation at drought-related candidate genes in wild tomatoes. Mol Ecol 19: 4144-4154

Xu T Lee K Gu L Kim J-I Kang H (2013) Functional characterization of a plastid-specific ribosomal
protein PSRP2 in *Arabidopsis thaliana* under abiotic stress conditions. Plant Physiol Biochem
73: 405-411

Yamamoto M, Uji S, Sugiyama T, Sakamoto T, Kimura S, Endo T, Nishikawa S (2020) ERdj3B-
mediated quality control maintains anther development at high temperatures. Plant Physiol 182:
1979-1990

Yang ZH (1997) PAML: a program package for phylogenetic analysis by maximum likelihood. Comput
Appl Biosci 13: 555-556

Yang Z (2007) PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol 24: 1586-1591

Yates SA, Chernukhin I, Alvarez-Fernandez R, Bechtold U, Baeshen M, Baeshen N, Mutwakil MZ,
Sabir J, Lawson T, Mullineaux PM (2014) The temporal foliar transcriptome of the perennial C_3
desert plant *Rhazya stricta* in its natural environment. BMC Plant Biol. 14: 2

Yu X, Michaels SD (2010) The *Arabidopsis* Paf1c complex component CDC73 participates in the
modification of FLOWERING LOCUS C chromatin. Plant Physiol 153: 1074-1084

Zhang X, Zhang M, Takano T, Liu, S (2011) Characterization of an *AtCCX5* gene from *Arabidopsis*
thaliana that involves in high-affinity K^+ uptake and Na^+ transport in yeast. Biochemi Biophys Res
Commun 414: 96-100

Zhang X, Lei L, Lai J, Zhao H, Song W (2018a) Effects of drought stress and water recovery on
physiological responses and gene expression in maize seedlings. BMC Plant Biol 18: 68

Zhang L, Chen L, Yu D (2018b) Transcription factor WRKY75 interacts with DELLA proteins to affect
flowering. Plant Physiol 176: 790-803

Zhao Q, Shen J, Gao C, Cui Y, Wang Y, Cui J, Cheng L, Cao W, Zhu Y, Huang S, et al. (2019)
RST1 is a FREE1 suppressor that negatively regulates vacuolar trafficking in *Arabidopsis*. Plant
Cell 31: 2152-2168

Zhou N, Robinson SJ, Huebert T, Bate NJ and Parkin IA (2007) Comparative genome organization
reveals a single copy of CBF in the freezing tolerant crucifer *Thlaspi arvense*. Plant Mol Biol 65:
693-705

Zhou D, Zhou J, Meng L, Wang Q, Xie H, Guan Y, Ma Z, Zhong Y, Chen F, Liu J (2009) Duplication
and adaptive evolution of the *COR15* genes within the highly cold-tolerant *Draba* lineage
(Brassicaceae). Gene 441: 36-44

Zhou Y, Zhu J, Tong T, Wang J, Lin B and Zhang J (2019) A statistical normalization method and
differential expression analysis for RNA-seq data between different species. BMC Bioinformatics
20: 163

Zhu JK, Whited J, Seluanov A, Gorbunova V, Kasahara M, Amdam GV, Ulanovsky N, Feng G, Brunet
A, Margoliash D (2015) The next top models. Cell 163: 18-20

Supporting Information

Fig. S1. Transcriptome sequencing and hybrid assembly workflow.

Fig. S2. An example of a three-day heat wave event in the Dead Sea valley during April 2008.

Fig. S3. Clustering dendrogram of module eigenvalues for *A. thaliana* and *A. hierochuntica*
transcriptome profiles under and heat stress conditions.

Fig. S4. Validation of "between species" RNA-seq analysis.

Fig. S5. Basal (control) expression of 15 orthologous *A. thaliana* and *A. hierochuntica* housekeeping
genes.

Fig. S6. GO-term enrichment analysis of positively selected genes.

Dataset S1. *A. hierochuntica* transcriptome functional annotation.

Dataset S2. *A. thaliana* and *A. hierochuntica* raw read data plus GO-terms, and DEGs.

Dataset S3. Assignment of ortholog pairs modes of expression (log$_2$ fold-change data and WGCNA
modules).

Dataset S4. Early heat-response modules (rlog expression data).

Dataset S5. *A. thaliana* early heat module (Thistle1) gene list.

Dataset S6. *A. thaliana* late heat module (Ivory) gene list.

Dataset S7. *A. hierochuntica* early heat module (Black) gene list.

Dataset S8. *A. hierochuntica* early heat module (Floralwhite) gene list.

Dataset S9. GO-term enrichment of *A. thaliana* and *A. hierochuntica* morning heat-response modules.

Dataset S10. Difference in percent change orthologous gene expression (based on TPM) between *A.*

heirochuntica and *A. thaliana*.

Dataset S11. GO-term enrichment of genes that are more responsive to heat in *A. hierochuntica* (see Fig.

5C).

Dataset S12. Species included in the custom *Brassicaceae* CDS database for the *Brassicaceae*

phylogenomic analysis.

Dataset S13. CODEML positive selected genes ($q<0.05$) in the all extremophyte species run with the

branch-site model.

Dataset S14. CODEML positive selected genes ($q<0.05$) in *A. hierochuntica* run with the branch-site

model.

Dataset S15. CODEML positive selected genes ($q<0.05$) in *E. salsugineum* run with the branch-site

model.

Dataset S16. CODEML positive selected genes ($q<0.05$) in *S. parvula* run with the branch-site model.

Dataset S17. CODEML positive selected genes ($q<0.05$) in *A. thaliana* run with the branch-site model.

Dataset S18. GO-terms over-represented in the all extremophyte species 194 positive selected genes.

Dataset S19. GO-terms over-represented in *A. hierochuntica* 120 positive selected genes.

Dataset S20. GO-terms over-represented in *E. salsugineum* 131 positive selected genes.

Dataset S21. GO-terms over-represented in *S. parvula* 99 positive selected genes.

Dataset S22. GO-terms over-represented in *A. thaliana* 112 positive selected genes.

Dataset S23. 109 most conserved orthologs ($\geq$ 98% query coverage, $\geq$ 99% nucleotide identity, $\geq$ BLAST

value of e-100) between *A. thaliana* and *A. hierochuntica***.**

Dataset S24. PCR primers used in this study.

Methods S1

1012

Figure legends

Figure 1. Geographic distribution of *A. hierochuntica* and *de-novo* reference transcriptome. (A),

Geographic distribution data are based on *Anastatica* L. in the Global Biodiversity Information Facility

database (GBIF Secretariat (2019). GBIF Backbone Taxonomy. Checklist dataset

https://doi.org/10.15468/39omei accessed via GBIF.org on 2021-03-12). Average temperature data for

this region are from 1948 to Feb 2021 acquired by the Physical Sciences Laboratory (Fan and van den

Dool, 2008); (B), Lab-grown and wild *A. hierochuntica* plants. Panels: (i) 40 d-old lab-grown plant. Note

the axillary inflorescence at each branch point; (ii) large mature plant from the Ovda valley population

in the Negev desert. Ruler length = 30 cm; (iii) young seedlings growing near the dead mother plant from

a Neot Smadar population in the Negev desert; (iv) a large population of *A. hierochuntica* in the Ovda

valley with high variation in plant size due to spatial and temporal variations in water availability; (v) *A.*

hierochuntica seedling already beginning to flower after producing four true leaves (Neot Smadar); (vi)

two tiny dead plants (white arrows) from a population near the Dead Sea valley, having already dispersed

their few seeds; (C), Transcript length distribution and *A. hierochuntica* assembly descriptive statistics;

(D), Assessment of reference transcriptome assembly completeness using the Benchmarking Universal

Single-Copy Orthologs (BUSCO) tool (Simão et al., 2015). The percentages of 1,375 single-copy genes,

conserved among land plants, identified in the *A. hierochuntica* transcriptome are shown.

1030

Figure 2. *A. thaliana* and *A. hierochuntica* exhibit similar transcriptome adjustment to heat stress.

(A), Experimental design for *A. thaliana* and *A. hierochuntica* control and heat stress conditions. Control

plants were harvested the day before the initiation of heat stress and on the last day of the experiment

(indicated by vertical arrows) from a parallel 23 °C control chamber. Red and blue circles represent

samples harvested 1.5 h (morning) or 7 h (afternoon) respectively, after onset of light/heat. Each circle

represents 3 independent experiments, each comprising 6 pooled plants. (B and C), Effect of heat stress

on *A. thaliana* and *A. hierochuntica* leaf area (B) and fresh weight (C). Data are mean ± S.D. (n = 5) and

are representative of two independent experiments. Letters above bars indicate significant difference at

$p < 0.05$ (Tukey HSD test). Blue shading, control conditions; Pink shading, heat conditions. (D), Principal

component analysis (PCA) of *A. thaliana* and *A. hierochuntica* transcript levels. Each point represents

one biological replicate and the three replicates for each condition are depicted with the same symbol.

Symbols are explained in the legend box and refer to the experimental design shown in (A); (E),

Comparison of the abundance of 27,416 and 30,670 protein-coding *A. thaliana* and *A. hierochuntica*

transcripts, respectively. Asterisks represent significant difference at $p < 0.05$ (Wilcoxon rank sum test)

between the treatment compared to its respective control. Black vertical line within plots is median

expression; (F), Percent of *A. thaliana* and *A. hierochuntica* differentially expressed genes (DEGs) in

response to heat stress. In total, 17,989 *A. thaliana* and 19,443 *A. hierochuntica* genes were differentially

expressed in response to heat stress in at least one condition (Dataset S2), and % DEGs was calculated

based on 27,416 and 30,670 protein-coding genes for *A. thaliana* and *A. hierochuntica*, respectively. CM,

control morning; CN, control afternoon; HW1M, heat wave 1 morning; HW1N, heat wave 1 afternoon;

HW2N, heat wave 2 afternoon; HW3N, heat wave 3 afternoon; R1N, day 1 recovery from heat stress

afternoon; CR2N, control plants parallel to the R2N time point afternoon; R2N, day 2 recovery from heat

stress afternoon; Union, DEGs identified under either HW1M or HW1N or HW2N or HW3N.

1054

Figure 3. The *A. hierochuntica* transcriptome does not exist in a "stress-ready" state but exhibits a

lower basal expression and higher fold-change expression than *A. thaliana* in response to heat

stress. (A), Modes of expression of ortholog pairs between *A. thaliana* and *A. hierochuntica* in response

to heat stress. WGCNA followed by DESeq2 was used to assign orthologs to response modes (Fig. S3;

Dataset S3). Differences in absolute transcripts levels were identified by comparing TPM minimum or

maximum expression values (Student's *t*-test, $p \leq 0.05$). The green (*A. thaliana*) and orange (*A.*

hierochuntica) lines indicate idealized expression patterns of the ortholog pairs in each species under

control and heat conditions, compared to the *A. thaliana* control (dashed line). Ctrl, control; + heat, heat

stress treatment; SA, genes associated with GO-terms for abiotic stress (Methods S1); (B), Transcript

abundance of 17,962 *A. thaliana* and *A. hierochuntica* ortholog pairs under control conditions. (C),

Combined violin and box plots showing absolute $\log_2$ fold-changes of *A. thaliana* and *A. hierochuntica*

DEGs in response to heat stress (Dataset S2). The median $\log_2$ fold-change is shown as a black square

inside each box plot; (D), Number of DEGs, median $\log_2$ fold-change values and *p*-values for panel C.

CM, control morning; CN, control afternoon; CR2N, control plants parallel to the R2N (day 2 recovery

from heat stress afternoon) time point; HW1M, heat wave 1 morning; HW1N, heat wave 1 afternoon;

HW2N, heat wave 2 afternoon; HW3N, heat wave 3 afternoon; Union, DEGs identified under either

HW1M or HW1N or HW2N or HW3N. Asterisks represent significant difference at $p < 0.05$ (Wilcoxon

rank sum test) between *A. thaliana* and *A. hierochuntica*.

1073

Figure 4. *A. hierochuntica* shared- and unique-expressed orthologs in specific functional groups

display lower basal and greater heat-mediated % change in expression than in *A. thaliana*. All

genes used in this analysis possess a unique AGI code (putative *A. hierochuntica* orthologs were assigned

A. thaliana AGI codes). Genes were chosen based on their association with GO terms for their respective

categories (Methods S1). Basal expression levels were based on CM conditions. % change in expression

from basal level was calculated based on the maximum rlog expression levels of upregulated genes

(abiotic stress [A]) or minimum rlog expression levels of downregulated genes (abiotic stress [B],

photosynthesis [C], cell cycle [D]) in response to heat stress over the three heat waves. Basal and %

change in expression values for all genes in each category are in Dataset S2. For box and whisker plots,

the median (thick black line), the mean (cross below the median line) and interquartile range (IQR) of
the observed differences are shown. Whiskers indicate the maximum/minimum range. Open circles
correspond to extreme observations with values >1.5 times the IQR. Underlined numbers above the
circles indicate the number of shared or unique expressed genes. Letters above the circles indicate
significant differences at *p* < 0.05 (Student's *t*-test). Numbers next to boxes are median values. *A.t*,
Arabidopsis thaliana; *A.h*, *Anastatica hierochuntica*.

1089

Figure 5. The *A. hierochuntica* **early heat response transcriptome displays lower basal and greater**
heat-mediated % change in expression than in *A. thaliana***.** (A), Expression profiles of *A. thaliana*
(left two panels) and *A. hierochuntica* (right two panels) morning and afternoon early heat-response
modules. These modules were assigned standard color-based names by WGCNA (e.g. Thistle, Ivory etc.;
Fig. S3; Datasets S4-S8). Transcript levels were scaled to visualize patterns of expression. The relative
intensity of gene expression (red, upregulated; green, downregulated) is shown in the scale bar. Gene
expression in each condition represents the average of three biological replicates. The number of genes
in each module is shown under the respective module. CM, control morning; CN, control afternoon;
HW1M, heat wave 1 morning; HW1N, heat wave 1 afternoon; HW2N, heat wave 2 afternoon; HW3N,
heat wave 3 afternoon; R1N, day 1 recovery from heat stress afternoon; CR2N, control plants parallel to
the R2N time point afternoon; R2N, day 2 recovery from heat stress afternoon; Blue lines above heat
map, control conditions; Pink lines, heat conditions. (B), Expression of orthologs associated with abiotic
stress GO terms (Dataset S2; Methods S1) Underlined numbers above the circles indicate the number of
shared- or unique-expressed genes. Letters above the circles indicate significant differences at *p* < 0.05
(Student's T-test). Numbers next to boxes are median values. *A.t*, *Arabidopsis thaliana*; *A.h*, *Anastatica*
hierochuntica; (C), Functional clusters enriched among *A. hierochuntica* orthologous genes exhibiting a
higher heat-induced % change in expression than *A. thaliana*. For full reactive gene list see Dataset 10.
Clustering was performed with the GOMCL tool (https://github.com/Guannan-Wang/GOMCL) (Wang
et al., 2020; Methods S1) Clusters are colored differently and labelled with the representative functional
term (Dataset 11). Each node represents a gene ontology (GO) term and node size signifies the number
of genes in the test set assigned to that functional term; the number of genes in each cluster is in
parentheses. The shade of each node represents the *p*-value assigned by the enrichment test (false
discovery rate (FDR)-adjusted *p* < 0.05) with darker shades indicating smaller *p*-values. GO-terms
sharing > 50% of genes are connected by edges. Only selected clusters are highlighted, the rest are greyed
out.

Figure 6. Phylogenomic and comparative positive selection analyses of *A. hierochuntica* and other representative *Brassicaceae* genomes. (A), Maximum-likelihood tree topology based on supermatrix analysis of 13,806 ortholog groups that contain an amino acid sequence from at least four taxa. All nodes are 100% supported by 100 rapid bootstrapping repeats. Red asterisks, extremophyte species; (B), Comparison of the number of positively selected genes among species. Positively selected genes in each species were identified using the AGALMA-PAML pipeline; (C), Comparative GO-term enrichment analysis of positively selected genes. The red color intensity corresponds to the number of positively selected genes assigned with that GO term (the numbers are indicated within the cells). Cells with a white color correspond to GO terms that were not significantly enriched. The *A. thaliana* genome was used as the background gene set and significance (q-value < 0.05) of enrichment was assessed via the Fisher's exact test. For the full list of enriched GO-terms see Fig. S6 and Datasets S18 to S22).

Figure 7. Expression of positively selected genes in response to heat stress. Gene expression was assessed by RNA-seq transcriptome analysis of *A. thaliana* and *A. hierochuntica* plants grown under control conditions or exposed to heat stress (see Fig. 2A for experimental design). Expression is expressed as transcripts per kilobase million (TPM) normalized gene expression. (A), Positively selected genes from the "all extremophyte species" and *A. hierochuntica* analyses that are associated with abiotic stress responses (Table 1); (B), Positively selected *A. hierochuntica* genes that function in photoperiodic flowering, regulation of meristems, and control of morphology (Table 1). Data are mean ± S.D. (n = 3) Asterisks indicate significant difference at $p < 0.05$ between *A. thaliana* and *A. hierochuntica* at the same time point and condition (Student's *t*-test). CM, control morning; CN, control afternoon; HW1M, heat wave 1 morning; HW1N, heat wave 1 afternoon; HW2N, heat wave 2 afternoon; HW3N, heat wave 3 afternoon; R1N, day 1 recovery from heat stress afternoon; CR2N, control plants parallel to the R2N time point afternoon; R2N, day 2 recovery from heat stress afternoon; Blue shading, control conditions; Pink shading, heat conditions.

Table 1. Positively selected genes[a], with a potential role in adaptation to extreme environments. Selected genes from five CODEML branch-site model analyses are indicated based on their *A. thaliana* ortholog identifier. TF, Transcription Factor.

Positively selected gene	Function	References
All extremophyte species (*A. hierochuntica, E. salsugineum, S. parvula, T. arvense* and *A. halleri*)		
AKS2 (At1g05805)	TF; facilitates stomatal opening, ABA response	Takahashi et. al. 2013
ASF1B (At5g38110)	Histone H3/H4 chaperone; repair of UV-B-induced DNA damage, basal and acquired thermotolerance	Lario et al., 2013; Nie et al., 2014; Weng et al., 2014

Gene	Description	References
MYB52 (At1g17950)	TF; ABA response, drought tolerance, involved in the regulation of secondary wall formation, seed mucilage	Park et al., 2011; Cassan-Wang et al., 2013; Shi et al., 2018
PHR1/UVR2 (At1g12370)	Photolyase enzyme; repair of UV-B-induced DNA damage	Ahmad et al., 1997; Landry et al., 1997; Jiang et al., 1997
WRKY75 (At5g13080)	TF; Pi starvation, root development, GA-mediated flowering, defense response	Devaiah et al., 2007; Velasco et al., 2016; Guo et al., 2017; Zhang et al., 2018b

A. hierochuntica

APX6 (At4g32320)	Hydrogen peroxide-scavenging enzyme; alleviation of ROS damage	Chen et al., 2014
bZIP1 (At5g49450)	TF, light and nitrogen sensing, salt and drought tolerance	Obertello et al., 2010; Sun et al., 2012; Para et al., 2014
CYP71 (At3g44600)	Cyclophilin; silencing of homeotic genes; meristem development, interacts with FAS1 and the floral repressor LHP1	Li et al., 2007; Li and Luan, 2011
FAS1 (At1g65470)	Subunit of CaF-1; organization of apical meristems, cellular differentiation, DNA repair	Leyser & Furner, 1992; Kaya et al., 2001; Hisanaga et al., 2013
FBH2 (At4g09180)	TF; photoperiodic flowering	Ito et al., 2012
SBI1/LCMT1 (At1g02100)	Leucine carboxylmethyltransferase; brassinosteroid signaling; flowering, stress responses	Di Rubbo et al., 2011; Wu et al., 2011; Creighton et al., 2017
VIP5 (At1g61040)	PAF1c component; activates floral repressors and photoperiodic pathway regulators. regulation of N uptake	Oh et al., 2004; Yu and Michaels, 2010; Crevillen & Dean, 2011; Widiez et al., 2011; Lu et al., 2017

E. salsugineum

ATCES1/ACER (At4g22330)	Alkaline ceramidase; sphingolipid homeostasis, disease resistance, salt tolerance	Wu et al., 2015
GRXS13 (At1g03850)	Glutaredoxin; Chilling and photooxidative stress tolerance	Laporte et al., 2012; Hu et al., 2015
NCA1 (At3g54360)	Chaperone; regulates catalase 2 (ROS-scavenging enzyme) activity, salt, cold, high pH stresses	Li et al., 2015
PSRP2 (At3g52150)	Plastid-specific ribosomal protein; RNA chaperone activity, negative regulator of seed germination under abiotic stress	Xu et al., 2013
SLK2 (At5g62090)	Transcriptional adaptor; embryogenesis, organ development, repression of stress-responsive gene transcription	Bao et al., 2010; Lee et al., 2014; Shrestha et al., 2014

S. parvula

Atrab28 (At1g03120)	LEA protein; Li^+ tolerance	Borrell et al., 2002
CAX11/CCX5 (At1g08960)	Cation calcium exchanger; K^+ uptake, Na^+ transport in yeast	Zhang et al., 2011
PER1 (At1g48130)	Peroxiredoxin; ROS scavenging, enhances primary seed dormancy	Chen et al., 2020

A. thaliana

ATG6 (At3g61710)	AuTophGy-related protein; autophagy, pathogen defense	Patel & Dinesh-Kumar, 2008
ATL2 (At3g16720)	RING-H2 zinc-finger protein; pathogen defense	Serrano & Guzmán, 2004
ERDJ3B (At3g62600)	ER-localized DNAJ chaperone; anther development under heat stress, pathogen defense	Nekrasov et al., 2009; Yamamoto et al., 2020
RST1 (At3g27670)	ARM-repeat protein; RNA exosome cofactor, vacuolar trafficking, cuticular wax production, embryo development, pathogen defense	Chen et al., 2005; Mang et al., 2009; Lange et al., 2019; Zhao et al., 2019;
XLG2 (At4g34390)	Heterotrimeric G protein; pathogen defense	Liang et al., 2016

1141   [a]For log-likelihood values of the alternative and null models, log-likelihood ratio tests and p-values, see
1142   Datasets S13 to S17.

Figure 1

A

Temperature (°C)

B

(i) (ii) (iii) (iv) (v) (vi)

C

Number of putative protein-coding transcripts	30,670
Transcript N50 length	1,642 bp
Mean ORF length	1,323 bp
Minimum ORF length	106 bp
Maximum ORF length	10,843 bp
% mapped reads to reference assembly	88%
% detected complete BUSCOs	85.6%

A. thaliana
A. hierochuntica

Number of transcripts

Transcript length (bp)

D

% BUSCO genes

6.4 Missing
8.0 Fragmented
13.5 Complete (duplicated)
72.1 Complete (single copy)

Figure 2

Figure 3

A

(a) Stress-ready

	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica
# orthologs	38	183	124	411
# SA orthologs	0	21	7	23

(b) Shared response

	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica
# orthologs	6,054	3,708
# SA orthologs	487	179

(c) Unique response

	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica
# orthologs	209	258	64	64
# SA orthologs	18	22	5	7

(d) Opposite response

	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica
# orthologs	84	169
# SA orthologs	7	18

(e) No response

	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica	A. thaliana / A. hierochuntica
# orthologs	252	228	44
# SA orthologs	21	22	3

B

- A. thaliana
- A. hierochuntica

CM: 3.44 / 3.03 *
CN: 3.40 / 2.93 *
CR2N: 3.50 / 3.06 *

Density — Normalized expression ($\log_2$(TPM+1))

C

- A. thaliana
- A. hierochuntica

Absolute log2 fold-change — HW1M, HW1N, HW2N, HW3N, Union

D

Condition	No. DEGs		Median $\log_2$ fold-change		p-value
	A. thaliana	A. hierochuntica	A. thaliana	A. hierochuntica	
HW1M	10,595	12,109	0.68	0.82	4.35E-63
HW1N	11,635	11,988	0.72	0.75	1.26E-05
HW2N	12,636	14,057	0.76	0.85	5.12E-33
HW3N	12,343	12,612	0.76	0.79	3.38E-02
Union	17,989	19,433	0.89	0.98	2.37E-34

Figure 4

Figure 5

A

A. thaliana			*A. hierochuntica*	

Thistle1 **Heat shock morning**	**Ivory** **Heat shock afternoon**	**Black** **Heat shock morning**	**Floralwhite** **Heat shock afternoon**

CM CN HW1M HW1N HW2N HW3N R1N CR2N R2N

1,112 genes   **325 genes**   **3,001 genes**   **269 genes**

(color scale: 1 / 0 / -1)

B

Basal expression

Shared 63   **Unique** 137   399

Normalized counts (rlog)

Shared: A.t **a** 10.5 ; A.h **a** 9.2
Unique: A.t **a** 9.2 ; A.h **b** 8.0

Heat responsiveness

Shared 63   **Unique** 137   399

% change in gene expression

Shared: A.t **a** 38.6 ; A.h **b** 75.9
Unique: A.t **a** 35.9 ; A.h **b** 66.5

C

Functional clusters enriched in *A. hierochuntica* genes with higher heat mediated % change expression than *A. thaliana*

C1: organ development (1098)
C2: response to hormone stimulus (956)
C5: response to biotic stimulus (682)
C6: reproduction; vegetative to reproductive phase transition of meristem (534)
C7: phosphate metabolic process; regulation of stomatal movement (532)
C9: response to oxidative stress; response to metal ion (285)
C10: response to radiation (261)
C12: response to osmotic stress; response to salt stress (202)
C15: response to DNA damage stimulus (180)
C18: photosynthesis, light reaction (125)
C21: response to water deprivation (109)
C22: potassium ion transport (101)
C24: protein folding (89)
C25: response to wounding (79)
C27: protein modification by small protein conjugation or removal (62)

Figure 6

Figure 7