The mechanism underlying the organization of Borna disease virus inclusion

bodies is unique among mononegaviruses

Yuya Hirai[a*], Keizo Tomonaga[b,c,d*], Masayuki Horie[b,e,f*]

[a]Department of Biology, Osaka Dental University, 8-1, Kuzuha Hanazono-cho, Hirakata, Osaka 573-

1121, Japan.

[b]Laboratory of RNA viruses, Department of Virus Research, Institute for Frontier Life and Medical

Sciences (InFRONT), Kyoto University, 53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507,

Japan

[c]Laboratory of RNA Viruses, Department of Mammalian Regulatory Network, Graduate School of

Biostudies, Kyoto University, 53 Kawahara-cho, Shogoin, Sakyo-ku, 606-8507 Kyoto, Japan.

[d]Department of Molecular Virology, Graduate School of Medicine, Kyoto University, 53 Kawahara-

cho, Shogoin, Sakyo-ku, 606-8507 Kyoto, Japan.

[e]Hakubi Center for Advanced Research, Kyoto University, 53 Kawahara-cho, Shogoin, Sakyo-ku,

Kyoto 606-8507, Japan

[f]Laboratory of Veterinary Microbiology, Division of Veterinary Sciences, Graduate School of Life and

Environmental Sciences, Osaka Prefecture University, 1-58 Rinku-oraikita, Izumisano, Osaka 598-

8531, JAPAN

*Corresponding authors:

Yuya Hirai, PhD

E-mail: hirai-y@cc.osaka-dent.ac.jp

Keizo Tomonaga, DVM, PhD

E-mail: tomonaga@infront.kyoto-u.ac.jp

Masayuki Horie, DVM, PhD

Email: mhorie@vet.osakafu-u.ac.jp

Keywords

Borna disease virus; non-segmented negative-strand RNA viruses; mononegavirus,

nucleoprotein; phosphoprotein; liquid-liquid phase separation; intrinsically disordered

regions; biomolecular condensates; membraneless organelles; nucleus; liquid droplets

Abstract

Inclusion bodies (IBs) are characteristic biomolecular condensates organized by mononegaviruses. Here, we characterize the IBs of Borna disease virus 1 (BoDV-1), a unique mononegavirus that forms IBs in the nucleus, in terms of liquid-liquid phase separation (LLPS). The BoDV-1 phosphoprotein (P) alone induces LLPS and the nucleoprotein (N) is incorporated into the P droplet *in vitro*. In contrast, co-expression of N and P is required for the formation of IB-like structure in cells. Furthermore, while BoDV-1 P binds to RNA, an excess amount of RNA dissolves the liquid droplets formed by N and P. Notably, the N-terminal intrinsically disordered region of BoDV-1 P is essential to drive LLPS and bind to RNA, suggesting that both abilities could compete with one another. These features are unique among mononegaviruses, and thus this study will contribute to a deeper understanding of LLPS-driven organization and RNA-mediated regulation of biomolecular condensates.

Introduction

The order *Mononegavirales* contains many pathogens of great importance to public health, such as measles virus, Ebola virus, rabies virus, and human respiratory syncytial virus (RSV) [1]. These mononegaviruses form various types of inclusion bodies (IBs) in infected cells, which are thought to be the sites of viral replication [2]. The IBs of mononegaviruses are membraneless organelles, also called biomolecular condensates [3–6]. Growing evidence has shown that the IBs of several mononegaviruses, such as rabies virus [7], vesicular stomatitis virus (VSV) [8], measles virus [9,10], and RSV [11], are organized by liquid-liquid phase separation (LLPS).

The viral ribonucleoprotein (vRNP) complexes of mononegaviruses, which function as the fundamental units of transcription and replication, consist of viral genomic RNA, nucleoprotein (N), phosphoprotein (P), and large RNA-dependent RNA polymerase (L). Among the components of vRNP complexes, the expression of N and P is sufficient for the formation of IBs of rabies virus [7], measles virus [9], RSV [11], human metapneumovirus [12], and human parainfluenza virus 3 [13]. Previous studies of these IBs have provided insights into the viral replication strategies, thereby contributing to the control of viral infectious diseases. Additionally, since viruses exploit the host machinery for replication, further studies would be helpful to gain a deeper understanding of the molecular condensates in cells.

Borna disease virus 1 (BoDV-1) is a prototype virus of the family *Bornaviridae* of the order *Mononegavirales*, which causes fatal encephalitis in mammals, including humans [14]. BoDV-1 is a unique mononegavirus as replication occurs in the nucleus of the infected cell, whereas that of most mononegaviruses occurs in the cytoplasm [15]. Consequently, the IBs of cytoplasmic mononegaviruses are formed in the cytoplasm,

while those of BoDV-1, termed as viral speckle of transcripts (vSPOTs), are organized in

the nucleus [16]. The vSPOTs are membraneless spherical structures containing the viral

genome, antigenome, and several viral proteins. The BoDV-1 genome encodes six genes,

N, P, matrix protein (M), envelope glycoprotein (G), L, and accessory protein (X), among

which N, P, M, L, and X are localized in the vSPOTs [17,18].

Although present in the nucleus, the vSPOTs share common properties with those of

the other mononegaviruses, such as membraneless structures, and their components.

However, the surrounding environment is important for the formation of biomolecular

condensates. As described above, BoDV-1 forms IBs in the cell nucleus, where the

surrounding environment differs from that of the cytoplasm. Thus, the IBs of BoDV-1

may be formed by a different mechanism from those of other cytoplasmic

mononegaviruses. Nonetheless, the propensies and minimal components of BoDV-1 IBs

remain entirely elusive.

In this study, to clarify the mechanism of the formation of vSPOTs, we analyzed the

properties of the BoDV-1 N and P proteins from the aspect of phase-separated

condensates. We revealed that the BoDV-1 P protein alone causes LLPS *in vitro*,

depending on the intrinsically disordered region (IDR) present at the N-terminal region.

Although BoDV-1 N alone is not sufficient for the induction of LLPS, it was incorporated

into the BoDV-1 P droplets. Also, co-expression of BoDV-1 N and P is sufficient to induce

the formation of IB-like structures in cells. Further, P of BoDV-1, but not VSV or RSV,

binds to RNA. Interestingly, the N-terminal IDR of BoDV-1 P is involved in this RNA-

biding activity, which is overlapped with the region important for driving LLPS, while an

excess amount of RNA dissolved the liquid droplets consisting of BoDV-1 P alone and

BoDV-1 N and P. These findings are expected to help clarify the fundamental mechanism

of the morphological regulation of the IBs of BoDV-1 and the comprehensive

mechanisms underlying the formation of biomolecular condensates.

Materials and methods

DNA constructs and antibodies

Escherichia coli expression vectors were constructed as follows. Oligonucleotides of

codon-optimized BoDV-1 N, P, His-tagged enhanced green fluorescent protein (EGFP),

and His-tagged mCherry (strain He/80/FR) were synthesized by Thermo Fisher Scientific

(Waltham, MA, USA). The synthesized oligonucleotides of BoDV-1 N and P were

inserted into the *Xho*I and *Bam*HI restriction sites of the plasmid pET-15b using

NEBuilder HiFi DNA Assembly Master Mix (E2621; New England Bioscience, Ipswich,

MA, USA), which were designated as pET-15b-BoDV-1-Nco and pET-15b-BoDV-1-Pco,

respectively. The synthesized oligonucleotides of His-tagged mCherry and EGFP were

cloned into pET-15b using the *Nco*I and *Nde*I restriction sites, which were named pET-

15b-His-mCherry and pET-15b-His-EGFP, respectively. The codon-optimized N and P

sequences were amplified by PCR and then inserted into pET-15b-His-mCherry and pET-

15b-His-EGFP, respectively, using the *Nco*I site.

For mammalian expression vectors, the coding regions of the N and P genes of

BoDV-1 (strain He/80/FR) with the Kozak sequence were amplified by PCR, and then

cloned into the pCAGGS vector at the *Eco*RI and *Xho*I restriction sites using NEBuilder

HiFi DNA Assembly Master Mix. All of the primer and plasmid sequences are available

upon request.

The following antibodies were used in this study: anti-BoDV-1 N mouse monoclonal

antibody (HN132), anti-BoDV-1 P rabbit polyclonal antibody (HB03), anti-high mobility

group box 1 (HMGB1) antibody (sc-74085; Santa Cruz Biotechnology, Inc., Dallas, TX, USA), goat anti-mouse immunoglobulin G (IgG) (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 488 (A-11029; Thermo Fisher Scientific), goat anti-mouse IgG (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 568 (A-11031; Thermo Fisher Scientific), goat anti-rabbit IgG (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 488 (A-11034; Thermo Fisher Scientific), and goat anti-rabbit IgG (H+L) highly cross-adsorbed secondary antibody, Alexa Fluor 568 (A-11036; Thermo Fisher Scientific).

Protein expression and purification

N-terminal His-tagged recombinant proteins were expressed in Rosetta™ 2(DE3) pLysS competent cells (71403; Novagen, Inc., Madison, WI, USA) by induction with 0.5 mM isopropyl β- d-1-thiogalactopyranoside at 30°C for 4 h (N, P, and P'), at 25°C for 5 h (N-mCherry, VSV P, and RSV P), or at 16°C overnight (P-EGFP). The pelleted cells were resuspended in lysis buffer (20 mM Tris-HCl, pH 7.4, 500 mM NaCl, 10 mM imidazole) containing a protease inhibitor cocktail (25955-24; Nacalai Tesque, Inc., Kyoto, Japan). After sonication, the lysates were clarified by centrifugation at $20,000 \times g$ for 10 min at 4°C. Then, the supernatants were loaded onto a column containing TALON® Metal Affinity Resin (635501; Takara Bio, Inc., Shiga, Japan). The resins were washed with lysis buffer and the His-tagged proteins were eluted with elution buffer (20 mM Tris-HCl, pH 7.4, 500 mM NaCl, 300 mM imidazole), dialyzed with dialysis buffer (20 mM Tris-HCl, pH 7.4, 500 mM NaCl), and concentrated with Amicon® Ultra Centrifugal Filters (Merck KGaA, Darmstadt, Germany).

In vitro phase separation assay

The concentrated proteins were diluted to the indicated concentrations in phase separation buffer (20 mM Tris-HCl, pH 7.4, 150 mM NaCl, except for Fig. 1D). RNA and 1,6-hexanediol (H0099; Tokyo Chemical Industry, Tokyo, Japan) were added to the mixed reaction samples at final concentrations of 200 ng/µl and 6%, respectively. Fluorescently labeled liquid droplets were formed by mixing 9.5 µM P and 0.5 µM P-EGFP (P droplets), and 5 µM N, 4.5 µM P, and 0.5 µM P-EGFP (NP droplets). The mixed samples were transferred to the wells of 384-well EZVIEW® glass-bottom assay plates (5883-384; Iwaki Co., Ltd., Chiba, Japan) and observed at room temperature (RT) with a laser scanning confocal microscope (LSM 700; Carl Zeiss AG, Oberkochen, Germany) equipped with a Plan-Apochromat 63X objective lens (numerical aperture = 1.4). The size and intensity of each droplet was calculated using ImageJ software (https://imagej.nih.gov/ij/). The sizes and intensities are presented as the average of at least three separate fields.

In vitro transcription

DNA templates (a BoDV-1 minigenome containing the complementary sequence of Gaussia luciferase [Gluc] flanked by the leader and trailer sequences of the BoDV-1 genome [Fig. S3A] or the complementary sequence of Gluc) for *in vitro* transcription were amplified by PCR using primers containing the T7 promoter sequence at the 5'-end and pBDVmg-Gluc[19] as a template. The *in vitro* transcription reaction was performed using T7 RNA polymerase (2540A; Takara Bio, Inc.) in accordance with the manufacturer's protocol. To label RNA, Cy3-UTP (ENZ-42505; Enzo Life Sciences, Inc., Farmingdale, NY, USA) was added into the *in vitro* transcription reaction at a final

concentration of 0.05 mM. The transcribed RNA was treated with DNaseI, then purified

using the phenol/chloroform isolation method and resuspended in RNase-free water.

Electrophoretic mobility shift assay

Proteins at the indicated concentrations were mixed with 2 μg of RNA, incubated for 30

min at RT, and then separated by 1.5% agarose gel electrophoresis. RNA signals were

detected using GelGreen®, a highly sensitive, non-toxic green fluorescent nucleic acid

dye (41004; Biotium, Inc., Fremont, CA, USA).

Cell culture

Human oligodendroglioma (OL) cells persistently infected with BoDV-1 strain huP2br [20]

were cultured in high-glucose (4.5%) Dulbecco's modified Eagle's medium (DMEM)

(11965092; Thermo Fisher Scientific) supplemented with 5% fetal bovine serum. U-2

osteosarcoma (OS) cells (92022711; European Collection of Authenticated Cell Cultures,

Public Health England, London, England) were cultured in DMEM (08456; Nacalai

Tesque, Inc.) supplemented with 10% fetal bovine serum. The cells were transfected using

Lipofectamine 2000® Transfection Reagent (11668027; Thermo Fisher Scientific) or

Polyethylenimine "Max" (24765-1; Polysciences, Inc., Warrington, PA, USA).

Immunofluorescence microscopy

The cells were cultured on coverslips, fixed with 4% paraformaldehyde for 10 min,

blocked with 5% bovine serum albumin containing 0.5% Triton X-100 for 15 min, and

then probed with primary antibodies for 2 h at RT. Afterward, the cells were washed twice

with phosphate-buffered saline (PBS), incubated with the secondary antibodies and 4′,6′-

diamidino-2-phenylindole for 1 h at RT, then washed three times with PBS, mounted with

ProLong® Diamond Antifade Reagent (Life Technologies, P36961), and observed using

a laser scanning confocal microscope (LSM 700; Carl Zeiss AG) equipped with a Plan-

Apochromat 63× objective lens (numerical aperture = 1.4).

199 Time-lapse imaging of BoDV-1-infected cells

200 BoDV-1-infected OL cells transfected with a plasmid expressing EGFP-tagged P [21] were

201 cultured in a glass-bottom dish and imaged with a C1 confocal laser scanning microscope

202 (Nikon Corporation, Tokyo, Japan) equipped with a CFI Apo Lambda S 60× objective

203 lens (numerical aperture = 1.4), a 37°C heating stage, and lens heater. A series of images

204 were acquired at intervals of 10 s.

206 Fluorescence recovery after photobleaching (FRAP) analysis

207 To induce LLPS, 9 μM N, 1 μM N-mCherry, 9 μM P, and 1 μM P-EGFP were mixed.

208 FRAP analysis was performed using a laser scanning confocal microscope (LSM 700;

209 Carl Zeiss AG). After acquiring 10 images (97.75 ms per frame), the indicated spots were

210 bleached using a 488-nm laser with 15 iterations. Then, 590 images were captured. The

211 fluorescence intensity was quantified using ZEN software (Carl Zeiss AG).

213 Results

214 P triggers and is necessary for the formation of liquid droplets of BoDV-1 N and P

We first determined the viral protein(s) that forms liquid droplets *in vitro*. As described

above, the N and (partial) P proteins are necessary for the formation of liquid droplets by

other mononegaviruses [10,11]. In addition, the IDR is one of several factors that initiate

phase separation [4,22,23]. Therefore, we predicted the IDRs of the BoDV-1 N and P proteins

(Fig. 1A and S1). P was predicted to have a disordered amino acid sequence, especially

at the N-terminal region (Fig. 1A). Notably, the BoDV-1 genome encodes two P isoforms,

P and P', which are translated from alternative start codons [24]. The fully disordered region

was predicted to be present only in P, but not P' (Fig. 1A).

To investigate which proteins cause phase separation, we examined whether P, P',

and/or N form liquid droplets *in vitro*. At a near physiological salt concentration (150 mM

NaCl), P clearly formed liquid droplets at a protein concentration of 10 µM (Fig. 1B),

while N formed aggregate-like structures but not liquid droplets, and P' formed neither

(Fig. 1B). At a protein concentration of 5 µM, but not 2.5 µM, in 150 mM NaCl, P formed

liquid droplets (Fig. 1C). However, in 500 mM NaCl, P failed to initiate a phase separation,

even at a protein concentration of 20 µM (Fig. 1C).

In infected cells, N and P are colocalized within viral biomolecular condensates,

vSPOTs [16]. Therefore, we next investigated the effects of P on the aggregate-like

structures of N by mixing both N and P *in vitro*. Here we mixed fluorescently labeled

proteins (N-mCherry and P-EGFP, both were tagged at the C-terminus) to the reaction

mixture to visually distinguish between the N and P proteins. The fluorescent protein-

labeled N and P were incorporated into droplets formed by unlabeled N and P (Fig. 1D),

respectively, indicating that both fluorescent protein-labeled N and P exhibited the same

behaviors as the unlabeled proteins. When N and P were mixed, the liquid droplets were

observed, which include not only P but also N (Fig. 1D). However, a mixture of N and P'

formed distorted aggregate-like structures with few liquid droplets. (Fig. 1E).

These results suggest that P, but not P', alone can form liquid droplets via the IDR

at the N-terminal region and N forms liquid droplets with P, which is also dependent on

the IDR of P.

Viral condensates formed by BoDV-1 have liquid-like properties

BoDV-1 forms membraneless IBs containing BoDV-1 N and P in the nucleus of the infected cells. Also, we showed that BoDV-1 N and P formed liquid droplets *in vitro* (Fig. 1D and E). These properties are similar to those of cellular biomolecular condensates. However, the detailed characteristics of BoDV-1 IBs remain unclear. Therefore, we examined the potential liquid properties of BoDV-1 IBs. Time-lapse imaging of BoDV-1-infected cells revealed the fusion of two vSPOTs (Fig. 2A). Considering BoDV-1 P is continuously exchanged between vSPOTs [25], this result indicates that vSPOTs, like other biomolecular condensates, also have liquid properties.

Next, we investigated and compared the properties of the liquid droplets formed by P alone (P droplets) and by both N and P (NP droplets) by a simplified *in vitro* system. Fusion events were observed with the P and NP droplets (Fig. 2B and C), suggesting that both have liquid properties. However, the NP droplets required a much longer time for two droplets to form one spherical droplet as compared to the P droplets (Fig. 2B and C). Additionally, the P droplets were completely dissolved by 1,6-hexanediol (1,6-HD), which is an aliphatic alcohol that affects the formation of liquid droplets [26], while the NP droplets were relatively more resistant to 1,6-HD, although the size and solubility were reduced (Fig. 2D). FRAP analysis showed that the fluorescence recovery of P-EGFP was slower within the NP droplets than the P droplets (Fig. 2E). These results suggest that N decreases the fluidity and increases the stability of droplets.

Both N and P are required for the formation of the condensates in cells

The BoDV-1 nuclear IBs (i.e., vSPOTs) contain several viral proteins (N, P, X, M, and L),

viral RNA, and the host protein HMGB1 [16,17]. Although other mononegaviruses, such as

rabies virus, measles virus, RSV, metapneumovirus, and human parainfluenza virus 3,

require the expression of both N and P for the formation of viral IBs in uninfected cells,

the proteins needed for the formation of vSPOTs in cells remain unknown. Hence, we

expressed N alone, P alone, and both N and P in uninfected U-2 OS cells and observed

them by immunofluorescence staining and confocal microscopy. When N alone was

expressed, condensates were not observed in the nucleus, although aggregates were

observed in the cytoplasm (Fig. 3A). The expression of P alone did not induce the

formation of nuclear condensates (Fig. 3B). On the other hand, the expression of both N

and P induced the formation of nuclear condensates that were morphologically similar to

vSPOTs (Fig. 3C, arrowheads). The nuclear condensates induced by the expression of N

and P contained HMGB1, a host factor that is known to be localized to the vSPOTs of

infected cells (Fig. 3C). Note that the condensates containing both N and P were observed

in the nucleus as well as the cytoplasm (Fig. 3C, arrows). The expression of N and P' did

not induce the formation of condensates (Fig. 3D), which is consistent with the *in vitro*

result (Fig. 1E). These results suggest that N and P are sufficient for the formation of

BoDV condensates in cells, as also reported for other mononegaviruses, and P, but not P',

contributes to the formation of the condensates.

RNA controls the LLPS of BoDV-1 condensates.

The N of mononegaviruses encapsidates RNA and forms nucleocapsids. P, also a

component of RNPs, functions as a cofactor of L, which is a RNA-dependent RNA

polymerase [18,27]. From this perspective, there is a possible, although elusive, relationship

between P and RNA. Therefore, we investigated possible interactions between P and the

BoDV-1 minigenome RNA (mgRNA), which contains the leader and trailer of BoDV-1

and Gluc sequences (Fig. S3A) [19], with the use of the electrophoretic mobility shift assay.

A band shift was observed with an increasing concentration of P, suggesting interactions

with RNA (Fig. 4A). On the other hand, P' did not interact with RNA. Note that a band

shift was also observed when using RNA lacking the 3' leader and 5' trailer regions (Fig.

S3B), suggesting that P interacts with RNA in a sequence-independent manner. No

apparent shift was observed with the P proteins of other mononegaviruses (i.e., VSV and

RSV) (Fig. 4A).

We also investigated the possible incorporation of mgRNA into the droplets. We

observed that mgRNA was incorporated into both the P and NP droplets (Fig. 4B and C),

consistent with the RNA-binding activity of BoDV-1 P (Fig. 4A). However, the signal

intensity of RNA was much stronger in the NP than P droplets, which probably reflects

the role of BoDV-1 N in encapsidating viral genomic RNA.

RNA regulates the formation of liquid droplets formed by RNA-binding proteins [28–

31]. Therefore, we analyzed the effects of adding RNA to the P and NP droplets. The results

showed that mgRNA inhibited the formation of not only P but also NP droplets (Fig. 4B

and C, S1C). These results suggest that RNA is incorporated into the viral condensates in

cells, but an excess amount of RNA could begin to dissolve the viral condensates (see

Discussion).

Discussion

Many mononegaviruses form IBs in infected cells. These IBs are thought to be the sites

of viral transcription and replication, thus it is important to elucidate the properties and

organizations of viral IBs to gain a deeper understanding of the replication strategies of

viruses. Additionally, because viral IBs are biomolecular condensates formed by

exogenous materials in cells, it is helpful to understand the properties of biomolecular

condensates of organisms. In this study, to reveal the organization of condensates formed

by BoDV-1, we analyzed the basic properties of BoDV-1 N and P. Similar to other

cytoplasmic mononegaviruses, co-expression of BoDV-1 N and P was sufficient to

reconstruct the viral IB-like structures in cells (Fig. 3). On the other hand, BoDV-1 P

alone drove the formation of liquid droplets *in vitro*, which was dependent on the IDR

(Fig. 1). This feature is distinct from other mononegaviruses that needs both N and P to

form liquid droplets. Further, P of BoDV-1, but not other mononegaviruses (i.e., VSV and

RSV), binds to RNA and high concentrations of RNA dissolved droplets formed by

BoDV-1 N and P *in vitro* (Fig. 4 and S3). These findings revealed that although BoDV-1

shares common strategies with other mononegaviruses for the formation of IBs, at least

to a certain extent, BoDV-1 apparently adopted strategies different from those of other

mononegaviruses.

The unique feature of BoDV-1 P to form liquid droplets without N may be attributed

to the overlap of the region important for driving LLPS with the RNA-binding region.

We revealed that the N-terminal IDR is important for the RNA-binding activity of BoDV-

1 P (Fig. 4), which is overlapped with the region necessary to drive LLPS *in vitro* (Fig.

1). Further, an excess amount of RNA dissolved the BoDV-1 P droplets (Fig. 4). These

findings suggest that the LLPS and RNA-binding activities of BoDV-1 might be

competing against each other. Notably, a high concentration of RNA is reportedly present

in the cell nucleus [28]. Thus, BoDV-1 P may have a greater ability to drive LLPS than the

other P proteins of mononegaviruses to induce LLPS in the nuclear environment.

The transcription and replication of BoDV-1 might be regulated by the competitive

feature of BoDV-1 P. We revealed that an excess amount of RNA also dissolved the NP

droplets, even though RNA was efficiently incorporated into the NP droplet, which was

probably due to the ability of BoDV-1 N to encapsidate RNA (Fig. 4). Considering that

BoDV-1 may control transcription and replication to establish persistent infection in the

nucleus, we propose the following hypothetical mechanism underlying the regulation of

viral transcription and replication: if viral transcription and replication are excessive, the

excess RNA binds to the N-terminal IDR of BoDV-1 P, leading to dissolving the viral IBs

to arrest them. Although the detailed relationship between the LLPS driving and RNA-

binding abilities of BoDV-1 P remain unclear, this hypothesis could explain one of the

mechanisms needed to establish a persistent infection in the host cell.

Our data suggest that BoDV-1 N plays an important role in the formation of viral

IBs in cells. Although our data suggest that BoDV-1 P has greater ability to drive LLPS,

co-expression of BoDV-1 N was needed to induce the formation of viral IB-like structures

in cells (Fig. 3). Importantly, we could compare the properties of the P and NP droplets

in vitro because BoDV-1 P alone can form liquid droplets, which makes it possible to

speculate the effect of BoDV-1 N on viral condensates. Notably, the NP droplets had

lower fluidity and were more resistant to treatment with 1,6-HD than the P droplets (Fig.

2). Our previous studies revealed that BoDV-1 IBs had insoluble properties and that

BoDV-1 N has an immobile property [16,32]. Thus, besides the function as a nucleocapsid

protein (i.e., encapsidation of viral genomic RNA), BoDV-1 N might also convey the

properties of insolubility and/or viscosity to viral IBs, which might lead to stable

transcription and replication.

In this study, we did not test the effects of viral proteins other than BoDV-1 N and P.

As described above, other viral proteins, including X, M, L, are colocalized in the vSPOTs.

Although our data indicated that both N and P are sufficient for the formation of

condensates as phase-separated structures, other viral components may affect the

formation, structure, localization, and function of viral condensates as reported for other

mononegaviruses [33]. Indeed, some of the condensates formed by BoDV-1 N and P were

observed in the cytoplasm (Fig. 3), although cytoplasmic viral IBs were barely observed

in BoDV-1-infected cells. Hence, further analyses are necessary to elucidate the

organization and function of viral IBs.

LLPS is adopted not only in mononegaviruses but also in many other RNA viruses,

including severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). In SARS-

CoV-2, the nucleocapsid protein forms condensates by LLPS in an RNA-dependent

manner [30,31,34–38]. Remarkably, a higher amount of RNA was reported to dissolve the

condensates formed by the nucleocapsid protein and RNA [30,31,36,37], which is consistent

with the results of the present study (Fig. 4), as well as other cellular proteins [28]. This

suggests that regulation of the phase-separated condensates by RNA is a common

mechanism not only among both positive-strand and negative-strand RNA viruses but

also among cells. Further analyses focusing on LLPS are warranted to elucidate the

various phenomena in the life cycle of RNA viruses, leading to the development of anti-

viral reagents targeting phase-separated condensates.

Taken together, the results of this study will help to elucidate the mechanisms

underlying the formation of phase-separated condensates, which should be useful to

further develop antivirals and understand the cellular biology.

Acknowledgments

We are grateful to Takehiro Kanda and Chiaki Tanaka (Kyoto University) for technical

support. We would like to thank Enago (www.enago.jp) for the English language reveiw.

Funding

This work was supported in part by JSPS KAKENHI grant numbers JP19K16133 (YH),

JP18K19443 (MH), and JP21H01199 (MH); JP19K22530 (KT) and JP20H05682 (KT);

MEXT KAKENHI grant number JP 19H04833 (MH), JP16H06429 (KT), JP16K21723

(KT), JP16H06430 (KT); Hakubi Project at Kyoto University (MH), JSPS Core-to-Core

Program (KT); and the Joint Usage/Research Center Program on inFront, Kyoto

University.

Author contributions

YH and MH conceived and designed the study. YH and MH performed experiments. YH

and MH analyzed the data, and all authors discussed the data interpretation. YH wrote the

initial draft of the manuscript, and all authors revised the manuscript.

Competing interests

The authors declare no competing interests.

References

1.   Kuhn, J. H. *et al. 2020 taxonomic update for phylum Negarnaviricota (Riboviria:*

Orthornavirae), including the large orders Bunyavirales and Mononegavirales.

Archives of Virology **165**, (2020).

2.   Nevers, Q., Albertini, A. A., Lagaudrière-Gesbert, C. & Gaudin, Y. Negri bodies

and other virus membrane-less replication compartments. *Biochim. Biophys. Acta*

- Mol. Cell Res. **1867**, 118831 (2020).

3.   Brangwynne, C. P. *et al.* Germline P Granules Are Liquid Droplets That Localize

by Controlled Dissolution/Condensation. *Science (80-.).* **324**, 1729–1732 (2009).

4.   Molliex, A. *et al.* Phase Separation by Low Complexity Domains Promotes

Stress Granule Assembly and Drives Pathological Fibrillization. *Cell* **163**, 123–

133 (2015).

5.   Berry, J., Weber, S. C., Vaidya, N., Haataja, M. & Brangwynne, C. P. RNA

transcription modulates phase transition-driven nuclear body assembly. *Proc.*

Natl. Acad. Sci. **112**, E5237–E5245 (2015).

6.   Altmeyer, M. *et al.* Liquid demixing of intrinsically disordered proteins is seeded

by poly(ADP-ribose). *Nat. Commun.* **6**, (2015).

7.   Nikolic, J. *et al.* Negri bodies are viral factories with properties of liquid

organelles. *Nat. Commun.* **8**, 1–12 (2017).

8.   Heinrich, B. S., Maliga, Z., Stein, D. A., Hyman, A. A. & Whelan, S. P. J. Phase

transitions drive the formation of vesicular stomatitis virus replication

compartments. *MBio* **9**, 1–10 (2018).

9.   Zhou, Y., Su, J. M., Samuel, C. E. & Ma, D. Measles Virus Forms Inclusion

Bodies with Properties of Liquid Organelles. *J. Virol.* **93**, 1–18 (2019).

10. Guseva, S. *et al.* Measles virus nucleo- and phosphoproteins form liquid-like

phase-separated compartments that promote nucleocapsid assembly. *Sci. Adv.* **6**,

eaaz7095 (2020).

11. Galloux, M. *et al.* Minimal Elements Required for the Formation of Respiratory

Syncytial Virus Cytoplasmic Inclusion Bodies In Vivo and In Vitro. *MBio* **11**, 1–

16 (2020).

12. Derdowski, A. *et al.* Human metapneumovirus nucleoprotein and phosphoprotein interact and provide the minimal requirements for inclusion body formation. *J. Gen. Virol.* **89**, 2698–2708 (2008).

13. Zhang, S. *et al.* An amino acid of human parainfluenza virus type 3 nucleoprotein is critical for template function and cytoplasmic inclusion body formation. *J Virol* **87**, 12457–12470 (2013).

14. Kuhn, J. H. *et al.* Taxonomic reorganization of the family Bornaviridae. *Arch. Virol.* **160**, 621–632 (2015).

15. Cubitt, B. & de la Torre, J. C. Borna disease virus (BDV), a nonsegmented RNA virus, replicates in the nuclei of infected cells where infectious BDV ribonucleoproteins are present. *J. Virol.* **68**, 1371–1381 (1994).

16. Matsumoto, Y. *et al.* Bornavirus closely associates and segregates with host chromosomes to ensure persistent intranuclear infection. *Cell Host Microbe* **11**, 492–503 (2012).

17. Hirai, Y. *et al.* Borna disease virus assembles porous cage-like viral factories in the nucleus. *J. Biol. Chem.* **291**, 25789–25798 (2016).

18. Schneider, U., Naegele, M., Staeheli, P. & Schwemmle, M. Active borna disease virus polymerase complex requires a distinct nucleoprotein-to-phosphoprotein ratio but no viral X protein. *J. Virol.* **77**, 11781–9 (2003).

19. Reuter, A. *et al.* Synergistic antiviral activity of ribavirin and interferon-α against parrot bornaviruses in avian cells. *J. Gen. Virol.* **97**, 2096–2103 (2016).

20. Nakamura, Y. *et al.* Isolation of Borna disease virus from human brain tissue. *J. Virol.* **74**, 4601–11 (2000).

21.   Kobayashi, T. *et al.* Modulation of Borna disease virus phosphoprotein nuclear

localization by the viral protein X encoded in the overlapping open reading

frame. *J. Virol.* **77**, 8099–107 (2003).

22.   Lin, Y., Protter, D. S. W., Rosen, M. K. & Parker, R. Formation and Maturation

of Phase-Separated Liquid Droplets by RNA-Binding Proteins. *Mol. Cell* **60**,

208–219 (2015).

23.   Nott, T. J. *et al.* Phase Transition of a Disordered Nuage Protein Generates

Environmentally Responsive Membraneless Organelles. *Mol. Cell* **57**, 936–947

(2015).

24.   Kobayashi, T., Watanabe, M., Kamitani, W., Tomonaga, K. & Ikuta, K.

Translation initiation of a bicistronic mRNA of Borna disease virus: A 16-kDa

phosphoprotein is initiated at an internal start codon. *Virology* **277**, 296–305

(2000).

25.   Charlier, C. M. *et al.* Analysis of borna disease virus trafficking in live infected

cells by using a virus encoding a tetracysteine-tagged p protein. *J. Virol.* **87**,

12339–48 (2013).

26.   Lin, Y. *et al.* Toxic PR Poly-Dipeptides Encoded by the C9orf72 Repeat

Expansion Target LC Domain Polymers. *Cell* **167**, 789-802.e12 (2016).

27.   Perez, M., Sanchez, A., Cubitt, B., Rosario, D. & de la Torre, J. C. A reverse

genetics system for Borna disease virus. *J. Gen. Virol.* **84**, 3099–104 (2003).

28.   Maharana, S. *et al.* RNA buffers the phase separation behavior of prion-like RNA

binding proteins. *Science* **360**, 918–921 (2018).

29.   Zhang, H. *et al.* RNA Controls PolyQ Protein Phase Transitions. *Mol. Cell* **60**,

220–230 (2015).

30.   Lu, S. *et al.* The SARS-CoV-2 nucleocapsid phosphoprotein forms mutually

483   exclusive condensates with RNA and the membrane-associated M protein. *Nat.*

484   *Commun.* **12**, 502 (2021).

485   31.   Iserman, C. *et al.* Genomic RNA Elements Drive Phase Separation of the SARS-

486   CoV-2 Nucleocapsid. *Mol. Cell* **80**, 1078-1091.e6 (2020).

487   32.   Hirai, Y., Honda, T., Makino, A., Watanabe, Y. & Tomonaga, K. X-linked RNA-

488   binding motif protein (RBMX) is required for the maintenance of Borna disease

489   virus nuclear viral factories. *J. Gen. Virol.* **96**, 3198–203 (2015).

490   33.   Ringel, M. *et al.* Nipah virus induces two inclusion body populations:

491   Identification of novel inclusions at the plasma membrane. *PLoS Pathog.* **15**,

492   (2019).

493   34.   Perdikari, T. M. *et al.* SARS-CoV-2 nucleocapsid protein phase-separates with

494   RNA and with human hnRNPs. *EMBO J.* **39**, 1–15 (2020).

495   35.   Savastano, A., Ibáñez de Opakua, A., Rankovic, M. & Zweckstetter, M.

496   Nucleocapsid protein of SARS-CoV-2 phase separates into RNA-rich

497   polymerase-containing condensates. *Nat. Commun.* **11**, 6041 (2020).

498   36.   Carlson, C. R. *et al.* Phosphoregulation of Phase Separation by the SARS-CoV-2

499   N Protein Suggests a Biophysical Basis for its Dual Functions. *Mol. Cell* **80**,

500   1092-1103.e4 (2020).

501   37.   Cubuk, J. *et al.* The SARS-CoV-2 nucleocapsid protein is dynamic, disordered,

502   and phase separates with RNA. *Nat. Commun.* **12**, 1936 (2021).

503   38.   Zhao, M. *et al.* GCG inhibits SARS-CoV-2 replication by disrupting the liquid

504   phase condensation of its nucleocapsid protein. *Nat. Commun.* **12**, 2114 (2021).

505   39.   Mészáros, B., Erdős, G. & Dosztányi, Z. IUPred2A: Context-dependent

506   prediction of protein disorder as a function of redox state and protein binding.

507   *Nucleic Acids Res.* **46**, W329–W337 (2018).

Figure legends

Fig. 1 IDR-dependent formation of liquid droplets by BoDV-1 N and P.

A. A schematic diagram of the amino acid regions of p24 and p16. The IDRs were predicted by IUPred2A [39]. **B.** Differential interference contrast (DIC) images of the liquid droplet formation of BoDV-1 P (p24), P (p16), and N. **C.** DIC images of the liquid droplet formation of the indicated protein concentration of BoDV-1 P (p24) in 150 mM and 500 mM NaCl. **D.** DIC and fluorescence images of the liquid droplet formation of the indicated protein combination. **E.** DIC images of the liquid droplet formation of the indicated protein combination. The arrowheads indicate the distorted aggregate-like structures and the arrow indicate the droplets that were scarcely observed.

Fig. 2 The liquid-like properties the BoDV-1 condensates.

A. Time-lapse imaging of the fusion events of vSPOTs. P-EGFP was expressed in BoDV-1-infected OL cells and observed by confocal microscopy. **B** and **C**. Time-lapse DIC imaging of the fusion events of droplets formed by BoDV-1 P alone (**B**) or N and P (**C**). **D**. Dissolution of P and NP droplets by the addition of 6% 1,6-HD. The area and intensity of fluorescence per unit area were calculated. Statistical significance was calculated with the two-tailed unpaired *t*-test. **E.** Representative images in the FRAP experiment of P and NP droplets (left) and the recovery curves of the fluorescence intensity of P-EGFP after photobleaching (right). The intensities are presented as relative values for which the mean intensities before photobleaching were set to a value of 1.

Fig. 3 Condensates formed by BoDV-1 N and P in cells.

BoDV-1 N alone (**A**), P (p24) alone (**B**), N and P (p24) (**C**), and N and P (p16) (**D**) were

expressed in U-2 OS cells, which were subjected to immunofluorescence staining

followed by confocal microscopy. The magnified images of the areas indicated by the

rectangles are inserted (**C**, top).

Fig. 4 The regulation of the BoDV-1 liquid droplets by RNA.

A. The indicated concentration of BoDV-1 P (p24), P (p16), VSV P, or RSV P was mixed

with the BoDV-1 minigenome RNA and subjected to the electrophoretic mobility shift

assay. **B** and **C**. Fluorescence signals of P (**B**) and NP (**C**) droplets were observed by

confocal microscopy in the presence or absence of 200 ng/µl of Cy3-labeled BoDV-1

minigenome RNA. The area and intensity of fluorescence per unit area were calculated.

Statistical significance was calculated with the two-tailed unpaired *t*-test.

Fig.1

Bar: 5 µm

Fig.2

A P-EGFP, infected cells

0 s 10 s 20 s 30 s

40 s 50 s 60 s 70 s

Bar: 5 μm

B P droplets

0 s 1 s 2 s 3 s 4 s

5 s 6 s 7 s 8 s 9 s

Bar: 5 μm

C NP droplets

0 s 10 s 20 s 30 s 40 s 50 s

60 s 70 s 80 s 90 s 100 s 110 s

Bar: 5 μm

D Ctrl 1,6-HD

P

NP

Bar: 5 μm

NP droplets

Area (μm²) — Ctrl, 1,6-HD — *p* < 0.0001

NP droplets

Intensity (a.u.) — Ctrl, 1,6-HD — *p* < 0.0001

E pre-bleach 0 s 10 s 20 s 40 s

P

NP

Bar: 5 μm

Rerative intensity vs Time (s)
- P droplets
- NP droplets

Fig.3

Bar: 5 µm

Fig.4

A

B

C