Testing the maintenance of natural responses to survival-relevant calls in the conservation
breeding population of a critically endangered corvid (*Corvus hawaiiensis*)

Anne C. Sabol[1,2], Alison L. Greggor[1], Bryce Masuda[1], Ronald R. Swaisgood[1]

[1]Conservation Science and Wildlife Health, San Diego Zoo Wildlife Alliance, Escondido,
USA

[2]Department of Biological Sciences, Florida International University

Abstract:

Vocal communication serves an important role in driving animals' social interactions and
ultimately their survival. However, natural vocal behavior can erode in human care. Determining
if animals in conservation breeding programs exhibit and respond appropriately to species-specific
vocalizations is therefore important for ensuring their survival post-release. We tested whether
endangered 'alalā (*Corvus hawaiiensis*), which are extinct in nature, have retained their natural
responses to survival-relevant vocal calls. We conducted our studies on breeding populations
derived from a small number of founding 'alalā maintained in human care since their extinction in
the wild in 2002. We presented pairs of 'alalā with alarm, territorial intrusion, and two types of
control playback calls (a non-threatening territorial maintenance call and a novel heterospecific
call). 'Alalā were significantly more likely to approach the speaker following alarm call playback
than other call types, and were more likely to respond to territorial intrusion calls with the same
aggressive territorial calls. Males were more likely to make these aggressive calls than females,
mirroring their roles in territory defense. We also found individual consistency in the level of vocal
behavior response across all call types, indicating that some individuals are more vocal than others.
These results are encouraging, showing that 'alalā exhibit relevant, species-specific behaviors
despite generations under human care. It does illustrate, however, that all individuals do not
respond uniformly, so vocal response may be an important factor to consider in determining the
release suitability of individuals.

Keywords: anti-predator, captive breeding, communication, corvid, translocation, territory
defense, vocalization

Introduction

Animals have evolved mechanisms for communication that facilitate survival and
reproduction. For primarily vocal species, how individuals respond to conspecific vocal cues and
signals can have survival-relevant consequences. For instance, failing to respond to an alarm call
could result in predation, or failing to adequately broadcast territorial calls could cause a loss of
territory, mates, or offspring. Whether or not animals exhibit appropriate species-specific
communication is particularly important in conservation breeding programs. Animals that have
spent generations in human care need to retain their natural behaviors for successful return to the
wild (Rabin 2003; McPhee 2004 a, b). However, the loss of survival-relevant behaviors (McPhee
& Carlstead, 2010; Shier 2016), including losses in vocal diversity (Corfield et al., 2008, Digby et
al 2013) and dialect drift (Lewis et al., 2021), are a common byproduct of conservation breeding
settings. A combination of forces from evolution, culture (Brakes et al., 2019), and direct
experience interact to shape animals' vocal behaviors (Bradbury & Vehrencamp, 2011; Hollén &
Radford, 2009), all of which may be impacted by the altered environment in conservation breeding
facilities. Management of animals in human care, therefore, faces the challenge of providing
opportunities for animals to maintain and express these behaviors (Greggor et al., 2018).

Among the vocal signals that animals need to retain, anti-predator signals (alarm calls) are
particularly important because they are a component of an animal's defense against predators
(Hollen & Radford, 2009). Developmentally, they may be more canalized, with alarm call
production and responses emerging in development without direct experience with predators.
While learning is undoubtedly important to fine-tune production and responses to antipredator
calls (Griffin et al., 2000), selection should favor individuals that respond appropriately to alarm
calls without direct experience associating alarm calls with predation. However, some antipredator
behavior responsiveness can be lost over an individual's lifetime in the absence of predator
exposure (Muralidhar et al., 2019). The importance of antipredator alarm calling for conservation
breeding and translocation programs is underscored by the finding that predation is one of the
primary causes of mortality for animals after release to the wild (Fischer & Lindenmayer, 2000,
Moseby et al., 2011, Berger-tal et al., 2020).

Another important set of signals helps animals avoid conspecific conflict by alerting others
to their shared or defended resources. Territorial species, for instance, have evolved multiple forms

of communication that broadcast an intent to defend their territory. From ornamental displays, to
physical demonstrations (e.g., Decourcy & Jenssen, 1994), or auditory signals (Greenfield &
Minckley, 1993), there are a variety of ways that animals communicate territory occupancy to
avoid unnecessary conflict. These calls often serve to communicate motivation and a level of
severity of the territorial threat (Mager et al., 2012), ranging from maintenance calls to aggressive
intrusion calls. The ability to efficiently set up and defend a territory can be critical to breeding
and later reproductive success (Hiebert et al., 1989). Therefore, for territorial species that rely on
vocal communication, having an ability to understand and respond to different territorial calls may
be crucial for survival and reproduction in the wild. Any divergence in the production or perception
of these important calls from wild phenotypes may challenge social integration and the success of
translocation outcomes (Lewis et al., 2021).

Here we examine the communication system of the endangered 'alalā (*Corvus hawaiiensis*)
within conservation breeding facilities, with the goal of evaluating their responses to major
categories of conspecific vocal signals. Ultimately, we aim to determine if they have retained
important components of natural behaviors that will facilitate successful reintroduction. The 'alalā,
or Hawaiian crow, is the last remaining corvid species of the Hawaiian islands (Banko et al., 2002).
They are keystone species for the Hawaiian wet and mesic forests, and the only known seed
disperser for a number of native plants (Culliney et al., 2012). They went extinct in the wild in
2002, after decades of rapid decline, surviving only in conservation breeding facilities (USFWS,
2009). Efforts are ongoing to reintroduce 'alalā back into the wild, yet the birds still face a long
road to recovery and have not yet demonstrated successful breeding in the wild.

There have been but a few studies on 'alalā vocal behavior, yet we know they have a diverse
vocal repertoire (Tanimoto et al., 2017a). Based on the complexity of vocal communication in
other corvids and the role it plays in their social lives (Clayton & Emery, 2007), we would expect
'alalā calls to vary in meaning, with important fitness consequences. 'Alalā are fiercely territorial
as adults, but as juveniles they form flocks, and associate with members of both sexes. Historically,
breeding pairs, and especially males, would make a number of different, frequent territorial
broadcast calls on the edges of their large territories (Banko et al., 2002); however, the use of some
of these calls appears to have declined in breeding facilities (Tanimoto et al., 2017b). Additionally,
like many species, 'alalā are known to employ a range of alarm calls to warn others of danger

(Tanimoto et al., 2017a, Greggor et al., in review). While comparatively little was known about

ʻalalā calls when the species became extinct in the wild, evidence suggests that the frequency and

type of calls they make in the conservation breeding centers differ from their historical vocal

behavior in the wild (Tanimoto et al., 2017b). It is unclear whether these changes are due to the

erosion of natural behavior in the conservation breeding facilities, or due to a lack of context for

expressing wild-type calls. Additionally, given that they are housed a much higher densities than

wild ʻalalā and suffer reproductive consequences from close proximity of conspecifics (Flanagan

et al., 2020), it is possible individuals in the conservation breeding population have become

desensitized to their social surroundings, and no longer respond appropriately to these important

signals. Thus far there have been no studies examining behavioral responses to experimental

playbacks of vocal calls in ʻalalā.

To address this important knowledge gap in the behavioral biology of this near-extinct

species, we conducted an investigation of ʻalalā responses to vocal signals. Specifically, we

examined whether ʻalalā still respond appropriately to species-specific calls, despite their

generations removed from nature, or whether their responses to conspecific call playbacks are

indicative of behavioral erosion in human care, a call for concern that portends poor social

integration and antipredator defense upon release. We presented birds with recorded playbacks of

alarm and aggressive territory intrusion calls, alongside relevant control calls and sounds, and

measured how likely birds were to approach the calling sound and to respond to it with the same

call type or a different call type (Table 1). We chose these two call types in the context of preparing

birds for survival alongside predators and the maintenance of social skills necessary for setting up

and defending territories from conspecifics. We also presented ʻalalā with two control calls to rule

out the effect of the social novelty and auditory novelty of hearing a call outside of their aviary.

Namely, we played a non-threatening ʻalalā territorial maintenance call, which birds routinely

make from their existing territories, and should therefore not broadcast any threat, and a call from

a novel species that ʻalalā have never heard. If the environment of the conservation breeding

facility has dulled birds' responses to auditory stimuli generally, we would expect that ʻalalā

produce no response to any of the call types (neither approaching nor making calls of their own),

suggesting that the birds are over-stimulated. In contrast, if birds have retained their natural

responses, we would expect to see differences between the recorded playbacks of aggressive

territory intrusion calls and control call types. Specifically, birds should respond to territorial

intrusion calls with their own aggressive territorial intrusion calls (Bradbury & Vehrencamp, 2011)
, and this effect should be most pronounced in males, due to the larger role males historically
played in territorial defense (Banko et al.,, 2002). For alarm calls, the natural response is more
complicated. We would expect birds to approach the source of the call to investigate the potential
source of danger, and may respond with alarm calls (Manser et al., 2002, Hill 1986). Meanwhile
the birds should ignore the general territorial maintenance call, since it does not denote a threat,
and may show signs of neophobia, or a hesitancy to approach, the novel call, since 'alalā are highly
neophobic in other contexts (Greggor et al., 2020).

Methods

We conducted the experiment on 28 breeding pairs, i.e. 62% of the entire 'alalā breeding
population, at the Keauhou Bird Conservation Center (KBCC; n = 24 pairs) and the Maui Bird
Conservation Center (MBCC; n = 4 pairs) between September and November 2018. The birds at
both facilities are currently several generations removed from the wild. Birds were tested in their
home enclosure with their breeding partner. Each pair was housed in an outdoor aviary, with
covered areas for shelter and feeding, and had daily ad libitum access to food and water.
Throughout the facility, interaction with people is minimized to reduce the effects of human care
(see Greggor et al., 2018). The birds have auditory access to other breeding pairs and most do not
have visual access. For the few buildings with more than one pair in adjacent aviaries we only
tested one pair per building to minimize subjects' prior exposure to the experimental setup. It is
possible that birds could see the experimental setup at adjacent aviaries, so we put as much time
as possible between trials at neighboring aviaries. Four pairs were physically separated from each
other for husbandry reasons but both had visual access to each other during all of their trials.

Experimental design of playback study

We designed the playback study to determine how 'alalā respond to alarm and other social
and non-social acoustic stimuli. We hypothesized that responses to different call types will vary
according to the context in which they are used and putative call meaning (Tanimoto et al., 2017a).
The experiment entailed playback of four distinct acoustic stimuli: (1) conspecific alarm call, (2)

conspecific territorial intrusion call, (3) conspecific non-threatening territorial maintenance call,
and (4) heterospecific call from a novel bird species (Table 1). Alarm calls are high-pitched calls
given to warn other birds of danger (Tanimoto et al., 2017a), which are typical of corvid vocal
repertoires (e.g. Bila et al., 2016, Marzluff et al., 2010, McIvor et al., 2018). By contrast, territorial
intrusion calls are given when birds are actively and aggressively defending their territory, in
situations that can escalate into physical aggression. Meanwhile, the non-threatening, territorial
maintenance calls that ʻalalā make routinely on the edges of their territories in the absence of any
direct threat or aggression were included as a conspecific control stimuli.

We employed a repeated-measures design in which each subject received playback of each
call type in a balanced, random order ensuring that each type of call was played first, second, third,
or fourth an equal number of times across all trials. Individual and temporal differences in
responsiveness to the experimental setup and acoustic playback can be a source of response
variation that adds statistical noise in playback studies. To address this possibility, we included a
pre-trial period before call playbacks in which subjects were exposed to ambient forest noise
playback (details below). The same forest noise was used as a post-playback stimulus for all trials,
so that conditions were constant for comparisons of observed behavior in pre- and post-call
playback periods. We used this comparison to evaluate behavioral changes in the aftermath of call
playbacks.

Call type	Experimental design function	Predicted behavioral responses
Alarm	Evaluate whether ʻalalā retain anti-predator responsiveness to conspecific calls indicating danger or predation threat	Vigilance, alarm calls, cautious approach (predicted to coordinate social anti-predator response)

Territorial Intrusion	Discriminate between response to an alarming social stimulus versus an alarming antipredator stimulus	Aggressive approach, territorial intrusion calling
Territorial maintenance	Control: non-alarming conspecific stimulus	No specific call response predicted, but 'alalā could show interest and possible social attraction
Novel Control	Control: non-alarming heterospecific stimulus	No call response predicted; perhaps hesitancy to approach, based on neophobia.

Table 1. Explanation of stimuli types and predicted responses.

Call recording and playback methods

We collected all audio recordings with a Roland R-05 acoustic recorder and directional boom
microphone. Conspecific calls and forest noise controls were recorded at the KBCC. We collected
calls opportunistically but tried to capture certain situations where we expected birds to make the
calls in question. For example, for territorial intrusion calls we recorded when males were being
moved between aviaries for husbandry purposes, as this is a time when we would expect these
territorial intrusion calls. To address pseudoreplication, one set of calls were collected from three
separate males, resulting in three unique sets of conspecific calls. A follow up analysis revealed
no difference in birds' responses between the three sets of calls (see Supplementary Materials). We
assigned 'alalā call types initially by social context and subjective perception, then confirmed
assignment by examining spectrograms. Calls from unfamiliar species were collected
opportunistically at the Panaewa Zoo in Hilo, Hawaii from medium- or large-bodied birds that
breeding center 'alalā would never have experienced (peacock, cockatoo, and toucan). Since

several species of corvids are able to distinguish characteristics of conspecifics from calls (Woods

et al, 2018), each pair received all three call exemplars from the same male to avoid individual

differences in donor birds influencing responses to call types. As breeding center birds are likely

able to hear most conspecifics housed at the same facility, all subjects were likely somewhat

familiar with individual calls from nearby conspecifics. However, to minimize familiarity with

calls we selected playback calls from individuals that were not housed in adjacent aviaries and did

not repeat call playback sets at adjacent aviaries.

We ensured that calls in the same category from different males had similar delta frequency

and average power using Raven Pro, version 1.5. In order to standardize the duration and number

of calls between call types with different durations and pauses, we ascertained that each 30-second

stimulus contained 6-7 seconds of sound and between 6-20 individual calls, ensuring the natural

spacing that the bird used between individual sounds to the extent possible. Therefore, we did not

edit the calls themselves but did edit the spacing between calls and the number of calls to keep the

total sound duration to 6-7 seconds for all kinds of stimuli. We measured sound duration and

conducted all audio editing in Audacity. We also checked that all sounds in the same category

were played from the speaker (omni jacket ultra, Altec Lansing) at the same minimum and

maximum decibel level using the Decibel X app for iphones.

All playback trials were conducted between 9:30am and 11:30am, when morning

husbandry-related disturbance had finished. Also, this time window is not within the species-

specific period of peak vocal activity that have been historically reported (0600 to 0900 and 1500

to 1800; Banko et al., 2002), to attempt to avoid times where there could be background noise

from other aviaries that might otherwise cause a distraction. For each playback trial we placed the

speaker on the ground in a standardized location on the side of the aviary, along an opaque wall.

While the birds may have known a person was outside the aviary, they were unable to directly see

the observer placing the speaker. We allowed the subjects a 5-min habituation period after the

observer placed the speaker and moved to the observation location. Each of four trials consisted

of three different observation periods: pre-trial (3-min playback of forest noise to control for

general response to sound playback), stimulus presentation (0.5-min call playback), and post-trial

(3-min forest noise playback repeated). Following a 19-min intermission, we repeated the three-

period trial using a different call type. This playback routine was repeated for all four calls in

balanced, randomized order across individuals. This resulted in 30 minutes between stimulus
presentations. The observer scored trials live through a small window at the front of the aviary and
recorded trials by setting up a video camera at each window. The birds likely heard the observer
enter, but once the 5-minute habituation began, the observer stayed quiet and out of view as much
as possible. Therefore, it is possible that the birds were aware of the presence of the observer and
video camera, but we tried to minimize disturbance as much as possible. We scored trials live but
consulted video recordings to improve data accuracy. We recorded the following behaviors across
all three trial periods: (1) approach (bird approaches playback speaker); (2) latency to approach
(time at which bird first moved in the direction of the stimulus); (3) number and (4) type of all
calls made by each bird. Call types were classified into four categories: alarm, territorial intrusion,
subordinate begging and other. Territorial maintenance calls were never made in response to the
stimuli so they were not included in the models.

Analysis

All analyses were conducted in R version 3.4.1 (R core team, 2017). In order to determine
if our pre-trial conditions were consistent for each stimulus, we ran a generalized linear mixed
model (GLMM) with a negative binomial distribution and a log link function, using the package
glmmTMB (version 0.2.3; Brooks et al., 2017), with the number of all calls made during the pre-
trial as the response variable. The initial model included the main effects of stimulus type and
order as well as the interactions between stimulus type and order. We also included the random
effect of bird ID.

To investigate the degree of individual consistency in call responses, we ran an intraclass
correlation using the package irr (version 0.84; Gamer et al., 2012) for the number of calls each
individual produced during the stimulus and post-trial combined for each stimulus type.

In order to determine interest in different stimuli, we ran a cox proportional hazards model
using the package survival (version 2.38; Therneau 2015) on the likeliness of birds to approach
the stimulus for each stimulus type. This model included stimulus type, sex, and order as main
effects as well as the interaction between stimulus type and sex and stimulus type and order. We
clustered data around bird ID.

To evaluate the circumstances influencing call production, we used generalized linear

mixed models (GLMM) with a negative binomial distribution (as data were zero inflated) and a

log link function, using the package glmmTMB (version 0.2.3; Brooks et al., 2017) to separately

test whether all calls, alarm calls, territorial intrusion calls and begging calls were more likely to

occur during the stimulus and post-trial period depending on the playback type. All models initially

included the main effects of stimulus type, sex and trial order as well as the interactions between

stimulus type and sex and stimulus type and order. We also included the random effect of bird ID.

We used the territorial maintenance control call as the reference group when comparing the

different stimulus types as this should represent a conspecific call in a new location, so this

accounts for any calls or approaches simply due to the social novelty and not the call itself. The

GLMM with the number of territorial intrusion calls made during the stimulus and post-trial would

not converge properly when the model included interactions, so only main effects were tested.

For all GLMMs except for the model with territorial intrusion calls as the response variable,

we first determined if the interaction terms warranted inclusion in the models. Starting with the

interaction between stimulus type and order, as this was likely less biologically relevant, and then

continuing with the interaction between stimulus type and sex, we removed interaction terms if

their inclusion failed to decrease AIC values by > 2.. We did not simplify the model past

determining which interactions to include, as all remaining effects were important variables we

wanted to consider in the final model. See the supplementary materials for the full process of

model selection for each response variable. For all GLMMs we also visually inspected binned

plots of the expected versus residual values for the final model.

Ethical note

This work using animal subjects was approved by San Diego Zoo Wildlife Alliance's

IACUC committee (No. 16-009). Several permits were issued for the conservation breeding of

'alalā (USFWS Native Endangered Species Recovery Permit TE060179-5, State of Hawaii

Protected Wildlife Permit WL19-16).

Results

Pre-trial call frequency in response to playback of forest noise

As expected, we found no differences in call number among treatments before stimulus
exposure during the pre-trial period (alarm trial vs. territorial maintenance: GLMM, b = -1.008, z
= -0.942, *P* = 0.346; novel control trial vs. territorial maintenance: b = -0.836, z = -0.804, *P* =
0.421; territorial intrusion trial vs. territorial maintenance: b = -0.677, z = -0.665, *P* = 0.506).
Therefore, we have not weighted post-trial data as a function of pre-trial calling rates. However,
bird calls significantly increased across trials (b = 0.769, z = 2.179, *P* = 0.029); therefore, we
included trial order in all subsequent models.

Approach latency during stimuli presentation

Birds were more likely to approach after the alarm stimulus than the territorial intrusion
stimulus (Figure 1B; alarm vs. territorial maintenance: Cox Proportional Hazards Model,
coefficient = 1.293, z = 2.293, Pr = 0.0218) whereas the other stimuli had no significant effect on
the likelihood of approaching compared to the territorial maintenance stimulus (Figure 1A; novel
control compared to territorial maintenance: coefficient = 0.827, z = 1.608, Pr = 0.108; territorial
intrusion versus territorial maintenance: coefficient = 0.479, z = 0.892, Pr = 0.372). Trial order
was also significant, with birds more likely to approach in later trials (coefficient = 0.242, z =
1.997, Pr = 0.0458). Sex did not have a significant effect on the likelihood of approaching (effect
of sex: coefficient = 0.115, z = 0.376, Pr = 0.707)

Individual call consistency during stimuli presentation and post-trial

Birds were individually consistent in the total number of calls they each made during the
stimulus and post-trial periods across all different trial treatments (Intraclass correlation
coefficient, 0.367, CI = 0.231-0.515, P < 0.001); i,e., birds who made fewer calls in one type of
trial also made fewer calls in all trials.

Call responses

Even though birds differed individually in the number of calls they made across all stimuli,
no patterns emerged in how many calls birds made between stimuli types during the playback and
post-trial period combined (Figure 2; alarm vs. territorial maintenance: GLMM, b = 0.497, z =

0.737, P = 0.461; novel control vs. territorial maintenance: b = 0.271, z = 0.408, P = 0.683;

territorial intrusion vs. territorial maintenance: b = 1.007, z = 1.471, P = 0.141). There was also no

effect of sex (b = -0.598, z = -1.211, P = 0.226) or order (b = 0.099, z = 0.394, P = 0.693) on the

total number of calls birds made during the stimulus and post-trial.

When we looked at the production of specific calls, we found that the number of alarm calls

birds made during the stimulus and post-trial combined did not significantly vary between any of

the different stimuli (Figure 3; alarm stimulus vs. territorial maintenance: b = -1.584, z = -1.093,

P = 0.275; novel control stimulus vs. territorial maintenance: b = 0.169, z = 0.097, P = 0.923;

territorial stimulus vs. territorial maintenance: b = 0.390, z = 0.312, P = 0.755). All other terms

were not significant. There was also no difference in the number of begging calls birds made during

the stimulus and post-trial between any of the stimuli (alarm stimulus vs. territorial maintenance:

b = 0.820, z = 0.665, P = 0.506; novel control vs. territorial maintenance: b = -0.193, z = -0.170,

P = 0.865; territorial intrusion stimulus vs. territorial maintenance: b = -0.441, z = -0.409, P =

0.683). Birds never made the territorial maintenance call in response to any stimuli.

Overall, birds made significantly more territorial intrusion calls during the stimulus and

post-trial in response to the territorial intrusion stimulus than the territorial maintenance stimulus

(Figure 4; b = 3.825, z = 2.298, P = 0.022) whereas they did not make a significantly different

amount of territorial calls during the stimulus and post-trial in response to any other kind of

stimulus (Figure 4; alarm stimulus vs. territorial maintenance: b = 2.033, z = 1.064, P = 0.287;

novel control vs. territorial maintenance: b = -19.797, z = -0.002, P = 0.999). Males made

significantly more territorial intrusion calls during the stimulus and post-trial than females (Figure

5; b = 5.087, z = 3.106, P = 0.002).

Discussion

Species raised in human care have the potential to lose some of their survival-relevant

behaviors (Shier, 2016, McPhee & Carlstead, 2010), including responses to species-specific

vocalizations (Corfield et al., 2008, Digby et al., 2013, Tanimoto et al., 2017b). Conservation

breeding programs or managed care may also change selective forces operating across generations

or developmental processes operating during individuals' lifetimes, creating new cultural vocal

variants and response patterns (Lewis et al., 2021). Disrupting the normal signaler-receiver

interplay may result in a breakdown in the communication system, with fitness consequences

challenging conservation goals. To address these possibilities, we tested whether pairs of 'alalā

have maintained responses to survival-relevant classes of vocalizations. Although the conservation

breeding history of the species may have dulled some of their responses, we still saw encouraging

signs that 'alalā were able to distinguish between natural call types and demonstrate survival-

appropriate responses. Specifically, we found that 'alalā were more likely to approach in response

to alarm calls, indicating they may have been investigating a potential threat or seeking to

coordinate a defensive response with conspecifics. Also, we found that males in particular were

more likely to respond to territorial intrusion calls with territorial intrusion calls of their own,

suggesting they were willing to defend their breeding aviary. Finally, our results illustrate that

there is some individual consistency in the amount of calling over all treatments, but that individual

characteristics (e.g. personality or specific rearing history) does not fully explain the variation in

call response to the different treatments. A note of caution in interpreting these results is warranted

due to the absence of any quantitative and limited qualitative observations of 'alalā vocal behavior

before they went extinct in the wild, so we have few reference points other than comparisons with

and generalizations from other species.

As a corvid, flexible learning is expected throughout their life (Emery & Clayton, 2004),

but vocalizations often require exposure during times of parental care or sensitive periods, which

may be limited in human care (Corfield et al., 2008, Digby et al 2013, Tanimoto et al., 2017b).

Most 'alalā in this study were puppet-reared by human caretakers (Gregor et al. 2018), removing

any opportunities for learning vocal behavior from parents. Of known call types, alarm calls are

more likely to be preserved since they potentially share some innate characteristics that make them

harder for predators to locate (Maynard Smith & Harper 2003). In the case of 'alalā, we found that

birds were more likely to approach alarm call playbacks than other auditory stimuli. However,

they were not more likely to make alarm calls in response to alarm playbacks. This lack of vocal

response is consistent with what was reported in some pilot work on 'alalā anti-predator behavior

(Greggor et al., in review), suggesting that alarm calls may function to alert birds to the need to

gather additional information. It is also possible that because we presented alarm calls without

other stimuli that could indicate danger, they investigated but did not respond with their own alarm

calls when they saw no clear threat. Further testing of alarm calls with and without presenting a

dangerous context would be necessary to tease this apart. However, 'alalā's approach of the source

of the playback alone may still have had a social function. Animals, including corvids, will often

make alarm calls to elicit assistance from conspecifics in investigating or confronting a predator

(Coomes et al., 2019, Curio 1978). The fact that birds approached the alarm calls (and not just

conspecific playbacks in general) suggests they may still have a similar function in 'alalā: approach

of the caller may precede mobbing or other group antipredator defense. Although mobbing-like

behavior has been anecdotally reported in 'alalā (Greggor et al., in review), this study provides

some empirical support that the birds responded appropriately to alarm calls in the absence of other

signs of danger and have not become completely desensitized to them.

Territorial calls are also an important part of vocal communication, including in the

historical vocal repertoire of 'alalā (Banko et al., 2002, Tanimoto et al., 2017a). We played two

types of territorial calls, a neutral territorial maintenance call and an aggressive territorial intrusion

call. 'Alalā did not significantly differ in their response to the territorial maintenance call from

other call types. However, we found that 'alalā responded to a simulated territorial intrusion with

territorial intrusion calls significantly more than they did to any other playback stimuli. A vocal

territorial response is the natural reaction we would expect to a territorial challenge (Maynard

Smith & Harper 2003; Bradbury & Vehrencamp 2011), considering we presented the birds with

this stimulus in their home aviaries. We did not test any of the birds in a new location outside of

their artificial "territory" so we do not know if their responses are flexible and adaptive to context,

i.e. defending only their occupied territory. However, given that 'alalā exhibit fewer territorial

calls than their wild counterparts (Tanimoto et al., 2017b), we find it encouraging that 'alalā

displayed this behavioral response suggesting some normal signaler-receiver exchange.

Additionally, we found that males were more likely to make territorial intrusion calls, which

mirrors patterns noted previously in the wild (Banko et al., 2002). As a sexually dimorphic species

with respect to size, males are the larger and more aggressive sex, and thus their greater role in

territorial defense is expected (Archer, 1988).

While we found encouraging signs that some birds exhibited natural responses to our

playbacks in the absence of any other context or stimuli, many of the birds did not respond vocally

when faced with any type of conspecific call. Given that the birds are living in social densities that

are far higher than observed in the wild (Flanagan et al., 2020), the high levels of individual

variation we found suggest that some 'alalā may have become desensitized to the calls of

conspecifics vocalizations, perhaps a result of repeated exposure without consequence. While

'alalā in the conservation breeding facilities may not necessarily be less vocal than wild birds

(Tanimoto et al., 2017b), the lack of vocal behavior we saw in many individuals in response to

social cues suggests a potential decoupling of the meaning or relevance of conspecific calls.

Alternatively, perhaps there is additional context we did not adequately capture in our recordings

(e.g. individual caller ID, social status, etc.), that could have differentially influenced some

individuals more than others. Corvids are known to respond to individual qualities of callers, such

as their dominance status (Massen et al., 2014), and membership in the breeding colony (Woods

et al., 2018). Additionally, since calls were recorded opportunistically, there may be subtle

differences to the different calls (for example, the particular stimulus causing the alarm calls) that

we are unaware of. Therefore, we may have inadvertently been broadcasting information beyond

the content of the calls. However, we saw no differences in the number of calls birds made in

response to the three different call sets, suggesting that the identity of the caller was not a major

cause of variation. Regardless of the cause of the low responses of many individuals, there were

still other birds that clearly demonstrated survival-relevant responses.

Although we found no evidence of widespread erosion of vocal communication behavior

under human care, the individual differences that we saw in how birds responded to the playbacks

could indicate that not all individuals are equally as well equipped for release into the wild. Many

factors go into determining fitness for release, and these results suggest that we may need to

consider whether birds demonstrate adequate responses to conspecific calls as a criterion for

release. Additionally, future research could investigate how likely birds are to regain survival-

relevant responses to vocalizations if they are exposed to training that encourages associations

between conspecific calls and relevant responses. Given the critical conservation status of the

species, any technique that could limit the impact of the conservation breeding environment on

survival-relevant behavior is warranted.

Acknowledgements

We greatly appreciate the wildlife care specialists at KBCC and MBCC who accommodated this
research and thank Patrick Hart for lending us recording equipment. We would also like to thank
members of the Catenazzi and Cox labs for their feedback on an earlier draft and to Ramona
Rauber for helpful discussion about 'alalā call types.

Data accessibility

Data and code will be available at an online repository upon acceptance.

Figures

Figure 1. A. Inverted survival curves showing the likelihood that birds approach each stimulus
type. B. Inverted survival curves showing the likelihood that birds approach in response to only
the two significantly different stimuli, territorial maintenance and alarm.

Figure 2.   Violin plots and raw data showing the distribution of the number of all calls birds
made during the stimulus and post-trial period after each type of stimulus. The points are the raw
data.

Figure 3.   Violin plots and raw data showing the distribution of the number of alarm calls birds
made during the stimulus and post-trial period after each type of stimulus. The points are the raw
data.

Figure 4. Violin plots and raw data showing the distribution of the number of territorial intrusion
calls birds made during the stimulus and post-trial period after each type of stimulus. The points
are the raw data.

Figure 5. Violin plots and raw data showing the distribution of the number of territorial intrusion
calls males and females made during the stimulus and post-trial period after each type of
stimulus. The points are the raw data.

References

Archer, J., 1988. The Behavioural Biology of Aggression. Cambridge University Press,

Cambridge.

Banko, P. C., Ball, D. L. & Banko, W. E. 2002. Hawaiian crow Corvus hawaiiensus. In: The

Birds of North America, Vol 648 (Ed. by A. Poole & F. Gill), Cornell Lab of

Ornithology.

Berger-Tal, O., Blumstein, D. T. & Swaisgood, R. R. 2020. Conservation translocations: a

review of common difficulties and promising directions. Animal Conservation, 23, 121–

131.

Bílá, K., Beránková, J., Veselý, P., Bugnyar, T., Schwab, C., 2017. Responses of urban crows to

con- and hetero-specific alarm calls in predator and non-predator zoo enclosures. Animal

Cognition 20, 43-51.

Bradbury, J. W., & Vehrencamp, S. L. (2011). Principles of animal communication (2nd

ed.). Sunderland MA: Sinauer

Brakes, P., Dall, S.R.X., Aplin, L.M., Bearhop, S., Carroll, E.L., Ciucci, P., Fishlock, V., Ford,

J.K.B., Garland, E.C., Keith, S.A., McGregor, P.K., Mesnick, S.L., Noad, M.J., di Sciara,

G.N., Robbins, M.M., Simmonds, M.P., Spina, F., Thornton, A., Wade, P.R., Whiting,

M.J., Williams, J., Rendell, L., Whitehead, H., Whiten, A., Rutz, C., 2019. Animal

cultures matter for conservation. Science 363, 1032-1034.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Maechler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and

467   Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The

468   R Journal, 9(2), 378-400.

469 Clayton, N.S. and Emery, N. J. 2007. The social life of corvids. Current Biology, 17, 652–656.

470 Coomes, J.R., Mcivor, G.E., Thornton, A., 2019. Evidence for individual discrimination and

471   numerical assessment in collective antipredator behaviour in wild jackdaws (Corvus

472   monedula). Biol. Lett. 15, 20190380.

473 Corfield, J., Gillman, L. and Parsons, S. 2008. Vocalisations of the North Island Brown Kiwi

474   (Apteryx mantelli). Auk, 125:326–335.

475 Culliney, S., Pejchar, L., Switzer, R., Ruiz-gutierrez, V., Culliney, S., Pejchar, L., Switzer, R. &

476   Ruiz-gutierrez, V. 2012. Seed dispersal by a captive corvid: the role of the 'Alalā (

477   Corvus hawaiiensis) in shaping Hawai'i's plant communities. Ecological Applications,

478   22, 1718–1732.

479 Curio, W. E. 1978. The adaptive significance of avian mobbing II: cultural transmission of

480   enemy recognition in blackbirds effectiveness and some constraints. Zeitschrift für Tier-

481   psychologie, 48, 184–202.

482 Decourcy, K. R. & Jenssen, T. A. 1994. Structure and use of male territorial headbob signals by

483   the lizard Anolis carolinensis. Animal Behaviour, 47, 251–262.

484 Digby, A., Bell, B.D. and Teal, P.D. 2013. Vocal cooperation between the sexes in Little Spotted

485   Kiwi Apteryx owenii. Ibis, 155:229–245.

486 Emery, N. J. & Clayton, N. S. 2004. The mentality of crows: convergent evolution of

487   intelligence in corvids and apes. Science, 306, 1903–7.

488 Fischer, J. & Lindenmayer, D. B. 2000. An assessment of the published results of animal

489   relocations. Biological Conservation, 96, 1–11.

Flanagan, A. M., Rutz, C., Farabaugh, S., Greggor, A. L., Masuda, B. & Swaisgood, R. R. 2020.

Inter-aviary distance and visual access influence conservation breeding outcomes in a

territorial, endangered bird. Biological Conservation, 242, 108429.

Gamer, M., Lemon, J., Fellows, I., & Singh, P. 2012. irr: Various Coefficients of Interrater

Reliability and Agreement. R package version 0.84. https://CRAN.R-

project.org/package=irr

Greenfield, M. D. & Minckley, R. L. 1993. Acoustic dueling in tarbush grasshoppers: settlement

of territorial contests via alternation of reliable signals. Ethology, 95, 309-326.

Greggor, A.L., Masuda, B., Gaudioso-Levita, J.M., Nelson, J., White, T.H., Shier, D.M.,

Farabaugh, S., Swaisgood, R.R. (in review). Pre-release training, predator interactions

and evidence for persistence of anti-predator behavior in reintroduced 'alalā, Hawaiian

crow. Global Ecology and Conservation.

Greggor, A.L., Masuda, B.M., Sabol, A., Swaisgood, R.R. (in review). Designing anti-predator

training to maximize learning and efficacy assessments. Biological Conservation.

Greggor, A. L., Vicino, G. A., Swaisgood, R. R., Fidgett, A., Brenner, D., Kinney, M. E.,

Farabaugh, S., Masuda, B. & Lamberski, N. 2018. Animal Welfare in Conservation

Breeding: Applications and Challenges. Frontiers in Veterinary Science, 5, 1–6.

Greggor A.L., Masuda B, Flanagan A.M., Swaisgood RR. 2020. Age-related patterns of

neophobia in an endangered island crow: implications for conservation and natural

history. Animal Behaviour, 2020, 160, 61–68. doi:10.1016/j.anbehav.2019.12.002

Griffin, A. S., Blumstein, D. T. & Evans, C. S. 2000. Training captive-bred or translocated

animals to avoid predators. Conservation Biology, 14, 1317–1326.

Hiebert, S. M., Stoddard, P. K. & Arcese, P. 1989. Repertoire size, territory acquisition and

reproductive success in the song sparrow. Animal Behaviour, 37, 266–273.

Hill, G.E., 1986. The function of distress calls given by tufted titmice (Parus bicolor): an

experimental approach. Animal Behaviour 34, 590-598.

Hollén, L.I., Radford, A.N., 2009. The development of alarm call behaviour in mammals and

birds. Animal Behaviour 78, 791-800.Lewis, R.N., Williams, L.J., Gilman, R.T., 2021.

The uses and implications of avian vocalizations for conservation planning. Conservation

Biology 35, 50-63.

Mager, J. N., Walcott, C. & Piper, W. H. 2012. Male common loons signal greater aggressive

motivation by lengthening territorial yodels. Wilson Journal of Ornithology, 124, 73–80.

Manser, M.B., Seyfarth, R.M., Cheney, D.L., 2002. Suricate alarm calls signal predator class and

urgency. Trends in Cognitive Sciences 6, 55-57.

Massen, J. J. M., Pašukonis, A., Schmidt, J. & Bugnyar, T. 2014. Ravens notice dominance

reversals among conspecifics within and outside their social group. Nature

communications, 5, 3679.

Maynard Smith, J. & Harper, D. 2003. Animal signals. Oxford: Oxford University Press.

McIvor, G.E., Lee, V.E., Thornton, A., 2018. Testing social learning of anti-predator responses

in juvenile jackdaws: the importance of accounting for levels of agitation. Royal Society

open science 5, 171571.

McPhee, E.M., Silverman, E.D., 2004a. Increased behavioral variation and the calculation of

release numbers for reintroduction programs. Conservation Biology 18, 705-715.

McPhee, M.E., 2004b. Generations in captivity increases behavioral variance: considerations for

captive breeding and reintroduction programs. Biological Conservation 115, 71-77.

McPhee, M.E., Carlstead, K., 2010. The importance of maintaining natural behaviors in captive

mammals, in: Kleiman, D., Thompson, K. (Eds.), Wild Mammals in Captivity: Principles

and Techniques for Zoo Management. pp. 303–213

Moseby, K. E., Read, J., Paton, D., Copely, P., Hill, B. & Crisp, H. 2011. Predation determines

the outcome of 10 reintroduction attempts in arid South Australia. Biological

Conservation, 144, 2863–2872.

Muralidhar, A., Moore, F.L., Easton, L.J., Jamieson, I.G., Seddon, P.J., van Heezik, Y., 2019.

Know your enemy? Conservation management causes loss of antipredator behaviour to

novel predators in New Zealand robins. Animal Behaviour 149, 135-142.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rabin, L.A., 2003. Maintaining behavioral diversity in captivity for conservation: natural

behaviour management. Animal Welfare 12, 85-94.

Shier, D. M. 2016. Manipulating animal behavior to ensure reintroduction success. In:

Conservation Behavior: Applying Behavioral Ecology to Wildlife Conservation and

Management, (Ed. by O. Berger-Tal & D. Saltz), pp. 275–304. Cambridge: Cambridge

University Press.

Tanimoto, A. M., Hart, P. J., Pack, A. A. & Switzer, R. 2017a. Vocal repertoire and signal

characteristics of 'Alalā, the Hawaiian Crow (Corvus hawaiiensis). The Wilson Journal

of Ornithology, 129, 25–35.

Tanimoto, A. M., Hart, P. J., Pack, A. A., Switzer, R., Banko, P. C., Ball, D. L., Sebastián-

González, E., Komarczyk, L. & Warrington, M. H. 2017b. Changes in vocal repertoire of

the Hawaiian crow, Corvus hawaiiensis, from past wild to current captive populations.

Animal Behaviour, 123, 427–432.

Therneau, T. (2015). A Package for Survival Analysis in S. version 2.38,

<URL:https://CRAN.R-project.org/package=survival>.

US Fish and Wildlife Service. 2009. Revised Recovery Plan for the `Alalā (Corvus hawaiiensis).

Woods, R. D., Kings, M., McIvor, G. E. & Thornton, A. 2018. Caller characteristics influence

recruitment to collective anti- predator events in jackdaws. Scientific Reports, 7343.