1 **Title:** The *Phtheirospermum japonicum* isopentenyltransferase PjIPT1a regulates host
2 cytokinin responses in *Arabidopsis*
3

4 **Authors:** Anne Greifenhagen[1], Isabell Braunstein[1], Jens Pfannstiel[2], Satoko Yoshida[3], Ken
5 Shirasu[4,5], Andreas Schaller[1], Thomas Spallek[1*]
6

Affiliations: [1]Department of Plant Physiology and Biochemistry, University of Hohenheim,
Stuttgart, 70599, Germany
[2]Core Facility Hohenheim, Mass Spectrometry Unit, University of Hohenheim, Stuttgart,
70599, Germany
[3]Division of Biological Science, Graduate School of Science and Technology, Nara Institute of
Science and Technology, Ikoma, Nara, 630-0192, Japan
[4]RIKEN Center for Sustainable Resource Science, Yokohama, 230-0045, Japan
[5]Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan
15

16 **Corresponding Author:** Thomas Spallek, address as above, Phone: +49 711-459-22200,
17 email: thomas.spallek@uni-hohenheim.de
18

ORCID ID:
Anne Greifenhagen (Orcid ID: 0000-0003-2979-3203)
Isabell Braunstein (N/A)
Jens Pfannstiel (Orcid ID: 0000-0002-7651-4302)
Satoko Yoshida (Orcid ID: 0000-0002-9999-7861)
Ken Shirasu (Orcid ID: 0000-0002-0349-3870)
Andreas Schaller (Orcid ID: 0000-0001-6872-9576)
Thomas Spallek (Orcid ID: 0000-0003-4425-1015)

27
28
29

30 Total word count: 2989
31 4 main figures (all colour), 4 supplemental figures, 3 supplemental tables
32

Summary

The hemiparasitic plant *Phtheirospermum japonicum* is a nutritional specialist that supplements its nutrient requirements by parasitizing other plants through haustoria. During parasitism, the *Phtheirospermum* haustorium transfers hypertrophy-inducing cytokinins (CKs) to the infected host root. The CK biosynthesis genes required for haustorium-derived CKs and the induction of hypertrophy are still unknown. We searched for haustorium-expressed isopentenyltransferases (IPTs) that catalyse the first step of CK biosynthesis, confirmed the specific expression by *in vivo* imaging of a promoter-reporter, and further analysed the subcellular localization, the enzymatic function, and contribution to inducing hypertrophy by studying CRISPR-Cas9 induced *Phtheirospermum* mutants. *PjIPT1a* was specifically expressed in intrusive cells of the haustorium close to the host vasculature. PjIPT1a and its closest homolog PjIPT1b located to the cytosol and showed isopentenyltransferases activity *in vitro* with differences in substrate specificity. Mutating *PjIPT1a* abolished parasite-induced CK responses in the host. A homolog of *PjIPT1a* with shared characteristics was also identified in the related weed *Striga hermonthica*. With *PjIPT1a* we identified a *bona fide* parasitism gene. We propose that *PjIPT1a* exemplifies how parasitism-related functions evolve through gene duplications and neofunctionalization.

Keywords

Cytokinin, parasite, haustorium, intrusive cells, hypertrophy, *Striga*, neofunctionalization

Introduction

Most parasitic species maintain some level of photosynthesis (hemiparasites), but access soil

nutrients mainly by parasitizing other plants (Heide-Jorgensen, 2008; Brundrett & Tedersoo,

2018). The two main lineages of hemiparasitic plants, Santalales and the family of

Orobanchaceae in the Lamiales account for more than 90% of all parasitic species (Nickrent,

2020). The greatest in-family diversity of hemiparasitic plants with varying degrees of host-

dependency is found in the Orobanchaceae. In contrast to the agriculturally important weed

Striga hermonthica, non-weedy species such as the model parasitic plant *Phtheirospermum*

japonicum can propagate without being attached to a host (Clarke *et al.*, 2019).

Genetic adaptations to the parasitic lifestyle start to be revealed from the sequencing of

transcriptomes and genomes of parasitic plants (Yang *et al.*, 2015; Vogel *et al.*, 2018; Sun *et*

al., 2018; Yoshida *et al.*, 2019). At the centre of these studies is the haustorium, a multi-cellular

organ that facilitates infection and parasitism (Yoshida *et al.*, 2016; Kokla & Melnyk, 2018).

Comparative transcriptomics in three Orobanchaceae species with varying degrees of host

dependencies unearthed remarkable conservation of the transcriptional signatures during

infection (Yang *et al.*, 2015). Further studies confirmed these tendencies also in other lineages

(Vogel *et al.*, 2018; Sun *et al.*, 2018; Yoshida *et al.*, 2019). The term "core parasitism genes"

describes the shared transcriptional behaviour of a group of homologous genes within a

parasitic plant lineage (Yang *et al.*, 2015).

The concept of core parasitism genes may be extended to "core parasitism strategies".

Strategies that are commonly observed during plant parasitism include the secretion of cell wall

degrading enzymes and changes in hormone homeostasis (Brun *et al.*, 2020). For example,

cytokinins are produced in haustoria of *Cuscuta japonicum*, *Santalum album* and

Phtheirospermum japonicum (Zhang *et al.*, 2012; Furuhashi *et al.*, 2013; Spallek *et al.*, 2017).

Transfer of cytokinin from the parasite to the host was demonstrated using *Arabidopsis*

cytokinin biosynthesis mutants and transgenic *Phtheirospermum* roots that express a cytokinin-

degrading enzyme (Spallek *et al.*, 2017). With prolonged cytokinin transfer, infected host tissue

becomes hypertrophic, a symptom that is also induced by mistletoes, *Viscum album*

(Loranthaceae in the order Santalales), and *Alectra vogelii* (Orobanchaceae) (Heide-Jorgensen,

2008).

Cytokinins are N^6-substituted adenine derivatives, of which isoprenoid cytokinins are most

prevalent in plants (Sakakibara, 2006). N^6-prenylation is catalysed by adenosine phosphate-

isopentenyltransferases (IPTs). Plant IPTs prefer dimethylallyl diphosphate (DMAPP) as

isoprenoid donor, and ATP and ADP (ATP/ADP-type IPTs) or tRNA (tRNA-type IPTs) as

isoprenoid acceptors (Kakimoto, 2001; Sakakibara, 2006). The agrobacterial IPTs, Tmr and

Tzs, prefer hydroxy-methylbutenyl diphosphate and AMP as substrates. Tmr and Tzs are

required for virulence and crown-gall formation (Akiyoshi *et al.*, 1984; Heinemeyer *et al.*,

1987). Overexpression of IPTs in *Arabidopsis* causes ectopic cytokinin responses (Kakimoto,

2001) indicating that IPTs catalyse the rate limiting step in cytokinin biosynthesis. In

Arabidopsis, the major route of cytokinin biosynthesis relies on ATP/ADP-type IPTs targeted

to plastids (AtIPT1, AtIPT4, AtIPT5, AtIPT8), to mitochondria (AtIPT7), or residing in the

cytosol and nucleus (AtIPT3, AtIPT4). tRNA-type IPTs (AtIPT2, AtIPT9) also localize to the

cytosol and nucleus (Kasahara *et al.*, 2004; Kieber & Schaller, 2014). The biological relevance

of different subcellular IPT locations is not known.

Here, we report the function of *Phtheirospermum PjIPT1a* as a *bona fide* core parasitism gene.

PjIPT1a is an unconventional plant isopentenyltransferase with parasitism-related features.

PjIPT1a uses AMP and ATP as prenyl acceptors and is located in the cytosol. It is expressed

in intrusive cells of the *Phtheirospermum* haustorium. Targeted gene-knockouts of *PjIPT1a* by

CRISPR/Cas9 suppressed parasite-induced cytokinin responses in the host. Similarities to

isopentenyltransferases from *Striga* suggest that hemiparasitic Orobanchaceae use common

strategies to manipulate host plants.

Materials and Methods

Plant growth conditions and transformation

Phtheirospermum japonicum (Okayama, wild-type) seeds were sterilized in 70% ethanol,

0.05% SDS for 10 minutes, and then washed with 98% ethanol. For transformation, plants were

grown on Gamborg B5 medium (Duchefa) supplemented with 1% sucrose at 25°C under long-

day conditions (16 h light). Seven-days-old *Phtheirospermum* seedlings were transformed with

Agrobacterium rhizogenes as previously described (Ishida *et al.*, 2011). Approximately four

weeks after transformation, *Phtheirospermum* seedlings with hairy roots were transferred to

0.8 % agar containing 300 μg/ml cefotaxime. Two days later, *Arabidopsis* Ws *pARR5:GFP*

seedlings grown for one week on MS (Duchefa) plates were aligned next to *Phtheirospermum*

hairy roots. One week later, infected plants were transferred to half-strength MS plates and

grown for additional 21 days. *Nicotiana benthamiana* plants were grown in soil under long-day

conditions at 25°C.

Phylogenetic analysis

IPT amino acid sequences were retrieved from public databases (NCBI), and by blast search

using IPT protein sequences as query against published genomes (Table **S1**). CLC Main

Workbench Version 8 was used to align protein sequences and to construct a phylogenetic tree

based on the UPGMA and Jake-Cantor algorithm with a bootstrap analysis of 1000 replicates.

Cloning, plasmid design and genotyping

We cloned the 2.6 kb upstream sequence of the *PjIPT1a* gene while removing an internal

Bpi1site into a GoldenGate vector that was then used to generate 3xVenus-NLS-based reporter

as previously described (Engler *et al.*, 2014; Cui *et al.*, 2016). Primer sequences are given in

Table **S2**. Open reading frames (ORFs) of *PjIPT1a* and *PjIPT1b* were first cloned into

pENTR/D-TOPO entry vector (Thermo Fisher Scientific) to generate GFP fusion constructs

using the pGWB5 binary vector (Nakagawa *et al.*, 2007). The putative chloroplast transit

peptide was cloned into pART7 using XhoI and ClaI (Thermo Fisher Scientific) and then into

pART27 (Gleave, 1992). An empty pGREEN vector was used to express *p35S:GFP* (Hellens

et al., 2000). Goldengate-compatible vectors carrying *PjIPT1a*-specific gRNAs, *Cas9*, and the

transformation marker *p35S:DsRED* were assembled as previously described (Nekrasov *et al.*,

2013; Engler *et al.*, 2014). gRNAs were chosen based on their specificity to the intended target.

Individual hairy roots were genotyped by amplifying a 1651 bp long amplicon that contained

the 954 bp *PjIPT1a* gene. *PjIPT1a* alleles were cloned into pBluescript-II-SK(+) vectors for

sequencing. For expression in *E. coli*, ORFs of *PjIPT1a*, *PjIPT1b*, and *GST* were fused with a

C-terminal octa-His tag in the pET42 vector (Novagen).

Expression of proteins in *N. benthamiana* and *E. coli*

Agrobacterium tumefaciens GV3101 were infiltrated in a 10 mM MES (pH 5.6) and 10 mM

$MgCl_2$ buffer into approximately five-week-old *N. benthamiana* leaves, which were imaged

two days later. *PjIPT1a-His*, *PjIPT1b-His*, and *GST-His* in *E. coli* were expressed in BL21 RIL

cells and purified on NiNTA beads (Qiagen), then concentrated and desalted using Vivaspin

500 columns (Sartorius).

Microscopy

Hairy roots were imaged in water using a Zeiss Axio Image Z.1 fluorescence microscope.

PjIPT1a reporter activity was recorded with a Leica TCS SP5 II, the subcellular location of

fluorescent proteins with a Zeiss LSM 700 confocal microscope.

IPT-*in vitro* assays

IPT-*in vitro* assays were conducted as previously described (Brugière *et al.*, 2008). In brief, 8

µg of purified proteins were incubated with 1 mM DMAPP (Sigma) and 1 mM AMP or ATP

for two hours at 30°C in 12.5 mM Tris–HCl (pH 7.5), 37.5 mM KCl, 5 mM $MgCl_2$. The reaction

was stopped at 95°C and methanol precipitated. The supernatant was analysed by UHPLC-

MS/MS on a 1290 UHPLC system (Agilent) coupled to a Q-Exactive Plus Orbitrap mass

spectrometer equipped with a heated electrospray ionization (HESI) source (Thermo Fisher

Scientific). Analyte separation was achieved by a Phenomenex Luna Omega Polar C18 column.

Mobile phase A was 0.2% formic acid in water, and mobile phase B 0.2% formic acid in

acetonitrile. The HESI source was operated in positive ion mode, with a capillary voltage of

4.2 kV and an ion transfer capillary temperature of 360 °C. Mass spectra were acquired in MS

mode within the mass range of 150 to 800 m/z at a resolution of 70,000 FWHM using an

Automatic Gain Control (AGC) target of 3×106 of and 100 ms maximum ion injection time

(MIT). Data dependent MS/MS spectra in a mass range of 200 to 2000 m/z were generated for

the five most abundant precursor ions with a resolution of 17,500 FWHM using an AGC target

of 5×104, 64 ms MIT and a stepped collision energy of 15, 30 and 55. Xcalibur software

version 4.4.16.14 (Thermo Fisher Scientific) was used for instrument control and data

acquisition. Compound Discoverer software 3.2 (Thermo Fischer Scientific) was used for

analysing raw instrument data (.raw files) and peak area calculation.

Statistical analysis

The R packages Rmisc, readr, multcompView, ggplot2 and ggpubr were used for statistics (R

Core Team, 2020).

Results

PjIPT1a is expressed in intrusive cells of *Phtheirospermum* haustoria

CK-biosynthesis is tightly regulated by environmental and developmental cues and restricted

to distinct tissues and cells. We thus reasoned that CK production in *Phtheirospermum*

haustoria requires the haustorium-specific expression of at least one *IPT* gene (Fig. **1a**). We

surveyed published *Phtheirospermum* transcriptomes for infection-specific expression of *IPTs*

and identified one, hereinafter referred to as *PjIPT1a*, that is upregulated in the course of

infection and shows 25-fold higher expression in intrusive cells relative to the rest of the

haustorium (Fig. **1b**)(Ogawa *et al.*, 2020)(Kurotani *et al.*, 2020). *PjIPT1a* expression increases

until day 3 of the infection and continues at a relatively high level for the later time points (Fig.

1c). This expression profile coincides with the induction of cytokinin response markers in

parasite and host, which become detectable from 3 - 4 dpi (days post-infection) and remain

visible until at least 30 dpi (Spallek *et al.*, 2017). Using a 2.6 kb *PjIPT1a* promoter fragment to

drive the expression of a *3xVenus-NLS* reporter, we confirmed the specific expression of the

PjIPT1a gene in intrusive cells of *Phtheirospermum* haustoria at 4 dpi of *Arabidopsis* roots (Fig.

1d-f). *PjIPT1a* is thus expressed in cells in close, likely direct, contact to the host vasculature

(Fig. **1g-i**).

PjIPT1a encodes a cytosolic IPT with homologs in *Striga hermonthica*

To validate the phylogenetic relationship of PjIPT1a to other *Phtheirospermum* IPTs, we

compared the amino acid sequences of PjIPTs to known plant IPTs and two agrobacterial IPTs,

Tzs and Tmr (Fig. **2a**). PjIPT1a clustered with other plant IPT1s, and was more similar to

proteins from the related parasitic species *Striga hermonthica* (Orobanchaceae, Lamiales) and

Streptocarpus rexii (Gesneriaceae, Lamiales) than to IPTs from more distantly related non-

parasitic species such as common hop (*Humulus lupulus*, Cannabaceae, Rosales) and

Arabidopsis thaliana (Brassicaceae, Brassicales). Two close homologs were detected for

PjIPT1a and its *Striga* ortholog ShIPT1a. Like *PjIPT1a*, only *ShIPT1a* is induced upon

infections but not *ShIPT1b* (Fig. **S1**).

Bioinformatics tools predict a chloroplast transit peptide (CTP) in the first 22 amino acids of

PjIPT1a; however, at a lower score compared to AtIPT1 and SrIPT1 (Table **S3**). Putative CTPs

of PjIPT1a and ShIPT1 are less than half as long as the CTPs predicted for AtIPT1 (71 amino

acids) and SrIPT1 (46 amino acids) (Table **S3**; Fig. **S2**). Therefore, we experimentally tested

subcellular localization of PjIPT1a and functionality of its CTP. A PjIPT1a-GFP fusion protein

transiently expressed in *N. benthamiana* located to cytosol and nucleus (Fig. **2a**), with no apparent difference to the localization of free GFP (Fig. **2e**). Western-blots using anti-GFP antibodies confirmed the integrity of the PjIPT1a-GFP full-length protein (Fig. **S3**). The data thus suggest nuclear-cytoplasmic localization of PjIPT1a.

Nuclear-cytoplasmic localization is also observed for chloroplast-targeted AtIPT3, when it is post-translationally modified by C-terminal farnesylation (Galichet *et al.*, 2008). To exclude the possibility that post-translational modifications of PjIPT1a outside the putative CTP would affect its localization, we analysed the subcellular targeting of super-folder GFP fused to the predicted transit peptide (the first 22 amino acids) (Fig. **2b**). The localization of PjIPT1a(1-22)-sfGFP was indistinguishable from PjIPT1a-GFP and free GFP (Fig. **1a,e**). The same nuclear-cytoplasmic localization was also observed for PjIPT1b-GFP (Fig. **2d**). The putative CTP signal in *Striga* ShIPT1a is even shorter (11 amino acids) than the apparently non-functional CTP of PjIPT1a (Fig. **S2**; Table **S3**), while no CTP was predicted for ShIPT1b. Therefore, we assume a similar subcellular localization pattern of the homologous ShIPT1a and ShIPT1b proteins.

PjIPT1a drives CKs responses in host plants

Haustorium-derived CKs induce hypertrophy of the host root and can be visualized by a host-expressed cytokinin reporter (*pARR5:GFP*)(Spallek *et al.*, 2017). To address a role for *PjIPT1a* in the production of haustorium-derived CKs, we generated transgenic *Phtheirospermum* roots (hairy root, HR) that expressed the *Cas9* gene together with two *PjIPTa*-specific gRNAs (Fig. **3b**). The construct also contained *p35S:dsRED* as a fluorescent marker for identification of transformed roots. HR transformed with *p35S:dsRED* alone served as controls. In contrast to control HRs that triggered *pARR5:GFP* expression and hypertrophic growth in infected *Arabidopsis* roots, neither *pARR5:GFP* expression nor hypertrophic growth were observed after infection with HRs containing the *Cas9/PjIPT1a-gRNA*s cassette (Fig. **3a**). While the *PjIPT1a* locus was successfully PCR-amplified from control HR (Fig. **3c**), smaller bands were observed for HRs containing the *Cas9* gene (Fig. **3c**) resulting from Cas9-induced deletions ranging from single-nucleotides to 1100 bp (Fig. **3d**). Despite low abundance, we also identified one additional allele that carried a 40 bp deletion in the *PjIPT1a* gene for the hairy root HR3. With the more abundant alleles with larger deletions (893 bp and 1110 bp), this third allele revealed a chimeric pattern of *Cas9*-induced mutations in *Phtheirospermum* hairy roots. The presence of Cas9-induced deletions in *PjIPT1a* correlated perfectly with the absence of a

247 *Phtheirospermum*-induced cytokinin response in *Arabidopsis*. We therefore conclude that
248 *PjIPT1a* is an essential component in the CK transfer from parasite to host.
249

PjIPT1a and PjIPT1b differ in isopentenyltransferase activity
PjIPT1a and ShIPT1a share a unique triple-proline motif close to the nucleotide-binding site
that we only found in IPTa isoforms of parasitic Orobanchaceae species (Fig. **S4**). This raised
questions regarding their enzymatic activity. We purified His-tagged PjIPT1a and PjIPT1b
from *E. coli* and assayed for isoprenylation of AMP and ATP with DMAPP as the isoprenyl
donor and His-tagged GST as negative control (Fig. **4**). When AMP was used as substrate, the
total amount of isoprenylated products were similar (factor of 1.3) for PjIPT1a-His and PjIPTb-
His (Fig. **4a**). However, with ATP as substrate, the total amount of isoprenylated products was
about five times higher for PjIPT1b-His compared to PjIPT1a-His, indicating higher selectivity
for ATP over AMP (Fig. **4b**).

Discussion

Here, we studied parasite-induced CK responses in host plants as an example of a

neofunctionalized gene function in the Orobanchaceae species *Phtheirospermum japonicum*

and *Striga hermonthica*. We found one IPT gene, *PjIPT1a*, to be expressed in *Phtheirospermum*

haustoria. Reporter gene analysis revealed spatially restricted activity of the *PjIPT1a* promotor

in intrusive cells that are in contact with the host vasculature (Fig. **1g-i**). The proximity to the

host vasculature is likely sufficient to transfer cytokinins to the host and to trigger CK responses

above the haustorium attachment site (Spíchal, 2012). The parasitic tree *Santalum album* also

expresses an IPT1 homolog at a similar stage of infections; however, the exact location of

SaIPT1 expressing cells is not known (Zhang *et al.*, 2015).

Unlike in *Santalum album* and many other parasitic plants that are not amenable to reverse

genetic manipulation, we were able to use CRISPR/Cas9-based loss-of-function analysis to

demonstrate that *PjIPT1a* adopted new parasitism-related functions in *Phtheirospermum*.

While RNAi-based knock-down approaches have largely been replaced by the CRISPR/Cas9

technology, RNAi remained, until now, the only strategy to study parasitic plant genes

(Bandaranayake *et al.*, 2010; Alakonya *et al.*, 2012; Wakatake *et al.*, 2020). Here, we report the

first successful application of the CRISPR/Cas9 technology for studying loss-of-function

phenotypes in parasitic plants. CRISPR/Cas9 efficiently introduced deletions in *PjIPT1a*

resulting in the loss of up to 98% of the coding sequence, or in the disruption of its open reading

frame. In individual hairy roots, we found more than one, in one instance more than two,

mutated alleles of *PjIPT1a*, a result of the occasionally chimeric nature of *Agrobacterium*

rhizogenes-induced hairy roots (Wakatake *et al.*, 2018). All *Phtheirospermum* roots with non-

functional *PjIPT1a* genes failed to induce host CK responses (Fig. **3b,c**), showing that *PjIPT1a*

is indeed a *bona fide* parasitism gene with a specific parasitism-related physiological function.

As previously proposed for other parasitism genes in the Orobanchaceae, our phylogenetic

analysis of *IPTs* showed that the duplication event producing *PjIPT1a/ShIPT1a* and

PjIPT1b/ShIPT1b occurred in a common non-parasitic ancestor of *P. japonicum* and *S.*

hermonthica (Yang *et al.*, 2015). Surprisingly, PjIPT1a and ShIPT1a, which adopted a new

function as parasitism genes, are more similar to SrIPT1 from *Streptocarpus rexii* than PjIPT1b

and ShIPT1b (Fig. **2a**). The high similarity between parasitism-related and conventional IPTs

suggests that neofunctionalization of *PjIPT1a* mainly occurred at the level of its expression

profile. Enzymatic constraints may have limited divergence of the gene body and the enzymatic

functions of PjIPT1a and b are indeed comparable. However, preference for ATP was reduced in PjIPT1a; its preference for the energetically less costly AMP as isopentenyl donor may reflect an adaptation to the resource-limited parasitic lifestyle that is induced under nutrient-deficient conditions (Spallek *et al.*, 2017). A notable exception to the similarity of PjIPT1a and ShIPT1a to SrIPT1 is a characteristic triple-proline-motif that is only found in parasitism-related IPTs (Fig. **S4**).

Another unexpected feature of PjIPT1a is its nuclear-cytoplasmic localization (Fig. **2b,c**), which it shares with PjIPT1b (Fig. **2d**) and likely also with ShIPT1a and ShIPT1b (Table **S1**). *Arabidopsis* AtIPT1 and AtIPT3 feature functional N-terminal CTPs (Kasahara *et al.*, 2004). However, while AtIPT1 is located in plastids, AtIPT3 is dually targeted, either to the chloroplast, or to the nucleus/cytoplasm in a farnesylation-dependent manner (Galichet *et al.*, 2008). We can exclude a similar mechanism for PjIPT1a, as its predicted CTP appeared to be non-functional, as it was insufficient for targeting GFP to plastids (Fig. **2c**). The relevance of the nuclear-cytoplasmic localization is not clear, but given that it is shared between IPTs that are up-regulated during infection and other that are not, a functional link to parasitism seems unlikely.

In conclusion, we identified *PjIPT1a* as a *bona fide* parasitism gene that originated from a non-parasitic ancestral gene through gene duplication followed by neofunctionalization. Targeted knock-out of *PjIPT1a* showed its essential role for *Phtheirospermum*-induced CK-responses in the host. PjIPT1a has homologs in other parasitic plants. Targeting of neofunctionalized parasitism genes may allow for specific control of parasitic weeds leaving host plants unharmed.

Acknowledgements

This work supported by the German Research Foundation (DFG, 424122841) to T.S and MEXT

KAKENHI JP20H05909 to S.Y. and K.S. The authors thank Ursula Glück-Behrens and Ute

Bertsche for technical support.

Author Contribution

T.S., A.S., and K.S supervised the research. A.G. and I.B. analysed the subcellular location of

proteins, purified IPTs from *E. coli* and conducted the *in vitro* assays. J.P. measured the IPT *in*

vitro assays compounds by HPLC-MS/MS. T.S. performed the remaining experiments. A.S.,

P.S., and T.S. analysed data. Y.S. and K.S. provided crucial non-published data. T.S. and A.S.

wrote the paper.

References

Akiyoshi DE, Klee H, Amasino RM, Nester EW, Gordon MP. 1984. T-DNA of

Agrobacterium tumefaciens encodes an enzyme of cytokinin biosynthesis. *Proceedings of the*

National Academy of Sciences, USA **81**: 5994–5998.

Alakonya A, Kumar R, Koenig D, Kimura S, Townsley B, Runo S, Garces HM, Kang J,

Yanez A, David-Schwartz R, *et al.* 2012. Interspecific RNA interference of *SHOOT*

MERISTEMLESS-like disrupts *Cuscuta pentagona* plant parasitism. *Plant Cell* **24:** 3153–3166.

Bandaranayake PCG, Filappova T, Tomilov A, Tomilova NB, Jamison-McClung D, Ngo

Q, Inoue K, Yoder JI. 2010. A single-electron reducing quinone oxidoreductase is necessary

to induce haustorium development in the root parasitic plant *Triphysaria*. *Plant Cell* **22:** 1404–

1419.

Bardgett RD, Smith RS, Shiel RS, Peacock S, Simkin JM, Quirk H, Hobbs PJ. 2006.

Parasitic plants indirectly regulate below-ground properties in grassland ecosystems. *Nature*

439: 969–972.

Brugière N, Humbert S, Rizzo N, Bohn J, Habben JE. 2008. A member of the maize

isopentenyl transferase gene family, *Zea mays isopentenyl transferase 2* (*ZmIPT2*), encodes a

cytokinin biosynthetic enzyme expressed during kernel development. *Plant Molecular Biology*

67: 215–229.

Brun G, Spallek T, Simier P, Delavault P. 2020. Molecular actors of seed germination and

haustoriogenesis in parasitic weeds. *Plant Physiology* **185:** 1270-1281.

Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and global
host plant diversity. *New Phytologist* **220:** 1108–1115.

Clarke CR, Timko MP, Yoder JI, Axtell MJ, Westwood JH. 2019. Molecular dialog
between parasitic plants and their hosts. *Annual Review of Phytopathology* **57:** 279–299.

Cui S, Wakatake T, Hashimoto K, Saucet SB, Toyooka K, Yoshida S, Shirasu K. 2016.
Haustorial hairs are specialized root hairs that support parasitism in the facultative parasitic
plant *Phtheirospermum japonicum. Plant Physiology* **170:** 1492–1503.

Engler C, Youles M, Gruetzner R, Ehnert TM, Werner S, Jones JDG, Patron NJ,
Marillonnet S. 2014. A golden gate modular cloning toolbox for plants. *ACS Synthetic Biology*
3: 839–843.

Furuhashi T, Kojima M, Sakakibara H, Fukushima A, Hirai MY, Furuhashi K. 2013.
Morphological and plant hormonal changes during parasitization by *Cuscuta japonica* on
Momordica charantia. Journal of Plant Interactions **9:** 220–232.

Galichet A, Hoyerová K, Kamínek M, Gruissem W. 2008. Farnesylation directs AtIPT3
subcellular localization and modulates cytokinin biosynthesis in *Arabidopsis. Plant Physiology*
146: 1155–1164.

Gleave AP. 1992. A versatile binary vector system with a T-DNA organisational structure
conducive to efficient integration of cloned DNA into the plant genome. *Plant Molecular*
Biology **20:** 1203–1207.

Heide-Jorgensen H. 2008. *Parasitic Flowering Plants.* Leiden: Brill Academic Publishers.

Heinemeyer W, Buchmann I, Tonge DW, Windass JD, Alt-Moerbe J, Weiler EW, Botz T,
Schröder J. 1987. Two *Agrobacterium tumefaciens* genes for cytokinin biosynthesis: Ti
plasmid-coded isopentenyltransferases adapted for function in prokaryotic or eukaryotic cells.
Molecular and General Genetics **210:** 156–164.

Hellens RP, Edwards E a, Leyland NR, Bean S, Mullineaux PM. 2000. pGreen: a versatile
and flexible binary Ti vector for *Agrobacterium*-mediated plant transformation. *Plant*
Molecular Biology **42:** 819–832.

Ishida JK, Yoshida S, Ito M, Namba S, Shirasu K. 2011. *Agrobacterium rhizogenes*-
mediated transformation of the parasitic plant *Phtheirospermum japonicum. PloS one* **6:**
e25802.

Kakimoto T. 2001. Identification of plant cytokinin biosynthetic enzymes as dimethylallyl
diphosphate:ATP/ADP isopentenyltransferases. *Plant and Cell Physiology* **42:** 677–685.

Kasahara H, Takei K, Ueda N, Hishiyama S, Yamaya T, Kamiya Y, Yamaguchi S, Sakakibara H. 2004. Distinct isoprenoid origins of cis- and trans-zeatin biosyntheses in *Arabidopsis*. *Journal of Biological Chemistry* **279:** 14049–14054.

Kieber JJ, Schaller GE. 2014. Cytokinins. *The Arabidopsis Book* **12:** e0168.

Kokla A, Melnyk CW. 2018. Developing a thief: Haustoria formation in parasitic plants. *Developmental Biology* **442:** 53–59.

Kurotani K, Wakatake T, Ichihashi Y, Okayasu K, Sawai Y, Ogawa S, Suzuki T, Shirasu K, Notaguchi M. 2020. Host-parasite tissue adhesion by a secreted type of β-1,4-glucanase in the parasitic plant *Phtheirospermum japonicum*. *Communications Biology* **3:** 407.

Miyawaki K, Matsumoto-Kitano M, Kakimoto T. 2004. Expression of cytokinin biosynthetic isopentenyltransferase genes in *Arabidopsis*: tissue specificity and regulation by auxin, cytokinin, and nitrate. *The Plant Journal* **37:** 128–138.

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka K, Jinbo T, Kimura T. 2007. Development of series of gateway binary vectors, pGWBs, for realizing efficient construction of fusion genes for plant transformation. *Journal of Bioscience and Bioengineering* **104:** 34–41.

Nekrasov V, Staskawicz B, Weigel D, Jones JDG, Kamoun S. 2013. Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nature Biotechnology* **31:** 691–693.

Nickrent DL. 2020. Parasitic angiosperms: How often and how many? *TAXON* **69:** 5–27.

Ogawa S, Wakatake T, Spallek T, Ishida JK, Sano R, Kurata T, Demura T, Yoshida S, Ichihashi Y, Schaller A, Shirasu K., 2020. Subtilase activity in intrusive cells mediates haustorium maturation in parasitic plants. *Plant Physiology* **185:** 1381-1394.

R Core Team 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Sakakibara H. 2006. Cytokinins: activity, biosynthesis, and translocation. *Annual Review of Plant Biology* **57:** 431–49.

Spallek T, Melnyk CW, Wakatake T, Zhang J, Sakamoto Y, Kiba T, Yoshida S, Matsunaga S, Sakakibara H, Shirasu K. 2017. Interspecies hormonal control of host root morphology by parasitic plants. *Proceedings of the National Academy of Sciences*, USA **114:** 5283–5288.

Spíchal L. 2012. Cytokinins - Recent news and views of evolutionally old molecules. *Functional Plant Biology* **39:** 267–284.

Sun G, Xu Y, Liu H, Sun T, Zhang J, Hettenhausen C, Shen G, Qi J, Qin Y, Li J, et al.
2018. Large-scale gene losses underlie the genome evolution of parasitic plant *Cuscuta*
australis. *Nature Communications* **9:** 2683.

Vogel A, Schwacke R, Denton AK, Fischer K, Bolger A, Marie E, Gundlach H, Mayer
KFX, Weiss-Schneeweiss H, Eva M, et al. 2018. Footprints of parasitism in the genome of
the parasitic flowering plant *Cuscuta campestris*. *Nature Communications* **9:** 2515.

Wakatake T, Ogawa S, Yoshida S, Shirasu K. 2020. An auxin transport network underlies
xylem bridge formation between the hemi-parasitic plant *Phtheirospermum japonicum* and host
Arabidopsis. *Development* **147:** dev187781.

Wakatake T, Yoshida S, Shirasu K. 2018. Induced cell fate transitions at multiple cell layers
configure haustorium development in parasitic plants. *Development* **145:** dev164848.

Yang Z, Wafula EK, Honaas LA, Zhang H, Das M, Fernandez-Aparicio M, Huang K,
Bandaranayake PCG, Wu B, Der JP, et al. 2015. Comparative transcriptome analyses reveal
core parasitism genes and suggest gene duplication and repurposing as sources of structural
novelty. *Molecular Biology and Evolution* **32:** 767–790.

Yoshida S, Cui S, Ichihashi Y, Shirasu K. 2016. The Haustorium, a specialized invasive
organ in parasitic plants. *Annual Review of Plant Biology* **67:** 643–667.

Yoshida S, Kim S, Wafula EK, Tanskanen J, Kim Y-M, Honaas L, Yang Z, Spallek T,
Conn CE, Ichihashi Y, et al. 2019. Genome sequence of *Striga asiatica* provides insight into
the evolution of plant parasitism. *Current Biology* **29:** 3041-3052.

Zhang X, Berkowitz O, Teixeira da Silva JA, Zhang M, Ma G, Whelan J, Duan J. 2015.
RNA-Seq analysis identifies key genes associated with haustorial development in the root
hemiparasite *Santalum album*. *Frontiers in Plant Science* **6:** 661.

Zhang X, Teixeira da Silva JA, Duan J, Deng R, Xu X, Ma G. 2012. Endogenous hormone
levels and anatomical characters of haustoria in *Santalum album* L. seedlings before and after
attachment to the host. *Journal of Plant Physiology* **169:** 859–866.

Figure legends

Figure 1:   ***PjIPT1a* is specifically expressed in intrusive cells.** (a) Prenylation of

adenosine-tri-, -di- and -mono-phosphates (ATP, ADP, AMP) by isopentenyltransferase (IPT)

with di-methylallyl pyrophosphate (DMAPP) to isopentyl-ribosides (IPR) is the first step of

cytokinin (CK) biosynthesis. (b) RNA-Sequencing data showing increased expression of

PjIPT1a in intrusive cells of *Phtheirospermum* (Pj) haustoria five days-post infection (dpi) of

rice roots (*Oryza sativa*, Os), and (c) upregulation of *PjIPT1a* expression during infection of

Arabidopsis (*At*) roots with a maximum of 61 CPM (counts per million mapped reads) at 3 dpi.

(d - i) Confocal   images   show   a   *Phtheirospermum*   haustorium   expressing

pIPT1:PjIPT1:3xVenus-NLS during infection of Arabidopsis (At) at 4 dpi. Merged yellow

channel for Venus detection and bright-field is depicted in (d), individual Venus and bright-

field channels in (e) and (f), respectively. Zoomed-in scans shown in g - i correspond to the

region marked by dotted lines in (d – f). Intrusive cells are indicated with white arrows. Scale

bars = 100 μm.

Figure 2:   ***PjIPT1a* and its closest homolog are nuclear-cytoplasmic IPTs** Phylogenetic

analysis (UPGMA) based on the amino acid sequences of prokaryotic (*Agrobacterium*) and

plant IPTs positions IPT1a within a cluster of plant IPT1s and closest to the *Striga* ShIPT1a

protein. Bootstrap values are shown as percentage of 1000 repetitions. (b) PjIPT1a and PjIPT1b

(c) located in the cytoplasm and nucleus when expressed in *N. benthamiana*. The subcellular

localization pattern of the full-length PjIPT1a-GFP and PjIPT1b-GFP fusions matched the

pattern observed for PjIPTa[(1-22)]-sfGFP (d) and free GFP (e). Scale bar = 50 μm.

Figure 3:   ***PjIPT1a* drives host CK responses at the haustorium.** (a) Control (con.)

haustoria of hairy roots without the *Cas9/PjIPT1a-gRNA*s cassette (HR1, HR2) induced the

expression of host CK-response reporter *pARR5:GFP* and hypertrophic growth (arrows)

whereas haustoria with the *Cas9/PjIPT1a-gRNA*s cassette (HR3-4) did not. Images were taken

at 28 dpi, scale bars = 100 μm (b) Schematic representation of the *PjIPT1a* gene flanked by the

5' and 3' extension that were amplified in (c). Red colour highlights the target sequences of the

guide RNAs (c) PCR amplicons of the *PjIPT1a* and the *CAS9* genes in Phtheirospermum hairy

roots were sequenced to determine *CAS9-induced mutation (red)* as shown in (d). PAM

sequences are depicted in orange, the start codon in bold font.

Figure 4:   PjIPT1a and PjIPT1b are isopentyltransferases. Plots show average +/- SD

(n = 3 – 6) peak areas of extracted-ion chromatograms of individual and cumulative (sum)

quantifications of isoprenylated derivatives that were detected in reaction mixtures containing

DMAPP, equal amounts of *E. coli* purified GST-His, PjIPT1b-His or PjIPT1b-His, and (a)

AMP or (b) ATP as isoprenoid acceptors. Different lower-case letters indicate significant

differences according to Tukey HSD with 95% confidence levels.

a

DMAPP
+ ATP
+ ADP → IPT → IPRTP
+ AMP   IPRDP → CK
         IPRMP

b

PjIPT1a   5 dpi

Os

reads: 9.3 236

c

dpi: 0   1   2   3   5   7

At   At   At   At   At   At

CPM: 2.9 0.9 21.1 61.0 45.4 38.5

d Pj At g

e Pj At h

f Pj At i

g

h

i

Merged   Venus   Bright-field

a

100	Tzs	*Agrobacterium*
	Tmr	*Agrobacterium*

ShIPT1a *Striga*
PjIPT1a *Phtheirospermum*
SrIPT1 *Streptocarpus*
ShIPT1b *Striga*
PjIPT1b *Phtheirospermum*
HlIPT1 Hop
AtIPT4 *Arabidopsis*
AtIPT1 *Arabidopsis*
AtIPT7 *Arabidopsis*
AtIPT5 *Arabidopsis*
PjIPT5 *Phtheirospermum*
SrIPT3 *Streptocarpus*
PjIPT3 *Phtheirospermum*
AtIPT3 *Arabidopsis*
SrIPT2 *Streptocarpus*
PjIPT2 *Phtheirospermum*
AtIPT2 *Arabidopsis*
SrIPT9 *Streptocarpus*
AtIPT9 *Arabidopsis*

0.4

b PjIPT1a-GFP

c PjIPT1a$^{(1-22)}$-sfGFP

d PjIPT1b-GFP

e GFP