Rhizobiales commensal bacteria promote *Arabidopsis thaliana* root growth via host sulfated peptide pathway

Jana Hucklenbroich[1], Tamara Gigolashvili[2], Anna Koprivova[2], Philipp Spohr[3], Mahnaz Nezamivand Chegini[2,4], Gunnar W. Klau[3], Stanislav Kopriva[2], Ryohei Thomas Nakano[1,#]

[1]Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, Cologne, Germany

[2]Institute for Plant Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), University of Cologne, Cologne, Germany

[3]Algorithmic Bioinformatics, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich Heine University, Düsseldorf, Germany

[4]Institute of biotechnology, Shiraz university, Shiraz, Iran

[#]Corresponding author: nakano@mpipz.mpg.de

Short title:
Root manipulation by commensal bacteria

Abstract

Root-associated commensal bacteria that belong to the order Rhizobiales, which also contains symbiotic and pathogenic bacteria, promote primary root growth of *Arabidopsis thaliana*. Yet, its underlying molecular mechanism and physiological impact remained unclear. Here, we conducted a transcriptomic analysis of *A. thaliana* roots inoculated with root-associated commensal bacteria of Rhizobiales and sister lineages and revealed common and strain/lineage-specific transcriptional responsea, possibly mediated by WRKY and ANAC family of transcription factors. We revealed that the observed common response was also partially triggered by a wide range of non-pathogenic bacteria, fungi, and a multi-kingdom synthetic community (SynCom). This response was characterized by a down-regulation of genes related to intracellular redox regulation, suggesting distinctive redox status between pathogenic and non-pathogenic interactions. By integrating this analysis with developmental and cell biological analyses, we identified a crucial role for the sulfated peptide pathway mediated by TYROSYLPROTEIN SULFOTRANSFERASE (TPST) in Rhizobiales root growth promotion (RGP) activity. Conversely, none of the known sulfated peptide pathway appeared to be required for this activity, suggesting a novel sulfated protein pathway targeted by Rhizobiales RGP. Finally, we show that TPST is needed for RGP exerted by Rhizobiales but not Pseudomonadales isolates, delineating lineage-specific mechanisms to manipulate host root development.

Introduction

Various prokaryotic and eukaryotic microorganisms heavily colonize plant tissues in nature, collectively shaping the plant microbiota. Many of these do not engage in pathogenic or mutually symbiotic lifestyles, and are therefore frequently referred to as "commensal" microbes (Berg et al., 2020). Bacterial communities associated with roots (root bacterial microbiota) are significantly distinct from soil microbiota (Bulgarelli et al., 2012; Lundberg et al., 2013) and qualitatively highly conserved across different plant species at the family level (Hacquard et al., 2015; Thiergart et al., 2020). At the quantitative and strain levels, however, there appears to be host species/genotype specificity (Wagner et al., 2016; Zgadzaj et al., 2016; Wippel et al., 2021; Yu et al., 2021). These findings indicate the existence of a regulatory mechanism in plant and bacteria that enables taxonomically defined community assembly in a host-dependent manner. It also implies that, over the course of evolution, the extant microbiota

members have benefited the host plants and *vice versa*. Recent studies have provided

evidence that the root bacterial microbiota as a community is vital for plant health, for

example, by protecting the host from fungi (Wei et al., 2015; Duran et al., 2018) or by

facilitating nutrient acquisition (Castrillo et al., 2017; Harbort et al., 2020; Salas-

Gonzalez et al., 2021). Imbalance of the microbiota compositions ("dysbiosis") in roots

as well as in leaves results in growth penalty (Chen et al., 2020; Finkel et al., 2020),

illustrating the importance of eubiotic homeostasis of root microbiota (Paasch and He,

2021). Moreover, root bacterial microbiota members can interfere with root

development and immunity. For example, many if not all commensal bacteria isolated

from healthy *A. thaliana* roots are capable of either promoting or inhibiting host primary

root growth in mono-association (Finkel et al., 2020). Another subset of root

commensals have been shown to suppress root growth inhibition triggered by a

chronic exposure to flg22, a 22-amino acid immunogenic peptide derived from

bacterial flagella (thereby referred to as suppressive isolates)(Gomez-Gomez et al.,

1999; Garrido-Oter et al., 2018; Yu et al., 2019; Ma et al., 2021; Teixeira et al., 2021).

Overall, these studies have demonstrated the holobiome concept, an idea of

collectively considering plants and associated microbiota as a single physiological and

ecological unit. In contrast, our knowledge about the physiology of plants, including

developmental regulation of roots, has been primarily provided by the studies using

axenic plants. Toward ultimate understanding of the plant molecular physiology under

ecological conditions, it remains necessary to better understand the molecular dialog

between plants and commensal bacteria and how this influences host physiological

pathways.

Comparative analysis of root microbiota across a wide range of plant species

has identified bacterial lineages that are highly prevalent across and abundant within

root microbiota (Hacquard et al., 2015). These include Rhizobiales, which is a bacterial

order within the class Alphaproteobacteria, containing strains that engage in different

lifestyles. Bacterial species in the order Rhizobiales include rhizobia, a paraphyletic

group of bacteria that induce nodule organogenesis in legume roots and fix

atmospheric nitrogen into ammonium for the host growth, and Agrobacterium, a

monophyletic group of bacteria within the genus *Rhizobium* that invade plant tissues

to induce tumors and ultimately restrict plant growth. These lifestyles are highly

differentiated and require dedicated genes in bacteria and plants, such as *nod* and *nif*

genes in rhizobia, *vir* genes in Agrobacterium, and *NOD FACTOR RECEPTOR* (*NFR*)

genes in legumes. The vast majority of Rhizobiales order bacteria, including strains
that were isolated from soil or animals and therefore do not directly engage in
interaction with plant hosts, do not possess these genes (Garrido-Oter et al., 2018). A
recent study showed that host-associated and soil free-living lifestyles are frequently
interchangeable (Wang et al., 2020). We previously showed that the extant
commensal Rhizobiales bacteria are commonly capable of promoting primary root
growth of *Arabidopsis thaliana* Col-0 wild-type plants when gnotobiotically inoculated
with germ-free hosts. Bacterial isolates from the sister lineages, Sphingomonadales
and Caulobacterales, did not possess this host root growth promotion (RGP) activity,
implying that the RGP activity represents an ancestral trait specific to the order
Rhizobiales (Garrido-Oter et al., 2018). The wide prevalence among extant
Rhizobiales strains also suggests an important role of the RGP activity in interaction
between *A. thaliana* roots and Rhizobiales commensals. Therefore, revealing the
molecular mechanisms by which Rhizobiales bacteria interfere with host root growth
may provide a valuable insight into the root-commensal molecular dialog.

In the previous study, we showed that Rhizobiales RGP was associated with an
enlargement of the root meristematic zone (MZ), where cells undergo proliferation
without elongation or differentiation, while the cell size in the elongation zone (EZ) and
the differentiation zone (DZ) was comparable between axenic and R129_E-inocualted
roots (Garrido-Oter et al., 2018). Root cell growth and proliferation are regulated by
an orchestrated operation of various pathways: The mitotic activity of stem cell
daughters within the MZ is maintained by a gradient accumulation of PLEHTHORA
(PLT) transcription factors (TFs) under the control of auxin (Aida et al., 2004; Galinha
et al., 2007). Auxin accumulated within the MZ induces expression of
TYROSYLPROTEIN SULFOTRANSFERASE (*TPST*), which encodes an enzyme to
catalyze protein sulfation of proteins (Komori et al., 2009; Zhou et al., 2010), including
ROOT MERISTEM GROWTH FACTOR (RGFs) (Matsuzaki et al., 2010; Shinohara,
2021). Recent studies showed that perception of RGFs lead to activation of
MITOGEN-ACTIVATED PROTEIN (MAP) kinases and redistribution of reactive
oxygen species (ROS) to increase the PLT protein accumulation (Lu et al., 2020; Shao
et al., 2020; Yamada et al., 2020). Accumulation of auxin and PLT gradually decreases
along with the longitudinal axis of the roots and an auxin minimum is formed at the
transition zone (TZ) where cytokinin antagonizes with auxin to switch cell fate from
division to elongation (Dello Ioio et al., 2008; Di Mambro et al., 2017). This transition

process is also influenced by brassinosteroid and ROS distribution (Tsukagoshi et al.,
2010; Chaiwanon and Wang, 2015). After the cell fate transition, the root cells continue
growing within the EZ until they reach to the DZ. This elongation in the EZ is facilitated
by gibberellin and another group of sulfated peptides, PHYTOSULFOKINE (PSK) and
PLANT PEPTIDE CONTAINING SULFATED TYROSINE 1 (PSY1) (Matsubayashi et
al., 2006; Amano et al., 2007; Achard et al., 2009; Ubeda-Tomas et al., 2009; Ladwig
et al., 2015), whose sulfation is also dependent on TPST (Komori et al., 2009). The
tpst mutant of *A. thaliana* lacks all functional sulfated peptides, including but not limited
to RGFs, PSKs, and PSY1, as well as another class of sulfated peptides involved in
endodermal cell differentiation (CASPARIAN STRIP INTEGRITY FACTOR, CIFs;
Doblas et al., 2017), shows a highly disorganized root MZ and therefore exhibits
severe root growth defects, showing the crucial role of the sulfate peptides in root
growth regulation (Komori et al., 2009). It is plausible that Rhizobiales commensal
bacteria target one of the aforementioned processes that coordinately dictate the
primary root growth to exert the RGP activity. Yet, its underlying molecular mechanism
is unknown. Our previous study showed that genetic perturbation of auxin-cytokinin
antagonism in *arr1arr12* and *ahk3-3* mutants (Dello Ioio et al., 2008) did not disrupt
Rhizobiales RGP (Garrido-Oter et al., 2018), pointing to a possible role of other
components than auxin and cytokinin.
In this study, to disentangle the molecular interactions between *A. thaliana* and
commensal Rhizobiales, we examined the host transcriptional response in roots
inoculated with Rhizobiales and sister lineage commensal isolates. We identified
common transcriptional responses commonly triggered by these bacteria. We
revealed that a part of the observed common response was also triggered by a wide
range of non-pathogenic bacterial and fungal microbes. By analyzing linage-specific
responses and the developmental details of roots inoculated with Rhizobiales, we
show that TPST, but none of the known receptors for RGF and PSKs, is indispensable
for Rhizobiales RGP. Lastly, we show that many Rhizobiales but not
Pseudomonadales commensals require the TPST pathway to promote root growth,
illuminating the lineage-specific innovation of mechanisms to interfere with the host
root development.
154
155
156

Results

Host transcriptional reprogramming by root-associated Alphaproteobacteria
commensals

To obtain an overview of transcriptional responses to Rhizobiales commensals, we isolated and sequenced RNA from *A. thaliana* (Col-0 wild type) roots inoculated with a panel of root-associated commensals that belong to the class Alphaproteobacteria. Our previous RNAseq analysis revealed that the transcriptional responses triggered by R129_E were similar in roots at 4, 8, and 12 days post-inoculation (dpi), which were largely different from roots at 16 dpi (Garrido-Oter et al., 2018). A statistically insignificant or marginal RGP phenotype is typically observed at 12 dpi at the level of primary root length, while developmental manipulation at the cellular level (i.e. enlarged root MZ) already initiates at an earlier time point (at 4-8 dpi). Therefore, we decided to harvest roots at 12 dpi for our RNA-seq experiment to consider the impact of Rhizobiales RGP at the cellular level while minimizing the difference accounted for by different root growth. As inoculant bacteria, we used R129_E, R13_A, Root491, and Root142 from the order Rhizobiales and Root1497 and Root700 from sister lineages Sphingomonadales and Caulobacterales, respectively, as an outgroup control. Among these isolates, only Rhizobiales strains exhibit root growth promotion (RGP) activity, while only R129_E and Root1497 are able to suppress host flg22-triggered root growth inhibition (Garrido-Oter et al., 2018). We identified a total of 1,439 differentially expressed genes (DEGs) between axenic and commensal-inoculated roots, based on the false discovery rate (FDR; α = 0.05) and $\log_2$-scale fold changes ($|\log FC| > 1$), and a principal coordinates analysis (PCoA) was performed using their zero-centered expression values (Figure 1A). This revealed that these commensals triggered a significant shift in the host transcriptional profiles (R^2 = 0.662, *P* < 0.001, PERMANOVA) in a manner dependent on the taxonomic lineages they belonged to (filled *versus* open shapes; R^2 = 0.457, *P* = 0.0045, pairwise PERMANOVA between Rhizobiales and sister lineage isolates). We revealed a number of DEGs specific to each inoculation, as well as DEGs commonly identified in more than one inoculation (Supplemental Figure 1). On the other hand, transcriptional responses to these commensal bacteria were qualitatively similar and showed a strong correlation (Figures 1B and 1C), indicating that the taxonomy-dependency of host transcriptional reprogramming, as shown in Figure 1A, is explained by their quantitative differences.

Figure 1. Alphaproteobacteria commensals trigger qualitatively similar transcriptional responses in host roots.

(A) A principal coordinates analysis separated transcriptome of axenic roots (asterisks), roots inoculated with Rhizobiales commensals (closed shapes), and roots inoculated with sister lineage isolates (open shapes). (B) A heatmap showing cosine similarities between logFC upon Alphaproteobacteria commensal inoculations. (C) Heatmaps showing log₂-scale count per million (log2cpm), zero-centered expression values, log₂-scale fold changes (logFC), and significance (FDR < 0.05; |logFC| > 1), aligned according to their k-means cluster assignment (k = 9, based on Bayesian information criterion). (D) Boxplots showing logFC of each k-means cluster. Asterisks indicate statistical significance corresponding to Student's one-sample t test after Bonferroni's correction for multiple comparisons (μ = 0; α = 0.001) and letters indicate statistical significance corresponding to pairwise Wilcoxon's rank sum test after Bonferroni's correction for multiple comparisons within each cluster (α = 0.05). (E) A heatmap showing enrichment of GO terms in the k-means clusters. (F) A network plot of enriched GO terms. Each node represents each GO term, and GOs with overlapping gene contents are connected with edges. k-means assignment of composing genes is shown as a pie chart. We selected the terms with the best P values from each of the 9 clusters, with the constraint that there are no more than 15 terms per cluster and no more than 250 terms in total. Representative terms are manually curated and indicated with composing k-means clusters for each GO cluster.

To obtain functional insights, we classified all 1,439 DEGs into nine *k*-means
gene clusters based on the zero-centered expression values and tested whether Gene
Ontology (GO) categories were enriched in each cluster (Figures 1C to 1F and
Supplemental Figure 2). We assigned down-regulated genes into Clusters 1, 2, and 3
and up-regulated genes into Clusters 4 to 9. Clusters 2 and 5 illustrated the general
response to these commensal bacteria, and Clusters 3, 4, and 6, and Clusters 1, 7,
and 9 represented responses quantitatively characteristic to Rhizobiales and sister
lineage isolates, respectively. The commonly up-regulated Cluster 5 was enriched in
genes related to metal ion transport and ageing, while commonly down-regulated
Cluster 2 was enriched in genes related to response to hypoxia and to various abiotic
stresses, as well as genes associated with biosynthesis of terpenoid, phytohormone
signal transduction, and the detoxification pathway. Clusters characteristic to sister
lineage isolates (Clusters 1, 7, and 9) were enriched in unique GO categories, such
as photosynthesis (Cluster 1), response to cold and water deprivation (Cluster 7), and
cell wall rearrangement and immune responses (Cluster 9). Clusters characteristic to
Rhizobiales were characterized by up- and down-regulation of "Response to hypoxia"
and "detoxification" (Clusters 3 and 4), down-regulation of biosynthesis of ethylene,
glucosinolate, and flavonoid, and response to various abiotic stresses (Cluster 3), and
up-regulation of response bacteria, wounding, salicylic acid (SA), and jasmonic acid
(JA) (Clusters 4 and 6). Cluster 8 contained genes up-regulated by all isolates except
for Root142, which were enriched in immune-related genes, such as genes
responding to SA, bacteria, fungi, and hypoxia or related to calcium homeostasis, as
well as genes involved in response to abiotic stresses such as heat, drugs, and amino
acids. We previously reported that Root142 is one of the Rhizobiales strains that were
able to efficiently colonize *A. thaliana* roots (Garrido-Oter et al., 2018), implying the
ability of Root142 to partly evade recognition by the host or to interfere with these host
immune responses, albeit its inability to suppress flg22-triggered root growth inhibition.
In contrast, only a handful (4 up- and 6 down-regulated genes) of DEGs were found
to be common between suppressive R129_E and Root1497 (Supplemental Figure 1),
probably accounted for by different mechanisms employed by these isolates to
interfere with host immune responses. These findings illustrate that the tested
Alphaproteobacteria commensals commonly interfere with the host genetic pathways
associated with hypoxia, detoxification, and plant immunity, to a different extent
depending on the isolates. Lineage-specific DEGs (46 and 99 genes specific to

Rhizobiales and the sister lineage, respectively) and isolate-specific DEGs (from 44 to 164 genes) also demonstrated the presence of lineage/strain-specific responses, whose functional aspects remained unclear based on their GO annotation.

We then looked for TF binding motifs enriched in the 1-kb upstream sequences of the genes in each cluster. We identified several binding motifs, mainly from the TF families of WRKY and ANAC, as well as the AHL family of DNA-binding proteins (Figure 2A), suggesting a role of these TF families in commensal-triggered transcriptional reprogramming. We found that 20 WRKY and NAC family TFs were differentially expressed (differentially expressed TFs, DETFs), and used these DETFs as a list of potential regulators to infer a gene regulatory network (GRN) by the so-called GENIE3 algorithm (Figure 2B). This algorithm considers both positive and negative potential regulations to infer a GRN, which is a strong advantage over correlation-based approaches, such as k-means clustering and gene co-expression networks (Huynh-Thu et al., 2010). The inferred GRN was composed of two arbitrarily defined modules, characterized by the predominance of either WRKY or NAC family DETFs as hub nodes (module 1 and module 2, respectively; Figure 2B). Among DETFs, WRKY54 and ANAC060 showed the first and second highest centrality indices (degree, closeness, and betweenness centralities), suggesting their role in the observed transcriptional reprogramming. Genes commonly induced/suppressed by Alphaproteobacteria commensals (Clusters 1, 8, and 9) were enriched in module 1, whereas genes in the Rhizobiales-characteristic clusters (Clusters 3, 4, ad 6) were enriched in module 2 (Supplemental Figures 3A and 3B). We observed similar trends using all DETFs, independent of promoter motif enrichment analysis, as potential regulators to infer a GRN, albeit with a less clear GRN structure (Supplemental Figure 4). Importantly, in both GRNs, WRKY54 and ANAC060 were identified as candidate hub regulators based on the centrality indices and connected with genes whose transcriptional responses were commonly triggered by all isolates or characteristically by Rhizobiales isolates, respectively. Together, our RNAseq experiments demonstrated common and lineage-specific transcriptional responses in roots that were potentially mediated by WRKY54 and ANAC060, respectively, along with other TFs.

Figure 2. WRKY and ANAC transcription factors are important for transcriptional reprogramming.
(A) Heatmaps showing mean logFC of each cluster (left) and the enrichment of corresponding transcription factor binding motifs in their promoter regions (1-kb upstream from transcription initiation sites). AHL, AT-HOOK MOTIF CONTAINING NUCLEAR LOCALIZED; CCHC, CCHC-type zinc-finger motif containing protein; ATHB, ARABIDOPSIS THALIANA HOMEOBOX; FRS9, FAR1-RELATED SEQUENCE9. (B) GENIE3-inferred gene regulatory network using WRKY (green) and ANAC (magenta) family DETFs as potential regulators (colored; bigger node size). Arbitrarily defined modules are shown by dotted lines. (C) Degree, closeness, and betweenness centralities were computed based on the network in (B) for all potential regulators. Green and magenta dots correspond to WRKY and ANAC family TFs.

 Overlapping transcriptional responses to a wide range of non-pathogenic microbes

Our RNAseq experiments identified common as well as lineage-specific transcriptional

responses, both of which were enriched in similar GO terms, such as "detoxification"

and "response to hypoxia". We noted that these GO terms were also found among

genes responding to other root microbiota members (Hacquard et al., 2016; Ma et al.,

2021; Teixeira et al., 2021; Wippel et al., 2021). This suggests that a wide range of

root-associated microbes might broadly interfere with this response, and we aimed to

understand the extent to which the observed root transcriptional response is common

and specific to Alphaproteobacteria and Rhizobiales commensals. We compared our

data with publicly available transcriptomic datasets using the roots inoculated with

root-inhabiting microbes that engage in beneficial, commensal, and pathogenic

lifestyles under different nutrient conditions, in different experimental setups, at

different time points (Supplemental Table 1). These include beneficial and pathogenic

Colletotrichum fungi (*C. tofieldiae* and *C. incanum*; Hacquard et al., 2016; Hiruma et

al., 2016), a bacterial synthetic community (B-SynCom; Harbort et al., 2020), and a

multi-kingdom synthetic community (SynCom) composed of bacteria, fungi, and

oomycetes (BFO-SynCom; Hou et al., 2020). In addition, we sequenced RNA obtained

from roots inoculated with beneficial and pathogenic *Burkholderia* bacteria (*B.*

phytofirmans and *B. glumae* isolated from onion and rice, respectively), as well as

from roots inoculated with R129_E in the presence or absence of flg22 to consider the

impact of host immune responses. We compared and clustered responses to different

microbial inoculations based on the similarity of logFC (Figures 3A). LogFC within non-

pathogenic inoculations exhibited significantly greater similarities than between the

responses to pathogenic and non-pathogenic inoculations (Figure 3B), without any

clear sign of the taxonomy or the complexity of respective inocula (shapes in the

dendrogram). A PCoA based on the transcriptome-wide logFC similarities revealed a

clustering of non-pathogenic inoculations, irrespective of their taxonomy and

complexity (Figure 3C). A pairwise PERMANOVA between different lifestyles

(commensal, beneficial, pathogenic, and SynCom) revealed that all combinations of

lifestyles significantly contribute to the difference in the host transcriptional responses

(FDR-corrected P values of 0.002 ~ 0.01), while the R^2 values were larger between

pathogenic and non-pathogenic lifestyles than within non-pathogenic lifestyles

(Supplemental Table 2). Notably, minimizing batch effects by a canonical analysis of

Figure 3. Non-pathogenic microbial inoculation triggered similar transcriptional responses in the host roots.
(A) A heatmap showing cosine similarities between logFC upon different microbial inoculations. DEGs were selected based on the inoculation in rows (numbers of DEGs are shown in parentheses), whose logFC cosine similarity was calculated in comparison with inoculations in columns. For each row, all cosine similarity values were plotted on the right, whose colors indicate the lifestyle of the comparing inoculation. (B) Cosine similarities within non-pathogenic inoculations and within pathogenic inoculations were higher than those between pathogenic and non-pathogenic inoculations. (C) Principal coordinates analysis based on logFC cosine similarities of all commonly quantified genes (n = 21,815). Colors correspond to lifestyles. (D) Constrained analysis of principal coordinates (CAP) based on the logFC cosine similarities, using the constraining and conditioning factors indicated above. Colors correspond to inoculated microbial strains. (E) A heatmap showing logFC of genes assigned with the GO categories related to detoxification and response to hydrogen peroxide, reactive oxygen species, and hypoxia. Inoculations are aligned according to the dendrogram in (A) and genes are aligned according to k-means clustering as shown in Supplemental Figure 2B. Numbers indicated in the names of inoculation indicate the time point of harvesting (dpi).

principal coordinates (CAP) clustered all non-pathogenic interactions with microbes
isolated from *A. thaliana* (Figure 3D)*.*

Concatenated lists of DEGs (13,889 genes in total) were then classified into 21
k-means clusters and tested for their enriched GO categories (Supplemental Figures
5A and 5B). This identified the detoxification pathway to be strongly enriched in Cluster
11 (P = 5.1×10^{-18}), whose transcriptional suppression characterized the non-
pathogenic interactions. Our of 22 genes in Cluster 11 assigned with the GO term
"detoxification", 13 genes encode glutathione (GSH) *S*-transferases (GSTs), which
play an important role in cellular redox homeostasis. The hypoxia response is known
to trigger ROS production via RESPIRATORY BURST OXIDASE HOMOLOG D
(RbohD) in *A. thaliana* (Pucciariello et al., 2012), and is associated with strong immune
response induction. Therefore, we speculated that non-pathogenic interactions may
impact the root redox response. Among DEGs, we found a total of 327 genes assigned
with GO categories of response to hypoxia, response to ROS, and detoxification
pathway (Figure 3E). The hypoxia-triggered immune response includes transcriptional
up-regulation of genes crucial for plant immunity, such as *FLAVIN-DEPENDENT*
MONOOXYGENASE 1 (*FMO1*; (Hartmann et al., 2018), *CALMODULIN-BINDING*
PROTEIN 60-LIKE G (*CBP60G*; (Wang et al., 2009), *ELICITOR PEPTIDE 3*
PRECURSOR (*PROPEP3*; (Huffaker and Ryan, 2007), *CYTOCHROME P450*
FAMILY 82, SUBFAMILY C, POLYPEPTIDE 2 (*CYP82C2*; (Rajniak et al., 2015), and
PEROXIDASE4 (*PER4*; (Rasul et al., 2012), in a WRKY22-dependent manner (Hsu
et al., 2013). In our meta-analysis, these genes were mostly up-regulated in many
pathogenic and non-pathogenic interactions (Clusters 3/5/17), suggesting that the
genetic sector responding to commensal bacteria inoculation overlaps with the
hypoxia response. On the other hand, another subset of hypoxia-responsive genes,
such as *RbohD*, *HEAT SHOCK TRANSCRIPTION FACTOR A 2* (*HSFA2*),
ARABIDOPSIS NAC DOMAIN CONTAINING PROTEIN 102 (*ANAC102*), and
HYPOXIA RESPONSIVE ERF2 (*HRE2*), along with many *GST* genes, were found in
Cluster 11 and down-regulated by non-pathogenic microbes. Down-regulation of the
transcriptional levels of *GST*s was not limited to a subset of isoforms but broadly
observed across most of the *GST* genes encoded by the *A. thaliana* genome
(Supplemental Figure 5C), suggesting a link to the general redox response rather than
a specific secondary metabolic pathway (discussed below). Overall, these results
suggest that the host root redox status linked with the expression of genes related to

response to hypoxia and the detoxification pathway is associated with different types
of interactions between pathogenic and non-pathogenic microbe, rather than the
Rhizobiales- or Alphaproteobacteria-specific responses.
328
ANAC060 transcription factor characterized Rhizobiales-specific root transcriptional
response
Our initial RNAseq experiment identified the hypoxia response and detoxification
pathway in the genes enriched in Rhizobiales-characteristic gene clusters (Clusters 3
and 4 in Figure 1), while the subsequent meta-analysis revealed that these gene
pathways were rather associated with interaction with a wide range of microbes.
Clusters 3, 11, and 17 contained more than 40% of Alphaproteobacteria DEGs
(Supplemental Figure 5A, rank abundance plot) and contained genes related to
hypoxia-responsive immunity and detoxification pathways (Figure 3E), suggesting that
the hypoxia response and detoxification pathways are not necessarily associated with
the Rhizobiales RGP activity. In order to characterize the molecular mechanism
responsible for Rhizobiales-triggered RGP, we aimed to further explore the response
characteristic to Rhizobiales independently of k-means clustering and Gene Ontology
annotations. Given the uncharacterized nature of interaction included in the meta-
analysis, we again focused on our dataset targeted to Alphaproteobacteria
commensals. We identified a total of 46 genes that were significantly up- or down-
regulated by Rhizobiales but not by sister lineage isolates ("Rhizobiales DEGs"; Figure
4A), which were then mapped onto the GRN shown in Figure 2B. We found that
Rhizobiales DEGs were exclusively located on a sub-module within module 2 and
tightly associated with *ANAC060* (Figures 4B and 4C and Supplemental Figure 3C).
Given that *ANAC060* is one of the two DETFs with the highest centralities (Figure 3C)
and the only TF among Rhizobiales DEGs (Figure 4A), we speculate that ANAC060
is crucial for the Rhizobiales-specific transcriptional response, a part of which
ultimately triggers RGP phenotype.
353
354 *Plant sulfated peptides are required for R129_E RGP*
355 We previously reported that the Rhizobiales RGP is associated with the enlargement
356 of and increased cell number within the root MZ, but not with cell size, pointing to
357 facilitated cell division triggered by Rhizobiales commensals. We confirmed this by
358 staining cells undergoing S-phase by a thymidine analog 5-ethynyl-2'-deoxyuridine

Figure 4. Rhizobiales-specific DEGs are potentially regulated by ANAC060.

(A) Heatmaps showing log2cpm, zero-centered expression values, logFC, significance, and GO assignment of genes specifically regulated by Rhizobiales commensals but not by sister lineage isolates. (B) Enlarged view of the GENIE3-inferred GRN, shown in Figure 2B, colored for potential regulator TFs (left) or Rhizobiales-specific DEGs (right). (C) Same view as in (B), colored by their logFC upon different bacterial inoculation.

(EdU; Supplemental Figure 7A). This corroborated the scenario that Rhizobiales RGP

targets root MZ homeostasis. On the other hand, we did not find a dramatic increase

or decrease of auxin and cytokinin, at least not at the detectable level of DR5:ER-GFP

and TCSn:ER-GFP accumulation (Supplemental Figures 7B and 7C), which further

supported the potential role of other components than auxin-cytokinin antagonism. In

published single-cell RNA sequencing (scRNA-seq) data of *A. thaliana* roots (Denyer

et al., 2019), *ANAC060* showed relatively high expression in the cell population

characterized by the expression of meristematic genes ("cluster 11"; Supplemental

Figures 6A and 6B). We also found that the top 100 genes co-expressed with

ANAC060 across 4,727 sequenced root cells were enriched in the GO categories

related to root MZ regulation, such as "meristematic growth," "regulation of cell

division," and "response to gravity" (Supplemental Figure 6C). Interestingly, genes that

were best co-expressed with *ANAC060* in the scRNAseq dataset included *RGF3*

(14th), as well as *PLT1* (5th) and *PLT2* (63rd) (Supplemental Figures S6A). Moreover,

a pervious study reported that ANAC060 potentially represses the expression of PSY1

while activating the expression of PSY4, a sequence-related homologue of PSY1 (Yu

et al., 2020; Tost et al., 2021), overall suggesting a potential link between ANAC060

and the sulfated peptide pathway.

To test the link between Rhizobiales RGP and the plant sulfated peptide pathway,

we inoculated a Rhizobiales commensal R129_E with *tpst* mutant plants, depleted in

all classes of functional sulfated peptides. Strikingly, we identified a lack of the RGP

phenotype in the *tpst* mutant (Figure 5A), without a clear depletion of bacterial cells

from roots (Figure 5B). In contrast, we observed significant RGP in *rgi1;2;3;4;5-1*,

pskr1pskr2psy1r, and *sgn3* mutants, which are impaired in perception of RGF, PSK,

and CIF peptides, respectively (Figure 5C and Supplemental Figure 8A). Likewise, the

RGP activity of R129_E remained functional when the three major isoforms of RGF

peptides operating in the root MZ were genetically abolished in the *rgf123* mutant

(Supplemental Figure 8A) (Matsuzaki et al., 2010). Disorganized root MZ in the *tpst*

mutant is known to be rescued by external application of RGF1 peptide but not its

desulfated form (dRGF1). However, treatments with neither of these peptides rescued

the RGP activity of R129_E, despite the substantial recovery of root MZ under these

conditions (Supplemental Figures 8B and 8C). These findings demonstrate that

R129_E requires the host sulfotransferase enzyme TPST for its RGP activity but not

the RGF and PSK peptide pathways. Of note, the root MZ in the *rgi1;2;3;4;5* mutant

Figure 5. Plant sulfated peptide pathway is needed for R129_E RGP activity.
(A) Primary root growth and representative pictures of Col-0 and *tpst* plants grown axenically or inoculated with R129_E. Seeds were inoculated and grown under continuous light conditions for 12 days. (B) Bacterial abundance relative to plant biomass was quantified by DNA-based quantitative PCR using primers specific to *A. thaliana* and R129_E *16S rRNA* genes. No clear depletion of bacterial cells from *tpst* roots was observed. (C) Mutants impaired RGF1 perception (*rgi1;2,3;4;5-1*) or PSK perception (*pskr1pskr2psy1r*) did not phenocopy *tpst* when 7-day-old seedlings were inoculated with R129_E for 14 days under short-day conditions. (D) Col-0 root MZ accumulate higher amounts of proPLT1::PLT1-GFP when 7-day-old seedlings were inoculated with R129_E for 7 days under short-day conditions, whereas *tpst* root MZ accumulated a similar amount irrespective of R129_E inoculation. For each panel, a merged image of propidium iodide staining (gray) and PLT1-GFP (green) signals and an image only showing PLT1-GFP signals pseudo-colored as shown in the calibration bar are shown in left and right, respectively. Bars indicate 50 µm. Images are aligned with quiescent centers on the dotted line. Letters indicate statistical significance (α = 0.05) corresponding to Student's *t* test after fitting to a linear mixed model with biological and technical replicates as random factors, corrected for multiple comparison by the Benjamini-Hochberg method.

was even more severely disrupted than in the *tpst* mutant (Ou et al., 2016; Song et al.,
2016), while this mutant nonetheless exhibited the RGP activity when inoculated with
R129_E (Figure 5C). Combined with the observation that restoration of root MZ by
RGF1 did not rescue RGP activity, our findings point to a specific regulatory role for
sulfated peptides, making the potential confounding effects due to the pleiotropic effect
of highly disorganized root MZ in *tpst* less likely. Lastly, we inoculated wild-type and
tpst mutant plants expressing PLT1-GFP under its *cis*-regulatory sequence to test
whether R129_E showed any impact on PLT1 protein accumulation. This showed that
R129_E indeed stimulates the accumulation of PLT1-GFP in Col-0 root MZ, which was
not observed when the sulfated peptides were abolished in the *tpst* root MZ (Figure
5D). Overall, these findings indicate that the host sulfated peptide pathway is
indispensable for R129_E-stimulated accumulation of PLT1 protein, thereby
facilitating the cell division within the root MZ and promoting primary root growth.
406
The sulfated peptide pathway is needed by Rhizobiales but not Pseudomonadales
RGP commensals
We previously showed that the RGP activity is prevalent within the order Rhizobiales
(Garrido-Oter et al., 2018), while it remained unclear whether they promote root growth
by the same molecular mechanism. We, therefore, inoculated *tpst* mutants with two
additional Rhizobiales isolates, Root491 (*Agrobacterium*) and Root142
(*Sinorhizobium*), which were both used in the RNAseq analysis (Figure 1), and found
that none of these isolates were able to promote *tpst* mutant root growth (Figure 6A).
To further explore its prevalence, we inoculated *tpst* with a panel of Pseudomonadales
isolates, namely Root7, Root71, Root329, and Root569, which are also capable of
promoting *A. thaliana* root growth in a manner yet to be identified. Interestingly, these
isolates showed RGP activity even in the absence of the host TPST enzyme (Figure
6B). These findings indicate a specific and conserved contribution of the plant sulfated
peptide pathway to Rhizobiales RGP, but not Pseudomonadales RGP, illustrating the
clear distinction between these two RGP phenotypes at the molecular level. This also
implies that root-associated commensal bacteria have independently acquired RGP
capability more than once, pointing to a potential physiological relevance of this
phenotype in the context of plant-microbiota interactions.
425
426

Figure 6. TPST is needed for the RGP activity explicitly exerted by Rhizobiales RGP but not Pseudomonadales.

Seven-day-old seedling of Col-0 and *tpst* were inoculated with a panel of Rhizobiales commensals (A) or Pseudomonadales commensals (B) and grown for 14 days under short-day conditions. Letters indicate statistical significance (α = 0.05) corresponding to Student's t test after fitting to a linear mixed model with biological and technical replicates as random factors, corrected for multiple comparison by the Benjamini-Hochberg method.

Discussion

WRKY and ANAC family TFs in transcriptional responses to Alphaproteobacteria commensals

Our comparative transcriptomic analysis identified genes that were differentially regulated by a panel of Alphaproteobacteria commensals. This enabled us to identify common as well as strain/lineage-specific responses, which were qualitatively similar and functionally overlapped with each other. The 1-kb upstream sequences of these genes tend to possess binding motifs of WRKY and ANAC family TFs, suggesting the role of these families of TFs in Alphaproteobacteria commensal-triggered transcriptional reprograming. WRKY family TFs are widely known for their crucial role in defense-related stress responses and are often also involved in plant development such as leaf senescence (Phukan et al., 2016), while many ANAC family TFs are reported to be involved in stress-induced regulation of development (Shao et al., 2015) and also in plant immunity (Yuan et al., 2019). It is well known that the WRKY family of TFs form a highly rigid transcriptional network (Eulgem and Somssich, 2007), and disruption of a single gene is frequently insufficient for perturbation of the entire network. This also appears to be the case for the ANAC family of TFs (Kim et al., 2018), pointing to a complex transcriptional network that ultimately contribute to the commensal-triggered reprogramming. The TF families identified in our promoter enrichment analysis were also found in a recent study addressing the co-occurrence of multiple TF binding motifs in the genes responding to immunogenic stimuli in roots in a cell-type specific manner (Rich-Griffin et al., 2020), further supporting the idea of their key role in plant-microbe interactions in roots of *A. thaliana*. A detailed time-course transcriptomic analysis is needed to identify the key regulator of the entire response and to test its contribution to the observed transcriptomic shift.

Host redox status may play a role in the distinction between pathogenic and non-pathogenic microbes

Recent studies described transcriptional responses in roots and leaves to another set of a panel of *A. thaliana* root- and leaf-derived bacterial commensals (Maier et al., 2021; Nobori et al., 2021; Teixeira et al., 2021). These commensal bacteria triggered qualitatively similar responses in the host, which is consistent with our finding that the response to Alphaproteobacteria commensals were largely similar. Furthermore, we have developed a method to compare transcriptional responses analyzed in

independent RNAseq experiments. This allowed us to identify a substantial overlap

between root transcriptional responses to a wide range of bacteria, fungi, and

SynComs, inoculated in very different experimental setups, expanding the model of

common transcriptional responses to a broader range of interactions. This includes

up-regulation of hypoxia-responsive immune-related genes by both pathogenic and

non-pathogenic interactions even after weeks of inoculation without causing growth

restriction. This is contradictory with the growth-defense tradeoff model, according to

which a chronic activation of immunity results in growth inhibition (Gomez-Gomez et

al., 1999; Huot et al., 2014), suggesting the presence of an immune response sector

that is uncoupled from the tradeoff. We also identified genes specifically and

commonly down-regulated by non-pathogenic microbes, which were enriched in the

detoxification pathway. We noted that this was mostly driven by the down-regulation

of genes encoding GSTs. GSTs are often involved in secondary metabolic pathways,

for example, by conjugation of GSH to toxic intermediates (thereby annotated as

detoxification pathway components). Moreover, GSH is one of the critical components

in sensing and regulating the cellular redox status, where GSTs play a crucial role

(Noctor et al., 2012). We observed that the down-regulation of GSTs was not limited

to GST isoforms that were specific to the family Brassicaceae (Supplemental Figure

5D, shaded by blue), which are often involved in lineage-specific secondary

metabolism (Dixon et al., 2010), but were broadly observed among different classes

of GSTs, including GSTUs and GSTFs (Pislewska-Bednarek et al., 2018). These

results point to a possibility that pathogenic and non-pathogenic microbes differentially

influence host cellular redox status. The broad impact on the expression of multiple

GST genes also implies that their transcriptional down-regulation may be a

consequence of altered GSH homeostasis, rather than a cause of the transcriptional

reprogramming. A recent study suggested that the extracellular and cytoplasmic ROS

responses are mutually distinguishable, where the cytoplasmic response is more

prolonged (Arnaud et al., 2021). GST proteins in *A. thaliana* do not have predicted

signal peptides for secretion (SignalP-5.0) and are mainly localized in the intracellular

compartments (Dixon et al., 2009; Noctor et al., 2012; Lallement et al., 2014). Given

that we observed the influence of non-pathogenic inoculations on genes related to

cellular redox at relatively late time points (from 4 days to 5 weeks), it is plausible that

the cytoplasmic redox regulation is involved in the non-pathogenic interactions, while

apoplastic ROS production is involved in the defense response against pathogens.

Overall, based on these findings, we speculate that host root intracellular redox status

has a role in the discrimination between pathogenic and non-pathogenic interactions,

either as a determinant of the response or as a consequence of different types of

interactions. Yet, the key molecular signatures in plants and microbes that differentiate

pathogenic and non-pathogenic interactions remain to be identified.

Rhizobiales commensals interfere with host sulfated peptide pathway to promote root

growth

We here provide a clear genetic evidence for the role of plant sulfated peptides in

Rhizobiales RGP. Plant sulfated peptides characterized thus far mostly play a crucial

role in plant development, including cell division, growth, and differentiation

(Kaufmann and Sauter, 2019). In roots, RGFs, PSKs, and PSY1 orchestrate with a

phytohormonal network to coordinate cell division and growth constitutively as well as

in response to environmental conditions (Matsubayashi et al., 2006; Amano et al.,

2007; Matsuzaki et al., 2010; Cederholm and Benfey, 2015), while CIFs specifically

regulate endodermal cell differentiation (Doblas et al., 2017). Our mutant analysis

demonstrated that known receptors of these sulfated peptides were not essential for

Rhizobiales RGP. This suggests the presence of unidentified cognate receptors or yet

another class of sulfated peptides. Receptors for extracellular peptides are typically a

member of the leucine-rich repeat receptor-like kinase (LRR-RLK) superfamily, which

has been immensely diversified in land plant lineages (Shiu et al., 2004; Furumizu et

al., 2021). To date, only a handful of LRR-RLKs have been functionally characterized,

while the physiological function of many members, including their cognate ligands,

remain unknown. Of note, while the *pskr1pskr2psy1r* mutant is mostly insensitive to

PSKs, it still significantly responds to the PSY1 peptide albeit to a lesser extent than

wild-type plants (Pruitt et al., 2017). Given that the direct binding between PSY1 and

PSY1R has not yet been experimentally shown, it is sensible to assume the presence

of other LRR-RLKs that perceive PSY1 and related peptides (Tost et al., 2021).

Besides, a large variety of uncharacterized secreted peptides are encoded by the *A.*

thaliana genome, including several classes of peptides with a predicted motif for

tyrosyl sulfation (aspartate-tyrosin; DY motif; Ghorbani et al., 2015), pointing to an

unexplored repertoire of plant sulfated peptides. Further characterization of the

mechanisms by which Rhizobiales promote root growth in a manner dependent on the

host sulfated peptide pathway may lead to identification of a novel class of receptors

and/or ligand sulfated peptides.

Interestingly, a hemibiotrophic bacterial pathogen *Xanthomonas oryzae* pv.

oryzae (*Xoo*) also produces a sulfated peptide, whose amino acid sequence is highly

similar to that of plant PSY1. This peptide, called RaxX, is needed for *Xoo* to exert its

virulence in rice, which is on the other hand recognized by the host immune receptor

XA21 to initiate responses that limit *Xoo* infection (Pruitt et al., 2015; Pruitt et al., 2017).

An exogenous treatment of *A. thaliana* plants with RaxX promotes root growth, as

does PSY1, and, therefore, it has been speculated that RaxX interferes with the PSY1

pathway to interrupt host immunity. Although RaxX has been the only bacterial

sulfated peptides that have been molecularly characterized thus far, it is plausible that

other bacterial species, especially those interacting with eukaryotic hosts, have

acquired the ability to secrete sulfated peptides that mimic and/or interfere with host

extracellular peptide signaling. In fact, genes with high similarity to the sulfotransferase

RaxST in *Xoo* were found in *A. thaliana* root-associated bacterial genomes (20 out

194 genomes, based on blastp with an *E* value cutoff of 1.0; (Bai et al., 2015), mostly

in those belonging to the orders Sphingomonadales, Burkholderiales,

Pseudomonadales, and Xanthomonadales. It is also possible that, even despite the

lack of sequence-related RaxST homologues, other bacterial species have

convergently acquired another class of enzymes that catalyze sulfation of secreted

peptides. Yet, it is not highly likely that the Rhizobiales RGP is mediated by bacterial

sulfated peptides. It is important to note that the TPST enzyme catalyzes tyrosine *O*-

sulfation within the Golgi apparatus, most likely at the *cis*-side, before the substrate

(pro)peptides reach the apoplast after its translation in the endoplasmic reticulum

(Komori et al., 2009). For external peptides to be sulfated by the host TPST, they need

to be internalized into the host cells and transported to *cis*-Golgi, followed by tyrosine

sulfation by TPST and re-secretion back to the apoplast. Overall, it is probable that

the lack of TPST in the host would not directly affect *de novo* production of sulfated

peptides by bacterial or mediate sulfation of naïve bacterial peptides in the

extracellular space. Our results rather delineate the involvement of plant endogenous

sulfated peptides.

 Physiological relevance of Rhizobiales RGP

Manipulation of host root growth appears to be a common feature of root-associated commensal bacteria (Zamioudis et al., 2013; Finkel et al., 2020), while one of the most common and dominant pathways is auxin-triggered root growth inhibition (Finkel et al., 2020). Despite the known mechanisms to manipulate host abiotic stress responses via phytohormonal network (Bulgarelli et al., 2013), little is known about the mechanisms by which root commensal promote root growth. Moreover, it is yet to be seen whether root growth phonotypes observed in artificial gnotobiotic system is relevant in the ecological/physiological context. Due to the plasticity and high variability of root growth depending on the environmental conditions, it remains challenging to directly tackle this question using wild-type plants. Here, we provide genetic evidence for the crucial role of sulfated peptides in Rhizobiales RGP. Retained RGP exerted by Pseudomonadales commensals in the *tpst* mutant is indicative of a regulatory role of sulfated peptides, rather than a confounding effect due to a pleiotropic disorganization of the root MZ in this mutant. ANAC060 may also be involved in this interaction, although the evidence remains circumstantial. A genetic analysis of ANAC060 and other related ANAC family TFs is needed to disentangle the transcriptional regulatory machinery that is associated with and possibly responsible for Rhizobiales RGP activity.

Our results show that two independent lineages in Alpha- and Gammaproteobacteria classes employ independent mechanisms to promote host root growth, demonstrating that root-associated bacteria have likely acquired RGP capabilities more than once, independently. This, together with the wide prevalence of TPST-dependent RGP in extant Rhizobiales commensals, implies a benefit for bacteria in promoting host root growth, which has given positive selection on this trait. Whether this bacterial pathway actively aims to manipulate host growth or is passively utilized by host to alter their root growth remains to be addressed. A growing collection of microbial strains from a wide range of plant species, including those from an aquatic alga *Chlamydomonas reinhardtii* (Durán et al., 2021), provides a useful tool to address this question. Moreover, identification of bacterial genes responsible for RGP phenotype is of crucial importance to genetically test the impact of this trait on plant and host fitness. We here provide the basis to better understand the molecular dialog between *A. thaliana* roots and Rhizobiales commensals to genetically address the relevance of microbiota-influenced regulation of plant growth and development.

Methods

Plant materials and growth conditions

We used *A. thaliana* wild-type accession Col-0 (CS60000) and commensal bacterial strains isolated from *A. thaliana* grown in natural soils, as described previously (Bai et al., 2015; Garrido-Oter et al., 2018). Plants were grown under short-day conditions (10 h under light at 21°C and 14 h under dark at 19°C) in a square plate (120x120 mm^2) placed vertically within a climate-controlled light cabinet (MLR-352, PHCbi).

Bacterial culture

We used TY media (5 g/L tryptone, 3 g/L yeast extract, 10 mM CaCl$_2$) to grow R129_E, R13_A, Root491, and Root142, and half strength of Tryptic Soy Broth media (15 g/L TSB; Sigma-Aldrich) to grow Root1497 and Root700. Bacterial cells were stored in glycerol stocks (16% v/v) at -80°C and recovered on TY supplemented with 1.8% agar, or 50% TSB supplemented with 1.5% agar, and cultivated at 28°C under dark conditions.

Inoculation of plants with bacteria, root length quantification, and bacterial abundance quantification

Inoculation of plants with Alphaproteobacteria was performed as described previously (Garrido-Oter et al., 2018). Briefly, we cultivated bacteria in liquid TY or 50% TSB media for 2 days at 28°C with 180 rpm agitation and mixed bacterial cells into half-strength of Murashige and Skoog media (750 mM MgSO$_4$, 625 mM KH$_2$PO$_4$, 10.3 mM NH$_4$NO$_3$, 9.4 mM KNO$_3$, 1.5 mM CaCl$_2$, 55 nM CoCl$_2$, 53 nM CuCl$_2$, 50 mM H$_3$BO$_3$, 2.5 mM KI, 50 mM MnCl$_2$, 520 nM Na$_2$MoO$_4$, 15 mM ZnCl$_2$, 75 mM Fe-EDTA, 500 mM MES-KOH pH 5.5) supplemented with 1% Bacto Agar (Difco). Bacterial titer (OD$_{600}$) was adjusted to 0.00005 by mixing 5 µl of bacterial suspension (OD$_{600}$ = 0.5 in 10 mM MgCl$_2$) into 50 ml of MS media pre-cooled to around 40-50°C. *A. thaliana* seeds were directly placed onto bacterium-containing media or precultured on half-strength of Murashige and Skoog media with vitamin (Duchefa) supplemented with 1% Bacto Agar for 1 week and before seedlings were transferred to the bacterium-containing plates. The same media was supplemented with 1 µM flg22 (QRLSTGSRINSAKDDAAGLQIA; EZBiolab), 100 nM RGF1 [D-Y(SO$_3$H)-SNPGHHP-Hyp-RHN; Beijing Scilight Biotechnology], or dRGF1 [DYSNPGHHP-Hyp-RHN] when necessary. Plates were then recorded by a scanner for root growth quantification using Fiji (Schindelin et al., 2012). Obtained data were square root transformed and fitted to

a linear mixed model using biological and technical replicates as random factors, using

the lmer function of the *lme4* R package. Fitted data was diagnosed to ensure the

normal distribution of residuals and used for Student's *t* test followed by correction for

multiple comparison based on the Benjamini-Hochberg method. For bacterial biomass

quantification, roots were thoroughly washed twice in sterile water to avoid residual

agar pieces and frozen in a 2 ml screw-cap tubes with 1-mm zirconia beads in liquid

nitrogen. Quantitative PCR was performed using the genomic DNA extracted from

roots by DNeasy Plant Mini Kit (QIAGEN) and primer sets specific to *16S* rRNA gene

sequences of *A. thaliana* (5'-CAGGCGGTGGAAACTACCAAG-3' and 5'-

TACAGCACTGCACGGGTCGAT-3') and R129_E (5'-

CGAGCTAATCTCCAAAAGCCATC-3' and 5'-TGACCCTACCGTGGTTAGCTG-3')

with iQ SYBR Green Supermix (BIO-RAD), as described previously (Garrido-Oter et

al., 2018).

RNA extraction and sequencing

For transcriptomic analysis of roots inoculated with Alphaproteobacteria commensals,

roots inoculated with bacteria for 12 days, as described above, were harvested and

immediately frozen in liquid nitrogen. Roots from three plates (approximately 25

plants) were pooled into one sample. Three full-factorial biological replicates from

three fully independent experiments were prepared for each condition. Roots were

then homogenized with 1-mm zirconia beads (Carl Roth) and Precellys24 (6,300 rpm

for 30 sec, 2 times with 15-sec intervals; Bertin), and RNA was extracted and purified

with the RNeasy Plant Mini kit (QIAGEN). Quality control, library preparation, and

sequencing (on the Illumina HiSeq 3000 platform) were conducted at the Max Planck

Genome Centre (MPGC), Cologne. Approximately 20,000,000 reads (150-bp single-

end) were obtained per sample.

For transcriptomic analysis of roots inoculated with R129_E in the presence of flg22,

plants were inoculated as described above. Plants from one plate were pooled into

one sample (approximately 10 plants) and three independent samples were obtained

per condition. Two fully independent experiments were conducted and six samples

per condition were obtained. Quality control, library preparation, and sequencing (on

the Illumina HiSeq 3000 platform) were conducted at the Max Planck Genome Centre

(MPGC), Cologne. Approximately 6,000,000 reads (150-bp single-end) were obtained

per sample.

For transcriptomic analysis of roots inoculated with *B. phytofirmans* and *B. glumae*,

Col-0 plants were grown on vertical agarose plates with Long Ashton nutrient solution

adjusted for sulfate concentration of 750 µM (control) or 25 µM (low S) as described

before (Koprivova et al., 2019). Total RNA was isolated from 18-day-old (14 dpi) roots

and sequenced at the Genomics & Transcriptomics Laboratory (GTL) at Heinrich

Heine University, Düsseldorf. Approximately 40,000,000 reads (150-bp single-end)

were obtained per sample.

RNA-seq analysis

Obtained reads, as well as reads downloaded from the Sequence Read Archive (SRA)

database, were preprocessed by fastp (Chen et al., 2018) with default parameters and

aligned to *A. thaliana* Col-0 genome by HISAT2 (Kim et al., 2019) with default

parameters based on the latest gene annotation (Cheng et al., 2017). Reads per gene

were counted by featureCounts (Liao et al., 2014) with default parameters.

Subsequent statistical analyses were performed using R software (https://www.r-

project.org/) unless described otherwise. Differential expression analysis was

performed using the *edgeR* (Robinson et al., 2010) package. Library size was

normalized by the weighted trimmed mean of M-values (TMM) method, and

normalized read counts were fitted to a generalized linear model (GLM) with a negative

binomial distribution to identify significantly differentially expressed genes (DEGs).

Zero-centered expression values were calculated based on $\log_2$ counts per million

(log2cpm) by subtracting the mean for each gene across samples. For comparison

between datasets, TMM normalization, GLM fitting, and zero-centered expression

values were calculated within each dataset. Zero-centered expression values were

used to compute *k*-means clustering, where *k* was decided based on Bayesian

Information Criterion (BIC). The similarity between samples was inferred by computing

Pearson's Correlation Coefficients (PCC) based on zero-centered expression values

of DEGs (n = 1,439). Distance of PCCs to 1 was used as dissimilarity indices to

perform principal coordinates analysis (PCoA) by cmdscale function. GO enrichment

analysis was performed by Metascape online software (Zhou et al., 2019).

Transcription binding motif enrichment analysis was performed using the ame function

implemented in the MEME Suite (McLeay and Bailey, 2010). Gene regulatory network

inference was performed using the *GENIE3* package (Huynh-Thu et al., 2010) with

default parameters. We compared responses to different microbes by computing

logFC within each dataset and use it to compute cosine similarity by the cosine

function in the packge *lsa* and performed *k*-means clustering, whose *k* was determined

based on the Bayesian Information Criterion. Distance of cosine similarities to 1 was

used as dissimilarity indices to perform PCoA and CAP. Permutational analysis of

variance (PERMANOVA) and pairwise PERMANOVA were performed by the adonis2

and pairwise.perm.manova functions of the packages *vegan* and *RVAideMemoire*

packages. All plots were visualized by Metascape or by the *ggplot2*, *ggtree*, *patchwork*,

and *igraph* packages in R.

Single-cell RNA-seq analysis

T-distributed Stochastic Neighbor Embedding (t-SNE) and representation of cluster-

wise expression levels as a violin plot were created by The Plant Single Cell RNA-

Sequencing   Database   (https://www.zmbp-resources.uni-

tuebingen.de/timmermans/plant-single-cell-browser/). To perform co-expression

analysis, read count data of 27,629 genes from 4,727 cells was retrieved from the

NCBI Sequence Read Archive (SRA) under the accession number GSE123818,

followed by global-scaling normalization by the NormalizeData function in the *Seurat*

package (Hao et al., 2020). We then filtered for 22,017 highly variable genes and their

expression values were scaled and centered by the FindVariableGenes and

ScaleData functions. Normalized and scaled expression values were used to compute

mutual ranks as a co-expression index for all pairs of genes as described previously

(Obayashi and Kinoshita, 2009). The top 100 genes co-expressed with *ANAC060*

were selected based on the mutual ranks and their GO enrichment was analyzed by

Metascape.

Confocal microscopy

Root tips (1-2 cm) were excised on agar plates and mounted onto a slide glass with

Milli-Q water or 10 μg/ml propidium iodide (PI). PI and GFP fluorescence were

observed by a SP8 FALCON-DIVE confocal microscope (Leica) using the 2-photon

excitation at XXX nm for GFP and 1,045 nm for PI excitation and emission at XXX-

XXX for GFP and XXX-XXX nm for PI, or by a LSM780 confocal microscope (Zeiss)

using the 488-nm line of a 25-mW Argon ion laser for GFP.

For EdU staining, roots of seed-inoculated plants at 21 dpi were incubated with

EdU for two hours directly on agar plates, followed by fixation with 2% formaldehyde,

0.3% glutaraldehyde, 0.1% Triton X-100, 10 mM EGTA, and 5 mM $MgSO_4$ in 50 mM

sodium phosphate buffer (pH 7.2) for 1 hour, washing with phospho-buffered saline,

and staining with Alexa488-azide using the Click-iT EdU Cell Proliferation Kit (Thermo

Fisher). Stained roots were observed under a LSM780 confocal microscope (Zeiss)

using the 488-nm line of a 25-mW Argon ion laser.

All acquired images were processed and adjusted (equally within each

experiment) by Fiji (Schindelin et al., 2012).

Data and code availability

Raw sequencing reads obtained in this study are deposited to the European

Nucleotide Archive (ENA) under the accession numbers of PRJEB45043

(Alphaproteobacteria), PRJEB45044 (R129_E with flg22), and PRJEB45045

(*Burkholderia*). The scripts for the computational analyses described in this study are

available at https://github.com/rtnakano1984/RNAseq_Alphaproteobacteria and

https://github.com/rtnakano1984/RNAseq_compare to ensure replicability and

reproducibility of these results.

Acknowledgments

We would like to thank Paul Schulze-Lefert at the Max Planck Institute for Plant

Breeding Research for providing infrastructures and helpful advice for this study, Jiri

Friml at the Institute of Science and Technology Austria, Niko Geldner at the University

of Lausanne, Woei-Jiun Guo at the National Cheng Kung University, Yoshikatsu

Matsubayashi and Hidefumi Shinohara at the Nagoya University, Birgit Kemmerling at

the University of Tübingen, Jia Li at the Lanzhou University, Margret Sauter at the

University of Kiel, and Bruno Müller at the University of Zurich for providing *A. thaliana*

mutant and transgenic seeds, Bruno Hüttel and Mark Holstein at the MPGC and

Patrick Petzsch and Karl-Erich Köhler at the GTL for sequencing, and Christina Phillip

and Petra Köchner for their technical support. We also would like to thank Tatsuya

Nobori, Kenichi Tsuda, Stephane Hacquard, and Rozina Kardakaris for their critical

reading and help editing the manuscript. This work was supported by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under the Priority

Programme "2125 Deconstruction and Reconstruction of Plant Microbiota

(DECRyPT)" funded to RTN (402201269) and SK (401836049). This work was also

supported by the Cluster of Excellence on Plant Sciences (CEPLAS) funded by DFG

to TG, SK and GWK. The authors declare no conflicts of interest. In memory of Alina

Kuroczik, who provided technical support for this study.

Author contribution

RTN designed the research and performed experiments. JH, TG, and AK performed experiments partly under supervision of RTN and SK. PS, MNC and GWK contributed to the RNAseq data analysis. RTN analyzed and interpreted the data. RTN wrote the paper with inputs from the other authors.

References

Achard, P., Gusti, A., Cheminant, S., Alioua, M., Dhondt, S., Coppens, F., Beemster, G.T., and Genschik, P. (2009). Gibberellin signaling controls cell proliferation rate in Arabidopsis. Curr Biol **19,** 1188-1193.

Aida, M., Beis, D., Heidstra, R., Willemsen, V., Blilou, I., Galinha, C., Nussaume, L., Noh, Y.S., Amasino, R., and Scheres, B. (2004). The PLETHORA genes mediate patterning of the Arabidopsis root stem cell niche. Cell **119,** 109-120.

Amano, Y., Tsubouchi, H., Shinohara, H., Ogawa, M., and Matsubayashi, Y. (2007). Tyrosine-sulfated glycopeptide involved in cellular proliferation and expansion in Arabidopsis. Proc Natl Acad Sci U S A **104,** 18333-18338.

Arnaud, D., Deeks, M.J., and Smirnoff, N. (2021). Differences between apoplastic and cytosolic reactive oxygen species production in Arabidopsis during pattern-triggered immunity. bioRxiv**,** 2021.2004.2020.440614.

Bai, Y., Muller, D.B., Srinivas, G., Garrido-Oter, R., Potthoff, E., Rott, M., Dombrowski, N., Munch, P.C., Spaepen, S., Remus-Emsermann, M., Huttel, B., McHardy, A.C., Vorholt, J.A., and Schulze-Lefert, P. (2015). Functional overlap of the Arabidopsis leaf and root microbiota. Nature **528,** 364-369.

Berg, G., Rybakova, D., Fischer, D., Cernava, T., Verges, M.C., Charles, T., Chen, X., Cocolin, L., Eversole, K., Corral, G.H., Kazou, M., Kinkel, L., Lange, L., Lima, N., Loy, A., Macklin, J.A., Maguin, E., Mauchline, T., McClure, R., Mitter, B., Ryan, M., Sarand, I., Smidt, H., Schelkle, B., Roume, H., Kiran, G.S., Selvin, J., Souza, R.S.C., van Overbeek, L., Singh, B.K., Wagner, M., Walsh, A., Sessitsch, A., and Schloter, M. (2020). Microbiome definition re-visited: old concepts and new challenges. Microbiome **8,** 103.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., Ver Loren van Themaat, E., and Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. Annu Rev Plant Biol **64,** 807-838.

Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., Rauf, P., Huettel, B., Reinhardt, R., Schmelzer, E., Peplies, J., Gloeckner, F.O., Amann, R., Eickhorst, T., and Schulze-Lefert, P. (2012). Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature **488,** 91-95.

Castrillo, G., Teixeira, P.J., Paredes, S.H., Law, T.F., de Lorenzo, L., Feltcher, M.E., Finkel, O.M., Breakfield, N.W., Mieczkowski, P., Jones, C.D., Paz-Ares, J., and Dangl, J.L. (2017). Root microbiota drive direct integration of phosphate stress and immunity. Nature **543,** 513-518.

Cederholm, H.M., and Benfey, P.N. (2015). Distinct sensitivities to phosphate deprivation suggest that RGF peptides play disparate roles in Arabidopsis thaliana root development. New Phytol **207,** 683-691.

Chaiwanon, J., and Wang, Z.Y. (2015). Spatiotemporal brassinosteroid signaling and
antagonism with auxin pattern stem cell dynamics in Arabidopsis roots. Curr
Biol **25,** 1031-1042.
Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics **34,** i884-i890.
Chen, T., Nomura, K., Wang, X., Sohrabi, R., Xu, J., Yao, L., Paasch, B.C., Ma, L.,
Kremer, J., Cheng, Y., Zhang, L., Wang, N., Wang, E., Xin, X.F., and He,
S.Y. (2020). A plant genetic network for preventing dysbiosis in the
phyllosphere. Nature **580,** 653-657.
Cheng, C.Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and
Town, C.D. (2017). Araport11: a complete reannotation of the Arabidopsis
thaliana reference genome. Plant J **89,** 789-804.
Dello Ioio, R., Nakamura, K., Moubayidin, L., Perilli, S., Taniguchi, M., Morita,
M.T., Aoyama, T., Costantino, P., and Sabatini, S. (2008). A genetic
framework for the control of cell division and differentiation in the root meristem.
Science **322,** 1380-1384.
Denyer, T., Ma, X., Klesen, S., Scacchi, E., Nieselt, K., and Timmermans, M.C.P.
(2019). Spatiotemporal Developmental Trajectories in the Arabidopsis Root
Revealed Using High-Throughput Single-Cell RNA Sequencing. Dev Cell **48,**
840-852 e845.
Di Mambro, R., De Ruvo, M., Pacifici, E., Salvi, E., Sozzani, R., Benfey, P.N.,
Busch, W., Novak, O., Ljung, K., Di Paola, L., Maree, A.F.M., Costantino,
P., Grieneisen, V.A., and Sabatini, S. (2017). Auxin minimum triggers the
developmental switch from cell division to cell differentiation in the Arabidopsis
root. Proc Natl Acad Sci U S A **114,** E7641-E7649.
Dixon, D.P., Skipsey, M., and Edwards, R. (2010). Roles for glutathione transferases
in plant secondary metabolism. Phytochemistry **71,** 338-350.
Dixon, D.P., Hawkins, T., Hussey, P.J., and Edwards, R. (2009). Enzyme activities
and subcellular localization of members of the Arabidopsis glutathione
transferase superfamily. J Exp Bot **60,** 1207-1218.
Doblas, V.G., Smakowska-Luzan, E., Fujita, S., Alassimone, J., Barberon, M.,
Madalinski, M., Belkhadir, Y., and Geldner, N. (2017). Root diffusion barrier
control by a vasculature-derived peptide binding to the SGN3 receptor. Science
355, 280-284.
Duran, P., Thiergart, T., Garrido-Oter, R., Agler, M., Kemen, E., Schulze-Lefert,
P., and Hacquard, S. (2018). Microbial Interkingdom Interactions in Roots
Promote Arabidopsis Survival. Cell **175,** 973-983 e914.
Durán, P., Flores-Uribe, J., Wippel, K., Zhang, P., Guan, R., and Garrido-Oter, R.
(2021). Characterization of the Chlamydomonas reinhardtii phycosphere
reveals conserved features of the plant microbiota. bioRxiv,
2021.2003.2004.433956.
Eulgem, T., and Somssich, I.E. (2007). Networks of WRKY transcription factors in
defense signaling. Curr Opin Plant Biol **10,** 366-371.
Finkel, O.M., Salas-Gonzalez, I., Castrillo, G., Conway, J.M., Law, T.F., Teixeira,
P., Wilson, E.D., Fitzpatrick, C.R., Jones, C.D., and Dangl, J.L. (2020). A
single bacterial genus maintains root growth in a complex microbiome. Nature
587, 103-108.
Furumizu, C., Krabberød, A.K., Hammerstad, M., Alling, R.M., Wildhagen, M.,
Sawa, S., and Aalen, R.B. (2021). The Sequenced Genomes of Non-Flowering

Land Plants Reveal the (R)Evolutionary History of Peptide Signaling. bioRxiv, 2020.2006.2002.130120.

Galinha, C., Hofhuis, H., Luijten, M., Willemsen, V., Blilou, I., Heidstra, R., and Scheres, B. (2007). PLETHORA proteins as dose-dependent master regulators of Arabidopsis root development. Nature **449,** 1053-1057.

Garrido-Oter, R., Nakano, R.T., Dombrowski, N., Ma, K.W., AgBiome, T., McHardy, A.C., and Schulze-Lefert, P. (2018). Modular Traits of the Rhizobiales Root Microbiota and Their Evolutionary Relationship with Symbiotic Rhizobia. Cell Host Microbe **24,** 155-167 e155.

Ghorbani, S., Lin, Y.C., Parizot, B., Fernandez, A., Njo, M.F., Van de Peer, Y., Beeckman, T., and Hilson, P. (2015). Expanding the repertoire of secretory peptides controlling root development with comparative genome analysis and functional assays. J Exp Bot **66,** 5257-5269.

Gomez-Gomez, L., Felix, G., and Boller, T. (1999). A single locus determines sensitivity to bacterial flagellin in Arabidopsis thaliana. Plant J **18,** 277-284.

Hacquard, S., Garrido-Oter, R., Gonzalez, A., Spaepen, S., Ackermann, G., Lebeis, S., McHardy, A.C., Dangl, J.L., Knight, R., Ley, R., and Schulze-Lefert, P. (2015). Microbiota and Host Nutrition across Plant and Animal Kingdoms. Cell Host Microbe **17,** 603-616.

Hacquard, S., Kracher, B., Hiruma, K., Munch, P.C., Garrido-Oter, R., Thon, M.R., Weimann, A., Damm, U., Dallery, J.F., Hainaut, M., Henrissat, B., Lespinet, O., Sacristan, S., Ver Loren van Themaat, E., Kemen, E., McHardy, A.C., Schulze-Lefert, P., and O'Connell, R.J. (2016). Survival trade-offs in plant roots during colonization by closely related beneficial and pathogenic fungi. Nat Commun **7,** 11362.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zagar, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E.P., Jain, J., Srivastava, A., Stuart, T., Fleming, L.B., Yeung, B., Rogers, A.J., McElrath, J.M., Blish, C.A., Gottardo, R., Smibert, P., and Satija, R. (2020). Integrated analysis of multimodal single-cell data. bioRxiv, 2020.2010.2012.335331.

Harbort, C.J., Hashimoto, M., Inoue, H., Niu, Y., Guan, R., Rombola, A.D., Kopriva, S., Voges, M., Sattely, E.S., Garrido-Oter, R., and Schulze-Lefert, P. (2020). Root-Secreted Coumarins and the Microbiota Interact to Improve Iron Nutrition in Arabidopsis. Cell Host Microbe **28,** 825-837 e826.

Hartmann, M., Zeier, T., Bernsdorff, F., Reichel-Deland, V., Kim, D., Hohmann, M., Scholten, N., Schuck, S., Brautigam, A., Holzel, T., Ganter, C., and Zeier, J. (2018). Flavin Monooxygenase-Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic Immunity. Cell **173,** 456-469 e416.

Hiruma, K., Gerlach, N., Sacristan, S., Nakano, R.T., Hacquard, S., Kracher, B., Neumann, U., Ramirez, D., Bucher, M., O'Connell, R.J., and Schulze-Lefert, P. (2016). Root Endophyte Colletotrichum tofieldiae Confers Plant Fitness Benefits that Are Phosphate Status Dependent. Cell **165,** 464-474.

Hou, S., Thiergart, T., Vannier, N., Mesny, F., Ziegler, J., Pickel, B., and Hacquard, S. (2020). Microbiota-root-shoot axis modulation by MYC2 favours Arabidopsis growth over defence under suboptimal light. bioRxiv, 2020.2011.2006.371146.

Hsu, F.C., Chou, M.Y., Chou, S.J., Li, Y.R., Peng, H.P., and Shih, M.C. (2013). Submergence confers immunity mediated by the WRKY22 transcription factor in Arabidopsis. Plant Cell **25,** 2699-2713.

Huffaker, A., and Ryan, C.A. (2007). Endogenous peptide defense signals in
Arabidopsis differentially amplify signaling for the innate immune response.
Proc Natl Acad Sci U S A **104,** 10732-10736.
Huot, B., Yao, J., Montgomery, B.L., and He, S.Y. (2014). Growth-defense tradeoffs
in plants: a balancing act to optimize fitness. Mol Plant **7,** 1267-1287.
Huynh-Thu, V.A., Irrthum, A., Wehenkel, L., and Geurts, P. (2010). Inferring
regulatory networks from expression data using tree-based methods. PLoS
One **5**.
Kaufmann, C., and Sauter, M. (2019). Sulfated plant peptide hormones. J Exp Bot
70, 4267-4277.
Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat
Biotechnol **37,** 907-915.
Kim, H.J., Park, J.H., Kim, J., Kim, J.J., Hong, S., Kim, J., Kim, J.H., Woo, H.R.,
Hyeon, C., Lim, P.O., Nam, H.G., and Hwang, D. (2018). Time-evolving
genetic networks reveal a NAC troika that negatively regulates leaf senescence
in Arabidopsis. Proc Natl Acad Sci U S A **115,** E4930-E4939.
Komori, R., Amano, Y., Ogawa-Ohnishi, M., and Matsubayashi, Y. (2009).
Identification of tyrosylprotein sulfotransferase in Arabidopsis. Proc Natl Acad
Sci U S A **106,** 15067-15072.
Koprivova, A., Schuck, S., Jacoby, R.P., Klinkhammer, I., Welter, B., Leson, L.,
Martyn, A., Nauen, J., Grabenhorst, N., Mandelkow, J.F., Zuccaro, A., Zeier,
J., and Kopriva, S. (2019). Root-specific camalexin biosynthesis controls the
plant growth-promoting effects of multiple bacterial strains. Proc Natl Acad Sci
U S A **116,** 15735-15744.
Ladwig, F., Dahlke, R.I., Stuhrwohldt, N., Hartmann, J., Harter, K., and Sauter, M.
(2015). Phytosulfokine Regulates Growth in Arabidopsis through a Response
Module at the Plasma Membrane That Includes CYCLIC NUCLEOTIDE-
GATED CHANNEL17, H+-ATPase, and BAK1. Plant Cell **27,** 1718-1729.
Lallement, P.A., Brouwer, B., Keech, O., Hecker, A., and Rouhier, N. (2014). The
still mysterious roles of cysteine-containing glutathione transferases in plants.
Front Pharmacol **5,** 192.
Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general
purpose program for assigning sequence reads to genomic features.
Bioinformatics **30,** 923-930.
Lu, X., Shi, H., Ou, Y., Cui, Y., Chang, J., Peng, L., Gou, X., He, K., and Li, J. (2020).
RGF1-RGI1, a Peptide-Receptor Complex, Regulates Arabidopsis Root
Meristem Development via a MAPK Signaling Cascade. Mol Plant **13,** 1594-
1607.
Lundberg, D., Persson, I., and Ekberg, C. (2013). Crystal structure of [Eu(CyMe4-
BTBP)2kappa2O,O'-(NO3)](NO3)2.n-C8H17OH and its structure in 1-octanol
solution. Dalton Trans **42,** 3767-3770.
Ma, K.-W., Niu, Y., Jia, Y., Ordon, J., Copeland, C., Emonet, A., Geldner, N., Guan,
R., Stolze, S.C., Nakagami, H., Garrido-Oter, R., and Schulze-Lefert, P.
(2021). Coordination of microbe–host homeostasis by crosstalk with plant
innate immunity. Nature Plants.
Maier, B.A., Kiefer, P., Field, C.M., Hemmerle, L., Bortfeld-Miller, M.,
Emmenegger, B., Schafer, M., Pfeilmeier, S., Sunagawa, S., Vogel, C.M.,
and Vorholt, J.A. (2021). A general non-self response as part of plant immunity.
Nat Plants **7,** 696-705.

Matsubayashi, Y., Ogawa, M., Kihara, H., Niwa, M., and Sakagami, Y. (2006).
Disruption and overexpression of Arabidopsis phytosulfokine receptor gene
affects cellular longevity and potential for growth. Plant Physiol **142,** 45-53.
Matsuzaki, Y., Ogawa-Ohnishi, M., Mori, A., and Matsubayashi, Y. (2010).
Secreted peptide signals required for maintenance of root stem cell niche in
Arabidopsis. Science **329,** 1065-1067.
McLeay, R.C., and Bailey, T.L. (2010). Motif Enrichment Analysis: a unified
framework and an evaluation on ChIP data. BMC Bioinformatics **11,** 165.
Nobori, T., Cao, Y., Entila, F., Dahms, E., Tsuda, Y., Garrido-Oter, R., and Tsuda,
K. (2021). Dissecting the co-transcriptome landscape of plants and microbiota
members. bioRxiv, 2021.2004.2025.440543.
Noctor, G., Mhamdi, A., Chaouch, S., Han, Y., Neukermans, J., Marquez-Garcia,
B., Queval, G., and Foyer, C.H. (2012). Glutathione in plants: an integrated
overview. Plant Cell Environ **35,** 454-484.
Obayashi, T., and Kinoshita, K. (2009). Rank of correlation coefficient as a
comparable measure for biological significance of gene coexpression. DNA
Res **16,** 249-260.
Ou, Y., Lu, X., Zi, Q., Xun, Q., Zhang, J., Wu, Y., Shi, H., Wei, Z., Zhao, B., Zhang,
X., He, K., Gou, X., Li, C., and Li, J. (2016). RGF1 INSENSITIVE 1 to 5, a
group of LRR receptor-like kinases, are essential for the perception of root
meristem growth factor 1 in Arabidopsis thaliana. Cell Res **26,** 686-698.
Paasch, B.C., and He, S.Y. (2021). Toward understanding microbiota homeostasis in
the plant kingdom. PLoS Pathog **17,** e1009472.
Phukan, U.J., Jeena, G.S., and Shukla, R.K. (2016). WRKY Transcription Factors:
Molecular Regulation and Stress Responses in Plants. Front Plant Sci **7,** 760.
Pislewska-Bednarek, M., Nakano, R.T., Hiruma, K., Pastorczyk, M., Sanchez-
Vallet, A., Singkaravanit-Ogawa, S., Ciesiolka, D., Takano, Y., Molina, A.,
Schulze-Lefert, P., and Bednarek, P. (2018). Glutathione Transferase U13
Functions in Pathogen-Triggered Glucosinolate Metabolism. Plant Physiol **176,**
538-551.
Pruitt, R.N., Joe, A., Zhang, W., Feng, W., Stewart, V., Schwessinger, B., Dinneny,
J.R., and Ronald, P.C. (2017). A microbially derived tyrosine-sulfated peptide
mimics a plant peptide hormone. New Phytol **215,** 725-736.
Pruitt, R.N., Schwessinger, B., Joe, A., Thomas, N., Liu, F., Albert, M., Robinson,
M.R., Chan, L.J., Luu, D.D., Chen, H., Bahar, O., Daudi, A., De
Vleesschauwer, D., Caddell, D., Zhang, W., Zhao, X., Li, X., Heazlewood,
J.L., Ruan, D., Majumder, D., Chern, M., Kalbacher, H., Midha, S., Patil, P.B.,
Sonti, R.V., Petzold, C.J., Liu, C.C., Brodbelt, J.S., Felix, G., and Ronald,
P.C. (2015). The rice immune receptor XA21 recognizes a tyrosine-sulfated
protein from a Gram-negative bacterium. Sci Adv **1,** e1500245.
Pucciariello, C., Parlanti, S., Banti, V., Novi, G., and Perata, P. (2012). Reactive
oxygen species-driven transcription in Arabidopsis under oxygen deprivation.
Plant Physiol **159,** 184-196.
Rajniak, J., Barco, B., Clay, N.K., and Sattely, E.S. (2015). A new cyanogenic
metabolite in Arabidopsis required for inducible pathogen defence. Nature **525,**
376-379.
Rasul, S., Dubreuil-Maurizi, C., Lamotte, O., Koen, E., Poinssot, B., Alcaraz, G.,
Wendehenne, D., and Jeandroz, S. (2012). Nitric oxide production mediates
oligogalacturonide-triggered immunity and resistance to Botrytis cinerea in
Arabidopsis thaliana. Plant Cell Environ **35,** 1483-1499.

Rich-Griffin, C., Eichmann, R., Reitz, M.U., Hermann, S., Woolley-Allen, K., Brown, P.E., Wiwatdirekkul, K., Esteban, E., Pasha, A., Kogel, K.H., Provart, N.J., Ott, S., and Schafer, P. (2020). Regulation of Cell Type-Specific Immunity Networks in Arabidopsis Roots. Plant Cell **32,** 2742-2762.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics **26,** 139-140.

Salas-Gonzalez, I., Reyt, G., Flis, P., Custodio, V., Gopaulchan, D., Bakhoum, N., Dew, T.P., Suresh, K., Franke, R.B., Dangl, J.L., Salt, D.E., and Castrillo, G. (2021). Coordination between microbiota and root endodermis supports plant mineral nutrient homeostasis. Science **371,** eabd0695.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., and Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods **9,** 676-682.

Shao, H., Wang, H., and Tang, X. (2015). NAC transcription factors in plant multiple abiotic stress responses: progress and prospects. Front Plant Sci **6,** 902.

Shao, Y., Yu, X., Xu, X., Li, Y., Yuan, W., Xu, Y., Mao, C., Zhang, S., and Xu, J. (2020). The YDA-MKK4/MKK5-MPK3/MPK6 Cascade Functions Downstream of the RGF1-RGI Ligand-Receptor Pair in Regulating Mitotic Activity in Root Apical Meristem. Mol Plant **13,** 1608-1623.

Shinohara, H. (2021). Root meristem growth factor RGF, a sulfated peptide hormone in plants. Peptides **142,** 170556.

Shiu, S.H., Karlowski, W.M., Pan, R., Tzeng, Y.H., Mayer, K.F., and Li, W.H. (2004). Comparative analysis of the receptor-like kinase family in Arabidopsis and rice. Plant Cell **16,** 1220-1234.

Song, W., Liu, L., Wang, J., Wu, Z., Zhang, H., Tang, J., Lin, G., Wang, Y., Wen, X., Li, W., Han, Z., Guo, H., and Chai, J. (2016). Signature motif-guided identification of receptors for peptide hormones essential for root meristem growth. Cell Res **26,** 674-685.

Teixeira, P., Colaianni, N.R., Law, T.F., Conway, J.M., Gilbert, S., Li, H., Salas-Gonzalez, I., Panda, D., Del Risco, N.M., Finkel, O.M., Castrillo, G., Mieczkowski, P., Jones, C.D., and Dangl, J.L. (2021). Specific modulation of the root immune system by a community of commensal bacteria. Proc Natl Acad Sci U S A **118,** e2100678118.

Thiergart, T., Duran, P., Ellis, T., Vannier, N., Garrido-Oter, R., Kemen, E., Roux, F., Alonso-Blanco, C., Agren, J., Schulze-Lefert, P., and Hacquard, S. (2020). Root microbiota assembly and adaptive differentiation among European Arabidopsis populations. Nat Ecol Evol **4,** 122-131.

Tost, A.S., Kristensen, A., Olsen, L.I., Axelsen, K.B., and Fuglsang, A.T. (2021). The PSY Peptide Family-Expression, Modification and Physiological Implications. Genes (Basel) **12**.

Tsukagoshi, H., Busch, W., and Benfey, P.N. (2010). Transcriptional regulation of ROS controls transition from proliferation to differentiation in the root. Cell **143,** 606-616.

Ubeda-Tomas, S., Federici, F., Casimiro, I., Beemster, G.T., Bhalerao, R., Swarup, R., Doerner, P., Haseloff, J., and Bennett, M.J. (2009). Gibberellin signaling in the endodermis controls Arabidopsis root meristem size. Curr Biol **19,** 1194-1199.

Wagner, M.R., Lundberg, D.S., Del Rio, T.G., Tringe, S.G., Dangl, J.L., and
Mitchell-Olds, T. (2016). Host genotype and age shape the leaf and root
microbiomes of a wild perennial plant. Nat Commun **7,** 12151.
Wang, L., Tsuda, K., Sato, M., Cohen, J.D., Katagiri, F., and Glazebrook, J. (2009).
Arabidopsis CaM binding protein CBP60g contributes to MAMP-induced SA
accumulation and is involved in disease resistance against Pseudomonas
syringae. PLoS Pathog **5,** e1000301.
Wang, S., Meade, A., Lam, H.M., and Luo, H. (2020). Evolutionary Timeline and
Genomic Plasticity Underlying the Lifestyle Diversity in Rhizobiales. mSystems
5, e00438-00420.
Wei, Z., Yang, T., Friman, V.P., Xu, Y., Shen, Q., and Jousset, A. (2015). Trophic
network architecture of root-associated bacterial communities determines
pathogen invasion and plant health. Nat Commun **6,** 8413.
Wippel, K., Tao, K., Niu, Y., Zgadzaj, R., Guan, R., Dahms, E., Zhang, P., Jensen,
D.B., Logemann, E., Radutoiu, S., Schulze-Lefert, P., and Garrido-Oter, R.
(2021). Host preference and invasiveness of commensals in the Lotus and
Arabidopsis root microbiota. bioRxiv, 2021.2001.2012.426357.
Yamada, M., Han, X., and Benfey, P.N. (2020). RGF1 controls root meristem size
through ROS signalling. Nature **577,** 85-88.
Yu, B., Wang, Y., Zhou, H., Li, P., Liu, C., Chen, S., Peng, Y., Zhang, Y., and Teng,
S. (2020). Genome-wide binding analysis reveals that ANAC060 directly
represses sugar-induced transcription of ABI5 in Arabidopsis. Plant J **103,** 965-
979.
Yu, K., Liu, Y., Tichelaar, R., Savant, N., Lagendijk, E., van Kuijk, S.J.L., Stringlis,
I.A., van Dijken, A.J.H., Pieterse, C.M.J., Bakker, P., Haney, C.H., and
Berendsen, R.L. (2019). Rhizosphere-Associated Pseudomonas Suppress
Local Root Immune Responses by Gluconic Acid-Mediated Lowering of
Environmental pH. Curr Biol **29,** 3913-3920 e3914.
Yu, P., He, X., Baer, M., Beirinckx, S., Tian, T., Moya, Y.A.T., Zhang, X.,
Deichmann, M., Frey, F.P., Bresgen, V., Li, C., Razavi, B.S., Schaaf, G., von
Wiren, N., Su, Z., Bucher, M., Tsuda, K., Goormachtig, S., Chen, X., and
Hochholdinger, F. (2021). Plant flavones enrich rhizosphere
Oxalobacteraceae to improve maize performance under nitrogen deprivation.
Nat Plants **7,** 481-499.
Yuan, X., Wang, H., Cai, J., Li, D., and Song, F. (2019). NAC transcription factors in
plant immunity. Phytopathology Research **1,** 3.
Zamioudis, C., Mastranesti, P., Dhonukshe, P., Blilou, I., and Pieterse, C.M.
(2013). Unraveling root developmental programs initiated by beneficial
Pseudomonas spp. bacteria. Plant Physiol **162,** 304-318.
Zgadzaj, R., Garrido-Oter, R., Jensen, D.B., Koprivova, A., Schulze-Lefert, P., and
Radutoiu, S. (2016). Root nodule symbiosis in Lotus japonicus drives the
establishment of distinctive rhizosphere, root, and nodule bacterial
communities. Proc Natl Acad Sci U S A **113,** E7996-E8005.
Zhou, W., Wei, L., Xu, J., Zhai, Q., Jiang, H., Chen, R., Chen, Q., Sun, J., Chu, J.,
Zhu, L., Liu, C.M., and Li, C. (2010). Arabidopsis Tyrosylprotein
sulfotransferase acts in the auxin/PLETHORA pathway in regulating
postembryonic maintenance of the root stem cell niche. Plant Cell **22,** 3692-
3709.

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O.,
Benner, C., and Chanda, S.K. (2019). Metascape provides a biologist-oriented
resource for the analysis of systems-level datasets. Nat Commun **10,** 1523.
1107

1108

Tables

Figure and table legends

Figure 1. Alphaproteobacteria commensals trigger qualitatively similar
transcriptional responses in host roots.

(A) A principal coordinates analysis separated transcriptome of axenic roots
(asterisks), roots inoculated with Rhizobiales commensals (closed shapes), and roots
inoculated with sister lineage isolates (open shapes). (B) A heatmap showing cosine
similarities between logFC upon Alphaproteobacteria commensal inoculations. (C)
Heatmaps showing $\log_2$-scale count per million (log2cpm), zero-centered expression
values, $\log_2$-scale fold changes (logFC), and significance (FDR < 0.05; |logFC| > 1),
aligned according to their k-means cluster assignment (k = 9, based on Bayesian
information criterion). (D) Boxplots showing logFC of each k-means cluster. Asterisks
indicate statistical significance corresponding to Student's one-sample t test after
Bonferroni's correction for multiple comparisons (μ = 0; α = 0.001) and letters indicate
statistical significance corresponding to pairwise Wilcoxon's rank sum test after
Bonferroni's correction for multiple comparisons within each cluster (α = 0.05). (E) A
heatmap showing enrichment of GO terms in the k-means clusters. (F) A network plot
of enriched GO terms. Each node represents each GO term, and GOs with
overlapping gene contents are connected with edges. k-means assignment of
composing genes is shown as a pie chart. We selected the terms with the best P
values from each of the 9 clusters, with the constraint that there are no more than 15
terms per cluster and no more than 250 terms in total. Representative terms are
manually curated and indicated with composing k-means clusters for each GO cluster.

1132

1133 **Figure 2. WRKY and ANAC transcription factors are important for transcriptional**
1134 **reprogramming.**

1135 (A) Heatmaps showing mean logFC of each cluster (left) and the enrichment of
1136 corresponding transcription factor binding motifs in their promoter regions (1-kb
1137 upstream from transcription initiation sites). AHL, AT-HOOK MOTIF CONTAINING

NUCLEAR LOCALIZED; CCHC, CCHC-type zinc-finger motif containing protein; ATHB, ARABIDOPSIS THALIANA HOMEOBOX; FRS9, FAR1-RELATED SEQUENCE9. (B) GENIE3-inferred gene regulatory network using WRKY (green) and ANAC (magenta) family DETFs as potential regulators (colored; bigger node size). Arbitrarily defined modules are shown by dotted lines. (C) Degree, closeness, and betweenness centralities were computed based on the network in (B) for all potential regulators. Green and magenta dots correspond to WRKY and ANAC family TFs.

Figure 3. Non-pathogenic microbial inoculation triggered similar transcriptional responses in the host roots.

(A) A heatmap showing cosine similarities between logFC upon different microbial inoculations. DEGs were selected based on the inoculation in rows (numbers of DEGs are shown in parentheses), whose logFC cosine similarity was calculated in comparison with inoculations in columns. For each row, all cosine similarity values were plotted on the right, whose colors indicate the lifestyle of the comparing inoculation. (B) Cosine similarities within non-pathogenic inoculations and within pathogenic inoculations were higher than those between pathogenic and non-pathogenic inoculations. (C) Principal coordinates analysis based on logFC cosine similarities of all commonly quantified genes (n = 21,815). Colors correspond to lifestyles. (D) Constrained analysis of principal coordinates (CAP) based on the logFC cosine similarities, using the constraining and conditioning factors indicated above. Colors correspond to inoculated microbial strains. (E) A heatmap showing logFC of genes assigned with the GO categories related to detoxification and response to hydrogen peroxide, reactive oxygen species, and hypoxia. Inoculations are aligned according to the dendrogram in (A) and genes are aligned according to k-means clustering as shown in Supplemental Figure 2B. Numbers indicated in the names of inoculation indicate the time point of harvesting (dpi).

Figure 4. Rhizobiales-specific DEGs are potentially regulated by ANAC060.

(A) Heatmaps showing log2cpm, zero-centered expression values, logFC, significance, and GO assignment of genes specifically regulated by Rhizobiales commensals but not by sister lineage isolates. (B) Enlarged view of the GENIE3-inferred GRN, shown in Figure 2B, colored for potential regulator TFs (left) or

1171   Rhizobiales-specific DEGs (right). (C) Same view as in (B), colored by their logFC

1172   upon different bacterial inoculation.

1173

Figure 5. Plant sulfated peptide pathway is needed for R129_E RGP activity.

(A) Primary root growth and representative pictures of Col-0 and *tpst* plants grown

axenically or inoculated with R129_E. Seeds were inoculated and grown under

continuous light conditions for 12 days. (B) Bacterial abundance relative to plant

biomass was quantified by DNA-based quantitative PCR using primers specific to *A.*

thaliana and R129_E *16S rRNA* genes. No clear depletion of bacterial cells from *tpst*

roots was observed. (C) Mutants impaired RGF1 perception (*rgi1;2,3;4;5-1*) or PSK

perception (*pskr1pskr2psy1r*) did not phenocopy *tpst* when 7-day-old seedlings were

inoculated with R129_E for 14 days under short-day conditions. (D) Col-0 root MZ

accumulate higher amounts of proPLT1::PLT1-GFP when 7-day-old seedlings were

inoculated with R129_E for 7 days under short-day conditions, whereas *tpst* root MZ

accumulated a similar amount irrespective of R129_E inoculation. For each panel, a

merged image of propidium iodide staining (gray) and PLT1-GFP (green) signals and

an image only showing PLT1-GFP signals pseudo-colored as shown in the calibration

bar are shown in left and right, respectively. Bars indicate 50 µm. Images are aligned

with quiescent centers on the dotted line. Letters indicate statistical significance (α =

0.05) corresponding to Student's *t* test after fitting to a linear mixed model with

biological and technical replicates as random factors, corrected for multiple

comparison by the Benjamini-Hochberg method.

1193

Figure 6. TPST is needed for the RGP activity explicitly exerted by Rhizobiales

RGP but not Pseudomonadales.

Seven-day-old seedling of Col-0 and *tpst* were inoculated with a panel of Rhizobiales

commensals (A) or Pseudomonadales commensals (B) and grown for 14 days under

short-day conditions. Letters indicate statistical significance (α = 0.05) corresponding

to Student's t test after fitting to a linear mixed model with biological and technical

replicates as random factors, corrected for multiple comparison by the Benjamini-

Hochberg method.

1202

1203   **Supplemental   Figure   1.   Qualitative   comparison   of   DEGs   for**

1204   **Alphaproteobacteria inoculations**

Upset diagrams showing the number of DEGs specific to or shared (intersect) between different inoculations. Up-regulated and down-regulated DEGs were separately analyzed.

Supplemental Figure 2. Similarity in gene contents between GO categories enriched in Alphaproteobacteria DEGs based on *k*-means clustering.

For each cluster, significant enriched GO categories are shown in rows. Genes in the respective clusters with at least one significantly enriched GO category are shown in columns. Assignment of a gene to a given GO category is indicated by black filling color. GO categories with highly similar gene contents are manually marked and representative terms are indicated. Bold R and B in the labels indicate "response to" and "biosynthesis of", respectively.

Supplemental Figure 3. WRKY54 and ANAC060 are associated with responses characteristic to sister lineage and Rhizobiales isolates, respectively.

The same GRN as in Figure 2B, colored by k-means clusters (A), by logFC upon Alphaproteobacteria commensals (B; the full view of Figure 3C), and for Rhizobiales-specific DEGs (C; the full view of Figure 3B). Module 1 and 2 contain genes from Clusters 1, 8, and 9 and Clusters 3, 4, and 6, respectively, and Rhizobiales DEGs were exclusively found in the module 2. Part of the network presented in Figures 4B and 4C are indicated by a square.

Supplemental Figure 4. Robust centrality of WRKY54 and ANAC060 irrespective of the lists of potential regulators.

GENIE3-based inference of gene regulatory network using all differentially expressed transcription factors as potential regulators, colored for transcription factor families (A), by k-means clusters, corresponding to Figure 1B (B), for Rhizobiales-specific DEGs, corresponding to Figure 3A (C) or by logFC (D). Modules are arbitrarily defined. (E) Centrality indices of potential regulators in the inferred GRN.

Supplemental Figure 5. Non-pathogenic microbial inoculation triggered similar transcriptional responses in the host roots.

(A) A heatmap showing logFC of genes differentially expressed at least in one of the analyzed treatments (n = 13,889), aligned according to *k*-means clustering (*k* = 21, based on Bayesian information criterion). Assignment of genes to *k*-means cluster for Alphaproteobacteria dataset (Figure 1) are also shown above. (B) GO enrichment analysis of k-means clustering shown in (A), as in Figure 1F. (C) A heatmap showing logFC of GST genes, aligned according to their phylogeny. Shaded by blue are the genes specifically acquired within the family Brassicaceae (Pisléska-Bednarek et al., 2018).

Supplemental Figure 6. ANAC060 is expressed in the cell population within the root meristematic zone, where PLETHORA TFs and RGF peptides are also expressed to control meristematic mitotic activity.

(A) T-distributed stochastic neighbor embedding (t-SNE) of single cell RNAseq datasets reported by Denyer and Me et al. (2019), where colors correspond to the normalized expression levels of genes indicated above. (B) Cluster assignment of ANAC060, PLT1, PLT2, and PLT4, where the cluster 11 corresponds to the meristematic cell population. t-SNE and violin plots were created by The Plant Single Cell RNA-Sequencing Database. (C) GO enrichment analysis of 100 genes best co-expressed with ANAC060 in the single cell RNAseq datasets, based on Pearson's correlation-based Mutual Ranks. Data are visualized as in Figure 1F.

Supplemental Figure 7. DNA synthesis and auxin and cytokinin accumulation in axenic and R129_E-inoculated roots.

(A) Roots incubated with EdU for 2 hours were staining with Alexa488 after seeds were inoculated with R129_E for 21 days under short-day conditions. EdU/Alexa488 signals in green was merged with autofluorescence signals in magenta. (B) Roots expressing DR5:ER-GFP were imaged after seeds were inoculated with R129_E for 8 days under short-day condition. (C) Roots expressing TSCn:ER-GFP were imaged after 7-day-old seedlings were inoculated with R129_E for 10 days under short-day conditions. Bars, 50 μm.

Supplemental Figure 8. RGF and CIF peptide pathways were not essential for R129_E RGP activity.

(A) R129_E promoted primary root growth of Col-0, *rgf123*, and *sgn3* plants after 7-

day-old seedlings were inoculated for 14 days under short-day conditions. (B) External

application of 100 nM RGF1 or 100 nM desulfated RGF1 did not affect the ability of

R129_E to promote root growth of Col-0 and did not rescue the lack of RGP in *tpst*

mutant roots. Letters indicate statistical significance (α = 0.05) corresponding to

Student's *t* test after fitting to a linear mixed model with biological and technical

replicates as random factors, corrected for multiple comparison by the Benjamini-

Hochberg method. (C) Roots of *tpst* mutant plants were stained with 10 μg/mL

propidium iodide after 7-day-old seedlings were inoculated with R129_E for 14 days

in the presence or absence of 100 nM RGF1. Arrowheads indicate the site of transition

from cell division to cell elongation. Bars, 50 μm.

1282

1283   **Supplemental Table 1. RNAseq datasets analyzed in this study.**

1284   **Supplemental Table 2. Pairwise PERMANOVA of logFC by lifestyles**