Loss-of-function variants in the schizophrenia risk gene *SETD1A* alter neuronal

network activity in human neurons through cAMP/PKA pathway

Shan Wang[1], Jon-Ruben van Rhijn[1], Ibrahim Akkouh[2, 3], Naoki Kogo[4, 5], Nadine Maas[1],

Anna Bleeck[1], Irene Santisteban Ortiz[1], Elly Lewerissa[4], Ka Man Wu[4], Chantal

Schoenmaker[4], Srdjan Djurovic[2, 6], Hans van Bokhoven[1, 4], Tjitske Kleefstra[4], Nael

Nadif Kasri[1, 4, 7, #], Dirk Schubert[1, 7, #, *]

1.   Department of Cognitive Neurosciences, Radboudumc, Donders Institute for

Brain   Cognition and Behaviour , Nijmegen (the Netherlands)

2.   Department of Medical Genetics, Oslo University Hospital, Oslo (Norway)

3.   NORMENT, Institute of Clinical Medicine, University of Oslo, Oslo, (Norway)

4.   Department of Human Genetics, Radboudumc, Donders Institute for Brain

Cognition and Behaviour, Nijmegen (the Netherlands)

5.   Department of Biophysics, Donders Institute for Brain Cognition and Behaviour,

Nijmegen (the Netherlands)

6.   NORMENT, Department of Clinical Science, University of Bergen, Bergen,

(Norway)

7.   These authors contributed equally

[#]Lead contact

[*]Corresponding author: Dirk Schubert

Email: d.schubert@donders.ru.nl

Dept. Cognitive Neuroscience

Donders Institute for Brain, Cognition and Behaviour

Radboud University Nijmegen Medical Centre

Postbus 9101

6500 HB Nijmegen

The Netherlands

Phone: +31 24 3615039

Summary

Heterozygous loss-of-function (LoF) mutations in *SETD1A*, which encodes a subunit

of histone H3 lysine 4 methyltransferase, were shown to cause a novel

neurodevelopmental syndrome and increase the risk for schizophrenia. We generated

excitatory/inhibitory neuronal networks from human induced pluripotent stem cells with

a *SETD1A* heterozygous LoF mutation (*SETD1A*$^{+/-}$) using CRISPR/Cas9. Our data

show that *SETD1A* haploinsufficiency resulted in morphologically increased dendritic

complexity and functionally increased bursting activity. This network phenotype was

primarily driven by *SETD1A* haploinsufficiency in glutamatergic neurons. In

accordance with the functional changes, transcriptomic profiling revealed

perturbations in gene sets associated with glutamatergic synaptic function. At the

molecular level, we identified specific changes in the cAMP/PKA pathway pointing

toward a hyperactive cAMP pathway in *SETD1A*$^{+/-}$ neurons. Finally, by

pharmacologically targeting the cAMP pathway we were able to rescue the network

deficits in *SETD1A*$^{+/-}$ cultures. Our results demonstrate a link between SETD1A and

the cAMP-dependent pathway in human neurons.

Keywords

SETD1A; neurons; Schizophrenia; network activity; micro-electrode array; cAMP/PKA.

Introduction

Schizophrenia (SCZ) is a complex and heterogeneous syndrome with poorly defined neurobiology. It is a highly heritable disease (~80% heritability) with a substantial genetic component (Hilker et al., 2018; Legge et al., 2021a; Skene et al., 2018). In the past decade, considerable progress has been made to better understand the genetic burden related to SCZ. Genetic loci related to SCZ can either be common variants, which typically exert small effects, or rare variants, which can result in a large effect on individual risk (Legge et al., 2021b). One of the genes with such rare large-effect size variants is *SETD1A,* encoding SET domain-containing protein 1A*.* Accumulating studies demonstrate that loss-of-function (LoF) and missense mutations in *SETD1A* are associated with SCZ (Singh et al., 2016, 2020; Takata et al., 2014), but also found in individuals with disrupted speech development (Eising et al., 2019) and early-onset epilepsy (Yu et al., 2019). Furthermore, we presented a series of de novo *SETD1A* heterozygous LoF mutations and defined a novel neurodevelopmental syndrome based on a cohort of 15 individuals (age ranging from 34 months to 23 years old) (Kummeling et al., 2020). The core characteristics of these individuals include global developmental delay (such as speech delay or motor delay) and/or intellectual disability, facial dysmorphisms as well as behavior and psychiatric abnormalities, including psychotic episodes. In addition, abnormalities in brain structure, visual and hearing impairments have also been reported in some of these individuals (Kummeling et al., 2020). *SETD1A* mutations thus appear to cause biological vulnerability to a broad neurodevelopmental phenotypic spectrum.

SETD1A encodes a subunit of the human Set/COMPASS complex, which methylates histone H3 at position lysine-4 (H3K4me1, H3K4me2, H3K4me3) and participates in the regulation of gene expression. Mouse models with heterozygous LoF mutation of *Setd1a* (*Setd1a$^{+/-}$*) recapitulate SCZ-related behavioral abnormalities, such as deficits in working memory and social interaction (Mukai et al., 2019; Nagahama et al., 2020). At the cellular level, *Setd1a$^{+/-}$* mice display reduced axon branches and dendritic spines, increased neuronal excitability (Mukai et al., 2019) as well as impaired

excitatory synaptic neurotransmission (Nagahama et al., 2020). Furthermore,

Setd1a[+/-] mice show an altered transcriptomic profile in the medial prefrontal cortex

(mPFC), a highly relevant region for SCZ (Mukai et al., 2019). In other brain regions,

such as visual cortex (V1), *Setd1a*[+/-] mice exhibit aberrant ensemble activity and

gamma oscillations (Hamm et al., 2020). All of these studies suggest *SETD1A*

haploinsufficiency results in neuronal circuit dysfunction. However, the exact cellular

and molecular mechanisms of how *SETD1A* mutations lead to disrupted neuronal

connectivity causing such severe mental symptoms, especially in a human context,

remain poorly understood.

To investigate the role of SETD1A in neuronal network development and synaptic

organization, we generated an isogenic human induced pluripotent stem cell (hiPSC)

line with *SETD1A* haploinsufficiency through CRISPR/Cas9. Subsequently, hiPSCs

were differentiated into homogenous populations of glutamatergic and GABAergic

neurons (Mossink et al., 2021). In *in vitro* cultures containing defined compositions of

glutamatergic and GABAergic neurons, we comprehensively analyzed molecular,

structural and functional neuronal properties from single cells and neuronal networks

during development. The results presented here demonstrate that *SETD1A*

haploinsufficiency leads to key morphological, electrophysiological and transcriptional

alterations. At molecular level, we show that the *SETD1A*[+/-] network phenotype is

mediated by upregulated cyclic adenosine monophosphate (cAMP)/PKA pathway.

This has been further confirmed by showing that pharmacological inhibition of

cAMP/PKA rescues the *SETD1A*[+/-] network phenotype. Therefore, our results reveal

cAMP/PKA as a potential downstream pathway affected by *SETD1A* mutation,

opening novel therapeutics opportunities for patients carrying *SETD1A* variant.

Results

***SETD1A*[+/-] neuronal networks exhibit dysregulated functional organization.**

We used CRISPR/Cas9 to generate an isogenic hiPSC line with a heterozygous LoF

mutation of *SETD1A* by targeting exon 7 of *SETD1A* in a healthy hiPSC line (Miyaoka

et al., 2014) **(Figure 1a)**. We introduced a frameshift mutation in exon 7 leading to a LoF of the protein, mimicking a mutation reported in an individual diagnosed with SCZ (Takata et al., 2014). Two clones with indels in *SETD1A* were selected for further characterization, which carry 28 (clone 1) and 8 (clone 2) base pairs (bp) deletions on one allele, respectively **(Figure 1a, S1a)**, both predicting a premature stop codon. SETD1A mRNA and protein levels were approximately halved in both *SETD1A*$^{+/-}$ hiPSC clones, indicating that both variants represent LoF alleles **(Figure 1b-c, S1b)**. All selected clones showed positive expression of pluripotency markers (OCT4, TRA-1-81, NANOG and SSEA4) and karyotyping and off-target analysis was performed to confirm genetic integrity **(Figure S1c-e)**.

Disrupted neuronal connectivity has been reported in SCZ patients and animal models of SCZ (Owen et al., 2016). We evaluated whether LoF of *SETD1A* gene results in neuronal network impairment *in vitro*. To this end, we generated composite networks of glutamatergic (~75%) and GABAergic (~25%) neurons (excitatory/inhibitory cultures, E/I cultures) comprised of either control or *SETD1A*$^{+/-}$ hiPSCs, by forced expession of the transcription factor *Ngn2* or *Ascl1,* respectively, as recently described (**Figure 1d**) (Mossink et al., 2021). We did not detect any significant differences in the percentage of glutamatergic or GABAergic neurons between control and *SETD1A*$^{+/-}$ networks at days *in vitro* (DIV) 49 (**Figure 1e, S2a**). Moreover, following acute treatment with 100 µM Picrotoxin (PTX) at DIV49 on micro-electrode array (MEA), both mean firing rate and network burst duration went up for both control and *SETD1A*$^{+/-}$ networks (**Figure S2b-c**), indicating that at DIV49 GABAergic neurons exhibit robust inhibitory control (Mossink et al., 2021).

In order to assess if neuronal network activity differs between control and *SETD1A*$^{+/-}$ E/I cultures during development, we used MEA recordings (Frega et al., 2019) (**Figure S2b**). We recorded neuronal network activity once a week from DIV21 to DIV49. After 3 weeks of differentiation, both control and *SETD1A*$^{+/-}$ networks showed network burst activity **(Figure 1f)**, which indicates the neurons are functionally connected and integrated into a network. Network burst activity increased during development for both groups, and plateaued at around DIV42. Strikingly,

SETD1A[+/-] networks showed a significantly increased network burst activity compared

to control from DIV21 to DIV49, accompanied by a shorter network inter burst interval,

a decrease in network burst duration as well as lower spike rate within a burst **(Figure**

1g, S2d). Interestingly, the global activity (i.e., mean firing rate, **Figure 1g**) was

similar between control and *SETD1A*[+/-] networks, which implies there was a functional

re-organization of network connectivity in *SETD1A*[+/-] E/I cultures, rather than general

hyperactivity. Calcium imaging further confirmed the increased synchronized activity

in *SETD1A*[+/-] networks **(Figure S3a-d, Supplementary Video1-2)**.

We next evaluated if this functional network re-organization in *SETD1A*[+/-] E/I

cultures is related to changes in intrinsic properties and/or synaptic inputs using

single-cell patch clamp. Intrinsic properties were similar between control and

SETD1A[+/-] neurons at both DIV21 and DIV49 **(Figure S4, Table S1)**, in glutamatergic

and GABAergic neurons. We additionally measured general synaptic integration by

recording spontaneous excitatory postsynaptic currents (sEPSCs). Neither sEPSC

frequency nor amplitude were different between control and *SETD1A*[+/-] cultures at

DIV21 or DIV49 **(Figure 1h, Table S1)**. However, we did observe that the frequency

of temporally correlated bursts of synaptic inputs (sEPSC bursts) onto the

postsynaptic neuron was significantly increased at DIV49 in *SETD1A*[+/-] E/I cultures,

both on glutamatergic and GABAergic neurons **(Figure 1i)**. This is in line with the

increased network burst rate at the population level shown in the MEA recordings and

confirms that reduced SETD1A expression results in increased network activity. Since

analysis of sEPSCs reflects network activity rather than yield quantitative information

regarding synaptic connectivity, we next measured miniature postsynaptic currents

(mEPSCs), which are action-potential independent and can be indicative for the

reorganization of synaptic inputs, altered synapse numbers, and changes in release

probabilities (mEPSC frequency) and receptor abundance (mEPSC amplitude). Both

mEPSC frequency and amplitude were significantly increased in both glutamatergic

and GABAergic *SETD1A*[+/-] neurons at DIV49 (**Figure 1j**). This suggests that networks

comprised of *SETD1A*[+/-] neurons not only exhibit increased network activity but also

elevated synaptic connectivity, and this could be a major contributor to the network

phenotype we observed at the population level.

SETD1A^{+/-} neurons show aberrant somatodendritic morphology. _SETD1A_$^{+/-}$ neurons show aberrant somatodendritic morphology.

Based on the MEA results, we next sought to investigate whether _SETD1A_$^{+/-}$ neurons

exhibit changes in dendritic morphology and/or synapse formation. At DIV21, control

and _SETD1A_$^{+/-}$ E/I cultures did not differ from each other in any of the morphological

parameters for both glutamatergic and GABAergic neurons **(Figure 2a-b, e-f)**.

However, at DIV49, _SETD1A_$^{+/-}$ neurons displayed a significantly larger soma size,

accompanied by a longer dendritic length, more dendritic branches and a larger

covered area for both glutamatergic and GABAergic neurons **(Figure 2c-f)**. Finally,

we performed a Sholl analysis and confirmed that there were no significant

differences in the distribution of dendritic length at DIV21 for both neuronal types

(Figure 2a-b), whereas at DIV49, the dendritic length at multiple distances from the

soma was significantly longer in both glutamatergic and GABAergic _SETD1A_$^{+/-}$

neurons **(Figure 2c-d)**. These results collectively indicate that _SETD1A_

haploinsufficiency leads to a more complex somatodendritic morphology, but this only

becomes significant during later developmental stages.

To measure synapse formation, we immunostained neurons for pre- and

postsynaptic markers (Synapsin and Homer1 for glutamatergic synapses, VGAT and

Gephyrin for GABAergic synapses). We found no differences in the density of

Synapsin/Homer1 or VGAT/Gephyrin co-localised puncta between genotypes at both

DIV21 and DIV49 on excitatory and inhibitory neurons **(Figure 2g-h, S5)**. These data,

combined with the observation that all _SETD1A_$^{+/-}$ neurons show a longer dendritic

length at DIV49, imply that the total amount of synapses per neuron is higher in

SETD1A$^{+/-}$ neurons, which is reflected by the increase in mEPSC frequency at DIV49

(Figure 1j).

SETD1A haploinsufficiency leads to an altered transcriptomic profile.

To identify the molecular perturbations that underlie the network phenotypes caused by *SETD1A* haploinsufficiency, we conducted RNA-seq at DIV49. Principal component analysis (PCA) of transcriptional profiles exhibited clear clustering of biological replicates per genotype **(Figure 3a)**. We quantified the differentially expressed genes (DEGs) and found that 380 genes were downregulated and 539 genes were upregulated in *SETD1A*[+/-] E/I cultures compared with control cultures **(Figure 3b)**. Interestingly, when we performed a disease enrichment analysis using the DisGeNET database which consists of more than 10,000 disorders (Piñero et al., 2020), SCZ was identified as the top hit **(Figure 3c, Table S2)**. This indicates that *SETD1A* haploinsufficiency in our model captures some key genetic alterations typically linked to SCZ. In addition, we conducted Gene Ontology (GO) analysis to examine which biological functions are over-represented in the DEGs. Significant enrichment in GO terms relevant to "synaptic function", "morphogenesis", "ion channels" and "learning and memory" were identified **(Figure 3d, Table S3)**. Proteins encoded by these genes interact with each other as indicated in the protein-protein interaction analysis generated through STRING (Szklarczyk et al., 2021) **(Figure 3e)**. Interestingly, we noticed that many GO terms and DEGs relevant to glutamatergic neurons were identified, such as "glutamate receptor binding", "glutamate receptor signaling pathway" and "glutamatergic synapse" **(Figure 3d-f)**. This is consistent with the data from *Setd1a*[+/-] mouse models, where it was shown that SETD1A target genes are highly expressed in pyramidal neurons (Mukai et al., 2019). This also suggests that the altered network phenotype caused by *SETD1A* haploinsufficiency might be driven predominantly by perturbations in glutamatergic neurons.

To explore the cross-species validity of the SETD1A target genes, we compared our DEGs list with the published transcriptome of *Setd1a*[+/-] mouse model from Mukai et al (Mukai et al., 2019), and found in total 11 overlapping genes **(Table S4)**. One of these genes is *SLITRK4,* the gene showing the strongest upregulation in *SETD1A*[+/-] cultures (**Figure 3b**), which is involved in neurite outgrowth (Yim et al., 2013). Furthermore, we found a significant association between our DEGs and 78 SFARI genes (Fisher's exact: $p=1.1e^{-5}$), an autism-related genes database **(Table S5)**. This

is in line with the suggested genetic overlap between SCZ and autism (Carroll and

Owen, 2009). Taken together, these results indicate that transcription is profoundly

disturbed in *SETD1A*$^{+/-}$ neurons and DEGs are enriched in specific gene sets related

to SCZ and synaptic function, especially the ones relevant for glutamatergic synapse

function.

SETD1A$^{+/-}$ glutamatergic neuronal cultures recapitulate the SETD1A$^{+/-}$ E/I

network phenotype

Our RNA-seq results clearly establish a link between *SETD1A* haploinsufficiency and

glutamatergic synapse function. We therefore hypothesized that glutamatergic

neurons might be the main contributor to the network phenotype in *SETD1A*$^{+/-}$ E/I

cultures. To test this hypothesis, we set up homogeneous glutamatergic neuronal

cultures **(Figure 4a)**. We measured the glutamatergic neuronal network activity on

MEA between DIV21 to DIV49 **(Figure 4b)**. Consistent with the phenotype of E/I

networks, *SETD1A*$^{+/-}$ glutamatergic neuronal networks exhibited significantly higher

burst activity while decreased burst duration and burst spike rate at DIV49. Global

activity (i.e. mean firing rate) did not differ between two groups **(Figure 4c)**. Next, we

investigated if in glutamatergic cultures, there might be differences between control

and *SETD1A*$^{+/-}$ neurons at the single cell level **(Figure 4g-k, S8a-b, Table S6)**.

Similar to the E/I cultures, we observed an increased sEPSC burst rate at the single

cell level in particular at DIV49 **(Figure 4j-k, Table S6)**. Taken together, our data

show that E/I cultures and glutamatergic cultures are similarly affected when SETD1A

expression is reduced. This was further supported by the observation that in

glutamatergic neuronal networks, *SETD1A*$^{+/-}$ neurons showed a similar increase in

morphological complexity as in the E/I cultures **(Figure 4d-f, S7a-d)**. Finally, we

performed a transcriptome analysis for glutamatergic cultures at DIV49. We identified

in total 455 DEGs, with 177 downregulated and 278 upregulated genes **(Figure**

S6a-b). Notably, we again identified *SLITRK4* as a top hit **(Figure S6b)**. Enrichment

analysis detected GO terms such as "glutamatergic synapse", "synapse organization",

"synapse assembly", and many others related to synaptic function **(Figure S6d-f)**. In

disease association analysis, we found "bipolar disorder" and "SCZ" within the top hits

(Figure S6c). Overall, *SETD1A*⁺ᐟ⁻ glutamatergic cultures recapitulate the network

phenotype observed in the E/I cultures, indicating that SETD1A plays a major role in

glutamatergic neurons.

***SETD1A* haploinsufficiency leads to activation of cAMP/PKA/CREB pathway**

The changes in neuronal network activity observed in *SETD1A*⁺ᐟ⁻ cultures (E/I and

glutamatergic cultures) suggest that pathways related to regulation of neuronal

activity could be affected. Interestingly, in our transcriptome data we found that the

upregulated DEGs were enriched in several annotations related to second messenger

signaling, such as "G protein-coupled receptor signaling pathway, coupled to cyclic

nucleotide second messenger", "second-messenger-mediated signaling", and

"adenylate cyclase-modulating G protein-coupled receptor signaling pathway" **(Figure**

5a, Table S7). In agreement with this finding, we found that multiple enzymes

involved in the synthesis and degradation of cAMP were dysregulated, both in the E/I

and glutamatergic cultures. Specifically, genes coding for proteins involved in cAMP

production (*ADCY2, ADCY3* and *ADCY8)* (Pieroni et al., 1995; Qiu et al., 2016; Wang

and Storm, 2003) were significantly upregulated, whereas genes encoding for

phosphodiesterases (*PDE12, PDE7a* and *PDE1a*), important for cAMP degradation

(Epstein, 2017; Evripioti et al., 2019; Keravis and Lugnier, 2012), were downregulated

(Figure 5b). This strongly suggests that the cAMP pathway is hyperactive in

SETD1A⁺ᐟ⁻ neurons.

To test this hypothesis, we measured the phosphorylation of cAMP-response

element binding protein (pCREB), which is a downstream signaling protein in the

cAMP/PKA pathway (Tasken et al., 1993) **(Figure 5c)**. We immunostained neurons

from DIV49 E/I cultures for anti-pCREB, an antibody detecting endogenous levels of

CREB only when phosphorylated at serine[133]. As a positive control, we stimulated the

cAMP pathway on control E/I cultures with 10 µM forskolin (FSK), a well-known

adenylyl cyclase (AC) agonist and PKA/CREB pathway activator (Delghandi et al.,

2005; Namkoong et al., 2009; Schiller et al., 2010). As expected, FSK significantly

increased the level of pCREB in control neurons. In addition, we found that the level of

pCREB in *SETD1A*$^{+/-}$ neurons was significantly higher compared to control neurons

(Figure 5d-e). This result supports our hypothesis that the cAMP/PKA/CREB pathway

exhibits increased activity in *SETD1A*$^{+/-}$ neurons.

To further investigate the link between enhanced cAMP activation and the

neuronal network phenotype observed in *SETD1A*$^{+/-}$ neurons, we bidirectionally

modulated cAMP levels in control or *SETD1A*$^{+/-}$ neurons from E/I cultures.

Interestingly, we found that acute stimulation of DIV51 control neurons with 10 µM

FSK for 1 hour resulted in an increase in network bust rate, with a parallel reduction in

inter network burst interval **(Figure 5f-g, S9a-c)**, mimicking the network phenotype of

SETD1A$^{+/-}$ E/I cultures. In addition, burst spike rate and network burst duration were

decreased, although the difference was not significant **(Figure S9c)**. These results

show that control networks treated with forskolin resemble *SETD1A*$^{+/-}$ neuronal

networks. In a second set of experiments, we examined whether blocking cAMP using

AC inhibitor SQ22536 in *SETD1A*$^{+/-}$ networks (starting from DIV42) would be sufficient

to normalize the phenotype caused by *SETD1A* haploinsufficiency to control levels.

Indeed, after 8 days of treatment with 100 µM SQ22536, we found that the major

network parameters were normalized, including the network burst rate **(Figure 5h-i,**

S10a-b). We next investigated whether acute manipulation of the PKA pathway,

which is a downstream target of cAMP, would be sufficient to normalize *SETD1A*$^{+/-}$

network activity to control levels. Two independent chemical inhibitors of PKA, H89

and KT5720 (Murray, 2008; Song et al., 2015), were applied to *SETD1A*$^{+/-}$ networks at

DIV51 on MEA. A one-hour treatment with either H89 (2 µM) or KT5720 (1 µM) in

SETD1A$^{+/-}$ networks, was sufficient to normalize the major network parameters,

including the network burst rate **(Figure 5j-k, S11a-c)**. Taken together, our results

suggest that cAMP/PKA is the main molecular pathway through which *SETD1A*

haploinsufficiency leads to key neuronal network alteration.

Discussion

Heterozygous LoF variants in *SETD1A*, with high penetrance for SCZ, provides a unique opportunity to investigate neuronal dysfunction that might underlie the increased vulnerability to SCZ. Here, we reveal crucial molecular signatures and neurodevelopmental abnormalities caused by *SETD1A* haploinsufficiency in human neurons. We show that *SETD1A* haploinsufficiency leads to altered neuronal network organization, within which changes in glutamatergic neuronal function might be one of the main contributors. Furthermore, we identify increased cAMP/PKA activity as a molecular mechanism to the functional phenotype of *SETD1A*$^{+/-}$ networks.

One major consistent characteristic of hiPSC-derived *SETD1A*$^{+/-}$ networks is the increased network burst rate, which is reflected by an increase in synaptic connectivity at the single cell level. Synchronized burst activity plays an important role during early brain development (Khazipov and Luhmann, 2006) and is considered to be a fundamental mechanism for information processing, hence relevant for perception, memory and cognition (Lisman, 1997). Intriguingly, abnormalities in neural synchronization *in vivo*, which reflect neural circuit dysfunction, are suggested as one of the core pathophysiological mechanisms in SCZ (Ford et al., 2007; Uhlhaas and Singer, 2010). For example, increased oscillatory synchronization is found to be correlated with positive symptoms of SCZ (Lee et al., 2006; Spencer et al., 2009), while both increased and reduced oscillations are associated with negative symptoms (Lee et al., 2003; Uhlhaas and Singer, 2010). In the primary visual cortex of *Setd1a*$^{+/-}$ mice, Hamm et al identified reduced oscillations, providing circuit-level insight into the underlying neurobiology of sensory-processing abnormality seen in SCZ (Hamm et al., 2020). In this regard, both our human *in vitro* model and mouse model showed aberrant synchronized activity at different developmental stages, suggesting circuit disruption caused by *SETD1A* haploinsufficiency.

Disrupted glutamatergic signaling plays a critical role in the pathogenesis of SCZ (Tsai and Coyle, 2002). Consistently, on several levels, our data provide evidence

supporting the idea that glutamatergic signaling are one of the main contributors for

the network phenotype caused by *SETD1A* haploinsufficiency. First, through

transcriptional profiling of *SETD1A*$^{+/-}$ E/I cultures, we identified a genetic signature

suggestive of perturbed function of glutamatergic signaling. Dysregulated genes

include the ones encoding for metabotropic glutamate receptors (*GRM4, GRM3*),

ionotropic glutamate receptors subunits (*GRIK3, GRIN2A*), as well as proteins

involved in glutamatergic signaling (*SHISA6, SHISA7, CAMK4, PSD93, VAMP1*). This

is in line with the previous finding that target genes of SETD1A were mainly

expressed in pyramidal neurons (Mukai et al., 2019). In particular, *SHISA6* which is

important for AMPA receptor function has been strongly down-regulated, suggesting

dysregulated AMPA receptor activity (Klaassen et al., 2016). Furthermore, in L2/3

pyramidal neurons from the mPFC of *Setd1a*$^{+/-}$ mice, only excitatory synaptic

transmission was altered, whereas inhibitory synaptic transmission remained

unchanged (Nagahama et al., 2020). Second, in E/I networks, when GABAergic

inputs were blocked by PTX, the phenotype of increased networks burst frequency

still remained **(Figure S2c)**. In addition, measurements from single-cell

electrophysiology and MEA with homogenous glutamatergic neurons show the same

network signature as observed in E/I networks, indicating glutamatergic neurons are

one of the main driving factors in this phenotype. Third, from both E/I and

glutamatergic cultures, transcriptomic data highlight a striking upregulation of

SLITRK4, which is involved in neurite outgrowth and synaptogenesis, especially in

glutamatergic synapse formation (Yim et al., 2013). Moreover, it has been shown that

variants in *SLITRK4* are associated with neuropsychiatric disorders, such as

schizophrenia (Kang et al., 2016). *SLITRK4* has also been identified as a DEG from

Setd1a$^{+/-}$ mice (Mukai et al., 2019), emphasizing its important role as a downstream

target of SETD1A across species. Indeed, our data show that *SETD1A*

haploinsufficiency leads to increased dendritic complexity in human glutamatergic

neurons. This morphological phenotype however was not detected in the *SETD1A*$^{+/-}$

mice model from Mukai et al. where pyramidal neurons in mature prefrontal cortical

networks showed unchanged dendritic arborization (Mukai et al., 2019). This

difference can potentially be explained by the differences in the assessed

developmental stages or species related differences in SETD1A function. Taken

together, these results imply that SETD1A plays an essential role in the function of

glutamatergic signaling. However, dysfunction of GABAergic signaling might be

involved in the network phenotype as well. Our results show that *SETD1A*

haploinsufficiency leads to increased dendritic complexity of GABAergic neurons,

which likely affects the input connectivity and signal integration of these neurons.

Further experiments are needed to clearly dissect the role of SETD1A in GABAergic

neurons.

Our data identify several molecular pathways that may contribute to the increased

network burst frequency in *SETD1A*$^{+/-}$ networks. In particular, our transcriptomic data

indicated that *SETD1A* haploinsufficiency leads to increased cAMP/PKA, which has

been shown to enhance synaptic strength through different aspects, such as by

modulating different ion channels (Anderson et al., 2000; Beaumont and Zucker, 2000;

Connors et al., 2008; Gutierrez-Castellanos et al., 2017; Hell' et al., 1995; van der

Horst et al., 2020; Renner et al., 2017) and increasing neurite outgrowth (Aglah et al.,

2008; Chu et al., 2006; Wan et al., 2011). Intriguingly, a recent study demonstrated

that cAMP/PKA induces calcium influx through voltage-gated calcium channel, which

pushes the neuronal network towards a large-scale and synchronized burst activity

(Thornquist et al., 2021). Consistently, our calcium imaging data showed increased

synchronized activity in *SETD1A*$^{+/-}$ networks, suggesting that there could be a similar

cAMP-triggered mechanism in our model.

cAMP signaling has been implicated in neurodevelopmental and neuropsychiatric

disorders, including SCZ (Funk et al., 2012; Millar et al., 2005; Muly, 2002; Turetsky

and Moberg, 2009; Vacic et al., 2011; Wang et al., 2018) , bipolar disorder (Chang et

al., 2003; Ren et al., 2014), Fragile X syndrome (Berry-Kravis et al., 1995; Berry-

Kravis and Huttenlocher, 1992; Kelley et al., 2007) and autism (Kelley et al., 2008;

Zamarbide et al., 2019). For example, in patients with bipolar disorder, cAMP/PKA

signaling is upregulated (Chang et al., 2003), while in the case of Fragile X syndrome,

cAMP/PKA signaling is decreased (Berry-Kravis et al., 1995). These findings

emphasize the essential role of balanced cAMP activity in normal brain function,

making it an appealing therapeutic target. Interestingly, our results show that on

network level both blocking the PKA pathway and inhibiting AC can rescue the

phenotype. However, we cannot rule out the possibilities that other downstream

effectors of cAMP such as EPAC (a guanine-nucleotide-exchange factor) and

cyclic-nucleotide-gated ion channels (Sassone-Corsi, 2012), are also partly involved

in the phenotype caused by *SETD1A* haploinsufficiency. In addition, since cAMP/PKA

signaling functions in various cells types, therapeutic intervention of its activity

therefore needs further clarification of its cell type specific roles. In this context, future

studies should comprehensively investigate whether upregulated cAMP/PKA

signaling caused by *SETD1A* haploinsufficiency also applies to other brain regions or

other cell types related to SCZ or neurodevelopmental disorders, such as

serotonergic (Carvajal-Oliveros and Campusano, 2021) or dopaminergic neurons

(Eells, 2005), as well as glial cells (Blanco-Suárez et al., 2017).

Even though our current human neuronal *in vitro* model skips certain early

developmental stages, such as the neuro-progenitor cell stage, of neuronal networks

and it cannot capture the cellular complexity of fully mature cortical networks, the

defined composition of human neuronal networks allows the assessment of core

molecular and cellular mechanisms related to *SETD1A* haploinsufficiency. Taken

together, our data suggest glutamatergic synaptic dysfunction is one of the potential

pathogenic mechanisms of *SETD1A* haploinsufficiency-associated disorders and we

identified cAMP/PKA dysregulation as underlying mechanisms responsible for the

altered network phenotype in *SETD1A*$^{+/-}$ cultures. Future studies, using both rodent

models and human *in-vitro* neuronal models are required to further explore the impact

and therapeutic potential of cAMP/PKA in *SETD1A*-associated disorders.

Materials and Methods

Cell culture and neuronal differentiation

hiPSCs used in this study were obtained from reprogrammed fibroblasts (Miyaoka et al., 2014). hiPSCs were cultured on a 6-well plate pre-coated with 1:15 (diluted in DMEM/F12 medium) Matrigel (Corning, #356237) in Essential 8 Flex medium (Thermo Fisher Scientific) supplemented with primocin (0.1 g/ml, Invitrogen) at $37^{o}C/5\%CO_2$. hiPSCs were infected with *Ascl1* or *Ngn2* and *rtTA* lentivirus (The transfer vector used for the rtTA lentivirus is pLVX-EF1α-(Tet-On-Advanced)-IRES-G418(R); The transfer vector used for the Ngn2 lentivirus is pLVX-(TRE-thight)-(MOUSE) Ngn2-PGK-Puromycin(R); The transfer vector used for the Ascl1 lentivirus is pLV[TetOn]-Puro-TRE>mAscl1. All the plasmids are available upon request). Medium was supplemented with puromycin (0.5 g/ml) and G418 (50 g/ml). Medium was refreshed every 2-3 days and hiPSCs were passaged twice per week using an enzyme-free reagent (ReLeSR, Stem Cell Technologies).

The neuronal differentiation protocol used in this article was previously described (Mossink et al., 2021). Glutamatergic neurons were either cultured alone or in coculture with GABAergic neurons. When cocultured, GABAergic neurons were plated at days *in vitro* (DIV) 0 and transduced with AAV2-hSyn-mCherry (UNC Vector Core) for visualization. After 4 hours incubation, cultures were washed twice with DMEM/F12 (Thermo Fisher Scientific) after which glutamatergic neurons were plated into the well. hiPSCs were plated in E8 flex supplemented with doxycycline (4 µg/ml), Revitacell (1:100, Thermo Fisher Scientific) and Forskolin (10 µM, Sigma). At DIV 1 cultures were switched to DMEM/F12 containing Forskolin, N2 (1:100, Thermo Fisher Scientific), non-essential amino acids (1:100, Sigma), primocin (0.1 µg/ml, Invivogen), NT3 (10 ng/ml, PromoCell), BDNF (10 ng/ml, PromoCell), and doxycycline (4 µg/ml). To support neuronal maturation, freshly prepared rat astrocytes were added to the culture in a 1:1 ratio at DIV 2. At DIV 3 the medium was changed to Neurobasal

medium (Thermo Fisher Scientific) supplemented with Forskolin (10 µM, Sigma),

B-27 (Thermo Fisher Scientific), GlutaMAX (Thermo Fisher Scientific), primocin (0.1

µg/ml), NT3 (10 ng/ml), BDNF (10 ng/ml), and doxycycline (4 µg/ml). To remove the

proliferating cells from the culture, cytosine-b-D-arabinofuranoside (Ara-C, 2 µM,

Sigma) was added to the medium at DIV 3. From DIV 6 onwards half of the medium

was refreshed three times a week. From DIV 10 onwards, the medium was

additionally supplemented with 2.5% FBS (Sigma) to support astrocyte viability. After

DIV 13, Forskolin and doxycycline were removed from the medium. Cultures were

kept at least until DIV49.

CRISPR/Cas9 editing of *SETD1A*

CRISPR/Cas9 technology was used to create a heterozygous indel mutation in exon

7 of *SETD1A* in a healthy hiPSC line derived from a male, 30 years old (Mandegar et

al., 2016). In brief, sgRNAs were designed to specifically target *SETD1A*

(GTCCTTGGGGCCAGAGATAC AGG), and then cloned into pSpCas9(BB)-2A-Puro

(PX459) V2.0 (Addgene #62988). Single-cell suspension of hiPSCs was nucleofected

with 5 µg of the generated SpCas9-sgRNA plasmid using the P3 Primary Cell

4D-Nucleofector Kit (Lonza, #V4XP-3024) in combination with the 4D Nucleofector

Unit X (Lonza, #AAF-1002X), program CA-137. After nucleofection, cells were

resuspended in E8 Flex supplemented with Revitacell and seeded on Matrigel

pre-coated plates. One day after the nucleofection, 0.5 µg/ml puromycin was added

for 24 hours for selection. Puromycin-resistant colonies were manually picked and

Sanger Sequencing was performed to ensure heterozygous editing of exon 7. Two

positive clones were selected for further characterization. Cells were tested routinely

as mycoplasma-negative. The expression of pluripotency markers OCT4, NANOG,

SSEA4 and TRA1-81 were detected with immunocytochemistry. Karyotyping was

performed as a service by the Diagnostics Department at Radboud University Medical

Center. Potential off-target sites were checked by sanger sequencing.

Immunocytochemistry

Cells seeded on coverslips were fixed with 4% paraformaldehyde supplemented with

4% sucrose for 15 min at room temperature, followed by permeabilization with 0.2%

triton for 10 min. Nonspecific binding sites were blocked by incubation in 5% normal

goat serum for 1 hour at room temperature. Cells were incubated with primary

antibodies overnight at 4°C. The second day, secondary antibodies, conjugated to

Alexa-fluorochromes, were added and incubated for 1 hour at room temperature.

Hoechst 33342 was used to stain the nucleus before cells were mounted with

fluorescent mounting medium (DAKO, #S3023). The following primary antibodies

were used: Rabbit anti-MAP2 (1:1000, Abcam, #ab32454); Mouse anti-Gephyrin

(1:1000, Synaptic Systems 147011); Rabbit anti-VGAT (1:500, Synaptic systems

131013); Guinea pig anti-Synapsin 1/2 (1:1000, Synaptic Systems 106004); Mouse

anti-Homer1 (1:500, Synaptic Systems 160011); Rabbit anti-Phospho-CREB (1:500,

Cell Signaling 87G3 9198). Secondary antibodies that were used are: Goat-anti rabbit

Alexa 568 (1:1000, Invitrogen, A11036); Goat-anti-mouse Alexa 488 (1:1000,

Invitrogen, A11029); Goat anti-guinea pig Alexa Fluor 568 (1:2000,

Invitrogen, A11075); Goat anti-guinea pig Alexa Fluor 647 (1:1000, Invitrogen,

A21450). Cells were imaged at 63x magnification using the Zeiss Axio Imager Z1

equipped with ApoTome. Fluorescent images were analyzed using FIJI software

(Schindelin et al., 2012). Synapse puncta were counted manually and normalized to

the length of the dendritic branch where they reside.

Western Blot

To lyse the cells, medium was removed and the well was washed with 2 ml ice cold

PBS before 100 µl lysis buffer were applied (RIPA buffer supplemented with

PhosSTOP; Roche) and protease inhibitors (complete Mini, EDTA free; Roche).

Before blotting the protein, concentration was determined by means of a Pierce™

BCA protein assay (Thermo Scientific 23225). For each sample, the same amount of

protein around 15 µg was loaded and separated by SDS-PAGE. Depending on the

primary antibody, separated proteins were transferred to PVDF membrane (BioRad).

Primary antibodies were used are: Mouse anti-SET1A (1: 500, Santa Cruz 515590);

Mouse anti-γ-tubulin (1:1000, Sigma T5326). For visualization horseradish

peroxidase-conjugated secondary antibodies were used: Goat anti-mouse (1:20000;

Jackson Immuno Research Laboratories 115-035-062).

Neuron reconstruction

Reconstruction was performed using Neurolucida 360 (Version 2017.01.4,

Microbrightfield Bioscience). Neurons were fixed and labeled with MAP2 antibody. To

distinguish GABAergic neurons from glutamatergic neurons, we infected GABAergic

neurons with AAV2-hSyn-mCherry for (UNC Vector Core) visualization. We chose two

time points to fix the neurons: DIV21 and DIV49. This allows us to compare the

morphological phenotype at different developmental stages. Fluorescent images of

MAP2-labelled neurons were taken at 20x magnification using Zeiss Axio Imager Z1

equipped with ApoTome. The images were stitched using Fiji 2018 software with the

stitching plugin and followed by reconstruction using Neurolucida 360 (Version

2017.01.4, Microbrightfield Bioscience). The 3-dimensional reconstructions and

quantitative morphometrical analyses focused on the somatodendritic organization of

the neurons. The axon was excluded based on its long, thin properties and

far-reaching projections. Neurons that had at least two primary dendrites were

selected for reconstruction and further analysis. For morphometrical analysis, we

determined soma size, number of primary dendrites, dendritic nodes and ends and

total or mean dendritic length as well as covered surface by dendritic trees. In addition,

we also investigated dendritic complexity by performing Sholl analysis. Sholl profile

was obtained by applying a series of concentric circles at 20 μm interval from the

soma center, subsequently, dendritic length, number of intersections and number of

nodes of the neurons were measured for each distance interval.

MEA recordings and analysis

All recordings were performed using the 24-wells MEA system (Multichannel Systems,

MCS GmbH, Reutlingen, Germany). Recordings and analysis were performed

according to previous published protocols (Frega et al., 2019). Briefly, spontaneous

electrophysiological activity of hiPSC-derived neuronal networks cultured on MEA

was recorded for 10 minutes in a recording chamber which was constantly maintained

at 37^0C with 95%O_2 and 5%CO_2. Before recording, cultures on MEA were allowed to

adapt for 10 min in the recording chamber. The recording was sampled at 10 kHz, and

filtered with a high-pass filter with a 100 Hz cut-off frequency and a low-pass filter with

a 3500 Hz cut-off frequency. The spike detection threshold was set at ± 4.5 standard

deviations. Spike, burst and network burst detection was performed by a built-in

algorithm in Mulitwell Analzer software (Multichannel Systems), and a custom-made

MATLAB (The Mathworks, Natrick) code to extract parameters characterizing network

activity. Mean firing rate (MFR) was calculated as the average of the spike frequency

of all channels across one MEA well. From the burst detection, the number of bursting

channels (above threshold 0.4 burst/s and at least 5 spikes in burst with a minimal

inter-burst-interval of 100 ms) was determined. A network burst was defined when at

least 50% of the channels in one well exhibited a synchronous burst.

Chemicals

All reagents were prepared fresh into concentrated stocks, and stored frozen at

−20 °C. The following compounds were used in pharmacological experiments:

Picrotoxin (100 mM in DMSO, Tocris 1128); Forskolin (12 mM in DMSO, Sigma

F6886); SQ22536 (50 mM in DMSO, Sigma S153); H89 (5 mM in MQ, Tocris 2910);

KT5720 (1 mM in DMSO, Tocris 1283). For all experiment on MEA, before adding

chemical to the cultures, an aliquot of the concentrated stock was first diluted in DPBS

at room temperature. Then, the appropriate amount of working dilution was added

directly to wells on the MEA and mixing was primarily through diffusion into the (500 µl)

cell culture medium.

Pharmacological experiment

Control and *SETD1A*[+/-] networks on MEA were treated with Picrotoxin (PTX, 100 µM),

Forskolin (1 µM), H89 (2 µM) and KT5720 (1 µM) at DIV49 or DIV51, and SQ22536

(100 µM) at DIV42 after a 20 min recording of spontaneous activity. Then the

recording was stopped temporarily, and the compounds were added to the MEA. We

recorded neuronal network activity for 10 min after 5 min treatment of PTX, 60 min

treatment of Forskolin, KT5720, H89 and 8 days treatment of SQ22536.

Whole cell patch clamp

Whole cell patch clamp was performed as previously described (Mossink et al., 2021).

Coverslips were placed in the recording chamber of the electrophysiological setup,

continuously perfused with oxygenated (95% O_2/ 5% CO_2) ACSF at 32°C containing

(in mM) 124 NaCl, 1.25 NaH$_2$PO4, 3 KCl, 26 NaHCO$_3$, 11 Glucose, 2 CaCl$_2$, 1 MgCl$_2$.

Patch pipettes with filament (ID 0.86 mm, OD1.05 mm, resistance 6-8 MΩ) were

pulled from borosilicate glass (Science Products GmbH, Hofheim, Germany) using a

Narishige PC-10 micropipette puller (Narishige, London, UK). For all recordings of

intrinsic properties and spontaneous activity and mPSC activity, a potassium-based

intracellular solution containing (in mM) 130 K-Gluconate, 5 KCl, 10 HEPES, 2.5

MgCl$_2$, 4 Na$_2$-ATP, 0.4 Na$_3$-ATP, 10 Na-phosphocreatine and 0.6 EGTA was used,

with a pH of 7.2 and osmolality of 290 mOsmol/L. Resting membrane potential (Vrmp)

was measured immediately after generation of a whole-cell configuration. Further

analysis of active and passive membrane properties was conducted at a holding

potential of -60 mV. Passive membrane properties were determined via voltage steps

of -10 mV. Active intrinsic properties were measured with a stepwise current injection

protocol. Spontaneous postsynaptic currents (sPSCs) and miniature postsynaptic

currents (mPSCs) were measured by 10 min continuous recording at a holding

potential (V_h) of -60 mV. In our in-vitro cultures GABAergic spontaneous and

miniature events are very sparse (Mossink et al., 2021), thus detected PSCs were

considered as mainly reflecting glutamatergic excitatory postsynaptic currents

(EPSCs). sEPSC burst inputs were manually counted using clampfit 10.7. sEPSCs

were grouped as a burst if at least 3 consecutive events occurred within 50 ms, with at

least one of these events showing an amplitude above 100 pA. For the recording of

mEPSCs, 1 µM TTX was added to the recording medium. Recordings were

conducted at either DIV 21 and DIV 49 (intrinsic properties and sEPSCs) or only DIV

49 (mEPSCs). Cells were visualized with an Olympus BX51WI upright microscope

(Olympus Life Science, PA, USA), equipped with a DAGE-MTI IR-1000E (DAGE-MTI,

IN, USA) camera) and a CoolLED PE-200 LED system (Scientifica, Sussex, UK)

which aided in fluorescent identification of GABAergic neurons. A Digidata 1440A

digitizer and a Multiclamp 700B amplifier (Molecular Devices) were used for data

acquisition. Sampling rate was set at 20 kHz (intrinsic properties) or 10 kHz (sEPSCs

and mEPSCs) and a lowpass 1 kHz filter was used during recording. Recordings were

not corrected for liquid junction potential (±10 mV). Recordings were discarded if

series resistance reached >25 MΩ or dropped below a 10:0 ratio of membrane

resistance to series resistance. Intrinsic electrophysiological properties were analyzed

using Clampfit 11.2 (molecular devices, CA, USA), and sEPSCs were analyzed using

MiniAnalysis 6.0.2 (Synaptosoft Inc, GA, USA) as previously described (Mossink et al.,

2021). Regarding the analysis of the intrinsic properties: In brief, the adaptation ratio

was defined as the Δt action potential 8-9/Δt action potential 2-3. The

afterhyperpolarization time was defined as the time from which the repolarization

phase reaches the threshold potential to the time at which the most hyperpolarized

potential was reached. Action potential half-time was calculated as the time to reach

50% ΔmV of the action potential to the AHP peak.

Calcium imaging

For calcium imaging, cultures were incubated with 4 µg/ml Fluo-8-AM for 30 min at

37°C. After incubation, we removed the excess dye by washing the cells 3 times with

HHBS. The cells were then left in culture medium to recover for 15 min. To image the

cells, we placed them under the microscope (SliceScope Pro 2000, Scientifica). We

continuously perfused the recording chamber with oxygenated (95% O_2/ 5% CO_2) and

artificial cerebrospinal fluid (ACSF) that was composed of (in mM) 124 NaCl, 3 KCl,

1.25 NaH_2PO_4, 2 $CaCl_2$, 1 $MgCl_2$, 26 $NaHCO_3$, 10 Glucose and heated to 37°C.

Imaging was performed using a sCMOS camera (Prime BSI Express, Teledyne

Photometrics) controlled by Micro-Manager acquisition software (NIH). Fluo-8-AM in

the cells was excited at 470 nm by LED (KSL470, Rapp OptoElectronic). We recorded

the cells for 2 min with frame rate of 10 Hz. After recording, we analysed the video

using MATLAB (The Math Works, Inc. MATLAB. Version 2020b) with a home-made

script based on Sun et al (Sun and Südhof, 2020). Circular ROIs were chosen by

hand at the centre of the soma with a diameter of 10 μm. We obtained the fluorescent

change over time which is defined as $\Delta F/F = (F-F_0)/F_0$. Furthermore, the decay of

baseline intensity due to bleaching was corrected by exponential fitting to the baseline.

Additionally, the traces of each location were analysed for a synchronous firing rate

among the selected ROIs to determine network patterns.

RNA sequencing

Cells were harvested on DIV49 of neuronal differentiation. For RNA-seq, the prepared

samples were sequenced on an Illumina NovaSeq SP platform at an average depth of

~50 million reads per sample using a read length of 100 base pairs and an insert size

of 350 base pairs. Three biological replicates of control and *SETD1A*$^{+/-}$ E/I networks

and glutamatergic networks respectively (12 samples in total) using the NucleoSpin

RNA isolation kit (Machery Nagel, 740955.250) according to the manufacturer's

instructions. RNA yield was quantified with a NanoDrop Spectrophotometer

(NanoDrop Technologies, Wilmington, DE, USA) and RNA integrity was assessed

with Bioanalyzer 2100 RNA 6000 Nano Kit (Agilent Technologies, Santa Clara, CA,

USA). All samples had an RNA Integrity Number (RIN) > 9. Library preparation and

paired-end RNA-sequencing were carried out at the Norwegian High-Throughput

Sequencing Centre (www.sequencing.uio.no). Briefly, libraries were prepared with the

TruSeq Stranded mRNA kit from Illumina which involves Poly-A purification to capture

coding as well as several non-coding RNAs. The prepared samples were then

sequenced on an Illumina NovaSeq SP platform.

RNA-seq data processing

Raw sequencing reads were quality assessed with FastQC (Babraham Institute). To

pass the initial QC check, the average Phred score of each base position across all

reads had to be at least 30. Reads were further processed by cutting individual

low-quality bases and removing adapter and other Illumina-specific sequences with

Trimmomatic V0.32 using default parameters (Bolger et al., 2014). The trimming

process may result in some reads being discarded and their mates thereby unpaired,

therefore only reads that remained paired after trimming were used for downstream

analyses. Since the cultures contained both hiPSC-derived neurons as well as rat

astrocytes which were added to support neuronal maturation, sequencing reads were

separated according to their species of origin using the *in silico* RNA-seq read sorting

tool Sargasso (Qiu et al., 2018), which was able to successfully eliminate sequencing

reads stemming from rat astrocytes (**Figure S12**). To quantify gene expression levels,

reads mapped by Sargasso were summarized at the gene level using featureCounts

(Liao et al., 2014) guided by ENSEMBL annotations.

Differential expression (DE) analysis and over-representation test

To evaluate the species separation performance of Sargasso, bioinformatical

estimation of cell type abundances (deconvolution) was carried out with

CIBERSORTx (Newman et al., 2019) using expression signatures for human neurons

and rodent astrocytes (Zhang et al., 2016) (**Figure S12**). Before conducting the DE

analyses, genes with very low to zero expression were removed by filtering out any

gene with ≤1 counts per million (CPM) in 3 or more samples (the smallest group size).

DE analysis was performed using the statistical R package DESeq2 (Love et al.,

2014), which provides methods to test for differentially expressed genes by use of

negative binomial generalized models. The DESeq2 workflow begins by taking raw

read count data as input and applies an internal normalization method that corrects

for sequencing depth and RNA composition. The standard DE analysis consists of

size factor estimation, dispersion estimation, and model fitting, as well as an

independent filtering step that optimizes the number of significant DE genes. After the

pre-filtering and independent filtering steps, a total of 14,144 genes were retained and

examined in the DE analyses. A DE gene was considered significant if the FDR was

<0.05. Gene Ontology (GO) enrichment tests of significant DE gene sets were

conducted with the over-representation analysis tool clusterProfile (Yu et al., 2012)

using the enrichGO function. A GO term was considered significantly enriched if the

FDR was <0.05. Disease association analysis was performed with the R package

disgenet2r (Piñero et al., 2020) using both the CURATED database, which includes

more than 10,000 somatic and mental disorders, and the PSYGENET, which includes

109 mental diseases. In both cases, FDR<0.05 was used as the threshold to

determine significant disease association.

Data analysis

The statistical analysis of the data were performed using GraphPad Prism 8

(GraphPad Software, Inc., CA, USA). We ensured normal distribution using a

Shapiro-Wilk normality test. Analysis was done using unpaired Student's t tests when

comparing two variables at a single time point, or one-way ANOVA with sequential

post hoc Bonferroni corrections. Results with p values lower than 0.05 were

considered as significantly different. $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***). Data is

shown as mean ± standard error of the mean (SEM).

Data availability

All the data are available within this manuscript and its Supplementary information.

RNA-seq raw and processed data are available from GEO repository (GEO:

GSE180648). Additional information is available from the corresponding author upon

reasonable request.

Acknowledgements

This work was supported by grants from: ERA-NET NEURON-102 SYNSCHIZ grant

(NWO) 013-17-003 4538 (to D.S) and ERA-NET NEURON DECODE! grant (NWO)

013.18.001 (to N.N.K); The Netherlands Organization for Scientific Research,

NWO-CAS grant 012.200.001 (to N.N.K); the Netherlands Organization for Health

Research and Development ZonMw grant 91217055 (to. H.v.B and N.N.K); SFARI

grant 610264 (to N.N.K); Marie Curie Actions European Fellowship 794273 (to N.K);

South-Eastern Norway Regional Health Authority #2018094 (to S.D); China

scholarship council 201806210076 (to S.W).

Author contribution

S.W, D.S and N.N.K conceived and designed all the experiments. D.S, N.N.K and

H.v.B supervised the study. S.W, J.v.R, I.A, N.K, N.M, A.B, E.L, K.W, C.S performed

all experiments. S.W, J.v.R, I.A, N.K, N.M, A.B, D.S, N.N.K performed data analysis.

S.W, D.S, J.v.R, I.A and N.N.K wrote the manuscript. H.v.B, S.D and T.K edited the

manuscript.

Declaration of interests

The authors declare no competing interests.

Reference

Aglah, C., Gordon, T., and Posse de Chaves, E.I. (2008). cAMP promotes neurite

outgrowth and extension through protein kinase A but independently of Erk activation

in cultured rat motoneurons. Neuropharmacology *55*, 8–17.

Anderson, A.E., Adams, J.P., Qian, Y., Cook, R.G., Pfaffinger, P.J., and Sweatt, J.D.

(2000). Kv4.2 Phosphorylation by cyclic AMP-dependent protein kinase. J. Biol. Chem.

275, 5337–5346.

Beaumont, V., and Zucker, R.S. (2000). Enhancement of synaptic transmission by

cyclic AMP modulation of presynaptic I(h) channels. Nat. Neurosci. *3*, 133–141.

Berry-Kravis, E., Hicar, M., and Ciurlionis, R. (1995). Reduced cyclic AMP production

in fragile X syndrome: Cytogenetic and molecular correlations. Pediatr. Res. *38*,

638–643.

Berry‐Kravis, E., and Huttenlocher, P.R. (1992). Cyclic AMP metabolism in fragile X

syndrome. Ann. Neurol. *31*, 22–26.

Blanco-Suárez, E., Caldwell, A.L.M., and Allen, N.J. (2017). Role of

astrocyte–synapse interactions in CNS disorders. J. Physiol. *595*, 1903–1916.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for

Illumina sequence data. Bioinformatics *30*, 2114–2120.

Carroll, L.S., and Owen, M.J. (2009). Genetic overlap between autism, schizophrenia

and bipolar disorder. Genome Med. *1*, 102.

Carvajal-Oliveros, A., and Campusano, J.M. (2021). Studying the Contribution of

Serotonin to Neurodevelopmental Disorders. Can This Fly? Front. Behav. Neurosci.

14, 12.

Chang, A., Li, P.P., and Warsh, J.J. (2003). Altered cAMP-dependent protein kinase

subunit immunolabeling in post-mortem brain from patients with bipolar affective

disorder. J. Neurochem. *84*, 781–791.

Chu, M.S., Chang, C.F., Yang, C.C., Bau, Y.C., Ho, L.L.T., and Hung, S.C. (2006).

Signalling pathway in the induction of neurite outgrowth in human mesenchymal stem

cells. Cell. Signal. *18*, 519–530.

Connors, E.C., Ballif, B.A., and Morielli, A.D. (2008). Homeostatic regulation of Kv1.2

potassium channel trafficking by cyclic AMP. J. Biol. Chem. *283*, 3445–3453.

Delghandi, M.P., Johannessen, M., and Moens, U. (2005). The cAMP signalling

pathway activates CREB through PKA, p38 and MSK1 in NIH 3T3 cells. Cell. Signal.

17, 1343–1351.

Eells, J. (2005). The Control of Dopamine Neuron Development, Function and

Survival: Insights From Transgenic Mice and The Relevance to Human Disease. Curr.

Med. Chem. *10*, 857–870.

Eising, E., Carrion-Castillo, A., Vino, A., Strand, E.A., Jakielski, K.J., Scerri, T.S.,

Hildebrand, M.S., Webster, R., Ma, A., Mazoyer, B., et al. (2019). A set of regulatory

genes co-expressed in embryonic human brain is implicated in disrupted speech

development. Mol. Psychiatry *24*, 1065–1078.

Epstein, P.M. (2017). Different phosphodiesterases (PDEs) regulate distinct

phosphoproteomes during cAMP signaling. Proc. Natl. Acad. Sci. U. S. A. *114*,

7741–7743.

Evripioti, A.A., Ortega-Prieto, A.M., Skelton, J.K., Bazot, Q., and Dorner, M. (2019).

Phosphodiesterase-induced cAMP degradation restricts hepatitis B virus infection.

Philos. Trans. R. Soc. B Biol. Sci. *374*.

Ford, J.M., Krystal, J.H., and Mathalon, D.H. (2007). Neural Synchrony in

Schizophrenia: From Networks to New Treatments. Schizophr. Bull. *33*, 848–852.

Frega, M., Linda, K., Keller, J.M., Gümüş-Akay, G., Mossink, B., van Rhijn, J.R.,

Negwer, M., Klein Gunnewiek, T., Foreman, K., Kompier, N., et al. (2019). Neuronal

network dysfunction in a model for Kleefstra syndrome mediated by enhanced

NMDAR signaling. Nat. Commun. *10*, 1–15.

Funk, A.J., McCullumsmith, R.E., Haroutunian, V., and Meador-Woodruff, J.H. (2012).

Abnormal activity of the MAPK- and cAMP-associated signaling pathways in frontal

cortical areas in postmortem brain in schizophrenia. Neuropsychopharmacology *37*,

896–905.

Gutierrez-Castellanos, N., Da Silva-Matos, C.M., Zhou, K., Canto, C.B., Renner, M.C.,

Koene, L.M.C., Ozyildirim, O., Sprengel, R., Kessels, H.W., and De Zeeuw, C.I.

(2017). Motor Learning Requires Purkinje Cell Synaptic Potentiation through

Activation of AMPA-Receptor Subunit GluA3. Neuron *93*, 409–424.

Hamm, J.P., Shymkiv, Y., Mukai, J., Gogos, J.A., and Yuste, R. (2020). Aberrant

Cortical Ensembles and Schizophrenia-like Sensory Phenotypes in Setd1a+/− Mice.

Biol. Psychiatry *88*, 215–223.

Hell', J.W., Yokoyama2, C.T., Breeze, L.J., Chavkin, C., and Catterall', W.A. (1995).

Phosphorylation of presynaptic and postsynaptic calcium channels by

cAMP-dependent protein kinase in hippocampal neurons.

Hilker, R., Helenius, D., Fagerlund, B., Skytthe, A., Christensen, K., Werge, T.M.,

Nordentoft, M., and Glenthøj, B. (2018). Heritability of Schizophrenia and

Schizophrenia Spectrum Based on the Nationwide Danish Twin Register. Biol.

Psychiatry *83*, 492–498.

van der Horst, J., Greenwood, I.A., and Jepps, T.A. (2020). Cyclic AMP-Dependent

Regulation of Kv7 Voltage-Gated Potassium Channels. Front. Physiol. *11*.

Kang, H., Han, K., Won, S.Y., Kim, H.M., Lee, Y.H., Ko, J., and Um, J.W. (2016). Slitrk

missense mutations associated with neuropsychiatric disorders distinctively impair

slitrk trafficking and synapse formation. Front. Mol. Neurosci. *9*, 1–18.

Kelley, D.J., Davidson, R.J., Elliott, J.L., Lahvis, G.P., Yin, J.C.P., and Bhattacharyya,

A. (2007). The Cyclic AMP Cascade Is Altered in the Fragile X Nervous System. PLoS

One *2*, e931.

Kelley, D.J., Bhattacharyya, A., Lahvis, G.P., Yin, J.C.P., Malter, J., and Davidson,

R.J. (2008). The cyclic AMP phenotype of fragile X and autism. Neurosci. Biobehav.

Rev. *32*, 1533–1543.

Keravis, T., and Lugnier, C. (2012). Cyclic nucleotide phosphodiesterase (PDE)

isozymes as targets of the intracellular signalling network: Benefits of PDE inhibitors

in various diseases and perspectives for future therapeutic developments. Br. J.

Pharmacol. *165*, 1288–1305.

Khazipov, R., and Luhmann, H.J. (2006). Early patterns of electrical activity in the

developing cerebral cortex of humans and rodents. Trends Neurosci. *29*, 414–418.

Klaassen, R. V., Stroeder, J., Coussen, F., Hafner, A.-S., Petersen, J.D., Renancio,

C., Schmitz, L.J.M., Normand, E., Lodder, J.C., Rotaru, D.C., et al. (2016). Shisa6

traps AMPA receptors at postsynaptic sites and prevents their desensitization during

synaptic activity. Nat. Commun. 2016 71 *7*, 1–12.

Kummeling, J., Stremmelaar, D.E., Raun, N., Reijnders, M.R.F., Willemsen, M.H.,

Ruiterkamp-Versteeg, M., Schepens, M., Man, C.C.O., Gilissen, C., Cho, M.T., et al.

(2020). Characterization of SETD1A haploinsufficiency in humans and Drosophila

defines a novel neurodevelopmental syndrome. Mol. Psychiatry 1–12.

Lee, K.H., Williams, L., Haig, A., and Gordon, E. (2003). "Gamma (40 Hz) phase

synchronicity" and symptom dimensions in schizophrenia. Cogn. Neuropsychiatry *8*,

57–71.

Lee, S.H., Wynn, J.K., Green, M.F., Kim, H., Lee, K.J., Nam, M., Park, J.K., and

Chung, Y.C. (2006). Quantitative EEG and low resolution electromagnetic

tomography (LORETA) imaging of patients with persistent auditory hallucinations.

Schizophr. Res. *83*, 111–119.

Legge, S.E., Santoro, M.L., Periyasamy, S., Okewole, A., Arsalan, A., and Kowalec, K.

(2021a). Genetic architecture of schizophrenia: a review of major advancements.

Psychol. Med. 1–10.

Legge, S.E., Santoro, M.L., Periyasamy, S., Okewole, A., Arsalan, A., and Kowalec, K.

(2021b). Genetic architecture of schizophrenia: a review of major advancements.

Psychol. Med. 1–10.

Liao, Y., Smyth, G.K., and Shi, W. (2014). FeatureCounts: An efficient general

purpose program for assigning sequence reads to genomic features. Bioinformatics

30, 923–930.

Lisman, J.E. (1997). Bursts as a unit of neural information: making unreliable

synapses reliable. Trends Neurosci. *20*, 38–43.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Mandegar, M.A., Huebsch, N., Frolov, E.B., Shin, E., Truong, A., Olvera, M.P., Chan,

A.H., Miyaoka, Y., Holmes, K., Spencer, C.I., et al. (2016). CRISPR Interference

Efficiently Induces Specific and Reversible Gene Silencing in Human iPSCs. Cell

Stem Cell *18*, 541–553.

Millar, J.K., Pickard, B.S., Mackie, S., James, R., Christie, S., Buchanan, S.R., Malloy,

M.P., Chubb, J.E., Huston, E., Baillie, G.S., et al. (2005). Genetics: DISC1 and

PDE4B are interacting genetic factors in schizoprenia that regulate cAMP signaling.

Science (80-.). *310*, 1187–1191.

Miyaoka, Y., Chan, A.H., Judge, L.M., Yoo, J., Huang, M., Nguyen, T.D., Lizarraga,

P.P., So, P.L., and Conklin, B.R. (2014). Isolation of single-base genome-edited

human iPS cells without antibiotic selection. Nat. Methods *11*, 291–293.

Mossink, B., van Rhijn, J.-R., Wang, S., Linda, K., Vitale, M.R., Zöller, J.E.M., van

Hugte, E.J.H., Bak, J., Verboven, A.H.A., Selten, M., et al. (2021). Cadherin-13 is a

critical regulator of GABAergic modulation in human stem-cell-derived neuronal

networks. Mol. Psychiatry 1–18.

Mukai, J., Cannavò, E., Crabtree, G.W., Sun, Z., Diamantopoulou, A., Thakur, P.,

Chang, C.Y., Cai, Y., Lomvardas, S., Takata, A., et al. (2019). Recapitulation and

Reversal of Schizophrenia-Related Phenotypes in Setd1a-Deficient Mice. Neuron *104*,

471-487.e12.

Muly, C. (2002). Signal transduction abnormalities in schizophrenia: the cAMP system.

Psychopharmacol. Bull. *36*, 92–105.

Murray, A.J. (2008). Pharmacological PKA inhibition: All may not be what it seems.

Sci. Signal. *1*, re4–re4.

Nagahama, K., Sakoori, K., Watanabe, T., Kishi, Y., Kawaji, K., and Koebis, M. (2020).

Article Setd1a Insufficiency in Mice Attenuates Excitatory Synaptic Function and

Recapitulates Schizophrenia- Related Behavioral Abnormalities ll Setd1a

Insufficiency in Mice Attenuates Excitatory Synaptic Function and Recapitulates

Schizophrenia-Related . CellReports *32*, 108126.

Namkoong, S., Kim, C.K., Cho, Y.L., Kim, J.H., Lee, H., Ha, K.S., Choe, J., Kim, P.H.,

Won, M.H., Kwon, Y.G., et al. (2009). Forskolin increases angiogenesis through the

coordinated cross-talk of PKA-dependent VEGF expression and Epac-mediated

PI3K/Akt/eNOS signaling. Cell. Signal. *21*, 906–915.

Newman, A.M., Steen, C.B., Liu, C.L., Gentles, A.J., Chaudhuri, A.A., Scherer, F.,

Khodadoust, M.S., Esfahani, M.S., Luca, B.A., Steiner, D., et al. (2019). Determining

cell type abundance and expression from bulk tissues with digital cytometry. Nat.

Biotechnol. *37*, 773–782.

Owen, M.J., Sawa, A., and Mortensen, P.B. (2016). Schizophrenia. Lancet *388*,

86–97.

Pieroni, J.P., Harry, A., Chen, J., Jacobowitz, O., Magnusson, R.P., and Iyengar, R.

(1995). Distinct characteristics of the basal activities of adenylyl cyclases 2 and 6. J.

Biol. Chem. *270*, 21368–21373.

Piñero, J., Ramírez-Anguita, J.M., Saüch-Pitarch, J., Ronzano, F., Centeno, E., Sanz,

F., and Furlong, L.I. (2020). The DisGeNET knowledge platform for disease genomics:

2019 update. Nucleic Acids Res. *48*, D845–D855.

Qiu, J., Dando, O., Baxter, P.S., Hasel, P., Heron, S., Simpson, T.I., and Hardingham,

G.E. (2018). Mixed-species RNA-seq for elucidation of non-cell-autonomous control

of gene transcription. Nat. Protoc. *13*, 2176–2199.

Qiu, L., LeBel, R.P., Storm, D.R., and Chen, X. (2016). Type 3 adenylyl cyclase: A key

enzyme mediating the cAMP signaling in neuronal cilia. Int. J. Physiol. Pathophysiol.

Pharmacol. *8*, 95–108.

Ren, X., Rizavi, H.S., Khan, M.A., Bhaumik, R., Dwivedi, Y., and Pandey, G.N. (2014).

Alteration of cyclic-AMP response element binding protein in the postmortem brain of

subjects with bipolar disorder and schizophrenia. J. Affect. Disord. *152–154*,

326–333.

Renner, M.C., Albers, E.H., Gutierrez-Castellanos, N., Reinders, N.R., van Huijstee,

A.N., Xiong, H., Lodder, T.R., and Kessels, H.W. (2017). Synaptic plasticity through

activation of GluA3-containing AMPA-receptors. Elife *6*.

Sassone-Corsi, P. (2012). The Cyclic AMP pathway. Cold Spring Harb. Perspect. Biol.

4.

Schiller, M., Dennler, S., Anderegg, U., Kokot, A., Simon, J.C., Luger, T.A., Mauviel,

A., and Böhm, M. (2010). Increased cAMP levels modulate transforming growth

factor-β/smad-induced expression of extracellular matrix components and other key

fibroblast effector functions. J. Biol. Chem. *285*, 409–421.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source

platform for biological-image analysis. Nat. Methods *9*, 676–682.

Singh, T., Kurki, M.I., Curtis, D., Purcell, S.M., Crooks, L., McRae, J., Suvisaari, J.,

Chheda, H., Blackwood, D., Breen, G., et al. (2016). Rare loss-of-function variants in

SETD1A are associated with schizophrenia and developmental disorders. Nat.

Neurosci. *19*, 571–577.

Singh, T., Neale, B.M., and Daly, M.J. (2020). Exome sequencing identifies rare

coding variants in 10 genes which confer substantial risk for schizophrenia on behalf

of the Schizophrenia Exome Meta-Analysis (SCHEMA) Consortium*. MedRxiv

2020.09.18.20192815.

Skene, N.G., Bryois, J., Bakken, T.E., Breen, G., Crowley, J.J., Gaspar, H.A.,

Giusti-Rodriguez, P., Hodge, R.D., Miller, J.A., Muñoz-Manchado, A.B., et al. (2018).

Genetic identification of brain cell types underlying schizophrenia. Nat. Genet. *50*,

825–833.

Song, J., Cheon, S.Y., Lee, W.T., Park, K.A., and Lee, J.E. (2015). PKA Inhibitor H89

(N-[2-p-bromocinnamylamino-ethyl]-5-isoquinolinesulfonamide) Attenuates Synaptic

Dysfunction and Neuronal Cell Death following Ischemic Injury. Neural Plast. *2015*.

Spencer, K.M., Niznikiewicz, M.A., Nestor, P.G., Shenton, M.E., and McCarley, R.W.

(2009). Left auditory cortex gamma synchronization and auditory hallucination

symptoms in schizophrenia. BMC Neurosci. *10*, 85.

Sun, Z., and Südhof, T.C. (2020). A simple Ca2+-imaging approach to neural network

analysis in cultured neurons. BioRxiv 2020.08.09.243576.

Szklarczyk, D., Gable, A.L., Nastou, K.C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva,

N.T., Legeay, M., Fang, T., Bork, P., et al. (2021). The STRING database in 2021:

Customizable protein-protein networks, and functional characterization of

user-uploaded gene/measurement sets. Nucleic Acids Res. *49*, D605–D612.

Takata, A., Xu, B., Ionita-Laza, I., Roos, J.L., Gogos, J.A., and Karayiorgou, M. (2014).

Loss-of-Function Variants in Schizophrenia Risk and SETD1A as a Candidate

Susceptibility Gene. Neuron *82*, 773–780.

Tasken, Y., Skhlhegghl, S., Solberg$, R., Anderssonll, K.B., Taylor, S.S., Lean, T., Kiil

Blomhofnl, H., Jahnsens, T., and Hansson, V. (1993). THE JOURNAL OF

BIOLOGICAL CHEMISTRY Novel Isozymes of cAMP-dependent Protein Kinase Exist

in Human Cells Due to Formation of RIa-RIP Heterodimeric Complexes*.

Thornquist, S.C., Pitsch, M.J., Auth, C.S., and Crickmore, M.A. (2021). Biochemical

evidence accumulates across neurons to drive a network-level eruption. Mol. Cell *81*,

675-690.e8.

Tsai, G., and Coyle, J.T. (2002). Glutamatergic mechanisms in schizophrenia. Annu.

Rev. Pharmacol. Toxicol. *42*, 165–179.

Turetsky, B.I., and Moberg, P.J. (2009). An odor-specific threshold deficit implicates

abnormal intracellular cyclic AMP signaling in schizophrenia. Am. J. Psychiatry *166*,

226–233.

Uhlhaas, P.J., and Singer, W. (2010). Abnormal neural oscillations and synchrony in

schizophrenia. Nat. Rev. Neurosci. *11*, 100–113.

Vacic, V., McCarthy, S., Malhotra, D., Murray, F., Chou, H.H., Peoples, A., Makarov,

V., Yoon, S., Bhandari, A., Corominas, R., et al. (2011). Duplications of the

neuropeptide receptor gene VIPR2 confer significant risk for schizophrenia. Nature

471, 499–503.

Wan, G., Zhou, L., Lim, Q.E., Wong, Y.H., and Too, H.P. (2011). Cyclic AMP

signalling through PKA but not Epac is essential for neurturin-induced biphasic

ERK1/2 activation and neurite outgrowths through GFRα2 isoforms. Cell. Signal. *23*,

1727–1737.

Wang, H., and Storm, D.R. (2003). Calmodulin-regulated adenylyl cyclases:

Cross-talk and plasticity in the central nervous system. Mol. Pharmacol. *63*, 463–468.

Wang, H., Xu, J., Lazarovici, P., Quirion, R., and Zheng, W. (2018). cAMP Response

Element-Binding Protein (CREB): A Possible Signaling Molecule Link in the

Pathophysiology of Schizophrenia. Front. Mol. Neurosci. *11*, 255.

Yim, Y.S., Kwon, Y., Nam, J., Yoon, H.I., Lee, K., Kim, D.G., Kim, E., Kim, C.H., and

Ko, J. (2013). Slitrks control excitatory and inhibitory synapse formation with LAR

receptor protein tyrosine phosphatases. Proc. Natl. Acad. Sci. U. S. A. *110*,

4057–4062.

Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). ClusterProfiler: An R package for

comparing biological themes among gene clusters. Omi. A J. Integr. Biol. *16*,

284–287.

Yu, X., Yang, L., Li, J., Li, W., Li, D., Wang, R., Wu, K., Chen, W., Zhang, Y., Qiu, Z.,

et al. (2019). De Novo and Inherited SETD1A Variants in Early-onset Epilepsy.

Neurosci. Bull. *35*, 1045–1057.

Zamarbide, M., Mossa, A., Muñoz-Llancao, P., Wilkinson, M.K., Pond, H.L., Oaks,

A.W., and Manzini, M.C. (2019). Male-Specific cAMP Signaling in the Hippocampus

Controls Spatial Memory Deficits in a Mouse Model of Autism and Intellectual

Disability. Biol. Psychiatry *85*, 760–768.

Zhang, Y., Sloan, S.A., Clarke, L.E., Caneda, C., Plaza, C.A., Blumenthal, P.D., Vogel,

H., Steinberg, G.K., Edwards, M.S.B., Li, G., et al. (2016). Purification and

Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional

and Functional Differences with Mouse. Neuron *89*, 37–53.

1001

1002   **Figure legends**

1003

Figure 1. *SETD1A^{+/-}* neuronal networks exhibit dysregulated functional

organization. (a) Generation of *SETD1A* isogenic lines (*SETD1A^{+/-}*) using

CRISPR/Cas9. Schematic diagram showing the position of sgRNA sequence and

indels generated in *SETD1A^{+/-}* clone1 and *SETD1A^{+/-}* clone2. **(b-c)** qPCR and

Western Blot show reduced mRNA (b) and protein level of SETD1A (c). **(d)** Schematic

representation of neuronal differentiation workflow. **(e)** Immunofluorescence staining

of GABA (green) and MAP2 (red) at DIV49. Scale bar: 30 μm. **(f-g)** Analysis of

neuronal activity using MEA recording. **(f)** Representative raster plots (1 min) of

electrophysiological activity exhibited by control and *SETD1A^{+/-}* neuronal networks at

different time points during development. **(g)** Quantification of network parameters as

indicated. Sample size: control n = 40 MEA wells. *SETD1A^{+/-}* n = 63 MEA wells from 5

independent batches. Representative whole-cell voltage-clamp recordings and

quantitative analyses of **(h)** spontaneous postsynaptic currents (sEPSCs) and **(i)**

correlated synaptic inputs (sEPSC bursts) in glutamatergic and GABAergic neurons in

control and *SETD1A^{+/-}* E/I cultures at DIV49. Glutamatergic neurons: control (n = 15),

SETD1A^{+/-} (n = 13); GABAergic neurons: control (n = 15), *SETD1A^{+/-}* (n = 12). **(j)**

Representative whole-cell voltage-clamp recordings and quantitative analyses of

mEPSC activity in glutamatergic and GABAergic neurons. Glutamatergic neurons:

control (n = 17), *SETD1A^{+/-}* (n = 16); GABAergic neurons: control (n = 16),

SETD1A^{+/-} (n = 13). Data represent means ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$,

Students' T-test with Bonferroni correction for multiple testing (g, h-j) or

Kolmogorov-Smirnov test (h) for comparing control vs. *SETD1A^{+/-}* cultures.

1026

Figure 2. *SETD1A*$^{+/-}$ neurons show aberrant somatodendritic morphology.

(a-b) Representative somatodendritic reconstructions of glutamatergic (a) or

GABAergic (b) neurons and Sholl analysis in control and *SETD1A*$^{+/-}$ neurons at DIV21

(Glutamatergic neurons: n = 20 for control, n = 20 for *SETD1A*$^{+/-}$; GABAergic neurons:

n = 18 for control, n = 21 for *SETD1A*$^{+/-}$). **(c-d)** Representative somatodendritic

reconstructions of glutamatergic (c) or GABAergic (d) neurons and Sholl analysis in

control and *SETD1A*$^{+/-}$ networks at DIV49 (Glutamatergic neurons: n = 38 for control,

n = 47 for *SETD1A*$^{+/-}$; GABAergic neurons: n = 33 for control, n = 45 for *SETD1A*$^{+/-}$).

(e-f) Main morphological parameters in reconstruction for glutamatergic neurons (e)

and GABAergic neurons (f) at both DIV21 and DIV49. **(g)** Representative images of

immunocytochemistry stained for glutamatergic synapse: Synapsin as a presynaptic

marker and Homer1 as a postsynaptic marker, and GABAergic synapse: VGAT as a

pre-synaptic marker and Gephyrin as a post synaptic marker. Scale bar = 2 μm **(h)**

Quantification of the density of co-localised Synapsin/Homer1 and VGAT/Gephyrin

puncta (number per 10 μm). (Synapsin/Homer1: n = 25 for control, n = 21 for

SETD1A$^{+/-}$ at DIV21, n = 34 for control, n = 36 for *SETD1A*$^{+/-}$ at DIV49;

VGAT/Gephyrin: n = 18 for control, n = 21 for *SETD1A*$^{+/-}$ at DIV21, n = 34 for control, n

= 36 for *SETD1A*$^{+/-}$ at DIV49.) Data represent means ± SEM. $^{*}p < 0.05$, $^{**}p < 0.01$,

$^{***}p < 0.001$, two-way ANOVA with post hoc Bonferroni correction (a-d), unpaired

Student's t test (e-f) or one-way ANOVA with post hoc Bonferroni correction (h).

1047

Figure 3. *SETD1A* haploinsufficiency leads to an altered transcriptomic profile.
(**a**) PCA showing tight clustering of 3 replicates for each genotype. (**b**) Volcano plots showing differentially expressed genes (DEGs) between *SETD1A$^{+/-}$* and control E/I cultures. Relative to control, significantly up or down-regulated genes are shown in red. Top 3 upregulated and downregulated DEGs are labeled. (**c**) Disease terms of DisGeNET database associated with DEGs. (**d**) Gene Ontology (GO) term analysis of DEGs. (**e**) Diagram for dysregulated protein network showing the interactions among several synaptic functions using STRING. (**f**) Heatmap showing the fold change of DEGs compared to control in gene sets related to glutamatergic synapse.

Figure 4. *SETD1A$^{+/-}$* glutamatergic neuronal cultures recapitulate the *SETD1A$^{+/-}$* E/I network phenotype (a) Schematic of neuronal differentiation workflow. (**b**) Representative raster plots (1 min) of electrophysiological activity measured by MEA from control and *SETD1A$^{+/-}$* glutamatergic neuronal networks at different time points during development. (**c**) Quantification of network parameters as indicated at DIV49. Sample size: control n = 35 MEA wells. *SETD1A$^{+/-}$* n = 37 MEA wells from 4 independent batches. (**d-e**) Representative somatodendritic reconstructions of glutamatergic neurons and Sholl analysis in control and *SETD1A$^{+/-}$* networks at DIV 49 (n = 38 for control. n = 33 for *SETD1A$^{+/-}$*). Scale bar = 40 µm. (**f**) Main morphological parameters for reconstructions of glutamatergic neurons at DIV49. (**g**) Representative voltage-clamp recordings of spontaneous inputs (sEPSCs) onto glutamatergic neurons at DIV49 activity and the corresponding quantitative analyses of sEPSC frequency (**h**) and amplitude **(i). (j)** Representative voltage-clamp recordings of correlated synaptic inputs (sEPSC bursts) in control and *SETD1A$^{+/-}$* glutamatergic neurons at DIV49 and the corresponding quantitative analyses **(k)**. n = 24/17 for control/*SETD1A$^{+/-}$* DIV49. Data represent means ± SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, two-way ANOVA with post hoc Bonferroni correction (e), unpaired Student's t test (c, f) or Students' T-test with Bonferroni correction for multiple testing (h, i, k).

1077

1078

Figure 5. *SETD1A* **haploinsufficiency leads to activation of cAMP/PKA/CREB**

pathway. (a) GO terms related to second messenger signaling associated with

upregulated differentially expressed genes (DEGs) **(b)** Change of mRNA levels of

genes coding for adenylyl cyclase (AC) and phosphodiesterases. **(c)** Schematic of

cAMP/PKA/CREB pathway and drug target at different steps. **(d-e)** Representative

images showing pCREB expression and quantification of intensity of pCREB for

control (n = 105 cells), control + forskolin (n = 30 cells) and *SETD1A*$^{+/-}$ neurons (n =

122 cells from 2 independent batches). Scale bar = 10 µm. **(f-g)** Representative raster

plot for 30 sec recorded by MEA and quantification of network burst rate showing the

effect of AC agonist forskolin on control networks (sample size: control n=10 wells,

control + forskolin n = 12 wells, *SETD1A*$^{+/-}$ n = 28 wells). **(h-i)** Representative raster

plot for 30 sec recorded by MEA and quantification of network burst rate showing the

effect of AC inhibitor SQ22536 on *SETD1A*$^{+/-}$ E/I cultures (sample size: control n=11

wells, *SETD1A*$^{+/-}$ + SQ22536 n = 12 wells, *SETD1A*$^{+/-}$ n =13 wells). **(j-k)**

Representative raster plot for 30 sec recorded by MEA and quantification of network

burst rate showing the effect of PKA inhibitor H89 and KT5720 on *SETD1A*$^{+/-}$ E/I

cultures (sample size: control n=15 wells, *SETD1A*$^{+/-}$ + H89 n = 18 wells, *SETD1A*$^{+/-}$ +

KT5820 n = 18 wells, *SETD1A*$^{+/-}$ n = 22 wells). Data represent means ± SEM. *P

<0.05, **P < 0.01, ***P < 0.001, one-way ANOVA test and post-hoc Bonferroni

correction.