1

TnCentral: A Prokaryotic Transposable Element Database and Web Portal for Transposon
Analysis

Karen Ross[1§], Alessandro M. Varani[2§], Erik Snesrud[3*], Hongzhan Huang[4], Danillo Oliveira
Alvarenga[2], Jian Zhang[1], Cathy Wu[4], Patrick McGann[3], Mick Chandler[5]

[1]Protein Information Resource, Department of Biochemistry and Molecular & Cellular
Biology, Georgetown University Medical Center, Washington, DC, USA.
14
15 [2] School of Agricultural and Veterinary Sciences, Universidade Estadual Paulista, Jaboticabal,
16 Sao Paulo, Brazil.
17
18 [3]Multidrug-Resistant Organism Repository and Surveillance Network, Walter Reed Army
19 Institute of Research, Silver Spring, Maryland, USA.
20
21 [4] Protein Information Resource, Center for Bioinformatics and Computational Biology,
22 University of Delaware, USA
23
24 [5]Department of Biochemistry and Molecular & Cellular Biology, Georgetown University
25 Medical Center, Washington, DC, USA
26
27 [§]These authors contributed equally to this work.
28
29 *We would like to dedicate this article to the memory of Erik Snesrud who was
30 instrumental in initiating this work but was sadly unable to see it completed.

34 Keywords: Mobile Genetic Elements, genome evolution, antibiotic resistance, virulence

Importance

The ability of bacteria to undergo rapid evolution and adapt to changing environmental circumstances drives the public health crisis of multiple antibiotic resistance as well as outbreaks of disease in economically important agricultural crops and animal husbandry. Prokaryotic transposable elements (TE) play a critical role in this. Many carry "passenger genes" (not required for the transposition process) conferring resistance to antibiotics or heavy metals or causing disease in plants and animals. Passenger genes are spread by the normal TE transposition activities, by insertion into plasmids which then spread via conjugation within and across bacterial populations. Thus, an understanding of TE composition and transposition mechanisms is key to developing strategies to combat bacterial pathogenesis. Toward this end, we have developed TnCentral, a bioinformatics resource dedicated to describing and exploring the structural and functional features of prokaryotic TE and whose use is intuitive and accessible to users with or without bioinformatics expertise.

Abstract

We describe here the structure and organization of TnCentral (https://tncentral.proteininformationresource.org/), a web resource for prokaryotic transposable elements (TE). TnCentral currently contains ~400 carefully annotated TE, including transposons from the Tn*3,* Tn*7,* Tn*402* and Tn*554* families, compound transposons, integrons and associated insertion sequences (IS). These TE carry passenger genes, including genes conferring resistance to over 25 classes of antibiotics and nine types of heavy metal as well as genes responsible for pathogenesis in plants, toxin/antitoxin gene pairs, transcription factors and genes involved in metabolism. Each TE has its own entry page providing details about its transposition genes, passenger genes, and other sequence features required for transposition as well as a graphical map of all features. TnCentral content can be browsed and queried through text and sequence-based searches with a graphic output. We describe three use cases, which illustrate how the search interface, results tables, and entry pages can be used to explore and compare TEs.

TnCentral also includes downloadable software to facilitate user-driven identification, with manual annotation, of certain types of TE in genomic sequences.

65 Through the TnCentral homepage, users can also access TnPedia which provides
66 comprehensive reviews of the major TE families including an extensive general section, and
67 specialised sections with descriptions of insertion sequence and transposon families.
68 TnCentral and TnPedia are intuitive resources that can be used by clinicians and scientists to
69 assess TE diversity in clinical, veterinary and environmental samples.
70

71 **Introduction**
72
Transposable elements (TE) are key facilitators of bacterial evolution and adaptation
and central players in the emergence of antibiotic and heavy metal resistance and to the
transmission of virulence and pathogenic traits. Some TE can capture "passenger genes"
(genes not involved in the transposition process) encoding these traits and transmit them to
plasmids, where they accumulate and are then transferred within and between bacterial
populations by conjugation. TE also contribute significantly to the on-going reorganization of
bacterial genomes giving rise to new strains that are more adept at proliferating in clinical
and agricultural environments, as well as in natural ecosystems.
Understanding TE nature, distribution, and activity is therefore an indispensable part
of the struggle to cope with the public health crisis of multiple antibiotic resistance (ABR)
[1,2]. To understand the impact of TE on bacterial populations, it is essential to provide a
detailed description and catalog of TE structures and diversity. The simplest TE, known as
Insertion Sequences (IS), have a profound impact on genome organization and function (see
[3–7]) but do not themselves generally carry integrated passenger genes. There are a large
number of significantly more complex TE (Figure 1), arguably even more important in the
global emergence of ABR and other virulence and pathogenicity traits. These are generically
called transposons and may carry multiple passenger genes, including some of the most
clinically important antibiotic resistance genes. Like IS, these TE are grouped into a number
of distinct families with characteristic organizations [3]. Their transposition activities
facilitate the rapid spread of groups of antibiotic resistance genes and promote their
horizontal transfer. Yet another important aspect of their impact is their ability to assemble
passenger genes into resistance clusters [8,9]. While there appears to be a wide-spread
appreciation that mobile plasmids are responsible for the spread of antibiotic resistance, it

is less well-known that IS and transposons are the conduits that transfer this information
between chromosomes and plasmids.

There are a number of other bioinformatics resources that cover aspects of
prokaryotic TE biology. These include databases for TE passenger genes such as antibiotic
resistance (CARD [10], ARDB [11]) or toxin/antitoxin gene pairs (TADB [12], TASmania [13])
as well as the various classes of TE themselves such as insertion sequences (ISfinder [14]),
integrons (INTEGRALL [15]), integrative conjugative elements, ICE (ICEberg [16,17]),
plasmids (PlasmidFinder [18]) or more general databases which include a variety of these
genome components (ACLAME [19–21]). However, there is a need for a resource that
collects, compares and collates detailed information on the various different classes of TE
that are responsible for the transmission of medically and economically important
passenger genes in an intuitive and accessible way.

Here, we describe TnCentral (https://tncentral.proteininformationresource.org/), a
database of detailed structural and functional information on bacterial TE. Additionally,
TnCentral provides access to TnPedia (https://tnpedia.fcav.unesp.br/), a comprehensive
encyclopedia describing the current state of our knowledge of the biology of IS and
transposons. Together, TnCentral and TnPedia provide a detailed description of TE diversity
with easy-to-understand graphics outputs that are accessible to users without significant
bioinformatic knowledge. They allow users to rapidly analyse the landscape of TE in
genomes (chromosomes and plasmids) isolated from clinical, veterinary and environmental
samples.

Results

TnCentral Website Content

As of May 2021, TnCentral contains information on ~400 TE. About half of these TE are *Tn3*-
family transposons. The remainder are integrons, compound transposons, transposons from
the *Tn402*, *Tn554* and *Tn7* families, and IS that are associated with TE or are part of
compound transposons (Supplementary Table 1). They include TE with resistance to over 25
different classes of antibiotics and nine different heavy metals. The collection also contains
TE that carry a toxin/antitoxin system for bacterial plasmid maintenance [22–24] and TE
from xanthomonads carrying genes for plant pathogenicity.

TnCental Web Portal

The TnCentral home page is designed to give the user easy access to the contents of TnCentral with a number of options (Figure 2A), including:

TnCentral Search *(search of the TnCentral database),*

Sequence Search *(BLAST-like search for sequence similarities in the database),*

Browse Tn list *(view all TE in TnCentral),*

Tnfinder Software *(access to downloadable scripts for identifying potential TE in sequence databases),*

Documentation *(downloadable documentation for TnCentral),*

For Curators *(detailed curation guidelines),*

TnPedia *(TE Encyclopedia),*

Related links, *and*

Feedback.

TnCentral Search. The interface provides a variety of search functions divided into two search types: *Transposon search* and *Gene search* (Figure 2B).

Transposon search. The transposon collection can be searched using the transposon **name**, **synonyms** which may have been used in the literature, the **type** of mobile genetic element (e.g., insertion sequence, transposon or integron), the **family** and **subgroup** to which it belongs, the **host organism**, **country** of identification and **date** of identification. The latter three search terms are intended for use in epidemiological tracking. These search terms result in a table that can be sorted, customized and downloaded (See Use Case #1, below).

Gene search. It is also possible to search for TE-associated genes by name, by class (Transposase, Accessory Gene or Passenger Gene) or by function (Antibiotic Resistance, Heavy Metal Resistance) and to retrieve information on the transposons in which they are found (see Use Case #2, below).

Sequence Search. Sequence Search allows users to perform sequence similarity

searches using BLAST [26,27] (see Use Case #3, below). By default, the search database is
the TnCentral database, but the page also provides links to BLAST against the ISfinder
(https://isfinder.biotoul.fr/blast.php),   NCBI   https://blast.ncbi.nlm.nih.gov/Blast.cgi),
Comprehensive   Antibiotic   Resistance   Database   (CARD;
https://card.mcmaster.ca/analyze/blast) and the Toxin-Anitoxin (TADB;
https://bioinfo-mml.sjtu.edu.cn/TADB2/) databases. The BLAST tool automatically
distinguishes between DNA and protein query sequences.

Browse Tn list. This option allows the user to browse the entire TnCentral database.

The Transposon entry page. All of the search and browse options provide links to entry
pages for each TE (Figure 3), which provide detailed information about TE features and
origins. The   page   includes: 1)   host   information: host   species,   strain,   and
plasmid/chromosome in which the transposon was found as well as the date and geographic
location of the isolate; 2) a graphic representation of the annotated sequence with color-
coded features; 3) Terminal Inverted Repeats (IR); 4) DNA sequence; 5) internal
recombination sites (e.g. *res* sites) including their coordinates, length and DNA sequence; 6)
ORF summary, which includes all protein coding genes in the order in which they appear, 5'-
3', in the TE sequence, the element with which they are associated (important for nested TE
in which one TE is inserted into another), their coordinates, their class (e.g., Transposase,
Accessory Gene, Passenger Gene) and subclass (e.g., Antibiotic Resistance, Heavy Metal
Resistance) and their relative orientation within the TE; 7) a detailed ORF description
including the amino acid sequence; 8) if applicable, a table of Internal Transposable
Elements (TE inserted in the main element) including the name, type location and length; 9)
if applicable, a table of Internal Repeats (repeat elements, other than the terminal inverted
repeats, that are found within the TE), including the associated TE, coordinates and DNA
sequence; 10) Bibliographic references with direct links to PubMed [25]. Each section can be
collapsed using a button on the right-hand side of the section heading. Sections can be
viewed either by scrolling down on the page or by clicking on the section name in the menu
located on the left side of the page. Sequence files in FASTA and GenBank format can be
downloaded using the links on the left side of the page under the menu.

Tnfinder Software. This section provides three user-downloadable scripts written in-house

for identifying transposons. They provide users with local control over analyses and help

them screen datasets containing large numbers of genomic sequences using their own

servers for identifying potential candidates and which are then manually curated.

Tn3 Transposon Finder (Tn3_finder) performs the automatic prediction of

transposable elements of the Tn*3* family in bacteria and archaea. It compares user-provided

bacterial and archaeal genome sequences to custom Tn*3* transposase and resolvase

databases by BLAST alignments. The criteria for identifying potential transposon regions

according to similarity, coverage and distance values can be adjusted by the user. Additional

ORFs that might be related to passenger genes are also predicted, and flanking regions can

also be retrieved and analyzed. The automatic prediction results are written in report files

and pre-annotated GenBank files to help in subsequent manual curation. Tn*3*_finder allows

for the concurrent analysis of multiple genomes by multithreading.

Composite Transposon Finder (TnComp_finder) predicts the putative composite

transposons in bacterial and archaeal genomes based on insertion sequence replicas in a

relatively short span. It works by comparing nucleotide sequences from bacterial and

archaeal genomes to a custom transposon database and identifying duplicated transposons

in user-defined genomic regions from BLAST alignments. Similar to Tn3_finder,

multithreaded analyses of multiple genomes are available and parameters for similarity,

coverage, distance and flanking regions can be adjusted by the user. Results are written in

report files and pre-annotated GenBank files to help in subsequent manual curation.

Antibiotic Resistance Gene-associated IS Finder (ISAbR_finder) is an experimental

program for the automatic prediction of antibiotic resistance genes associated with known

IS elements derived from the ISfinder database and has yet to be tested extensively. It

works by comparing IS nucleotide sequences from bacterial and archaeal genomes to a

custom antibiotic resistance database based on the parsing of BLAST alignment results,

using a number of parameters that can be customized by the user for stricter or more

relaxed criteria and allowing multithreaded alignments of multiple genomes. ISAbR_finder

also produces report files and pre-annotated GenBank files on which the recommended
manual curation should be performed.

Documentation. This section, which can be downloaded as a pdf file, provides a short
background description of transposons and TnCentral together with a short description of
the **curation workflow** and of planned future developments.

For Curators. This section provides a detailed description of the curation workflow used in
generating the annotated TnCentral data.

TnPedia. TnCentral provides access from the homepage to TnPedia, an online knowledge
base which contains information concerning transposition in prokaryotes. TnPedia is
developed using MediaWiki (https://www.mediawiki.org) and can also be accessed directly
(https://tnpedia.fcav.unesp.br/). It is structured into three main sections: **General**
Information, IS Families and Transposon Families (Figure 4).

The **General Information** section provides a series of clickable sections with an
extensive bibliography and direct links to the articles in PubMed. It includes a historical
perspective, definitions and descriptions of a variety of prokaryotic TE, the basic
mechanisms involved in their movement and the enzymes involved in these processes. It
also contains information describing their impact on their host genomes and how their
activities are controlled.

The **IS Families** section consists of individual chapters describing each of the ~25 IS
families in detail and covers, where possible, the identification of the founding members,
their organisation, distribution, variability and phylogenetic relationships, regulation of their
transposition, impact on their host genomes, and their transposition mechanisms including
genetic, biochemical and structural studies.

The **Transposon Families** section describes each transposon family with similar
information to that included in the IS family descriptions but, in addition, including a
detailed description of their structures and the passenger genes which they may carry.

Examples of TnCentral Use

Use Case #1: Comparing Protein Coding Genes in Tn*554* Family Members

The Tn*554* family is a small family restricted to the Firmicutes. Members encode three

genes, *tnpA*, *tnpB* and *tnpC*, involved in transposition [28,29]

(https://tnpedia.fcav.unesp.br/index.php/Transposons_families/Tn554_family). TnpA and

TnpB both exhibit a C-terminal motif which shares all the important catalytic residues of a

typical tyrosine site-specific recombinase [28,29]. They insert in a sequence-specific way

into the DNA repair gene *radC* [30,31] and can also be found in a circular form [32–36]. To

compare the protein coding genes in Tn*554* family members side by side, we searched for

Tn*554* in the TE family field of the Transposon Search interface (Figure 5A). Fourteen Tn*554*

family members were found (of which only 10 are shown in Figure 5B). In order to perform a

side-by-side comparison of the protein-coding genes in these TE, we used the Customize

Display option on the search results page, to add the "All Gene Fields" columns, which

provide information about the protein coding genes, to the display and to remove several

columns (e.g., Host Organism, Country) (Figure 5B). Results for two of the Tn*554*

transposons (Tn*558.3* and Tn*559*) are shown in Figure 5C. Both transposons have the three-

part transposition module (*tnpA, tnpB, tnpC*) characteristic of the family. However, the two

transposons are quite diverse in their passenger genes. Tn*558.3* has gene called *fla*, which

contains a flavodoxin-like domain, and the ABR gene *fexA*, which confers resistance to

phenicol antibiotics. Tn*559* has just a single passenger gene, the ABR gene, *dfrK*, which

confers resistance to diaminopyrimidine antibiotics. As shown by this example, the flexible

search results page makes it easy to compare features across multiple transposons.

Use Case #2: Type II Toxin/Antitoxin Systems in Tn*3* Transposons

Toxin/Antitoxin (TA) systems are implicated in plasmid maintenance in bacterial populations

[37]. These systems are characterized by a stable toxin and an unstable antitoxin that binds

to the toxin and inhibits its lethal effect. Loss of a plasmid carrying a TA system will lead to

rapid depletion of the antitoxin, allowing the persistent toxin to kill the cell. Thus, only

members of a population that retain the plasmid will survive. Recently, a set of Tn*3*-family

transposons carrying TA systems were characterized and included in the TnCentral database

[22]. To explore these Tn, we used the TnCentral Gene Search function, selecting "Passenger

Gene" from the Gene Class pull-down menu and "Toxin" from the Gene Sub-Class pull-down

menu (Figure 6A, red box). The search results included eight different toxin genes (Gp49,

HEPN, PIN, PIN_3, *abiEii, higB, parE*, and zeta) found in 43 different transposons. Similarly,

transposons carrying antitoxin genes were identified using the Gene Search function with

the Gene Sub-Class menu set to "Antitoxin" (Figure 6B, red box). There were 44 transposons

carrying 11 different antitoxin genes. Combinations of toxin and antitoxin genes in individual

transposons were examined by going to the ORF Summary section of the entry pages for the

TA transposons. For example, Tn*Sku1* (Figure 6B, yellow box; Figure 6C) has a Gp49 toxin

gene and an antitoxin gene containing an HTH domain (referred to as HTH). Most

transposons have a single toxin/antitoxin gene pair except for Tn*Xca1*, which has two TA

pairs, and Tn*5501.5*, which has a *parD* antitoxin gene and no toxin gene. The majority of

Tn*5501* derivatives in TnCentral have a *parE* toxin gene as well as the *parD* antitoxin,

suggesting that Tn*5501.5* may have undergone a deletion in the region containing *parE*

(Supplementary Figure 1).

Use Case #3: Tn*21* and its Relatives

Tn*21* is the canonical member of a subfamily of Tn*3* transposons that confers a variety of

antibiotic resistances [38–40] and several analyses have proposed mechanisms to explain

how *Tn21* arose from simpler ancestor transposons (e.g., [40,41]). Tn*21* has a mercury

resistance operon at the 5'- (left) end, a *tnpA/tnpR* transposition module at the 3'-(right)

end, and a transposition-deficient integron (In2) carrying several ABR genes (a GCN5-related

N-acetyltransferase (GNAT_fam), *sul1, qacEdelta1*, and *aadA*) in the middle (Supplementary

Figure 2). These ABR genes confer resistance to aminoglycosides, sulfones, sulfonamides,

quaternary ammonium salts, and acridine dye. More recently, a transposon that lacks the

integron insertion but is otherwise identical to Tn*21* (the hypothetical Tn*21* backbone Tn21Δ

in [40]) was discovered [42]. This transposon, Tn*5060*, was proposed to be the ancestor of

Tn*21* [42]. Tn*21* also has numerous relatives that carry different combinations of antibiotic

resistance genes within and outside the integron. To explore the Tn*21* subfamily, we

performed a TnCentral Sequence Search (BLAST) using the putative ancestral Tn*5060*

sequence (Figure 7A). In addition to Tn*5060* itself, we identified ten transposons in the

database (Tn*20*, Tn*21*, Tn*21.1*, Tn*21.2,* Tn*5086,* Tn*2411,* Tn*2424,* Tn*4*, Tn*1935*, and Tn*As3*;

Supplementary Figure 2) that contain all (or nearly all) of the Tn*5060* sequence. With the

exception of Tn*20*, which is almost identical to Tn*5060* (99.5%), these transposons have two

or more discontinuous sub-regions that align to Tn*5060*. For example, Tn*21* has two sub-

regions, one of which is a close match to the left half of Tn*5060* and the other of which is a

close match to the right half of Tn*5060* (red bars in Figure 7B). This suggests that these

transposons arose from Tn*5060* via the insertion of other sequences.

We compared the antibiotic resistance profiles of the ten transposons by inspecting their

TnCentral entry pages. Tn*20*, like Tn*5060*, carries no ABR genes. The other nine transposons

carry ABR genes targeting aminoglycosides, sulfones, sulfonamides, and quaternary

ammonium salts (Figure 7C). Other resistances found in a subset of the six include acridine

dye (Tn*1935*, Tn*21*, Tn*2411*, Tn*4*, Tn*As3,* Tn*2424,* Tn*5086*), carbapenams (Tn*1935* and Tn*4*),

cephalosporins (Tn*1935* and Tn*4*), carbapenems (Tn*4*), monobactams (Tn*4*), phenicols

(Tn*As3*, Tn*2424*, Tn*21.1*, Tn*21.2*), diaminopyrimidines (Tn*5086*, Tn*21.1*, *Tn21.2*), and

tetracyclines (Tn*21.2*)**.** Interestingly, in some cases where the transposons have resistances

in common, they are conferred by different genes (Figure 7C). For example, phenicol

resistance is conferred by *CAT* in Tn*As3, catB2 in* Tn*2424*, and *cmlA6* in Tn*21.1* and Tn*21.2*.

Similarly, sulfonamide and sulfone resistance is conferred by *sul1* in all of the antibiotic-

resistant family members except for Tn*21.1* and Tn*21.2*, where those resistances are

conferred by *sul3*. Thus, even this closely related subfamily of transposons shows diversity in

its antibiotic resistance genes. This is partially due to the flexibility of the integron to

incorporate new antibiotic resistance gene cassettes but also to insertion of ABR-gene

containing elements outside of the integron region (e.g., Tn*3.1* in Tn*4*, Supplementary Figure

2).

Discussion

Here, we have described TnCentral, a user-friendly resource for exploration of prokaryotic

TE. TnCentral provides a flexible search interface, TE-specific entry pages with intuitive

graphics and detailed information about TE features, and a BLAST interface that allows users

to identify TE that carry features of interest or to identify TE that are present in sequences

of interest (e.g., plasmids). As shown in the use cases, the flexible search results page makes

it easy to compare features across multiple transposons, the detailed entry pages allow

exploration of TE passenger genes, such as ABR genes, and the Sequence Search enables

retrieval of TE with related sequences that could be used as a starting point for evolutionary

analyses. Moreover, TnCentral provides access to Tnfinder software for locating candidate

TE in sequence data and to TnPedia, a comprehensive review of the biology of selected TE

families.

As discussed in the Introduction, a variety of resources dedicated to aspects of prokaryotic

TE biology currently exist. TnCentral's unique contribution to this universe of resources lies

in its coverage of a variety of TE (e.g. different transposon families and compound

transposons with their associated IS and integrons) and its detailed focus on both core

transposition genes and passenger genes of clinical, environmental, and economic

importance. It has the additional feature of providing a clear graphic output for visualizing

the often complex structures of TE.

The next step beyond annotation of individual TE is to annotate and visualize the TE content

of prokaryotic chromosomes and plasmids. These studies are critical for understanding the

propagation of high impact passenger genes, such as those that confer antibiotic resistance.

Several tools that address this problem are available. For example, ISsaga [43], which is

integrated into ISfinder, annotates IS present in user-provided sequences. Other software

suites have been designed specifically to annotate IS in short read raw data (e.g. ISQuest

[44], Transposon Insertion Finder [45], ISMapper [46] and panISa [47]) using preassembled

libraries of TE and their components, while yet other approaches are based on *ab initio*

prediction (e.g., OASIS [48], ISseeker [49] ISEscan [50], or provide a comparative view of IS

mobilisation events (e.g. ISCompare [51]). These annotation tools are only as good as their

underlying TE databases. ISfinder, which includes nearly 6000 individual examples of IS

classified in distinct families and subfamilies according to their transposition mechanism and

structural organization, provides such a rigorous framework for IS and has been

incorporated into a number of annotation pipelines (e.g., ISsaga [43], MobileElementFinder

[52]). However, IS represent only a fraction of prokaryotic TE, and unlike transposons and

integrons, they rarely carry passenger genes. We hope that TnCentral will become a

benchmark for more complex TE as ISfinder is for IS.

TnCentral is an ongoing project, and we will continue to expand and update the content. In
addition to the exporting annotated TE in GenBank format, we plan to make all files
available in a SnapGene file format which will allow users to use SnapGene
(https://www.snapgene.com/), a commercial software tool (with a free viewer version) for
visualizing and documenting nucleotide sequences and their features, to analyze and
explore them. We also intend to enhance the visualization of TnCentral Sequence Search
(i.e., BLAST) results to better support the analysis of plasmid sequences that may carry a
complex complement of TE although it should be noted that the Sequence Search tool can
already accommodate analysis of large plasmids. Ultimately, we envision that TnCentral
could be used to analyze the TE content of a collection of sequences, such as patient,
veterinary and environmental samples from an antibiotic resistance outbreak, to
understand TE-driven evolution of the prokaryotic mobilome.

Methods

Methods

Curation Workflow

The TnCentral curation workflow is depicted in Figure 8. Curation is performed by members
of the TnCentral development team as well as by graduate students in bioinformatics
courses at Georgetown University Medical Center. TnFinder scripts are run against RefSeq
and other sequence databases and GenBank files potentially containing TE are retrieved. TE
sequences are isolated and annotated using SnapGene (https://www.snapgene.com).
Features of interest (i.e., protein coding genes, TE, repeat elements, and recombination
sites) are annotated according to detailed curation guidelines (provided in the "For
Curators" of TnCentral). Fully annotated features are saved in a SnapGene Custom Library.
New transposon sequences can be searched against this library, enabling detection of
features previously identified in other TEs. All annotated TE files are checked by a second
curator. An enhanced GenBank file containing all annotations is exported from SnapGene
and checked for common curation formatting errors using a custom Perl script. Detected
errors are manually corrected in the SnapGene file, which is then exported as a revised
enhanced GenBank file. Information from this GenBank file is used to populate the
TnCentral database, which, in turn, serves as the backend for the TnCentral web portal. An

image file showing a color-coded map of TE features is also exported from SnapGene and
displayed on the TE entry page.

Although we have adhered to the standard nomenclature for transposons extracted from
the literature, for the many transposons newly identified during TnCentral database-
building, we have temporarily used names indicating their source. In all cases, the
Transposon Registry [53] accession number is provided as a synonym. There is some
ambiguity in the literature concerning class 1 integrons and members of the Tn*402*
transposon family. Class 1 integrons appear to be derivatives of this transposon family and
include members with a range of Tn*402* transposition genes with varying degrees of
completeness. We have therefore elected to include all Class 1 integrons as members of the
Tn*402* family (Supplementary Table S1). ISfinder classification is used for the individual IS
and in the case of compound transposons, the group to which they are belong is defined by
the flanking IS.

Properties of protein coding genes are annotated with cross-references to database or
ontology identifiers whenever possible. Antibiotic resistance gene properties, including gene
name, sequence family, antibiotic resistance mechanism, and target drug classes are
annotated according to the Antibiotic Resistance Ontology (ARO) as presented in
Comprehensive Antibiotic Resistance Database (CARD) [10]. The Pfam [54] and InterPro
resources [55] are used to define sequence family information.

TnCentral Website implementation
TE features and sequence information are extracted from the enhanced GenBank files. TE
feature information is used for the search and the entry pages, and the TE DNA and protein
sequence information are used for the Sequence Search and display. The extracted data is
loaded into the TnCentral database, implemented using MySQL. The website is built on a
Linux server with Apache, and the web application is built on Perl CGI. Apache Lucene is
used to index the data for flexible and fast search and retrieval. JavaScript is used for the
interactive web-interface and display. BLAST is used for similarity search.

Acknowledgements

We would like to thank John Dekker (NIAID, NIH Bethesda, Md, USA), Fred Dyda and Alison Hickman (NIDDK, NIH Bethesda, Md, USA), Patricia Siguier (CNRS Toulouse, France), Susu He (Nanjing University Medical School, Nanjing, China), Laurence van Melderen (Université Libre de Bruxelles, Belgium), and Gipsi Lima-Mendez and Bernard Hallet (Université de Louvain la Neuve, Belgium) for helpful discussions; Ben Glick (University of Chicago, SnapGene) for his help with the SnapGene software; the student curators at Georgetown University Medical Center for their contributions to the annotation process; and the Protein Information Resource (University of Delaware, Georgetown University Medical Center) for informatics support. This project was supported the U.S. Department of Defense (DoD) Global Emerging Infections Surveillance (GEIS) Branch (P0020_18_WR) and by institutional resources at the Center for Bioinformatics and Computational Biology, University of Delaware.

Figure Legends

Figure 1. Structural arrangement of prokaryotic Transposable Elements. The TE is indicated

by a pale-yellow horizontal bar at the top of each section. Open reading frames are shown

as horizontal arrows with the arrowheads indicating the direction of expression: purple,

transposition-associated genes; red, antibiotic resistance genes; green, other passenger

genes. The inverted terminal repeats found at the ends of the majority of TE are shown as

grey arrows and the direct target repeats generally produced by insertion are indicated by

small black arrows. **A.** Insertion Sequence (IS), a short DNA segment encoding only the

mobilization protein (Transposase, TnpA), flanked by two imperfect Inverted Repeats (IRs),

and generally containing a short flanking directly repeated duplication (DR) on the target of

insertion. **B.** tIS (transporter IS) are structurally similar to an IS, but contain passenger genes.

They are presently restricted to the IS*1595* and IS*66* families. **C.** Compound transposons are

formed by two IS in either direct or inverted orientation, flanking a variety of passenger

genes including those for antibiotic resistance. **D.** Transposons are more heterogeneous

structures and include different sets of transposition-related genes which are specific to

each Tn family and multiple antibiotic resistances, virulence and other passenger genes. This

is an example of a Tn*3* family transposon with transposon, *tnpA*, and resolvase genes, *tnpR*.

Figure 2. A) TnCentral homepage showing clickable links to various TnCentral sections in the

box on the left. B) TnCentral search interface showing search choices for TE on the left and

for transposition-related and passenger genes on the left.

Figure 3. TnCentral TE Entry Page. #1-10: Sections of the entry page (see text for details).

Figure 4. The main sections of TnPedia, a TnCentral-related wiki compiling information on

prokaryotic transposable elements. Only three of the four sections (General Information, IS

families and Transposon families) are illustrated. The fourth section is a Transposition

Glossary, which is under construction.

Figure 5. Comparing Protein Coding Genes in Tn*554* Family Members. A) TnCentral Transposon Search interface, showing a search for Tn*554* in the TE family field. B) Interface for customizing the columns in the search results display. Clicking on "Customize Display" (red box) opens the interface. C) Partial Tn*554* family search results after customization to show information on protein coding genes (All Gene Fields).

Figure 6. Exploring Toxin/Antitoxin Genes in TnCentral. A) Partial results of searching TnCentral for toxin genes. The settings used to obtain these results are shown in the red box. Links to entry pages for the TE carrying the indicated genes are provided in the MGE Accession column (e.g., Tn*Sku1*-CP002358.1, yellow box). B) Partial search results for antitoxin genes in TnCentral. Settings are shown in the red box. C) ORF Summary section of the entry page for Tn*Sku1*-CP002358.1, showing the presence of a toxin/antitoxin gene pair (Gp49 toxin/HTH antitoxin).

Figure 7. Analysis of ABR in Tn*21* Relatives. A) TnCentral Sequence Search using the sequence of Tn*5060*, the proposed ancestor of Tn*21*, as a query. B) Sequence Search results. The query sequence is represented by the width of the Alignment column. The red bars represent regions of the matched transposons that are highly similar to regions of Tn*5060*. C) ABR genes and targeted antibiotic classes in Tn*21* relatives. Red shading in the table cells indicates that the transposon carries at least one gene targeting the antibiotic class; blue shading indicates that it does not. The ABR genes found in each transposon are indicated in the table cells.

Figure 8. TnCentral Curation Workflow (see text for description).

Supplementary Figures

Supplementary Figure 1. Maps of Tn*5501* and Tn*5501.5* showing the loss of *parE* toxin gene in Tn*5501.5*. Feature color code: yellow--TE; purple--transposition genes; dark orange--toxin/anti-toxin genes; light orange--other open reading frames; grey--repeat elements; green--recombination sites. Maps were created with SnapGene.

Supplementary Figure 2. Maps of Tn*21* and its relatives. The feature color code is the same

as in Supplementary Figure 1. Maps were created with SnapGene. Note that the different

transposon derivatives are not to scale but their individual lengths are included.

Supplementary Table S1. The table displays the entire collection of TE at present in the

database (May 2021) with columns indicating their **TnCentral accession numbers**, their

names, **synonyms** from the literature and/or the Transposon Registry [53], TE **Type, Family**

Group), **Host Organism** and **Molecular Source** (e.g., plasmid or chromosome). If no

information is provided in the Molecular Source column, the source is chromosomal or

unknown.

Bibliography

1. Spellberg B, Guidos R, Gilbert D, Bradley J, Boucher HW, Scheld WM, Bartlett JG,
Edwards J, Infectious Diseases Society of America. The epidemic of antibiotic-resistant
infections: a call to action for the medical community from the Infectious Diseases
Society of America. Clin Infect Dis. 2008 Jan 15;46(2):155–164. PMID: 18171244

2. O'Neill J. Tackling Drug-Resistant Infections Globally:' ' Final Report and
Recommendations. London?: Welcome Foundation HM Government UK; 2016 May.

3. Craig, Chandler, Gellert, Lambowitz, Rice, Sandmeyer, editors. Mobile DNA III.
American Society of Microbiology; 2015.

4. Mahillon J, Chandler M. Insertion sequences. Microbiol Mol Biol Rev. 1998
Sep;62(3):725–774. PMCID: PMC98933

5. Siguier P, Gourbeyre E, Varani A, Ton-Hoang B, Chandler M. Everyman's guide to
bacterial insertion sequences. Microbiol Spectr. 2015 Apr;3(2):MDNA3–0030. PMID:
26104715

6. Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic
impact and diversity. FEMS Microbiol Rev. 2014 Sep;38(5):865–891. PMCID:
PMC7190074

7. Vandecraen J, Chandler M, Aertsen A, Van Houdt R. The impact of insertion sequences
on bacterial genome plasticity and adaptability. Crit Rev Microbiol. 2017
Nov;43(6):709–730. PMID: 28407717

8. He S, Chandler M, Varani AM, Hickman AB, Dekker JP, Dyda F. Mechanisms of
Evolution in High-Consequence Drug Resistance Plasmids. MBio. 2016 Dec 6;7(6).
PMCID: PMC5142620

9. He S, Hickman AB, Varani AM, Siguier P, Chandler M, Dekker JP, Dyda F. Insertion
Sequence IS26 Reorganizes Plasmids in Clinically Isolated Multidrug-Resistant Bacteria
by Replicative Transposition. MBio. 2015 Jun 9;6(3):e00762. PMCID: PMC4471558

10. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A, Huynh W,
Nguyen A-LV, Cheng AA, Liu S, Min SY, Miroshnichenko A, Tran H-K, Werfalli RE, Nasir
JA, Oloni M, Speicher DJ, Florescu A, Singh B, Faltyn M, Hernandez-Koutoucheva A,
Sharma AN, Bordeleau E, Pawlowski AC, Zubyk HL, Dooley D, Griffiths E, Maguire F,
Winsor GL, Beiko RG, Brinkman FSL, Hsiao WWL, Domselaar GV, McArthur AG. CARD
2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance
database. Nucleic Acids Res. 2020 Jan 8;48(D1):D517–D525. PMCID: PMC7145624

11.   Liu B, Pop M. ARDB--Antibiotic Resistance Genes Database. Nucleic Acids Res. 2009
553   Jan;37(Database issue):D443–7. PMCID: PMC2686595

554   12.   Xie Y, Wei Y, Shen Y, Li X, Zhou H, Tai C, Deng Z, Ou H-Y. TADB 2.0: an updated
555   database of bacterial type II toxin-antitoxin loci. Nucleic Acids Res. 2018 Jan
556   4;46(D1):D749–D753. PMCID: PMC5753263

557   13.   Akarsu H, Bordes P, Mansour M, Bigot D-J, Genevaux P, Falquet L. TASmania: A
558   bacterial Toxin-Antitoxin Systems database. PLoS Comput Biol. 2019 Apr
559   25;15(4):e1006946. PMCID: PMC6504116

560   14.   Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference
561   centre for bacterial insertion sequences. Nucleic Acids Res. 2006 Jan 1;34(Database
562   issue):D32–6. PMCID: PMC1347377

563   15.   Moura A, Soares M, Pereira C, Leitão N, Henriques I, Correia A. INTEGRALL: a database
564   and search engine for integrons, integrases and gene cassettes. Bioinformatics. 2009
565   Apr 15;25(8):1096–1098. PMID: 19228805

566   16.   Bi D, Xu Z, Harrison EM, Tai C, Wei Y, He X, Jia S, Deng Z, Rajakumar K, Ou H-Y. ICEberg:
567   a web-based resource for integrative and conjugative elements found in Bacteria.
568   Nucleic Acids Res. 2012 Jan;40(Database issue):D621–6. PMCID: PMC3244999

569   17.   Liu M, Li X, Xie Y, Bi D, Sun J, Li J, Tai C, Deng Z, Ou H-Y. ICEberg 2.0: an updated
570   database of bacterial integrative and conjugative elements. Nucleic Acids Res. 2019
571   Jan 8;47(D1):D660–D665. PMCID: PMC6323972

572   18.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller
573   Aarestrup F, Hasman H. In silico detection and typing of plasmids using PlasmidFinder
574   and plasmid multilocus sequence typing. Antimicrob Agents Chemother. 2014
575   Jul;58(7):3895–3903. PMCID: PMC4068535

576   19.   Leplae R, Lima-Mendez G, Toussaint A. ACLAME: a CLAssification of Mobile genetic
577   Elements, update 2010. Nucleic Acids Res. 2010 Jan;38(Database issue):D57–61.
578   PMCID: PMC2808911

579   20.   Leplae R, Hebrant A, Wodak SJ, Toussaint A. ACLAME: a CLAssification of Mobile
580   genetic Elements. Nucleic Acids Res. 2004 Jan 1;32(Database issue):D45–9. PMCID:
581   PMC308818

582   21.   Leplae R, Lima-Mendez G, Toussaint A. A first global analysis of plasmid encoded
583   proteins in the ACLAME database. FEMS Microbiol Rev. 2006 Nov;30(6):980–994.
584   PMID: 17064288

22. Lima-Mendez G, Oliveira Alvarenga D, Ross K, Hallet B, Van Melderen L, Varani AM,
Chandler M. Toxin-Antitoxin Gene Pairs Found in Tn3 Family Transposons Appear To
Be an Integral Part of the Transposition Module. MBio. 2020 Mar 31;11(2). PMCID:
PMC7157771

23. Szuplewska M, Czarnecki J, Bartosik D. Autonomous and non-autonomous Tn3-family
transposons and their role in the evolution of mobile genetic elements. Mob Genet
Elements. 2014 Dec;4(6):1–4. PMCID: PMC4588545

24. Loftie-Eaton W, Yano H, Burleigh S, Simmons RS, Hughes JM, Rogers LM, Hunter SS,
Settles ML, Forney LJ, Ponciano JM, Top EM. Evolutionary Paths That Expand Plasmid
Host-Range: Implications for Spread of Antibiotic Resistance. Mol Biol Evol. 2016
Apr;33(4):885–897. PMCID: PMC4840908

25. Sayers EW, Agarwala R, Bolton EE, Brister JR, Canese K, Clark K, Connor R, Fiorini N,
Funk K, Hefferon T, Holmes JB, Kim S, Kimchi A, Kitts PA, Lathrop S, Lu Z, Madden TL,
Marchler-Bauer A, Phan L, Schneider VA, Schoch CL, Pruitt KD, Ostell J. Database
resources of the National Center for Biotechnology Information. Nucleic Acids Res.
2019 Jan 8;47(D1):D23–D28. PMCID: PMC6323993

26. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990 Oct 5;215(3):403–410. PMID: 2231712

27. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
BLAST+: architecture and applications. BMC Bioinformatics. 2009 Dec 15;10:421.
PMCID: PMC2803857

28. Murphy E, Huwyler L, de Freire Bastos M do C. Transposon Tn554: complete
nucleotide sequence and isolation of transposition-defective and antibiotic-sensitive
mutants. EMBO J. 1985 Dec 1;4(12):3357–3365. PMCID: PMC554666

29. Bastos MC, Murphy E. Transposon Tn554 encodes three products required for
transposition. EMBO J. 1988 Sep;7(9):2935–2941. PMCID: PMC457088

30. Krolewski JJ, Murphy E, Novick RP, Rush MG. Site-specificity of the chromosomal
insertion of Staphylococcus aureus transposon Tn554. J Mol Biol. 1981 Oct
15;152(1):19–33. PMID: 6279864

31. Murphy E, Reinheimer E, Huwyler L. Mutational analysis of att554, the target of the
site-specific transposon Tn554. Plasmid. 1991 Jul;26(1):20–29. PMID: 1658834

32. Haroche J, Allignet J, El Solh N. Tn5406, a new staphylococcal transposon conferring
resistance to streptogramin a and related compounds including dalfopristin.
Antimicrob Agents Chemother. 2002 Aug;46(8):2337–2343. PMCID: PMC127375

33.   Kehrenberg C, Schwarz S. Florfenicol-chloramphenicol exporter gene fexA is part of
the novel transposon Tn558. Antimicrob Agents Chemother. 2005 Feb;49(2):813–815.
PMCID: PMC547298

34.   Kadlec K, Schwarz S. Identification of the novel dfrK-carrying transposon Tn559 in a
porcine methicillin-susceptible Staphylococcus aureus ST398 strain. Antimicrob Agents
Chemother. 2010 Aug;54(8):3475–3477. PMCID: PMC2916295

35.   Li D, Li X-Y, Schwarz S, Yang M, Zhang S-M, Hao W, Du X-D. Tn6674 Is a Novel
Enterococcal optrA-Carrying Multiresistance Transposon of the Tn554 Family.
Antimicrob Agents Chemother. 2019 Aug 23;63(9). PMCID: PMC6709469

36.   Schwendener S, Perreten V. New transposon Tn6133 in methicillin-resistant
Staphylococcus aureus ST398 contains vga(E), a novel streptogramin A, pleuromutilin,
and lincosamide resistance gene. Antimicrob Agents Chemother. 2011
Oct;55(10):4900–4904. PMCID: PMC3187003

37.   Hayes F, Van Melderen L. Toxins-antitoxins: diversity, evolution and function. Crit Rev
Biochem Mol Biol. 2011 Oct;46(5):386–408. PMID: 21819231

38.   De La Cruz F, Grinsted J. Genetic and Molecular Characterization of Tn21, a Multiple
Resistance Transposon from R100.1. J Bacteriol. 1982;151:222–228. PMCID: 6282806

39.   Grinsted J, de la Cruz F, Schmitt R. The Tn21 subgroup of bacterial transposable
elements. Plasmid. 1990 Nov;24(3):163–189. PMID: 1963947

40.   Liebert CA, Hall RM, Summers AO. Transposon Tn21, flagship of the floating genome.
Microbiol Mol Biol Rev. 1999 Sep;63(3):507–522. PMCID: PMC103744

41.   Tanaka M, Yamamoto T, Sawai T. Evolution of complex resistance transposons from an
ancestral mercury transposon. J Bacteriol. 1983 Mar;153(3):1432–1438. PMCID:
PMC221794

42.   Kholodii G, Mindlin S, Petrova M, Minakhina S. Tn5060 from the Siberian permafrost is
most closely related to the ancestor of Tn21 prior to integron acquisition. FEMS
Microbiol Lett. 2003 Sep 26;226(2):251–255. PMID: 14553919

43.   Varani AM, Siguier P, Gourbeyre E, Charneau V, Chandler M. ISsaga is an ensemble of
web-based methods for high throughput identification and semi-automatic annotation
of insertion sequences in prokaryotic genomes. Genome Biol. 2011 Mar 28;12(3):R30.
PMCID: PMC3129680

44.   Biswas A, Gauthier DT, Ranjan D, Zubair M. ISQuest: finding insertion sequences in
prokaryotic sequence fragment data. Bioinformatics. 2015 Nov 1;31(21):3406–3412.
PMID: 26116929

45.   Nakagome M, Solovieva E, Takahashi A, Yasue H, Hirochika H, Miyao A. Transposon
654   Insertion Finder (TIF): a novel program for detection of de novo transpositions of
655   transposable elements. BMC Bioinformatics. 2014 Mar 14;15:71. PMCID: PMC4004357

656   46.   Hawkey J, Hamidian M, Wick RR, Edwards DJ, Billman-Jacobe H, Hall RM, Holt KE.
657   ISMapper: identifying transposase insertion sites in bacterial genomes from short read
658   sequence data. BMC Genomics. 2015 Sep 3;16:667. PMCID: PMC4558774

659   47.   Treepong P, Guyeux C, Meunier A, Couchoud C, Hocquet D, Valot B. panISa: ab initio
660   detection of insertion sequences in bacterial genomes from short read sequence data.
661   Bioinformatics. 2018 Nov 15;34(22):3795–3800. PMID: 29931098

662   48.   Robinson DG, Lee M-C, Marx CJ. OASIS: an automated program for global investigation
663   of bacterial and archaeal insertion sequences. Nucleic Acids Res. 2012
664   Dec;40(22):e174. PMCID: PMC3526298

665   49.   Adams MD, Bishop B, Wright MS. Quantitative assessment of insertion sequence
666   impact on bacterial genome architecture. Microb Genom. 2016 Jul 18;2(7):e000062.
667   PMCID: PMC5343135

668   50.   Xie Z, Tang H. ISEScan: automated identification of insertion sequence elements in
669   prokaryotic genomes. Bioinformatics. 2017 Nov 1;33(21):3340–3347. PMID: 29077810

670   51.   Mogro EG, Ambrosis N, Lozano MJ. Easy identification of insertion sequence
671   mobilization events in related bacterial strains with ISCompare. BioRxiv. 2020 Oct 16;

672   52.   Johansson MHK, Bortolaia V, Tansirichaiya S, Aarestrup FM, Roberts AP, Petersen TN.
673   Detection of mobile genetic elements associated with antibiotic resistance in
674   Salmonella enterica using a newly developed web tool: MobileElementFinder. J
675   Antimicrob Chemother. 2021 Jan 1;76(1):101–109. PMCID: PMC7729385

676   53.   Tansirichaiya S, Rahman MA, Roberts AP. The Transposon Registry. Mob DNA. 2019
677   Oct 9;10:40. PMCID: PMC6785933

678   54.   El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson
679   LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE,
680   Finn RD. The Pfam protein families database in 2019. Nucleic Acids Res. 2019 Jan
681   8;47(D1):D427–D432. PMCID: PMC6324024

682   55.   Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang H-Y,
683   El-Gebali S, Fraser MI, Gough J, Haft DR, Huang H, Letunic I, Lopez R, Luciani A,
684   Madeira F, Marchler-Bauer A, Mi H, Natale DA, Necci M, Nuka G, Orengo C,
685   Pandurangan AP, Paysan-Lafosse T, Pesseat S, Potter SC, Qureshi MA, Rawlings ND,
686   Redaschi N, Richardson LJ, Rivoire C, Salazar GA, Sangrador-Vegas A, Sigrist CJA, Sillitoe

687   I, Sutton GG, Thanki N, Thomas PD, Tosatto SCE, Yong S-Y, Finn RD. InterPro in 2019:

688   improving coverage, classification and access to protein sequence annotations. Nucleic

689   Acids Res. 2019 Jan 8;47(D1):D351–D360. PMCID: PMC6323941

A

About
TnCentral Search
Sequence Search
Browse Tn List
TnFinder Software
Documentation
For Curators
TnPedia (TE Encyclopedia)
Related Links
Feedback

TnCentral

An annotated prokaryotic transposon database

GEORGETOWN UNIVERSITY
School of Medicine

PIR

unesp
UNIVERSIDADE ESTADUAL PAULISTA
"JÚLIO DE MESQUITA FILHO"

B

TnCentral Search

Transposon Search

Any field: [   ]

TE Name: [contains ▾] [   ]
TE Synonym: [contains ▾] [   ]
TE type: [All ▾]
TE family: [contains ▾] [   ]
TE group: [contains ▾] [   ]

TnCentral accession: [contains ▾] [   ]
Host organism: [contains ▾] [   ]

Country: [   ]
Date of Isolation: [   ]

[Submit] [Reset]

Gene Search

By Gene Name:
[   ]
[Submit]

By Gene Class:
[-- Select one ---------------- ▾]
[Submit]

By Gene Function:
[-- Select one ---------------- ▾]
[Submit]

1

Transposon

Name: Tn5501

Family: Tn3 **Group:** Tn3000

Evidence of Transposition: Yes

Host

Host Organism: Delftia sp. KV29

Molecular Source: plasmid pKV29

Date of Isolation: 1998

2

Map

Tn5501–JN648090.1
7928 bp

3

Terminal Inverted Repeats (IR)

IRL (Length: 38 bp) → GGGGTTCTTAAGCCGGAACCGCCGAAAATTCCGTCAGCC

IRR (Length: 38 bp) → GGGGTTCTTAAGCCAGAACCGCCGAAATTTCCGTCATCC

4

Sequence

DNA Sequence   Length 7928

```
--------10 --------20 --------30 --------40 --------50 --------60 --------70 --------80 --------90 -------100
GGGGTTCTAA GCCGGAACCG CCGAAAATTC CGTCAGCCGA TCAACGTGGC TTGTCCGCGA CCCGGTCGAT GGGGTAGACC CAACAGTCGT GTCACTAGCC 100
GCCATTTCGA TCACGGCAAT GCCAGCCGCA CGTCACGTCC AGATTGTTCC GGGTTGGCGG AGCTCTGGATG TGACGGGTGG CATACAACTG 200
CTGTGAGTCC TGCAGGGGGG CAGCTGCTG ACCGGACGGC GAGCATCGAC CCATCTCATG TATTAGTCAT GTCAGCTTTG ACACTGCGCA CGCGACGGCA 300
CCCGACCCGT TGCAGACCCC CAGACATATG GAAAGCTGAC AGCCATCTG GAGTTAGCCG GCGGACGCG GCTTCATCGC GCAGCGTCCG TGTTGATGGA 400
```

5

Recombination Sites

Name	Coordinates	Length	Sequence
res	4139-4269	131	GCCTGTCGGA AAACATTTGT TTTTCGACAG GCCTTCAACG GTCCTCTGCA CCAACCTCCG AGTGGCCGCA AAATTGTGCG GAAAACTCTG TCGCCAGACG CTACCATACG GAAACCTCGT CTTAATGGTT T

6

ORFs

ORF Summary

Gene Name	Associated TE	Coordinates	Class	Sub Class	Orientation
tnp	ISDIsp1	462-1436	Transposase		+
cupin2	Tn5501	1626-2111	Passenger Gene	Other	-
tnpA	ISGNB1-1	2150-3097	Transposase		-

ORF Details

Gene Name	Protein Name	Associated TE	Length	Coordinates	Strand
tnp	Tnp	ISDIsp1	975	462-1436	+

7

Class: Transposase

Transposase Chemistry: DDE

Protein Sequence:
```
MLTGMKQSSL ELNLSTRKTR KQELLAQMDR VVPWAALVEL IAPYYPEGKN GRPPFALEAM LRVHCMQQWF TLSDLAMEEA FFDTPIYREF AGLDAHGRMP
DESTILRFRH RLEKHRLAEQ ILATVNDLLA ARGLLLKAGT AVDATLIAAP SSTKNKDRKR DPEMHSSQKG NEWHFGMKAH IGVDADSGLV HIVGTSGNV
ADVTEGNSLL HGEETDAFGD AGYQGAHKRP DARKDVTWHV AMRPGKRKEL DKENNPVDAL IDQVEKIKAS IRAKVEHPFR VIKRQFGYTK VRYRGLKKNT
LQLKTLFALS NLWMVRHQLL GAQG
```

8

Internal Transposable Elements (TE)

TnCentral Accession	TE Name	Type	Coordinates	Length
ISDIsp1-JN648090.1	ISDIsp1	Insertion Sequence	409-1592	1184
ISGNB1-1-EF628291	ISGNB1	Insertion Sequence	2097-3143	1047

9

Internal Repeat Elements

Name	Associated Mobile Element	Coordinates	Sequence (Top Strand)
IRL	ISDIsp1	409-428	GGAAATCCTG CAAAAACCTCG
IRR	ISDIsp1	1574-1592	GCTCAACAAG TCCTGTAGG

10

References

Stolze Y, Eikmeyer F, Wibberg D, Brandis G, Karsten C, Krahn I, Schneiker-Bekel S, Viehover P, Barsch A, Keck M, Top EM, Niehaus K, Schluter A. *IncP-1beta plasmids of Comamonas sp. and Delftia sp. strains isolated from a wastewater treatment plant mediate resistance to and decolorization of the triphenylmethane dye crystal violet.* Microbiology. 2012 Aug;158(Pt 8):2060-2072. doi: 10.1099/mic.0.059220-0. Epub 2012 May 31. PubMed ID: 22653947

Main Page

Welcome to TnPedia, the TnCentral Wiki

This Tnpedia has been written in an attempt to assemble a body of information (including many of the historical articles) generally dispersed in the literature as an aid to understanding how knowledge has been built up to our present view of the key role played by transposable elements (TE) in prokaryotes – both in influencing gene expression, in driving genome evolution and in facilitating horizontal gene transfer (HGT). It is divided into four sections:

General Information on Prokaryotic Elements

A section providing general information about Transposition and **T**ransposable **E**lements (TE) with emphasis on prokaryotic elements. It was originally written for ISfinder (https://www-is.biotoul.fr/index.php⧉) with contributions from P. Siguier⧉ and E. Gourbeyre. It contains historical, mechanistic and genetic information.

Insertion Sequence (IS) Families

A second section describing each **I**nsertion **S**equence (IS) family in some detail. This section has been entirely updated compared to that which was included in a previous ISfinder⧉ version. It contains information from a number of reviews and from the primary literature together with analyses undertaken in the framework of ISfinder⧉ and TnCentral⧉. P. Siguier⧉ and E. Gourbeyre provided a large proportion of these analyses.

Transposon families [In progress]

A third section presenting detailed descriptions of transposon and transposon families written in the framework of TnCentral⧉.

A

Transposon Search

Any field: [____________]

TE Name: [contains ▾] [____________]
TE Synonym: [contains ▾] [____________]
TE type: [All ▾]
TE family: [contains ▾] [Tn554]
TE group: [contains ▾] [____________]

TnCentral accession: [contains ▾] [____________]
Host organism: [contains ▾] [____________]

Country: [____________]
Date of Isolation: [____________]

[Submit] [Reset]

B

Transposon Search

Transposon Search Output Page

[search ▶] [MGE Family ▾] [AND ▾] [Any Field ▾] [+ add input box]
[Tn554] [____________] [− del input box]

Page Size [10 items/page ▾]

Fields Not in Display
Country
Date of Isolation
Evidence of Transposition
First Isolate

Fields In Display
All Gene Fields
MGE Family
MGE Name

[apply ✓]
[cancel ✕]

14 entries | 2 pages | 10 / page | [1] 2

[↓ Select]

 Customize Display   Save Result As: ⟲TABLE | ⟲FASTA

	TnCentral Accession	MGE Name	MGE Synonyms	MGE Type	MGE Family	MGE Group	Host Organism	Country	Date of Isolation
☐	Tn5406-AF186237	Tn5406		transposon	Tn554		Staphylococcus aureus MRSA	Spain	January 2011
☐	Tn554-X03216.1.1	Tn554		transposon	Tn554		Staphylococcus aureus	USA	1979
☐	Tn554Cad-AB037671.1	Tn554Cad		transposon	Tn554		Staphylococcus aureus 85_2082	New Zealand	2001
☐	Tn558-AJ715531.1	Tn558		transposon	Tn554		Staphylococcus lentus	USA	2004
☐	Tn558.1-AM408573.1	Tn558.1		transposon	Tn554		Staphylococcus warneri	Denmark	2006
☐	Tn558.2-MH018573	Tn558.2		transposon	Tn554		Enterococcus Enterococcus avium strain C674	China	2018
☐	Tn558.3-CP025122	Tn558.3		transposon	Tn554		Bacillus sp. HBCD-sjtu	China	2017
☐	Tn559-FN677369	Tn559		transposon	Tn554		Staphylococcus aureus ST398	Germany	2010
☐	Tn6133-FR772051.1	Tn6133		transposon	Tn554		Staphylococcus aureus subsp. aureus ST398	Switzerland	2011
☐	Tn6188-HF565366	Tn6188		transposon	Tn554		Listeria monocytogenes 6179	Austria	2013

C

	TnCentral Accession	MGE Name	MGE Family	Gene Name	Gene Class	Gene Function	ORF Target	Gene Length (bp)	Protein Length (aa)	Protein Molecular Function	Sequence Family
☐	Tn558.3-CP025122	Tn558.3	Tn554	tnpA	Transposase			1086	361		
				tnpB	Transposase			1920	639		
				tnpC	Accessory Gene	Helper		366	121		Tn554_family
				fla	Passenger Gene	Other		417	138		flavodoxin
				fexA (ARO:3002704)	Passenger Gene	Antibiotic Resistance	phenicol antibiotic (ARO:3000387)	1428	475	antibiotic efflux (ARO:0010000)	major facilitator superfamily (MFS) antibiotic efflux pump (ARO:0010002)
☐	Tn559-FN677369	Tn559	Tn554	tnpA	Transposase			1086	361		
				tnpB	Transposase			1893	630		
				tnpC	Accessory Gene	Helper		378	125		Tn554_family
				dfrK (ARO:3002869)	Passenger Gene	Antibiotic Resistance	diaminopyrimidine antibiotic (ARO:3000171)	492	163	antibiotic target replacement (ARO:0001002)	trimethoprim resistant dihydrofolate reductase dfr (ARO:3001218)

Gene Search

Gene Search Output Page

Gene Class: Passenger Gene **Gene Sub Class:** Toxin **Gene Target:** -- Select one -----------------

8 items | 1 page | 10 / page |

↓ Select — Customize Display — Save As ▾TABLE

Gene Name	Gene Class	Gene Function	ORF Target	MGE Accession	MGE Name	Host Organism	Country
☐ Gp49	Passenger Gene	Toxin	ribosome associated mRNA	TnpPGH1-Y09450.1	TnpPGH1	Pseudomonas putida	
				TnSku1-CP002358.1	TnSku1	Sulfuricurvum kujiense DSM 16994	Japan
				Tn4662a.1-AY831462.1	Tn4662a.1	Pseudomonas putida GJ31	
				Tn4662a-NC_014124.1	Tn4662a	Pseudomonas putida HS	U.S.A
				Tn5501.12-CP017294.1	Tn5501.12	Pseudomonas aeruginosa PA83	Germany
☐ HEPN	Passenger Gene	Toxin	RNA	TnSod9-NC_004349	TnSod9	Shewanella oneidensis MR-1	
☐ PIN	Passenger Gene	Toxin	single stranded RNA	TnXca1-NC_007507	TnXca1	Xanthomonas campestris pv. vesicatoria	
				TnPsy42-KX009060.1	TnPsy42	Pseudomonas syringae pv. actinidiae RT594	Japan
				TnXax1.1-NC_016053	TnXax1.1	Xanthomonas arboricola pv. pruni CFBP 55306	

Gene Class: Passenger Gene **Gene Sub Class:** Antitoxin **Gene Target:** -- Final --

11 items | 2 pages | 10 / page | 1 | 2

↓ Select — Customize Display — Save As ▾TABLE

Gene Name	Gene Class	Gene Function	ORF Target	MGE Accession	MGE Name	Host Organism	Country
☐ HTH	Passenger Gene	Antitoxin		TnSku1-CP002358.1	TnSku1	Sulfuricurvum kujiense DSM 16994	Japan
☐ HTH_37	Passenger Gene	Antitoxin		Tn4662a.1-AY831462.1	Tn4662a.1	Pseudomonas putida GJ31	
				Tn4662a-NC_014124.1	Tn4662a	Pseudomonas putida HS	U.S.A
				Tn5501.12-CP017294.1	Tn5501.12	Pseudomonas aeruginosa PA83	Germany
				TnpPGH1-Y09450.1	TnpPGH1	Pseudomonas putida	
☐ PIN_12	Passenger Gene	Antitoxin		TnBth3-CP003766	TnBth3	Bacillus thuringiensis HD-789	

ORF Summary

Gene Name	Associated TE	Coordinates	Class	Sub Class	Orientation
HTH	TnSku1	126-416	Passenger Gene	Antitoxin	+
Gp49	TnSku1	419-712	Passenger Gene	Toxin	+
tnpR	TnSku1	713-1336	Accessory Gene	Resolvase	-
tnpA	TnSku1	1492-4410	Transposase		+

Sequence Search (BLAST)

A

Job Name: [   ]

Database

TnCentral

Other Sequence Search Services:

iSfinder: ○   CARD: ○   Toxin/Antitoxin: ○   NCBI: ○

Query Sequence

Enter a sequence here:

```
>Tn5060
GGGGGCACCTCAGAAAACGGAAAATAAAGCACGCTAAGGCATAGCTGACCTTGCCAGGCCTGCTTCGCCCTGTAGTGACGCGATCAACGGGCAG
GAAACATTCCCCTTTCGTGCATGGCAGGCGCACACGAGTTCAGACAGCACGGTTTCCATGCGCGCCAAGTCGGCCATCTTCTCGCGCACGTCCT
TGAGCTTGTGTTCGGCCAGGCTGCTGGCCTCCTCGCAGTGGGTGCCATCGTCGAGCCGCAACAGCTCGGCAATCTCGTCCAGACTGAACCCCAG
CCGCTGTGCCGATTTCACGAATTTCACCCGAACCACGTCCCGCTCTCCCATAGCGGCGATGCTGCCGTAAGGCTTGTCCGGTTCCGGCAACAGG
CCCTTGCGCTGATAGAAGCGGATTGTCTCCACGCTGACCCCGGCCGCCTTGGCAAAAACGCCAATGGTCAGGTTTTCCAAATTATTTTCCATAT
CGCTTGACTCCGTACATGAGTACGGAAGTAAGGTTACGCTATCCAATCCAAATTCAAAAGGGCCAACGTATGTCTGAACCACAAAACGGGCGCG
GTGCGCTCTTCGCCGGCGGGCTGGCCGCCATTCTTGCATCGACCTGCTGCCTGGGGCCGCTAGTACTGGTCGCCCTGGGCTTCTCCGGTGCTTG
GATCGGCAACCTGACGGTGCTGGAACCCTATCGACCGTTGTTCATCGGCGCGGCGCTAGTGGCGCTGTTCTTCGCCTGGAAGCGGATTTACCGG
CCCGTGCAGGCATGCAAGCCAGGTGAGGTCTGCGCGATTCCGCAGGTGCGCGCCACCTACAAGCTGATTTTCTGGATCGTGGCCGTGCTGGTCC
```

Or, upload file (Fasta format) : [Choose File] No file chosen

B

TnCentral Accession ⇕	TE Name ⇕	TE Length ⇕	Host Organism ⇕	TE Family ⇕	Alignment (by score) ⇕
Tn5060-AJ551280.1	Tn5060	8667	Pseudomonas sp. A19-1	Tn3	
☐ Tn20-AF457211.1	Tn20	8644	Escherichia coli	Tn3	
☐ Tn21-AF071413	Tn21	19672	Shigella flexneri	Tn3	
☐ TnAs3-CP000645.1	TnAs3	18735		Tn3	
☐ Tn21.2-MH626558	Tn21.2	35400	Salmonella enterica subsp. enterica serovar Typhimurium	Tn3	
☐ Tn2424-UGCJ01000005	Tn2424	26008	Escherichia coli NCTC11186	Tn3	
☐ Tn5086-CP054343	Tn5086	15341	Escherichia coli SCU-164	Tn3	
☐ Tn1935-MK797990	Tn1935	23364	Salmonella enterica subsp. enterica serovar Wien ZM3	Tn3	
☐ Tn21.1-MH257753	Tn21.1	21668	Salmonella enterica subsp. enterica serovar Typhimurium	Tn3	
☐ Tn2411-FN554766	Tn2411	18055	Escherichia coli 042	Tn3	
☐ Tn4-KY749247.1	Tn4	23009	Salmonella enterica subsp. enterica serovar Paratyphi B	Tn3	

C

	Tn1935	Tn21	Tn2411	Tn4	TnAs3	Tn2424	Tn5086	Tn21.1	Tn21.2
aminoglycosides	GNAT_fam, aadA, aph3'-Ia	GNAT_fam, aadA				GNAT_fam, AAC(6')-Ia, aadA3	GNAT_fam	aadA, aadA2	
sulfonamides	sul1							sul3	
sulfones									
quaternary ammonium salts	qacEdelta1							qacL	
acridine dye									
phenicols					CAT	catB2		cmlA6	
diaminopyrimidines							dfrA7	dfrA12	
tetracyclines								tetR, tet(B), tetC_p	
penams	bla-OXA-1								
cephalosporins				bla-TEM-1					
penems									
monobactams									

TnFinder Scripts, Literature

Transposon Feature Map (Image file)

GenBank File

SnapGene® 5.2.4

New DNA File...

New Protein File...

Open

Open Recent File

Import

TnCentral Database

TnCentral
An annotated prokaryotic transposon database

TnCentral Website

Enhanced GenBank File

Curation Guidelines

QC: Validation by Second Curator Automated Format Checking (Perl Script)