Single-cell phenotypic characteristics of phenotypic resistance under recurring antibiotic exposure in

Escherichia coli.

Silvia Kollerová[1], Lionel Jouvet[1], Julia Smelková [1], Sara Zunk-Parras[2], Alexandro Rodríguez-Rojas[2],

Ulrich K. Steiner[1,2*]

[1]Department of Biology, University of Southern Denmark, Odense, Denmark

[2]Biological Institute, Freie Universität Berlin, Berlin, Germany

[*]Corresponding author

Email: ulrich.steiner@fu-berlin.de

Keywords: single-cell, microfluidics, antibiotic resistance, β-lactam antibiotics, growth, survival,

persistence, tolerance

Short title: Phenotypic characteristics of phenotypic antibiotic resistant cells

Abstract:

Despite increasing interest, non-heritable, phenotypic drug resistance, such as tolerance and

persistence towards antibiotics, remains less characterized compared to genetic resistance. Non-

heritable drug resistance challenges antibiotic treatment and have implications towards heritable

resistance evolution. Phenotypically resistant cells have commonly been characterized as growth

arrested cells prior and during antibiotic application that quickly resume growth post-application.

Here, we add novel combinations of characteristics of phenotypic resistant *E. coli* single cells—that

are of particular interest towards genetically fixed resistance—, and contrast those to susceptible

cells of the isoclonal initial population by exposure to different levels of recurrent antibiotic. We find

that phenotypic resistant cells reduced their growth rate by about 50% compared to growth rates

prior to antibiotic exposure, but cells do not go into near growth arrest. The growth reduction is

induced by antibiotic exposure and not caused by a stochastic switch or predetermined state as

frequently described. Cells exhibiting constant intermediate growth survived best under antibiotic

exposure and, contrary to expectations, selection did not primarily act on fast growing cells. Our

findings support diverse modes of phenotypic resistance, and we revealed resistant cell

characteristics that supports acclaims of an underappreciated role of phenotypic resistant cells

towards resistance evolution.

Introduction

Resistance to antibiotics, in combination with slow progress in discovering novel antibiotics, poses a major challenge to modern medicine (1). Antibiotic resistant cells sustain bactericidal concentrations of antibiotics through heritable, often well-known and explored, genetically fixed mutations. In contrast, non-heritable phenotypic resistant cells, also often referred to as antibiotic persistent or tolerant cells are in a non-heritable drug resistant phenotypic state. This state allows them to survive periods of bactericidal concentration of antibiotics and is classically characterized by non- or very slow growth prior and during exposure to the drug (2–5). More recently 'specialized' persister cells have been described that show less growth dependent persistence (6–12). Identifying persister cells is usually done through a characteristic biphasic killing curve that is observed at the population level and caused by differential selection of persister cells of a susceptible population during antibiotic exposure (2,13), in contrast tolerant cells do not show a biphasic killing curve as they are assumed to be a population wide response and not a subpopulation one (2–5,9). However, definitions on persistence and tolerance are not consistent as persistence is also referred to as a special case of tolerance, or is defined as hetero-tolerance (4). The persister cell stage can be environmentally induced or stochastically switched to, hence, cells can dynamically alter states, and progeny of persiter cells are as susceptible as the parental population from which they derived (4,14). Quantitatively, below 0.1% of cells in wild-type susceptible populations are persister cells, though variation in persister fractions exists among experiments, strains, growth media, and environmental conditions (15). Uncertainty in accurately quantifying persister cell fractions based on killing curves comes from highly sensitive determination of slopes and intercepts of killing curves (10).

Tolerance is a population level response and leads to a slowed killing curve, tolerant cells are therefore much more common than persister cells (2,9,16). Tolerant cells tolerate antibiotic exposures by mainly non- or very slow growing under antibiotic exposure, characteristics they share with persister cells (2). Though compared to persister cells, tolerant cells can only sustain shorter

durations of antibiotic exposure (4). Under antibiotic application, tolerant cell populations show

prolonged minimum duration of killing (MDK) without a change in the minimum inhibitory

concentration (MIC), that is, they die slowly under antibiotic application even when exposed to very

high concentrations.

A third class of cells that can sustain bactericidal concentration of antibiotics by phenotypic

resistance are viable but non-culturable cells (VBNC/ deep dormant cells / sleeper cells) (Dong et al

2020). In contrast to persister and tolerant cells, VBNC cells do not regain growth 3-4 hours post-

antibiotic exposure, but instead remain at low metabolic activity in a deeper dormancy state and do

not easily resuscitate, potentially due to a lack of intracellular ATP (17). Persister and tolerant cells

are therefore more susceptible to recurring exposure to antibiotics compared to VBNC cells (17).

VBNC cells are further phenotypically characterized by being small (2.3 ± 0.2μm for E. *coli*) and less

elongated compared to persister cells (18).

Differentiating among these different phenotypic resistant cell types can be difficult, in particular

when single-cell data is not available (3,4,7,19). A growing number of studies on persistence work at

the single-cell level and primarily aim at a mechanistic understanding of phenotypic resistance

(8,18,20–24), yet in depth analyses of single-cell phenotypic characteristics of phenotypic resistant

cells are still rare (8,10,25). Many of these single-cell studies use persister mutants, and start with

growth-arrested, stationary phase cells, as these conditions increase the fraction of persister cells.

Some recent studies suggest that persister cells arising of exponential growing populations are

predominantly not growth arrested prior to antibiotic exposure (10). Despite evidence for diverse

modes of phenotypic resistance, overall, these single-cell level studies highlight that that persister

cells show limited growth when exposed to β-lactam or other antibiotics that inhibit cell wall

formation or cell division (6,7,26), and that switching to persister stages can be irreversible or a

reversible dynamic stochastic process (6,10). Earlier studies reported that the stochastic switch to

the rare persister state occurred for growth-limited cells within exponentially growing *E. coli*

populations, with ~80% of persisters arising from the non-growing sub-population prior to

treatment, while the remaining persister cells originated from cells that were growing prior to

treatment (11,27). Other high throughput experiments, not working on persister mutant strains,

suggest that large fractions of persister cells arise from non-dormant cells when selected of an

exponentially growing population (10). Quantitative high-quality single-cell data on phenotypic

characteristics of phenotypic resistant cells are limited, as persister cells are so rare and selecting for

them remains challenging (10). However, characterizing single-cell growth, division and mortality of

phenotypically resistant cells would provide deeper understanding about the evolutionary

importance of non-heritable resistant cells for the evolution of heritable (genetic) resistance and

efficacy of antibiotic treatment, as high frequencies of non-heritable tolerant cells provide

evolutionary reservoirs and increase the probability of genetically fixed resistance mutations to

occur (28–37).

Here, we use a demographic and phenomenological approach to quantify the fraction of cells in a

phenotypic resistant stage among susceptible stage cells, we investigate how phenotypic resistant

cells differ in their single-cell characteristics compared to susceptible cells, and we study what

triggers changes in their phenotypes. To increase the number of cells that sustained antibiotic

exposure, we reduce exposure periods to recurring 1.5 h periods compared to a typical single 3-5h

exposure period. Whether the resulting selected cells should be referred to as hetero-tolerant cells,

or persiter cells depends on definitions, we follow with the term persister. We started with a

susceptible clonal exponentially growing population of cells that we expose to recurrent antibiotic

treatment at sub- to supra level antibiotic concentrations, we track each cell for their growth,

division, and survival in a microfluidic device called a mother machine (38–40). By exposing cells to a

β-lactam, ampicillin, that corrupts cell wall synthesis, we expected quickly-growing cells to be heavily

selected against, leaving primarily slow-growing phenotypic resistant cells that also arose from the

slowest-growing population of cells prior to antibiotic exposure (3). Phenotypic resistant cells,

compared to genetically resistant cells, should not be sensitive to antibiotic concentrations beyond

the MIC of the susceptible cell population (3), and are expected to regain normal growth rates within

3-4 hours post-antibiotic treatment, as long as they are not VBNC cells. We expected few cells to

stochastically switch to tolerant resistance stages or switch by environmental induction, as we

expected phenotypic resistant stages to be mainly predetermined prior to antibiotic exposure. To

evaluate how our findings at the single-cell level might scale to cell culture dynamics, we exposed

cell cultures to similar recurring exposures of antibiotics and counted colony forming units (CFU)

before and after each exposure to antibiotics. In addition, we performed an experiment by

measuring the optical density dynamics of constantly exposed cell cultures to the same levels of

antibiotics. We discuss why scaling and quantitative comparison between single-cell characteristics

and experiments, and cell culture level exploration can be challenging.

MATERIAL METHODS

We conducted four set of experiments, three of them with three recurring antibiotic (ampicillin)

exposure periods of 90 minutes; exposure periods started 60 min, 330 min and 1080 min past onset

of the experiments. Two of these sets collected single-cell data using a mother machine microfluidic

device and time-lapse microscopy (Movie SI) (38,39,41) and one set collected cell culture level data,

i.e., CFU counts (details SI). Cells were derived of an exponentially growing susceptible isoclonal *E.*

coli strain K12-MG1655, ATCC® 700926TM, population. Exposure was at one of eight different

concentrations ranging from sub-MIC to supra-MIC levels (0-128μg/ml; details SI). Cell death in the

microfluidic device was determined using propidium iodide. The fourth experiment, which was at

the cell culture level, continuously exposed cells to antibiotics and served as a control experiment.

All experiments, single-celled and cell culture experiments, were done in supplemented minimal

media (M9), and all cultures were derived from the same *E. coli* strain, that have a replication time

of ~25-30 minutes in M9 medium at 37°C (details SI). We chose this strain as it is well adapted and

explored and has been used in previous work on persister cells (11,18,26,42). More details on

bacterial cell culturing, microchip casting and mounting, microchip loading, life cell imaging, image

analysis, experimental treatment are provided in the SI. We remain brief in our description here as

most methods follow previously published protocols (39,43).

The resulting data was subsequently analysed in program R (44) using general linear, generalized

linear, and nonlinear (Generalized Additive Models, GAM) models. Models were compared based on

information criteria (AIC)(45). We considered an ΔAIC >2 as better support between competing

models. Survival analyses (Kaplan-Meier Fig. 1a) are computed with R package survival (survfit

function) and compared amongst them with a Cox proportional hazard model. Probability of death

curves (Fig. 1b, Fig. 3b), division rate curves (Fig. 2a), and size curves (Fig. 2b) were plotted with

ggplot2 package and a loess smoothing (geometric_smooth function; package ggplot2), growth

curves were fitted with loess smoothed GAM models (Fig. 2c), competing GAM models (Table 1) are

fitted with the restricted maximum likelihood method (REML). Details on the statistical testing,

exclusion of extreme values, and model specifics are given in the SI, which also include details on

comparisons of growth differences before death (GLM's; post-hoc Tukey test using glht function of

mutlcomp package), and testing for the robustness of growth rates (Fig. 4; GAM).

For the cell culture experiments, to collect the number of colony forming unit (CFU) data and the cell

culture density data (OD_{600}), we initially seeded these experiments using cells from an exponential

growing culture. Each experiment had 8 replicates, the experiments were done in 96 well-plates, and

cells grew at 37°C and vigorous shaking in their respective well (Supplementary File 2). Recurrent

exposure to different antibiotic concentrations was achieved through serial dilution, centrifuging,

removing super natant, resuspending and culturing cells, followed by serial dilution and plating on

agar plates and CFU were counted after 16h of incubation, more details are provided in the SI.

RESULTS

Mortality and selection for tolerance

As expected, the probability of cell death, i.e., the force of mortality, increased after the first

exposure of the susceptible parental population to the antibiotic, when exposed to MIC or higher

concentrations (Fig. 1b, Movie SI). Selection for phenotypic resistant cells was strong with about two

thirds to three quarter of cells dying before the second exposure period, when exposed to MIC or

higher concentrations (Fig. 1a, Movie SI). The survivorship patterns (Fig. 1a) somewhat resemble a

biphasic killing curve as frequently observed at the cell culture level of persister cells, though we

highlight that direct comparison among cell culture level killing curves and single-cell survival can

only be made in a crude way. Mortality peaked 30 to 60 minutes after the first 90-minute antibiotic

exposure period, which suggests that the force of selection peaked with a lag time of 2 to 2.5 hours

after the start of the first exposure. Survival patterns were similar at, or above, the MIC of 4 µg/ml

(Fig. 1a, Cox proportional hazard model Likelihood ratio test 477.1, $p<0.0001$, n= 3496; for post hoc

testing see SI Table S1). The subsequent 90-minute exposure period triggered a less pronounced

mortality force compared to the first exposure period with a similar lag time of about 30 to 90

minutes after the second antibiotic exposure. Note, cell numbers became small (~20-30 cells per

concentration) and estimates of mortality rates less reliable towards the third exposure period. For

the control, 0 µg/ml, cells died, as expected, at a low rate and in agreement with experiments aiming

at senescence patterns of aging single cells (40).

Fig. 1: Survivorship curves (a) and probability of death, i.e., force of mortality (b) of cells exposed to

different levels of antibiotics across the duration of the experiment. The grey curves are control group

cells that were not exposed to any antibiotics. The grey vertical bars mark the exposure period to the

antibiotic. Kaplan-Meier survivorship curves (a) are shown with 95% CI. Probability of death curves

(b) are loess smoothed with 95% CI. Note, the two panels (a & b) are built from the same single-cell

mortality data.

Division rate, size, and growth responses

In contrast to the control group of no antibiotic exposure, cells that had been exposed to MIC or

higher antibiotic concentrations reduced their division rate after the first exposure period, changing

from ~2 to ~1 division per hour (Fig. 2a, Table 1). Most of this reduction in reproductive rate was

reached ~150-200 minutes after the onset of the first exposure (300-360 minutes after onset of

experiment). The division rates tended to only slowly decline after the second exposure, patterns

that might also arise due to senescence and not only antibiotic exposure (41).

Fig. 2: Cell division rates (a), cell size (in pixels with 100 pixels~2.4μm) (b), and growth rates (c) of

cells exposed to different levels of antibiotics across the duration of the experiment. The grey vertical

bars mark the exposure period to the antibiotics. Division rate (a) size (b) and growth rate (c) curves

are with 95% CI. Note, size data (b) for the control group (0μg/ml) was not available.

Table 1: Model comparison among competing GAM models.

		DF	AIC
Division rate	**S(Age) + Concentr.**	**15.8**	**455232**
	S(Age)	8.4	487667
	Concentr.	8.0	455263
Size	**S(Age) + Concentr.**	**16.8**	**2455427**
	S(Age)	10.8	2460359
	Concentr.	8.0	2458502
Growth rate	**S(Age) + Concentr.**	**17.7**	**-1091065**
	S(Age)	10.5	-1073598
	Concentr.	9.0	-1087172

For each of the three response variables (Division rates, Size, and Growth rates), we compared three competing models that had two explanatory variables (Age, i.e. time since onset of experiment, and antibiotic concentration). The best supported of these three models is highlighted in bold.

Average cell size remained approximately constant within a given antibiotic concentration. Although

we found a surprising amount of variation in size among cells exposed to different levels of

antibiotics (Fig. 2b, Table 1), there was no relationship between size and the levels of antibiotics. Cell

growth rates responded as predicted by declining during and after the first antibiotic exposure for

cells exposed to the MIC or above. We highlight here that growth rates only approximately halved

(Fig. 2c, Table 1), which contrasts our predictions based on previous findings where persister cells

were growth arrested or near growth arrested (4). The induced growth reduction started about 30

minutes after the onset of antibiotic exposure, which is a lag time that corresponds to approximately

the doubling rate (division time) of the parental population. The lowered growth rates stabilized

about 4-5 hours after onset of the initial exposure period, suggesting substantial variance in lag time

among individual cells' growth response. Induced growth reductions depended partly on antibiotic

concentrations with higher levels of antibiotics leading to lower growth rates. However, even with

the highest antibiotic concentration, the growth rates did not drop below 1.02-1.03/4 minutes, i.e.,

2-3% cell elongation within 4 minutes. Further, growth did not resume to prior exposure levels rates

within 3-4 hours after release of exposure.

Growth specific selection

The growth rates as discussed above (Fig. 2c) are average growth rates at a given point in time, and

even though variance in growth rates (see CI Fig. 2c) does not increase substantially during the

experiment (beyond the expected increase due to decreasing number of cells towards the end of the

experiment), selection might still act preferentially against cells with high growth as expected from a

β-lactam antibiotic and as previously described for phenotypic resistant cells (14). To evaluate the

possibility of such differential selection based on cell growth we analysed together all cells exposed

to MIC or above, and then separated them out into three growth categories: cells that did not grow

or shrank (growth arrested cells with growth rates <1.0), cells that grew at low to intermediate rates

(growth rates 1.0-1.1), and cells that grew quickly (growth rates >1.1) (Fig. 3a). Note that cells can

switch dynamically among these categories throughout the experiment and that mortality and

survival is associated only with the current growth rate of a cell and not a cell's growth rate averaged

across time. As expected, quickly growing cells showed increased probabilities of death upon

exposure to antibiotics, but contrasting to our expectations, the probability of death of growth

arrested cells increased substantially upon exposure to antibiotics, whereas the probability of death

of slow-intermediate growing cells did not change (Fig. 3b, see also SI Table S2 and Supplementary

file 1C).

Fig. 3: Cell growth rate distribution divided into three categories: growth arrested (shrinking or non-growing; green), intermediate growth (blue), and fast growth (red) (a), and probability of death for these three groups of growth arrested, intermediate, and quickly-growing cells across the duration of the experiment (b). Confidence intervals (95% CI) are shown in shading in panel (b). The grey vertical bars in panel (b) mark the exposure period to the antibiotics. Note that cells can switch among growth categories throughout the experiment, that is cells are categorized according to their current cell growth rate. For this analysis, we have combined cells across levels of antibiotic exposure ≥4 µg/ml and excluded all cells exposed to 2 µg/ml. A statistical model comparison showed that mortality differed substantially among the three growth category groups: ΔAIC 25.8 comparing a growth category model and a null model, see also Supplemental Information. Classical p-value post-hoc analyses showed that quickly growing and growth arrested groups did not differ from each other in age at death, but the intermediate group did.

To verify the robustness of our finding that selective mortality is dependent on growth rates, and to broaden our understanding of how growth influences mortality beyond the current growth rate (Fig. 3b), we did three comparisons between average growth rates of cells dying during specific periods in time. First, we compared the average growth rate (averaged across time) of cells that survived the first exposure period to those that died before the first exposure period. Second, we compared the average growth rate of cells dying during or shortly after the first exposure period (cut off at peak of

first mortality) to those surviving past the first peak of mortality, and third we compared similarly for

cells that died between the first peak of mortality and the second peak of mortality to those that

survived past the second peak of mortality (Table 2). Before the first exposure, cells that died had

lower average growth rates compared to those that survived past the first exposure, showing that

without selection of antibiotics slower growing cells are preferentially selected against. Cells that

died during the first exposure or shortly after, i.e., during times of strong selection, had slightly

higher growth rates than those that survived past this first increased selection period. This pattern of

growth dependent selection remained for the next period of increased selection after the second

antibiotic exposure. These findings support our initial prediction that faster growing cells would be

preferentially selected against under antibiotic exposure. However, note that the effect sizes

(growth differences) among surviving and dying cells are small, even though they are significantly

different. Combined with the findings shown in Fig. 3, we interpret our findings on growth specific

selection under antibiotic exposure as not primarily acting on fast-growing cells.

Table 2: Differences in average growth rates of cells surviving past or dying during or after specific

points in time.

PERIOD	0 – 60 MIN	60 – 328 MIN	328 – 600 MIN	+600 MIN
GOWTH RATE OF CELLS DYING DURING PERIOD (±STDEV)	1.047 (±0.039)	1.067 (±0.032)	1.057 (±0.030)	1.045 (±0.027)
GROWTH RATE OF CELLS SURVIVING PERIOD (±STDEV)	1.072 (0.0412)	1.060 (±0.035)	1.050 (±0.028)	
AIC GROWTH DURING/AFTER	**-8020.8**	**-8816.3**	**-3170.8**	
AIC NULL MODEL	-7985.4	-8772.0	-3161.4	
ΔAIC	35.4	44.3	9.4	

Model comparison among a null model (intercept only model) and a model with the two groups of

cells dying during a particular period or surviving that period as explanatory variable and the mean

growth rate (per 4 min) as response variable. Three sets of models were fit, one for each period [0

min – 60 min; 60 min – 328 min; 328 min – 600 min]. A Gaussian error structure was used to fit these

models.

Robustness of growth rates within cells

In Fig. 2c, we investigated the average growth rate at a given time point. As cells can dynamically

alter growth rates between the current and a future point in time, we correlated a cell's current

growth rate to its growth rate in the future, which provides a measure of robustness of growth.

Overall correlation between an individual's cell growth rate at a given time t and its growth rate in

the immediate future (time t+1 [4 minutes later]) or future time points (time t+3 [12 min later]; time

t+5 [20 minutes later]) was weak (Fig.4, Table 3), that is, growth rates are little robust and a cell's

current growth rate does not well predict its future growth rate. However, there was the exception

for cells with intermediate growth rates (1.0-1.1). Cells growing in that range (Fig. 4, x-axis range 1.0-

1.1) showed a strong correlation between current and future growth, illustrating robust constant

growth for these cells. Growth arrested cells (growth rate <1 at time t) tended to increase in their

growth rates at future points in time, while quickly-growing cells (growth rates >1.1 at time t) tended

to decrease their growth rates (Fig. 4). Overall, we observed substantial temporal dynamics in

growth rates with less robust growing cells (growth arrested and quickly-growing cells) being more

affected by the concentration of antibiotics, which suggests that the growth rate responses to

different concentrations of antibiotics shown in Fig. 2c are mainly influenced by the fastest growing

and growth arrested cells.

Fig. 4: Correlation in cell growth rates at a given time t, compared to growth rates at time t+1 (4 minutes later) (a), t+3 (12 minutes later) (b), and t+5 (20 minutes later) separated out into populations recurrently exposed to different levels of antibiotics (as described in the main experiment). All growth rate correlations are fitted with a GAM with 95% CI. The dashed black line illustrates the direct correlation among time points.

Table 3: Model comparison among competing GAM models, for correlation of current growth rates to growth rates at future points in time.

	MODEL	DF	AIC
GROWTH T+1	**Concentr. + S(Growth by Concentr)**	**70.3**	**-441151**
	Concentr. + S(Growth)	17.0	-440468
	S(Growth)	11.0	-438626
	Concentr.	8.0	-425891
GROWTH T+3	**Concentr. + S(Growth by Concentr)**	**69.2**	**-421883**
	Concentr. + S(Growth)	17.0	-421503
	S(Growth)	11.0	-420235
	Concentr.	8.0	-407188
GROWTH T+5	**Concentr. + S(Growth by Concentr)**	**69.0**	**-412563**

	Concentr. + S(Growth)	17.0	-412180
	S(Growth)	11.0	-410925
	Concentr.	8.0	-398709

Best supported model for each response variable (Growth t+1, t+3, t+5 respectively) is highlighted in

bold. GAM models are fitted with the restricted maximum likelihood method, a smoothing parameter

for Growth rate at time t with a shrinkage version of a cubic regression spline as smoothing

parameter (bs=cs) and/or the concentration, including a model that accounts for the interaction of

the two explanatory variables, growth rate and concentration of antibiotics. Growth rates are fitted

with a Gaussian error structure and an inverse link function.

Population level responses

Much research investigating phenotypic resistance has been done in cell cultures, i.e. at the

population level, which makes it challenging to partition mortality, reproduction and growth

responses, characteristics that need single-cell level evaluation. Further, density dependence and

resource depletion need consideration at the cell culture level but are controlled for by the

microfluidics setup at the single-cell level. Scaling and comparing single-cell characteristics to

population level patterns can therefore be challenging. In the following section we report on

'control' experiments done in cell cultures. For these cell culture experiments we took two

approaches: CFUs were counted before and after each exposure period to antibiotics in order to

mimic the recurrent application of antibiotics as in the single-cell experiments, or we evaluated cell

densities (OD) under constant antibiotic concentrations to understand longer exposure to

antibiotics. Cell cultures exposed up to the MIC showed similar qualitative dynamics in terms of

numbers of CFUs when recurrently exposed to antibiotics (Fig. 5a). The higher the antibiotic

concentration the stronger the reduction in CFUs, and the stronger the increase after relieving

exposure to antibiotics, with dampened responses to the second and third exposure and subsequent

releases. These patterns align with predictions of selection for phenotypic resistant cells as revealed

for the single-cell findings. During the long recovery time after the second exposure, most cultures,

except those previously exposed to 64 and 128 μg/ml, reached densities that likely imposed growth

limitation due to resource limitation (i.e., the cultures approached stationary phase). The decline in

CFU during the recurrent exposure shows the continued susceptibility of the cells after series of

selection due to antibiotic exposure. When populations were permanently exposure to antibiotics

(Fig. 5b), they, as expected, initial increased in density before densities were reduced to very low

levels for all cultures exposed to MIC or supra-MIC concentrations, illustrating the susceptibility of

the initial population and no occurrence of resistance mutations during the experiment.

Fig. 5: Colony forming units (CFU) (a), and optical density, OD_{600} (b) of cell cultures exposed to different levels of antibiotics across the duration of the experiment (a) or constant exposure to antibiotics throughout the experiment (b). The grey vertical bars (a) mark the exposure period to the antibiotics. CFU (a) are plotted as boxplots with line connecting mean number of CFU per time when CFU were counted. Single data points are outliers. Note, a jitter is added (a) for better visibility. For the OD curves (b) the dotted lines show the different replicates. the negative control (b) is no bacteria added to medium. CFU (a) and OD (b) are plotted on a log scale.

DISCUSSION

We report on phenological characteristics of single-cells in *E. coli* in response to recurring exposure

to a β-lactam antibiotic. Our findings suggest additional diversity of modes of phenotypic resistance,

given that we show previously undescribed combination of characteristics. Compared to previous

persister studies that quantify fractions of <0.1% persisters (2–4,8,10,25), we reveal a substantial

larger fraction (~25%) of cells that survived 1.5 hours of antibiotic exposure, independent of the level

of antibiotics (Fig. 1a, see also findings at cell culture for <64 μg/ml; Fig. 5a). This considerably larger

fraction we find is best explained by the shorter exposure period to antibiotics we used to select for

phenotypically resistant cells compared to the commonly used 3-5h exposure periods to select for

persisters. We choose the exposure period as 1.5 hours exceed pharmacokinetics actions of

antibiotics (46), illustrated for instance by the time to kill a cell by antibiotics often being comparable

with bacterial generation time (16). For our conditions division time is much shorter than the 1.5 h

exposure period and other studies report on substantially reduced mortality rates after 2h of

exposure (16). Our single-cell survival curves (Fig. 1a) resemble those of a biphasic culture level

killing curve, as is characteristic for persister cells. However the biphasic pattern cannot be directly

compared, as the second phase of our findings concerns a higher fraction of cells and this phase

might also be influenced by the release of the antibiotics after 1.5 h (as killing curves in other

persister studies might be influenced by release time). Fractions of persisters varied among previous

persister studies as selection conditions varied with respect to drug concentration, growth stage

before selection (stationary exponential), preselection (priming of lineages, mutations), or growth

media, to name a few. All of these conditions have been shown to alter persister fractions

(2,3,15,29). In line with previous findings and as expected by selecting on persisters and against

susceptible cells, the MIC marked a survival threshold, illustrating that the initial population was

susceptible and not genetically resistant (Fig. 1), this susceptibility was also confirmed when

evaluated at the cell culture level (Fig. 5). Our results show that the susceptible clonal parental

population consists of phenotypically heterogeneous cells, whereby the cells have different

capacities to sustain exposure to antibiotics, leaving predominantly phenotypic resistant cells after

the first period of exposure.

We found reduced cell growth and division rates under antibiotic exposure, but this reduction was

quantitatively different compared to most previously described persister cells. Growth reduction in

our cells was about halved when exposed to antibiotics, while many other studies have reported

growth arrest or near growth arrest of persister cells during antibiotic exposure (2,4,7,26). Single

studies highlight that fast growth phenotypes with high efflux activity can avoid accumulation of

antibiotics, findings that do not align well with the fact that antibiotics of exponentially growing cells

accumulate within minutes in exposed cells (8,21). In contrast to previously described characteristics

of persiter cells, after release from antibiotics, we did not observe resumption of normal growth

within 2-4 hours (3,4). Our findings hence support studies that explored persisters derived from

exponentially growing cells that showed longer lag times to regain growth, note these cells had been

growth arrest during drug exposure, or highlighting substantial variation in lag times after antibiotic

exposure release (10,25,47). A lack of growth resumption, as we found, is also a characteristic of

persister type II cells and VBNC cells, however VBNC are found in complete growth arrest and are

characterized by a small cell size, and persister type II cells are not environmentally induced but

predetermined (18,26), features not found in our study and other studies (10). If classified, our

phenotypic resistant cells fall best under specialized antibiotically induced persister cell definitions

(9). Quantitative comparison to understand the resumed population growth at the cell culture level

(Fig. 5) in light of the single cell growth reduction (Fig. 2) are challenging, as it remains opaque

whether population growth dynamics are driven by altered cell division rates, altered cell growth, or

altered mortality rates (48). In addition, with respect to heterogeneous selection and cell size, the

average cell size remained constant over the course of the experiment (Fig. 2b) suggesting that

selection did not favour small cells like VBNC cells or small birth sizes as described for persister cells

(18,49), alternatively increased cell size should have been observed if increased frequencies in

filamentation of cells occurred; here we only observed rare filamentation. Finally, variation in cell

size did not increase substantially across the course of the experiment, which would have been

predicted had higher proportions of small cells (VBNC) and filamentous cells simultaneously

emerged, we also did not find other extreme morphological changes as reported for non-dormant

persisters (10).

Our observations differ from genetically fixed resistant cells in various ways. First, at the single-cell

level, unlike resistant cells, we did not find graded survival to different levels of antibiotics (Fig. 1 &

2a), which supports studies that show insensitivity of fractions of persisters up to ~100 µg/ml (15),

although at the cell-culture level we observed graded responses in CFUs (Fig. 5). Second, our cells

approximately halved their growth rates (~50% reduction in cell elongation rates) under drug

exposure (Fig. 2c), whereas a 20% reduction in population growth (evaluated in cell cultures) is

already considered to be an extreme growth cost for genetically fixed resistant cells (50). Third, in

contrast to genetic resistant cells, our cells showed continued susceptibility to repeated exposure to

antibiotics (Fig. 1, Fig. 5a). Finally, resistance is not expected for our single-cell findings, as our

population of cells was directly derived from a susceptible, standard *E. coli* lab strain K-12 MG1655

(see also Fig. 5b). A heteroresistant population, i.e., one composed of multiple sub-populations that

differ in their genetically fixed resistance to antibiotics (exemplified by different MIC), should have

left, post exposure, a resistant population with graded responses to increased levels of antibiotic

concentrations; we found none of these survival, division or growth rate patterns typical for

resistant cells (Fig. 1&2) (50). Despite the isogeneic origin, our initial population was heterotolerant

with a high fraction of cells (~25%) surviving the first exposure and continued showing dampened

sensitivities with respect to mortality for following exposure periods (Fig. 2a). Such continued, but

dampened, sensitivity could be explained by stochastic switching of smaller fractions of cells out of

the tolerant state, which might be related to stochastic gene expression prior, during, and after

exposure to antibiotics (51).

The growth characteristics of cells we describe in arising from exponential growing cells and not

going into growth arrest have important implications for the evolution of genetic resistance.

"Classical" persister cells only contribute to resistance development through repeated rounds of

population expansion by their offspring, as they are growth arrested prior to and during antibiotic

exposure, whereas our phenotypic resistant cells contribute before, during, and after antibiotic

exposure to population expansion, and therefore provide continued opportunities for the

accumulation of mutations towards resistance (31,37). Our phenotypic resistant cells cannot, *a*

priori, be distinguished from susceptible cells by their reduced growth rate (26). Rare types of

persister cells, that have previously described, did not arising from stationary phase cells and

showed slow growth resumption post-exposure, these cells were almost growth arrested under

antibiotic exposure (7,10), and we did not find such growth arrest for our cells.

Our finding that intermediate growing cells, that exhibited robust growth rates, had higher survival

rates compared to growth arrested and quickly-growing cells (Fig. 3 & 4) is surprising to us, as we

had predicted growth arrested cells to be the least affected by β-lactam exposure (2,4,14,26). Our

findings might indicate that phenotypic resistant cells that exhibit high survival have more balanced

and robust intrinsic metabolism (homeostasis) and less dynamic biochemical changes, which

contrasts with molecular level findings that link phenotypic resistance to a perturbed biological

network (33).

The phenotypic resistant state that our cells are characterized by is an induced state, and not an

instantaneous switch, a pre-existing state (4), or a rare occasional stochastic switch to a persistence

or tolerant state out of exponentially growing cells(26). Similar induced switching as we find has

been found for different persistence (10,52) and aligns with findings where increased phenotypic

resistance were induced by antimicrobial peptide exposure (7,53), though our cells seem not to

instantaneously switch to their persistence stage, or at least require different lag times to do so. The

emergence of our phenotypic resistant cells out of an exponentially growing population suggests

that the evolution, or the maintenance, of phenotypic resistance is not conditional on nutrient

limiting condition as has previously been suggested to be the favorable evolutionary route (54).

Our finding that cells cannot be easily identified prior to antibiotic exposure by their reduced growth

rates also supports findings that there are various evolutionary routes to phenotypic resistance

(4,5,9,10,33,55). This contrasts with our knowledge of the much better understood evolution of

genetically fixed resistance, which tends to arise from mutations in a few defined genes that relate

directly to the mechanism by which the antibiotic acts (33). The diversity of evolutionary paths to

phenotypic resistance is illustrated by experimental evolution studies that evolved high phenotypic

resistance by repeated exposure to ampicillin and showed that resistance was achieved through

increased lag time (56); other experimental evolution studies, which exposed stationary phase cells,

evolved phenotypic resistance without prolonged lag times (57), and still other experimental

evolution studies found again other, lag time independent, mutations (33,58).

Future investigations should explore varying exposure time and varied inter exposure time periods

to better understand how selection and characteristics depend on exposure periods. Prolonged

treatment cycles can cause tolerant strains to attain resistance at higher rates (28), while multi-drug

treatment can prolong time to persitence evolution (59), another dimension that could easily be

explored in the future. Different strains should be explored to understand how specific the diverse

phenotypic resistance mechanisms are, but such exploration should be systematic, as part of the

diverse findings on phenotypic resistance might be caused by small differences in experimental

settings not related to the main target (33). A rigorous phenomenological exploration, as we have

done, can provide synergistic insights for tailored molecular exploration. Such molecular exploration

will be interesting as of the expected diversity of molecular and evolutionary paths to phenotypic

resistance. For instance, our cells did not show alterations in cell size, which suggests that our

persistence might not be related to known resistance mechanisms to β-lactams, where filamentation

rate is increased, which in turn stops ampicillin from efficiently targeting the cells (7,58,60). Our

approach also illustrates the challenge and potential of scaling from single cell to the population

level, a link that will provide a deeper understanding of the evolutionary potential towards

resistance, but that is still rarely explored in detail (25). Our single-cell study is a cohort study and

not a cross-cohort study like cell culture experiments, however cross-cohort studies can be achieved

at the single cell level. Other differences, such as inter cellular communication, quorum sensing,

nutrient limitation, or density dependence, among single-cell and cell culture explorations are more

challenging to circumvent but these differences can reveal important conditions that help to reveal

potential resistance mechanisms. The diversity and partly contrasting findings on phenotypic

resistance characteristics, illustrates that we are still on the verge of understanding the role and

mechanisms of such non-genetically determined resistance, and standardized and comparative

studies are needed to solidify and advance our current understanding. Nevertheless, the importance

of phenotypic resistance is reflected in our cell culture experiments, where population crashes were

prevented by fractions of resistant cells when the population was exposed recurrently to antibiotics.

Our results support the idea of diverse modes towards phenotypic resistance, where there is no pre-

determined slow growth, no growth arrest during drug exposure, and no quick resumption of

growth. These characteristics have implications for the evolution of resistance in providing

evolutionary reservoirs and increased mutant supply; this combination of characteristics might even

call for a wider definition of what persister cells are. As for applied implications on medical

treatment, we need to be cautious about interpretation, because directly relating *in vitro* to *in vivo*

results for antibiotic treatment remains challenging.

Acknowledgements
We thank Jens Rolff and Sophie Armitage for comments and discussions.

Funding
This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) – 430170797

References

1. Ventola CL. The antibiotic resistance crisis: causes and threats. P T J [Internet]. 2015 [cited
2021 Feb 24];40(4):277–83. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/25859123%5Cnhttp://www.pubmedcentral.nih.gov/a
rticlerender.fcgi?artid=PMC4378521%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/25859123%
5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4378521

2. Brauner A, Fridman O, Gefen O, Balaban NQ. Distinguishing between resistance, tolerance
and persistence to antibiotic treatment. Vol. 14, Nature Reviews Microbiology. Nature
Publishing Group; 2016. p. 320–30.

3. Brauner A, Shoresh N, Fridman O, Balaban NQ. An Experimental Framework for Quantifying
Bacterial Tolerance. Biophys J [Internet]. 2017 Jun 20 [cited 2018 Nov 5];112(12):2664–71.
Available from: https://www.sciencedirect.com/science/article/pii/S0006349517305519

4. Balaban NQ, Helaine S, Lewis K, Ackermann M, Aldridge B, Andersson DI, et al. Definitions
and guidelines for research on antibiotic persistence. Nat Rev Microbiol. 2019 Jul
1;17(7):441–8.

5. Kaplan Y, Reich S, Oster E, Maoz S, Levin-Reisman I, Ronin I, et al. Observation of universal
ageing dynamics in antibiotic persistence. Nature [Internet]. 2021 Dec 9 [cited 2022 Jan
5];600(7888):290–4. Available from: https://doi.org/10.1038/s41586-021-04114-w

6. Wakamoto Y, Dhar N, Chait R, Schneider K, Signorino-Gelo F, Leibler S, et al. Dynamic
persistence of antibiotic-stressed mycobacteria. Science (80-) [Internet]. 2013 Jan 4 [cited
2021 Feb 24];339(6115):91–5. Available from: https://pubmed.ncbi.nlm.nih.gov/23288538/

7. Goormaghtigh F, Van Melderen L. Single-cell imaging and characterization of Escherichia coli
persister cells to ofloxacin in exponential cultures. Sci Adv. 2019;5(6).

8. Łapińska U, Voliotis M, Lee KK, Campey A, Stone RML, Tuck B, et al. Fast bacterial growth
reduces antibiotic accumulation and efficacy. Elife. 2022;11.

9. Urbaniec J, Xu Y, Hu Y, Hingley-Wilson S, McFadden J. Phenotypic heterogeneity in persisters:
A novel "hunker" theory of persistence [Internet]. Vol. 46, FEMS Microbiology Reviews.
Oxford University Press; 2022 [cited 2023 Jan 4]. p. 1–16. Available from:
https://doi.org/10.1093/femsre/fuab042

10. Umetani M, Fujisawa M, Okura R, Nozoe T, Suenaga S, Nakaoka H, et al. Observation of non-
dormant persister cells reveals diverse modes of survival in antibiotic persistence. bioRxiv
[Internet]. 2022 May 3 [cited 2023 Jan 4];2021.10.28.466227. Available from:
https://doi.org/10.1101/2021.10.28.466227

11. Orman MA, Brynildsen MP. Dormancy is not necessary or sufficient for bacterial persistence.
Antimicrob Agents Chemother [Internet]. 2013 Jul [cited 2021 Mar 8];57(7):3230–9. Available
from: https://pubmed.ncbi.nlm.nih.gov/23629720/

12. Greulich P, Scott M, Evans MR, Allen RJ. Growth-dependent bacterial susceptibility to
ribosome-targeting antibiotics. Mol Syst Biol [Internet]. 2015 Mar 19 [cited 2023 Jan
5];11(3):796. Available from: https://onlinelibrary.wiley.com/doi/10.15252/msb.20145949

13. Vaupel JW, Yashin AI. Heterogeneity's Ruses: Some Surprising Effects of Selection on
Population Dynamics. Am Stat [Internet]. 1985 Aug [cited 2014 Jun 19];39(3):176. Available
from:

http://apps.webofknowledge.com/full_record.do?product=UA&search_mode=GeneralSearc
h&qid=3&SID=P2GPN9Uc9CCP2wpN8k4&page=1&doc=2

14.   Lewis K. Persister cells, dormancy and infectious disease [Internet]. Vol. 5, Nature Reviews
529   Microbiology. Nature Publishing Group; 2007 [cited 2021 Feb 24]. p. 48–56. Available from:
530   www.nature.com/reviews/micro

531   15.   Salcedo-Sora JE, Kell DB. A quantitative survey of bacterial persistence in the presence of
532   antibiotics: Towards antipersister antimicrobial discovery. Antibiotics [Internet]. 2020 Aug 1
533   [cited 2023 Jan 4];9(8):1–36. Available from: https://pubmed.ncbi.nlm.nih.gov/32823501/

534   16.   Svenningsen MS, Svenningsen S Lo, Sørensen MA, Mitarai N. Existence of log-phase
535   Escherichia coli persisters and lasting memory of a starvation pulse. Life Sci Alliance
536   [Internet]. 2022 Feb 1 [cited 2023 Jan 4];5(2). Available from: /pmc/articles/PMC8605324/

537   17.   Pu Y, Li Y, Jin X, Tian T, Ma Q, Zhao Z, et al. ATP-Dependent Dynamic Protein Aggregation
538   Regulates Bacterial Dormancy Depth Critical for Antibiotic Tolerance. Mol Cell. 2019 Jan
539   3;73(1):143-156.e4.

540   18.   Bamford RA, Smith A, Metz J, Glover G, Titball RW, Pagliara S. Investigating the physiology of
541   viable but non-culturable bacteria by microfluidics and time-lapse microscopy. BMC Biol.
542   2017 Dec 21;15(1).

543   19.   Kim JS, Wood TK. Persistent Persister Misperceptions. Front Microbiol [Internet]. 2016 [cited
544   2022 Jan 5];7(DEC):2134. Available from: /pmc/articles/PMC5187198/

545   20.   Bergmiller T, Andersson AMC, Tomasek K, Balleza E, Kiviet DJ, Hauschild R, et al. Biased
546   partitioning of the multidrug efflux pump AcrAB-TolC underlies long-lived phenotypic
547   heterogeneity. Science (80-) [Internet]. 2017 Apr 21 [cited 2023 Jan 5];356(6335):311–5.
548   Available from: https://www.science.org/doi/10.1126/science.aaf4762

549   21.   Cama J, Voliotis M, Metz J, Smith A, Iannucci J, Keyser UF, et al. Single-cell microfluidics
550   facilitates the rapid quantification of antibiotic accumulation in Gram-negative bacteria. Lab
551   Chip [Internet]. 2020 Jul 28 [cited 2023 Jan 5];20(15):2765–75. Available from:
552   https://pubs.rsc.org/en/content/articlehtml/2020/lc/d0lc00242a

553   22.   Rhia M, Stone L, Łapin´ska U, Łapin´ska Ł, Pagliara S, Masi M, et al. Fluorescent macrolide
554   probes – synthesis and use in evaluation of bacterial resistance. RSC Chem Biol [Internet].
555   2020 Dec 8 [cited 2023 Jan 5];1(5):395–404. Available from:
556   https://pubs.rsc.org/en/content/articlehtml/2020/cb/d0cb00118j

557   23.   Goode O, Smith A, Łapińska U, Bamford R, Kahveci Z, Glover G, et al. Heterologous Protein
558   Expression Favors the Formation of Protein Aggregates in Persister and Viable but
559   Nonculturable Bacteria. ACS Infect Dis [Internet]. 2021 Jun 11 [cited 2023 Jan 5];7(6):1848–
560   58. Available from: https://pubs.acs.org/doi/full/10.1021/acsinfecdis.1c00154

561   24.   Dewachter L, Bollen C, Wilmaerts D, Louwagie E, Herpels P, Matthay P, et al. The dynamic
562   transition of persistence toward the viable but nonculturable state during stationary phase is
563   driven by protein aggregation. MBio [Internet]. 2021 Aug 1 [cited 2023 Jan 5];12(4). Available
564   from: https://journals.asm.org/doi/10.1128/mBio.00703-21

565   25.   Bakshi S, Leoncini E, Baker C, Cañas-Duarte SJ, Okumus B, Paulsson J. Tracking bacterial
566   lineages in complex and dynamic environments with applications for growth control and
567   persistence. Nat Microbiol 2021 66 [Internet]. 2021 May 20 [cited 2023 Jan 6];6(6):783–91.
568   Available from: https://www.nature.com/articles/s41564-021-00900-4

26.   Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a phenotypic

switch. Science (80-) [Internet]. 2004 Sep 10 [cited 2021 Mar 4];305(5690):1622–5. Available
from: www.sciencemag.org/cgi/content/full/1099390/DC1

27.   Roostalu J, Jõers A, Luidalepp H, Kaldalu N, Tenson T. Cell division in Escherichia coli cultures
monitored at single cell resolution. BMC Microbiol [Internet]. 2008 Apr 23 [cited 2021 Mar
8];8(1):68. Available from: http://bmcmicrobiol.biomedcentral.com/articles/10.1186/1471-
2180-8-68

28.   Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic tolerance
facilitates the evolution of resistance. Science (80-). 2017 Feb 24;355(6327):826–30.

29.   Levin-Reisman I, Brauner A, Ronin I, Balaban NQ. Epistasis between antibiotic tolerance,
persistence, and resistance mutations. Proc Natl Acad Sci U S A. 2019;116(29):14734–9.

30.   Bakkeren E, Huisman JS, Fattinger SA, Hausmann A, Furter M, Egli A, et al. Salmonella
persisters promote the spread of antibiotic resistance plasmids in the gut. Nature [Internet].
2019 Sep 12 [cited 2021 Mar 11];573(7773):276–80. Available from:
https://doi.org/10.1038/s41586-019-1521-8

31.   Barrett TC, Mok WWK, Murawski AM, Brynildsen MP. Enhanced antibiotic resistance
development from fluoroquinolone persisters after a single exposure to antibiotic. Nat
Commun [Internet]. 2019 Dec 1 [cited 2021 Feb 24];10(1):1–11. Available from:
https://doi.org/10.1038/s41467-019-09058-4

32.   Liu J, Gefen O, Ronin I, Bar-Meir M, Balaban NQ. Effect of tolerance on the evolution of
antibiotic resistance under drug combinations. Science (80-). 2020 Jan 10;367(6474):200–4.

33.   Sulaiman JE, Lam H. Evolution of Bacterial Tolerance Under Antibiotic Treatment and Its
Implications on the Development of Resistance. Front Microbiol [Internet]. 2021 Feb 26 [cited
2021 Mar 11];12:617412. Available from:
https://www.frontiersin.org/articles/10.3389/fmicb.2021.617412/full

34.   El Meouche I, Dunlop MJ. Heterogeneity in efflux pump expression predisposes antibiotic-
resistant cells to mutation [Internet]. Vol. 362, Science. 2018. Available from:
http://science.sciencemag.org/

35.   Van Den Bergh B, Michiels JE, Wenseleers T, Windels EM, Boer P Vanden, Kestemont D, et al.
Frequency of antibiotic application drives rapid evolutionary adaptation of Escherichia coli
persistence. Nat Microbiol 2016 15 [Internet]. 2016 Mar 7 [cited 2023 Jan 5];1(5):1–7.
Available from: https://www.nature.com/articles/nmicrobiol201620

36.   Windels EM, Michiels JE, Fauvart M, Wenseleers T, Van den Bergh B, Michiels J. Bacterial
persistence promotes the evolution of antibiotic resistance by increasing survival and
mutation rates. ISME J [Internet]. 2019 May 1 [cited 2021 Mar 11];13(5):1239–51. Available
from: https://doi.org/10.1038/s41396-019-0344-9

37.   Witzany C, Regoes RR, Igler C. Assessing the relative importance of bacterial resistance,
persistence and hyper-mutation for antibiotic treatment failure. Proc R Soc B Biol Sci
[Internet]. 2022 Nov 9 [cited 2023 Jan 11];289(1986). Available from:
https://royalsocietypublishing.org/doi/10.1098/rspb.2022.1300

38.   Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, et al. Robust growth of Escherichia
coli. Curr Biol [Internet]. 2010 Jun 22 [cited 2011 Jul 18];20(12):1099–103. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2902570&tool=pmcentrez&ren
dertype=abstract

39.   Jouvet L, Rodríguez-Rojas A, Steiner UK. Demographic variability and heterogeneity among

individuals within and among clonal bacteria strains. Oikos. 2018;127(5):728–37.

40.   Steiner UK, Lenart A, Ni M, Chen P, Song X, Taddei F, et al. Two stochastic processes shape
diverse senescence patterns in a single-cell organism. Evolution (N Y) [Internet]. 2019 Apr 14
[cited 2019 Aug 20];73(4):847–57. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/evo.13708

41.   Steiner UK, Lenart A, Ni M, Chen P, Song X, Taddei F, et al. Two stochastic processes shape
diverse senescence patterns in a single-cell organism. Evolution (N Y) [Internet]. 2019 Aug 14
[cited 2018 Aug 15];online ear:10.1111/evo.13708. Available from:
https://www.biorxiv.org/content/early/2018/08/14/105387

42.   Roostalu J, Jõers A, Luidalepp H, Kaldalu N, Tenson T. Cell division in Escherichia coli cultures
monitored at single cell resolution. BMC Microbiol [Internet]. 2008 [cited 2021 May 7];8.
Available from: https://pubmed.ncbi.nlm.nih.gov/18430255/

43.   Conde DA, Staerk J, Colchero F, da Silva R, Schöley J, Maria Baden H, et al. Data gaps and
opportunities for comparative and conservation biology. Proc Natl Acad Sci U S A.
2019;116(19).

44.   R Core Team RA language and environment for statistical computing. R: A language and
environment for statistical computing. [Internet]. Team RDC, editor. R: A language and
environment for statistical computing. Vienna, Austria: R Foundation for Statistical
Computing; 2022. p. 409. (R Foundation for Statistical Computing; vol. 1). Available from:
http://www.r-project.org

45.   Burnham K, Anderson DR. Model selection and multimodel inference: a practical
information-theoretic approach. Second. New York: Springer; 1998.

46.   Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. Pharmacodynamic
functions: a multiparameter approach to the design of antibiotic treatment regimens.
Antimicrob Agents Chemother [Internet]. 2004 Oct [cited 2023 Feb 3];48(10):3670–6.
Available from: https://pubmed.ncbi.nlm.nih.gov/15388418/

47.   Wilmaerts D, Focant C, Matthay P, Michiels J. Transcription-coupled DNA repair underlies
variation in persister awakening and the emergence of resistance. Cell Rep. 2022 Mar
1;38(9):110427.

48.   Knudsen GM, Fromberg A, Ng Y, Gram L. Sublethal concentrations of antibiotics cause shift to
anaerobic metabolism in Listeria monocytogenes and induce phenotypes linked to antibiotic
tolerance. Front Microbiol. 2016 Jul 12;7(JUL):1091.

49.   Hingley-Wilson SM, Ma N, Hu Y, Casey R, Bramming A, Curry RJ, et al. Loss of phenotypic
inheritance associated with ydcI mutation leads to increased frequency of small, slow
persisters in Escherichia coli. Proc Natl Acad Sci U S A [Internet]. 2020 Feb 25 [cited 2023 Jan
5];117(8):4152–7. Available from: /pmc/articles/PMC7049120/

50.   Melnyk AH, Wong A, Kassen R. The fitness costs of antibiotic resistance mutations. Evol Appl
[Internet]. 2015 Mar 1 [cited 2021 Mar 4];8(3):273–83. Available from:
/pmc/articles/PMC4380921/

51.   Rossi NA, El Meouche I, Dunlop MJ. Forecasting cell fate during antibiotic exposure using
stochastic gene expression. Commun Biol [Internet]. 2019 Dec 1 [cited 2021 Jun 2];2(1):1–7.
Available from: https://www.nature.com/articles/s42003-019-0509-0

52.   Dörr T, Lewis K, Vulić M. SOS response induces persistence to fluoroquinolones in Escherichia
coli. PLoS Genet. 2009;5(12):e1000760.

53. 　Rodríguez-Rojas A, Baeder DY, Johnston P, Regoes RR, Rolff J. Bacteria primed by
antimicrobial peptides develop tolerance and persist [Internet]. Vol. 17, PLOS Pathogens.
PLoS Biol; 2021 [cited 2021 May 5]. p. e1009443. Available from:
https://doi.org/10.1371/journal.ppat.1009443

54. 　Windels EM, Van den Bergh B, Michiels J. Bacteria under antibiotic attack: Different strategies
for evolutionary adaptation. Silverman N, editor. PLOS Pathog [Internet]. 2020 May 7 [cited
2021 Mar 11];16(5):e1008431. Available from:
https://dx.plos.org/10.1371/journal.ppat.1008431

55. 　Balaban NQ, Liu J. Evolution under antibiotic treatments: Interplay between antibiotic
persistence, tolerance, and resistance. In: Persister Cells and Infectious Disease [Internet].
Springer International Publishing; 2019 [cited 2021 Apr 19]. p. 1–17. Available from:
https://link.springer.com/chapter/10.1007/978-3-030-25241-0_1

56. 　Fridman O, Goldberg A, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time underlies
antibiotic tolerance in evolved bacterial populations. Nature. 2014 Sep 18;513(7518):418–21.

57. 　Van Den Bergh B, Michiels JE, Wenseleers T, Windels EM, Boer P Vanden, Kestemont D, et al.
Frequency of antibiotic application drives rapid evolutionary adaptation of Escherichia coli
persistence. Nat Microbiol [Internet]. 2016 Mar 7 [cited 2021 Apr 19];1(5):1–7. Available
from: https://www.nature.com/articles/nmicrobiol201620

58. 　Sulaiman JE, Lam H. Proteomic Investigation of Tolerant Escherichia coli Populations from
Cyclic Antibiotic Treatment. J Proteome Res [Internet]. 2020 Feb 7 [cited 2021 Apr
19];19(2):900–13. Available from: https://pubmed.ncbi.nlm.nih.gov/31920087/

59. 　Khare A, Tavazoie S. Extreme Antibiotic Persistence via Heterogeneity-Generating Mutations
Targeting Translation. mSystems [Internet]. 2020 Jan 21 [cited 2021 Apr 19];5(1). Available
from: http://msystems.asm.org/

60. 　Zahir T, Wilmaerts D, Franke S, Weytjens B, Camacho R, Marchal K, et al. Image-Based
Dynamic Phenotyping Reveals Genetic Determinants of Filamentation-Mediated β-Lactam
Tolerance. Front Microbiol [Internet]. 2020 Mar 13 [cited 2021 Mar 9];11:374. Available from:
https://www.frontiersin.org/article/10.3389/fmicb.2020.00374/full