Single-cell phenotypic characteristics of tolerance under recurring antibiotic exposure in *Escherichia*

coli.

Silvia Kollerová[1], Lionel Jouvet[1], Julia Smelková [1], Sara Zunk-Parras[2], Alexandro Rodríguez-Rojas[2], Ul-

rich K. Steiner[1,2*]

[1]Department of Biology, University of Southern Denmark, Odense, Denmark

[2]Biological Institute, Freie Universität Berlin, Berlin, Germany

[*]Corresponding author

Email: ulrich.steiner@fu-berlin.de

Keywords: single-cell, microfluidics, evolution, antibiotic resistance, β-lactam, growth, survival,

metabolism, persister, tolerance

Short title: Phenotypic characteristics of antibiotic tolerant cells

Abstract:

Non-heritable drug resistance, such as tolerance and persistence towards antibiotics, is little

characterized compared to genetic resistance. Tolerance and persistence allow cells to survive

application of antibiotics that are bactericidal to non-tolerant cells. Non-heritable drug resistance

challenges antibiotic treatment, particularly of recurrent infections, and have implications towards

heritable resistance evolution. Tolerant cells have commonly been characterized as growth arrested

cells prior and during antibiotic application that quickly resume growth post-application. Here, we

explore characteristic of tolerant and susceptible *E. coli* single bacteria cells to different levels of

recurrent antibiotic exposure and quantify their occurrence. In using a high throughput single-cell

microfluidic device, we find that tolerant cells reduce their growth rate by about 50%, but contrary to

previous findings do not go into growth arrest or near growth arrest. The growth reduction is induced

by antibiotic exposure and not caused by a stochastic switch or predetermined state as previously

described. Cells exhibiting constant intermediate growth survived best under antibiotic exposure and

selection did not primarily act on fast growing cells, as expected for a β-lactam antibiotic. Control

experiments on population cultures confirmed and challenged scaling of single cell findings to

population level processes. Our findings suggest a prevalent type of tolerance that differs from

previously described tolerance and persister characteristics. Our described characteristics and its high

frequency of occurrence supports acclaims of an underappreciated role of tolerant cells towards

resistance evolution.

Author Summary

Resistance to antibiotics is a major cause of treatment failure in modern medicine. In addition to

genetic resistance, bacterial cells can sustain periods of bactericidal application in a so-called tolerant

state. Previously tolerant cells were characterized by growth arrest prior and during exposure to

antibiotics. Here, we show a distinctive type of tolerance that arises out of normally growing cells,

where during antibiotic exposure, cell growth is slowed but does not go into growth arrest. Cells with

an intermediate growth have the best survival, which is contrary to the assumptions of exposure to a

β-lactam antibiotic where selection is assumed to act against the fastest growing cells and the least

against non-growing cells. The novel characteristics of tolerance that we have uncovered provides

bacteria with another route to evade antibiotics and that could be a considerable threat to the

efficiency of antibiotic treatment.

Introduction

Resistance to antibiotics, in combination with slow progress in discovering novel antibiotics, poses a major challenge to modern medicine [1]. Antibiotic resistant cells sustain bactericidal concentrations of antibiotics through heritable genetically fixed mutations. In contrast, persistent and tolerant cells are in a non-heritable drug resistant phenotypic state. This state allows them to survive periods of bactericidal concentration of antibiotics by non- or very slowly growing[2–5]. They primarily arise from a non- or very slowly-growing sub-population of a susceptible population prior to antibiotic exposure [2–5]. The persister cell stage can be environmentally induced or stochastically switched to, hence, cells can dynamically alter states, and progeny of persiter cells are as susceptible as the parental population from which they derived [4,6]. Quantitatively, below 0.1% of cells in wild-type susceptible populations are persister cells.

Tolerant cells are more common than persister cells and they can also tolerate antibiotic exposures, and are also non- or very slow growing under antibiotic exposure [2]. Though compared to persister cells, tolerant cells can only sustain shorter durations of antibiotic exposure [4]. Under antibiotic application, tolerant cell populations show prolonged minimum duration of killing (MDK) without a change in the minimum inhibitory concentration (MIC), that is, they die slowly under antibiotic application even when exposed to very high concentrations.

A third class of cells that can sustain bactericidal concentration of antibiotics are viable but non-culturable cells (VBNC). In contrast to persister and tolerant cells, VBNC cells do not regain growth 3-4 hours post-antibiotic exposure, but instead remain at low metabolic activity in a deeper dormancy state and do not easily resuscitate, potentially due to a lack of intracellular ATP [7]. Persister and tolerant cells are therefore more susceptible to recurring exposure to antibiotics compared to VBNC cells [7]. VBNC cells are further phenotypically characterized by being small (2.3 ± 0.2μm for E. *coli*) and less elongated compared to persister cells [8].

The definition of these different cell types is not always consistent, and it can be difficult to distinguish

among them empirically, in particular when single-cell data is not available [3,4,9,10]. Only few studies

on resistance, tolerance, and persistence work at the single-cell level, and most of these few studies

evaluated persistence. Overall, these single-cell level studies highlight that that persister cells show

limited growth when exposed to β-lactam or other antibiotics that inhibit cell wall formation or cell

division [10–12], and that switching to persister stages can be irreversible or a reversible dynamic

stochastic process [12]. The stochastic switch to the rare persister state has been observed for growth-

limited cells within exponentially growing *E. coli* populations, with ~80% of persisters arising from the

non-growing sub-population prior to treatment, while the remaining persister cells originated from

cells that were growing prior to treatment [13,14]. Even though some high-quality single-cell data

exists on phenotypic characteristics of tolerant and persister cells, the quantification of the fractions

of susceptible, non-genetic tolerant and resistant cells remains scarce—including quantitative

understanding of cell type specific mortality rates and an accurate quantitative evaluation of cell

growth and division response to antibiotic application. However, such insights would provide deeper

understanding about the evolutionary importance of non-heritable tolerant cells for the evolution of

heritable resistance and efficacy of antibiotic treatment, as high frequencies of non-heritable tolerant

cells provide evolutionary reservoirs and increase the probability of genetically fixed resistance

mutations to occur [15–21].

Here, we use a quantitative demographic and phenomenological approach to quantify the fraction of

cells in a tolerant stage among susceptible stage cells, investigate how tolerant cells differ in their

single-cell characteristics compared to susceptible cells, and study what triggers changes in their

phenotypes. The focus is on tolerant cells, as persister cells are too rare to include in such quantitative

comparison at the single-cell level. Starting with a susceptible clonal population of cells that we expose

to recurrent antibiotic treatment at different concentrations, we track each cell for their growth,

division, and survival in a microfluidic device called a mother machine [22–24]. By exposing cells to a

β-lactam, ampicillin, that corrupts cell wall synthesis, we expected quickly-growing cells to be heavily

selected against, leaving primarily slow-growing tolerant cells that also arose from slowest-growing

population of cells prior to antibiotic exposure[3]. Tolerant cells should not be sensitive to antibiotic

concentrations beyond the MIC [3]. Tolerant cells are expected to regain normal growth rates within

hours post-antibiotic treatment, and such tolerant cells are expected to be the dominating cell type,

compared to VBNC cells, that do not regain growth past antibiotic application. We expected few cells

to switch by environmental induction to tolerant stages, i.e. antibiotic induced switching, as we

expected tolerant stages to be mainly predetermined prior to antibiotic exposure. We also expected

rare stochastic switching. To evaluate how our findings at the single-cell level might scale to cell

culture dynamics, we exposed cell cultures to similar recurring exposures of antibiotics and counted

colony forming units (CFU) before and after each exposure to antibiotics. In addition, we performed

an experiment by measuring the optical density dynamics of constantly exposed cell cultures to the

same levels of antibiotics.

RESULTS

Mortality and selection for tolerance

To select for and quantify characteristics of antibiotic tolerant cells, we exposed single-cells in a first

set of experiments to recurring antibiotic exposure of one of eight different concentrations. As

expected, the probability of cell death, i.e., the force of mortality, increased after the first exposure of

the susceptible parental population to the antibiotic, when exposed to MIC or higher concentrations

(Fig. 1b, Movie SI). Selection for tolerant cells was strong with about two thirds to three quarter of

cells dying before the second exposure period, when exposed to MIC or higher concentrations (Fig.

1a, Movie SI). Mortality peaked 30 to 60 minutes after the first 90-minute antibiotic exposure period,

which suggests that the force of selection peaked with a lag time of 2 to 2.5 hours after the start of

the first exposure. Survival patterns were similar at or above the MIC of 4 µg/ml (Fig. 1a, Cox

proportional hazard model Likelihood ratio test 477.1, p<0.0001, n= 3496; for post hoc testing see SI

Table S1). Neither of the two subsequent 90-minute exposure periods triggered a mortality force that

was as high as the first exposure period. After the initial rise in mortality triggered by the antibiotic

exposure, the probability of death declined, and increased again towards the end of the second

exposure period, though in a less synchronized way among cells exposed to different antibiotic

concentrations. The second peak force of mortality was reached about 30 to 90 minutes after the

second antibiotic exposure, resulting in a similar lag time to the first exposure, i.e., 2 to 3 hours.

Mortality rates continued to decline after the second peak of force of selection and fluctuated for the

remainder of the experiment. Note, cell numbers became small (~20-30 cells per concentration) and

estimates of mortality rates less reliable towards the end of the experiment.

Fig. 1: Survivorship curves (a) and probability of death, i.e., force of mortality (b) of cells exposed to

different levels of antibiotics across the duration of the experiment. The grey curves are control group

cells that were not exposed to any antibiotics. The grey vertical bars mark the exposure period to the

antibiotic. Kaplan-Meier survivorship curves (a) are shown with 95% CI. Probability of death curves (b)

are loess smoothed with 95% CI. X-axis units are given per 4 minutes to reflect the life imaging

frequency.

Division rate, size, and growth responses

In contrast to the control group, cells that had been exposed to MIC or higher antibiotic concentrations

halved their division rate after the first exposure period, changing from ~2 to ~1 division per hour (Fig.

2a, Table 1). This reduction in reproductive rate was reached around 5.5 hours after the onset of the

first exposure. The division rates remained almost constant after the second exposure for another 6

to 7 hours, and then dropped off towards the end of the experiment.

Fig. 2: Cell division rates (a), cell size (in pixels with 100 pixels~2.4μm) (b), and growth rates (c) of cells

exposed to different levels of antibiotics across the duration of the experiment. The grey vertical bars

mark the exposure period to the antibiotics. Division rate (a) size (b) and growth rate (c) curves are

with 95% CI. Note, size data (b) for the control group (0μg/ml) was not available.

Table 1: Model comparison among competing GAM models.

		DF	AIC
Division rate	**S(Age) + Concentr.**	**15.8**	**455232**
	S(Age)	8.4	487667
	Concentr.	8.0	455263
Size	**S(Age) + Concentr.**	**16.8**	**2455427**
	S(Age)	10.8	2460359

	Concentr.	8.0	2458502
Growth rate	**S(Age) + Concentr.**	**17.7**	**-1091065**
	S(Age)	10.5	-1073598
	Concentr.	9.0	-1087172

For each of the three response variables (Division rates, Size, and Growth rates), we compared three competing models. The best supported of these three models is highlighted in bold.

Average cell size remained approximately constant within a given antibiotic concentration. Although we found a surprising amount of variation in size among cells exposed to different levels of antibiotics (Fig. 2b, Table 1), there was no relationship between size and the levels of antibiotics. Cell growth rates responded as predicted by declining during and after the first antibiotic exposure for cells exposed to the MIC or above. We highlight here that growth rates only approximately halved (Fig. 2c, Table 1), which starkly contrasts our predictions based on previous findings where persister or tolerant cells were growth arrested or near growth arrested [4]. The induced growth reduction started about 30 minutes after the onset of antibiotic exposure, which is a lag time that corresponds to approximately the doubling rate (division time) of the parental population. The lowered growth rates stabilized about 4.5 hours after onset of the initial exposure period, suggesting substantial variance in lag time among individual cells' growth response. Induced growth reductions depended partly on antibiotic concentrations with higher levels of antibiotics leading to lower growth rates. However, even with the highest antibiotic concentration, the growth rates did not drop below 1.02-1.03, i.e., 2-3% cell elongation within 4 minutes. Further, after the second antibiotic exposure, growth did not resume in the 11-hour phase without antibiotic application.

Growth specific selection

The growth rates as discussed above (Fig. 2c) are average growth rates at a given point in time, and

even though variance in growth rates does not increase substantially during the experiment (beyond

the expected increase due to decreasing number of cells towards the end of the experiment), selection

might still act preferentially against cells with high growth as expected from a β-lactam antibiotic and

as previously described for persister and tolerant cells [6]. To evaluate the possibility of such

differential selection based on cell growth we analysed together all cells exposed to MIC or above, and

then separated them out into three growth categories: cells that did not grow or shrank (growth

arrested cells with growth rates <1.0), cells that grew at low to intermediate rates (growth rates 1.0-

1.1), and cells that grew quickly (growth rates >1.1) (Fig. 3a). Note that cells can switch dynamically

among these categories throughout the experiment and that mortality and survival is associated only

with the current growth rate of a cell and not a cell's growth rate averaged across time. As expected,

quickly growing cells showed increased probabilities of death upon exposure to antibiotics, but

contrasting to our expectations, the probability of death of growth arrested cells increased

substantially upon exposure to antibiotics, whereas the probability of death of slow-intermediate

growing cells did not change (Fig. 3b, see also SI Table S2 and Table S3).

Fig. 3: Cell growth rate distribution divided into three categories: growth arrested (shrinking or non-

growing; green), intermediate growth (blue), and fast growth (red) (a), and probability of death for

these three groups of growth arrested, intermediate, and quickly-growing cells across the duration of

the experiment (b). The grey vertical bars in panel (b) mark the exposure period to the antibiotics. Note

that cells can switch among growth categories throughout the experiment, that is cells are categorized

according to their current cell growth rate. For this analysis, we have combined cells across levels of

antibiotic exposure ≥4 µg/ml and excluded all cells exposed to 2 µg/ml. A statistical model comparison

showed that mortality differed substantially among the three growth category groups: ΔAIC 25.8

comparing a growth category model and a null model, see also Supplemental Information. Classical p-

value post-hoc analyses showed that quickly growing and growth arrested groups did not differ from

each other in age at death, but the intermediate group did.

To verify the robustness of our finding that selective mortality is dependent on growth rates, and to

broaden our understanding of how growth influences beyond the current growth rate (Fig. 3b), we

did three comparisons between average growth rates of surviving to or dying before specific point in

time during the experiment. First, we compared the average growth rate (averaged across time) of

cells that survived the first exposure period to those that died before the first exposure period.

Second, we compared the average growth rate of cells dying during or shortly after the first exposure

period (cut off at peak of first mortality) to those surviving past the first peak of mortality, and third

we compared similarly for cells that died between the first peak of mortality and the second peak of

mortality to those that survived past the second peak of mortality (Table 2). Before the first exposure,

cells that died had lower average growth rates compared to those that survived past the first

exposure, showing that without selection of antibiotics slower growing cells are preferentially selected

against. Cells that died during the first exposure or shortly after, i.e., during times of strong selection,

had slightly higher growth rates than those that survived past this first increased selection period. This

pattern of growth dependent selection remained for the next period of increased selection after the

second antibiotic exposure. These findings support our initial prediction that faster growing cells

would be preferentially selected against under antibiotic exposure. However, note that the effect sizes

(growth differences) among surviving and dying cells are small, even though they are significantly

different. We interpret our findings on growth specific selection under antibiotic exposure as not primarily acting on fast-growing cells.

Table 2: Differences in average growth rates of cells surviving past or dying during or after specific points in time.

PERIOD	0 – 60 MIN	60 – 328 MIN	328 – 600 MIN	+600 MIN
GOWTH RATE OF CELLS DYING DURING PERIOD (±STDEV)	1.047 (±0.039)	1.067 (±0.032)	1.057 (±0.030)	1.045 (±0.027)
GROWTH RATE OF CELLS SURVIVING PERIOD (±STDEV)	1.072 (0.0412)	1.060 (±0.035)	1.050 (±0.028)	
AIC GROWTH DURING/AFTER	**-8020.8**	**-8816.3**	**-3170.8**	
AIC NULL MODEL	-7985.4	-8772.0	-3161.4	
ΔAIC	35.4	44.3	9.4	

Model comparison among a null model (intercept only model) and a model with the two groups of cells dying during a particular period or surviving that period as explanatory variable and the mean growth rate (per 4 min) as response variable. Three sets of models were fit, one for each period [0 min – 60 min; 60 min – 328 min; 328 min – 600 min]. A Gaussian error structure was used to fit these models.

Robustness of growth rates within cells

In Fig. 2c, we investigated the average growth rate at a given time point. As cells can dynamically alter growth rates between the current and a future point in time, we correlated a cell's current growth rate to its growth rate in the future. Overall correlation between an individual's cell growth rate at a given time t and its growth rate in the immediate future (time t+1 [4 minutes later]) or future time points (time t+3 [12 min later]; time t+5 [20 minutes later]) was weak (Fig.4, Table 3), that is a cell's current growth rate does not well predict its future growth rate. However, there was the exception for cells with intermediate growth rates (1.0-1.1), that showed a strong correlation between current and future growth (Fig. 4), illustrating robust constant growth. Growth arrested cells (growth rate <1 at time t) tended to increase in their growth rates in the at future points in time, while quickly-growing

cells (growth rates >1.1 at time t) tended to decrease their growth rates (Fig. 4). Overall, intermediate

growing cells showed robust and stable growth rates as illustrated by them following tightly the direct

correlation line (Fig. 4), while growth arrested and quickly-growing cells deviate from this direct

correlation. Across cells, we observed substantial temporal dynamics in growth rates. Growth arrested

and quickly-growing cells seemed to be more affected by the concentration of antibiotics, which

suggests that the growth rate responses to different concentrations of antibiotics shown in Fig. 2c are

mainly influenced by the fastest growing and growth arrested cells.

Fig. 4: Correlation in cell growth rates at a given time t, compared to growth rates at time t+1 (4

minutes later) (a), t+3 (12 minutes later) (b), and t+5 (20 minutes later) separated out into populations

recurrently exposed to different levels of antibiotics (as described in the main experiment). All growth

rate correlations are fitted with a GAM with 95% CI. The dashed black line illustrates the direct

correlation among time points.

Table 3: Model comparison among competing GAM models, for correlation of current growth rates to

growth rates at future points in time.

	MODEL	DF	AIC
GROWTH T+1	**Concentr. + S(Growth by Concentr)**	**70.3**	**-441151**
	Concentr. + S(Growth)	17.0	-440468

	S(Growth)	11.0	-438626
	Concentr.	8.0	-425891
GROWTH T+3	**Concentr. + S(Growth by Concentr)**	**69.2**	**-421883**
	Concentr. + S(Growth)	17.0	-421503
	S(Growth)	11.0	-420235
	Concentr.	8.0	-407188
GROWTH T+5	**Concentr. + S(Growth by Concentr)**	**69.0**	**-412563**
	Concentr. + S(Growth)	17.0	-412180
	S(Growth)	11.0	-410925
	Concentr.	8.0	-398709

Best supported model for each response variable (Growth t+1, t+3, t+5 respectively) is highlighted in

bold. GAM models are fitted with the restricted maximum likelihood method, a smoothing parameter

for Growth rate at time t with a shrinkage version of a cubic regression spline as smoothing parameter

(bs=cs) and/or the concentration, including a model that accounts for the interaction of the two

explanatory variables, growth rate and concentration of antibiotics. Growth rates are fitted with a

Gaussian error structure and an inverse link function.

Population level responses

Much research investigating antibiotic tolerance and persistence has been done at the population

level, which makes it challenging to understand mechanisms and describe characteristics operating at

the single-cell level. At the same time, scaling single-cell characteristics to population level can be

challenging, in the following section we therefore report on the control experiments done in cell

cultures. For these cell culture experiments we took two approaches that were suitable as control

experiments: CFUs were counted before and after each exposure period to antibiotics in order to

mimic the recurrent application of antibiotics as in the single-cell experiments, or we evaluated cell

densities (OD) under constant antibiotic concentrations to understand longer exposure to antibiotics.

Cell cultures exposed up to the MIC showed similar dynamics in terms of numbers of CFUs when

recurrently exposed to antibiotics (Fig. 5a). For higher levels of exposure, the higher the antibiotic

concentration the stronger the reduction in CFUs, and the stronger the increase after the first

exposure to antibiotics was relieved. The response to the second and third exposure and antibiotic

release was dampened, which aligns with predictions of selection for tolerant cells as revealed for the

single-cell findings. During the long recovery time after the second exposure, most cultures, except

those previously exposed to 64 and 128 µg/ml, reached densities that likely imposed growth limitation

due to resource limitation (i.e., the cultures approached stationary phase). The decline in CFU during

the recurrent exposure shows the continued susceptibility of the cells after series of selection due to

antibiotic exposure. When populations were permanently exposure to antibiotics (Fig. 5b), they, as

expected, initial increased in density before densities were reduced to very low levels for all cultures

exposed to MIC or above MIC concentrations, illustrating the susceptibility of the initial population.

Fig. 5: Colony forming units (CFU) (a), and optical density, OD_{600} (b) of cell cultures exposed to different levels of antibiotics across the duration of the experiment (a) or constant exposure to antibiotics throughout the experiment (b). The grey vertical bars (a) mark the exposure period to the

antibiotics. CFU (a) are plotted as boxplots with line connecting mean number of CFU per time when CFU were counted. Single data points are outliers. Note, a jitter is added (a) for better visibility. For the OD curves (b) the dotted lines show the different replicates. NA in panel (b) is the negative control, i.e., no bacteria added to medium. CFU (a) and OD (b) are plotted on a log scale.

DISCUSSION

We report a previously undescribed, combination of phenological single-cell characteristics in *E. coli*

in response to recurring exposure to a β-lactam antibiotic. A substantial fraction (>25%) of cells

survived antibiotic exposure independent of the level of antibiotics (Fig. 1a, see also findings at cell

culture for <64 µg/ml; Fig. 5a), this is a considerably larger fraction than previously described for

persister cells (<0.1%) at similar concentrations [2–4]. In line with previous findings and as expected

the MIC marked a survival threshold, illustrating that the initial population was susceptible and not

resistant (Fig. 1), this susceptibility was also confirmed when evaluated at the cell culture level (Fig.

5). Our results show that the susceptible clonal parental population consists of phenotypically

heterogeneous cells, whereby the cells have different capacities to sustain exposure to antibiotics,

leaving predominantly tolerant cells after the first period of exposure.

We found reduced cell growth and division rates under antibiotic exposure, but this reduction was

quantitatively different compared to previously described tolerant cells. Growth reduction in our cells

was only halved, while other studies have reported growth arrest or near growth arrest of tolerant

and persister cells [2,4,10,11]. In contrast to previously described characteristics of tolerant and

persister cells, after release from antibiotics, we did not observe resumption of normal growth within

2-4 hours [3,4]. Such a lack of growth resumption is characteristic of some persister cells (type II) and

VBNC cells, however VBNC are found in complete growth arrest and are characterized by a small cell

size, and these specific persister (type II) cells are not environmentally induced but predetermined

[8,11], features not found in our study. Quantitative comparison to understand the resumed

population growth at the cell culture level (Fig. 5) in light of the single cell growth reduction (Fig. 2)

are challenging, as it remains opaque whether population growth dynamics are driven by increased

cell division rates, increased cell growth, or altered mortality rates [25]. In addition, with respect to

heterogeneous selection and cell size, the average cell size remained constant over the course of the

experiment (Fig. 2b) suggesting that selection did not favor small cells like VBNC cells, or increased cell

size as observed for increased frequencies in filamentation of cells. Filamentation in bacteria is the

continued cell elongation without dividing. Finally, variation in cell size did not increase substantially

across the course of the experiment, which would have been predicted had higher proportions of

small cells (VBNC) and filamentous cells simultaneously emerged.

Our observations differ from genetically fixed resistant cells in various ways. First, at the single-cell

level, unlike resistant cells, we did not find graded survival to different levels of antibiotics (Fig. 1 &

2a), although at the cell-culture level graded responses in CFUs were observed (Fig. 5). Second, our

cells halved their growth rates (50% reduction in cell elongation rates) under exposure (Fig. 2c),

whereas a 20% reduction in population growth (evaluated in cell cultures) is already considered to be

an extreme growth cost for resistant cells [26]. Third, in contrast to resistant cells, our cells showed

continued susceptibility to repeated exposure to antibiotics (Fig. 1, Fig. 5a). Finally, resistance is not

expected for our single-cell findings, as our population of cells was directly derived from a susceptible,

standard *E. coli* lab strain K-12 MG1655 (see also Fig. 5b). A heteroresistant population, i.e., one

composed of multiple sub-populations that differ in their genetically fixed resistance to antibiotics

(exemplified by different MIC), should have left, post exposure, a resistant population with graded

responses to increased levels of antibiotic concentrations. However these were patterns that we did

not find in survival, division or growth rates (Fig. 1&2) [26]. Despite the isogeneic origin, our initial

population was heterotolerant with a high fraction of cells (>25%) surviving the first exposure and

continued showing dampened sensitivities with respect to mortality for following exposure periods

(Fig. 2a). Such continued, but dampened, sensitivity could be explained by stochastic switching of

smaller fractions of cells out of the tolerant state, which might be related to stochastic gene expression

prior, during, and after exposure to antibiotics [27].

In contrast to previously described tolerant cells, our tolerant cells did not arise from stationary phase

cells, they were also not growth-arrested or slow growing prior to exposure [2,4,6,11]. These growth

characteristics have important implications for the evolution of resistance. "Classical" tolerant and

persister cells only contribute to resistance development through repeated rounds of population

expansion by their offspring, as they are growth arrested prior to and during antibiotic exposure,

whereas our tolerant cells contribute before, during, and after antibiotic exposure to population

expansion, and therefore provide continued opportunities for the accumulation of mutations towards

resistance [19]. Our tolerant cells cannot, *a priori*, be distinguished from non-tolerant cells by their

reduced growth rate [11]. Rare types of persister cells, that have previously described, did not arising

from stationary phase cells and showed slow growth resumption post-exposure, these cells were

almost growth arrested under antibiotic exposure [10], and we did not find such growth arrest for our

cells.

Our finding that intermediate growing cells that exhibited robust growth rates, had higher survival

rates compared to growth arrested, and quickly-growing cells (Fig. 3 & 4), is surprising to us as we had

predicted growth arrested cells to be the least affected by β-lactam exposure [2,4,6,11]. Our findings

might indicate that tolerant cells with high survival have more balanced and robust intrinsic

metabolism (homeostasis) and less dynamic biochemical changes, which contrasts with molecular

level findings that link tolerance to a perturbed biological network [15].

The tolerant state that our cells are characterized by is an induced state, and not an instantaneous

switch, a pre-existing state [4], or a rare occasional stochastic switch to a persistence or tolerant state

out of exponentially growing cells as previously described [11]. Similar induced switching as we find

has been found for fluoroquinolone-induced persistence [28] and aligns with findings where increased

tolerance and persistence were induced by antimicrobial peptide exposure [10,29]. The switch to the

induced tolerant state we observe takes on average 2 to 3 hours, with substantial among-cell

variation, emphasizing that it is not an instantaneous switch. This lag time spans a minimum of 2 to 3

divisions or a time during which the initial cell's biomass increases by about 4-8 times, and this lag

time is independent of the levels of antibiotic exposure. The emergence of our tolerant cells out of an

exponentially growing population suggests that the evolution, or at least the maintenance, of

tolerance is not conditional on nutrient limiting condition as has previously been suggested to be the

favorable evolutionary route [30].

Our finding that cells cannot be easily identified prior to antibiotic exposure by their reduced growth

rates also supports findings that there are various evolutionary routes to tolerance [4,5,15,31]. This

contrasts with our knowledge of the much better understood evolution of genetically fixed resistance,

which tends to arise from mutations in a few defined genes that relate directly to the mechanism by

which the antibiotic acts [15]. The diversity of evolutionary paths to tolerance is illustrated by

experimental evolution studies that evolved high tolerance by repeated exposure to ampicillin and

showed that tolerance was achieved through increased lag time [32]; other experimental evolution

studies, which exposed stationary phase cells, evolved tolerance without prolonged lag times [33],

and still other experimental evolution studies found again other mutations [15,34].

Our findings contrast in surprising ways with previous findings on tolerance and persistence. We

therefore need to ask why we find such differences. For instance characteristics and frequency of

persisters depend on culture conditions and growth phase, even when the same species, strains, and

antibiotics are used [45]. Studies that enriched persisters by preselecting dormant cells [14,46] could

not detect our cell types, as we did not find clearly predefined characteristics [10]. Results might also

be biased when working with high-frequency persister mutants, such as hip mutants, as they show

altered characteristics compared to wild type cells [11]. Growth medium can influence phenotypes,

such as the filamentation rate, or the recovery rate from filamentous states [22,24]. Exposure time to

antibiotics applies different selection pressure for tolerance, with our exposure time of 90 minutes

being shorter compared to other experiments that select cells by exposing them to ~5 hours [10,11].

The quantitative depth of our results, reporting on several hundred tolerant cells selected from a few

thousand cells, also differs from other studies. The exciting core findings of Goormaghtigh and Van

Melderen [10] rely on 23 persister cells selected out of more than 100,000 cells. In sum, the

characteristics we have described are unlikely to be explained solely by the experimental setup and

growth conditions and are more likely novel tolerance characteristics.

Future investigations should explore varying exposure time and varied inter exposure time periods.

Prolonged treatment cycles can cause tolerant strains to attain resistance at higher rates [16], while

multi-drug treatment can prolong time to tolerance evolution [47], another dimension that could

easily be explored in the future. Different strains can be explored, but such exploration should be

systematic, as part of the diverse findings on tolerance might be caused by small differences in

experimental settings not related to the main target [15]. A rigorous phenomenological exploration,

as we have done, can provide synergistic insights for tailored molecular exploration. Such molecular

exploration will be interesting as of the expected diversity of molecular and evolutionary paths to

tolerance. This molecular diversity likely results in a diversity of phenomenological characteristics of

tolerance. For instance, our cells did not show alterations in cell size, which suggests that our tolerance

might not be related to known tolerance mechanisms to β-lactams, where filamentation rate is

increased, which in turn stops ampicillin from efficiently targeting the cells [10,34,48]. Our approach

also illustrates the challenge and potential of scaling from single cell to the population level, a link that

will provide a deeper understanding of the evolutionary potential towards resistance, but which is still

rarely explored in detail. Our single-cell study is a cohort study and not a cross-cohort study like cell

culture experiments, however cross-cohort studies can be achieved at the single cell level. Other

differences, such as inter cellular communication, quorum sensing, nutrient limitation, or density

dependence, among single-cell and cell culture explorations are more challenging to circumvent but

these differences can reveal important conditions that help to reveal potential tolerance mechanisms.

Microfluidic systems prevent density dependent and nutrient limiting regulations, single-cell level

understanding affords a distinction among mortality, division, size, and growth effects, which are

surmised by the cell culture level measurements of CFU or OD. The diversity and partly contrasting

findings on persister and tolerant cells, illustrates that we are still on the verge of understanding the

role and mechanisms of tolerance, and standardized and comparative studies are needed to solidify

and advance our current understanding. Nevertheless, the importance of tolerance is reflected in our

cell culture experiments, where population crashes were prevented by tolerant cells when the

population was exposed recurrently to antibiotics. Our results support the idea of a novel form of

tolerance, where there is no pre-determined slow growth, no growth arrest, and no quick resumption

of growth. These characteristics have implications for the evolution of resistance in providing

evolutionary reservoirs and increased mutant supply; this novel combination of characteristics might

even call for a wider definition of what tolerant cells are. As for applied implications on medical

treatment, we need to be cautious about interpretation, because directly relating *in vitro* to *in vivo*

results for antibiotic treatment remains challenging.

MATERIALS AND METHODS

We conducted four experiments, three of them with three antibiotic exposure periods of 90 minutes,

two of these experiments collected single-cell data and one cell culture level data, i.e. CFU counts. The

fourth experiment, which was at the cell culture level, continuously exposed cells to antibiotics and

served as a control experiment. All experiments, single-celled and cell culture experiments, were done

in supplemented minimal media (M9), and all cultures were derived from *E. coli* strain MG1655, a K12

strain. We chose antibiotic concentrations similar to previously explored ranges, that spanning from

below MIC to >30 times MIC, and exposed cells to antibiotics for periods that would cover three to

four divisions under non-antibiotic conditions. Post-exposure recovery times were also chosen on

previous results reporting on regained growth of persister cells within 3-4 hours post-treatment [10–

12]. We choose ampicillin as antibiotic, a β-lactamase that inhibits bacterial cell wall synthesis,

because this antibiotic has been extensively used in previous persister studies [11]. Comparing growth

dynamics, size, division dynamics, and mortality in response to recurring exposure to antibiotics allows

us to differentiate among tolerant and VBNC cells, and in comparing lag times we differentiate

antibiotic induced and stochastic switching to tolerance.

For the experiments with recurrent exposure, the first exposure period was 60-150 minutes past onset

of the experiments, the second exposure period was 330-420 minutes past onset of the experiments,

i.e. 180 minutes past the first exposure period, and the last exposure period was 1080-1170 minutes

past onset of the experiments, i.e. 660 minutes past the second exposure period. For the single-cell

experiments concentrations of ampicillin during the exposure periods were set to 2, 4, 16, 24, 32, 64

or 128µg/ml. As comparison and reference, data from another experiment was added as control

group, where cells were never exposed to antibiotics [23]. For the experiments at the cell culture level,

the concentrations of 0, 2, 4, 8, 16, 32, 64 or 128µg/ml were used, and in addition a negative control

testing for potential contamination was done.

To collect phenological data at the single-cell level, each cell was tracked by time-lapse life-imaging

(15 frames/hour) in a microfluidic (PDMS) chip, a mother machine [23], that was placed under a

temperature-controlled (37°C) inverted microscope. The cells loaded into the microfluidic chip came

from an exponentially growing culture. Given that the cell culture the loaded cells originated from was

in exponential growth phase, the vast majority of cells loaded were in an exponential growth phase

[24]. The microfluidic chip the cells grew in contained four main laminar flow channels (feeding

channels) and hundreds of dead-end side channels, in the latter the rod-shaped bacteria-cells grew.

Only one focal cell, the one closest to the dead-end of the side channel, was tracked for its growth

(cell elongation), size, time of each division, and death. Cell death was determined by propidium iodide

(PI) that emits a strong fluorescent signal when the cell wall lysis—the moment the PI can enter the

cell—and bonds to the cells' DNA. Image analysis was performed using a customized software and the

resulting data was subsequently analysed in program R [49] using general linear, generalized linear,

and nonlinear (Generalized Additive Models, GAM) models. Models were compared based on

information criteria (AIC)[50]. Kaplan-Meier survivorship curves (Fig. 1a) are computed with the

survival package (survfit function), probability of death curves (Fig. 1b) are plotted with a loess

smoothing function (geometric_smooth function; package ggplot2). Competing GAM models (Table

1) are fitted with the restricted maximum likelihood method (REML), a smoothing parameter for age

with a shrinkage version of a cubic regression spline as smoothing parameter (bs=cs) and/or the

concentration. Division rates (Fig.2a) are fitted with a loess smoothing function (geometric_smooth;

package ggplot2) using a binomial error structure with a logit link function, size data (Fig. 2b) is fitted

with a GAM model applying a Gaussian error structure and an identity link function, and the Growth

rates (Fig. 2c) are fitted with GAM model using a Gamma error structure and an inverse link function.

Note, smoothing functions (GAM) are highly sensitive at the extremes as partly illustrated by the

diverse initial average cell sizes (Fig. 2b) of cells that came from the same parental population and that

had not yet been exposed to antibiotics during the 60-minute lasting acclimation phase. Probability of

death curves (Fig. 3b) are plotted with a loess smoothing function with 95% CI (geometric_smooth;

package ggplot2). More details on the microchip production, loading of the chip, the bacteria growth,

imaging, and analyses on the single-cell experiments are given in the supplemental material.

To collect the number of colony forming unit (CFU) data and the cell culture density data (OD_{600}), we

initially seeded these experiments using cells from an exponential growing culture. Each experiment

had 8 replicates, the experiments were done in 96 well-plates, and cells grew at 37°C and vigorous

shaking in their respective well (see Fig. S1 for details). To achieve the sequence of antibiotic exposure,

the cell culture plates were centrifuged, super-natant was discarded and cells were resuspended in

fresh media with or without antibiotics depending on treatment. The concentration changed to match

those used for the single-celled experiments and CFU were counted by serial dilution plating on LB

agar plates when antibiotic exposure changed. To track cell culture density at constant antibiotic

exposure, the initial 8 replicates for each concentration of the antibiotic exposure were incubated at

37°C and shaking in a 96-well-plate and the optical density (OD600) was measured every 20 minutes

for 24 h in a plate reader Synergy H1 (Biotek, Germany). Fig. 5b the OD curves are fitted with a loess

smoothing function with 95% CI (geometric_smooth with a span of 0.1; package ggplot2). Further

information on these cell culture experiments is given in the supplemental material.

Acknowledgements

We thank Jens Rolff and Sophie Armitage for comments and discussions.

References

1.   Ventola CL. The antibiotic resistance crisis: causes and threats. P T J. 2015;40: 277–83.
doi:Article

2.   Brauner A, Fridman O, Gefen O, Balaban NQ. Distinguishing between resistance, tolerance
and persistence to antibiotic treatment. Nature Reviews Microbiology. Nature Publishing
Group; 2016. pp. 320–330. doi:10.1038/nrmicro.2016.34

3.   Brauner A, Shoresh N, Fridman O, Balaban NQ. An Experimental Framework for Quantifying
Bacterial Tolerance. Biophys J. 2017;112: 2664–2671. doi:10.1016/J.BPJ.2017.05.014

4.   Balaban NQ, Helaine S, Lewis K, Ackermann M, Aldridge B, Andersson DI, et al. Definitions
and guidelines for research on antibiotic persistence. Nat Rev Microbiol. 2019;17: 441–448.
doi:10.1038/s41579-019-0196-3

5.   Kaplan Y, Reich S, Oster E, Maoz S, Levin-Reisman I, Ronin I, et al. Observation of universal
ageing dynamics in antibiotic persistence. Nature. 2021;600: 290–294. doi:10.1038/s41586-
021-04114-w

6.   Lewis K. Persister cells, dormancy and infectious disease. Nature Reviews Microbiology.
Nature Publishing Group; 2007. pp. 48–56. doi:10.1038/nrmicro1557

7.   Pu Y, Li Y, Jin X, Tian T, Ma Q, Zhao Z, et al. ATP-Dependent Dynamic Protein Aggregation
Regulates Bacterial Dormancy Depth Critical for Antibiotic Tolerance. Mol Cell. 2019;73: 143-
156.e4. doi:10.1016/j.molcel.2018.10.022

8.   Bamford RA, Smith A, Metz J, Glover G, Titball RW, Pagliara S. Investigating the physiology of
viable but non-culturable bacteria by microfluidics and time-lapse microscopy. BMC Biol.
2017;15. doi:10.1186/s12915-017-0465-4

9.   Kim JS, Wood TK. Persistent Persister Misperceptions. Front Microbiol. 2016;7: 2134.
doi:10.3389/FMICB.2016.02134

10.   Goormaghtigh F, Van Melderen L. Single-cell imaging and characterization of Escherichia coli
persister cells to ofloxacin in exponential cultures. Sci Adv. 2019;5.
doi:10.1126/sciadv.aav9462

11.   Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a phenotypic
switch. Science (80-). 2004;305: 1622–1625. doi:10.1126/science.1099390

12.   Wakamoto Y, Dhar N, Chait R, Schneider K, Signorino-Gelo F, Leibler S, et al. Dynamic
persistence of antibiotic-stressed mycobacteria. Science (80-). 2013;339: 91–95.
doi:10.1126/science.1229858

13.   Roostalu J, Jõers A, Luidalepp H, Kaldalu N, Tenson T. Cell division in Escherichia coli cultures
monitored at single cell resolution. BMC Microbiol. 2008;8: 68. doi:10.1186/1471-2180-8-68

14.   Orman MA, Brynildsen MP. Dormancy is not necessary or sufficient for bacterial persistence.
Antimicrob Agents Chemother. 2013;57: 3230–3239. doi:10.1128/AAC.00243-13

15.   Sulaiman JE, Lam H. Evolution of Bacterial Tolerance Under Antibiotic Treatment and Its
Implications on the Development of Resistance. Front Microbiol. 2021;12: 617412.
doi:10.3389/fmicb.2021.617412

16.   Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic tolerance
facilitates the evolution of resistance. Science (80-). 2017;355: 826–830.
doi:10.1126/science.aaj2191

17.   Levin-Reisman I, Brauner A, Ronin I, Balaban NQ. Epistasis between antibiotic tolerance,
persistence, and resistance mutations. Proc Natl Acad Sci U S A. 2019;116: 14734–14739.
doi:10.1073/pnas.1906169116

18.   Bakkeren E, Huisman JS, Fattinger SA, Hausmann A, Furter M, Egli A, et al. Salmonella
persisters promote the spread of antibiotic resistance plasmids in the gut. Nature. 2019;573:
276–280. doi:10.1038/s41586-019-1521-8

19.   Barrett TC, Mok WWK, Murawski AM, Brynildsen MP. Enhanced antibiotic resistance
development from fluoroquinolone persisters after a single exposure to antibiotic. Nat
Commun. 2019;10: 1–11. doi:10.1038/s41467-019-09058-4

20.   Liu J, Gefen O, Ronin I, Bar-Meir M, Balaban NQ. Effect of tolerance on the evolution of
antibiotic resistance under drug combinations. Science (80-). 2020;367: 200–204.
doi:10.1126/science.aay3041

21.   El Meouche I, Dunlop MJ. Heterogeneity in efflux pump expression predisposes antibiotic-
resistant cells to mutation. Science (80-). 2018. doi:10.1126/science.aar7981

22.   Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, et al. Robust growth of Escherichia
coli. Curr Biol. 2010;20: 1099–103. doi:10.1016/j.cub.2010.04.045

23.   Jouvet L, Rodríguez-Rojas A, Steiner UK. Demographic variability and heterogeneity among
individuals within and among clonal bacteria strains. Oikos. 2018;127: 728–737.
doi:10.1111/oik.04292

24.   Steiner UK, Lenart A, Ni M, Chen P, Song X, Taddei F, et al. Two stochastic processes shape
diverse senescence patterns in a single-cell organism. Evolution (N Y). 2019;73: 847–857.
doi:10.1111/evo.13708

25.   Knudsen GM, Fromberg A, Ng Y, Gram L. Sublethal concentrations of antibiotics cause shift to
anaerobic metabolism in Listeria monocytogenes and induce phenotypes linked to antibiotic
tolerance. Front Microbiol. 2016;7: 1091. doi:10.3389/FMICB.2016.01091/BIBTEX

26.   Melnyk AH, Wong A, Kassen R. The fitness costs of antibiotic resistance mutations. Evol Appl.
2015;8: 273–283. doi:10.1111/eva.12196

27.   Rossi NA, El Meouche I, Dunlop MJ. Forecasting cell fate during antibiotic exposure using
stochastic gene expression. Commun Biol. 2019;2: 1–7. doi:10.1038/s42003-019-0509-0

28.   Dörr T, Lewis K, Vulić M. SOS response induces persistence to fluoroquinolones in Escherichia
coli. PLoS Genet. 2009;5: e1000760. doi:10.1371/journal.pgen.1000760

29.   Rodríguez-Rojas A, Baeder DY, Johnston P, Regoes RR, Rolff J. Bacteria primed by
antimicrobial peptides develop tolerance and persist. PLOS Pathogens. PLoS Biol; 2021. p.
e1009443. doi:10.1371/journal.ppat.1009443

30.   Windels EM, Van den Bergh B, Michiels J. Bacteria under antibiotic attack: Different strategies
for evolutionary adaptation. Silverman N, editor. PLOS Pathog. 2020;16: e1008431.
doi:10.1371/journal.ppat.1008431

31.   Balaban NQ, Liu J. Evolution under antibiotic treatments: Interplay between antibiotic
persistence, tolerance, and resistance. Persister Cells and Infectious Disease. Springer
International Publishing; 2019. pp. 1–17. doi:10.1007/978-3-030-25241-0_1

32.   Fridman O, Goldberg A, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time underlies
antibiotic tolerance in evolved bacterial populations. Nature. 2014;513: 418–421.
doi:10.1038/nature13469

33.   Van Den Bergh B, Michiels JE, Wenseleers T, Windels EM, Boer P Vanden, Kestemont D, et al.
Frequency of antibiotic application drives rapid evolutionary adaptation of Escherichia coli
persistence. Nat Microbiol. 2016;1: 1–7. doi:10.1038/nmicrobiol.2016.20

34.   Sulaiman JE, Lam H. Proteomic Investigation of Tolerant Escherichia coli Populations from
Cyclic Antibiotic Treatment. J Proteome Res. 2020;19: 900–913.
doi:10.1021/acs.jproteome.9b00687

35.   Lee JH, Lee J. Indole as an intercellular signal in microbial communities. FEMS Microbiology
Reviews. Blackwell Publishing Ltd; 2010. pp. 426–444. doi:10.1111/j.1574-6976.2009.00204.x

36.   Diez S, Ryu J, Caban K, Gonzalez RL, Dworkin J. The alarmones (p)ppGpp directly regulate
translation initiation during entry into quiescence. Proc Natl Acad Sci U S A. 2020;117: 15565–
15572. doi:10.1073/pnas.1920013117

37.   Vega NM, Allison KR, Khalil AS, Collins JJ. Signaling-mediated bacterial persister formation.
Nat Chem Biol. 2012;8: 431–433. doi:10.1038/nchembio.915

38.   Maisonneuve E, Gerdes K. Molecular mechanisms underlying bacterial persisters. Cell. Cell
Press; 2014. pp. 539–548. doi:10.1016/j.cell.2014.02.050

39.   Van Acker H, Van Dijck P, Coenye T. Molecular mechanisms of antimicrobial tolerance and
resistance in bacterial and fungal biofilms. Trends in Microbiology. Elsevier Ltd; 2014. pp.
326–333. doi:10.1016/j.tim.2014.02.001

40.   Olsen I. Biofilm-specific antibiotic tolerance and resistance. European Journal of Clinical
Microbiology and Infectious Diseases. Springer Verlag; 2015. pp. 877–886.
doi:10.1007/s10096-015-2323-z

41.   Harms A, Maisonneuve E, Gerdes K. Mechanisms of bacterial persistence during stress and
antibiotic exposure. Science (80-). 2016;354. doi:10.1126/science.aaf4268

42.   Michiels JE, Van den Bergh B, Verstraeten N, Michiels J. Molecular mechanisms and clinical
implications of bacterial persistence. Drug Resist Updat. 2016;29: 76–89.
doi:10.1016/j.drup.2016.10.002

43.   Page R, Peti W. Toxin-antitoxin systems in bacterial growth arrest and persistence. Nat Chem
Biol. 2016;12: 208–214. doi:10.1038/nchembio.2044

44.   Wilmaerts D, Windels EM, Verstraeten N, Michiels J. General Mechanisms Leading to
Persister Formation and Awakening. Trends in Genetics. Elsevier Ltd; 2019. pp. 401–411.
doi:10.1016/j.tig.2019.03.007

45.   Harms A, Fino C, Sørensen MA, Semsey S, Gerdes K. Prophages and growth dynamics
confound experimental results with antibiotic-tolerant persister cells. MBio. 2017;8.
doi:10.1128/mBio.01964-17

46. Shah D, Zhang Z, Khodursky A, Kaldalu N, Kurg K, Lewis K. Persisters: A distinct physiological
state of E. coli. BMC Microbiol. 2006;6: 1–9. doi:10.1186/1471-2180-6-53

47. Khare A, Tavazoie S. Extreme Antibiotic Persistence via Heterogeneity-Generating Mutations
Targeting Translation. mSystems. 2020;5. doi:10.1128/msystems.00847-19

48. Zahir T, Wilmaerts D, Franke S, Weytjens B, Camacho R, Marchal K, et al. Image-Based
Dynamic Phenotyping Reveals Genetic Determinants of Filamentation-Mediated β-Lactam
Tolerance. Front Microbiol. 2020;11: 374. doi:10.3389/fmicb.2020.00374

49. R Core Team RA language and environment for statistical computing. R: A language and
environment for statistical computing. Team RDC, editor. R Foundation for Statistical
Computing. Vienna, Austria: R Foundation for Statistical Computing; 2016. p. 409.
doi:10.1007/978-3-540-74686-7

50. Burnham K, Anderson DR. Model selection and multimodel inference : a practical
information-theoretic approach. Second. New York: Springer; 1998.