1 **Title: Complex structure of the *Bacillus subtilis* CdaA c-di-AMP cyclase domain and the**
2 **phosphoglucomutase GlmM**
3

4 **Monisha Pathania[a], Tommaso Tosi[a], Charlotte Millership[a], Fumiya Hoshiga[a], Rhodri M. L.**
5 **Morgan[b] Paul S. Freemont[c,d,e,#], Angelika Gründling[a,#]**
6
7
[a] Section of Molecular Microbiology and Medical Research Council Centre for Molecular Bacteriology
and Infection, Imperial College London, London SW7 2AZ, United Kingdom
[b] Department of Life Sciences, Imperial College London, London SW7 2AZ, United Kingdom
[c] London Biofoundry, Imperial College Translation and Innovation Hub, White City Campus, 80 Wood
Lane, London, W12 0BZ, United Kingdom
[d] Section of Structural and Synthetic Biology, Department of Infectious Disease, Imperial College
London, London, SW7 2AZ, United Kingdom
[e] UK Dementia Research Institute Centre for Care Research and Technology, Imperial College London,
London, United Kingdom
17
18
19 [#]To whom correspondence should be addressed:
20 Paul S. Freemont – p.freemont@imperial.ac.uk,
21 Angelika Gründling – a.grundling@imperial.ac.uk

22
23
24 **Running title:** *B. subtilis* CdaA:GlmM complex
25 **Key words:** Bacillus, cyclic dinucleotide, signalling, crystallography, protein structure
26
27
28
29

Abstract

Cyclic-di-adenosine monophosphate (c-di-AMP) is an important nucleotide signalling molecule, which plays a key role in osmotic regulation in bacteria. Cellular c-di-AMP levels are tightly regulated, as both high and low levels have a negative impact on bacterial growth. Here, we investigated how the activity of the main *Bacillus subtilis* c-di-AMP cyclase CdaA is regulated by the phosphoglucomutase GlmM. c-di-AMP is produced from two molecules of ATP by proteins containing a deadenylate cyclase (DAC) domain. CdaA is a membrane-linked cyclase with an N-terminal transmembrane domain followed by the cytoplasmic DAC domain. Here we show, using the soluble catalytic *B. subtilis* CdaA$_{CD}$ domain and purified full-length GlmM or the GlmM$_{F369}$ variant lacking the C-terminal flexible domain 4, that the cyclase and phosphoglucomutase form a stable complex *in vitro* and that GlmM is a potent cyclase inhibitor. We determined the crystal structure of the individual *B. subtilis* CdaA$_{CD}$ and GlmM proteins, both of which form dimers in the structures, and of the CdaA$_{CD}$:GlmM$_{F369}$ complex. In the complex structure, a CdaA$_{CD}$ dimer is bound to a GlmM$_{F369}$ dimer in such a manner that GlmM blocks the oligomerization of CdaA$_{CD}$ and formation of active head-to-head cyclase oligomers, thus providing molecular details on how GlmM acts as cyclase inhibitor. The function of a key amino acid residue in CdaA$_{CD}$ in complex formation was confirmed by mutagenesis analysis. As the amino acids at the CdaA$_{CD}$:GlmM interphase are conserved, we propose that the observed inhibition mechanism of CdaA by GlmM is conserved among Firmicutes.

Introduction

Nucleotide signalling molecules play important roles in helping bacteria to rapidly adapt to changing environmental conditions (1,2). One such signalling nucleotide, cyclic-di-adenosine monophosphate (c-di-AMP), which was discovered a little more than a decade ago (3), plays an important function in the osmotic regulation of bacteria by controlling potassium and osmolyte uptake (4-8). c-di-AMP also plays an important function in regulating cell size, either directly or indirectly through its function in osmotic regulation, cell-wall integrity and susceptibility to beta-lactam antibiotics, which target the synthesis of the peptidoglycan cell wall (9-12).

The function of c-di-AMP has been most extensively studied in a range of Firmicutes bacteria including the Gram-positive model organism *Bacillus subtilis* and Gram-positive bacterial pathogens such as *Staphylococcus aureus, Listeria monocytogenes* and several *Streptococcus* species (9,10,13-17). From these studies, it has become apparent that the cellular level of c-di-AMP needs to be tightly regulated as both an excess and a lack of c-di-AMP can negatively impact bacterial growth, physiology and virulence (17,18). To achieve the optimal level, a dynamic equilibrium must exist between the synthesis of c-di-AMP via diadenylate cyclases and its degradation into 5′-phosphadenylyl-adenosine (pApA) or two molecules of AMP by phosphodiesterases (18-20). As part of the current study, we investigated how the activity the *B. subtilis* c-di-AMP cyclase CdaA is regulated by GlmM, a phosphoglucomutase enzyme required for the synthesis of an essential peptidoglycan precursor.

c-di-AMP is formed from two molecules of ATP by enzymes containing a diadenylate cyclase (DAC) domain. These enzymes have been extensively characterized structurally as well as biochemically, but how their activity is regulated is an aspect that remains poorly understood. *B. subtilis* codes for three diadenylate cyclase enzymes (3,21-24). CdaA (also referred to as DacA in some bacteria) is a membrane-bound cyclase with three predicted N-terminal transmembrane helices and a cytoplasmic catalytic DAC domain. CdaA (DacA) is the "housekeeping" c-di-AMP cyclase in Firmicutes, as it is conserved and often the sole c-di-AMP cyclase in phylum (25,26). The two other *B. subtilis* c-di-AMP cyclases, DisA and CdaS, are soluble proteins, not as widely distributed among bacteria and have more specialized functions, with DisA involved in controlling DNA-repair processes during sporulation or spore outgrowth and CdaS also specifically expressed during the sporulation process (22,23,27). While there are no publications on the 3D-structures of the *B. subtilis* c-di-AMP cyclases, structures are available for the cytoplasmic enzymatic domains of the *L. monocytogenes* and *S. aureus* CdaA/DacA homologs (21,28), the DisA homolog from *Thermotoga maritima* (3) and the CdaS homolog from *B. cereus* (PDB 2FB5). These studies revealed that DAC domains have a mixed αß-fold, with highly conserved DGA and RHR amino acid motifs required for ligand binding (3,21,29). Formation of c-di-AMP requires a head-to-head conformation of two DAC domains. This was first demonstrated in the crystal structure of DisA, a protein which forms an octamer with four DAC domain dimers in the active head-to-head conformation (3). While also the *L. monocytogenes* and *S. aureus* CdaA/DacA catalytic

domains and the CdaS protein, where present as dimers and hexamers, respectively, they were in an
inactive conformation. These proteins therefore either need to rearrange or more likely form higher
oligomers in order to yield active enzymes with DAC domains in the head-to-head dimer conformation
(24,28). Recently, another structure of the cytoplasmic catalytic domain of the *L. monocytogenes* CdaA
enzyme (Δ100CdaA) has been reported with a c-di-AMP bound between two monomers, which based
on the crystal cell packing, were arranged in an active dimer of dimer configuration (30). These finding
are consistent with the idea that CdaA (DacA) enzymes will need to form higher oligomers to achieve
an active enzyme configuration. Hence, factors influencing the ability of c-di-AMP cyclases to rearrange
into an active conformation or to form higher oligomers will be able to regulate the activity of these
enzymes.
95   The genetic arrangement and operon structure coding for the "housekeeping" c-di-AMP cyclase
CdaA (DacA) is conserved in Firmicutes (29,31). Two genes, coding for the membrane-linked CdaA
regulator CdaR (also named YbbR in some bacteria) and cytoplasmically-located peptidoglycan
precursor synthesis enzyme GlmM, are found downstream and in an operon with *cdaA* (29,31). Through
recent studies in *B. subtilis, L. monocytogenes, Lactococcus lactis* and *S. aureus,* it is has become
apparent that these three genes are not only co-transcribed but that the encoded proteins also from a
complex and that CdaR and GlmM can regulate the activity of the c-di-AMP cyclase CdaA (29,32,33).
While CdaR has been reported to function as both an activator and repressor of CdaA activity depending
on the growth conditions, GlmM has been shown to be a potent inhibitor of the cyclase activity
(17,29,31,32,34). However, the molecular mechanisms on how the CdaA cyclase activity is regulated
by these proteins are not yet known and this was further investigated as part of this study.
106   GlmM is a phosphoglucomutase enzyme catalysing the conversion of glucosamine-6-phosphate
to glucosamine-1-phosphate, which is subsequently used to produce the essential peptidoglycan
precursor UDP-N-acetyl-glucosamine (35). In *B. subtilis, L. monocytogenes* and *L. lactis* a protein-
protein interaction between CdaA and GlmM has been detected using bacterial two-hybrid assays
performed in *Escherichia coli* (29,31,33). In *B. subtilis* this interaction has been further confirmed by *in*
vivo protein cross-linking and pulldown assays (29) and in *L. monocytogenes* by the co-elution of
purified proteins (33). The first evidence that that GlmM serves as negative regulator of CdaA/DacA
activity came from an *L. lactis* strain that produces a GlmM variant that is thought to form a stronger
interaction with CdaA; this strain produces lower cellular c-di-AMP levels than the bacteria expressing
wildtype GlmM (31). Furthermore, the activity of the soluble recombinant *S. aureus* DacA catalytic
domain (DacA$_{CD}$) could be blocked almost completely by the addition of purified GlmM protein in *in*
vitro assays and the recombinant proteins were shown to form a stable complex that could be purified
via size-exclusion chromatography (28). Additional mass-spectrometry and small-angle X-ray
scattering data (SAXS) analyses suggested that the complex is composed of a DacA$_{CD}$ dimer and a
GlmM dimer (28). Crystal structures of the individual *S. aureus* DacA$_{CD}$ and GlmM dimers revealed
that the *S. aureus* DacA$_{CD}$ protein assumed an "inactive" dimer conformation. GlmM had the typical
four-domain fold of phosphoglucomutases with a flexible C-terminal domain 4 and the dimer was "M-
shaped", characteristic for this class of enzymes (28). However, a high-resolution structure of the
complex could not be obtained and only a model for the complex could be proposed by fitting the
individual DacA$_{CD}$ and GlmM dimer structures into the SAXS envelope (28). Based on this, a model
was proposed in which GlmM could potentially block the activity of the DacA$_{CD}$ cyclase by preventing
the formation of higher oligomers.
128   Here, we set out to provide atomic resolution information on the CdaA:GlmM complex to gain
insight into the molecular mechanism how GlmM can control the activity of the c-di-AMP cyclase
enzyme. Using the purified *B. subtilis* CdaA catalytic domain (CdaA$_{CD}$) and purified full-length GlmM
or the truncated GlmM$_{F369}$ variant lacking the flexible C-terminal domain 4, we show that the two
proteins form a stable complex *in vitro* and that GlmM and GlmM$_{F369}$ are potent inhibits of the cyclase.
Crystal structures of the *B. subtilis* CdaA$_{CD}$ cyclase, the GlmM phosphoglucomutase and the
CdaA$_{CD}$:GlmM$_{F369}$ complex were obtained, revealing dimer conformations of the individual proteins as
well as a dimer of dimer conformation in the complex structure. More importantly, from the complex
structure the mechanism by which binding of GlmM inhibits the cyclase activity becomes apparent, that
is by preventing the oligomerisation of CdaA and formation of active head-to-head cyclase oligomers.
138
139

Results
The *B. subtilis* phosphoglucosamine GlmM interacts with and inhibits the activity of the c-di-AMP cyclase CdaA$_{CD}$

Using the purified *S. aureus* DacA$_{CD}$ catalytic domain and GlmM, it has been shown that the proteins form a stable complex *in vitro* and that GlmM is a potent inhibitor of the c-di-AMP cyclase without requiring any additional factors (28). To examine if this is also the case for the *B. sutbilis* proteins, the full-length *B. subtilis* GlmM protein as well as the truncated GlmM$_{F369}$ variant were expressed and purified along with the soluble catalytic domain of the *B. subtilis* c-di-AMP cyclase CdaA$_{CD}$. The GlmM$_{F369}$ variant lacks the flexible C-terminal domain 4 and was constructed to aid subsequent structural investigations. The proteins were expressed as His-tagged proteins in *E. coli* and purified individually via Ni-NTA affinity chromatography followed by size exclusion chromatography (Fig. 1). To test for a CdaA-GlmM interaction, lysates of strains producing CdaA$_{CD}$ and GlmM (Fig. 1A) or CdaA$_{CD}$ and GlmM$_{F369}$ (Fig. 1B) were mixed prior to affinity and size-exclusion chromatography. The elution profiles and analysis of the retention volumes revealed that CdaA$_{CD}$ formed a complex with GlmM and with GlmM$_{F369}$ that eluted as a single, higher-mobility species compared to the individual proteins (Fig. 1). The peak fractions of each complex were further analysed by SDS-PAGE, confirming the presence of both proteins (Fig. 1, inserts). To determine if the *B. subtilis* GlmM protein impacts the activity of CdaA$_{CD}$, *in vitro* cyclase activity assays were performed, and the conversion of ATP (spiked with a small amount of α-^{32}P-labelled ATP) into c-di-AMP assessed. The purified *B. subtilis* CdaA$_{CD}$ protein was enzymatically active in the presence of the divalent metal ion Mn^{2+} but showed only limited activity in the presence of Co^{2+} or Mg^{2+} (Fig. 2A) and after 4 h incubation approximately 50% of the ATP substrate was converted to c-di-AMP (Fig. 2B). Addition of GlmM or GlmM$_{F369}$ at a 2:1 molar ratio over CdaA$_{CD}$, led to a significant reduction in the conversion of ATP to c-di-AMP (Fig. 2C). Taken together, these data show that the purified *B. subtilis* CdaA$_{CD}$:GlmM as well as CdaA$_{CD}$:GlmM$_{F369}$ proteins form a stable complex *in vitro* and that both the full-length and truncated GlmM variant, inhibit the activity of the c-di-AMP cyclase CdaA$_{CD}$.

Crystal structures of the *B. subtilis* CdaA$_{CD}$ and GlmM proteins

To gain atomic level details of the CdaA$_{CD}$ and GlmM protein complex, we started off by determining the crystal structures of the individual proteins. The tag-less *B. subtilis* CdaA$_{CD}$ protein was crystallised and the structure solved at 2.8 Å (Table 1 and Fig. 3). The protein displayed the expected diadenylate cyclase protein fold, with a central β-sheet made up of 6 antiparallel strands flanked by 5 helices (Fig. 3A). However, it lacked the seventh β strand that was seen in the structures of CdaA homologs of other bacteria (21,28). In *B. subtilis* CdaA$_{CD}$ the residues corresponding to this β-strand are instead in a loop that adapts a very similar confirmation to the β strand observed in other CdaA structures. Superposition of the *B. subtilis* CdaA$_{CD}$ structure with the *L. monocytogenes* Δ100CdaA (PDB 4RV7) (21), and *S. aureus* DacA$_{CD}$ (PDB 6GYW) (28) structures, all lacking the N-terminal transmembrane helices, gave r.m.s.ds of 0.79 and 0.75, respectively, highlighting the overall structural similarities of these enzymes (Fig. 3B). The *B. subtilis* CdaA$_{CD}$ structure was solved as a dimer in the asymmetric unit with hydrogen-bonding interactions observed at the interaction interface (Fig 3C). Interactions were observed between the side chains of amino acid residues Asn166, Thr172 and Leu174 (site 1) and residues Leu150, Lys153 and Met155 (site 2) (Fig 3C). Similar hydrogen-bonding interactions were also identified in the *S. aureus* DacA$_{CD}$ and *L. monocytogenes* Δ100CdaA structures with amino acid residues in site 1 being absolutely conserved (28,30) (Fig. S1). Analysis of the interface with PDBePISA (36) indicated a buried surface of 1400 Å^2, which is similar to the value of 1460 Å^2 previously reported for the *S. aureus* DacA$_{CD}$ protein, indicative of a stable dimer formation. In this dimer confirmation, the active sites face opposite directions and hence cannot be engaged in a catalytically active head-to-head conformation (Fig. 3A). Taken together, these data indicate that the conformationally-inactive dimerization interface is conserved among different CdaA homologs in Gram-positive bacteria and that the enzyme needs to form higher oligomers for catalysis.

The His-tagged *B. subtilis* GlmM protein was crystallised and the structure solved by molecular replacement using the *B. anthracis* GlmM structure (PDB 3PDK; (37)) as the search model (Table 1 and Fig. 3D). The *B. subtilis* GlmM protein displayed a four-domain architecture typical for phosphoglucosamine mutase proteins (28,37) (Fig. 3D). Domains 1-3 are comprised of α-β mixed cores linked via a flexible loop to domain 4, which displays a 3-stranded β sheet fold surrounded by two α-

helices (Fig. 3D). While one GlmM molecule was present in the asymmetric unit, the typical "M-shaped" GlmM dimer arrangement was observed in the crystal cell packing (Fig. S2). Interactions were formed between domains 1, leading to the formation of a large groove at the top of the dimer molecule, mostly formed by domain 2 and the active site of each monomer subunit facing the opposite direction. Two different structures were solved for the *B. subtilis* GlmM protein at 2.9 and 3.0 Å resolutions with a superposition r.m.s.d. score of 0.29 (Table 1 and Fig. S2). One of the crystal structures was obtained with a divalent cation bound to the catalytic serine residue, which during catalysis is thought to be converted to a phosphoserine residue and the metal ion playing an important role during catalysis (Fig. S2). The exact type of metal ion could not be deduced due to the limitation of the structural resolution. However, we speculate that it is a magnesium ion, as magnesium was present in the crystallisation conditions and this metal ion is usually also bound in fully active enzymes. Furthermore, when a magnesium ion was modelled into the structure and analyzed using the program CheckMyMetal (38), a better fit was observed as compared to zinc or calcium ions, which could also fill the density. In the second structure, a phosphate molecule (PO_4) was bound to Arg419, located within a loop region in domain 4 (Fig S2B) at a similar location as observed in the *B. anthracis* GlmM structure (37). The superimposition of the *B. subtilis* GlmM structure with the *S. aureus* (6GYZ; (28)) and *B. anthracis* (3PDK; (37)) GlmM structures, gave small r.m.s.d. values of 1.0261 and 1.0668, respectively (Fig. 3E), indicating high similarity. However, the inter-residue distance between Arg419 in the phosphate binding site in domain 4 and the catalytic Ser100 in domain 1 was 20.22 Å in the phosphate bound *B. subtilis* GlmM structure compared to 18.4 Å in the *S. aureus* GlmM (PDB 6GYZ) or 15.18 Å in the *B. anthracis* GlmM (PDB 3PDK) structures (Fig. 3F). This highlights the flexibility of domain 4 in GlmM enzymes and also reveals that the *B. subtilis* GlmM protein was captured in most open state of the enzyme reported so far in a crystal structure.

Structure of the *B. subtilis* CdaA$_{CD}$:GlmM$_{F369}$ complex
To understand how GlmM interacts and inhibits the CdaA, we next aimed to obtain the structure of the complex. Any crystals obtained for the *B. subtilis* CdaA$_{CD}$:GlmM complex diffracted poorly. On the other hand, diffracting crystals were obtained for the CdaA$_{CD}$:GlmM$_{F369}$ complex, in which the GlmM protein lacks the flexible C-terminal domain 4. The crystals were obtained under two different conditions (see experimental procedures section), and the structure of the CdaA$_{CD}$:GlmM$_{F369}$ complex could be solved at 3.6 Å (Complex 1) and at 4.2 Å (Complex 2) by molecular replacement using the *B. subtilis* CdaA$_{CD}$ and GlmM (dimer) structures as search models (Table 1 and Fig. 4, Fig. S3). While obtained under two different conditions, complex 1 and complex 2 were nearly identical and overlapped with an r.m.s.d of 0.22 Å (Fig. S4A). Furthermore, in both complex structures, three complex molecules were obtained in the asymmetric unit and each complex was composed of a GlmM$_{F369}$ dimer and a CdaA$_{CD}$ dimer in the inactive dimer configuration (Fig. 4 and Fig. S3). The three complexes obtained in the asymmetric unit were almost identical to each other, as indicated by the superposition r.m.s.d. of 0.15 Å to 0.20 Å for complex 1 (Fig. S4B) and of 0.15 Å to 0.16 Å for complex 2 (Fig. S4C). Since the obtained complex structures were basically identical, all further descriptions are based on the higher resolution complex 1 structure. In the complex, a CdaA$_{CD}$ dimer was positioned in the large groove at the top of the GlmM$_{F369}$ dimer and formed interactions with domain 2 of GlmM (Fig 4A-4C). The complex was asymmetric, with one of the CdaA$_{CD}$ monomer, CdaA$_{CD}$(2) (shown in light blue in Fig. 4) placed in the center of the GlmM$_{F369}$ groove and the other monomer, CdaA$_{CD}$(1) (shown in dark blue in Fig. 4) projecting towards the solvent. Similarly, most of the interactions of the GlmM$_{F369}$ dimer with the CdaA$_{CD}$ dimer were made by GlmM$_{F369}$(1) (shown in dark pink in Fig 4). PDBePISA analysis revealed an average buried surface area of 996 Å^2 in the interface between GlmM$_{F369}$(1) and the CdaA$_{CD}$ dimer, which was stabilized by 4 hydrogen bond and 4 ionic bond interactions between GlmM$_{F369}$(1) and CdaA$_{CD}$(1) and 5 hydrogen bond and 3 ionic bond interactions with CdaA$_{CD}$(2) (Table S1 and Fig. S5). On the other hand, only an average 220.3 Å^2 surface area is occluded in GlmM$_{F369}$(2) (shown in light pink in Fig. 4). Based on the PDBePISA analysis, GlmM$_{F369}$(2) only formed two hydrogen bond interactions with the CdaA$_{CD}$(2) monomer but no interaction with CdaA$_{CD}$(1) (Table S1 and Fig. S5). A more detailed analysis of the interface showed that several interactions are made between two α-helices from domain 2 of GlmM$_{F369}$(1), α1 and α2, with the CdaA$_{CD}$(1) and CdaA$_{CD}$(2) monomers, respectively (Fig 4C and 4D). The main interactions in the complex were formed between three residues, D151, E154, and D194 of domain 2 in GlmM$_{F369}$(1) and residue R126 in each of the CdaA$_{CD}$ monomers. More

specifically, ionic bonds were formed between residue D195 in GlmM$_{F369}$(1) and residue R126 in
CdaA$_{CD}$(2). In addition, salt bridges were formed between residues D151 and E154 in GlmM$_{F369}$(1) and
residue R126 but this time from CdaA$_{CD}$(1) (Fig 4C and 4D, Table S2 and Fig. S5). The data suggest
that residue R126 in CdaA$_{CD}$ is potentially one of the most critical residues for complex formation, as it
contributes to a number of ionic as well as hydrogen-bond interactions and even though the complex is
asymmetric, it contributes to interactions in both CdaA$_{CD}$ monomers.
256
CdaA$_{CD}$ cannot form active oligomers in complex with GlmM
To gain insight how GlmM inhibits the activity of the c-di-AMP cyclase, we inspected the location of
the active sites of CdaA$_{CD}$ in the complex. The active site of DAC-domain enzymes is characterized by
DGA and RHR motifs, corresponding to residues D$_{171}$GA and R$_{203}$HR in *B. subtilis* CdaA (Fig. 5; areas
highlighted in yellow and green in the CdaA$_{CD}$ monomers). The active site in CdaA$_{CD}$(2) was completely
occluded upon interaction with the GlmM$_{F369}$ dimer (Fig. 5, dark blue CdaA$_{CD}$ monomer with active site
region highlighted in yellow) but the active site in CdaA$_{CD}$(1) appeared at least partially exposed (Fig.
5; light blue CdaA$_{CD}$ monomer with active site region highlighted in green). For CdaA$_{CD}$ to produce c-
di-AMP active head-to-head dimers need to be formed (3). The crystal structure of the *L. monocytogenes*
Δ100CdaA cyclase was recently determined with a c-di-AMP molecule bound in the catalytic site and
an active head-to-head dimer conformation seen in the crystal packing (30). Using the *L. monocytogenes*
Δ100CdaA structure (PDB 6HVL) as model, an active *B. subtilis* CdaA$_{CD}$ dimer was modelled and
superimposed on CdaA$_{CD}$(1) in the complex structure (Fig. 5). Although the active site of the CdaA$_{CD}$(1)
was exposed and accessible in the complex with GlmM$_{F369}$, in an active dimer conformation, parts of
the second CdaA$_{CD}$ molecule would collide and overlap with GlmM$_{F369}$, highlighting that also
CdaA$_{CD}$(1) cannot form active head-to-head oligomers in the complex (Fig. 5). Taken together, these
data indicate that in the complex, the interaction of GlmM with CdaA$_{CD}$ will prevent the formation of
functional diadenylate cyclase enzyme oligomers, which is essential for the formation of c-di-AMP. The
crystal structure of the CdaA$_{CD}$:GlmM$_{F369}$ complex therefore provides insight on an atomic level on the
catalytic inhibition of the diadenylate cyclase CdaA$_{CD}$ by the phosphoglucosamine enzyme GlmM.
277
Small angle X-ray scattering analysis of *B. subtilis* CdaA$_{CD}$:GlmM complex
To determine whether the full-length *B. subtilis* GlmM protein interacts with CdaA$_{CD}$ in a similar
manner as observed for GlmM$_{F369}$, a structural characterisation of the CdaA$_{CD}$:GlmM complex was
performed via small-angle X-ray scattering (SAXS). To this end, the individual purified *B. subtilis*
CdaA$_{CD}$ and GlmM proteins as well as the purified CdaA$_{CD}$:GlmM complex were passed over an
analytical size-exclusion column, followed by continuous automated SAXS data collection throughout
the run (Fig. 6, Fig S6 and Table S2). For CdaA$_{CD}$ and GlmM, the reconstructed maps were consistent
with the proteins forming dimers and the maps were a good fit for the *B. subtilis* CdaA$_{CD}$ dimer (Fig.
6A) and GlmM dimer (Fig. 6B) structures, respectively. The map reconstructed for the CdaA$_{CD}$:GlmM
complex was bigger in volume and dimensions (V$_c$: 890.1, R$_g$: 44.65 Å and d$_{max}$: 161 Å) as compared
to the individual maps calculated for GlmM (V$_c$: 625.8; R$_g$: 37.29 Å, d$_{max}$: 122 Å) and CdaA$_{CD}$ (V$_c$:
373.2, R$_g$: 26.94 Å, d$_{max}$: 88 Å), which is consistent with the formation of a complex. From the Guinier
plot analysis, the molecular weight of the *B. subtilis* CdaA$_{CD}$:GlmM complex was calculated to be 130
kDa, which is consistent with the theoretical molecular weight of 144.46 kDa for a complex made of
two CdaA$_{CD}$ and two GlmM molecules. To fit a CdaA$_{CD}$:GlmM dimer complex into the reconstructed
map, a model of the complex with full-length GlmM was first produced, by superimposing the crystal
structure of full-length GlmM onto the CdaA$_{CD}$:GlmM$_{F369}$ complex structure. The resulting complex
model was subsequently fitted in the reconstituted SAXS envelope of the complex. A good fit of the
CdaA$_{CD}$:GlmM dimer model complex into the reconstructed envelope was observed, however an
elongated density on one side remained unoccupied (Fig. 6C). It is plausible that the flexible C-terminal
domain 4 of the GlmM protein is responsible for this extra density. These data suggest that the full-
length GlmM protein likely forms a dimer-of-dimer complex with the c-di-AMP cyclase CdaA$_{CD}$ and
could assume a similar arrangement as observed for the CdaA$_{CD}$:GlmM$_{F369}$ complex.
301
302

Arginine 126 in *B. subtilis* CdaA_CD is essential for complex formation

The complex structure highlighted key interactions between residues D194 and residues D151/E154 in GlmM with residue R126 in each of the CdaA_CD monomers (Fig. 5D). To confirm our structural findings, a site-directed mutagenesis analysis was performed. To this end, D195A, D151A/E154A and D151A/E154A/D191A alanine substitution GlmM variants were created. Furthermore, residue R126 in CdaA_CD, which in both monomers makes contacts with GlmM, was mutated to an alanine. The different alanine substitution variants were expressed and purified from *E. coli* and complex formation assessed by size-exclusion chromatography. While our initial experiments using the GlmM single, double and triple alanine substitution variants appeared not or only marginally to affect complex formation with CdaA_CD (Fig. S7), no complex-specific peak was observed when the interaction between the CdaA_CD-R126A variant and GlmM was assessed. Instead, two peaks were observed for the CdaA_CD-R126A:GlmM sample, one corresponding to the retention volume of GlmM and the another to CdaA_CD-R126A (Fig. 7A). Analysis of the elution fractions from the CdaA_CD-R126A/GlmM sample by SDS-PAGE and Coomassie staining showed that only a very small fraction of the CdaA_CD-R126A protein co-eluted with GlmM (Fig. 7A). These data highlight that, consistent with the structural data, residue R126 in CdaA_CD plays a key role for the complex formation with GlmM. Based on these data, it can be predicted that the cyclase activity of the CdaA_CD-R126A variant should now longer be inhibited by GlmM. To test this experimentally, *in vitro* cyclase enzyme activity assays were performed. The CdaA_CD-R126A variant was active, although the activity was reduced as compared to wild-type CdaA_CD (Fig. 7B). Importantly and in contrast to wild-type CdaA_CD, the enzyme activity of this variant was no longer inhibited by the addition of GlmM (Fig. 7B). These data show that residue Arg126 in *B. subtilis* CdaA_CD plays a critical role for complex formation and that GlmM can only inhibit the activity of the c-di-AMP cyclase after the formation of a stable complex.

Discussion

In this study, we show that the *B. subtilis* GlmM and CdaA_CD cyclase domain form a stable dimer-of-dimer complex. GlmM acts through this protein-protein interaction as a potent inhibitor of the c-di-AMP cyclase without requiring any additional factors. Based on the atomic-resolution complex structure data, we suggest that GlmM inhibits the activity of CdaA_CD by preventing the formation of active head-to-head cyclase oligomers.

For CdaA to produce c-di-AMP, two monomers need to be arranged in an active head-to-head conformation. As part of this study, we determined the structure of the *B. subtilis* CdaA_CD protein and show that it has the typical DAC domain fold. While the protein was also found as a dimer in the structure, the dimer was in an inactive conformation, with the two active sites facing in opposite directions. The interface creating the inactive dimer conformation is conserved among CdaA proteins. The *L. monocytogenes* and *S. aureus* homologs, for which structures are available, were found in the same inactive dimer conformation even though the proteins crystallized under different conditions and were found in different space groups (21,28,30). This makes it less likely that a crystallographic symmetry artefact is responsible for the observed inactive dimer configuration. In addition to the inactive dimer configuration within an asymmetric unit, an active dimer conformation was observed in the *L. monocytogenes* Δ100CdaA protein by inspecting adjacent symmetry units (30). However, no such active dimer head-to-head conformations were identified for the *B. subtilis* CdaA_CD protein across different symmetry units in the current structure. While not further investigated as part of this study, previous work on the *S. aureus* homolog indicated that the inactive dimer conformation is very stable, and in order for the protein to produce an active enzyme, the protein needs to form higher-level oligomers (28). Given the similarity in the interaction interface, this is likely also the case for the *B. subtilis* CdaA_CD enzyme and we would suggest that the *B. subtilis* CdaA_CD dimer observed in the structure is unlikely to rearrange into an active dimer conformation.

We also solved the structure of the *B. subtilis* GlmM enzyme. The protein assumed the typical 4-domain architecture previously reported for GlmM enzymes (28,37) and the "M shape" in the dimer conformation, which in the case of the *B. sutbtilis* GlmM protein was formed across two adjacent crystallographic units. The *B. subtilis* GlmM structure further highlighted the flexibility of the most C-terminal domain 4, which was found in the most open conformation seen in any GlmM protein structure up to date. The conformational flexibility of domain 4 is probably also a main factor why we were unsuccessful in determining the structure of a complex between CdaA_CD and full-length GlmM.

However, GlmM domain 4 is not required for the interaction with and inhibition of CdaA$_{CD}$, since a *B.*
subtilis GlmM variant lacking domain 4 formed a complex and inhibited the activity of *B. subtilis*
CdaA$_{CD}$. Futhermore, by using the *B. subtilis* GlmMF$_{F369}$ variant lacking the flexible domain 4, we were
able to obtain the structure of the CdaA$_{CD}$:GlmM$_{F369}$ complex, revealing for the first time structural
details at the atomic level for this complex, thereby identifying the amino acids important for the
interaction between the two proteins. Several electrostatic interactions were detected between
CdaA$_{CD}$:GlmM$_{F369}$: between the negatively charged residues D151, E154 and D195 in domain 2 of *B.*
subtilis GlmM$_{F369}$ with the positively charged residue R126 in CdaA$_{CD}$. We could further show that
replacing residue R126 in CdaA$_{CD}$ with an alanine abolished complex formation and the activity of the
CdaA$_{CD}$-R126 variant was no longer inhibited by GlmM. A direct protein-protein interaction between
CdaA and GlmM has now been reported for these proteins in several Firmicutes bacteria, and hence the
amino acids required for the interaction might be conserved. Indeed, a ConSurf (39) analysis using 250
CdaA protein sequences, showed that the residue corresponding to R126 in *B. subtilis* CdaA is conserved
between the different homologs (Fig. S8). Likewise, all the three negatively charged residues, D151,
E154 and D195 in GlmM, which mediate the primary electrostatic interactions with R126 of CdaA$_{CD}$,
are highly conserved (Fig. S8). In previous work, we have shown, that the *S. aureus* DacA$_{CD}$ (the CdaA$_{CD}$
homolog) does not interact with GlmM proteins from *E. coli* and *Pseudomonas aeruginosa,* two Gram-
negative bacteria (28). While negatively charged amino acids corresponding to residues E154 and D195
in *B. subtilis* GlmM are also found in the GlmM protein from the Gram-negative bacteria (D153 and
E194 in *E. coli* and D152 and E193 in *P. aeruginosa*), the amino acids at the equivalent position of
D151 in *B. subtilis* GlmM is an arginine residue (R150 in *E. coli* and R149 in *P. aeruginosa*), which
may hinder the complex formation between CdaA (DacA), and GlmM proteins of Gram-negative
bacteria (28).
In a previous work, we proposed a model whereby GlmM inhibits the activity of CdaA cyclase
activity by preventing the formation of active head-to-head oligomers (28). In the absence of an actual
atomic resolution structure of the complex, this model was based on SAXS envelope data and fitting
individual protein structures. The model predicted that the likely interaction site between CdaA and
GlmM proteins is domain 2 of GlmM (28). The structure of *B. subtilis* CdaA$_{CD}$:GlmM$_{F369}$ complex we
present here now provides experimental evidence for such a model and shows that GlmM indeed inhibits
the activity of the CdaA cyclase *in vitro* by preventing the formation of active head-to-head oligomers.
The level of c-di-AMP is regulated by a fine balance between the activities of the cyclase, which
synthesizes c-di-AMP, and the phosphodiesterases, which break it down. Interestingly, these two classes
of enzyme appear to be regulated very differently; whereas the activity of several phosphodiesterases
has been shown to be regulated by small molecules, cyclase activity appears to be regulated through
protein-protein interaction. For example, the stringent response alarmone (p)ppGpp has been shown to
inhibit the activity both GdpP and PgpH enzymes (11,20). Furthermore, binding of heme to the Per-
ARNT-Sim (40) signalling domain in GdpP (which is separate from its DHH/DHHA1 enzymatic
domain that is responsible for the degradation of c-di-AMP) was shown in *in vitro* enzyme assays to
lead to reduced phosphodiesterase activity (40). Interestingly, the ferrous form of heme bound to GdpP
could form a pentacoordinate complex with nitric oxide (NO), resulting in increased c-di-AMP
phosphodiesterase activity. Based on these data it has been suggested that GdpP is a heme or NO sensor,
resulting in decreased or increased activity respectively (41). The function consequence and impact of
(p)ppGpp, heme or NO binding to the phosphodiesterases on bacterial physiology has not yet been fully
investigated. However, from these data it is clear that the activity of the c-di-AMP phosphodiesterases
can be regulated by small molecule ligands.
On the other hand, several proteins have been found to interact with and regulate the activity of
c-di-AMP cyclases. The *B. subtilis* DisA protein is involved in monitoring the genomic stability
ensuring that damaged DNA is repaired before cells progress with the sporulation process or exit from
spores (27,42). DisA is encoded in a multi-gene operon and the gene immediately upstream of *disA*
codes for RadA (also referred to as SMS). *B. subtilis* RadA possesses 5' to 3' DNA helicase activity,
contributes to DNA repair and DNA transformation processes in *B. subtilis* and has been shown to
interact and negatively impact the activity of DisA (27). However, the mechanistic basis of how RadA
binding to DisA inhibits the cyclase is currently not known. There is now ample evidence that the
activity of the "house-keeping" membrane-linked c-di-AMP cyclase CdaA is impacted by two
interacting proteins, the membrane-linked regulator protein CdaR and the cytoplasmic

phosphoglucomutase enzyme GlmM (29,33). We have shown provided experimental evidence for the
mechanistic basis by which GlmM inhibits the activity of CdaA, that is by preventing the formation of
active higher-level oligomers. How CdaR regulates the cyclase activity of CdaA remains unclear.
Recent work on the homologous proteins in *L. monocytogenes* indicated that the interaction of CdaA
with CdaR takes place via the N-terminal transmembrane region of CdaA, GlmM has been shown to
interact directly with the cytoplasmic cyclase domain of CdaA in *S. aureus* (28,33). Here, we show that
this is also the case for the *B. subtilis* GlmM protein, which can bind without the requirement of any
additional factor to the catalytic CdaA$_{CD}$ domain. In future works, it will be interesting to determine the
structure of the full-length CdaA enzyme, which might provide further insight into how the enzyme
forms higher oligomers for activity as well as how it interacts with CdaR. Furthermore, it will be
interesting to further investigate the interaction between GlmM and CdaR with CdaA within bacterial
cells to determine if this interaction is dynamic and which stimuli will promote or prevent complex
formation to fine-tune the synthesis of c-di-AMP. Identifying how interacting proteins regulate the
activity of these important cyclases, will provide important insight how bacterial cells maintain proper
levels of c-di-AMP under different growth conditions and in different environments.
428
Experimental procedures
Bacterial strains and plasmid construction
All bacterial strains and primers used in this work are listed in Tables S3 and S4, respectively.
pET28b-derived plasmids were constructed for the overproduction of the C-terminal catalytic domain
of the *Bacillus subtilis* CdaA enzyme starting from amino acid Phe97 and referred to as CdaA$_{CD}$, GlmM
and the GlmM$_{F369}$ variant comprising residues Met1 to Phe369 but lacking the C-terminal domain 4. To
this end, the corresponding DNA fragments were amplified by PCR using *B. subtilis* strain 168
chromosomal DNA as template and primer pairs ANG2760/ANG2761 (*cdaA$_{CD}$*), ANG2762/ANG2763
(*glmM*) and ANG2762/ANG2764 (*glmM$_{F369}$*). The PCR products were purified, digested with
NheI/BamHI (*cdaA$_{CD}$*) or NcoI/XhoI (*glmM* and *glmM$_{F369}$*) and ligated with pET28b, which had been
cut with the same enzyme. CdaA$_{CD}$ was cloned in frame with an N-terminal thrombin cleavable 6-
histidine tag, while GlmM and GlmM$_{F369}$ were cloned in frame with a C-terminal 6-histidine tag and a
thrombin cleavage site was introduced in front of the His-tag as part of the primer sequence. The
resulting plasmids pET28b-*his-cdaA$_{CD}$*, pET28b-*glmM-his* and pET28b-*glmM$_{F369}$-his* were initially
recovered in *E. coli* XL1-Blue, yielding the strains ANG4583, ANG4584 and ANG4585 and
subsequently transformed for protein expression into strain *E. coli* BL21(DE3), yielding strains
ANG4597, ANG4598 and ANG4599, respectively. Plasmids pET28b-*his-cdaA$_{CD}$-R126*, pET28b-
glmM-D194A-his, pET28b-*glmM-D151A/E154A-his* were constructed for the expression of CdaA$_{CD}$
and GlmM alanine substitution variants. The plasmids were constructed by QuikChange mutagenesis
using pET28b-*his-cdaA$_{CD}$* and primer pair ANG3373/ANG3374 or plasmid pET28b-*glmM-his* and
primer pairs ANG3381/ANG3382 and ANG3383/3384, respectively. The plasmids were initially
recovered in *E. coli* XL1-Blue, yielding strains ANG5933, ANG5937, ANG5938 and subsequently
introduced for protein expression into *E. coli* strain BL21(DE3) yielding strains ANG5940, ANG5944,
ANG5945. In addition, plasmid pET28b-*glmM-D151A/E154A/D194A-his* for expression of a GlmM
variant with a triple Asp151 (D151), *Glu154 (E154) and Asp195 (D195)* alanine substitution variant
was constructed by QuikChange mutagenesis using plasmid pET28b-*glmM-D194A-his* as template and
primer pair ANG3383/3384 to introduce the D151A and E154A mutations. Plasmid pET28b-*glmM-*
D151A/E154A/D194A-his was recovered in *E. coli* strain XL1-Blue, yielding strain ANG5939 and
subsequently introduced for protein expression into strain BL21(DE3) yielding strain ANG4946. The
sequences of all plasmid inserts were verified by fluorescent automated sequencing at Eurofins.
459
Protein expression, purification and quantification
Proteins CdaA$_{CD}$, GlmM and GlmM$_{F369}$ were expressed and purified from 1 L cultures as previously
described (28). Briefly, when bacterial cultures reached an OD$_{600}$ of approximately 0.6, protein
expression was induced with 1 mM IPTG (final concentration) for 3 hours at 37°C. Cells were harvested
by centrifugation, suspended in 20 mL of 50 mM Tris pH 7.5, 500 mM NaCl buffered and lysed using
a French Press system. Lysates were clarified by centrifugation and the supernatant loaded onto a gravity
flow column with 3 mL of Ni-NTA resin. Immobilized proteins were washed with 20 mL of 50 mM
Tris pH 7.5, 500 mM NaCl, 50 mM imidazole buffer and eluted in 5 x 1 ml fractions using 50 mM Tris

pH 7.5, 200 mM NaCl, 500 mM imidazole buffer. Fractions containing the proteins were pooled and
loaded onto a preparative Superdex 200 HiLoad 16/60 column equilibrated with 1 column volume of 30
mM Tris pH 7.5, 150 mM NaCl buffer. When appropriate, the purified proteins were concentrated using
10 mL 10 kDa cutoff Amicon concentrators for downstream applications. For the purification of the
$CdaA_{CD}$:GlmM and $CdaA_{CD}$:$GlmM_{F369}$ complexes or complexes of $CdaA_{CD}$ and GlmM alanine-
substitution variants, cell lysates of strains overproducing the respective proteins were mixed after the
French press step, then the same protein purification procedure steps as described above used for the
purification of individual proteins were performed. Protein concentrations were determined using the
BCA assay kit (Pierce™ BCA Protein Assay Kit). For each sample, the readings were taken in triplicates
and then averaged to obtain the protein concentration. Purified proteins were also separated on 12%
SDS PAGE gels and detected by Coomassie staining.
479
Protein crystallisation, data processing and analysis
For crystallisation, the histidine tag was removed from the purified *B. subtilis* $CdaA_{CD}$ protein. This was
done by incubating 10 mg purified protein with 20 U thrombin overnight at 4˚C with agitation. The
following day, the tag less $CdaA_{CD}$ was purified by size-exclusion chromatography as described above.
The $CdaA_{CD}$ protein was crystallized at a concentration of 4 mg/ml in 0.1M sodium cacodylate pH 6.5,
0.1M ammonium sulfate, 0.3 M sodium formate, 6% PEG 8000, 3% γ-PGA via the vapour diffusion
method. The crystal screens for *B. subtilis* GlmM (including the His tag) were set up at a concentration
of 10 mg/ml and protein crystals were obtained in two different conditions. The structure with bound
PO_4 ($GlmM:PO_4$) was obtained in the Morpheus screen containing 0.1 M buffer system 1 (Imidazole;
MES, pH 6.5), 0.09 M NPS ($NaNO3$; Na_2HPO_4; $(NH_4)_2SO_4$) and 37.5% MPD_P1K_P3350 (75% MPD,
PEG 1K, PEG 3350) and the divalent-ion bound crystal structure (GlmM:metal ion) was obtained in
0.05 M Magnesium chloride hexahydrate, 0.1 M HEPES pH 7.5, 30% v/v Polyethylene glycol
monomethyl ether 550 buffer. The crystals for the $CdaA_{CD}$:$GlmM_{F369}$ complex were set up at a protein
concentration of 10 mg/ml and crystals were obtained in 0.12 M alcohols, 0.1 M buffer system
(Imidazole; MES, pH 6.5) and 30% GOL_P4K (60% glycerol, PEG 4K) (Complex 1) and 0.1 M
carboxylic acids, 0.1 M buffer system 1 (Imidazole; MES, pH 6.5) and 30% GOL_P4K (60% glycerol,
PEG 4K) (Complex 2). The crystals were fished and stored in liquid nitrogen to test for diffraction at
the I03 beamline at the Diamond Light Source (Harwell Campus, Didcot, UK). Data were reduced with
DIALS (43) and scaled and merged with AIMLESS (44). The structures of $CdaA_{CD}$ and GlmM were
solved by the molecular replacement method using the program MR-PHASER (45) in the Phenix suite
(46), using the *L. monocytogenes* CdaA structure (PDB 4RV7; (21)) and *B. anthracis* GlmM structure
(PDB 3PDK; (37)) as the search models, respectively. To solve the phase problem for the structure of
the $CdaA_{CD}$:$GlmM_{F369}$ complex, dimers of *B. subtilis* $CdaA_{CD}$ and GlmM (each) were used as the search
models using the MR-PHASER program in Phenix. The models were manually built using COOT (47)
followed by iterative cycles of structure refinement using the Phenix-Refine program (48). The final
refined structures were analysed using the PDBePISA server (36) to identify buried interface areas for
each protein. To search for conserved residues among the phylogenetically related homologs, a protein
BLAST search was performed using *B. subtilis* $CdaA_{CD}$ and GlmM amino acid sequences as query
sequences and a multiple sequence alignment (MSA) of the top 2500 homologs found in Firmicutes was
prepared. The MSA was then used to identify conserved residues among the homologs using the
ConSurf server (49).
511
$CdaA_{CD}$ activity and inhibition assays
To assess the activity of the *B. subtilis* $CdaA_{CD}$ enzymes, 20 μl enzyme reactions were set up in 100 mM
NaCl, 40 mM HEPES pH 7 buffer containing 10 mM $MnCl_2$ (or 10 mM $MgCl_2$ or 10 mM $CoCl_2$ for
metal dependent assays), 100 mM ATP spiked with a-P^{32}-labelled ATP (Perkin Elmer; using 0.4 μl of
a 3.3 μM, 250 μCi solution per 20 μl reaction) and 5 μM $CdaA_{CD}$ enzyme. The mixture was incubated
at 37˚C for 4 hours, followed by heat inactivation at 95˚C for 5 minutes. After centrifugation for 10
minutes at 21,000 x g, 1 μl of the mixture was deposited onto a polyethylenimine-modified cellulose
TLC plate (Millipore) and nucleotides separated by running the plate for 20 minutes using a 3.52 M
$(NH_4)_2SO_4$ and 1.5 M KH_2PO_4 buffer system mixed at a 1:1.5 v/v ratio. Radioactive signals for ATP and
the c-di-AMP reaction product were detected using a Typhoon FLA-700 phosphor imager. The bands
were quantified using the ImageQuant program and the obtained values used to calculate the percent

conversion of ATP to c-di-AMP. For the time course experiment, a 100 μl reaction mixture was prepared
as described above and incubated at 37˚C. Ten μl aliquots were removed at the indicated time points
and the enzyme reactions stopped by incubation the removed aliquots at 95˚C for 5 minutes. To assess
the activity of $CdaA_{CD}$ in the presence of GlmM or $GlmM_{F369}$, the full length GlmM protein or C-
terminally truncated GlmM variant were added to the reaction mixture at a 1:2 ($CdaA_{CD}$: GlmM or
$CdaA_{CD}$: $GlmM_{F369}$) molar ratio and the reactions incubated at 37˚C for 4 hours, stopped and analysed
as described above. The enzyme activity assays were performed in triplicates with two independently
purified protein preparations.
531
SAXS sample preparation and analysis
For the SAXS analysis, purified $CdaA_{CD}$, GlmM and $CdaA_{CD}$:GlmM complex protein samples where
purified a second time by size exclusion chromatography as described above and subsequently
concentrated to 5 mg/ml for $CdaA_{CD}$ and 24 mg/ml each for GlmM and the $CdaA_{CD}$:GlmM complex.
Next, 50 μl protein samples were loaded on a high pressure Shodex column (KW403: range 10 kDa to
700 kDa) fitted to an Agilent 1200 HPLC system at the B21 beamline at the Diamond Light Source
(Didcot, UK). The size-exclusion column was equilibrated with 30 mM Tris pH 7.5, 150 mM NaCl
buffer prior to loading the protein sample and the data were collected continuously throughout the
protein elution. The analysis of the datasets was done via ScÅtter (50) using the scattering frames
corresponding to the elution peaks. The ab-initio analysis of the SAXS data to reconstruct a low-
resolution shape of the model was done using DAMMIF (51). The program performed 13 *ab-initio* runs
to generate models from each run that were averaged to determine the most persistent three-dimensional
shape of the protein. The cross-correlation Normalised Spatial Discrepancy (NSD) values were
calculated using SUPCOMB from each of the 13 generated models. The mean NSD values calculated
for each of the protein were: 0.591 ± 0.088 ($CdaA_{CD}$), 0.598 ± 0.014 (GlmM) and 0.661 ± 0.064
($DacA_{CD}$:GlmM Complex). Chimera (52) was used to visualise the reconstructed SAXS maps. The
crystal structures of $CdaA_{CD}$ and GlmM dimers were fitted in the respective SAXS envelopes in Chimera
using the one-step fit function. For the $CdaA_{CD}$: GlmM complex data, a structural model of the $CdaA_{CD}$
full-length GlmM complex was generated based on the crystal structure of $CdaA_{CD}$: $GlmM_{F369}$ complex,
which was then fitted into the SAXS envelope data using Chimera.
552
Data availability
The structure coordinates of the *B. subtilis* proteins were deposited in the Protein Database
(https://www.rcsb.org), under PDB codes 6HUW ($CdaA_{CD}$), 7OJR (GlmM: PO_4 bound), 7OML
(GlmM: metal bound) and 7OLH ($CdaA_{CD}$:$GlmM_{F369}$ Complex 1) and 7OJS ($CdaA_{CD}$:$GlmM_{F369}$
Complex 2). SAXS models were deposited in the SASBDB database, under the accession codes
SASDL25 (Complex $CdaA_{CD}$:GlmM), SASDLZ4 (GlmM) and SASDLY4 ($CdaA_{CD}$).
559
560 **Supporting Information**
561 This article contains a supporting information file with the following additional references
562 (36,49,50,53,54).
563

564 **Acknowledgments.** We would like to thank Christiaan van Ooij for helpful comments on the
565 manuscript.
566
Author contribution statement
Monisha Pathania: Conceptualization, Investigation, Data analysis, Visualization, Writing – original
draft preparation. **Tommaso Tosi:** Conceptualization, Investigation, Data analysis, Writing – review &
editing. **Charlotte Millsership:** Investigation, Data analysis, Writing – review & editing. **Fumiya
Hoshiga:** Investigation, Data analysis, Writing – review & editing. **Rhodri M. L. Morgan:** Data
analysis, Supervision, Writing – review & editing. **Paul S. Freemont:** Conceptualization, Funding
acquisition, Supervision, Data analysis, Writing – review & editing. **Angelika Gründling:**
Conceptualization, Funding acquisition, Data analysis, Supervision, Writing – original draft preparation.

575
576

Funding and additional information
This work was funded by the MRC grant MR/P011071/1 to AG and PSF. The crystallization facility at Imperial College was funded by the BBSRC (BB/D524840/1) and the Wellcome Trust (202926/Z/16/Z). X-Ray datasets for CdaA$_{CD}$, GlmM and the CdaA$_{CD}$:GlmM complex 2 were collected at the I03 beamline and the CdaA$_{CD}$:GlmM complex 1 dataset was collected at the I04 beamline at the Diamond Light Source (Didcot, UK). The SAXS data were collected at the B21 beamline at the Diamond Light Source (Didcot, UK).

Conflict of Interest
The authors declare no conflicts of interest in regard to this manuscript.

References

1. Hengge, R., Gründling, A., Jenal, U., Ryan, R., and Yildiz, F. (2016) Bacterial Signal Transduction by Cyclic Di-GMP and Other Nucleotide Second Messengers. *J Bacteriol* **198**, 15-26

2. Krasteva, P. V., and Sondermann, H. (2017) Versatile modes of cellular regulation via cyclic dinucleotides. *Nat Chem Biol* **13**, 350-359

3. Witte, G., Hartung, S., Buttner, K., and Hopfner, K. P. (2008) Structural biochemistry of a bacterial checkpoint protein reveals diadenylate cyclase activity regulated by DNA recombination intermediates. *Mol Cell* **30**, 167-178

4. Bai, Y., Yang, J., Zarrella, T. M., Zhang, Y., Metzger, D. W., and Bai, G. (2014) Cyclic di-AMP impairs potassium uptake mediated by a cyclic di-AMP binding protein in *Streptococcus pneumoniae*. *J Bacteriol* **196**, 614-623

5. Whiteley, A. T., Garelis, N. E., Peterson, B. N., Choi, P. H., Tong, L., Woodward, J. J., and Portnoy, D. A. (2017) c-di-AMP modulates *Listeria monocytogenes* central metabolism to regulate growth, antibiotic resistance and osmoregulation. *Mol Microbiol* **104**, 212-233

6. Huynh, T. N., Choi, P. H., Sureka, K., Ledvina, H. E., Campillo, J., Tong, L., and Woodward, J. J. (2016) Cyclic di-AMP targets the cystathionine beta-synthase domain of the osmolyte transporter OpuC. *Mol Microbiol* **102**, 233-243

7. Corrigan, R. M., Campeotto, I., Jeganathan, T., Roelofs, K. G., Lee, V. T., and Gründling, A. (2013) Systematic identification of conserved bacterial c-di-AMP receptor proteins. *Proc Natl Acad Sci U S A* **110**, 9084-9089

8. Gundlach, J., Herzberg, C., Kaever, V., Gunka, K., Hoffmann, T., Weiss, M., Gibhardt, J., Thurmer, A., Hertel, D., Daniel, R., Bremer, E., Commichau, F. M., and Stülke, J. (2017) Control of potassium homeostasis is an essential function of the second messenger cyclic di-AMP in *Bacillus subtilis*. *Sci Signal* **10**

9. Corrigan, R. M., Abbott, J. C., Burhenne, H., Kaever, V., and Gründling, A. (2011) c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. *PLoS Pathog* **7**, e1002217

10. Luo, Y., and Helmann, J. D. (2012) Analysis of the role of *Bacillus subtilis* sigma(M) in beta-lactam resistance reveals an essential role for c-di-AMP in peptidoglycan homeostasis. *Mol Microbiol* **83**, 623-639

11. Corrigan, R. M., Bowman, L., Willis, A. R., Kaever, V., and Gründling, A. (2015) Cross-talk between two nucleotide-signaling pathways in *Staphylococcus aureus*. *J Biol Chem* **290**, 5826-5839

12. Dengler, V., McCallum, N., Kiefer, P., Christen, P., Patrignani, A., Vorholt, J. A., Berger-Bächi, B., and Senn, M. M. (2013) Mutation in the c-di-AMP cyclase *dacA* affects fitness and resistance of methicillin resistant *Staphylococcus aureus*. *PLoS One* **8**, e73512

13. Woodward, J. J., Iavarone, A. T., and Portnoy, D. A. (2010) c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science* **328**, 1703-1705

14. Witte, C. E., Whiteley, A. T., Burke, T. P., Sauer, J. D., Portnoy, D. A., and Woodward, J. J. (2013) Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall homeostasis, and establishment of infection. *mBio* **4**, e00282-00213

15. Oppenheimer-Shaanan, Y., Wexselblatt, E., Katzhendler, J., Yavin, E., and Ben-Yehuda, S. (2011) c-di-AMP reports DNA integrity during sporulation in *Bacillus subtilis*. *EMBO Rep* **12**, 594-601

16. Devaux, L., Sleiman, D., Mazzuoli, M. V., Gominet, M., Lanotte, P., Trieu-Cuot, P., Kaminski, P. A., and Firon, A. (2018) Cyclic di-AMP regulation of osmotic homeostasis is essential in Group B *Streptococcus*. *PLoS Genet* **14**, e1007342

17. Mehne, F. M., Gunka, K., Eilers, H., Herzberg, C., Kaever, V., and Stülke, J. (2013) Cyclic di-AMP homeostasis in *Bacillus subtilis*: both lack and high level accumulation of the nucleotide are detrimental for cell growth. *J Biol Chem* **288**, 2004-2017

18. Commichau, F. M., Heidemann, J. L., Ficner, R., and Stülke, J. (2019) Making and Breaking of an Essential Poison: the Cyclases and Phosphodiesterases That Produce and Degrade the Essential Second Messenger Cyclic di-AMP in Bacteria. *J Bacteriol* **201**

19. Rao, F., See, R. Y., Zhang, D., Toh, D. C., Ji, Q., and Liang, Z. X. (2010) YybT is a signaling
protein that contains a cyclic dinucleotide phosphodiesterase domain and a GGDEF domain
with ATPase activity. *J Biol Chem* **285**, 473-482
20. Huynh, T. N., Luo, S., Pensinger, D., Sauer, J. D., Tong, L., and Woodward, J. J. (2015) An
HD-domain phosphodiesterase mediates cooperative hydrolysis of c-di-AMP to affect bacterial
growth and virulence. *Proc Natl Acad Sci U S A* **112**, E747-756
21. Rosenberg, J., Dickmanns, A., Neumann, P., Gunka, K., Arens, J., Kaever, V., Stülke, J., Ficner,
R., and Commichau, F. M. (2015) Structural and biochemical analysis of the essential
diadenylate cyclase CdaA from *Listeria monocytogenes*. *J Biol Chem* **290**, 6596-6606
22. Zheng, C., Ma, Y., Wang, X., Xie, Y., Ali, M. K., and He, J. (2015) Functional analysis of the
sporulation-specific diadenylate cyclase CdaS in *Bacillus thuringiensis*. *Front Microbiol* **6**, 908
23. Müller, M., Deimling, T., Hopfner, K. P., and Witte, G. (2015) Structural analysis of the
diadenylate cyclase reaction of DNA-integrity scanning protein A (DisA) and its inhibition by
3'-dATP. *Biochem J* **469**, 367-374
24. Mehne, F. M., Schroder-Tittmann, K., Eijlander, R. T., Herzberg, C., Hewitt, L., Kaever, V.,
Lewis, R. J., Kuipers, O. P., Tittmann, K., and Stülke, J. (2014) Control of the diadenylate
cyclase CdaS in *Bacillus subtilis*: an autoinhibitory domain limits cyclic di-AMP production. *J
Biol Chem* **289**, 21098-21107
25. Fahmi, T., Port, G. C., and Cho, K. H. (2017) c-di-AMP: An Essential Molecule in the Signaling
Pathways that Regulate the Viability and Virulence of Gram-Positive Bacteria. *Genes (Basel)* **8**
26. Corrigan, R. M., and Gründling, A. (2013) Cyclic di-AMP: another second messenger enters
the fray. *Nat Rev Microbiol* **11**, 513-524
27. Torres, R., Carrasco, B., Gandara, C., Baidya, A. K., Ben-Yehuda, S., and Alonso, J. C. (2019)
Bacillus subtilis DisA regulates RecA-mediated DNA strand exchange. *Nucleic Acids Res* **47**,
5141-5154
28. Tosi, T., Hoshiga, F., Millership, C., Singh, R., Eldrid, C., Patin, D., Mengin-Lecreulx, D.,
Thalassinos, K., Freemont, P., and Gründling, A. (2019) Inhibition of the *Staphylococcus
aureus* c-di-AMP cyclase DacA by direct interaction with the phosphoglucosamine mutase
GlmM. *PLoS Pathog* **15**, e1007537
29. Gundlach, J., Mehne, F. M., Herzberg, C., Kampf, J., Valerius, O., Kaever, V., and Stülke, J.
(2015) An essential poison: Synthesis and degradation of cyclic di-AMP in *Bacillus subtilis*. *J
Bacteriol* **197**, 3265-3274
30. Heidemann, J. L., Neumann, P., Dickmanns, A., and Ficner, R. (2019) Crystal structures of the
c-di-AMP-synthesizing enzyme CdaA. *J Biol Chem* **294**, 10463-10470
31. Zhu, Y., Pham, T. H., Nhiep, T. H., Vu, N. M., Marcellin, E., Chakrabortti, A., Wang, Y.,
Waanders, J., Lo, R., Huston, W. M., Bansal, N., Nielsen, L. K., Liang, Z. X., and Turner, M.
S. (2016) Cyclic-di-AMP synthesis by the diadenylate cyclase CdaA is modulated by the
peptidoglycan biosynthesis enzyme GlmM in *Lactococcus lactis*. *Mol Microbiol* **99**, 1015-1027
32. Pham, T. H., Liang, Z. X., Marcellin, E., and Turner, M. S. (2016) Replenishing the cyclic-di-
AMP pool: regulation of diadenylate cyclase activity in bacteria. *Curr Genet* **62**, 731-738
33. Gibhardt, J., Heidemann, J. L., Bremenkamp, R., Rosenberg, J., Seifert, R., Kaever, V., Ficner,
R., and Commichau, F. M. (2020) An extracytoplasmic protein and a moonlighting enzyme
modulate synthesis of c-di-AMP in *Listeria monocytogenes*. *Environ Microbiol* **22**, 2771-2791
34. Rismondo, J., Gibhardt, J., Rosenberg, J., Kaever, V., Halbedel, S., and Commichau, F. M.
(2016) Phenotypes Associated with the Essential Diadenylate Cyclase CdaA and Its Potential
Regulator CdaR in the Human Pathogen *Listeria monocytogenes*. *J Bacteriol* **198**, 416-426
35. Barreteau, H., Kovac, A., Boniface, A., Sova, M., Gobec, S., and Blanot, D. (2008) Cytoplasmic
steps of peptidoglycan biosynthesis. *FEMS Microbiol Rev* **32**, 168-207
36. Krissinel, E., and Henrick, K. (2007) Inference of macromolecular assemblies from crystalline
state. *J Mol Biol* **372**, 774-797
37. Mehra-Chaudhary, R., Mick, J., and Beamer, L. J. (2011) Crystal structure of *Bacillus anthracis*
phosphoglucosamine mutase, an enzyme in the peptidoglycan biosynthetic pathway. *J Bacteriol*
193, 4081-4087

38.   Zheng, H., Cooper, D. R., Porebski, P. J., Shabalin, I. G., Handing, K. B., and Minor, W. (2017)
698   CheckMyMetal: a macromolecular metal-binding validation tool. *Acta Crystallogr D Struct*
699   *Biol* **73**, 223-233
700   39.   Celniker, G., Nimrod, G., Ashkenazy, H., Glaser, F., Martz, E., Mayrose, I., Pupko, T., and
701   Ben-Tal, N. (2013) ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about
702   Protein Function. *Isr J Chem* **53**, 199-206
703   40.   Tan, E., Rao, F., Pasunooti, S., Pham, T. H., Soehano, I., Turner, M. S., Liew, C. W., Lescar,
704   J., Pervushin, K., and Liang, Z. X. (2013) Solution structure of the PAS domain of a
705   thermophilic YybT protein homolog reveals a potential ligand-binding site. *J Biol Chem* **288**,
706   11949-11959
707   41.   Rao, F., Ji, Q., Soehano, I., and Liang, Z. X. (2011) Unusual heme-binding PAS domain from
708   YybT family proteins. *J Bacteriol* **193**, 1543-1551
709   42.   Raguse, M., Torres, R., Seco, E. M., Gandara, C., Ayora, S., Moeller, R., and Alonso, J. C.
710   (2017) *Bacillus subtilis* DisA helps to circumvent replicative stress during spore revival. *DNA*
711   *Repair (Amst)* **59**, 57-68
712   43.   Winter, G., Waterman, D. G., Parkhurst, J. M., Brewster, A. S., Gildea, R. J., Gerstel, M.,
713   Fuentes-Montero, L., Vollmar, M., Michels-Clark, T., Young, I. D., Sauter, N. K., and Evans,
714   G. (2018) DIALS: implementation and evaluation of a new integration package. *Acta*
715   *Crystallogr D Struct Biol* **74**, 85-97
716   44.   Evans, P. R., and Murshudov, G. N. (2013) How good are my data and what is the resolution?
717   *Acta Crystallogr D* **69**, 1204-1214
718   45.   Bunkoczi, G., Echols, N., McCoy, A. J., Oeffner, R. D., Adams, P. D., and Read, R. J. (2013)
719   Phaser.MRage: automated molecular replacement. *Acta Crystallogr D Biol Crystallogr* **69**,
720   2276-2286
721   46.   Liebschner, D., Afonine, P. V., Baker, M. L., Bunkoczi, G., Chen, V. B., Croll, T. I., Hintze,
722   B., Hung, L. W., Jain, S., McCoy, A. J., Moriarty, N. W., Oeffner, R. D., Poon, B. K., Prisant,
723   M. G., Read, R. J., Richardson, J. S., Richardson, D. C., Sammito, M. D., Sobolev, O. V.,
724   Stockwell, D. H., Terwilliger, T. C., Urzhumtsev, A. G., Videau, L. L., Williams, C. J., and
725   Adams, P. D. (2019) Macromolecular structure determination using X-rays, neutrons and
726   electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877
727   47.   Emsley, P., Lohkamp, B., Scott, W. G., and Cowtan, K. (2010) Features and development of
728   Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501
729   48.   Headd, J. J., Echols, N., Afonine, P. V., Grosse-Kunstleve, R. W., Chen, V. B., Moriarty, N.
730   W., Richardson, D. C., Richardson, J. S., and Adams, P. D. (2012) Use of knowledge-based
731   restraints in phenix.refine to improve macromolecular refinement at low resolution. *Acta*
732   *Crystallogr D Biol Crystallogr* **68**, 381-390
733   49.   Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N. (2016)
734   ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation
735   in macromolecules. *Nucleic Acids Research* **44**, W344-W350
736   50.   Förster, S., Apostol, L., and Bras, W. (2010) Scatter: software for the analysis of nano- and
737   mesoscale small-angle scattering. *Journal of Applied Crystallography* **43**, 639-646
738   51.   Franke, D., and Svergun, D. I. (2009) DAMMIF, a program for rapid ab-initio shape
739   determination in small-angle scattering. *J Appl Crystallogr* **42**, 342-346
740   52.   Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C.,
741   and Ferrin, T. E. (2004) UCSF Chimera--a visualization system for exploratory research and
742   analysis. *J Comput Chem* **25**, 1605-1612
743   53.   Burkholder, P. R., and Giles, N. H., Jr. (1947) Induced biochemical mutations in *Bacillus*
744   *subtilis*. *Am J Bot* **34**, 345-348
745   54.   Laskowski, R. A., and Swindells, M. B. (2011) LigPlot+: multiple ligand-protein interaction
746   diagrams for drug discovery. *J Chem Inf Model* **51**, 2778-2786
747
748

749   **Tables**

750   **Table 1:** Crystallographic data and refinement statistics

	$CdaA_{CD}$	GlmM (PO$_4$ bound)	GlmM (metal bound)	$CdaA_{CD}$ and $GlmM_{F369}$ Complex 1	$CdaA_{CD}$ and $GlmM_{F369}$ Complex 2
Data collection					
Space group	P 4$_3$ 2$_1$ 2	P 3$_2$ 2 1	P 3$_2$ 2 1	P 1 2$_1$ 1	P 1 2$_1$ 1
Cell dimensions					
a,b,c (Å)	62.88, 62.88, 187.32	134.87, 134.87, 69.18	134.41, 134.41, 69.01	62.20, 227.56, 151.60	62.80, 228.53, 153.19
α,β,γ (°)	90.00, 90.00, 90.00	90.00, 90.00, 120.00	90.00, 90.00, 120.00	90.00, 99.66, 90.00	90.00, 99.86, 90.00
Resolution (Å)	93.66 (2.8)*	48.290 (3.0)*	48.15 (2.9)*	61.32 (3.65)*	76.18 (4.2)*
Unique reflections	9974 (1396)*	14081 (2497)*	15955 (2503)*	45925 (4441)*	31005 (4490)*
Rpim	0.05 (0.52)*	0.13 (2.58)*	0.09 (1.46)*	0.019 (0.15)*	0.06 (0.22)*
CC(1/2)	0.99 (0.97)*	0.54 (0.54)*	0.99 (0.39)*	0.99 (0.53)*	0.78 (0.39)*
$I / \sigma(I)$	18.2 (5.9)*	6.9 (2.4)*	12.6 (4.4)*	5.5 (1.2)*	2.1 (0.9)*
Completeness (%)	100.0 (100.0)*	99.9 (99.5)*	98.8 (97.3)*	99.6 (99.2)*	99.9 (99.9)*
Redundancy	23.2 (24.1)*	19.6 (17.3)*	20.2 (17.5)*	6.9 (7.1)	3.8 (3.8)*
Refinement					
Resolution (Å)	2.8	3.0	2.9	3.6	4.2
Rwork / Rfree(%)[a]	0.2545/ 0.2966	0.1996/ 0.2589	0.1868/ 0.2348	0.2418/0.2615	0.2473/ 0.2976
Rms deviations					
Bond lengths (Å)	0.008	0.016	0.0015	0.003	0.003
Bond angles (°)	1.192	1.533	1.396	0.658	0.703
Ramachandran plot (%) Most favoured/ outliers	89.82/ 2.11	95.71/ 0.68	87.36/ 1.81	92.49/1.44	92.94/ 0.88
PDB code	6HUW	7OJR	7OML	7OLH	7OJS

751   * refers to the highest resolution shell

752   [a]R$_{free}$ was computed same as R$_{work}$, using a test set (~5%) of randomly selected reflections that
753   were omitted from the refinement

758
759
Figure 1: **The *B. subtilis* GlmM and CdaA$_{CD}$ proteins form a complex *in vitro***. *A,* FPLC
chromatograms and SDS-PAGE gel analysis of the *B. subtilis* CdaA$_{CD}$ and GlmM complex. The *B.*
subtilis proteins CdaA$_{CD}$, GlmM and the CdaA$_{CD}$/GlmM complex were purified by Ni-affinity
chromatography and size-exclusion chromatography. The FPLC elution profiles, recorded at a
wavelength of 280 nm, are shown for CdaA$_{CD}$ (blue), GlmM (cyan) and the CdaA$_{CD}$/GlmM complex
(pink). Proteins from the peak fractions of the CdaA$_{CD}$.GlmM complex were separated on a 12% SDS
PAGE gel and proteins visualized by Coomassie staining (shown in the insert). *B,* FPLC chromatograms
and SDS-PAGE gel analysis of the *B. subtilis* CdaA$_{CD}$ and GlmM$_{F369}$ complex. Same as in (A) but using
the C-terminally truncated GlmM$_{F369}$ variant in place of the full-length GlmM protein. The experiments
were performed in triplicates and representative chromatograms and gel images are shown.
770
771

773
774
Figure 2: Enzymatic activity of the *B. subtilis* CdaA$_{CD}$ enzyme is inhibited by GlmM or GlmM$_{F369}$.
A, Metal-dependency of the *B. subtilis* CdaA$_{CD}$ enzyme. The metal-dependency of the *B. subtilis*
CdaA$_{CD}$ was assessed by performing enzyme reactions using 5 μM of purified CdaA$_{CD}$ in buffer
containing 1 mM Mn^{+2}, Mg^{+2} or Co^{+2} ions. After 4 h of incubation, the reactions were stopped, separated
by TLC and the percentage conversion of radiolabelled ATP to c-di-AMP determined. The average
values and standard deviations (SDs) from three experiments were plotted. *B,* CdaA$_{CD}$ activity time-
course experiment. Enzyme reactions were set up with the *B. subtilis* CdaA$_{CD}$ enzyme in buffer
containing 1 mM Mn^{+2}, aliquots removed, and reactions stopped at the indicated time points and
separated by TLC. The percentage conversion of radiolabelled ATP to c-di-AMP was determined and
the average values and SDs from three independent experiments plotted. *C,* Enzyme activity of the *B.*
subtilis CdaA$_{CD}$.enzyme in the presence of *B. subtilis* GlmM. The enzyme activity of CdaA$_{CD}$ was
measured in the absence or presence of GlmM or GlmM$_{F369}$ at a molar ratio of 2:1 GlmM or GlmM$_{F369}$
to CdaA$_{CD}$. After 4 h of incubation, the reactions were stopped, separated by TLC and the percentage
conversion of radiolabelled ATP to c-di-AMP was determined. The average values and SDs from six
independent experiments plotted. One-way ANOVA tests followed by Dunnett's multiple comparison
were performed to identify statistically significant differences in cyclase activity in the absence or
presence of GlmM or GlmM$_{F369}$. *** indicates $p < 0.0001$.
792
793
794

Figure 3: Crystal structures of the *B. subtilis* CdaA$_{CD}$ and GlmM enzymes. *A, B. subtilis* CdaA$_{CD}$ structure in cartoon representation. The CdaA$_{CD}$ protein crystallized as dimer and individual CdaA$_{CD}$ monomers are shown in blue and grey, with active site DGA and RHR motifs highlighted in green. *B,* Superimposition of the *B. subtilis, L. monocytogenes* and *S. aureus* CdaA$_{CD}$ (DacA$_{CD}$) structures. Monomers of the *B. subtilis* CdaA$_{CD}$ (grey), *L. monocytogenes* CdaA$_{CD}$ (yellow; PDB ID 4RV7) and *S. aureus* DacA$_{CD}$ (pink; PDB ID 6GYW) protein structures were superimposed using COOT (47). *C,* CdaA$_{CD}$ dimer interface and zoomed in view showing residues Leu150, Lys153, Asn161 and Leu169, which form hydrogen-bonds, in stick representation. *D, B. subtilis* GlmM structure in cartoon representation. The GlmM protein crystalized as a dimer (Fig. S2) with one monomer in the asymmetric unit. The GlmM monomer displayed the typical 4 domain architecture, and the individual domains are represented in different colours. A metal-ion was bound to the protein and is shown as yellow sphere. *E,* Superposition of the *B. subtilis, S. aureus* and *B. anthracis* GlmM structures. *B. subtilis* GlmM (grey), *B. anthracis* GlmM (yellow; PDB 3PDK) and *S. aureus* GlmM (pink; PDB 6GYX) monomer structures were superimposed using COOT (47). *F,* Zoomed in view of the superimposed *B. subtilis* (grey; metal-ion bound structure)*, S. aureus* (pink) and *B. anthracis* (yellow) GlmM structures showing the distances between the indicated Arg residue within the phosphate binding site and the active site Ser residue. Images were generated using PyMOL (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC).

819
820
Figure 4: Crystal structure of the *B. subtilis* CdaA$_{CD}$:GlmM$_{F369}$ complex 1. (*A-D*) Structure of the
B. subtilis CdaA$_{CD}$:GlmM$_{F369}$ complex 1 shown in cartoon representation. The *B. subtilis*
CdaA$_{CD}$:GlmM$_{F369}$ complex crystallized as a dimer of dimers with individual GlmM$_{F369}$ monomers
shown in dark pink [GlmM$_{F369}$(1)] and light pink [GlmM$_{F369}$(2)] and individual CdaA$_{CD}$ monomers
shown in dark blue [CdaA$_{CD}$(1)] and light blue [CdaA$_{CD}$(2)], respectively. The complex is show in *A,* in
front view, *B,* in side view (rotated 90º along the y-axis) and *C,* in top-side view rotated at the angle as
indicated with respect to (A). *D,* a zoom in view of the CdaA$_{CD}$/GlmM$_{F369}$ interface is shown. Residue
Arg126 from CdaA$_{CD}$(1) forms H-bond and ionic interactions with Asp 151 and Glu154 of GlmM$_{F369}$(1)
(residues shown in yellow), and residue Arg126 from CdaA$_{CD}$(2) forms ionic interactions with Asp195
in GlmM$_{F369}$(1) (residues shown in green). The images were prepared in PyMOL (The PyMOL
Molecular Graphics System, Version 2.0 Schrödinger, LLC).
832
833
834

835
836
837
Figure 5: **Formation of an active CdaA$_{CD}$ dimer is blocked in the complex.** (*A-B*) Structure of the *B.*
subtilis CdaA$_{CD}$ dimer from complex 1 (top panels) and the complete CdaA$_{CD}$:GlmM$_{F369}$ complex 1
(bottom panels) shown as space filling models. The active site DGA and RHR motifs in CdaA$_{CD}$(1)
(dark blue protomer) are shown in yellow and in CdaA$_{CD}$(2) (light blue protomer) in green. *A*, Bottom
panel, complex model showing that the active site (green residues) in CdaA$_{CD}$(2) (light blue protomer)
are completely occluded in the complex and *B,* the active site (yellow residues) of CdaA$_{CD}$(1) (dark blue
protomer) remains partially exposed. However, the modelling of another CdaA$_{CD}$ protomer (in red) that
forms an active dimer with CdaA$_{CD}$(1) (dark blue protomer) results in protein-protein clashes with the
GlmM$_{F369}$(1) monomer in the complex. The active CdaA$_{CD}$ dimer was modelled based on the *L.*
monocytogenes Δ100CdaA (PDB 6HVL) structure (30).
848
849

Figure 6: Small Angle X-ray Scattering (SAXS) data of the *B. subtilis* CdaA$_{CD}$, GlmM proteins and the *B. subtilis* and *S. aureus* CdaA$_{CD}$:GlmM complex. *A-C,* SAXS envelopes with fitted protein structures. For the SAXS experiment, 50 µl of CdaA$_{CD}$ (5 mg/ml), GlmM (24 mg/ml) and the complex CdaA$_{CD}$:GlmM (24 mg/ml) were injected onto a high pressure Shodex column coupled to the B21 Small-Angle X-Ray beamline at the Diamond Light Source (Didcot, UK). The data analysis and envelope reconstruction were performed using ScÅtter (50). The program Chimera (52) was used to visualise the reconstructed envelopes as maps into which the protein structures were fitted. *A,* SAXS envelope of the *B. subtilis* CdaA$_{CD}$ protein with a CdaA$_{CD}$ dimer protein structure fitted into the envelop. *B,* SAXS envelope of the *B. subtilis* GlmM protein with the GlmM protein dimer structure fitted into the envelop. *C,* SAXS envelope of the *B. subtilis* CdaA$_{CD}$:GlmM complex with a CdaA$_{CD}$:GlmM dimer model structure fitted into the envelope.

Figure 7: **Residue Arg126 in *B. subtilis* CdaA$_{CD}$-R126A is important for complex formation with GlmM**. *A,* FPLC chromatograms and SDS-PAGE gel analysis of the *B. subtilis* CdaA$_{CD}$-R126A, GlmM and a mix of both proteins. The *B. subtilis* CdaA$_{CD}$-R126A, GlmM and the CdaA$_{CD}$-R126A/GlmM protein mixture were purified by Ni-affinity chromatography and size-exclusion chromatography. The FPLC elution profiles, recorded at a wavelength of 280 nm, are shown for CdaA$_{CD}$-R126A (blue), GlmM (cyan) and the CdaA$_{CD}$-R126A/GlmM mixture (pink). Proteins from the peak fractions of the CdaA$_{CD}$-R126A/GlmM sample were separated on a 12% SDS PAGE gel and proteins visualized by coomassie staining (shown in the insert). The experiments were performed in triplicates and representative chromatograms and gel images are shown. *B,* Enzymatic activity of wild type CdaA$_{CD}$ and the CdaA$_{CD}$-R126A protein variant in the absence or presence of GlmM. Enzyme activity assays were performed with 5 μM purified CdaA$_{CD}$ or CdaA$_{CD}$-R126 in absence or presence of 10 μM GlmM protein. After 4 h of incubation, the reactions were stopped, separated by TLC and the percentage conversion of radiolabelled ATP to c-di-AMP determined. The average values and SDs from six experiments were plotted. T-tests were performed to determine statistically significant differences in enzyme activity of CdaA$_{CD}$ or CdaA$_{CD}$-R126A in the absence or presence of GlmM. ns indicates not statistically significant; *** indicates $p < 0.001$.