The geometry of decision-making

Vivek Hari Sridhar[a,b,c,1]**, Liang Li**[a,b,c,1]**, Dan Gorbonos**[a,b,c]**, Máté Nagy**[a,b,c,d,e,f]**, Bianca R. Schell**[g]**, Timothy Sorochkin**[h,i]**, Nir S. Gov**[i] **and Iain D. Couzin**[a,b,c,1]

[a]Department of Collective Behaviour, Max Planck Institute of Animal Behavior, Konstanz, 78464 Konstanz, Germany

[b]Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78464 Konstanz, Germany

[c]Department of Biology, University of Konstanz, 78464 Konstanz, Germany

[d]MTA-ELTE Statistical and Biological Physics Research Group, Hungarian Academy of Sciences, 1117, Budapest, Hungary

[e]MTA-ELTE 'Lendület' Collective Behaviour Research Group, Hungarian Academy of Sciences, 1117, Budapest, Hungary

[f]Department of Biological Physics, Eötvös Loránd University, 1117, Budapest, Hungary

[g]Department of Chemistry, University of Konstanz, 78464 Konstanz, Germany

[h]Department of Physics & Astronomy, University of Waterloo, Waterloo, ON N2L 3G1, Canada

[i]Department of Chemical and Biological Physics, Weizmann Institute of Science, Rehovot 76100, Israel

Choosing among spatially-distributed options is a central challenge for animals, from deciding among alternative potential food sources or refuges, to choosing with whom to associate. Using an integrated theoretical and experimental approach (employing immersive virtual reality), we consider the interplay between movement and vectorial integration during decision-making regarding two, or more, options in space. In computational models of this process we reveal the occurrence of spontaneous and abrupt "critical" transitions (associated with specific geometrical relationships) whereby organisms spontaneously switch from averaging vectorial information among, to suddenly excluding one, among the remaining options. This bifurcation process repeats until only one option—the one ultimately selected—remains. Thus we predict that the

brain repeatedly breaks multi-choice decisions into a series of binary decisions in space-time. Experiments with fruit flies, desert locusts, and larval zebrafish reveal that they exhibit these same bifurcations, demonstrating that across taxa and ecological context, there exist fundamental geometric principles that are essential to explain how, and why, animals move the way they do.

Animals constantly face the need to make decisions, and many such decisions require choosing among multiple spatially-distributed options. Despite this, most studies have focused on the outcome of decisions (1–3) (i.e. which option among alternatives is chosen), as well as the time taken to make decisions (4–6), but seldom on the movement of animals throughout the decision-making process. Motion is, however, crucial in terms of how space is represented by organisms during spatial decision-making; the brains of a wide range of species, from insects (7, 8) to vertebrates (9, 10), have been shown to represent egocentric spatial relationships, such as the position of desired targets, via explicit vectorial representation (11, 12). Such neuronal representations must, and do, change as animals move through space. Thus, while the movement of an animal may, initially, appear to simply be a readout of the decision made by the brain—and consequently not particularly informative—this view overlooks important dynamical properties introduced into the decision-making process that result from the inevitable time-varying geometrical relationships between an organism and spatially-distributed options (i.e. potential 'targets' in space).

Due to a dearth of existing studies, and with the objective to develop the necessary foundational understanding of the 'geometry' of decision-making, we focus here—first theoretically and then experimentally—on the consequences of the recursive interplay between movement and (collective) vectorial integration in the brain during relatively simple spatial decisions. We

employ immersive virtual reality to investigate decision-making regarding multiple (two or

more) options in both invertebrate (the fruit fly *Drosophila melanogaster*, and desert locust

Schistocerca gregaria) and vertebrate (larval zebrafish *Danio rerio*) models. Doing so allows

us to reveal the emergence of geometric principles that transcend the study organism and the

decision-making context, and thus are expected to be broadly relevant across taxa. In support

of this finding we also explore how these principles extend to collective decision-making in

mobile animal groups, allowing us to gain insights across three scales of biological

organisation, from neural dynamics, to both individual and collective decision-making.

Modelling decision-making on the move

Congruent with neurobiological studies of the invertebrate and vertebrate brain, we consider

organisms to have an egocentric vectorial representation of spatial options (11–13). We then

consider the collective dynamics of vector integration in the brain assuming there exists

reinforcement (excitation/positive feedback) among neural ensembles that have similar

directional representations (goal vectors), and global inhibition and/or negative feedback (both

produce broadly similar results, see SI Appendix and Fig. S1) among neural ensembles that

differ in vectorial representation. This captures, in a simple mathematical formulation, the

essence of both explicit ring-attractor networks (as found in insects (7)), and computation

among competing neural groups (as in the mammalian brain (14)). The animal's relative

preference for a target is given by activity of neurons that encode direction to that target relative

to activity of neurons that encode direction to other targets, and the angular sensitivity of the

neural representations (angular difference at which excitation no longer occurs) is specified by

a neural tuning parameter, ν. The network then computes a unique 'consensus' vector ('activity

bump') that, along with some angular noise, represents the animal's desired direction of

movement (Fig. S2). This is then translated into motor output (see SI Appendix for model

details \cite{pinkoviezky_collective_2018}). Stochasticity in neural dynamics is implemented

here as the neural noise parameter, T.

While capturing known, generic features of neural integration, our model is deliberately

minimal. This serves multiple purposes: firstly, following principles of maximum parsimony

we seek to find a simple model that can both predict and explain, the observed phenomena;

secondly, we aim to reveal general principles and thus consider features that are known to be

valid across organisms irrespective of inevitable difference in structural organization of the

brain; thirdly, it provides a convenient means to implement neural noise, and can be mapped

to the class of neural ring-attractor models widely-used in neuroscience (15–18) (see SI

Appendix for details). In addition, our results are shown to be extremely robust to model

assumptions, suggesting that it provides an appropriate low-level description of essential

system properties.

Deciding between two options

Beginning with the simplest case, we consider the feedback between motion and internal

vectorial-computation when an animal is presented with two equally-attractive, but spatially-

discrete, options. In this case the activity of neurons encoding option 1, N_1 will be equal to

those encoding option 2, N_2 (Fig. 1A). Our model predicts that an animal moving, from a

relatively distant location, towards the two targets, will spontaneously compute the average

directional preference, resulting in corresponding motion in a direction oriented between the

two targets. As it approaches the targets, however, upon reaching a certain angular difference

between the options, the internal network undergoes a sudden transition in which it

spontaneously selects one, or the other, target (Fig. 1C). This results in an abrupt change in

trajectory, the animal being redirected towards the respective 'selected' target (Fig. 1C; see

also Fig. S3A for the same phenomenon occurring for a wide range of starting positions).

Our model therefore predicts that despite the fact that the egocentric geometrical relationship

between the animal and the targets changes continuously, upon approaching the targets, there

exists a location whereby a further, very small, increase in angular difference between the

targets will result in a sudden change in system (neural) dynamics, and consequently in motion,

and thus decision-making. Such spatio-temporal dynamics do not occur if individuals were to

simply integrate noisy vectorial information or choose their travel direction from a summed

distribution of the location of targets in their sensory field (19), points we will return to later.

In numerical analysis of our model we find that irrespective of starting position, as the animal

reaches the respective angle in space, it will relatively suddenly select one of the options (Fig.

S3A). While the specific angular difference at which this phenomenon occurs is dependent on

neural tuning, ν (Fig. S3C), and the starting configuration (due to an interplay between the two

timescales involved—for movement and for neural consensus, see Fig. S3B), it is always

present as long as the neural noise, T remains below a critical firing rate, T_c (although even for

$T < T_c$, these bifurcations may be difficult to see for small values of ν due to inherent noise in

real biological systems; see Fig. S4 for simulations where vectorial representations of targets

include directional error).

To gain a deeper insight into the mechanism underlying the observed spatiotemporal dynamics,

we constructed a mean-field approximation of our model (see SI Appendix) since this has the

advantage of allowing us to conduct formal analyses of patterns realized in the simulated

trajectories.

Geometric principles of decision-making

The mean-field analysis of our model shows that below a critical level of neural noise, animals

will adopt the average among options as they approach the targets, until a critical phase

transition upon which the system spontaneously switches to deciding among the options (Figs.

1B and S5A). Thus despite varying in its exact location (Fig. 1B), the sudden transition

observed is an inevitable consequence of the system dynamics and will always occur.

Such sudden transitions correspond to 'bifurcations' in the mathematical study of dynamical

systems. A bifurcation is said to occur when a smooth change in an external parameter, in this

case perceived angular difference between the options, causes a sudden qualitative change in

the system's behavior, here corresponding to a literal bifurcation (or branching) in physical

space.

When dynamical systems undergo such a phase, or quasi-phase, transition they exhibit a

remarkable universal property: close to the transition, at the "critical-point" or "tipping-point",

the system spontaneously becomes extremely sensitive to very small perturbations (e.g. to

small differences in preference between options (20, 21)). This is true of both physical (e.g.

magnetic (22)) and biotic (e.g. cellular (23, 24)) systems undergoing a phase transition.

Correspondingly, we find that below a critical level of neural noise, the mean-field model

exhibits a sudden increase in susceptibility as the animal approaches the critical point,

immediately prior to the decision being made (Fig. S5A). This will not occur in previously-

considered models where an animal is assumed to choose its direction of travel based on the

summed distribution of targets in its sensory field, also known as probability density function,

or PDF, sum-based models (19). Thus, as animals approach targets we predict they will pass

through a window of space (corresponding to the critical angle for the respective geometry they

are experiencing) in which their brain spontaneously becomes capable of discriminating

between very small differences between options (e.g. a very small difference in neuronal

activity being in 'favor' of one option; see Fig. S3D and SI Appendix for details). This highly-

valuable property (for decision-making) is not built into the model, but is rather an emergent

property of the inherent collective dynamics.

In many real biological systems, including the ones we consider here, the (neural) system size

is typically not large enough to consider true phase transitions (which only occur for very large

systems, as per the mean-field approximation), but rather 'phase-transition-like', or 'quasi-

phase transition', behavior. Even though real biological systems are not necessarily close to the

infinite size limit of the mean-field approximation, we see very similar dynamics for both small

and large system sizes (Fig. S6).

Decision-making beyond two options

While the majority of decision-making studies consider only two options (due to both

theoretical and experimental tractability (14, 25, 26)), animals moving in real space frequently

encounter a greater number than this. Here we consider how animals will be expected to select

among three, or more, options (possible targets) in space. First we begin with three identical

options ($N_1 = N_2 = N_3$) since this gives us the clearest insight into the relationship between

motion and decision-making dynamics. Then we relax these assumptions and consider

differences between options (Fig. S3E) as well as a greater number of options (Fig. 2). Note

that we do not modify our model in any way prior to introducing these additional complexities.

Below T_c (see SI Appendix and Fig. S7 for considerations when $T > T_c$), we once again find

that the direction in which the animal moves is a function of the angular difference between

the targets. When relatively far from the targets, it moves in the average of these three

directions. Upon reaching a critical angular threshold between the leftmost and rightmost

option (from the animal's perspective), however, the neural system spontaneously eliminates

one of them and the animal begins moving in the direction average between the two remaining

options (Fig. 1D,E). It continues in this direction until a second critical angle is reached, and

now the animal eliminates one of the two remaining options and moves towards the only

remaining target (Figs. 1F and S5B). Thus we predict that the brain repeatedly breaks multi-

choice decisions into a series of binary decisions in space-time. Such bifurcation dynamics are

not captured in models of decision-making that do not include the required feedbacks, such as

if individuals simply sum noisy vectors (or PDFs) to targets in their sensory field (19). For the

case of three targets, vectors/votes to the leftmost option would tend to cancel those that favor

the rightmost option, resulting in the selection of the central option, an issue we will return to

later when considering collective animal behavior. Simulating a larger number of options (Fig.

2) and varying environmental geometries (Figs. S8 and S9) demonstrate the robustness of this

mechanism in the face of environmental complexity and the more complex spatial dynamics

that emerge as organisms undergo repeated bifurcations.

Experimental tests of our predictions

Since the decision-process is predicted to be sequential, and dependent on the geometry with

respect to the targets from an egocentric perspective, it should be possible to visualize it directly

from the trajectories taken by animals when making spatial decisions. In this respect, our

theoretical studies make a key testable prediction: if neural groups within the decision-making

ensemble exhibit relatively local excitation, and long-range/global inhibition, we should

observe bifurcations in the animals' trajectories as they choose among identical options; and

that if animals face three (or more) such options, then the complex decision task should be

broken down to a series of binary decisions.

Since the geometrical principles revealed above are expected to be both robust and generic, we

use immersive virtual reality (27) (Fig. S10) to test our predictions by investigating both two-

and three-choice decision-making in three evolutionarily highly-divergent brains under

ecologically-relevant scenarios: fruit flies (*Drosophila melanogaster*) and desert locusts

(*Schistocerca gregaria*) deciding which among multiple vertical objects to approach (e.g. to

perch), and zebrafish (*Danio rerio*) choosing with which conspecific(s) to school.

Like many other insects (28–31), fruit flies (32) and desert locusts (33) exhibit a natural

tendency to orient and move towards high-contrast vertical features (potential landing sites or

indicators of vegetation) in their environment. We exploit this tendency, presenting multiple

identical black pillars as targets in an otherwise white environment. We record trajectories of

our focal animals (solitary flies or locusts) as they choose to move towards one of these pillars,

thus obtaining a behavioral readout of the decision-making process (see SI Appendix for

experimental details; Figs. S11 and S12 show raw trajectories of flies and locusts respectively).

As predicted by our theory (Fig. 1B,C), we find that, in the two-choice case, most flies and

locusts that choose one of the presented targets initially move in the average of the egocentric

target directions until a critical angular difference (Fig. S13), at which point they select

(randomly) one, or the other, option and move towards it (randomization test where

y −coordinates between trajectories were swapped showed that the bifurcation fit to our

experimental data was highly significant; $p < 0.01$ for both flies and locusts; Figs. 1G and

S13). Here, we note that there may be multiple factors that affect the animals' direction of

movement. For example, it could be that animals repeatedly switch between fixating on each

of the two options before reaching the critical angular difference, following which it selects

one. However, quantification of their heading relative to the targets, and to the average

direction between the targets (Fig. S13), finds no evidence for this; instead, prior to the

bifurcation, both flies and locusts exhibit a heading towards the average of the egocentric target

directions. In the three-choice case, the animals' movements are also consistent with our

theory; as predicted (Fig. 1E,F) they break the three-choice decision into two sequential binary

decisions ($p < 10^{-4}$ for both flies and locusts; Fig. 1H). For both animals, the observed angle

of bifurcation (~110° for flies and ~90° for locusts) is much larger than their visual spatial

resolution (~8° and ~2° for flies (34) and locusts (35, 36), respectively). We note ~30% of

animals in our experiments (both flies and locusts) did not exhibit the sequential bifurcations

(see Figs. S11 and S12) described above, and instead moved directly towards one of the

presented targets (Figs. S11 and S12). Such variability in response is expected in animals, and

is consistent with recent work on the visual response of flies, which demonstrates a link

between stochastic (non-heritable) variation in brain wiring within the visual system and

strength of visual orientation response to a vertical stripe target (37). Furthermore, flies that

experience high temperatures during development appear to exhibit a particularly strong

orientation tendency, exhibiting the most direct paths to targets while flies that experience low

developmental temperatures exhibit wandering paths to targets (38). In our model such

differences can be accounted for by variation in directional tuning of the neural groups, with

high directional tuning (low ν) being associated with a strong orientational response, and such

individuals exhibiting direct tracks to targets from the outset (see Fig. S14).

A further, non-mutually exclusive, possibility, is that a subset of insects exhibit "handedness".

For example, in (39), it was shown that approximately 25% of *Drosophila* were either strongly

left-biased or right-biased when moving on a Y-maze, and that these consistent differences

among flies were similarly non-heritable. This experimental design did not assess whether a

further subset are biased to go directly forwards if offered three directional choices (such as

could occur in a hypothetical Ψ maze). In such cases, it is certainly possible that these intrinsic

directional biases break symmetry (Fig. S3D,E), resulting in directed paths to different targets.

We note that individuals predisposed to exhibit direct paths to targets would be expected to

make faster, yet less accurate, decisions, a prediction we plan to test in future studies.

Our zebrafish experiments consider spatial decision-making in a social context. We present

virtual conspecifics (see SI Appendix for methodological details) that move back-and-forth in

the arena parallel to each other as targets (Figs. 3A and S15A) and behave (Fig. S16), and are

responded to (Fig. S17), in the same way as real fish. Because they are social, the real fish

respond to these virtual fish by tending to follow at a (relatively) fixed distance behind them

(Fig. S15E). Our data are best represented within this moving frame of reference (the virtual

fish; Fig. S15). Theoretically we predict that for two virtual fish we should see a single

bifurcation, where the real fish will suddenly switch from averaging the target directions to

deciding among them (i.e. swimming predominantly with one of the virtual fish), as a function

of increasing the lateral distance, L, between the virtual fish (Figs. 3B and S18; see SI Appendix

for details of model implementation). The existence of this bifurcation is clearly seen in our

experiments (Fig. 3C). When considering three moving virtual conspecifics, the model predicts

that real fish will spontaneously break the three-choice decision to two binary decisions, and a

comparison of the theoretical prediction and experimental results demonstrates this to be the

case (c.f. Fig. 3E,F).

We also test predictions under conditions where there is an asymmetric geometry whereby two

fish swim closer to each other than the central one does to the third fish (Fig. 4A). As predicted

by our theory (Fig. 4B), the real fish tends to swim between the two closely-associated fish, or

close to the third, more distant, fish (Fig. 4B). Note that, also as predicted, the real fish spends

a similar amount of time in each of the two locations.

Although detailed models considering the specifics of each system would be expected to

provide additional quantitative fits (at the expense of losing some degree of generality and

analytical tractability), our results are broadly independent of the model implementation

details. Thus, we find that the key predictions of our model are validated in fruit flies, desert

locusts and larval zebrafish in distinct, yet ecologically relevant contexts.

Model features that determine network behavior

There are key features that are essential to produce the bifurcation patterns observed in our data

i.e. for any decision-making system to break multi-choice decisions to a series of binary

decisions.

• Feedback processes that provide the system directional persistence, and drive such

bifurcations, are crucial to exhibit the observed spatio-temporal dynamics. In the neural

system, this is present in the form of local excitation and long-range/global inhibition

(7, 16, 17). However, as shown in our model of collective animal behavior below, we

expect that similar dynamics will be observed if the necessary feedbacks are also

incorporated into other models of decision-making, such as to PDF-sum-based models,

for example (19).

• Observing similar decision dynamics requires a recursive (embodied) interplay

between neural dynamics, and motion in continuous space. Here, the animal's

geometrical relationship with the targets changes as it moves through physical space.

Since neural interactions depend on this changing relationship, space provides a

continuous variable by which the individual traverses the time-varying landscape of

neural firing rates.

These essential features, along with the observed animal trajectories in the two-choice context,

are reminiscent of collective decision-making in animal groups (models (40–44), fish schools

(45), bird flocks (46) and baboon troops (25)). Below, we consider an established model of

collective decision-making (40) to draw links between these two scales of biological

organization—decision-making in the brain, and decision-making in animal groups.

A link to collective decision-making

In order to draw a link between individual decision-making and collective decision-making in

animal groups, we consider an animal group with equal number of individuals exhibiting

preference for each target (see SI Appendix for methodological details). A long-standing

approach in the study of animal collectives is to consider them integrating vectorial information

from neighbors (47, 48), and there are a great number of publications of such "flocking",

"schooling" or "herding" behaviors (47–49). Individuals within groups may also have

preferences to reconcile this local vector-averaging with goal-oriented behavior, such as a

desired direction of travel (40, 45), and such models have made effective predictions regarding

how the number of individuals with a common desired direction of travel influences the

accuracy of group motion towards targets (25) and how the weighting of the internal 'goal-

oriented' vector representing the desired direction of travel, influences the capacity and

accuracy for individuals to act as leaders and to influence the directon taken by the group as a

whole (50).

We demonstrate here, however, that while ubiquitous, such models of collective animal

behavior fail to account for the known capability for animal groups to make decisions among

spatially discrete targets (see Fig. S19A,B). To do so, it is essential that the necessary

feedbacks, as described above for collective decision-making among neurons, are incorporated.

While these feedbacks are inherent to our neural model, they can also be included in other

models in the form of social interactions, or in the animals' response to their environment (51).

For example, one way feedback can be introduced here is via 'informed' individuals (those

with a desired direction of travel) associating with 'uninformed' or 'unbiased' individuals

(individuals that exhibit social interactions but have no specific desired direction of travel) (40,

45); 'uninformed' individuals are effectively recruitable by those with a desired direction of

travel (providing local positive feedback), but are also in finite supply, creating what is

effectively a competition among informed subsets that differ in their preferred direction of

travel (a form of longer-range inhibition between informed subsets). However, because

'uninformed' individuals tend to average the direction of all 'informed' individuals that recruit

them, we find that this type of feedback functions more as a social glue, and is only able to

explain bifurcations when the group is choosing between two options. In a decision-making

context with three options, this type of feedback, alone, results in the group almost always

moving towards the central target (Fig. S19D).

A means of resolving this issue is for individuals to change the strength of their goal-
orientedness as a function of their experienced travel direction; for example, individuals that
find themselves consistently moving in a (group) direction that differs from their preferred
target direction could weaken the strength of their preference over time (a form of
forgetting/negative feedback, effectively resulting in long-range/global inhibition; and once
this preference is lost, they will tend to spontaneously reinforce the majority-selected direction
(45), a form of positive feedback). We find that this biologically-plausible mechanism (40) will
allow individuals within the group to recover the capability to come to consensus even in the
absence of uninformed individuals (Fig. 5), and for a greater number of options than two (Fig.
5B).

Despite considerable differences in details between this model and that of neural dynamics
described above, with the former involving individual components that change neighbor-
relationships over time and where inhibition emerges from a different biological process, the
predictions regarding motion during decision-making are extremely similar (c.f. Fig. 5 and Fig.
1 for a comparison between predictions for animal groups and neural groups, respectively).
Thus, we find that similar principles may underlie spatial decision-making across multiple
scales of biological organization. Furthermore by presenting social interactions in a decision-
making context, our zebrafish experiments elucidate the neural basis of schooling allowing us
to glean insights across three scales of biological organization—from neural dynamics to
individual decisions, and from individual decisions to collective movement.

Conclusions

369 We demonstrate that, across taxa and contexts, explicitly considering the time-varying
370 geometry during spatial decision-making provides new insights that are essential to understand

how, and why, animals move the way they do. The features revealed here are highly robust,

and we predict that they occur in decision-making processes across various scales of biological

organization, from individuals to animal collectives (see Figs. 5 and S19, and SI Appendix),

suggesting they are fundamental features of spatiotemporal computation.

References

1. C. E. J. Kennedy, J. A. Endler, S. L. Poynton, H. McMinn, Parasite load predicts mate choice in guppies. *Behav Ecol Sociobiol* **21**, 291–295 (1987).

2. K. Summers, R. Symula, M. Clough, T. Cronin, Visual mate choice in poison frogs. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **266**, 2141–2145 (1999).

3. E. Forsgren, Predation Risk Affects Mate Choice in a Gobiid Fish. *The American Naturalist* **140**, 1041–1049 (1992).

4. R. Kiani, L. Corthell, M. N. Shadlen, Choice Certainty Is Informed by Both Evidence and Decision Time. *Neuron* **84**, 1329–1342 (2014).

5. J. M. Beck, *et al.*, Probabilistic Population Codes for Bayesian Decision Making. *Neuron* **60**, 1142–1152 (2008).

6. A. K. Churchland, R. Kiani, M. N. Shadlen, Decision-making with multiple alternatives. *Nature Neuroscience* **11**, 693–702 (2008).

7. S. S. Kim, H. Rouault, S. Druckmann, V. Jayaraman, Ring attractor dynamics in the Drosophila central brain. *Science* **356**, 849–853 (2017).

8. J. D. Seelig, V. Jayaraman, Neural dynamics for landmark orientation and angular path integration. *Nature* **521**, 186–191 (2015).

9. J. S. Taube, R. U. Muller, J. B. Ranck, Head-direction cells recorded from the postsubiculum in freely moving rats. II. Effects of environmental manipulations. *Journal of Neuroscience* **10**, 436–447 (1990).

10. A. Finkelstein, *et al.*, Three-dimensional head-direction coding in the bat brain. *Nature* **517**, 159–164 (2015).

11. A. Sarel, A. Finkelstein, L. Las, N. Ulanovsky, Vectorial representation of spatial goals in the hippocampus of bats. *Science* **355**, 176–180 (2017).

12. Ø. A. Høydal, E. R. Skytøen, S. O. Andersson, M.-B. Moser, E. I. Moser, Object-vector coding in the medial entorhinal cortex. *Nature* **568**, 400–404 (2019).

13. R. Wehner, B. Michel, P. Antonsen, Visual navigation in insects: coupling of egocentric
and geocentric information. *Journal of Experimental Biology* **199**, 129–140 (1996).

14. A. Bahl, F. Engert, Neural circuits for evidence accumulation and decision making in
larval zebrafish. *Nature Neuroscience* **23**, 94–102 (2020).

15. K. Zhang, Representation of spatial orientation by the intrinsic dynamics of the head-
direction cell ensemble: a theory. *J. Neurosci.* **16**, 2112–2126 (1996).

16. X. Sun, M. Mangan, S. Yue, An Analysis of a Ring Attractor Model for Cue Integration in
Biomimetic and Biohybrid Systems, Lecture Notes in Computer Science., V. Vouloutsi,
et al., Eds. (Springer International Publishing, 2018), pp. 459–470.

17. L. C. York, M. C. W. van Rossum, Recurrent networks with short term synaptic
depression. *J Comput Neurosci* **27**, 607–620 (2009).

18. S. I. Wiener, J. S. Taube, *Head Direction Cells and the Neural Mechanisms of Spatial
Orientation* (MIT Press, 2005).

19. B. Collignon, A. Séguret, J. Halloy, A stochastic vision-based model inspired by zebrafish
collective behaviour in heterogeneous environments. *Royal Society Open Science* **3**,
150473 (2006).

20. L. Benedetti-Cecchi, L. Tamburello, E. Maggi, F. Bulleri, Experimental perturbations
modify the performance of early warning indicators of regime shift. *Current Biology* **25**,
1867–1872 (2015).

21. A. Gelblum, *et al.*, Ant groups optimally amplify the effect of transiently informed
individuals. *Nature Communications* **6** (2015).

22. B. M. McCoy, T. T. Wu, *The two-dimensional Ising model: second edition* (Courier
Corporation, 2014).

23. J.-M. Choi, A. S. Holehouse, R. V. Pappu, Physical Principles Underlying the Complex
Biology of Intracellular Phase Transitions. *Annual Review of Biophysics* **49**, 107–133
(2020).

24. A. A. Hyman, K. Simons, Beyond Oil and Water—Phase Transitions in Cells. *Science* **337**,
1047–1049 (2012).

25. A. Strandburg-Peshkin, D. R. Farine, I. D. Couzin, M. C. Crofoot, Shared decision-making
drives collective movement in wild baboons. *Science* **348**, 1358–1361 (2015).

26. R. Ratcliff, P. L. Smith, S. D. Brown, G. McKoon, Diffusion Decision Model: Current Issues
and History. *Trends in Cognitive Sciences* **20**, 260–281 (2016).

27. J. R. Stowers, *et al.*, Virtual reality for freely moving animals. *Nature Methods* **14**, 995–
1002 (2017).

28. T. Poggio, W. Reichardt, A theory of the pattern induced flight orientation of the fly
Musca domestica. *Kybernetik* **12**, 185–203 (1973).

29. D. Varju, Stationary and dynamic responses during visual edge fixation by walking
insects. *Nature* **255**, 330–332 (1975).

30. W. Reichardt, T. Poggio, A theory of the pattern induced flight orientation of the fly
Musca domestica II. *Biological Cybernetics* **18**, 69–80 (1975).

31. P. K. Kaushik, M. Renz, S. B. Olsson, Characterizing long-range search behavior in Diptera
using complex 3D virtual environments. *PNAS* **117**, 12201–12207 (2020).

32. E. Horn, R. Wehner, The mechanism of visual pattern fixation in the walking fly,
Drosophila melanogaster. *Journal of Comparative Physiology A* **101**, 39–56 (1975).

33. G. K. Wallace, Some experiments on form perception in the nymphs of the desert locust,
Schistocerca gregaria forskål. *Journal of Experimental Biology* **35**, 765–775 (1958).

34. P. T. Gonzalez-Bellido, T. J. Wardill, M. Juusola, Compound eyes and retinal information
processing in miniature dipteran species match their specific ecological demands.
Proceedings of the National Academy of Sciences **108**, 4224–4229 (2011).

35. O. Faivre, M. Juusola, Visual coding in locust photoreceptors. *Plos One* **3**, e2173 (2008).

36. M. Wilson, Angular sensitivity of light and dark adapted locust retinula cells. *Journal of
Comparative Physiology* **97**, 323–328 (1975).

37. G. A. Linneweber, *et al.*, A neurodevelopmental origin of behavioral individuality in the
Drosophila visual system. *Science* **367**, 1112–1119 (2020).

38. F. R. Kiral, *et al.*, Variable brain wiring through scalable and relative synapse formation in
Drosophila. *bioRxiv* (2021) https:/doi.org/10.1101/2021.05.12.443860.

39. S. M. Buchanan, J. S. Kain, B. L. de Bivort, Neuronal control of locomotor handedness in
Drosophila. *PNAS* **112**, 6700–6705 (2015).

40. I. D. Couzin, J. Krause, N. R. Franks, S. A. Levin, Effective leadership and decision-making
in animal groups on the move. *Nature* **433**, 4 (2005).

41. B. Nabet, N. E. Leonard, I. D. Couzin, S. A. Levin, Dynamics of Decision Making in Animal
Group Motion. *J Nonlinear Sci* **19**, 399–435 (2009).

42. N. E. Leonard, *et al.*, Decision versus compromise for animal groups in motion.
Proceedings of the National Academy of Sciences **109**, 227–232 (2012).

43. R. Gray, A. Franci, V. Srivastava, N. E. Leonard, Multiagent Decision-Making Dynamics
Inspired by Honeybees. *IEEE Transactions on Control of Network Systems* **5**, 793–806
(2018).

44. A. Bizyaeva, A. Franci, N. E. Leonard, A General Model of Opinion Dynamics with Tunable
Sensitivity. *arXiv:2009.04332 [cs, eess, math]* (2020) (July 26, 2021).

45. I. D. Couzin, *et al.*, Uninformed individuals promote democratic consensus in animal
groups. *Science* **334**, 1578–1580 (2011).

46. D. Biro, D. J. T. Sumpter, J. Meade, T. Guilford, From compromise to leadership in pigeon
homing. *Current Biology* **16**, 2123–2128 (2006).

47. I. D. Couzin, J. Krause, R. James, G. D. Ruxton, N. R. Franks, Collective Memory and
Spatial Sorting in Animal Groups. *Journal of Theoretical Biology* **218**, 1–11 (2002).

48. C. W. Reynolds, Flocks, herds and schools: A distributed behavioral model in *Proceedings
of the 14th Annual Conference on Computer Graphics and Interactive Techniques*,
SIGGRAPH '87., (Association for Computing Machinery, 1987), pp. 25–34.

49. T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, O. Shochet, Novel Type of Phase Transition in
a System of Self-Driven Particles. *Phys. Rev. Lett.* **75**, 1226–1229 (1995).

50. C. C. Ioannou, M. Singh, I. D. Couzin, Potential Leaders Trade Off Goal-Oriented and
Socially Oriented Behavior in Mobile Animal Groups. *The American Naturalist* **186**,
284–293 (2015).

51. A. Perna, *et al.*, Individual Rules for Trail Pattern Formation in Argentine Ants
(Linepithema humile). *PLOS Computational Biology* **8**, e1002592 (2012).

52. I. Pinkoviezky, I. D. Couzin, N. S. Gov, Collective conflict resolution in groups on the
move. *Physical Review E* **97** (2018).

53. S.-I. Amari, Homogeneous nets of neuron-like elements. *Biol. Cybernetics* **17**, 211–220
(1975).

54. S. Amari, Dynamics of pattern formation in lateral-inhibition type neural fields. *Biol.
Cybern.* **27**, 77–87 (1977).

55. K. S. Kakaria, B. L. de Bivort, Ring Attractor Dynamics Emerge from a Spiking Model of
the Entire Protocerebral Bridge. *Front. Behav. Neurosci.* **0** (2017).

56. L. Peliti, *Statistical Mechanics in a Nutshell* (Princeton University Press, 2011) (July 27,
2021).

57. J. J. Hopfield, Neural networks and physical systems with emergent collective
computational abilities. *PNAS* **79**, 2554–2558 (1982).

58. K. G. Wilson, The renormalization group and critical phenomena. *Rev. Mod. Phys.* **55**,
583–600 (1983).

59. E. Schneidman, M. J. Berry, R. Segev, W. Bialek, Weak pairwise correlations imply
strongly correlated network states in a neural population. *Nature* **440**, 1007–1012
(2006).

60. S. Takagi, R. Benton, Animal Behavior: A Neural Basis of Individuality. *Current Biology* **30**,
R710–R712 (2020).

Materials and Methods

We construct a simple, spatially-explicit model of neural decision-making to study how the

brain reduces choice in the presence of numerous spatial options (adapted from (52)).

Theoretical predictions obtained were then tested experimentally by exposing invertebrate

(fruit flies and desert locusts) and vertebrate systems (zebrafish) to spatial choice tests in virtual

reality. To identify unifying principles of spatiotemporal computation across scales of

biological organisation, we also reproduce the obtained decision-making patterns with an

established model of collective decision-making in animal groups.

Neural decision-making model. We construct a computational model of neural decision-

making that takes in a representation of directions to the different targets as input, and outputs

a collective vectorial representation of the agent's future velocity (adapted from (52)). This

provides us explicit predictions for animal trajectories, allows us to determine which target is

reached in each realization of the simulation, and facilitates direct comparison with

experimental tests. Our model is within the class of widely-employed neural ring-attractor

models (see SI Appendix), which like neural field models (53, 54), and attractor network

models more generally (15, 16, 55), consider the collective firing activity of the neurons, or the

firing rate, as opposed to the microscopic state of each firing neuron.

In our model, the brain is composed of individual components, called "spins", that, collectively,

as a "spin system", represent neural activity. Spin systems, which have been long-studied in

physics due to their ability to give insight into a wide range of collective phenomena, from

magnetic to quantum systems (56), were first introduced in the study of neurobiology by

Hopfield in a landmark paper (57) that provided considerable insights into principles

underlying unsupervised learning and associative memory. In its simplest (and most common)

formulation, as in Hopfield networks, a spin system is comprised of entities, spins, that can

each be in state 0 or 1, or in the terminology of physics either 'up' or 'down'. Spin systems

have consistently provided deep insights into complex collective phenomena, from spin and

molecular systems, to neural systems, undergoing phase transitions (58, 59) (see SI Appendix

for details and discussion).

Here, the animal's brain is characterized by a system of N spins. Each spin i encodes

direction to one of the presented goals $\hat{p}^\_i$, and exists in one of two states:

$\sigma_i=0$ or $\sigma_i=1$. We do not imply that a spin is equivalent to a neuron, but

rather, as we show via a mathematical derivation, that the collective properties of interacting

spins in our model is equivalent to the firing rate in the neural ring attractor model (see SI

Appendix for details). Consequently, we refer to the individual components with which we

model our system as "spins", and "neural activity" as a term to represent this "firing rate"

equivalent. The energy of the system (for any given configuration) is given by its Hamiltonian,

H.

548
$$H = -\frac{k}{N} \sum_{i \neq j} J_{ij} \sigma_i \sigma_j$$

549   where, k is the number of options available to the individual and J_{ij} is the interaction strength

550   between neurons i and j. Here, J_{ij} is given by

551
$$J_{ij} = \cos\left(\pi \left(\frac{|\theta_{ij}|}{\pi}\right)^{\nu}\right)$$

where, θ_{ij} is the angle between preferred directions of neurons i and j, and ν represents the

neural tuning parameter. For $\nu = 1$, the interactions become "cosine-shaped" $J_{ij} = \cos(\theta_{ij})$,

and the network has a Euclidean representation of space (Fig. S1). For $\nu < 1$, the network has

more local excitation and encodes space in a non-Euclidean manner (Fig. S1). System

dynamics are implemented by energy minimization using the Metropolis-Hastings algorithm

(similar to other Ising spin models) and the agent then moves with a velocity $\vec{V}$ determined by

the normalized sum of goal vectors $\hat{p}_i$ of all active neurons.

559
$$\vec{V} = \frac{v_0}{N} \sum_{i=1}^{N} \hat{p}_i \, \sigma_i$$

where v_0 is the proportionality constant. The goal vector $\hat{p}_i$ now points from the agent's

updated location to the neuron's preferred goal with directional noise chosen from a circularly

wrapped Gaussian distribution centered at 0 with a standard deviation σ_e. As in the mean-field

approximation of the model, the timescale of movement (defined by the typical time to reach

the target) in the numerical simulations was set to be much greater than the timescale of neural

firing (the typical time between two consecutive changes in the neural states σ_i).

Collective decision-making model. We reproduce results from our neural decision-making

model in a model that describes spatial decision-making at a different scale of biological

organization (refer (40) for methodological details). To highlight the features that are key to

producing the observed bifurcation patterns, we run simulations with and without feedback on

the strength of goal-orientedness of individuals.

Fly virtual reality experiments. All experiments were conducted on 3- to 5-day old female

wild-type CS strain *Drosophila melanogaster* raised at $26°C$ on a 12 hr light, 12 hr dark cycle.

Experiments were conducted in a flyVR setup procured from loopbio GmbH. 60 tethered

Drosophila melanogaster were exposed to either a two-choice or a three-choice decision task

(30 and 30 individuals, respectively) in the virtual reality environment. Each experimental trial

lasted 15 min where flies were exposed to five sets of stimuli—three experimental sets and two

control sets. The experimental stimuli sets consisted of two or three black cylinders (depending

on the experimental condition) that were presented to the animal in an otherwise white

environment. A control stimulus with a single pillar was presented before and after the

experimental conditions. We rotated all trajectories such that the x −axis points from the

origin, to the centre of mass of the targets. To visualise trajectories in the various experimental

conditions, we created time-normalised (proportion of maximum across a sliding time window)

density maps. We then folded the data about the line of symmetry, $y = 0$ and applied a density

threshold to the time-normalised density map. A piecewise phase transition function was then

fit to quantify the bifurcation.

588
$$y = \begin{cases} 0 & x \le x_c \\ A|x - x_c|^\alpha & x > x_c \end{cases}$$

where x_c is the critical point, α is the critical exponent, and A is the proportionality constant.

We also performed randomisation tests for each bifurcation where we conducted the exact fit

procedure described above to data where the trajectories were randomised by keeping the x-

coordinates, and swapping the y-coordinates with values from other random events.

Randomizations show that the resultant fit to our experimental data were highly significant

($p < 0.01$ for binary choice and $p < 10^{-4}$ for the three-choice case).

Based on the amount of time it took flies to reach one of the available targets, we also classified

individual fly tracks into one of two categories—direct tracks and non-direct tracks (60) (see

Fig. S11A,H for details). In our model, the direct tracks were also accounted for by varying the

directional tuning of spins. A high neural tuning (low v) results in more directed tracks (Fig.

S14).

Locust virtual reality experiments. All experiments were conducted on 156 instar 5 desert locusts (*Schistocerca gregaria*; 57 individuals for two-choice and 99 individuals for three-choice experiments, respectively) raised in the Animal Research Facility of the University of Konstanz. Based on our filtering criteria, 122 out of the 156 locusts were used in our analyses. Experiments were conducted in a locustVR setup procured from loopbio GmbH (27). The experimental procedure was identical to the one described above for flies, except now, each experimental trial lasted 48 min—three experimental sets (12 min each) and two control sets (6 min each). Analyzing bifurcations in locust trajectories using the same methods described above showed that the resultant bifurcations fit to our experimental data were highly significant ($p < 0.01$ for binary choice and $p < 10^{-4}$ for the three-choice case).

Similar to the flies, the locust trajectories were also classified as direct, or non-direct tracks. However, because the locustVR system allowed the animals to stop and reconsider movement during the decision-making process, we added an additional category to classification of individual locust tracks *viz.* the wandering tracks (see Fig. S12A,J for details).

Fish virtual reality experiments. All experiments were conducted on 1 cm ± 0.1 cm long zebrafish (*Danio rerio*) of age 24 to 26 days post-fertilisation raised in a room at 28 ℃ on a 16 hr light, 8 hr dark cycle. 440 fish were tested in total. Of these, 198 fish were exposed to decision-making with two virtual targets, 39 fish were exposed to decision-making with three equidistant virtual targets, and 50 fish were exposed to decision-making with three targets in asymmetric geometry (see SI Appendix for more details). Experiments were conducted in a fishVR setup procured from loopbio GmbH (refer (27) for details). Once a fish was introduced in the arena, it was given 20 min to acclimatize to the environment. This was followed by a 10 min control where it was presented a single virtual conspecific circling the arena in a circle of

radius 8 cm. After this, for experiments in symmetric geometries, the real fish was exposed to

choice experiments that lasted 90 min with the virtual fish initialized with random lateral

distances between them and random swim direction. To visualize the bifurcations, we

normalized (proportion of maximum) and stacked the marginal distributions along the direction

of the virtual fish's motion for various lateral distances. For experiments in asymmetric

geometries, the real fish was exposed to choice experiments where distance between the center

virtual fish and its closer neighbor was 0.03 m and its distance to the other neighbor was 0.09

m (Fig. 4). All experiments were conducted in accordance with the animal ethics permit

approved by Regierungspräsidium Freiburg, G-17/170.

Acknowledgements. We thank all members of the Department of Collective Behaviour who

assisted with the project: Renaud Bastien for modelling discussions and help setting up the VR

experiments, Guy Amichay for providing control data of two real fish swimming together, and

Paul Szyszka for showing V.H.S. how to tether flies. We thank the 'Itai Cohen Lab' for the

fruit fly image used in Fig. \ref{fig:insects}, Jitin Thomas for help with visualizations in Unity

and Andreas Poehlmann, John Stowers and Max Hofbauer from loopbio GmbH for technical

support with the VR systems. V.H.S., L.L., B.R.S. and I.D.C. are also grateful to the animal

care at the University of Konstanz including Christine Bauer, Jayme Weglarski and Dominique

Leo for help in conducting the experiments. They also acknowledge the efforts of the scientific

and technical staff at the University of Konstanz including Michael Mende, Markus Miller,

Mäggi Hieber Ruiz and Daniel Piechowski. V.H.S. acknowledges the International Max Planck

Research School (IMPRS) for Organismal Biology for the graduate school community and

access to courses and resources. I.D.C. acknowledges support from the NSF (IOS-1355061),

the Office of Naval Research grant (ONR, N00014-19-1-2556), the Struktur- und

Innovationsfonds für die Forschung of the State of Baden-Württemberg, the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy-EXC 2117-422037984 and the Max Planck Society. N.S.G. is the incumbent of the Lee and William Abramowitz Professorial Chair of Biophysics and acknowledges support by the Minerva Foundation (grant no. 712601). M.N acknowledges support from the Eötvös Loránd Research Network (a grant to the MTA-ELTE 'Lendület' Collective Behaviour Research Group, grant number 95152, and MTA-ELTE Statistical and Biological Research Group) and Eötvös Loránd University. Additionally, all authors thank two anonymous referees for their constructive comments during the review process.

Author contributions. V.H.S. and I.D.C. designed the study; V.H.S., D.G., T.S., N.S.G. and I.D.C. constructed the model; D.G. and N.S.G. constructed the mean-field approximation; V.H.S. and I.D.C. designed the fly experiments; V.H.S. conducted these experiments and analyzed the data with L.L. and M.N.; V.H.S., B.R.S. and I.D.C. designed the locust experiments; B.R.S. conducted these experiments and V.H.S. analyzed the data with M.N.; L.L. and I.D.C. designed the fish experiments; L.L. conducted these experiments and analyzed this data with V.H.S. and M.N.; V.H.S. and I.D.C. drafted the manuscript with significant contributions from all authors.

Competing interests. The authors declare that they have no competing interests.

Figures.

Fig. 1. Geometrical principles of two-choice and three-choice decision-making. (A) Schematic of the binary decision-making experiments. This simplified representation shows that a sharp transition in the animal's direction of travel is expected near a critical angle, θ_c. (B) A phase diagram describing the 'critical' transition exhibited while moving from compromise to decision between two options in space. The shaded area (also in E) represents the region in parameter space where both the compromise, and the decision solutions exist. (C) Density plot showing trajectories predicted by the neural model in a two-choice context. The axes represent $x-$ and $y-$coordinates in Euclidean space. The black line (also in G) presents a piecewise phase-transition function fit to the bifurcation. (D) Schematic of three-choice decision-making experiments, where the central target is on the angle bisector of the angle subtended by the other two targets. (E) A phase diagram describing the first 'critical' transition when the individual chooses among three options. Once the individual eliminates one of the outermost targets, it can decide between the two remaining options, similar to the two-choice phase diagram described in B. (F) Theoretical predictions for decision-making in a three-choice context. The dashed line (also in H) is the bisector of the angle subtended by center target and the corresponding side target on the first bifurcation point. See Table S1 for parameters used in C and F. (G) and (H) Density plots from experiments conducted with flies and locusts

690   choosing among two and three options, respectively. Note that the density plots presented here

691   are for the non-direct tracks, which constitute the majority type of trajectory adopted by both

692   flies and locusts (Figs. S11 and S12). However, our conclusions do not differ if we use all,

693   unfiltered, data (Figs. S11G,N and S12I,R).

Fig. 2. Decision-making for a larger number of targets. Density plots of simulated trajectories

for four- (A), five- (B), six- (C) and seven-choice (D) decision-making when targets are placed

equidistant and equiangular from the agent. Thee axes represent $x-$ and $y-$coordinates in

Euclidean space. Geometrical configurations are also varied to place the targets on the same

side of the agent (A and B) or in radial symmetry (C and D). See Table S1 for parameters used

in A–C. In D, all parameters used are identical except the system size $N = 70$.

Fig. 3. Decision-making in a moving frame-of-reference. (A) Schematic of the two-choice decision-making experiments conducted with larval zebrafish. In these experiments (also in the three-choice experiments depicted in D), the virtual fish swim parallel to each other while maintaining a fixed lateral distance, L between them. We only consider data where the real fish swims behind the virtual fish, i.e., it follows the virtual fish (see SI Appendix and Fig. S15 for details). (B) Normalized probability distribution (proportion of maximum) of simulated positions of an agent following two moving targets, and corresponding experiments (C) conducted with larval zebrafish following two virtual conspecifics. (D) Schematic representation of the three-choice decision-making experiments. (E) Normalized probability distributions of simulated positions of an agent following three moving targets, and corresponding experiments (F) conducted with larval zebrafish following three virtual conspecifics. See Table S1 for model parameters used in B and E.

Fig. 4. Decision-making with the targets in an asymmetric geometry. (A) Schematic of the
asymmetric choice test presented to larval zebrafish. In these experiments, the virtual fish
swim parallel to each other while maintaining a fixed lateral distance, L between them. To
create asymmetry in the geometry, the center fish swims closer to one of the side fish than the
other ($L_{12} = 0.09$ m and $L_{23} = 0.03$ m). (B) The upper panel shows the probability density
function of simulated positions of an agent following three moving targets in an asymmetric
geometry corresponding to the experiments. The simulated agent occupies a position of $y =$

724   ± 0.04 m while following the targets ($v = 0.7$; $\sigma_\theta = 0.3$). The lower panel shows the

725   probability density function of the position of the real fish along the axis perpendicular to its

726   direction of motion. As predicted by our model, the real fish considers the two virtual

727   conspecifics closer to each other as a single target and adopts one of two positions behind the

728   virtual fish.

Fig. 5. Consensus decision-making in simulations of animal groups follow the same geometrical principles. Results for two- (A) and three-choice (B) decision-making in a model of animal collectives. The density plots show trajectories adopted by the centroid of the animal group for 500 replicate simulations where the groups don't split. The axes represent $x-$ and $y-$coordinates in Euclidean space. The black lines show a piecewise phase-transition function fit to the trajectories. For the three-choice case (B), the dashed line is the bisector of the angle subtended by center target and the corresponding side target on the first bifurcation point. See Table S2 for parameters used.