<u>Comparison of linear combination modeling strategies for GABA-edited MRS</u>
<u>at 3T</u>

Helge J. Zöllner[1,2], Sofie Tapper[1,2], Steve C. N. Hui[1,2], Peter B. Barker[1,2], Richard A. E. Edden[1,2],

Georg Oeltzschner[1,2,*]

[1] *Russell H. Morgan Department of Radiology and Radiological Science, The Johns Hopkins*

University School of Medicine, Baltimore, MD, United States

[2] *F. M. Kirby Research Center for Functional Brain Imaging, Kennedy Krieger Institute, Balti-*

more, MD, United States

*Corresponding author:

Georg Oeltzschner, Ph.D.

Division of Neuroradiology, Park 367G

The Johns Hopkins University School of Medicine

600 N Wolfe St

Baltimore, MD 21287

goeltzs1@jhmi.edu

Running title: Linear combination modeling of GABA-edited MRS

Word count: 7239

Graphical Abstract

102 strategies to model GABA-edited MRS with linear combination modeling were evaluated to quantify GABA and GABA+ in Osprey. Significantly different GABA and GABA+ estimates were found when a well-parameterized macro-molecule at 3 ppm was included. The findings suggest that linear combination modeling needs to be adapted for quantification of GABA-edited MRS.

Abstract

Abbreviations

γ-aminobutyric-acid – GABA; Linear combination modeling – LCM; macromolecule – MM; GABA + MM - GABA+; homocarnosine – HCar; glutamate – Glu; glutamine – Gln; glutathione – GSH; N-acetylaspartylglutamate – NAAG; N-acetylaspartate – NAA; Hankel singular value decomposition – HSVD; full-width at half-maximum – FWHM; creatine – Cr; negative creatine methylene – -CrCH$_2$; phosphocreatine – PCr; SD – standard deviation; Akaike Information Criterion – AIC; coefficients of variation – CVs;

Abbreviations

γ-aminobutyric-acid – GABA; Linear combination modeling – LCM; macromolecule – MM; GABA + MM - GABA+; homocarnosine – HCar; glutamate – Glu; glutamine – Gln; glutathione – GSH; N-acetylaspartylglutamate – NAAG; N-acetylaspartate – NAA; Hankel singular value decomposition – HSVD; full-width at half-maximum – FWHM; creatine – Cr; negative creatine methylene – -CrCH$_2$; phosphocreatine – PCr; SD – standard deviation; Akaike Information Criterion – AIC; coefficients of variation – CVs;

<u>Keywords</u>

40   • Magnetic resonance spectroscopy

41   • Linear combination modeling

42   • GABA-edited MEGA-PRESS

Abstract

Purpose

J-difference-edited spectroscopy is a valuable approach for the in vivo detection of γ-aminobu-

tyric-acid (GABA) with MRS. A recent expert consensus article recommends linear combination

modeling (LCM) of edited MRS, but does not give specific details of implementation. This

study explores different modeling strategies to adapt LCM for GABA-edited MRS.

Methods

61 medial parietal lobe GABA-edited MEGA-PRESS spectra from a recent 3T multi-site study

were modeled using 102 different strategies combining six different approaches to account for

co-edited macromolecules, three modeling ranges, three baseline knot spacings, and the use of

basis sets with or without homocarnosine. The resulting GABA and GABA+ estimates (quanti-

fied relative to total creatine), the residuals at different ranges, SDs and CVs, and Akaike infor-

mation criteria, were used to evaluate the models' performance.

Results

Significantly different GABA+ and GABA estimates were found when a well-parameterized

MM_{3co} basis function was included in the model. The mean GABA estimates were significantly

lower when modeling MM, while the CVs were similar. A sparser spline knot spacing led to

lower variation in the GABA and GABA+ estimates, and a narrower modeling range – only in-

cluding the signals of interest – did not substantially improve or degrade modeling performance.

Additionally, results suggest that LCM can separate GABA and the underlying co-edited MM_{3co}.

Incorporating homocarnosine into the modeling did not significantly improve variance in

GABA+ estimates.

Conclusion

GABA-edited MRS is most appropriately quantified by LCM with a well-parameterized co-ed-

ited MM_{3co} basis function with a constraint to the non-overlapped $MM_{0.93}$, in combination with a

sparse spline knot spacing and a modeling range between 0.5 and 4 ppm.

Introduction

A recent expert consensus paper recommended that linear combination modeling (LCM) should be used for the quantification of edited MRS data[1] , stating that standard fitting approaches originally optimized for short-TE MRS should be adapted for edited MRS. Further, it was recommended that quantum-mechanical simulations should be used to confirm the co-edited profile of all metabolites in the edited spectrum, and contributions from macromolecule (MM) signals should be specified. Despite these recommendations, little detail was given regarding several unique features of edited spectra, and how they should be appropriately modeled. These features include:

1) The MEGA-PRESS experiment is well-known to co-edit MM signals with coupled spins at 1.7 and 3 ppm, causing substantial contamination of the edited GABA signal, and forcing researchers to report the composite measure GABA+MM (GABA+)[1]. Because the co-edited MM signal is poorly characterized, there is currently no consensus or recommendation on how to appropriately account for it during spectral modeling. Instead, the most widely used analysis algorithms implement entirely different strategies to fit the composite 3-ppm signal. For example, the Gannet software uses a single Gaussian model[2], while a double-Gaussian is used in Tarquin[3], and LCModel[4] defaults to a basis set that only includes the GABA basis function.

2) Another co-edited compound contributing to the 3 ppm signal is homocarnosine (HCar), a dipeptide of GABA and histidine. While the 3 ppm multiplets of GABA and homocarnosine are separated by just 0.05 ppm (which are therefore unlikely to be successfully separated), inclusion of a homocarnosine basis function may be warranted based on its reported concentration in vivo (~0.5 mmol/kg [5], compared to ~1-2 mmol/kg for GABA), but it has not been investigated whether doing so has a stabilizing or destabilizing effect on the modeling[6].

3) Unedited spectra are typically modeled over a restricted frequency-domain range covering the visible upfield peaks, including macromolecular and lipid resonances between 0 and 1 ppm, but usually avoiding the water suppression window above ~4 ppm. The choice of frequency-domain modeling range for edited spectra is less obvious. Since

the main advantage of spectral editing is the isolation of a single target resonance,

modeling signals outside the immediate surrounding of the target may dilute the resolving

power of editing. On the other hand, increasing the modeling range may offer useful

constraints to stabilize the solution of the modeling problem. The difference is

highlighted by the different strategies encountered in common software tools – while the

Gannet software fits the GABA-edited difference spectrum over a narrow range (only

including the 3-ppm GABA+ and 3.75 ppm glutamate and glutamine peaks), the

LCModel recommendation is to include the strong co-edited signals from glutamate

(Glu), glutamine (Gln), glutathione (GSH), N-acetylaspartylglutamate (NAAG), and N-

acetylaspartate (NAA), which heavily overlap with GABA around 2.25 ppm. The effects

of limiting the modeling range have not been assessed systematically to date.

4) Linear combination modeling methods commonly include terms to account for smooth

baseline curvature, usually parametrized from cubic B-spline or polynomial functions, or

by smoothing residuals. The flexibility of the baseline model substantially affects

metabolite estimates from unedited spectra[7]; while baseline terms are necessary to

account for e.g. lipid contamination, poor water suppression etc., they are potential

sources of overfitting if awarded too many degrees of freedom. Baseline modeling may

have an even greater influence when modeling difference spectra, since only *co-edited*

lipid and MM signals contribute to the smooth background variation. Importantly, the co-

edited MM background of the GABA-edited difference spectrum has not been

appropriately characterized (e.g., through metabolite-nulled acquisition), suggesting that

the choice of baseline flexibility can drastically influence modeling results through two

highly susceptible regions of the spectrum. First, in the absence of an appropriate model

for the co-edited broad MM signal at 3 ppm, this signal may be absorbed into the baseline

depending on its flexibility. Second, strong MM and lipid signals in the region between

0.5 and 2.5 ppm may be affected by the 1.9 ppm editing pulse (either directly through

saturation or indirectly through coupling), likely leading to an unknown, but substantial,

MM contribution in this spectral region[8,9]. This is especially important considering that

the co-edited signals from NAA, NAAG, Glu, Gln, and GSH overlap with GABA in this

region. Overly rigid baselines may provide insufficient flexibility to capture these signals,

in turn compromising the accuracy of the estimation of the co-edited metabolites.

The aim of this study was to evaluate different strategies for linear combination modeling of GABA-edited MEGA-PRESS difference spectra, and to establish initial 'best practices' substantiating the recommendations of the expert consensus on spectral editing. To this end, different approaches to account for co-edited MM signals, various modeling ranges and baseline knot spacings, as well as the inclusion of homocarnosine were compared. In the absence of a 'gold standard', the performance of each modeling strategy was assessed by comparing descriptive statistics of the metabolite estimates, calculating the Akaike information criteria, and assessing the fit residuals.

Methods

Study participants & data acquisition

In this study, 61 publicly available GABA-edited MEGA-PRESS datasets originating from 7 sites from a recent 3T multi-center study[10] were analyzed (see **Supplementary Material 1** for subject list). All datasets were acquired on Philips 3T scanners with the following acquisition parameters: TR/TE = 2000/68 ms; 320 excitations (10m 40s scan time); 16-step phase-cycle; 2 kHz spectral width; 2000 samples; 27-ml cubic voxel volume in the medial parietal lobe. For this heuristic approach of exploring the GABA modeling the data homogeneity (SNR, FWHM, tissue composition, and absence of fat contamination) was increased while reducing the overall number of subjects by including only 61/298 subjects of the original dataset[10]. All sites except for P8 used a similar sequence implementation with interleaved water referencing for prospective frequency correction[11]. For the edit-ON transients, the editing pulses with 15 ms pulse duration and 82.5 Hz inversion bandwidth (FHWM) were applied at a frequency of 1.9 ppm to refocus the coupling evolution of the GABA spin system. For the edit-OFF transients, the editing pulses were applied at a frequency of 7.5 ppm. Edit-ON and edit-OFF transients were acquired in alternating order. An additional water reference scan was acquired for each dataset using interleaved water referencing [11], i.e. one excitation with water suppression and editing pulses deactivated every 40 water-suppressed excitations (total of 8 averages).

Data pre-processing

Data were analyzed in MATLAB using Osprey[12,13] (v.1.0.1.1), a recently published open-source MRS analysis toolbox. Raw data were eddy-current-corrected [14] based on the water reference, and individual transients were aligned separately within the edit-ON and edit-OFF conditions using the robust spectral registration algorithm[15]. Averaged edit-ON and edit-OFF spectra were aligned by optimizing relative frequency and phase such that the water signal in the difference spectrum was minimized. The final difference spectra for quantification were generated by subtracting the edit-OFF from the edit-ON spectra. Finally, any residual water signal was removed with a Hankel singular value decomposition (HSVD) filter[16] to improve data quality in the edit-OFF spectra and to reduce residual baseline roll in the difference spectra.

Basis set

The basis set used for modeling was generated from a fully localized 2D density-matrix simula-

tion of a 101 x 101 spatial grid (voxel size: 30 mm x 30 mm x 30 mm; field of view: 45 mm x 45

mm x 45 mm) implemented in a MATLAB based simulation toolbox FID-A [17], using vendor-

specific refocusing pulse shape and duration, sequence timings, and phase cycling. It contains 17

metabolite basis functions (ascorbate, aspartate, creatine (Cr), negative creatine methylene (-

$CrCH_2$), GABA, glycerophosphocholine, GSH, Gln, Glu, water, myo-inositol, lactate, NAA,

NAAG, phosphocholine, phosphocreatine (PCr), phosphoethanolamine, scyllo-inositol, and tau-

rine) and 8 Gaussian MM and lipid resonances ($MM_{0.94}$, $MM_{1.22}$, $MM_{1.43}$, $MM_{1.70}$, $MM_{2.05}$,

Lip09, Lip13, Lip20, details in **Supplementary Material 2** with similarly defined parametriza-

tion as described in the LCModel software manual[18]) for the edit-OFF spectrum.

For the difference spectrum, $MM_{0.94}$ and the co-edited macromolecular signal at 3 ppm (MM_{3co})

were parametrized as Gaussian basis functions ($MM_{0.94}$: 3-proton signal; chemical shift 0.915

ppm, full-width at half-maximum (FWHM) 11 Hz; MM_{3co}: 2-proton signal; chemical shift 3

ppm; FWHM 14 Hz). The MM_{3co} amplitude was defined under the assumptions of a pseudo-dou-

blet GABA signal at 3 ppm and the MM_{3co} contribution to the 3-ppm GABA peak to be around

50%[1,6,8,19]. The optimum FWHM used to parametrize the MM_{3co} basis function was determined

to be 14 Hz by fitting the mean difference spectrum of all datasets with a composite GABA+ ba-

sis function (GABA + MM_{3co}) with varying FHWM (between 1 and 20 Hz). The parameterized

Gaussian MM_{3co} basis function was integrated into the modeling process using different assump-

tions and constraints described in the following paragraphs.

Linear combination modeling of GABA-edited difference spectra

193
Osprey's frequency-domain linear combination model was used to determine the metabolite esti-

mates. Model parameters include metabolite basis function amplitudes, frequency shifts,

zero/first order phase correction, Gaussian and Lorentzian linebroadening, and cubic spline base-

line coefficients. All parameters are determined by Levenberg-Marquardt[20,21] non-linear least-

squares optimization, using a non-negative least-squares (NNLS) fit [22–24] to determine the metab-

olite amplitudes and baseline coefficients at each iteration of the non-linear optimization.

Amplitude ratio soft constraints are imposed on MM and lipid amplitudes, as well as selected

pairs of metabolite amplitudes, as defined in the LCModel manual[4,18]. The strength factor of the

amplitude ratio soft constraint λ is set to 0.05 by default.

Figure 1 – Different linear combination modeling strategies for GABA-edited spectra. (A) Different co-edited MM3co modeling approaches derived from a Gaussian function at 3.0 ppm (B)

All combinations of basis set composition, modeling range, spline knot spacing, and MM$_{3co}$ modeling leading to 102 different modeling strategies.

A range of modeling strategies for the GABA-edited difference spectrum was included in this

study, covering various aspects of the modeling process (**Figure 1**). The different parametriza-

tions and soft constraints to account for the co-edited MM$_{3co}$ signal are shown in **Figure 1A**. All

possible combinations for the modeling strategies: i) inclusion of homocarnosine in the basis set;

ii) different parametrizations and soft constraints to account for the co-edited MM$_{3co}$ signal; iii)

different modeling ranges and iv) different baseline spline knot spacings are tabulated graph-

ically in **Figure 1 B**. Each modeling aspect is described in detail below:

<u>*Including homocarnosine in the basis set*</u>

To assess the effects of including homocarnosine in the linear combination model, we repeated

all analysis steps with two different basis sets: the default Osprey basis set *with* and *without* an

additional HCar basis function. Chemical shift and scalar coupling parameters describing the

HCar spin system were taken from literature[6].

<u>*Varying the modeling range and baseline knot spacing*</u>

Two aspects of linear combination modeling are suggested to have a considerable influence on

metabolite estimates[7,25]. First, the choice of the modeling range, i.e., the frequency interval that

defines the part of the frequency-domain spectrum that is considered to calculate the least-

squares difference between model and data. Second, the baseline knot spacing, i.e., the frequency

difference between two adjacent knots of the cubic spline basis that is used to approximate the

smooth baseline.

Three different modeling range scenarios were considered, reflecting common choices in the lit-

erature and widely used software tools: a) a wide modeling range typically used to analyze uned-

ited spectra, including all signals in the GABA-edited difference spectrum (0.5 to 4 ppm –

"wide"); b) an intermediate modeling range excluding signals below 1.9 ppm (e.g. co-edited li-

pids and macromolecules), but including strong co-edited signals from NAA, NAAG, Glu, Gln,

and GSH (1.85 and 4.1 ppm, *"intermediate"*), similar to the range recommended in LCModel's

dedicated 'mega-press-3' option; and c) a narrow modeling range only including the co-edited

signals from GABA+ and Glx (2.79 – 4.2 ppm, *"narrow"*), the default modeling range in Gan-

net[2].

Three spline knot spacings were included in the analysis, with 0.4 ppm being the default Osprey

option, shown to create reproducible and comparable metabolite estimates for conventional MRS

[26], as well as sparser (0.55 ppm) and denser (0.25 ppm) spline knot spacings.

238
239
Co-edited macromolecule models
Seven different strategies to model the GABA-edited difference spectrum were implemented

(**Figure 1 A**). The trivial approach – not accounting for the co-edited signal MM_{3co} at all – is la-

beled **none**. The other six modeling strategies all include a dedicated parametrized Gaussian

MM_{3co} basis function. This basis function is given different degrees of freedom in the different

strategies, e.g. hard- or soft-constrained relative to the amplitude of the GABA or the $MM_{0.94}$ ba-

sis functions, and with a fixed or free width. Here, strategies with fewer degrees of freedom re-

flect the frequently made assumption that the GABA-to-MM ratio (and the MM background it-

self) is relatively stable across subjects and anatomical region, and assumed to be known, while

strategies with more degrees of freedom or soft constraints relax these assumptions:

• The **GABA_hard** model uses a single composite GABA+MM basis function by adding the

GABA and MM_{3co} (initial FWHM of the basis function = 14 Hz) basis functions with a fixed

1:1 amplitude ratio. The 1:1 ratio reflects the widely used empirical assumption that 50% of

the 3-ppm signal in a conventional GABA-edited difference spectrum can be attributed to co-

edited macromolecules[6,19].

• The **GABA_soft** model uses separate GABA and MM_{3co} (initial FWHM of the basis function =

14 Hz) basis functions and imposes a soft constraint on the ration of the amplitudes of both

basis functions during the optimization (1:1 ratio).

• The **Gauss_fixed** model uses separate GABA and MM_{3co} (initial FWHM of the basis function =

14 Hz) basis functions. No further constraints are imposed. This means possible changes in

the contributions to the 3-ppm GABA peak are modeled.

• The **Gauss_free** model uses separate GABA and MM_{3co} basis functions. In contrast to the

Gauss_fixed model, the FWHM of the Gaussian MM_{3co} signal is represented by an additional

model parameter. This means that the MM_{3co} basis function itself is not static, but

dynamically modified during optimization.

• The **MM09$_{hard}$** model uses separate GABA and MM basis functions. The MM_{3co} basis

function is replaced by a composite $MM_{0.94} + MM_{3co}$ basis function (i.e., the $MM_{0.94}$ (initial

FWHM of the basis function = 11 Hz) and MM_{3co} (initial FWHM of the basis function = 14

Hz) basis functions are added in a 3:2 ratio). Resulting in a single composite basis function

for $MM_{0.94}$ and MM_{3co}, adapted from the soft constraint model described in literature [9].

• The **MM09$_{soft}$** model uses separate GABA, $MM_{0.94}$ and MM_{3co} basis functions. In contrast to

the MM09$_{hard}$ model, soft constraints enforce a ~3:2 amplitude ratio for the $MM_{0.94}$ and

MM_{3co} amplitudes during optimization. Resulting in two separate basis functions for $MM_{0.94}$

and MM_{3co}, which is similar to previously described implementations [9].

The models MM09$_{hard}$ and MM09$_{soft}$ [27] as well as Gauss$_{fixed}$ [28] correspond to models previously

investigated using the LCModel software and the amplitude assumptions were derived empiri-

cally. It is worth noting that each basis function receives a separate Lorentzian linebroadening,

frequency shift, and amplitude parameter during the optimization. For the Gauss$_{free}$ model, the

MM_{3co} basis function is dynamically updated as an explicit modeling parameter during the opti-

mization, therefore the MM_{3co} basis function has effectively two separate parameters to account

for its linewidth (the Lorentzian linebroadening term and the FWHM of the MM_{3co} basis func-

tion). Finally, the composite models (GABA$_{hard}$), which do not have separate GABA and MM

functions, only have one linebroadening, one frequency, and one amplitude parameter compared

to twice the parameters for the soft constraint counterparts.

Combining the various MM_{3co} models (5 + 2 that were used for the wide modeling range only),

modeling ranges (3), baseline spline knot spacings (3), and basis sets (2), a total of 102 different

modeling strategies were investigated in this study. All models were implemented in Osprey[12]

and are available on GitHub[13].

Quantification, visualization, and statistics

Quantification

For the basis set without homocarnosine, GABA refers to the model amplitude estimate for the GABA basis function, which is of course only available for the modeling strategies with separate basis functions for GABA and MM_{3co} (none, $GABA_{soft}$, $Gauss_{fixed}$, $Gauss_{free}$, $MM09_{soft}$). GABA+ refers to the sum of the amplitude estimates for GABA and MM_{3co} ($GABA_{soft}$, $Gauss_{fixed}$, $Gauss_{free}$, $MM09_{hard}$, $MM09_{soft}$) or the amplitude estimate for the composite basis function including both MM and GABA ($GABA_{hard}$) and is therefore calculated for all strategies with an explicit MM_{3co} model. For comparison, the GABA amplitude for the `none` strategy is included in the figures reporting GABA + MM_{3co}. However, it still refers to a GABA-only estimate.

For the basis set that included homocarnosine (HCar), the difference in GABA and MM_{3co} estimates between the modeling strategies with and without HCar (ΔGABA and ΔMM_{3co}, respectively) were investigated to evaluate whether the inclusion of HCar has a systematic effect on the estimation of those signals with which it overlaps. All estimates were quantified relative to the total creatine (Cr + PCr) amplitude from the edit-OFF spectrum with the wide modeling range and a spline knot spacing of 0.4 ppm. Differences in GABA(+)/tCr between modeling strategies are therefore only related to the modeling of the difference spectra, but not to the reference compound modeling. No further tissue or relaxation corrections were applied.

Further, the relative contributions of MM_{3co} to the GABA+ estimate and the relative contributions of HCar to the sum of GABA+ and HCar estimate were calculated.

Visualization

The modeling performance and systematic characteristics of each modeling strategy were visually assessed through the mean spectra, mean fit, mean residual, and mean models of GABA+, GABA, MM_{3co}, HCar (if included) and the baseline, i.e., averaged across all datasets.

The metabolite estimate distributions were visualized as violin plots including boxplots with median, 25^{th}/75^{th} quartile ranges, and smoothed distributions to identify systematic differences between modeling strategies. In addition, the mean value of the 'none' model across the three spline knot spacings was added for each modeling range as a dashed horizontal line. Bar plots

were created to visualize quality metrics, including the standard deviation if appropriate. All

plots were generated with R[29] (Version 3.6.1) in RStudio (Version 1.2.5019, RStudio Inc.) using

SpecVis[26,30], an open-source package to visualize linear combination modeling results with the

ggplot2 package[31]. All scripts and results are publicly available[32].

Statistics

Significant differences in the mean and the variance of the GABA, GABA+, and MM_{3co} esti-

mates were assessed between all modeling strategies. The statistical tests were set up as paired

without any further inference. Differences of variances were tested with Fligner-Killeen's test,

with a post-hoc pair-wise Bonferroni-corrected Fligner-Killeen's test. The means were compared

with an ANOVA or a Welch's ANOVA, depending on whether variances were different or not.

Post-hoc analysis was performed with a paired t-test with equal or non-equal variances, respec-

tively.

Additionally, Pearson's correlation was used to investigate the impact of including HCar in the

basis set. The strength of the correlation was considered substantial for $R > 0.25$.

<u>Model evaluation criteria</u>

The performance of each modeling strategy was evaluated in different ways, including the im-

pact of the different modeling strategies on the GABA, GABA+ , and MM_{3co} estimates, as well

as several quality measures:

1) Visual inspection: Mean model, residual, and baseline were assessed for characteristic

341   features.

342   2) SD fit quality: The SD of the residual was determined, and then normalized by the noise

343   level (calculated as the SD of the noise between -2 and 0 ppm). This is done over the

344   entire modeling range of the difference spectrum and termed **residual**_{SD range}.

345   3) Amplitude fit quality: the difference between the maximum and minimum of the residual

346   was determined, and then normalized by the noise level [25] (similarly calculated as in the

347   second criterion).This was done over the entire modeling range of the difference

348   spectrum and termed **residual**_{ampl range}.

4) Amplitude 3-ppm peak fit quality: Similar to the third criterion, the residual was

calculated over the range of 3.027 ± 0.15 ppm to assess the fit quality of the 3-ppm

GABA peak and termed **residual$_{ampl\ 3ppm}$**.

5) Consistency of metabolite estimates: The across-subject coefficients of variation (CV =

SD/mean) for all metabolite estimates (GABA/tCr, GABA+/tCr) were calculated for each

modeling strategy.

6) Akaike Information Criterion (AIC): The Akaike information criterion [33], which takes the

number of model parameters into account, is defined as follows:

357
$$logLikelihood_i = -0.5 * N_i * (\log(2*\pi) + 1 - \log(N_i) + \log(SSE_i))$$

359
$$AIC_i = -2 * \frac{1}{N_i} * logLikelihood_i + 2K_i$$

Here, N_i is the number of points in the modeling strategy i, SSE_i is the sum of squared

error (i.e., squared residual) of that strategy, and K_i is the number of free model parame-

ters for that strategy. The $logLikelidhood_i$ was divided by the number of points N_i to

reduce the strong weighting of the datapoints and to make the AIC_i values comparable

for different modeling ranges. Soft constraint model parameters were included with a

value of 0.5. Lower AIC_i values indicate a more appropriate model. Subsequently, $\boldsymbol{\Delta AIC_i}$

scores were calculated as the difference of AIC_i of modeling strategy i and the model

with the lowest AIC_{min}:

368
$$\Delta AIC_i = AIC_i - AIC_{min}$$

Results

All 61 datasets were successfully processed and modeled with all 102 modeling strategies. No

data were excluded from further analysis. The data quality assessment indicated consistently

high spectral quality for all spectra ($NAA_{SNR} = 272 \pm 70$; $NAA_{FWHM} = 5.29 \pm 1.09$ Hz) without

lipid contamination (data overview in **Supplementary Material 4**). Other data quality measures

were extracted from the Gannet[2] analysis performed in recent multi-site studies[10,34]. For the

Philips-only subset of datasets in the present study, the tissue composition ($fGM = 0.60 \pm 0.04$;

$fWM = 0.27 \pm 0.03$; $fCSF = 0.13 \pm 0.04$) and across-subject CV ($GABA+/Cr = 9.99\%$) indicate

consistency in the dataset and the modeling. across-subject CV was interpreted as a measure of

modeling performance, assuming that increased CVs are mainly introduced by variability in the

modeling and do not reflect biologically meaningful variance of GABA+ estimates.

Summary and visual inspection of the modeling results

Figure 2 – Mean modeling results for all modeling strategies without homocarnosine. A substantial structured residual is apparent at 3 ppm if no MM modeling strategy is included. All three modeling ranges (columns), three spline knot spacings (rows), and MM$_{3co}$ model (color-coded) are presented with mean residuals and fits, as well as the GABA+, GABA, MM$_{3co}$, and spline baseline models. The mean data is included in black. The dashed lines indicate the range of the residual across one row. The arrows indicate the range of values for a specific modeling range and spline knot spacing with the color corresponding to the MM$_{3co}$ model with minimum/maximum value.

Figure 2 shows the mean modeling results for all modeling strategies without homocarnosine.

Not including MM_{3co} leads to a substantial structured residual around 3 ppm for all knot spacings

and modeling ranges. In contrast, all modeling strategies with MM_{3co} appear to reflect the line-

shape of the 3-ppm signal more accurately, with very similar results for the complete fit (metabo-

lites, MMs, and baseline) and the individual components. Modeling strategies with the interme-

diate and wide modeling range further show strong residuals around 2 ppm, suggesting slightly

inaccurate lineshape modeling of the methyl singlets from NAA and NAAG, or inaccurate mod-

eling of co-edited MM signals in this region. Structured residuals appear also in the region of the

3.75 ppm Glx signals, although they are much less pronounced in strategies with the narrow

modeling range, suggesting that including the 2.25 ppm multiplets (and underlying baseline fluc-

tuation) has a considerable impact on phase estimation.

In general, the residuals are consistent between different MM_{3co} models for any given knot spac-

ing and modeling range. Notably, residuals tend to be smaller on an absolute scale for denser

knot spacing and narrower modeling range.

Mean GABA models agree well between all strategies with a separate MM_{3co} model. The

$GABA_{hard}$ strategy appears to produce a larger signal as its GABA basis function includes the

MM_{3co} signal, but does not model it separately, while the strategies that do so produce compara-

ble mean MM_{3co} models.

The mean baseline is consistently flatter around 3 ppm for modeling strategies with an explicit

MM_{3co} model, while absorbing substantially more signal for the 'none' approach without an MM

model. This behavior is particularly obvious for the dense knot spacing (0.25 ppm) over the wide

modeling range. Baseline curvature generally increases for denser knot spacings around 2.2 ppm

for the intermediate and wide range.

Metabolite level distribution

Figure 3 – Distribution and across subject coefficients of variation (CVs) of GABA+ estimates for all modeling strategies. Including a MM_{3co} model significantly increases the mean estimates for all modeling strategies, while giving similar or reduced CVs. The mean estimates across the three spline knot spacings of the 'none' approach are indicated as a dashed line for each modeling range. All three modeling ranges (column) and three spline knot spacings (within each column), and co-edited MM models (color-coded) are presented. Distributions are shown as half-violins (smoothed distribution), box plots with median, interquartile range, and $25^{th}/75^{th}$ quartile. The median lines of the box plots are connected to visualize trends within a specific baseline knot spacing. CVs are summarized as bar plots. Minimum/maximum CVs for each modeling range are indicated as downwards/upwards triangle in the color corresponding to the MM_{3co} model. Minimum/maximum CVs for each baseline knot spacing within a specific modeling range are reported on the right side of each column. Global minimum and maximum CVs across all models are added as text.

Figure 3 shows distributions and coefficients of variation (CVs) of the GABA+ estimates for all modeling strategies. Table 1 summarizes the mean and SD GABA/GABA+ estimates as well as the statistics. GABA+ estimates are significantly higher than GABA-only estimates of the 'none' modeling strategy for all modeling ranges and knot spacings, supporting the notion from **Figure 2** that not including an MM model leaves a considerable fraction of the edited 3-ppm signal un-modeled, resulting in substantial residuals or increased baseline amplitude flexion.

Notably, all modeling strategies with MM_{3co} return comparable mean estimates and CVs within the same knot spacing (see Minimum/Maximum column of **Figure 3**). In addition, sparser knot spacing leads to lower CVs. The intermediate modeling range does not appear to perform more consistently than both other modeling ranges.

Table 1 – GABA+ mean and SDs for all modeling strategies (ratios to tCr). Significant differences (p < .05) between the corresponding model and the 'none' are shaded in gray.

modeling range		narrow			intermediate			wide		
knot spacing (ppm)		0.55	0.4	0.25	0.55	0.4	0.25	0.55	0.4	0.25
none	[GABA]	0.22 *.062*	0.213 *.071*	0.193 *.057*	0.297 *.048*	0.186 *.061*	0.204 *.044*	0.284 *.029*	0.288 *.034*	0.158 *.066*
GABA$_{hard}$	[GABA+]	0.328 *.041*	0.291 *.04*	0.312 *.046*	0.414 *.048*	0.296 *.062*	0.329 *.041*	0.389 *.041*	0.414 *.047*	0.3 *.068*
GABA$_{soft}$	[GABA+]	0.298 *.029*	0.267 *.036*	0.295 *.03*	0.417 *.053*	0.279 *.058*	0.323 *.031*	0.375 *.039*	0.397 *.042*	0.271 *.035*
Gauss$_{fixed}$	[GABA+]	0.306 *.04*	0.272 *.037*	0.304 *.034*	0.412 *.052*	0.298 *.055*	0.332 *.04*	0.369 *.05*	0.382 *.052*	0.343 *.04*
Gauss$_{free}$	[GABA+]	0.305 *.032*	0.277 *.031*	0.3 *.035*	0.409 *.052*	0.3 *.058*	0.331 *.04*	0.364 *.05*	0.382 *.045*	0.341 *.043*
MM09$_{hard}$	[GABA+]	-	-	-	-	-	-	0.396 *.039*	0.422 *.055*	0.345 *.084*
MM09$_{soft}$	[GABA+]	-	-	-	-	-	-	0.367 *.047*	0.386 *.045*	0.337 *.053*

444   <u>Model evaluation</u>

Figure 4 – Evaluation of all modeling strategies. Comparably low residual$_{ampl\ ange}$ are related to the high data quality without artifacts between 0.5 and 2 ppm. Including an MM$_{3co}$ model reduces the 3-ppm residual by ~30% without significant impact on the ΔAIC. All three modeling ranges (column) and three spline knot spacings (within each column), and co-edited MM models (color-coded) are presented. Bar plots represent mean values; SD is indicated by whiskers where appropriate. Minimum/maximum values for each modeling range are indicated as downwards/upwards triangle in the color corresponding to the MM$_{3co}$ model.

Minimum/maximum values for each baseline knot spacing within a specific modeling range are reported on the right side of each column. Global minimum and maximum values across all models are added as text.

Figure 4 summarizes the metrics used for model evaluation. The residual over the modeled fre-

quency range (residual$_{\text{SD range}}$ and residual$_{\text{ampl range}}$) is lowest for the narrow modeling range. For

the intermediate and wide modeling ranges, residual$_{\text{ampl range}}$ is substantially higher, largely driven

by the 2-ppm region (see also **Figure 2**). Consequentially, residual$_{\text{ampl range}}$ is comparable be-

tween MM modeling strategies for a given knot spacing (see Minimum/Maximum column of

Figure 4).

The residual around the GABA+ peak (residual$_{\text{ampl 3ppm}}$) is consistently reduced by up to 30% if a

MM$_{\text{3co}}$ model is included, in line with the reduction of structured residual in **Figure 2**. This ef-

fect is less pronounced for the dense knot spacing (0.25 ppm), indicating that a flexible baseline

is to some degree capable of accounting for otherwise unmodeled MM signal. Together, these

findings again support the notion that omitting an explicit MM$_{\text{3co}}$ model does not capture the

whole edited 3-ppm signal, which remains unmodeled (in the residual) or gets partially absorbed

by the baseline or interpreted incorrectly as GABA signal.

The strategy with the lowest AIC is the 'none' model with the intermediate modeling range and

sparse knot spacing, reflecting the low number of model parameters: there is no separate basis

function for MM, and the low number of splines. The ΔAIC (the difference between the lowest

AIC and the individual model's AIC) consequently increases for larger modeling ranges, as more

splines are included. Similarly, ΔAIC increases for denser knot spacings, and in fact, this in-

crease is much stronger compared to the resulting reduction in both residual measures, suggest-

ing that the increased flexibility and reduction of the residual does not justify the greater number

of model parameters.

For any given knot spacing and modeling range, ΔAIC values are comparable between MM$_{\text{3co}}$

models, with moderate increases when more parameters are estimated. Together with its low CV

(9.8% compared to the minimum CV value 9.7% for the GABA$_{\text{soft}}$ value with a narrow fit range)

for GABA+, the ΔAIC for the MM09$_{\text{hard}}$ model over the wide modeling range with sparse knot

spacing (ΔAIC = 3.1) indicates a good performance of this particular model without introducing

overfitting. Despite the slightly higher ΔAIC, it is beneficial to opt for the MM09$_{hard}$ model,

since the MM$_{0.94}$ peak provides an 'external', non-overlapped reference anchor point for the am-

plitude of the expected MM$_{30}$ peak – the MM landscape is thought to be relatively stable across

healthy subjects in a narrow age range, at least in the absence of pathology [8]. Furthermore, the

MM09$_{hard}$ model does not impose any amplitude assumptions or constraints on the target metab-

olite GABA.

Figure 5 - Distribution of GABA and MM$_{3co}$ estimates, relative contribution of GABA to GABA+ and Pearson's R between GABA and MM$_{3co}$ for all modeling strategies. All three modeling ranges (column) and three spline knot spacings (within each column), and MM$_{3co}$ models (color-coded) are presented. Distributions are shown as half-violins (smoothed distribution), box plots with median, interquartile range, and 25th/75th quartile. The median lines of the box plots are connected to visualize trends within a specific baseline knot spacing. The mean estimates across the three spline knot spacings of the 'none' approach are indicated as a dashed line for each modeling range. Across subject CVs are summarized as bar plots. Minimum/maximum CVs for each modeling range are indicated as downwards/upwards triangle in the color corresponding to the MM$_{3co}$ model. Minimum/maximum CVs for each baseline knot spacing within a specific modeling range are reported on the right side of each column. Global minimum and maximum CVs across all models are added as text.

Figure 5 shows the distributions and CVs of the separate GABA and MM_{3co} estimates of all modeling strategies. Including a separate MM_{3co} basis function significantly decreases GABA estimates, suggesting that not doing so may lead to GABA overestimation, as MM signal is mistakenly modeled as GABA. As was seen for the composite GABA+ estimates in **Figure 3**, sparser knot spacing appears to stabilize modeling, leading to lower CVs of GABA. This becomes especially obvious for the wide modeling range, where GABA CVs exceed 50% for dense knot spacing.

MM_{3co} estimates are stable across the different knot spacings, suggesting that the different parametrizations accurately account for most of the co-edited MM signal at 3 ppm.

The GABA model, in combination with a wide modeling range and 0.55 ppm knot spacing, ex-
hibits the lowest CV for GABA (10.4%). However, the $MM09_{hard}$ model in combination with the
same knot spacing and modeling range has only slightly higher GABA CVs (17.3%) with the
corresponding MM_{3co} CVs being 16.2% ($MM09_{hard}$). Again, despite slightly higher CV values it
is beneficial to use a modeling strategy with a constraint to an 'external' reference peak
($MM09_{hard}$) instead of the highly overlapped MM_{3co} peak of $GABA_{soft}$ (CV = 12.8%), or entirely
omitting MM_{3co}. Additionally, the correlation between the GABA and the MM_{3co} estimates is
lower for the $MM09_{hard}$ model, potentially implying a better separation of GABA and MM_{3co}.
However, a separation of GABA and co-edited macromolecules remains difficult with a low-to-
moderate correlation between GABA and MM_{3co} estimates for all but one modeling strategy
($MM09_{hard}$ for the wide fit range and 0.4 ppm baseline knot spacing). Supplementary Material **5**
reports the mean and SDs of the GABA and MM_{3co} estimates as well as the statistics. Significant
differences between the mean or the SD compared to the corresponding model omitting co-ed-
ited MMs are indicated as shaded box.
493

Figure 6 – Impact of including homocarnosine in the basis set. The directionality of the correlation indicates that HCar absorbs GABA signal specifically for the intermediate and wide modeling range and absorbs MM₃co signal for all modeling ranges. Correlation analysis between the differences between GABA/ MM₃co estimates with and without HCar in the basis set and the

HCar estimates. All three modeling ranges (A-C) and three spline knot spacings (within each subplot) were investigated. A summary bar plot with the correlation coefficient R is shown in the beginning of each row. Pearson's correlation was calculated for each MM_{3co} model (color-coded). Asterisks indicate significant correlations with $p < 0.05 = *$, $p < 0.01 = **$, and $p < 0.001 = ***$.

Finally, **Figure 6** shows the impact of including HCar into the basis set with the difference in

GABA and MM_{3co} estimates between the modeling strategies with and without HCar (ΔGABA

and ΔMM_{3co}, respectively). Interestingly, clear differences in the systematic effects of HCar are

evident between the modeling ranges:

For the narrow modeling range (**Figure 6 A**), HCar estimates correlate positively with ΔGABA,

but the correlation is *only* substantial (R > 0.25) for strategies with a *separate* MM basis

function. For precisely these strategies, HCar estimates correlate negatively with ΔMM_{3co}. These

observations suggest that HCar is likely to account for MM_{3co} in the narrow modeling range. In

contrast, HCar and ΔGABA correlate negatively for most strategies in the intermediate and wide

modeling ranges (**Figure 6 B** and **C**). The negative correlations between HCar and ΔMM_{3co} are

notably weaker for these modeling ranges, indicating that HCar is more likely to substitute for

GABA signal instead of MM.

This behavior can possibly be explained by the HCar signal shape for each modeling range

(**Supplementary Material 6**). For the narrow modeling range, the HCar basis function offers the

model an additional degree of freedom to account for deviations of the actual edited 3-ppm

signal from pure GABA and the symmetric Gaussian MM3co component, as no resonances

below 2.78 ppm are considered. As a result, HCar shows a high correlation with the difference in

MM_{3co}. For the intermediate and wide range, the HCar difference spectrum basis function more

nearly resembles its GABA counterpart since other resonances are included, thereby more

effectively coupling GABA and HCar estimates to each other. Perhaps unsurprisingly, HCar

estimates are significantly higher for 'none' modeling strategy, and are substantially lower for

more flexible baselines, supporting the notion that HCar rather serves as a substitute for an

explicit MM signal, in particular if the baseline cannot absorb the latter (**Supplementary**

Material 6). Within a given knot spacing and modeling range, HCar estimates are comparable

between different MM_{3co} models, a behavior observed for GABA estimates as well.

The GABA+ plus homocarnosine estimates show a slight increase compared to the GABA+

estimates without HCar (**Supplementary Material 7**). For the 'none' model, stronger changes

occur as HCar accounts for MM signal (see also Figure 6). There was no improvement in the

CVs observed when including HCar in the model. The relative contribution of HCar to GABA+

ranged between 2.2% and 19.1% for modeling strategies with an MM_{3co} basis function and

between 18% and 36% for the 'none' model.

Discussion

The application of linear combination modeling to edited difference spectra is neither straightfor­ward nor intuitive. The conceptual advantage of spectral editing arises from isolating a resolved target resonance, i.e. reducing the overlap of the target metabolite with other signals, as well as the number of signals in the spectrum in general[1]. LCM, on the other hand, benefits from maxim­izing the use of prior knowledge to solve the spectral modeling problem, i.e. using all available information for meaningful constraint, including from overlapping signals. The specific case of GABA-edited MRS at 3T poses unique and unresolved challenges. Firstly, a compromise must be drawn between maximizing the prior knowledge by increasing the modeling range and reduc­ing the impact of co-edited and unwanted signals. Secondly, an appropriate parametrization of poorly characterized co-edited signals must be found, and possible interactions with the target metabolite GABA must be evaluated. Thirdly, effects of baseline modeling must be studied, again a consequence of the macromolecular background signal in the GABA-edited difference spectrum not being determined to this date. In this study, a total of 102 linear combination mod­eling strategies were compared for GABA-edited difference spectra, each with different model­ing ranges, parametrizations of co-edited signals, and baseline model flexibility. The key find­ings are:

- Including a dedicated basis function for co-edited MM improves fit residuals, reduces CVs of GABA and GABA+ estimates, and avoids overestimation of GABA.

- Reducing the modeling range does not substantially stabilize or destabilize modeling, while removing potentially valuable information ($MM_{0.93}$ and 2-ppm NAA peak) from the optimization.

- Sparser baseline spline knot spacing leads, on average, to the lowest CV across all modeling ranges.

There is surprisingly little systematic investigation into linear combination modeling of GABA-edited difference spectra. To the best of our knowledge, there is only one conference abstract studying MM parametrization in GABA-edited MRS with the LCModel software[27]. The results from this preliminary investigation indicate that including a specific MM basis function

significantly reduces GABA estimates and were confirmed in the subsequent study [28], which is

substantiated by our findings.

Although the substantial contribution of broad MM signals to the 3-ppm peak in the GABA-ed-

ited spectrum is widely known[1,35], it is rarely explicitly addressed in linear combination model-

ing. Instead, it is assumed that either an incomplete model (without explicit MM term) will still

provide an accurate GABA estimate, or that baseline modeling will account for the MM signal.

The current results provide evidence that including an appropriately parametrized MM model is

a preferrable and easily implemented strategy, reducing the residual over the 3-ppm signal range

by up to 30%, with similar or lower CVs for GABA+. In contrast, not including an MM model

likely causes systematic overestimation of GABA, as the least-squares optimization attempts to

minimize the model-data difference with an inadequate set of basis functions (only GABA), par-

ticularly when a rigid baseline is chosen. Including MM_{3co} is a justified and reasonable measure

without overfitting (reflected by AIC), and stable mean estimates and CVs of MM_{3co} suggest an

adequately parametrized model.

The different MM models in this study were based on certain assumptions, including the relative

contribution of MM_{3co} to the 3-ppm GABA peak to be around 50%[1,6,8,19]. Levels of $MM_{0.93}$ have

been found to be stable across the whole brain[36] and are thought to be stable across healthy sub-

jects. Under these assumptions, the $MM09_{hard}$ model with a rigid amplitude coupling between

MM_{3co} and the non-overlapped $MM_{0.93}$ peak is a suitable strategy, supported by favorable CVs

and ΔAIC. Further studies need to be performed to investigate the distribution and correlation

between $MM_{0.93}$ and MM_{3co} in the brain.

Unedited MRSI data measured at 7T indicates significant differences between white and gray

matter for several macromolecules in the healthy brain[36]. Changes in the MM concentrations dur-

ing disease may also affect the relative contribution to the 3-ppm peak, and therefore render

models with prior amplitude assumptions inaccurate. If there is reason to expect strong fluctua-

tions of MM_{3co}, a modeling strategy with fewer assumptions about amplitude ratios between the

metabolite of interest GABA or the $MM_{0.93}$ signal and the MM_{3co} signal is preferable to the

$MM09_{hard}$ strategy. Here, the $Gauss_{free}$ and $Gauss_{fixed}$ strategies could be used to account for

changes in the MM_{3co} contribution more freely, as their mean estimates of GABA and GABA+

were in good agreement with the more constrained approaches, although they led to increased

CVs and ΔAICs. In addition, the less-constrained models might be more appropriate for investi-

gating changes in MM_{3co} due to age[37] or disease, or for exploring frequency-drift-related effects

on the co-edited MM signal[1,8,38]. Another potential way to model the co-edited MM signal is to

include lysine in the simulated basis set, as it has been identified as the potential source of the

signal[6], although this approach would require appropriate broadening and incorporation of chem-

ical shift and coupling values from protein databases[39].

Overall, results did not differ drastically between modeling ranges, although it is noteworthy that

the effects of baseline flexibility were less pronounced for the narrow modeling range, likely be-

cause the complex interaction of the overlapping 2.25 ppm GABA and Glx signals with the un-

derlying baselines is omitted. Furthermore, there was no evidence that the intermediate modeling

range, which is proposed in the LCModel manual[18] to avoid frequently occurring co-edited lipid

signals, improved quantification substantially compared to both other modeling ranges, although

it should be mentioned that this particular dataset did not suffer from severe lipid contamination.

Taken together, the choice of modeling range does not impact quantitative results as substantially

as the inclusion of an MM model.

Baseline models are included in most LCM algorithms to account for signals not otherwise mod-

eled, e.g. residual water tails or unparametrized macromolecules and lipids. Compared to con-

ventional short-TE spectra, water and non-co-edited MMs are removed upon subtraction in the

GABA-edited spectrum, which is therefore frequently modeled with a stiffer baseline[4,18]. Our re-

sults show that sparser knot spacing (0.55 ppm) leads to lower CVs in metabolite estimates. A

more flexible baseline (0.25 ppm) improves local and global residuals, but not enough to justify

the additional model parameters (as per the AICs). More importantly, an overly flexible baseline

may absorb edited signal, although it appeared that it did not do so excessively even for the 0.25-

ppm strategies. The exception was the 'none' model, where the baseline was the only available

part of the model to take up signal, underlining the inadequacy of the default LCModel approach.

Taken together, a relatively rigid baseline with a parametrized MM basis function is preferable

for LCM of GABA-edited spectra. A caveat to this recommendation is the observation of

structural baseline fluctuations underneath the 2.25 ppm signals from GABA, Glx, GSH, NAA

and NAAG, particularly for the 0.25 ppm knot spacing and a relatively broad increase in the

baseline between 2.7 and 3 ppm. These were observed previously[27], and are likely signals from

un-parametrized MMs directly and indirectly affected by the editing pulse. Rigid baselines may

force a wrong metabolite model in that region and interfere with accurate estimation of GABA

and Glx. In fact, the structural Glx residual at 3.75 ppm suggests a systematic misestimation of

the Glx phase, likely driven by the 2.25 ppm signals. While beyond the scope of this investiga-

tion, it is conceivable that more informed parametrization (or, ideally, direct measurement) of

this unexplored MM background may benefit the modeling of the entire difference spectrum. Al-

ternatively, hitherto unexplored approaches with variable baseline knot spacing may be worth

investigating.

The HCar molecule has a GABA moiety with similar chemical shifts and is therefore co-edited.

Evidence regarding in-vivo HCar levels in the human brain is inconclusive – early work deter-

mined HCar levels to be 0.5 mM[5] (compared to ~1 mM for GABA), while a recent hybrid up-

field/downfield inversion-recovery method determined the HCar/GABA ratio as 17%[40]. There-

fore, we tested the impact of adding HCar to the basis set without additional constraints. Includ-

ing HCar systematically affected GABA and MM_{3co} estimates, in a way that strongly depended

on the choice of modeling range. HCar estimates themselves ranged from 2.2% to 19.1% of the

GABA+ signal, depending strongly on the degree of baseline flexibility. The results suggest that

the overlap between the three model terms (HCar, GABA, MM_{3co}) is too substantial for reliable

three-way separation, particularly in the presence of a highly flexible baseline. A minor increase

in "GABA+ plus HCar" estimates compared to GABA+ estimates was observed and the inclu-

sion of HCar did not substantially improve the CVs. Additionally, the disagreement between the

model and the data at 2.9 ppm indicates that a simple unconstrained addition of HCar to the mod-

eling is not justified.

Symmetric GABA-editing (edit-ON frequency at 1.9 ppm and edit-OFF frequency at 1.7 ppm) is

commonly used eliminate the MM_{3co} contamination of the 3-ppm GABA+ signal. In practice, B_0

instabilities lead to residual MM_{3co} components with variable polarity [11]. The $Gauss_{free}$ and

$Gauss_{fixed}$ MM_{3co} models could potentially be used to account for those variable MM_{3co}

contributions in those spectra. Modeling of those spectra with the current strategies that do have

a non-negative model component as constraint would be challenging. Those modeling strategies

could potentially be adapted by using the B_0 history during the experiment [38] to predict the polar-

ity and relative amplitude of the MM_{3co} signal, and include those as a soft constraint relative to

the MM_{09} signal ($MM09_{hard}$ or $MM09_{soft}$) or the GABA signal ($GABA_{hard}$ or $GABA_{soft}$).

<u>Limitations</u>

A limitation of this study is the high spectral quality (SNR, linewidth, no apparent subtraction

artefacts, or lipid contaminations) of the dataset analyzed. We did not investigate model para-

metrizations of movement or drift, which may introduce systematic changes to the co-edited MM

signal. While our results suggest that using the wide modeling range with a rigid baseline is ben-

eficial, strong co-edited lipid signals are likely to not be modeled appropriately, and the interme-

diate modeling range may be more suitable. Further studies of the possible impact of changes in

spectral quality need to be performed to validate the modeling strategies under suboptimal condi-

tions.

Another limitation is that there is no 'gold standard' of metabolite level estimation in GABA-ed-

ited MRS to validate the results against. The performance of different algorithms or in this study

modeling strategy is often judged by the level of variance [26]. A lower variance does, of course,

not necessarily reflect greater modeling accuracy, but under the assumption that the homogene-

ous study population and data acquisition contribute comparably little biological and instrumen-

tal variance, CVs will predominantly reflect variance introduced by the modeling approach. Re-

cently, the field is witnessing increasing efforts to generate simulated spectra with known ground

truth as a gold standard, although these approaches can only be successful to the extent that those

spectra are truly representative of in-vivo data. Further, such gold standard studies with a known

ground truth could be used to validate whether a correct separation of GABA and MM_{3co} is

achievable by advanced LCM. This study indicates a low-to-moderate correlation between the

GABA and MM_{3co} estimates, suggesting that the two components are not reliably separated.

However, some of the modeling strategies appeared to have a lower association between both es-

timates and could possibly be validated further on a synthetic dataset with known GABA and

MM_{3co} concentrations.

AIC as a measure of the goodness of fit can be used for linear and non-linear approaches if the

log-likelihood is obtained similarly. However, there are two potential limitations for the for the

application of AIC in this study. First, for linear-combination modeling of MRS data, as imple-

mented in Osprey, a non-linear optimization is followed by a linear optimization during each it-

eration. Parameters are treated equally in the calculation of the AIC regardless of whether they

are non-linear (e.g., a phase parameter) or linear (an amplitude parameter). Second, the AIC pe-

nalizes complex models, but does not measure effects of soft constraints and is likely to prefer

models without a soft constraint as those should have a reduced likelihood [41]. Here, we intro-

duced a rather arbitrary correction term of 0.5 per soft constraint for those models to reduce this

effect. Therefore, the resulting ΔAIC values in this study should be interpreted with care and

considered as only one among several metrics to evaluate model performance.

Conclusion

This study proposed and compared different modeling strategies for LCM of GABA+-edited difference spectra from a multi-site MEGA-PRESS dataset. Introducing a parametrized model for co-edited macromolecules reduces fit residuals, while maintaining low coefficients of variation of GABA+ estimates. A rigid baseline was found to be beneficial, while using a narrower modeling range did not significantly improve the modeling. The overall modeling results suggest that GABA-edited data are reliably modeled with an adequately parametrized MM_{3co} model, constrained by the non-overlapped 0.93-ppm MM resonance, in combination with a full modeling range and sparse knot spacing. Incorporating homocarnosine into the modeling did not significantly improve the GABA+ estimates and did not allow for a stable separation of GABA and HCar.

References

1. Choi I-Y, Andronesi OC, Barker P, et al. Spectral editing in 1H magnetic resonance spectroscopy: Experts' consensus recommendations. *NMR Biomed*. 2021;n/a(n/a):e4411. doi:https://doi.org/10.1002/nbm.4411

2. Edden RAE, Puts NAJ, Harris AD, Barker PB, Evans CJ. Gannet: A batch-processing tool for the quantitative analysis of gamma-aminobutyric acid-edited MR spectroscopy spectra. *J Magn Reson Imaging*. 2014;40(6):1445-1452. doi:10.1002/jmri.24478

3. Wilson M, Reynolds G, Kauppinen RA, Arvanitis TN, Peet AC. A constrained least-squares approach to the automated quantitation of in vivo 1H magnetic resonance spectroscopy data. *Magn Reson Med*. 2011;65(1):1-12.

4. Provencher SW. Automatic quantitation of localized in vivo1H spectra with LCModel. *NMR Biomed*. 2001;14(4):260-264. doi:10.1002/nbm.698

5. Petroff OAC, Hyder F, Rothman DL, Mattson RH. Topiramate Rapidly Raises Brain GABA in Epilepsy Patients. *Epilepsia*. 2001;42(4):543-548. doi:https://doi.org/10.1046/j.1528-1157.2001.18800.x

6. Deelchand DK, Marjańska M, Henry P-G, Terpstra M. MEGA-PRESS of GABA+: Influences of acquisition parameters. *NMR Biomed*. 2019;n/a(n/a):e4199. doi:https://doi.org/10.1002/nbm.4199

7. Marjańska M, Terpstra M. Influence of fitting approaches in LCModel on MRS quantification focusing on age-specific macromolecules and the spline baseline. *NMR Biomed*. November 2019. doi:10.1002/nbm.4197

8. Shungu DC, Mao X, Gonzales R, et al. Brain γ-aminobutyric acid (GABA) detection in vivo with the J-editing (1) H MRS technique: a comprehensive methodological evaluation of sensitivity enhancement, macromolecule contamination and test-retest reliability. *NMR Biomed*. 2016;29(7):932-942. doi:10.1002/nbm.3539

9. Bhagwagar Z, Wylezinska M, Jezzard P, et al. Reduction in Occipital Cortex γ-Aminobutyric Acid Concentrations in Medication-Free Recovered Unipolar Depressed and Bipolar Subjects. *Biol Psychiatry*. 2007;61(6):806-812. doi:10.1016/j.biopsych.2006.08.048

10. Mikkelsen M, Barker PB, Bhattacharyya PK, et al. Big GABA: Edited MR spectroscopy at 24 research sites. *NeuroImage*. 2017;159:32-45. doi:10.1016/j.neuroimage.2017.07.021

11. Edden RAE, Oeltzschner G, Harris AD, et al. Prospective frequency correction for macromolecule-suppressed GABA editing at 3T. *J Magn Reson Imaging JMRI*. 2016;44(6):1474-1482. doi:10.1002/jmri.25304

12. Oeltzschner G, Zöllner HJ, Hui SCN, et al. Osprey: Open-source processing, reconstruction & estimation of magnetic resonance spectroscopy data. *J Neurosci Methods*. 2020;343:108827. doi:10.1016/j.jneumeth.2020.108827

13.   Osprey GitHub repository. Osprey GitHub repository. https://github.com/schorschinho/os-
prey. Published 2020. Accessed May 27, 2020.

14.   Klose U. In vivo proton spectroscopy in presence of eddy currents. *Magn Reson Med*.
1990;14(1):26-30. doi:10.1002/mrm.1910140104

15.   Mikkelsen M, Tapper S, Near J, Mostofsky SH, Puts NAJ, Edden RAE. Correcting fre-
quency and phase offsets in MRS data using robust spectral registration. *NMR Biomed*. July
2020:e4368. doi:10.1002/nbm.4368

16.   Barkhuijsen H, de Beer R, van Ormondt D. Improved algorithm for noniterative time-do-
main model fitting to exponentially damped magnetic resonance signals. *J Magn Reson
1969*. 1987;73(3):553-557. doi:10.1016/0022-2364(87)90023-0

17.   Simpson R, Devenyi GA, Jezzard P, Hennessy TJ, Near J. Advanced processing and simu-
lation of MRS data using the FID appliance (FID-A)—An open source, MATLAB-based
toolkit. *Magn Reson Med*. 2017;77(1):23-33. doi:10.1002/mrm.26091

18.   Provencher S. LCModel & LCMgui User's Manual. LCModel & LCMgui User's Manual.
http://s-provencher.com/pub/LCModel/manual/manual.pdf. Published 2020. Accessed April
28, 2020.

19.   Henry PG, Dautry C, Hantraye P, Bloch G. Brain gaba editing without macromolecule con-
tamination. *Magn Reson Med*. 2001;45(3):517-520. doi:10.1002/1522-
2594(200103)45:3<517::AID-MRM1068>3.0.CO;2-6

20.   Levenberg K. A method for the solution of certain non-linear problems in least squares. *Q
Appl Math*. 1944;2(2):164-168. doi:10.1090/qam/10666

21.   Marquardt DW. An Algorithm for Least-Squares Estimation of Nonlinear Parameters. *J Soc
Ind Appl Math*. 1963;11(2):431-441. doi:10.1137/0111030

22.   Becker S. LBFGSB (L-BFGS-B) mex wrapper - File Exchange - MATLAB Central.
LBFGSB (L-BFGS-B) mex wrapper - File Exchange - MATLAB Central.
https://www.mathworks.com/matlabcentral/fileexchange/35104-lbfgsb-l-bfgs-b-mex-wrap-
per. Published February 23, 2015. Accessed March 3, 2021.

23.   Byrd RH, Lu P, Nocedal J, Zhu C. A Limited Memory Algorithm for Bound Constrained
Optimization. *SIAM J Sci Comput*. 1995;16(5):1190-1208. doi:10.1137/0916069

24.   Zhu C, Byrd RH, Lu P, Nocedal J. Algorithm 778: L-BFGS-B: Fortran subroutines for
large-scale bound-constrained optimization. *ACM Trans Math Softw*. 1997;23(4):550-560.
doi:10.1145/279232.279236

25.   Wilson M, Andronesi O, Barker PB, et al. *Methodological Consensus on Clinical Proton
MRS of the Brain: Review and Recommendations*. Vol 82. John Wiley and Sons Inc.; 2019.
doi:10.1002/mrm.27742

26.   Zöllner HJ, Považan M, Hui SCN, Tapper S, Edden RAE, Oeltzschner G. Comparison of
774   different linear-combination modeling algorithms for short-TE proton spectra. *NMR Bio-*
775   *med*. 2021;n/a(n/a):e4482. doi:https://doi.org/10.1002/nbm.4482

776   27.   Murdoch JB, Dydak U. Modeling MEGA-PRESS macromolecules for a better grasp of
777   GABA. In: *19th Annual Meeting of the International Society for Magnetic Resonance in*
778   *Medicine (ISMRM)*. ; 2011. https://scholar.google.com/scholar_lookup?title=Model-
779   ing%20MEGA-PRESS%20macromolecules%20for%20a%20bet-
780   ter%20grasp%20of%20GABA&publication_year=2011&author=J.B.%20Murdoch&au-
781   thor=U.%20Dydak. Accessed July 1, 2020.

782   28.   Dydak U, Jiang YM, Long LL, et al. In vivo measurement of brain GABA concentrations
783   by magnetic resonance spectroscopy in smelters occupationally exposed to manganese. *En-*
784   *viron Health Perspect*. 2011;119(2):219-224. doi:10.1289/ehp.1002192

785   29.   R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria:
786   R Foundation for Statistical Computing; 2017. https://www.R-project.org/.

787   30.   SpecVis GitHub repository. SpecVis GitHub repository.
788   https://github.com/hezoe100/SpecVis. Published 2020. Accessed May 27, 2020.

789   31.   Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York;
790   2009. http://ggplot2.org.

791   32.   Zöllner HJ. Comparison of linear-combination modeling strategies for GABA-edited MRS
792   at 3T. https://osf.io/aqm8f/. Published April 30, 2021. Accessed April 30, 2021.

793   33.   Akaike H. A new look at the statistical model identification. *IEEE Trans Autom Control*.
794   1974;19(6):716-723. doi:10.1109/TAC.1974.1100705

795   34.   Mikkelsen M, Rimbault DL, Barker PB, et al. Big GABA II: Water-referenced edited MR
796   spectroscopy at 25 research sites. *NeuroImage*. 2019;191:537-548. doi:10.1016/J.NEU-
797   ROIMAGE.2019.02.059

798   35.   Cudalbu C, Behar KL, Bhattacharyya PK, et al. Contribution of macromolecules to brain
799   1H MR spectra: Experts' consensus recommendations. *NMR Biomed Revis*. 2020.

800   36.   Považan M, Strasser B, Hangel G, et al. Simultaneous mapping of metabolites and individ-
801   ual macromolecular components via ultra-short acquisition delay 1H MRSI in the brain at
802   7T. *Magn Reson Med*. 2018;79(3):1231-1240. doi:10.1002/mrm.26778

803   37.   Marjańska M, Deelchand DK, Hodges JS, et al. Altered macromolecular pattern and con-
804   tent in the aging human brain. *NMR Biomed*. 2018;31(2):e3865. doi:10.1002/nbm.3865

805   38.   Veen JW van der, Marenco S, Berman KF, Shen J. Retrospective correction of frequency
806   drift in spectral editing: The GABA editing example. *NMR Biomed*. 2017;30(8):e3725.
807   doi:https://doi.org/10.1002/nbm.3725

39.   Borbath T, Manohar SM, Henning A. Towards a Fitting Model of Macromolecular Spectra:
Amino Acids. In: *27th Annual Meeting of the International Society for Magnetic Resonance*
in Medicine (ISMRM). Montreal, Canada; 2019.

40.   Landheer K, Prinsen H, Petroff OA, Rothman DL, Juchem C. Elevated homocarnosine and
GABA in subject on isoniazid as assessed through 1H MRS at 7T. *Anal Biochem*.
2020;599:113738. doi:10.1016/j.ab.2020.113738

41.   Endres DM, Chiovetto E, Giese M. Model selection for the extraction of movement primi-
tives. *Front Comput Neurosci*. 2013;7. doi:10.3389/fncom.2013.00185

Supplementary Material

Table of Contents

Supplementary Material 1 – List of included subjects. All datasets are available at https://www.ni-trc.org/projects/biggaba/

site	subjects	Σ
P01	S01,S03,S04,S05,S08	5
P03	S02,S03,S04,S07,S08,S09,S10,S11,S12	9
P05	S01,S02,S03,S05,S06,S07,S08	7
P06	S01,S02,S03,S04,S05,S06,S07,S08,S09	9
P07	S02,S03,S04,S09,S10,S11,S12	7
P08	S01,S02,S03,S04,S05,S06,S07,S08,S09,S10,S11,S12	12
P09	S01,S02,S03,S04,S05,S06,S07,S08,S09,S10,S11,S12	12
Σ = 7		Σ = 61

Supplementary Material 2. *Properties of the Gaussian functions of the broad macromolecule and lipid resonances included in the basis sets, taken from section 11.7 of the LCModel manual. The amplitude values are scaled relative to the CH_3 singlet of creatine with amplitude 3.*

Name	Frequencies [ppm]	FWHM [ppm]	Amplitude
edit-OFF spectrum basis set			
$MM_{0.94}$	0.915	0.14	3.00
$MM_{1.22}$	1.22	0.15	2.00
$MM_{1.43}$	1.43	0.17	2.00
$MM_{1.70}$	1.67	0.15	0.20
$MM_{2.05}$	2.08	0.15	1.33
	2.25	0.20	0.33
	1.95	0.15	0.33
	3.00	0.20	0.40
Lip09	0.89	0.14	3.00
Lip13	1.28	0.15	2.00
	1.28	0.089	2.00
Lip20	2.04	0.15	1.33
	2.25	0.15	0.67
	2.80	0.20	0.87
Difference spectrum basis set			
$MM_{0.94}$	0.915	0.14	3
MM_{3co}	3	**14 Hz**	2

Data overview

data (mean +- SD)

data (individual spectra)

ppm

4   3   2   1

Supplementary Material 4 – Overview of the processed data including the mean ± SD and individual data.

Supplementary Material 5 – GABA and MM_{3co} mean and SDs for all modeling strategies (ratios to tCr). Significant differences ($p < .05$) between the corresponding model and the 'none' are shaded in gray. Differences in MM_{3co} are compared between the corresponding model and the $GABA_{soft}$ model and significant differences are also shaded in gray .

modeling range		narrow			intermediate			wide		
knot spacing (ppm)		0.55	0.4	0.25	0.55	0.4	0.25	0.55	0.4	0.25
none	[GABA]	0.22 *.062*	0.213 *.071*	0.193 *.057*	0.297 *.048*	0.186 *.061*	0.204 *.044*	0.284 *.029*	0.288 *.034*	0.158 *.066*
GABA$_{hard}$	[GABA]	0.164 *.021*	0.146 *.02*	0.156 *.023*	0.207 *.024*	0.148 *.031*	0.165 *.021*	0.194 *.021*	0.207 *.023*	0.15 *.034*
GABA$_{hard}$	[MM$_{3co}$]	-	-	-	-	-	-	-	-	-
GABA$_{soft}$	[GABA]	0.108 *.024*	0.102 *.027*	0.108 *.025*	0.207 *.048*	0.13 *.039*	0.148 *.03*	0.203 *.026*	0.204 *.027*	0.115 *.049*
GABA$_{soft}$	[MM$_{3co}$]	0.19 *.026*	0.165 *.025*	0.186 *.027*	0.21 *.038*	0.148 *.033*	0.175 *.024*	0.172 *.039*	0.194 *.032*	0.156 *.032*
Gauss$_{fixed}$	[GABA]	0.074 *.031*	0.08 *.031*	0.081 *.035*	0.174 *.055*	0.106 *.045*	0.118 *.045*	0.136 *.031*	0.145 *.032*	0.108 *.068*
Gauss$_{fixed}$	[MM$_{3co}$]	0.233 *.039*	0.192 *.038*	0.223 *.041*	0.237 *.048*	0.192 *.03*	0.215 *.035*	0.233 *.054*	0.237 *.045*	0.235 *.053*
Gauss$_{free}$	[GABA]	0.079 *.034*	0.078 *.032*	0.081 *.033*	0.18 *.054*	0.107 *.05*	0.119 *.045*	0.137 *.033*	0.145 *.031*	0.107 *.069*
Gauss$_{free}$	[MM$_{3co}$]	0.226 *.038*	0.198 *.036*	0.22 *.045*	0.229 *.044*	0.193 *.031*	0.213 *.035*	0.227 *.057*	0.237 *.042*	0.234 *.053*
MM09$_{hard}$	[GABA]	-	-	-	-	-	-	0.195 *.034*	0.189 *.032*	0.106 *.06*
MM09$_{hard}$	[MM$_{3co}$]	-	-	-	-	-	-	0.201 *.032*	0.232 *.043*	0.24 *.04*
MM09$_{soft}$	[GABA]	-	-	-	-	-	-	0.152 *.03*	0.154 *.031*	0.098 *.072*
MM09$_{soft}$	[MM$_{3co}$]	-	-	-	-	-	-	0.216 *.051*	0.232 *.037*	0.24 *.045*

Supplementary Material 6 – Mean modeling results and homocarnosine estimates for all modeling strategies with homocarnosine. A substantial structured residual is visible at 3 ppm if for all modeling strategies and for the narrow and intermediate modeling range the homocarnosine concentrations are significantly lower compared to omitting the co-edited MM, especially for knot spacings <= 0.4 ppm . All

three modeling ranges (columns), three spline knot spacings (rows), and MM$_{3co}$ model (color-coded) are presented with mean residuals and fits, as well as the GABA, MM$_{3co}$, homocarnosine (HCar) and spline baseline models. The mean data is included in black. The arrows indicate the range of values for a specific modeling range and spline knot spacing with the color corresponding to the MM$_{3co}$ model with minimum/maximum value.

Supplementary Material 7 - Distribution of GABA+ plus HCar and HCar estimates and the relative contribution of HCar to GABA+ plus HCar for all modeling strategies. The mean estimates of GABA+ plus HCar across the three spline knot spacings of the 'none' approach are indicated as an dashed line for each modeling range. All three modeling ranges (column) and three spline knot spacings (within each column), and MM$_{3co}$ models (color-coded) are presented. Distributions are shown as half-violins

(smoothed distribution), box plots with median, interquartile range, and $25^{th}/75^{th}$ quartile. The median lines of the box plots are connected to visualize trends within a specific baseline knot spacing. CVs are summarized as bar plots. Minimum/maximum CVs for each spline knot spacing are indicated as downwards/upwards triangle in the color corresponding to the MM_{3co} model. Global minimum and maximum CVs across all models are added as text.

Declaration of competing interests

The authors have nothing to declare.

Acknowledgement

This work is supported by NIH grants P41 EB031771, R01 EB016089, R01 EB023963, R01 EB028259, R21 AG060245, and K99/R00 AG062230.

CRediT authorship contribution statement

Helge J. Zöllner: Software, Formal Analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Visualization. **Sofie Tapper**: Investigation, Writing – Review & Editing. **Steve C. N. Hui**: Investigation, Writing – Review & Editing. **Richard A. E. Edden**: Conceptualization, Formal Analysis, Writing – Review & Editing, Supervision, Project administration, Funding acquisition. **Peter B. Barker**: Writing – Review & Editing, Supervision, Funding acquisition. **Georg Oeltzschner**: Conceptualization, Methodology, Software, Investigation, Formal Analysis, Writing – Review & Editing, Supervision, Funding acquisition.