ERK1/2 is an ancestral organising signal in spiral cleavage

Océane Seudre[1,2], Allan M. Carrillo-Baltodano[1,2], Yan Liang[1], José M. Martín-Durán[1*]

[1] School of Biological and Chemical Sciences. Queen Mary University of London. Mile End

Road, E1 4NS, London. United Kingdom.

[2] These authors contributed equally to this work.

[*] Correspondence: chema.martin@qmul.ac.uk

SUMMARY

Embryonic organisers are signalling centres that instruct the establishment of body plans

during animal embryogenesis, thus underpinning animal morphological diversity. In spiral

cleavage – a stereotypic developmental programme ancestral to 14, nearly half, of the major

animal groups (e.g., molluscs, annelids and flatworms), a cell known as the D-quadrant

organiser defines cell fates and the body axes. ERK1/2 specifies the embryonic organiser in

molluscs, yet how this signalling cascade exerts organising activity and whether this role is

conserved in other spiral cleaving groups is unclear. Here, we demonstrate that ERK1/2

promotes the specification and inductive activity of the D-quadrant organiser in *Owenia*

fusiformis, an early-branching annelid exhibiting ancestral developmental traits. In this

species, active di-phosphorylated ERK1/2 mediated by FGF receptor activity localises to the

4d micromere, establishing the bilateral symmetry and specifying the hindgut and trunk

mesodermal progenitor. Accordingly, impairing FGFR and ERK1/2 activity, as well as cell

communication results in embryos developing anteroventrally radialised. Differential

transcriptomic profiling shows the ParaHox *cdx* and the Notch ligand *delta* as FGFR/ERK1/2

downstream targets in 4d, further revealing that 4d specification instructs the expression of

mesodermal and posterodorsal genes in neighbouring cells, putatively via the Notch pathway.

The instructing role of ERK1/2 in the D-quadrant organiser is thus shared between *O.*

fusiformis and molluscs, representing an ancestral trait of spiral cleavage. Altogether, our

study begins to dissect the gene network promoting axial patterning and posterior growth in

spiral cleavage, revealing extensive mechanistic diversification in body plan specification

despite overall conservation of cleavage patterns in Spiralia.

Keywords: spiral cleavage, Spiralia, embryonic organiser, D-quadrant organiser, annelid,

ERK1/2, MAPK pathway, FGF pathway, Notch pathway.

Introduction

Embryonic organisers are cells or groups of cells able to instruct neighbouring tissues towards certain fates and behaviours during animal embryogenesis [1, 2] (Figure 1A). Embryonic organisers thus play a critical role in patterning tissues and organs, promoting morphogenetic processes that establish animal body plans, thereby emerging as potential drivers of morphological evolution [3-5]. Spiralia is a clade of bilaterally symmetrical animals (Bilateria) comprising nearly half of the major animal clades, including morphologically disparate groups such as molluscs, flatworms and annelids [6, 7]. A stereotypic early developmental programme termed spiral cleavage, characterised by alternating clockwise and counterclockwise oblique cell divisions from the 4-cell stage onwards [8, 9] is however widespread, and likely ancestral to Spiralia [8, 10, 11] (Figure 1A). How spiral cleavage generates the vast diversity of spiralian body plans is still unclear (Figure 1A), albeit this being fundamental to understand the evolution of animal development and morphology.

Spiral cleavage involves the formation of an embryonic organiser that instructs cell fates around four embryonic quadrants, named A to D, which roughly correspond to the left, ventral, right and dorsal sides of the embryo respectively [8, 9]. It is a cell belonging to the D-quadrant – e.g., the 3D or 4d blastomeres in molluscs [12] – that acts as embryonic organiser, and thus this signalling centre is named D-quadrant organiser (Figure 1A). The most common and probably ancestral mode to specify the D-quadrant and its organising cell is through inductive signals at around the 32-cell stage [13, 14]. This type of spiral cleavage is termed conditional; or equal, since early embryos exhibit quadri-radial symmetry during early stages. However, in some mollusc and annelid species, the specification of the D-quadrant organiser is largely cell-autonomous, as two asymmetric zygotic divisions segregate maternal mRNAs to one (the D) blastomere at the 4-cell stage [9, 13, 14]. This autonomous mode of spiral

cleavage is also referred to as unequal, because the D founder cell at the 4-cell stage is larger.

Regardless of the specification mode, the ERK1/2 signalling pathway, an evolutionarily

conserved intracellular cascade of kinases involved in an array of cellular processes [15], is

active and necessary in the D-quadrant organiser in molluscs [16-20] (Table S1). However,

what activates ERK1/2 and how ERK1/2 orchestrates the instructing role of the D-quadrant

organiser in this group is not understood (Figure 1A), and thus how embryonic organisers

define the spiralian body plan is still an unresolved question.

The role of ERK1/2 signalling during spiral cleavage in groups other than Mollusca is in

addition ambiguous and only explored in Annelida (Table S1). As in molluscs, the conditional

cleaving annelid *Hydroides hexagonus* exhibits active ERK1/2 in the 4d cell [17], which is

often called "mesentoblast" and cell lineage analyses identify as the trunk mesodermal

precursor in most annelid embryos [21-23]. Yet the role of ERK1/2 in the 4d cell of *H.*

hexagonus is unknown. In autonomous cleaving annelids, the patterns of ERK1/2 activity

differ markedly between species (Table S1), but in general ERK1/2 is not active in 4d and its

inhibition does not result in axial polarity defects in any of the studied species [24-27].

Indeed, the organiser activity occurs in the 2d blastomere and $2d^{11}$ + 4d in the autonomous

cleaving annelids *Capitella teleta* and *Tubifex tubifex* respectively [26, 28]; and its inductive

role relies on Activin/Nodal activity, and not on ERK1/2 signalling in the autonomous species

C. teleta and *Chaetopterus pergamentaceus* [24, 29]. Therefore, the cellular and molecular

basis underpinning D-quadrant organiser specification and activity varies considerably within

Annelida, which ultimately prevents inferring whether ERK1/2 activity in the D-quadrant

organiser is a homologous trait to Spiralia or a molluscan innovation that also appeared

convergently in some annelid lineages.

The annelid worm *Owenia fusiformis* belongs to Palaeoannelida – the sister group to all

remaining annelids [30] – and exhibits embryonic traits considered ancestral to Annelida,

including conditional spiral cleavage, formation of a coeloblastula, gastrulation by

invagination and the development of a planktotrophic trochophore-like larva [31] (Figure 1B,

C). While inductive signals appear to control the establishment of the axial polarity in *O.*

fusiformis [32], which blastomere acts as D-quadrant organiser and how this signalling centre

becomes specified and controls body patterning is unknown. Therefore, investigating the

nature of the D-quadrant organiser and the role of the ERK1/2 signalling pathway in *O.*

fusiformis can, by comparison with other annelid and molluscan lineages, help to infer

ancestral character states to Annelida and Spiralia as a whole, ultimately contributing to our

understanding of the evolution of spiral cleavage and spiralian body plans.

Results

***Di-phosphorylated ERK1/2 is enriched in the 4d cell in* O. fusiformis**

The annelid *O. fusiformi*s has a single ERK1/2 ortholog (Figure S1A), which is expressed at

low levels in active oocytes (Figure S1B). The expression of *erk1/2* is stable during early

cleavage divisions and up to 4 hours post fertilisation (hpf; ~32 cell stage), increasing

thereafter and plateauing at the larval stage (Figure S1B). A cross-reactive antibody against

the active di-phosphorylated form of ERK1/2 (di-P-ERK1/2) [16, 17] shows weak ubiquitous

signal in the zygote and up to 3 hpf (~16 cell stage) (Figure S1C). At 4 hpf, however, the four

most animal/apical micromeres ($1q^{111}$) exhibit enriched di-P-ERK1/2 signal, which disappears

at 5 hpf (coeloblastula) (Figure 1D, Figure S1C). At this stage, a single cell in the vegetal

pole, which positionally corresponds to a fourth quartet micromere (4q), shows strong signal

for di-P-ERK1/2 (Figure 1D, Figure S1C). In *O. fusiformis*, the fourth micromere quartet (4q)

starts to form at 4 hpf, when the 3Q macromeres sharing the vegetal cross furrow divide

(Figure 1E). This creates an asymmetry in the timing and order of appearance of the 4q

micromeres that is also observed during cleavage of the 3q micromeres, which divide in a

counterclockwise order after all 3Q macromeres have cleaved (Figure 1E). By 5 hpf however,

the vegetal pole exhibits again a symmetrical arrangement (Figure 1E). At 6 hpf, with the

onset of gastrulation [31], di-P-ERK1/2 signal is enriched in seven vegetal cells forming a

bilaterally symmetrical pattern that includes one 4q micromere and the $2q^{12}$ and $2q^{22}$

blastomeres of the other three embryonic quadrants (Figure 1D, Figure S1C). At this stage,

the 4Q macromeres have cleaved synchronously into the 5Q and 5q cells and all but the 4q

micromere enriched in di-P-ERK1/2 have divided, becoming the earliest morphological sign

of bilateral symmetry in *O. fusiformis* embryos. This di-P-ERK1/2 positive 4q micromere

divides into two large cells only after ingression during early gastrulation at 7 hpf (Figure 1E),

with the later progeny occupying a bilateral position on the posterodorsal side of the

archenteron once gastrulation completes at 9 hpf (Figure 1E). The behaviour of this 4q

micromere resembles the cellular dynamics described for the 4d micromere in other

conditional spiral cleaving annelids [33-36], suggesting that the di-P-ERK1/2 enriched 4q

micromere at 5 and 6 hpf is the 4d cell and its two large daughter cells are the MR/ML

mesoteloblasts (Figure 1D, E). Therefore, ERK1/2 signalling activates first in the $1q^{111}$ animal

micromeres and subsequently in the 4d micromere in the annelid *O. fusiformis*, which is

similar to the spatiotemporal dynamics of ERK1/2 activity reported in spiral cleaving

molluscs (Table S1).

***ERK1/2 signalling controls axial polarity in* O. fusiformis**

To examine the role of the ERK1/2 signalling during *O. fusiformis* development, we treated

embryos with brefeldin A (BFA), an inhibitor of intracellular protein trafficking previously

used to block the induction of the organiser in other spiral cleaving embryos [32, 37], and

U0126, a selective inhibitor of MEK1/2 and ERK1/2 di-phosphorylation [16, 38] (Figure 2A).

For both drugs, treatment from fertilisation (~0.5 hpf) to 5 hpf, when di-P-ERK1/2 is enriched

in 4d, effectively blocks activation of ERK1/2 (Figure 2B; Figures S2A; Table S2) and causes

the loss of bilateral symmetry, posterior structures (e.g., chaetae and hindgut) and larval

muscles in 100% of the embryos at a 10 µM concentration (Figure 2C; Figure S2B; Table S3,

S4). Compared to control samples, 0.5 to 5 hpf treated embryos lack a fully formed apical tuft

and apical organ, showing reduced ectodermal expression of the apical organ marker *six3/6*

[39] and just a few apical cells positive for the neuronal marker *synaptotagmin-1* (*syt-1*) [31]

(Figure 2D). In addition, treated embryos lack expression of hindgut (*cdx*) and trunk

mesodermal (*twist*) markers [39], exhibit expanded expression of the oral ectodermal marker

gene *gsc* [39] around the single gut opening (Figure 2D; Figure S2C), and retain expression of

the midgut endodermal marker *GATA4/5/6b* [39] (Figure S2C). We deem this phenotype as

antero-ventrally radialised (or Radial; Figure 2C, E). Therefore, activation of ERK1/2

signalling in the 4d cell at the coeloblastula stage relies on inductive cell-cell communication

signals (impaired by BFA treatment) required to specify and develop posterior and dorsal

structures during *O. fusiformis* embryogenesis.

To dissect the exact timing of induction and activity of ERK1/2 during *O. fusiformis* cleavage,

we treated embryos with 10µM BFA/U0126 in overlapping time windows from fertilisation

to early gastrulation (Figure 2E; Table S5). Blocking protein secretion with BFA from

fertilisation to the 8-cell stage does not affect normal development. However, BFA treatment

between the 8-cell stage and 4 hpf results in larvae with all morphological landmarks of a

typical mitraria larva and normal expression of tissue-specific markers, but with a compressed

morphology (Figure 2F; Figure S2C). Only treatment with BFA from 4 hpf to 6 hpf, and

hence spanning the formation of 4d, causes a radial phenotype, with treatments after 5 hpf

being lethal (Figure 2E, Figure S2C). This suggests that the intercellular communication event

inducing the activation of ERK1/2 in 4d happens between 4 and 5 hpf, right during 4q

micromere formation (Figure 1E). Unexpectedly, preventing ERK1/2 di-phosphorylation with

U0126 from the 2-cell stage until 4 hpf, when this induction event might begin, also causes a

radial phenotype (Figure 2E), with just slight differences between certain timepoints; embryos

treated between 2-cells and 3 hpf show expression of the mesodermal marker *twist* in a few

discrete cells, and embryos treated from 4 to 6 hpf are seemingly more elongated along the

apical-oral axis (Figure S2C). Therefore, ERK1/2 activity is essential for normal embryonic

patterning and posterodorsal development throughout most spiral cleavage in *O. fusiformis*.

However, the combination of BFA and U0126 phenotypes suggest that ERK1/2 acts

autonomously from the 2-cell stage to 4 hpf, while it requires of inductive cell-to-cell

communication signals for its enrichment in 4d.

ERK1/2 signalling activates posterodorsal and mesodermal genes

To investigate the mechanisms through which ERK1/2 controls posterodorsal development in

O. fusiformis, we hypothesised that genome-wide profiling of gene expression in BFA and

U0126 treated embryos would uncover upstream regulators and downstream targets of

ERK1/2 activity. We thus treated embryos with either 10 µM BFA or 10 µM U0126 from

fertilisation to 5 hpf (to cover both the autonomous and conditional phases of ERK1/2

activity) and performed RNA-seq transcriptome profiling in treated and controlled embryos

collected right after di-P-ERK1/2 enrichment in the 4d cell (coeloblastula; 5.5 hpf) and at the

larval stage (Figure 3A). Principal component analysis (PCA) and sample-to-sample distance

matrices clustered transcriptomic profiles according to developmental stage and biological

replica (Figure S3A, B). Differential expression analyses revealed 90 and 268 differentially

expressed genes (DEGs; Log Fold Change < -1.5 and adjusted p-value < 0.05) in BFA treated

coeloblastula and larva respectively, and 132 (coeloblastula) and 373 (larva) DEGs after

U0126 treatment (Figure 3B). When considering all comparisons and removing redundancies,

we detected a total of 628 DEGs, 414 (65.92%) of which were functionally annotated and

enriched in gene ontology terms related to regulation of transcription, development and cell

fate specification (Table S6, S7, S8). Most of these DEGs were downregulated (Figure 3B, C;

Figure S3C, D), with 26 genes downregulated under both drug treatments at the coeloblastula

stage and 154 genes commonly downregulated at the larval stage. Only three DEGs (*cdx*, *fer3*

and *foxH*) appeared systematically downregulated in both drugs and in the two developmental

time points, while we did not identify any common upregulated DEG (Figure 3B; Figure

S3D). Therefore, our approach revealed a confident and relatively small set of genes whose

expression is dependent on ERK1/2 activity and early inductive signals in *O. fusiformis*

embryos.

To validate the set of genes affected by 4d misspecification, we selected 22 candidate genes

for further gene expression analyses (Figure 3D, Table S9), including a variety of

transcription factors recurrently involved in axial embryonic patterning (e.g., *six3/6*, *gsc*, *cdx*,

AP2, *foxQ2*), mesoderm development (e.g., *twist*, *hand2*, *foxH*) and neurogenesis (e.g., *POU4,*

irxA), Wnt ligands (*wnt1*, *wntA* and *wnt4),* TGF-β modulators (*noggin* and *BAMBI*), and

Notch signalling components (*delta* and *notch-like*) (Table S8). Stage-specific RNA-seq data

covering twelve developmental time-points, from the unfertilized oocyte to the mature larva,

confirmed that for all candidate genes, their expression upregulates at the time of or just after

4d specification and di-P-ERK1/2 enrichment in this cell (Figure 3D). These genes are

expressed either in apical/anterior domains (*noggin*, *BAMBI*, *foxQ2*, *POU4*, *six3/6*, *gsc*), the

posterior larval tip and chaetae (*fer3*, *lhx1/5*, *wnt1*, *notch*, *msx2-a*, *irxA*, *AP2*, *wnt4*, *delta*), the

hindgut (*cdx*), or mesodermal derivatives (*foxH*, *rhox*, *wntA*, *POU3*, *hand2*, *twist*) (Figure 3D,

Figure S4). Analysis of the expression of these genes in control and treated embryos at the

coeloblastula (5.5hpf) and larva (24hpf) stages confirmed the expression domains of these

genes disappear after treatment with either BFA or U0126 (Figure 3E, Figure S4), thus

validating our RNA-seq approach. Altogether, these results define a set of co-regulated genes

that act downstream of ERK1/2 signalling and/or early inductive events, and that are involved

in the development of apical, posterodorsal and mesodermal structures in *O. fusiformis*.

ERK1/2 signalling specifies and patterns the D-quadrant

Our RNA-seq study and candidate gene screening revealed nine genes expressed at the

vegetal pole at 5.5 hpf and whose expression was affected by either direct inhibition of

ERK1/2 di-phosphorylation (*cdx*, *delta*, *foxH*, *wnt1*, *wntA*, *rhox*, *fer3* and *AP2*) or impairing

cell-to-cell inductive signals (*gsc*) (Figure 3E, Figure S4). None of these genes are expressed

at 5 hpf, at the onset of ERK1/2 activity in the 4d micromere (Figure 4A). Instead, the early

endodermal marker *GATA4/5/6a* [39], whose expression is unaffected by BFA and U0126

treatment and thus is cell-autonomous, is symmetrically expressed in the gastral plate

(including 4d) at this stage (Figure 4A; Figure S5A). At 5.5 hpf, half an hour after the initial

activation of ERK1/2 signalling in 4d, the vegetal pole becomes bilaterally symmetrically

patterned, but only two of the ERK1/2 dependent genes (*cdx* and *delta*) are expressed in the

4d micromere (Figure 4A). The ParaHox gene *cdx*, which is detected in the hindgut of *O.*

fusiformis at the larval stage [39] (Figure 3E; Figure S5B), becomes expressed in 4d, and later

on in two cells of the MR and ML progeny at the gastrula stage (Figure 4B), as expected from

the behaviour of 4d at these stages (Figure 1E). Similarly, the Notch ligand *delta* (Figure S5C,

D) is expressed in 4d at 5.5 hpf, but also in most of the descendants of 1d at the animal pole,

plus 1a–c[1121] and ectodermal derivatives of the C and D quadrant at the vegetal pole (Figure

4A, C). To investigate how ERK1/2 might control activation of *cdx* and *delta*, we used Assay

for Transposase-Accessible Chromatin using sequencing (ATAC-seq) data at 5 hpf and

identified known bound transcription factors in the accessible regions associated with these

genes (Figure 4D), which include DNA binding motifs for transcription factors known to be

modulated by ERK1/2 phosphorylation, such as ETS, RUNX and GATA factors [15].

Therefore, ERK1/2 di-phosphorylation in 4d seems to delay cell cycle progression in this cell

with respect to 4a–c (Figure 1E) and control the activity of transcriptional regulators that

induce posterior fates (*cdx*) and cell-cell communication genes (*delta*).

The other six additional ERK1/2-dependent genes pattern the micromeres surrounding 4d at

5.5 hpf, defining mesodermal and posterior ectodermal domains (Figure 4A). The

transcription factor *foxH* (Figure S5E), which regulates mesoderm development during

gastrulation in vertebrate embryos [40-43], is detected in the $3c^{22}$ and $3d^{22}$ micromeres

adjacent to 4d (Figure 4A), whose progeny contributes to lateral ectomesoderm in the annelid

Urechis caupo [36]. Later in development, *foxH* is expressed in a posterior V-shaped pattern

of putative mesodermal precursors during axial elongation, fading away in larval stages

(Figure S5B). The ligands *wnt1* and *wntA* (Figure S5), expressed in the posterior region and

D-quadrant during development in the annelid *Platynereis dumerilii* [44], are expressed in

two bilaterally symmetrical columns of micromeres ($3c^{22}$, $3c^{21}$, $3c^{12}$, $3c^{11}$, $3d^{22}$, $3d^{21}$, $3d^{12}$,

$3d^{11}$) plus the micromeres $2d^{222}$ and $2d^{112}$ in *wntA* (Figure 4A). The homeobox *rhox* (Figure

S5G), which is expressed in male and female primordial germ cells in vertebrates [45], is

detected in the micromeres $2d^{112}$ and $2d^{212}$ (Figure 4A) and thereafter in two small cells inside

the blastocele in the gastrula (Figure S4). The transcription factors *fer3* and *AP2* are expressed

in two single micromeres ($1c^{222}$ and $1d^{221}$) and in a broader posterior ectodermal domain

($1c^{222}$, $1d^{221}$, $2d^{122}$, $2d^{212}$, $2d^{2111}$, $2d^{2112}$, $2d^{1112}$, $2d^{1111}$) respectively, becoming restricted to a

small expression domain at or near the posterior chaetal sac of the larva (Figure S4; Figure

262 S5B). The larger size of 4d, which stays as a big posterior cell at 5.5 hpf, allows it to establish

263 almost as double direct cell-cell contacts with its surrounding cells than each of the daughter

264 cells of 4a–c, including most of the cells expressing *foxH*, *wnt1*, *wntA*, *rhox*, and *AP2* (Figure

265 4E). This data demonstrates the inductive role of the 4d in defining mesodermal and

266 posterodorsal fates, and together with the upregulation of the Notch-ligand *delta* in 4d after

267 ERK1/2 activation, suggests that this role might occur by direct cell-cell communication

268 mediated by the Notch signalling pathway.

270 The homeobox gene *gsc* is the only candidate expressed outside the D-quadrant at 5.5 hpf,

271 being detected in micromeres $2a^{11}$–$2c^{11}$, $2a^{12}$–$2c^{12}$, $2a^{21}$–$2c^{21}$, and $2a^{22}$–$2c^{22}$ (Figure 4A).

272 Consistent with its location outside the D-quadrant, *gsc* expression is independent of ERK1/2

273 activity, but requires of inductive signals, as demonstrated by being downregulated after BFA

274 treatment (Table S3). Accordingly, *gsc* expression disappears in BFA-treated coeloblastulae,

275 while inhibition of ERK1/2 activity with U0126 expands *gsc* domains, which becomes

276 detected in all embryonic quadrants in a radial fashion (Figure 4F). Indeed, in ERK1/2

277 inhibited embryos all 4q micromeres cleave into $4q^1$ and $4q^2$, as no 4q becomes the 4d cell

278 (Figure 4F). Therefore, all quadrants adopt antero-ventral fates (*gsc*), preventing the

279 expression of posterior marker genes such as *AP2* (Figure 4F). These results demonstrate that

280 the radial phenotype observed after U0126 is a consequence of misspecifying the 4d cell, and

281 not a result of the D-quadrant becoming specified but not developing further. In addition, this

282 data also demonstrates that specification of the 4d cell through ERK1/2 activity represses

283 anterior fates, as shown by limiting *gsc* expression.

FGF signalling regulates ERK1/2 di-phosphorylation in 4d

In *O. fusiformis*, *cdx* and *delta* are expressed in 4d under regulation of ERK1/2, and their expression is associated with hindgut and trunk mesoderm development, as well as posterior growth (Figure 2C). In vertebrate embryos, FGF signalling mediated by ERK1/2 activity regulates the expression of *cdx* and *delta* in the developing hindgut and presomitic mesoderm during posterior elongation [46] and FGF signalling promotes mesoderm development in the spiralian groups Brachiopoda and Phoronida [47]. We thus hypothesised that FGF signalling might be the upstream regulator driving ERK1/2 activity in 4d (Figure 5A). *O. fusiformis* has a single FGF receptor (FGFR; Figure S6A) with high amino acid conservation at key functional residues compared to the human FGFR1 ortholog (Figure 6B). While it is transcriptionally upregulated at the gastrula stage, *FGFR* appears weakly expressed at the gastral plate at the coeloblastula stage in *O. fusiformis* (Figure S6C, D), as observed in brachiopods and phoronids [47]. At this stage, FGF ligands and VEGFR, the other tyrosine kinase receptor inhibited by SU5402, are only weakly expressed (Figure S6C) and not detected by *in situ* hybridisation (Figure S6D). Treatment with 30 µM SU5402, a selective inhibitor of FGFR phosphorylation and activation [48] (Figure 5A), prevents di-P-ERK1/2 enrichment at 4d cell in 96% of the treated embryos at 5 hpf (Figure 5B, Figure S6E, Table S2). Indeed, treatment with SU5402 during the specification of the 4d micromere (4 to 6 hpf and 3 to 7 hpf; Figure 5C) results in an anteriorly radialised phenotype, with embryos developing into larvae lacking posterior and reduced mesodermal structures and showing radially expended oral ectodermal fates (*gsc*) (Figure 5D, Table S3). Conversely, treatment before di-P-ERK1/2 enrichment in 4d (3 to 5 hpf) causes a slightly compressed phenotype (Figure S6F). Therefore, inhibition of FGFR blocks ERK1/2 activation and phenocopies the effect of U0126 and BFA (Figure 2C), suggesting that FGF signalling activity is upstream of and necessary for 4d specification in *O. fusiformis*.

Discussion

Our study provides compelling evidence that FGFR and ERK1/2 signalling specify and

control the organising activity of the 4d micromere in the conditional spiral cleaving annelid

O. fusiformis (Figure 6A). FGFR/ERK1/2 signalling at around 5 hpf is necessary for

instructing one of the recently emerged 4q micromeres into the D fate (Figure 2E, Figure 5C,

Figure 6A). However, 4q micromeres form already asymmetrically at 4 hpf in *O. fusiformis*

(Figure 1E), yet whether there is an exact correspondence between the order of formation and

the fate of each 4q remains elusive. Asynchronies in the cleavage of 3Q have not been

described for other conditional cleaving annelids [33-35], but are more common in molluscs,

where 4d can be either the first or the last micromere to form [3, 12, 14]. This asynchrony in

3Q cleavage, together with the fact that ERK1/2 signalling affects axial polarity

independently of cell communication (i.e., unaffected by BFA) earlier than 4 hpf (Figure 2E,

Figure S2C) suggest that cell-autonomous ERK1/2-dependent mechanisms might act before

the inductive specification of 4d in *O. fusiformis*. In other systems, MAPK controls unequal

protein segregation during mitosis via pericentrosomal material [49], which is the subcellular

region where maternal mRNAs condense before asymmetric inheritance during spiral

cleavage in the mollusc *Tritia obsoleta* [50], and MAPK controls mRNA stability during early

zygotic divisions in the annelid *Helobdella robusta* [51]. Therefore, basal levels of ERK1/2

signalling during early cleavage might control the activity or differential segregation of

determinants required to respond adequately to the inductive signals specifying the 4d cell in

O. fusiformis. Although further work is needed to test this scenario, our findings indicate that

both autonomous and conditional mechanisms are essential to specify the D-quadrant

organiser in *O. fusiformis*, challenging traditional views that conditional/equal spiral cleaving

species rely mostly on inductive signals to establish their body pattern.

The specification of 4d micromere has at least four immediate consequences during *O.*

fusiformis embryogenesis (Figure 6A). First, and contrary to its role in promoting cell

proliferation [15], ERK1/2 activity correlates with 4d cell cycle arrest, breaking the quatri-

radial symmetry of the embryo as the 4a–4c micromeres cleave (Figure 1E; Figure 6A). In

some molluscs, activation of ERK1/2 in the 3D blastomere also correlates with cleavage

delays [12, 17], and thus a potential role of ERK1/2 signalling in arresting, instead of

promoting, cell division appears to be common in the context of the embryonic organiser in

spiralian embryos. Second, specification of 4d and the D-quadrant represses anterior fates to

the A–C quadrants, as observed by the expression of the BFA-affected oral ectodermal gene

gsc in all four quadrants in ERK1/2 inhibited embryos (Figure 4F). These findings suggest

that two as yet unknown opposing signals – one specifying anterior fates and promoting *gsc*

expression, and another driving 4d specification via FGFR/ERK1/2 – might act

simultaneously at 4–5 hpf (Figure 6A). Considering that *gsc* expression is eventually

recovered in BFA treated larvae (Figure S2C), we hypothesise that this anteriorising signal

must come from cell lineages already specified by 4–5 hpf, such as the endoderm/midgut

(Figure 4A) or the apical organ (Figure 2D). Third, FGF/ERK1/2 activity upregulates the

hindgut gene *cdx* in 4d (Figure 3C; Figure 4A; Figure 6A). Inhibition of both FGF and

ERK1/2 activity affects hindgut but not foregut and midgut development (Figure 2C; Figure

5D), indicating that unlike most annelids [21] but similar to molluscs [3, 52], 4d contributes to

both endoderm/hindgut and mesoderm in *O. fusiformis*. Fourth, FGF/ERK1/2 activity also

upregulates the Notch ligand *delta* in 4d, as well as a set of mesodermal and posterior genes,

Wnt ligands and TGF-β modulators in its surrounding cells, most of which establish a direct

cell-to-cell contact with 4d (Figure 4A, E; Figure 6A). Therefore, 4d instructs cell fates in the

D and C quadrant at the vegetal pole, probably through the Notch signalling pathway, as also

proposed for the mollusc *Tritia obsoleta* [53]. Altogether, our study demonstrates the

organising properties of the 4d cell (endomesodermal precursor) in *O. fusiformis*, defining a

comprehensive, testable model to study the establishment of the body plan in this and other

spiral cleaving organisms.

Enrichment of di-phosphorylated ERK1/2 in the 4d micromere is common to *O. fusiformis*

and *H. hexagonus*, the two conditional spiral cleaving annelids studied so far [17], which

contrasts with the variability in ERK1/2 activity observed among autonomous cleaving

annelids [25-27]. Likewise, the 4d cellular identity of the D-quadrant organiser in *O.*

fusiformis differs from the autonomous annelid species *C. teleta* and *T. tubifex*, which

concentrate the organising activity in the 2d and $2d^{11}$ + 4d cells respectively [26, 28].

However, the 4d cell identity observed in *O. fusiformis* resembles the molluscan condition,

where ERK1/2 is active in 3D and maintained in 4d, both of which can exert organising

activity [9, 12]. Moreover, *cdx* and *delta*, the downstream targets of ERK1/2 in the 4d of *O.*

fusiformis, are also expressed in 4d in the mollusc *Tritia obsoleta* [52, 53], further suggesting

conservation in both signalling cascades and transcriptional outputs between D-quadrant

organisers in annelids and molluscs. Therefore, our study supports that ERK1/2 activity in the

cell acting as embryonic organiser is a shared homologous trait between Annelida and

Mollusca and given our current understanding of Spiralia phylogeny [6, 7], a probable

ancestral character to at least the internal node Lophotrochozoa. Under this scenario,

heterochronic shifts to confer organising activity to 2d micromeres as seen in *C. teleta* and *T.*

tubifex [26, 28] and the use of Activin/Nodal signalling instead of ERK1/2 to establish the

body plan [24, 29] are thus derived conditions.

More broadly, our study uncovers a role of the FGF signalling pathway, an evolutionarily

conserved upstream regulator of mesoderm development in bilaterians [47, 54, 55], in the

specification of 4d, the D-quadrant organiser and endomesodermal precursor in *O. fusiformis*

(Figure 5B–D). FGFR activity in 4d results in the activation of *cdx*, Notch signalling (*delta*)

and Wnt signalling (*wnt1*, *wntA*) in vegetal micromeres of the D-quadrant (Figure 4A) that

will occupy the posterodorsal blastoporal lip as gastrulation begins, supporting that

blastopore-associated axial organisers also occur during spiral cleavage [4]. Furthermore,

FGF, Wnt and Notch signalling, together with *cdx* are all conserved gene modules regulating

posterior axis elongation in an array of bilaterian embryos (Figure 6B) [39, 46, 47, 56-58].

Classically, axial elongation in spiralian embryos has been proposed to occur by proliferation

of the 2d clone [59, 60], which express *delta* and *wntA* in *O. fusiformis* (Figure 4A), further

suggesting that specification of 4d also results in the deployment of broadly conserved genetic

mechanisms involved in posterior growth that eventually promote axial (and blastopore)

elongation in *O. fusiformis* (Figure 2C; Figure 6B). Altogether, our study demonstrates that

the establishment of the axial identities (A–D quadrants), endomesoderm specification,

posterodorsal patterning and axial elongation are all intimately connected events that depend

on the instruction of a single cell fate (the 4d cell) and use evolutionarily conserved

mechanisms in *O. fusiformis*, a conditional spiral cleaving annelid. Decoupling these

embryonic organising processes through maternal determinants (e.g., as in autonomous spiral

cleavers), heterochronic shifts in organiser specification and/or signalling pathway rewiring

(e.g., co-option of Activin/Nodal) might have thus contributed to the emergence of the vast

diversity of body plans observed today in Spiralia.

Acknowledgements

We thank the staff at Station Biologique de Roscoff for their help with collections and animal

supplies, as well as the Oxford Genomics Centre at the Wellcome Centre for Human Genetics

(funded by Wellcome Trust grant reference 203141/Z/16/Z) for the generation and initial

processing of RNA-seq and ATAC-seq sequencing data. We thank all members of the Martín-

Durán lab for support and valuable comments on the manuscript. This work was funded by a

European Research Council Starting Grant (action number 801669) to J.M.M.-D. and a Queen

Mary, University of London studentship to O.S.

Author Contributions

417   O.S., A.M.C.-B. and J.M.M.-D. conceived and designed the study. O.S., A.M.C.-B., Y.L. and

418   J.M.M.-D. performed all experimental approaches and critically analysed the data. O.S.,

419   A.M.C.-B. and J.M.M.-D. wrote the manuscript. All authors read and commented on the

420   manuscript.

Declaration of Interests

423   The authors declare no competing interests.

Materials and Methods

Animal husbandry and embryo collections

Adult specimens of *O. fusiformis* Delle Chiaje, 1844 were collected from the coast near the

Station Biologique de Roscoff (France) during the reproductive season (May to July). In the

lab, animals were kept in artificial seawater (ASW) at 15°C. *In vitro* fertilizations were

conducted as previously described [31, 39] and embryos develop in glass bowls with filtered

ASW at 19 °C until the desired embryonic stage.

Drug treatments

Embryos were treated with either brefeldin A (BFA; Sigma-Aldrich, #B7651),

U0126 (Sigma-Aldrich, #U120), or SU5402 (Sigma-Aldrich, #572630). All drug stocks were

made in dimethyl sulfoxide (DMSO) and diluted in ASW prior use, with optimal working

concentrations (10 µM for BFA and U0126 and 30 µM for SU5402) established after initial

titration. For all drug treatments, equivalent volumes of DMSO were used as negative control,

and in all cases, treatments were initiated after ~0.5 hours post fertilisation (hpf), right after

fertilisation and sperm removal. Treatments were stopped by washing out the drug with at

least three washes in ASW and embryos were either collected for downstream analyses or

raised until larval stage. Samples collected for immunohistochemistry and gene expression

analyses were fixed in 4% paraformaldehyde (PFA) in ASW for 1h at room temperature (RT).

Larvae were relaxed in 8% magnesium chloride ($MgCl_2$) prior fixation and fixed in 4% PFA

in 8% $MgCl_2$ for 1h at RT. After fixation, samples were washed several times with 0.1%

Tween-20 phosphate-buffered saline (PTw), and either stored in PTw at 4°C for

immunohistochemistry or dehydrated to 100% methanol and stored at -20°C for whole mount

in situ hybridisation. Treated and control embryos and larvae collected for RNA-seq analyses

were snap frozen in liquid nitrogen and stored at -80°C prior total RNA extraction.

Immunohistochemistry

F-actin labelling, and antibody staining were conducted as previously described [31], with the

modification that samples collected for anti-di-P-ERK1/2 staining were washed in PTw

supplemented with 1:100 dilution of phosphatase inhibitors (Cell Signalling, #5872) after

fixation. The primary antibodies mouse anti-double phosphorylated ERK1/2 (Sigma-Aldrich,

#M8159, 1:200) and mouse anti-acetylated α-tubulin (clone 6-11B-1, Millipore-446 Sigma,

#MABT868, 1:500) were diluted in 5% normal goat serum (NGS) in 0.1% Triton X-100

phosphate-buffered saline (PTx) and incubated overnight (ON) at 4 °C. After several washes

in 1% bovine serum albumin (BSA) in PTx, samples were incubated with either an anti-

mouse peroxidase (POD) conjugated antibody (Millipore-Sigma, #11207733910, 1:100) or an

AlexaFluor594 conjugated antibody (ThermoFisher Scientific, #A32731) diluted in 5% NGS

in PTx ON at 4ºC. For samples incubated with mouse anti-double phosphorylated ERK1/2

and anti-mouse POD conjugated antibody, signal was developed using either

diaminobenzidine (DAB) (Vector Laboratories, #SK-4100) or Tyramide Signal Amplification

(TSA) kit (Akoya Biosciences, #NEL742001KT) following manufacturer's

recommendations. DAB-developed samples were stored in 70% glycerol and counterstained

with 4',6-diamidino-2- phenylindole (DAPI, ThermoFisher Scientific, #D3571, 1:1000)

nuclear marker. Samples for laser scanning confocal microscopy (LSCM) were

counterstained with DAPI diluted in 1% bovine serum albumin (BSA) in PTx for 1h at RT

and mounted in 70% glycerol.

Gene isolation and gene expression analyses

Candidate genes were amplified by two rounds of nested PCR using gene specific primers and

a T7 universal primer on cDNA from mixed developmental stages as initial template (Table

S9). Riboprobes were synthesized with the T7 enzyme (Ambion's MEGAscript

kit, #AM1334) and stored in hybridisation buffer at a concentration of 50 ng/µl.

Colorimetric and fluorescent whole-mount *in situ* hybridisation were performed following an

established protocol [31, 39]. After probe washes, samples for fluorescent *in situ*

hybridisation were incubated with anti-DIG-POD Fab fragments (ROCHE, #11633716001)

and a mouse anti-acetylated α-tubulin antibody (clone 6-11B-1, Millipore-446 Sigma,

#MABT868, 1:500) to co-stain cell boundaries. After signal development with the TSA kit

(Akoya Biosciences, #NEL742001KT), these samples were washed several times in PTx and

treated for secondary antibody incubation as described above.

Genome sequencing, gene annotation and RNA-seq developmental time course

High molecular weight genomic DNA was obtained from a single *O. fusiformis* male

following the agar plug extraction method and sequenced on a PacBio Sequel II platform to a

~200x coverage. Initial assembly was conducted with Canu [61], subsequently scaffolded

with BioNano optical mapping data generated on an Irys platform and finally de-haploidised

with purge_haplotigs [62]. To annotate protein coding genes, we used available

transcriptomic data (SRA accession number SRR1222288) and generated replicated stage-

specific RNA-seq data covering twelve developmental time-points, from the unfertilized

active oocyte until the mature mitraria larva. RNA-seq reads were mapped with STAR [63]

and gene models were generated combining Mikado v.2.1 [64], Augustus [65] and PASA [66]

pipelines. Gene completeness was assessed with BUSCO [67] (97.5% complete, 1.5%

fragmented, 1% missing). Trinotate [68] and Panther [69] were used to functionally annotate

gene predictions and obtain initial gene orthology assignments. To identify gene expression

dynamics during development, we used the stage specific RNA-seq generated for annotation,

representing the average expression of the two replicates with the package pheatmap v.1.0.12

available in R, were colour intensity shows the z-score value for each gene. All raw

sequencing data is available under the project with accession number XXXXXX in the

European Nucleotide Archive (ENA).

Assay for Transposase-Accessible Chromatin using sequencing (ATAC-seq)

Two replicated ATAC-seq libraries were generated from >50,000 cells at 5 hpf (850-900

coeloblastulae) following the Omni-ATAC protocol [70], employing a home-made Tn5

transposase [71] and amplifying for 12 cycles. Libraries were sequenced on an Illumina

Hiseq4000 platform, 75 base paired-end output mode. After adaptor removal with cutadapt

v.2.5 [72], reads were mapped with NextGenMap v.0.5.5 [73] and duplicates removed with

samtools v.1.9 [74]. BAM files were shifted using deepTools2 v.3.4.3 [75] and MACS2

v2.2.7.1 [76, 77] was used for peak calling with parameters -f BAMPE --min-length 100 --

max-gap 75 -q 0.01. Reproducible peaks ($p < 0.05$) were identified applying Irreproducible

Discovery Rate package v2.0.4.2 [78]. Transcription factor motif enrichment was performed

with HOMER v.4.11 [79] and transcription factor foot printing with TOBIAS v.0.12.0 [80].

To identify bound transcription factors on peaks of *cdx* and *delta*, we intersected the genomic

positions of peaks and bound transcription factors with BEDTools v.2.29.2 [81]. Genomic

tracks were plotted with pyGenomeTracks v.2.1 [82].

RNA-seq profiling and differential gene expression analyses

Total RNA was extracted using the Monarch total RNA Miniprep Kit (New England

Biolabs, #T2010) and used for Illumina stranded mRNA library prep. Sequencing was

performed on an Illumina HiSeq 4000 platform in 75 bases paired-end output mode,

generating ~20 million reads per sample. Adapter sequences and low-quality bases were

removed using Trimmomatic v.0.36 [83]. Cleaned reads were then mapped to *O.*

fusiformis reference genome annotation using Kallisto v.0.44.0 to generate gene expression

counts [84]. Differential gene expression analyses were performed independently for each

drug and computed in pairwise comparisons of different stages and conditions using the R

package DEseq2 v.1.38.0 [85]. Significancy threshold was adjusted to a p-value equal or

lower than 0.05 and a Log2 fold change greater than 1.5 or lower than -1.5 [86]. PCAs and

hierarchical clustering were plotted using ggplot2 and pheatmap R packages,

respectively [87]. Volcano plots were obtained with the package EnhancedVolcano, available

in R. Candidate genes for further gene expression analyses were selected based on being

differentially expressed in both drug treatments (U0126 and BFA) or highly differentially

expressed at 5.5 hpf in U0126 treated embryos. We further refined our selection based on

gene ontology (GO) categories, focusing on transcription factors and developmental genes

(Table S7).

GO and KEGG enrichment analyses

For GO mapping, the GO terms of differentially expressed genes (DEGs) associated with

each comparison were extracted from *O. fusiformis* reference genome annotation. GO

enrichment analyses were implemented using the GOseq R package, correcting for gene

length bias [88], and analysing up-regulated and down-regulated genes independently. GO

terms with corrected p-values lower than 0.05 were considered to be significantly enriched

(Table S7). A similar approach was followed for Kyoto Encyclopaedia of Genes and

Genomes (KEGG) pathway enrichment analyses, mapping differentially represented KO

numbers to their respective KEGG pathway [89] (Table S8).

Orthology assignment and domain architecture analyses

Generally, gene orthology was based on the functional annotation of *O. fusiformis* gene

repertoire, based on BLAST searches against SwissProt database and Panther searches [69,

90]. Protein sequences for ERK1/2 and ERK5, WNT ligands, RHOX and related

homeoboxes, DLL and JAGGED, FOXH and FOXF genes and FGFR and VEGFR were

retrieved by mining publicly available transcriptomes and databases. Multiple protein

alignments (Supplementary File 1) were constructed with MAFFT v.7 in automatic mode [91]

and poorly aligned regions were removed with gBlocks version 0.91b [92]. Maximum

likelihood trees were constructed with either RAxML v.8.2.11 [93] or FastTree [94] and

visualised with FigTree (https://github.com/rambaut/figtree/). InterProScan 5 [95] was used to

perform the protein domain composition of DLL.

Imaging

Representative embryos and larvae from colorimetric analyses were imaged with a Leica

DMRA2 upright epifluorescent microscope equipped with an Infinity5 camera (Lumenera),

using bright field and differential interference contrast optics. Fluorescently stained samples

were scanned with a Leica SP5 LSCM. Confocal stacks were analysed with Fiji and

brightness/contrast and colour balance were adjusted in Pixelmator Pro (v. 2.0.3) or

Photoshop (Adobe), applying changes always to the whole image and not parts of it. Final

figures were designed using Illustrator (Adobe).

References

1. Martinez Arias, A., and Steventon, B. (2018). On the nature and function of organizers. Development *145*.

2. Anderson, C., Khan, M.A.F., Wong, F., Solovieva, T., Oliveira, N.M.M., Baldock, R.A., Tickle, C., Burt, D.W., and Stern, C.D. (2016). A strategy to discover new organizers identifies a putative heart organizer. Nat Commun *7*, 12656.

3. Lyons, D.C., Perry, K.J., Lesoway, M.P., and Henry, J.Q. (2012). Cleavage pattern and fate map of the mesentoblast, 4d, in the gastropod *Crepidula*: a hallmark of spiralian development. Evodevo *3*, 21.

4. Kraus, Y., Aman, A., Technau, U., and Genikhovich, G. (2016). Pre-bilaterian origin of the blastoporal axial organizer. Nat Commun *7*, 11694.

5. Lapraz, F., Haillot, E., and Lepage, T. (2015). A deuterostome origin of the Spemann organiser suggested by Nodal and ADMPs functions in Echinoderms. Nat Commun *6*, 8434.

6. Laumer, C.E., Bekkouche, N., Kerbl, A., Goetz, F., Neves, R.C., Sorensen, M.V., Kristensen, R.M., Hejnol, A., Dunn, C.W., Giribet, G., et al. (2015). Spiralian phylogeny informs the evolution of microscopic lineages. Curr Biol *25*, 2000-2006.

7. Marlétaz, F., Peijnenburg, K., Goto, T., Satoh, N., and Rokhsar, D.S. (2019). A New Spiralian Phylogeny Places the Enigmatic Arrow Worms among Gnathiferans. Curr Biol *29*, 312-318 e313.

8. Martín-Durán, J.M., and Marletaz, F. (2020). Unravelling spiral cleavage. Development *147*.

9. Henry, J.Q. (2014). Spiralian model systems. Int J Dev Biol *58*, 389-401.

10. Hejnol, A. (2010). A twist in time--the evolution of spiral cleavage in the light of animal phylogeny. Integr Comp Biol *50*, 695-706.

592   11.   Schleip, W. (1929). Die Determination der Primitiventwicklung, (Leipzig:

593   Akademische Verlagsgesellschaft).

594   12.   Lambert, J.D. (2008). Mesoderm in spiralians: the organizer and the 4d cell. J Exp

595   Zool B Mol Dev Evol *310*, 15-23.

596   13.   Freeman, G., and Lundelius, J.W. (1992). Evolutionary implications of the mode of D

597   quadrant specification in coelomates with spiral cleavage. Journal of Evolutionary

598   Biology *5*, 205-247.

599   14.   Van Den Biggelaar, J.A.M., Van Loon, A.E., and Damen, W.G.M. (1995).

600   Mesentoblast and Trochoblast Specification in Species With Spiral Cleavage Predict

601   Their Phyletic Relations. Netherlands Journal of Zoology *46*, 8-21.

602   15.   Lavoie, H., Gagnon, J., and Therrien, M. (2020). ERK signalling: a master regulator of

603   cell behaviour, life and fate. Nat Rev Mol Cell Biol *21*, 607-632.

604   16.   Lambert, J.D., and Nagy, L.M. (2001). MAPK signaling by the D quadrant embryonic

605   organizer of the mollusc *Ilyanassa obsoleta*. Development *128*, 45-56.

606   17.   Lambert, J.D., and Nagy, L.M. (2003). The MAPK cascade in equally cleaving

607   spiralian embryos. Dev Biol *263*, 231-241.

608   18.   Lartillot, N., Lespinet, O., Vervoort, M., and Adoutte, A. (2002). Expression pattern of

609   *Brachyury* in the mollusc *Patella vulgata* suggests a conserved role in the

610   establishment of the AP axis in Bilateria. Development *129*, 1411-1421.

611   19.   Henry, J.J., and Perry, K.J. (2008). MAPK activation and the specification of the D

612   quadrant in the gastropod mollusc, *Crepidula fornicata*. Dev Biol *313*, 181-195.

613   20.   Koop, D., Richards, G.S., Wanninger, A., Gunter, H.M., and Degnan, B.M. (2007).

614   The role of MAPK signaling in patterning and establishing axial symmetry in the

615   gastropod *Haliotis asinina*. Dev Biol *311*, 200-212.

616   21.   Ozpolat, B.D., Handberg-Thorsager, M., Vervoort, M., and Balavoine, G. (2017). Cell

617   lineage and cell cycling analyses of the 4d micromere using live imaging in the marine

618   annelid *Platynereis dumerilii*. Elife *6*.

619   22.   Gline, S.E., Nakamoto, A., Cho, S.J., Chi, C., and Weisblat, D.A. (2011). Lineage

620   analysis of micromere 4d, a super-phylotypic cell for Lophotrochozoa, in the leech

621   *Helobdella* and the sludgeworm *Tubifex*. Dev Biol *353*, 120-133.

622   23.   Meyer, N.P., Boyle, M.J., Martindale, M.Q., and Seaver, E.C. (2010). A

623   comprehensive fate map by intracellular injection of identified blastomeres in the

624   marine polychaete *Capitella teleta*. Evodevo *1*, 8.

625   24.   Lanza, A.R., and Seaver, E.C. (2020). Activin/Nodal signaling mediates dorsal-ventral

626   axis formation before third quartet formation in embryos of the annelid *Chaetopterus*

627   *pergamentaceus*. Evodevo *11*, 17.

628   25.   Pfeifer, K., Schaub, C., Domsch, K., Dorresteijn, A., and Wolfstetter, G. (2014).

629   Maternal inheritance of *twist* and analysis of MAPK activation in embryos of the

630   polychaete annelid *Platynereis dumerilii*. PLoS One *9*, e96702.

631   26.   Amiel, A.R., Henry, J.Q., and Seaver, E.C. (2013). An organizing activity is required

632   for head patterning and cell fate specification in the polychaete annelid *Capitella*

633   *teleta*: new insights into cell-cell signaling in Lophotrochozoa. Dev Biol *379*, 107-122.

634   27.   Kozin, V.V., Filimonova, D.A., Kupriashova, E.E., and Kostyuchenko, R.P. (2016).

635   Mesoderm patterning and morphogenesis in the polychaete *Alitta virens* (Spiralia,

636   Annelida): Expression of mesodermal markers *Twist*, *Mox*, *Evx* and functional role for

637   MAP kinase signaling. Mech Dev *140*, 1-11.

638   28.   Nakamoto, A., Nagy, L.M., and Shimizu, T. (2011). Secondary embryonic axis

639   formation by transplantation of D quadrant micromeres in an oligochaete annelid.

640   Development *138*, 283-290.

29.   Lanza, A.R., and Seaver, E.C. (2020). Functional evidence that Activin/Nodal

signaling is required for establishing the dorsal-ventral axis in the annelid *Capitella*

teleta. Development *147*.

30.   Weigert, A., Helm, C., Meyer, M., Nickel, B., Arendt, D., Hausdorf, B., Santos, S.R.,

Halanych, K.M., Purschke, G., Bleidorn, C., et al. (2014). Illuminating the base of the

annelid tree using transcriptomics. Mol Biol Evol *31*, 1391-1401.

31.   Carrillo-Baltodano, A.M., Seudre, O., Guynes, K., and Martin-Duran, J.M. (2021).

Early embryogenesis and organogenesis in the annelid *Owenia fusiformis*. Evodevo

12, 5.

32.   Gonzales, E.E., van der Zee, M., Dictus, W.J., and van den Biggelaar, J. (2007).

Brefeldin A and monensin inhibit the D quadrant organizer in the polychaete annelids

Arctonoe vittata and *Serpula columbiana*. Evol Dev *9*, 416-431.

33.   Treadwell, A.L. (1901). The cytogeny of *Podarke obscura* Verrill. J. Morph. *17*, 399-

478.

34.   Mead, A.D. (1897). The early development of marine annelids. J. Morph. *13*, 227-326.

35.   Woltereck, R. (1904). Beiträge zur praktischen Analyse der *Polygordius*-Entwicklung

nach dem 'Nordsee' - und dem 'Mittelmeer-typus'. I. Der für beide Typen gleich

verlaufende Entwicklungsabschnitt: vom Ei bis zum jüngsten Trochophora-Stadium.

Arch. f. Entwicklungsmech. *18*, 377-403.

36.   Newby, W.W. (1940). The embryology of the echiuroid worm *Urechis caupo*,

Volume 16, (Philadelphia: The American Philosophical Society).

37.   Gonzales, E.E., van der Zee, M., Dictus, W.J., and van den Biggelaar, J. (2007).

Brefeldin A or monensin inhibits the 3D organizer in gastropod, polyplacophoran, and

scaphopod molluscs. Dev Genes Evol *217*, 105-118.

38.   Favata, M.F., Horiuchi, K.Y., Manos, E.J., Daulerio, A.J., Stradley, D.A., Feeser,

666   W.S., Van Dyk, D.E., Pitts, W.J., Earl, R.A., Hobbs, F., et al. (1998). Identification of

667   a novel inhibitor of mitogen-activated protein kinase kinase. J Biol Chem *273*, 18623-

668   18632.

669   39.   Martin-Duran, J.M., Passamaneck, Y.J., Martindale, M.Q., and Hejnol, A. (2016). The

670   developmental basis for the recurrent evolution of deuterostomy and protostomy. Nat

671   Ecol Evol *1*, 5.

672   40.   Hoodless, P.A., Pye, M., Chazaud, C., Labbe, E., Attisano, L., Rossant, J., and Wrana,

673   J.L. (2001). FoxH1 (Fast) functions to specify the anterior primitive streak in the

674   mouse. Genes Dev *15*, 1257-1271.

675   41.   Pogoda, H.M., Solnica-Krezel, L., Driever, W., and Meyer, D. (2000). The zebrafish

676   forkhead transcription factor FoxH1/Fast1 is a modulator of nodal signaling required

677   for organizer formation. Curr Biol *10*, 1041-1049.

678   42.   Yamamoto, M., Meno, C., Sakai, Y., Shiratori, H., Mochida, K., Ikawa, Y., Saijoh, Y.,

679   and Hamada, H. (2001). The transcription factor FoxH1 (FAST) mediates Nodal

680   signaling during anterior-posterior patterning and node formation in the mouse. Genes

681   Dev *15*, 1242-1256.

682   43.   Watanabe, M., and Whitman, M. (1999). FAST-1 is a key maternal effector of

683   mesoderm inducers in the early Xenopus embryo. Development *126*, 5621-5634.

684   44.   Pruitt, M.M., Letcher, E.J., Chou, H.C., Bastin, B.R., and Schneider, S.Q. (2014).

685   Expression of the wnt gene complement in a spiral-cleaving embryo and trochophore

686   larva. Int J Dev Biol *58*, 563-573.

687   45.   MacLean, J.A., 2nd, and Wilkinson, M.F. (2010). The Rhox genes. Reproduction *140*,

688   195-213.

689   46.   Hubaud, A., and Pourquié, O. (2014). Signalling dynamics in vertebrate segmentation.

690   Nat Rev Mol Cell Biol *15*, 709-721.

691   47.   Andrikou, C., and Hejnol, A. (2021). FGF signaling acts on different levels of

692   mesoderm development within Spiralia. Development.

693   48.   Sun, L., Tran, N., Liang, C., Tang, F., Rice, A., Schreck, R., Waltz, K., Shawver, L.K.,

694   McMahon, G., and Tang, C. (1999). Design, Synthesis, and Evaluations of Substituted

695   3-[(3- or 4-Carboxyethylpyrrol-2-yl)methylidenyl]indolin-2-ones as Inhibitors of

696   VEGF, FGF, and PDGF Receptor Tyrosine Kinases. Journal of Medicinal Chemistry

697   *42*, 5120-5130.

698   49.   Fuentealba, L.C., Eivers, E., Geissert, D., Taelman, V., and De Robertis, E.M. (2008).

699   Asymmetric mitosis: Unequal segregation of proteins destined for degradation. Proc

700   Natl Acad Sci U S A *105*, 7732-7737.

701   50.   Lambert, J.D., and Nagy, L.M. (2002). Asymmetric inheritance of centrosomally

702   localized mRNAs during embryonic cleavages. Nature *420*, 682-686.

703   51.   Gonsalves, F.C., and Weisblat, D.A. (2007). MAPK regulation of maternal and

704   zygotic Notch transcript stability in early development. Proc Natl Acad Sci U S A *104*,

705   531-536.

706   52.   Johnson, A.B., and Lambert, J.D. (2021). The Caudal ParaHox gene is required for

707   hindgut development in the mollusc *Tritia* (a.k.a. *Ilyanassa*). Dev Biol *470*, 1-9.

708   53.   Gharbiah, M., Nakamoto, A., Johnson, A.B., Lambert, J.D., and Nagy, L.M. (2014).

709   *Ilyanassa* Notch signaling implicated in dynamic signaling between all three germ

710   layers. Int J Dev Biol *58*, 551-562.

711   54.   Dorey, K., and Amaya, E. (2010). FGF signalling: diverse roles during early

712   vertebrate embryogenesis. Development *137*, 3731-3742.

55.   Green, S.A., Norris, R.P., Terasaki, M., and Lowe, C.J. (2013). FGF signaling induces

mesoderm in the hemichordate *Saccoglossus kowalevskii*. Development *140*, 1024-

1033.

56.   McGregor, A.P., Pechmann, M., Schwager, E.E., and Damen, W.G. (2009). An

ancestral regulatory network for posterior development in arthropods. Commun Integr

Biol *2*, 174-176.

57.   Martin, B.L., and Kimelman, D. (2009). Wnt signaling and the evolution of embryonic

posterior development. Curr Biol *19*, R215-219.

58.   Fritzenwanker, J.H., Uhlinger, K.R., Gerhart, J., Silva, E., and Lowe, C.J. (2019).

Untangling posterior growth and segmentation by analyzing mechanisms of axis

elongation in hemichordates. Proc Natl Acad Sci U S A *116*, 8403-8408.

59.   Dictus, W.J.A.G., and Damen, P. (1997). Cell-lineage and clonal-contribution map of

the trochophore larva of *Patella vulgata* (Mollusca). Mechanisms of Development *62*,

213-226.

60.   Lyons, D.C., Perry, K.J., and Henry, J.Q. (2015). Spiralian gastrulation: germ layer

formation, morphogenesis, and fate of the blastopore in the slipper snail *Crepidula*

fornicata. Evodevo *6*, 24.

61.   Koren, S., Walenz, B.P., Berlin, K., Miller, J.R., Bergman, N.H., and Phillippy, A.M.

(2017). Canu: scalable and accurate long-read assembly via adaptive k-mer weighting

and repeat separation. Genome Res *27*, 722-736.

62.   Roach, M.J., Schmidt, S.A., and Borneman, A.R. (2018). Purge Haplotigs: allelic

contig reassignment for third-gen diploid genome assemblies. BMC Bioinformatics

19, 460.

63.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,

Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.

Bioinformatics *29*, 15-21.

64.   Venturini, L., Caim, S., Kaithakottil, G.G., Mapleson, D.L., and Swarbreck, D. (2018).

Leveraging multiple transcriptome assembly methods for improved gene structure

annotation. Gigascience *7*.

65.   Stanke, M., Schoffmann, O., Morgenstern, B., and Waack, S. (2006). Gene prediction

in eukaryotes with a generalized hidden Markov model that uses hints from external

sources. BMC Bioinformatics *7*, 62.

66.   Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Jr., Hannick,

L.I., Maiti, R., Ronning, C.M., Rusch, D.B., Town, C.D., et al. (2003). Improving the

Arabidopsis genome annotation using maximal transcript alignment assemblies.

Nucleic Acids Res *31*, 5654-5666.

67.   Seppey, M., Manni, M., and Zdobnov, E.M. (2019). BUSCO: Assessing Genome

Assembly and Annotation Completeness. Methods Mol Biol *1962*, 227-245.

68.   Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru,

D., Lee, T.J., Leigh, N.D., Kuo, T.H., Davis, F.G., et al. (2017). A Tissue-Mapped

Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors.

Cell Rep *18*, 762-776.

69.   Thomas, P.D., Campbell, M.J., Kejariwal, A., Mi, H., Karlak, B., Daverman, R.,

Diemer, K., Muruganujan, A., and Narechania, A. (2003). PANTHER: a library of

protein families and subfamilies indexed by function. Genome Res *13*, 2129-2141.

70.   Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong,

N.A., Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., et al. (2017). An

760   improved ATAC-seq protocol reduces background and enables interrogation of frozen

761   tissues. Nat Methods *14*, 959-962.

762 71.   Picelli, S., Bjorklund, A.K., Reinius, B., Sagasser, S., Winberg, G., and Sandberg, R.

763   (2014). Tn5 transposase and tagmentation procedures for massively scaled sequencing

764   projects. Genome Res *24*, 2033-2040.

765 72.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

766   sequencing reads. 2011 *17*, 3.

767 73.   Sedlazeck, F.J., Rescheneder, P., and von Haeseler, A. (2013). NextGenMap: fast and

768   accurate read mapping in highly polymorphic genomes. Bioinformatics *29*, 2790-

769   2791.

770 74.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,

771   Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The

772   Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

773 75.   Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,

774   S., Dundar, F., and Manke, T. (2016). deepTools2: a next generation web server for

775   deep-sequencing data analysis. Nucleic Acids Res *44*, W160-165.

776 76.   Gaspar, J.M. (2018). Improved peak-calling with MACS2. bioRxiv, 496521.

777 77.   Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,

778   Nusbaum, C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of

779   ChIP-Seq (MACS). Genome Biol *9*, R137.

780 78.   Li, Q., Brown, J.B., Huang, H., and Bickel, P.J. (2011). Measuring reproducibility of

781   high-throughput experiments. The Annals of Applied Statistics *5*, 1752-1779, 1728.

782 79.   Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X.,

783   Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-

determining transcription factors prime cis-regulatory elements required for

macrophage and B cell identities. Mol Cell *38*, 576-589.

80. Bentsen, M., Goymann, P., Schultheis, H., Klee, K., Petrova, A., Wiegandt, R., Fust,

A., Preussner, J., Kuenne, C., Braun, T., et al. (2020). ATAC-seq footprinting unravels

kinetics of transcription factor binding during zygotic genome activation. Nat

Commun *11*, 4267.

81. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for

comparing genomic features. Bioinformatics *26*, 841-842.

82. Lopez-Delisle, L., Rabbani, L., Wolff, J., Bhardwaj, V., Backofen, R., Gruning, B.,

Ramirez, F., and Manke, T. (2021). pyGenomeTracks: reproducible plots for

multivariate genomic datasets. Bioinformatics *37*, 422-423.

83. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for

Illumina sequence data. Bioinformatics *30*, 2114-2120.

84. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal

probabilistic RNA-seq quantification. Nat Biotechnol *34*, 525-527.

85. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

86. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A

Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical

Society. Series B (Methodological) *57*, 289-300.

87. Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis, (Springer-Verlag

New York).

88. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology

analysis for RNA-seq: accounting for selection bias. Genome Biol *11*, R14.

89.   Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2016).

KEGG as a reference resource for gene and protein annotation. Nucleic Acids Res *44*,

D457-462.

90.   UniProt, C. (2021). UniProt: the universal protein knowledgebase in 2021. Nucleic

Acids Res *49*, D480-D489.

91.   Katoh, K., Rozewicki, J., and Yamada, K.D. (2019). MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. Brief Bioinform

20, 1160-1166.

92.   Talavera, G., and Castresana, J. (2007). Improvement of phylogenies after removing

divergent and ambiguously aligned blocks from protein sequence alignments. Syst

Biol *56*, 564-577.

93.   Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics *30*, 1312-1313.

94.   Price, M.N., Dehal, P.S., and Arkin, A.P. (2010). FastTree 2--approximately

maximum-likelihood trees for large alignments. PLoS One *5*, e9490.

95.   Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,

Maslen, J., Mitchell, A., Nuka, G., et al. (2014). InterProScan 5: genome-scale protein

function classification. Bioinformatics *30*, 1236-1240.

Figures

Figure 1. Di-phosphorylated ERK1/2 is enriched in the 4d cell in *O. fusiformis*. (**A**)

Spiralia is a major bilaterian lineage exhibiting spiral cleavage as the most probably

synapomorphy. Differently from other bilaterian lineages, body patterning and embryonic

organisation is controlled by a cell in the D-quadrant, which in molluscs exhibits and requires

active di-phosphorilated ERK1/2. Whether the molluscan condition is ancestral to Spiralia is

unclear. (**B**) *O. fusiformis* is a member of Palaeoannelida, the sister group to all remaining

annelids, and undergoes typical equal spiral cleavage (**C**). The transition from the 4-cell to the

8-cell stage is characterised by a dextral shift of the mitotic spindle at a 45° angle relative to

the animal–vegetal axis and generates a quartet of slightly larger animal cells (the

micromeres, 1q) and a quartet of vegetal cells (the macromeres, 1Q). (**D**) Active di-

phosporylated-ERK1/2 (yellow) is enriched in four animal cells ($1q^{111}$) at 4 hours post

fertilisation (hpf), in a single cell (the 4d micromere) at 5hpf, and in six cells of the 2a-c

lineage and 4d at 6 hpf. DAPI: cyan. (**E**) Dynamics of cleavage and quadrant identities during

the onset and early division of 4d. The fourth micromere quartet emerges asymmetrically

(early 4 hpf), which is soon followed by the asymmetric division of one of the 3q micromeres

(arrowheads) (late 4 hpf). The vegetal pole is however quatri-radial at 5 hpf, with the 4d

arresting its division with respect to 4a–c, and only cleaving into MR and ML after ingression

at 7 hpf. F-actin (grey) shows cell boundaries and DAPI stains the nuclei. False colouring

indicates cell or quadrant identities. Asterisks in (**C**) and (**D**) point to the animal pole, and to

part of the 4d progeny in (**E**). Descriptions for (**D**) and (**E**) are based on at least 10 embryos

per stage, except at 7 hpf (n = 3). Drawings are not to scale. Scale bars are 50 μm.

Figure 2. ERK1/2 signalling controls axial polarity in *O. fusiformis*. (**A**) ERK1/2

signalling cascade responds to cell-cell communication and involves the di-phosphorylation

of ERK1/2 (di-P-ERK1/2) via MEK. Brefeldin A (BFA), a protein trafficking inhibitor, and

U0126, a specific MEK inhibitor, can block ERK1/2 di-phosphorylation. (**B**) Immunostaining

against di-P-ERK1/2. BFA and U0126 treatments from 0.5 hours post fertilisation (hpf) to 5

hpf prevent di-P-ERK1/2 enrichment in the 4d cell at 5 hpf. (**C**) BFA and U0126 treatment

(0.5 hpf to 5 hpf) result in anteroventrally radialised larvae that fail to form posterior

structures (e.g., hindgut, mesodermal bands and chaetae). Main panels are lateral z-stack

projections of confocal images and insets in treated embryos are ventral views. Cilia is

labelled with anti-acetylated tubulin. (**D**) Whole mount *in situ* hybridisation on control and

U0126 treated (0.5 to 5 hpf) larvae. Treated radial larvae show reduced apical markers

(*six3/6*, and *syt-1* in the inset), radially expanded anterior markers (*gsc*), and absent posterior

endomesodermal (*cdx*, *twist*) but not a midgut gene (GATA4/5/6b). (**E**) Schematic

representation of drug treatment windows and observed larval phenotypes (WT, wild type; R,

radial). (**F**) Morphological analysis of the compressed larval phenotype obtained after BFA

treatment from 0.5 hpf until 3–4 hpf. Treated larvae have a reduced apical organ and apical

tuft (first row) and reduced expression of apical gene markers (*six3/6*, and *syt-1* in the inset),

but other tissues are not affected. The larval blastocele is somehow obliterated. Asterisks

point to the apical pole. Schematic drawings are not to scale. ao: apical organ; an: anus; ch:

chaetae; mg: midgut; mo: mouth; pt: prototroch. Scale bars are 50 μm.

Figure 3. ERK1/2 signalling activates posterodorsal and mesodermal genes. (**A**)

Experimental design for RNA-seq collections. Solid colour bars show period of control

(DMSO), BFA or U0126 treatments. Red circles show sample collection, at the coeloblastula

stage (5.5 hpf) and at the larval stage (24 hpf). (**B**) Number of differentially expressed (DE)

genes in different conditions and comparisons, where red bars are downregulated genes and

blue bars are upregulated genes. (**C**) Volcano plots for BFA and U0126 treated coeloblastula.

Red dots show DE genes, with candidate genes labelled for each comparison. (**D**) Temporal

and spatial time course of expression of candidate genes affected by 4d misspecification and

associated with apical/anterior, chaetae/posterior, hindgut or mesoderm development. The

heatmap depicts the normalised z-score value of expression for each gene throughout

development, with red blocks indicating high levels of expression and blue blocks indicating

low levels of expression. The horizontal dotted line highlights the timing of 4d specification.

On the right, whole mount *in situ* hybridisation of a subset of these 22 candidate genes at the

coeloblastula (5 hpf), gastrula (9 hpf) and larval stages (24 hpf). (**E**) Only three of the

candidate genes (*cdx*, *foxH* and *fer3*) are downregulated in all stages and treatment

comparisons. Validation via *in situ* hybridisation demonstrates the expression is lost after

drug treatment. Pictures for *cdx* at the coeloblastula and larva in (**D**) and (**E**) are the same.

Asterisks point to the animal/apical pole, and arrowheads to the domains of expression.

Schematic drawings are not to scale. bp: blastopore; mo: mouth.

Figure 4. ERK1/2 signalling specifies and patterns the D-quadrant. (**A**) Whole mount *in*

situ hybridisation of a subset of genes (ERK1/2-dependent and independent) patterning the

vegetal pole at (5 hpf) and after (5.5 hpf) 4d specification. The row below is a schematic

representation of the vegetal pole depicting gene expression (in green) at 5.5 hpf. (**B**) The

ParaHox gene *cdx* (yellow) is expressed in the 4d cell at the coeloblastula stage and in the

progeny of MR and ML at the gastrula stage, corroborating the endodermal sublineage of 4d.

(**C**) The Notch ligand *delta* (yellow) is expressed in 4d, multiple cells from the D-quadrant,

and a few animal micromeres from the other three quadrants at the coeloblastula stage. In (**B**,

C) Nuclei (DAPI) are shown in cyan and cell boundaries (acetylated tubulin) in magenta.

Asterisks point to the animal pole. (**D**) Distribution of known bound transcription factors in

open chromatin regions around *cdx* (upper part) and *delta* (lower part) loci at 5 hpf. Dark and

light blue depict the two ATAC-seq replicates. (**E**) Number of cellular contacts for each of the

4q micromeres at 5.5 hpf based on the analysis of eight embryos. While $4a^1$–$4c^2$ establish 8

direct cell contacts, 4d touches 13 cells of the D and C quadrants. (**F**) Inhibition of ERK1/2

di-phosphorylation by U0126 prevents D-quadrant specification, as shown by the expansion

of the anterior marker *gsc* and loss of the posterior marker *AP2*. The 4d micromere is no

longer recognised, as all 4q micromeres cleave at the same time (asterisks). However, *gsc* is

lost in BFA treated embryos, and thus needs of inductive signals. Gene expression is shown in

yellow, nuclei (DAPI) in cyan and cell boundaries (acetylated tubulin) in magenta or grey. In

all panels, arrowheads point to expression domains. Schematic drawings are not to scale. gp:

gastral plate.

Figure 5. FGFR signalling regulates ERK1/2 di-phosphorylation in 4d. (**A**) The drug

SU5402 specifically inhibits receptors tyrosine kinase of the type FGF/VEGF, which can

signal through the ERK1/2 signalling cascade. (**B**) Whole mount immunohistochemistry

against di-phosphorylated ERK1/2 (di-P-ERK1/2). SU5402 treated embryos lose ERK1/2

enrichment in a vegetal micromere at 5 hours post fertilisation (hpf). (**C**) Schematic

representation of drug treatment windows and observed larval phenotypes (WT, wild type).

(**D**) Morphological and molecular patterning analysis of SU5402 treated embryos (4 to 6 hpf)

fixed at 24 hpf. The first column shows diagrams of lateral views of the wild type in DMSO

controls and radial larvae in treated embryos. Second and third columns show larvae labelled

with DAPI (cyan) and F-actin (magenta) plus acetylated tubulin staining (yellow),

respectively. Fourth column onwards are whole mount in situ hybridisations. SU5402

treatment from 4 hpf to 6 hpf results in phenocopies of U0126 and BFA radial larvae. Insets

are ventral views. Asterisks point to the apical pole. Schematic drawings are not to scale. ao:

apical organ; an: anus; ch: chaetae; mg: midgut; mo: mouth; pt: prototroch.

Figure 6. ERK1/2 activity in the D-quadrant embryonic organiser is an ancestral

spiralian trait. (**A**) Schematic model of FGFR/ERK1/2 activity in the 4d micromere during

late spiral cleavage in *O. fusiformis*. At 4 hours post fertilisation (hpf), di-phosphorylated

ERK1/2 (di-P-ERK1/2) is enriched in the apical most micromeres ($1q^{111}$). At 5 hpf, an

unidentified signal activates FGF receptor (FGFR) which di-phosphorylates ERK1/2 in the 4d

micromere. This activates *cdx* and *delta* in 4d (becoming the endomesodermal precursor) and

correlates with 4d arresting its cell cycle progression, as 4a–4c cleave. ERK1/2 activation in

4d is required to limit an anteriorising signal (Signal X) to the A–C quadrants, which activates

gsc expression in these quadrants. In addition, ERK1/2 activity in 4d induces the expression

of a battery of genes in the D-quadrant and part of the C-quadrant that probably induce lateral

mesodermal and posterodorsal fates, ultimately controlling axial elongation. This inductive

activity might happen via the Notch pathway. (**B**) The similarities in ERK1/2 activation

patterns between *O. fusiformis* and other molluscan embryos suggest this is a shared ancestral

trait to spiral cleavage. More broadly, ERK1/2 activity induces the expression of a set of

genes and signalling pathways that are repeatedly involved in posterior specification and

anteroposterior growth in bilaterian embryos. Drawings are not to scale.

Supplemental Information

Figure S1. ERK1/2 activity during *O. fusiformis* early embryogenesis. (A) Maximum

likelihood orthology assignment for ERK1/2, using ERK5 as outgroup. Only bootstrap values

for major nodes are shown. (**B**) DEseq2 normalised expression of *erk1/2* during *O. fusiformis*

development, from the active oocyte to the mitraria larval stage. The time of the specification

of 4d is highlighted with a schematic drawing. Coloured dots indicate values of expression for

each replicate and the dotted red line shows the mean value of expression. (**C**) Whole mount

immunohistochemistry against di-phosphorylated-ERK1/2 (dark orange) during spiral

cleavage in *O. fusiformis*. ERK1/2 is active at low levels during early divisions, and later

enriched in the four 1q[111] animal micromeres at 4 hours post fertilisation (hpf), the 4d

micromere at the time of its specification (5 hpf) and finally in six vegetal cells forming a

bilaterally symmetrical pattern (arrowheads) at 6 hpf. Insets in (**C**) show either nuclear

staining (blue) or lateral views. Asterisks point to the animal pole.

Figure S2. The effect of BFA and U0126 treatment in *O. fusiformis* development. (A)

Percentage of coeloblastula showing di-phosphorylated ERK1/2 (di-P-ERK1/2) enrichment in

the 4d micromere after 0.5 to 5 hours post fertilisation (hpf) treatment with brefeldin A

(BFA), U0126 or DMSO (control). (**B**) Percentage of radial phenotypes obtained when

treating embryos from 0.5 to 5 hpf with a range of BFA and U0126 concentrations. (**C**)

Whole mount in situ hybridisation of apical (*six3/6*), oral (*gsc*), posterior/hindgut (*cdx*),

midgut (*GATA4/5/6b*) and trunk mesodermal (*twist*) gene markers in control and treated

larvae. Radial phenotypes have a reduction of the apical marker, radial expansion of an oral

gene, loss of posterior structures (chaetae) and posterior/trunk mesoderm gene expression and

retain the expression of a midgut gene. Radial larvae treated with U0126 from the 2-cell stage

to 3 hpf show scattered *twist* expression in some embryos (arrowheads). The compressed

larval phenotype after BFA treatment from 0.5 to 4 hpf have a reduced apical organ and

apical tuft (first row) and reduced expression of apical markers (*six3/6*), but otherwise normal

morphology besides an obliterated internal blastocele. Asterisks indicate the apical pole and

arrowheads point to expression domains. Insets are ventral views. mo: mouth.

Figure S3. Differential expression analyses in BFA and U0126 treated embryos. (**A**)

Hierarchical pairwise correlation matrix between brefeldin A (BFA) and U0126 treated and

control samples. (**B**) Principal component analysis (PCA) plot for BFA (left) and U0126

(right) RNA-seq analyses. (**C**) Volcano plots for BFA and U0126 treated larva. Red dots

show differentially expressed (DE) genes and analysed candidate genes are labelled in each

comparison. (**D**) Venn diagrams showing the number of up- (top) and downregulated

(bottom) DE genes shared between different conditions. (**E**) Ten most overrepresented KEGG

pathways in each of the four conditions analysed by RNA-seq. The size of the coloured dots

is proportional to the number of overrepresented sub-categories nested into a given KEGG

pathway.

Figure S4. Validation of the differential expression analyses. (**A**–**G**) Whole mount in situ

hybridisation of control and treated embryos showing in addition a wild type gastrula stage to

reconstruct the time course of expression of each gene. In all cases, the expression patterns

disappear in BFA and U0126 treated embryos. Insets are ventral views. Asterisk point to the

animal/apical pole. Arrowheads point to 4d in (**A**) and to the apical organ in (**F**). bp:

blastopore; gp: gastral plate; mo: mouth.

Figure S5. Genes patterning the D-quadrant after the specification of the 4d-organiser

cell. (**A**) Schematic drawing of the cellular arrangements and cell nomenclature of the vegetal

pole just after the specification of the 4d micromere in *O. fusiformis* embryo at 5.5 hours post

fertilisation (hpf). Different colours depict different quadrants. (**B**) Time course of expression

of *cdx*, *foxH* and *fer3* via whole mount *in situ* hybridisation from 6 hpf to 18 hpf during *O.*

fusiformis embryogenesis. While *cdx* is expressed in the cells forming the hindgut, *foxH*

appears to upregulate at the earliest stages of cells giving rise to mesodermal derivatives. The

gene *fer3* is expressed in two posterior most cells of unknown function in the gastrula and

larva. Insets are lateral views. (**C**) The protein domain architecture of the Notch ligand

DELTA in *O. fusiformis* in comparisons with the domain structures of the Delta orthologs in

D. melanogaster and *H. sapiens*. (**D**–**G**) Maximum likelihood orthology assignments of

DELTA, FOXH, WNT ligands, and RHOX in *O. fusiformis*. Only bootstrap values supporting

each major clade are shown. bp: blastopore; mo: mouth.

1008

Figure S6. The FGF pathway and FGFR activity during 4d specification in *O.*

***fusiformis*.** (**A**) Maximum likelihood orthology assignment of fibroblast growth factor

receptor (FGFR) and vascular endothelial growth factor receptor (VEGFR). Only bootstrap

values for key nodes are shown. (**B**) Multiple protein alignment between *O. fusiformis* and *H.*

sapiens FGFR and VEGFR orthologs indicating conserved residues for the interaction of the

drug SU5402 with these receptors. (**C**) Temporal time course of expression of members of the

FGF and VEGF pathways in *O. fusiformis*. The heatmap depicts the normalised z-score value

of expression for each gene, with red blocks indicating high levels of expression and blue

blocks indicating low levels of expression. The horizontal dotted line highlights the

specification of the 4d micromere. (**D**) Whole mount in situ hybridisation for *FGFR* and

VEGFR at the coeloblastula stage of *O. fusiformis*. *FGFR* is expressed weakly at the gastral

plate at the time of 4d-instructed posterior re-patterning. (**E**) Proportion of treated and control

embryos exhibiting di-phosphorylated ERK1/2 enrichment in the 4d cell at 5 hpf after

SU5402 drug treatment from 0.5 to 5 hours post fertilisation (hpf). (**F**) Morphological and

molecular characterisation of SU5402 treated embryos fixed at the larval stage. The first

column shows diagrams of lateral views of the wild type, compressed and radial phenotypes.

From the second column onwards, whole mount *in situ* hybridisations of control and SU5402

treated larvae. The compressed larvae are phenocopies of brefeldin A (BFA) compressed

larvae, with reduced apical organ and apical tuft (second row) and reduced expression of

apical markers (*six3/6*), but other tissues are not affected. Radial larvae are phenocopies of

BFA and U0126 radial larvae with reduction of apical markers (*six3/6*), radial expansion of

oral genes (*gsc*), loss of posterior structures (chaetae) and expression of posterior

endomesodermal markers (*cdx*, *twist*), but normal expression of an endodermal gene

(*GATA4/5/6b*). Insets are ventral views. Asterisks point to the animal/apical pole. ao: apical

organ; an: anus; ch: chaetae; gp: gastral plate; mg: midgut; mo: mouth; pt: prototroch. Scale

bars are 50 μm.

Supplementary Tables and Files

Table S1. Di-P-ERK1/2 activation in spiralians.

Table S2. Di-P-ERK1/2 activation at 5 hpf in control, BFA and U0126 treated embryos.

Table S3. Larval phenotypes in control, BFA and U0126 treatments.

Table S4. Larval phenotypes at different concentrations of BFA and U0126.

Table S5. Larval phenotypes during staggered time windows of BFA and U0126 treatments.

Table S6. List of all differentially expressed genes at 5.5 hpf and 24 hpf.

Table S7. List of enriched gene ontology terms.

Table S8. List of enriched KEGG pathways.

Table S9. List of candidate genes and their gene IDs in differential gene expression analyses.

File S1. Multiple protein alignments

A

Embryonic organisers & body patterning

Bilateria

Deuterostomia
Vertebrates, sea urchin

X. laevis — Spemann-Mangold organiser — V, D
P. lividus — V, Ventral ectoderm, D

Ecdysozoa
Nematode, fly

C. elegans — ABa, ABb, EMS, P2
D. melanogaster — A, P

Spiral cleavage

? Embryo patterning

Lophotrochozoa

Gnathifera
Platyhelminthes
Lophophorata
Annelida
Mollusca
Nemertea

V, D — D-quadrant organiser → di-P-ERK1/2 (P, P)

? Spiralia ?

B

Annelida — Sedentaria, Errantia, Palaeoannelida

Owenia fusiformis

C

O. fusiformis early development

Fertilisation — 3 — 6 — 9 hpf

2-cells → 16-cells
4-cells: 1q, 1Q, Quadrant *
8-cells *
>32-cells *
Coeloblastula: Blastocele *
Gastrula: Archenteron, Blastopore *

D

>32-cell stage — 4 hpf
Coeloblastula — 5 hpf
Early gastrula — 6 hpf

Lateral view

1q^{111}, *
3d^{?}, 2d^{11}, 4a, 4d, 4D
*, 2d^{11}, 4d

Animal-Vegetal view

Animal: 1q^{111}
3c, 4C, 4D, 3d, 1d^{222}, 2d^{11}, 2d^{12}
Vegetal: 1d^{222}, 1d^{221}

Vegetal: 2b^{11}, 2b^{12}, 4b, 4a^{12}, 4a^{11}, 2a^{11}, 2a^{12}, 4c, 2c^{12}, 2c^{11}, 4d

di-P-ERK1/2 / DAPI

E

Early 4 hpf
4q, 3q, 3Q, 2d^{11}, 1q^{111}
Late 4 hpf
4q, 3q, 3Q, 3q^{?}, 2d^{21}, 1q^{?}

5 hpf
4q, 4Q
6 hpf
4q, 5b, 3c, 3d, 4d, 3q

7 hpf
4q, 4d, MR, ML
9 hpf
4q, MR, ML

F-actin / DAPI

C-quadrant · D-quadrant · 4q micromere · 4d and progeny

A

	Autonomous	di-P-ERK1/2 dependent								di-P-ERK1/2 independent
	GATA4/5/6a	*cdx*	*delta*	*foxH*	*wnt1*	*wntA*	*rhox*	*fer3*	*AP2*	*gsc*

5 hpf
4d (di-P-ERK1/2)

5.5 hpf

DAPI AcTub

Schematic expression

Cell identity:
- *GATA4/5/6a*: 5A-D, 5a-5d, 4a¹-4c¹, 4a²-4c², 4d
- *cdx*: 4d
- *delta*: 1c²²², 1d²²¹, 2d²¹¹¹, 2d²¹¹², 2d¹¹¹², 2d¹¹¹¹, 4d
- *foxH*: 3c²², 3d²²
- *wnt1*: 3c²¹, 3c²², 3d²¹, 3d²², 3c¹¹, 3c¹², 3d¹¹, 3d¹²
- *wntA*: 2d¹¹², 2d²¹², 3c²¹, 3c²², 3d²¹, 3d²², 3c¹¹, 3c¹², 3d¹¹, 3d¹²
- *rhox*: 2d¹¹², 2d²¹²
- *fer3*: 1c²²², 1d²²¹
- *AP2*: 1c²²², 1d²²¹, 2d¹¹², 2d²¹¹², 2d²¹¹¹, 2d¹¹¹², 2d¹¹¹¹
- *gsc*: 2a¹¹²-2c¹¹², 2a¹²-2c¹², 2a²¹²-2c²¹², 2a²²-2c²²

B

Coeloblastula — *cdx* DAPI AcTub — Lateral — gp, 4d
Gastrula (9 hpf) — Animal — MR, ML

C

Coeloblastula — *delta* DAPI AcTub
Animal
Lateral — 4d

D

cdx ~13 kb
Gene
ATAC-seq (324.6)
Bound motifs: CDX, FOX, RUNX, SOX, OCT, GATA, E2A, ETS

delta ~56 kb
Gene
ATAC-seq (955.1)
Bound motifs: E2A, ETS, HNF4, bHLH, SOX, FOX, TBX, NF-Y, FOSL2

E

No cellular contacts
6 8 10 12 14

4a¹
4a²
4b¹
4b²
4c¹
4c²
4d

F

Control — *gsc* DAPI AcTub | BFA — *gsc*
U0126 — *gsc* | U0126 — 4q AcTub
Control — *AP2* | U0126 — *AP2* — 4q

A

ERK1/2
- *H. sapiens* ERK2
- *M. musculus* ERK2
- *M. musculus* ERK1
- *H. sapiens* ERK1
- *S. purpuratus* ERK1/2
- *N. vectensis* ERK1/2
- *S. maritima* ERK1/2
- *C. gigas* ERK1/2
- *L. gigantea* ERK1/2
- *H. robusta* ERK1/2
- *O. fusiformis* ERK1/2
- *C. teleta* ERK1/2
- *L. anatina* ERK1/2
- *B. floridae* ERK1/2

ERK5
- *L. gigantea* ERK5
- *C. gigas* ERK5
- *O. fusiformis* ERK5
- *C. teleta* ERK5
- *L. anatina* ERK5
- *B. lanceolatum* ERK5
- *S. purpuratus* ERK5
- *N. vectensis* ERK5
- *M. musculus* ERK5
- *H. sapiens* ERK5

0.2

B

di-P-ERK1/2
in 4d cell

O. fusiformis erk1/2

Replicate 1
Replicate 2

2 hpf 4 hpf

Normalised expression

0 hpf 5 hpf 9 hpf 13 hpf 18 hpf 27 hpf

Developmental time

C

di-P-ERK1/2

Zygote * — Lateral
2-cell * — Lateral
4-cell * — Animal
8-cell * — 1q, 1Q — Lateral

3 hpf — Animal
4 hpf * 1q¹¹¹ — Animal
5 hpf * 4d — Vegetal
6 hpf * 4d — Vegetal