An Arginine-Rich Motif in the ORF2 Capsid Protein Regulates the Hepatitis E Virus Lifecycle and Interactions with the Host Cell

Kévin Hervouet,[1,¶] Martin Ferrié,[1,¶] Maliki Ankavay,[1,¶,#] Claire Montpellier,[1] Charline Camuzet,[1] Virginie Alexandre,[1] Aïcha Dembélé,[1] Cécile Lecoeur,[1] Arnold Thomas Foe,[1] Peggy Bouquet,[2] David Hot,[2] Thibaut Vausselin,[1] Jean-Michel Saliou,[2] Sophie Salomé-Desnoulez,[2] Alexandre Vandeputte,[2] Laurent Marsollier,[3] Priscille Brodin,[1,2] Marlène Dreux,[4] Yves Rouillé,[1] Jean Dubuisson,[1] Cécile-Marie Aliouat-Denis,[1] Laurence Cocquerel,[1,*]

[1]University of Lille, CNRS, INSERM, CHU Lille, Pasteur Institute of Lille, U1019-UMR 9017-CIIL- Center for Infection and Immunity of Lille, Lille, France

[2]Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, UMR2014 - US41 - PLBS-Plateformes Lilloises de Biologie & Santé, Lille, France.

[3]Equipe ATIP AVENIR, CRCINA, INSERM, University of Angers, Angers, France.

[4]CIRI - Centre International de Recherche en Infectiologie, Univ Lyon, Université Claude Bernard Lyon 1, Inserm-U1111, CNRS-UMR5308, ENS-Lyon, Lyon, France.

¶ These authors contributed equally to this work.

Present address: Division of Gastroenterology and Hepatology, Institute of Microbiology, Lausanne, Switzerland

* Corresponding author, laurence.cocquerel@cnrs.fr

Abstract

Hepatitis E virus (HEV) infection is the most common cause of acute viral hepatitis worldwide. Hepatitis E is usually asymptomatic and self-limiting but it can become chronic in immunocompromised patients and is associated with increased fulminant hepatic failure and mortality rates in pregnant women. HEV genome encodes three proteins including the ORF2 protein that is the viral capsid protein. Interestingly, HEV produces 3 isoforms of the ORF2 capsid protein which are partitioned in different subcellular compartments and perform distinct functions in the HEV lifecycle. Notably, the infectious ORF2 (ORF2i) protein is the structural component of virions, whereas the genome-free secreted and glycosylated ORF2 proteins likely act as a humoral immune decoy. Here, by using a series of ORF2 capsid protein mutants expressed in the infectious genotype 3 p6 HEV strain as well as chimeras between ORF2 and the CD4 glycoprotein, we demonstrated how an Arginine-Rich Motif (ARM) located in the ORF2 N-terminal region controls the fate and functions of ORF2 isoforms. We showed that the ARM controls ORF2 nuclear translocation, promoting regulation of host antiviral responses. This motif also regulates the dual topology and functionality of ORF2 signal peptide, leading to the production of either cytosolic infectious ORF2i or reticular non-infectious glycosylated ORF2 forms. It serves as maturation site of glycosylated ORF2 by cellular proteases, and promotes ORF2-host cell membrane interactions. The identification of ORF2 ARM as a unique central regulator of the HEV lifecycle uncovers how viruses settle strategies to condense their genetic information and hijack cellular processes.

Author summary

Hepatitis E virus (HEV) is the major cause of acute viral hepatitis worldwide. Although infection with HEV is usually self-resolving, it can cause up to 30% mortality in pregnant women in the third trimester. There is no specific treatment nor universal vaccine to fight against HEV. In our study, we focused on the HEV ORF2 capsid protein which is produced as different forms that perform distinct functions in the HEV lifecycle. The infectious ORF2i form is the structural component of virions, while the other forms likely act as an immune decoy. Herein, we deciphered the molecular determinants of ORF2 multifunctionality. We identified an Arginine-Rich Motif (ARM) located in the ORF2 N-terminal region that controls the subcellular localization, the fate and functions of ORF2 forms. It also promotes ORF2-host cell interactions. Our observations highlight the ORF2 ARM as a unique central regulator of ORF2 addressing that finely controls the HEV lifecycle.

Introduction

Hepatitis E virus (HEV) is the most common cause of acute viral hepatitis worldwide and is an emerging problem in industrialized countries. This virus causes about 20 million infections annually [1]. While HEV infection is asymptomatic for most patients, some human populations including pregnant women and immunocompromised patients have higher risk to develop severe forms and chronic infections, respectively. HEV strains infecting humans have been classified into 4 main distinct genotypes (gt) belonging to a single serotype [2]. Gt1 and gt2 that infect humans only, are primarily transmitted through contaminated drinking water and are responsible for waterborne hepatitis outbreaks in developing countries. In contrast, gt3 and gt4 are zoonotic and are largely circulating in industrialized countries. They are mainly transmitted by contact with swine and consumption of inadequately heated pork products [3]. There is no specific treatment nor universal vaccine to fight against HEV.

HEV is a quasi-enveloped [4,5], positive-sense RNA virus expressing three open reading frames (ORFs): ORF1, ORF2 and ORF3 [6]. ORF1 encodes the ORF1 non-structural polyprotein that is the viral replicase [7]. ORF2 encodes the ORF2 viral capsid protein and ORF3 encodes a small protein that is involved in virion morphogenesis and egress [8,9] . Since studying the HEV lifecycle has long been hampered by the absence of efficient systems to amplify HEV, many steps of the HEV lifecycle remain poorly understood [10]. By combining the gt3 p6 strain [11] and a highly transfectable subclone of PLC/PRF/5 cells (PLC3 cells), we previously described an efficient HEV cell culture system [12]. This model notably enabled the pioneering demonstration that, during its lifecycle, HEV produces at least 3 forms of the ORF2 capsid protein: infectious ORF2 (ORF2i), glycosylated ORF2 (ORF2g), and

cleaved ORF2 (ORF2c). The ORF2i protein is the structural component of infectious particles. It is not glycosylated and is likely derived from the assembly of the intracellular ORF2 (ORF2intra) form present in the cytosolic compartment. Importantly, we showed that a fraction of the ORF2intra form is translocated into the nucleus of infected cells [13]. Nuclear localization of ORF2 was also described by Lenggenhager et al. in naturally infected human hepatocytes [14], but the significance of ORF2 nuclear localization remains to be elucidated. The ORF2g and ORF2c proteins (herein referred to as ORF2g/c) are highly glycosylated and secreted in large amounts in culture supernatant (i.e., about 1000x more than ORF2i [15]) and are the most abundant antigens detected in patient sera [12]. These proteins likely act as a humoral immune decoy that inhibits antibody-mediated neutralization [15]. How the different forms of ORF2 are generated during the HEV lifecycle has not yet been fully investigated. However, their sequence and post-translational modifications suggest that they might be produced either by a distinct addressing into the secretory pathway and the nucleus [12,13], and/or by a differential translation process[15].

Here we investigated the mechanisms by which the ORF2 forms are produced and differentially addressed to cell compartments. We demonstrated that HEV has set up a nucleo-cytoplasmic transport mechanism of its capsid protein to modulate cell host immune responses. In addition, we found that during the HEV lifecycle, a fine-tuning of ORF2 partitioning occurs between cytosolic, reticular and nuclear compartments. Importantly, we identified a stretch of 5 amino acid residues (herein referred to as ARM, Arginine-Rich Motif) in the N-terminal region of the ORF2 protein that drives nuclear translocation and tightly modulates the stoichiometry between the different ORF2 forms, especially by regulating the functionality of the ORF2 signal peptide and interactions with the host cell.

Results

The ORF2 protein transits through the nucleus in the early phase of infection.

We and others previously showed that the ORF2 protein is translocated into the nucleus of infected cells of patient liver biopsies [14] and in cell culture system [13]. Here, we first analyzed the ORF2 expression in PLC3 cells electroporated with the infectious p6 strain (PLC3/HEV-p6), at different time post-electroporation (p.e.) by immunofluorescence (**Fig 1**). ORF2 staining and quantification of nuclear fluorescence showed that nuclear translocation of ORF2 takes place at early time points p.e. (*i.e.,* 18h) and is then followed by a nuclear export process, indicating that HEV has developed mechanisms for ORF2 nuclear import and export.

The ORF2 protein displays an Arginine-Rich Motif (ARM) that functions as a Nuclear Localization Signal (NLS).

To decipher the molecular mechanisms of ORF2 nuclear import, we first analyzed its amino acid (aa) sequence with the NLSTradamus prediction program [16]. We identified a potential Nuclear Localization Signal (NLS) corresponding to a conserved Arginine-Rich Motif (ARM, 5 Arginine residues: RRRGRR) in the N-terminal region of ORF2, downstream of its signal peptide (SP) (**Fig 2A**). We next generated a series of ORF2 mutants in the p6 strain that are depicted in **Fig 2A**. We characterized their expression and subcellular localization (**Figs 2B, 2C, S1** and **S2**), and their impact on the HEV lifecycle (**Fig 2D**).

The replacement of arginine residues by alanine (3R/3A, 2R/2A and 5R/5A mutants) led to a drastic reduction of ORF2 nuclear localization compared to the wt protein (**Figs 2B** and **2C**, Nuclear extract), indicating that the ARM is likely a functional NLS.

Interestingly, the reduced nuclear localization of these mutants was associated with an

accumulation of ORF2 in the Golgi apparatus (**S1 Fig**) and a reduced association with

cellular membranes (**Fig 2C**, Membrane extract), indicating that these mutated

proteins are likely soluble in the Golgi lumen. In addition, high molecular weight forms

of ORF2intra in the soluble fraction as well as an increase of ORF2g/c secretion were

observed (**Fig 2C**), suggesting a higher translocation into the secretory pathway

associated to an improved functionality of ORF2 SP. Quantification of intracellular

RNAs showed that replication was not altered in these mutants (**Fig 2D**). In addition,

ARM mutations did not affect ORF3 expression (**Fig 2C**). However, these mutants no

longer produced infectious viral progeny (**Fig 2D**). Thus, our results suggest that the

ARM drives the ORF2 nuclear translocation and plays important functions in the HEV

lifecycle, notably in the assembly of infectious particles.

We also generated mutants for which Pro25, Ser26 and Gly27 residues were replaced

by 3 arginine residues (PSG/3R mutant), and alternatively SP was fully or partially

deleted (ΔSP and ΔSP1 mutants, respectively) (**Fig 2A**). The PSG/3R mutant showed

a marked nuclear localization (**Figs 2B** and **2C**, Nuclear extract) but was impaired in

ORF2g/c secretion (**Fig 2C,** Supernatant), as observed for the SP deletion mutants,

indicating that the addition of arginine residues strengthens the NLS function of ARM

but inhibits the functionality of ORF2 SP. The PSG/3R mutant expressed the ORF3

protein but displayed lower intracellular replication levels and was no longer infectious

(**Fig 2D**). The increased nuclear localization of this mutant is therefore likely

responsible for the reduction of HEV RNA replication and assembly of infectious

particles.

The full (ΔSP) or partial (ΔSP1) deletion of the ORF2 SP led to a total inhibition of

ORF2 secretion (**Fig 2C**, Supernatant), as expected due to the absence of reticular

translocation. Interestingly, the ORF2 protein still exhibited a nuclear localization (**Fig**

2B), indicating that the nuclear translocation process is independent of the reticular

translocation. Because ORF2 and ORF3 are overlapping, and ORF3 is essential to

particle secretion, the SP deletion mutants did not express the ORF3 protein (**Fig 2C**)

and displayed reduced extracellular titers (**Fig 2D**). Intracellular titers were also

lowered in SP mutants (**Fig 2D**), indicating that the ORF2 SP likely plays an important

role in the assembly of infectious particles.

Lastly, the highly conserved Gly31 residue was also mutated (**Fig 2A**, G/A mutant).

This mutant displayed a subcellular distribution similar to that of wt, and expressed the

ORF3 protein. Although to a lesser extent than wt, G/A mutant produced intracellular

particles, but showed reduced extracellular RNA and infectious levels. This indicates

that the G/A mutation affects particle secretion.

We next carried out a comparative study of NLS sequences in viral proteins and their

importin. We found that the ORF2 ARM is similar to the Epstein-Barr nuclear antigen

leader protein (EBNA-LP) arginine-rich NLS (RRVRRR) that interacts with Importin-α1

[17]. Interestingly, ORF2 and Importin-α1 co-localized in the nucleus of infected cells

with a Pearson correlation coefficient (PCC) of 0.670, and the mutation of arginine

residues drastically reduced this colocalization (**S2 Fig**).

Taken together, our results indicate that ORF2 colocalizes with Importin-α1 thanks to

its ARM that serves as a functional NLS. In addition, these results suggest that the

ARM is involved in the fine-tuning of the addressing and stoichiometry of the ORF2

protein between the nuclear, cytosolic and reticular pathways. This stoichiometry is

likely essential to the HEV lifecycle.

The nuclear translocation of ORF2 down-regulates the NF-κB-related signaling.

Our results demonstrated that ORF2 localizes in the nucleus (ORF2ni) and the ARM is pivotal for nuclear translocation. Our results also suggested that ORF2ni is readily detected as early as 18 h.p.e. in PLC3/HEV-p6 cells while its nuclear targeting is transient and starts to decrease after 48h (**Fig 1**). This observation prompted us to address the impact of early nuclear translocation of ORF2 on the regulation of host genes. We performed a transcriptomic analysis by microarrays (Agilent SurePrint Technology) in PLC3/HEV-p6-wt, PLC3/HEV-p6-5R/5A, PLC3/HEV-p6-ΔORF3 and PLC3 mock cells at 18 h.p.e. Interestingly, in PLC3/HEV-p6-wt and PLC3/HEV-p6-ΔORF3 cells, we observed a significant inhibition of expression of 7 genes related to the TNFα, IL-17 and NF-κB-mediated signaling as well as inflammatory responses (*i.e.,* NOD-like receptor-induced response) (**Figs 2E** and **S3**). In contrast, no gene expression inhibition was observed in PLC3/HEV-p6-5R/5A cells, reflecting the significance of ORF2 nuclear translocation in the observed inhibition. Of note, while some reports suggested that ORF3 expression modulates the host responses [18–23], no marked difference was observed when comparing ΔORF3 mutant to wt.

Altogether our results suggest that the ORF2 ARM, which notably regulates ORF2 nuclear translocation, is a pivotal viral determinant for the modulation of host pathways and, especially, genes of the NF-κB-induced signaling upon infection. Further studies will be required to define precisely the impact of this HEV-driven host regulation on immune cell responses.

Nuclear export of the ORF2 protein

The observation that ORF2 nuclear targeting is transient and decreases after 48h (**Fig 1**) then prompted us to investigate the mechanisms of ORF2 nuclear export. We

treated PLC3/HEV-p6 cells with nuclear export inhibitors, Leptomycin B (LepB) and Verdinexor (Verd). These compounds are irreversible (LepB) and reversible (Verd) inhibitors of the ubiquitous transport receptor chromosome maintenance protein 1 (CRM1/Exportin 1), which recognizes hydrophobic leucine-rich export signals [24]. Treated cells displayed a highly significant nuclear accumulation of ORF2, as compared to control cells (**Fig 3A**). In addition, co-localization studies revealed that ORF2 partially or transiently colocalizes with CRM1 in untreated cells whereas they significantly colocalize upon treatment with nuclear export inhibitors (**S4 Fig**). These results indicate that ORF2 undergoes a nuclear export to the cytoplasm by a CRM1-dependent mechanism.

CRM1 recognizes hydrophobic leucine-rich export signals. By analyzing the ORF2 sequence, we identified at least 12 potential nuclear export signals (NES) into the ORF2 sequence. We replaced the hydrophobic residues in these conserved motifs by alanine residues and characterized the generated mutants as described above. Most of mutations did not affect ORF2 partitioning, whereas three mutants named NES9, NES10 and NES12 (**Fig 3B**) led to a highly significant accumulation of ORF2 inside the nucleus (**Figs 3C** and **3D**), as observed for cells treated with nuclear export inhibitors (**Fig 3A**). NES9 and NES10 mutants were no longer infectious, whereas the NES12 mutant with the lowest nuclear accumulation (**Fig 3C**) still exhibited some intracellular and extracellular infectivity (**Fig 3E**). Of note, intracellular replication was not altered by NES mutations (**Fig 3F**), indicating that the loss of infectious particle assembly is due to the differential subcellular localization of the mutants and not to a replication defect.

These results indicate that HEV has set up a nuclear export system for its ORF2 capsid protein. This mechanism involves CRM1 which recognizes three NES on the ORF2

sequence. Moreover, these results suggest again that a fine balance between the

nuclear, cytosolic and reticular pathways is likely essential to the HEV lifecycle.

Translocation and maturation of the glycosylated ORF2 forms

Next, we investigated the mechanisms of translocation and maturation of the highly

secreted and glycosylated ORF2g/c isoforms. First, we treated PLC3/HEV-p6 and

Mock cells with Mycolactone, an inhibitor of Sec61 translocon, the membrane

embedded protein complex responsible for the translocation of newly synthetized

polypeptides into the ER lumen [25]. Interestingly, we observed a dose-dependent

reduction of ORF2g/c secretion in Mycolactone-treated PLC3/HEV-p6 cell

supernatants (**Fig 4A**), indicating that reticular translocation of the ORF2g/c forms is

Sec61-dependent.

Previously, we demonstrated that the first residues of ORF2i, ORF2g and ORF2c

proteins are Leu14, Ser34 and Ser102, respectively [12,13] (**Fig 3B**). Therefore, the

first 20 aa of the ORF2i protein are not present in the ORF2g/c isoforms. Furthermore,

the ORF2i protein is not glycosylated whereas ORF2g/c proteins are highly

glycosylated [13] (**Fig 4B**). Thanks to these features, we generated a murine

monoclonal antibody (P1H1) that recognizes the N-terminus of ORF2i (**Fig 4B**). P1H1

specifically immunoprecipitates the ORF2i protein without cross-reacting with the

highly secreted and glycosylated ORF2g/c proteins (**Fig 4C**, SN). We also generated

the P3H2 antibody that recognizes the different isoforms of ORF2. Both antibodies

recognize the intracellular ORF2 form (**Fig 4C,** Cells).

These antibodies were used to evaluate the effects of three furin inhibitors and related

proprotein convertases (PC) on ORF2g/c maturation. PC cleave the multibasic motifs

R/K-X_n-R/K (where X_n corresponds to a 0-, 2-, 4-, 6-amino-acid spacer) in the precursor

proteins[26]. The presence of ARM and RRR motif upstream of the ORF2g/c N-termini

(**Fig 3B**), respectively, suggests that a PC might be involved in the maturation of these

ORF2 forms. Therefore, we treated PLC3/HEV-p6 cells with three potent furin/PC

inhibitors (decanoyl-RVKR-chloromethylketone [CMK], hexa-D-arginine amide [D6R],

and SSM3 trifluoroacetate [SSM3]) [27] and immunoprecipitated ORF2 proteins in cell

supernatants with P1H1 and P3H2 antibodies. Intracellular contents were probed by

WB for ORF2intra, cleavage of cellular αV-pro-integrin (a substrate of intracellular

furin) and tubulin (**Figs 4D-F**). In these experiments, immunoprecipitation of ORF2g by

P1H1 antibody was used as a read-out of the inhibition of ORF2g maturation (**Fig 4B,**

ORF2g*). In treated cells, we observed a dose-dependent immunoprecipitation of

ORF2g* by P1H1 (**Figs 4D-F**), indicating that furin/PC inhibitors abrogated ORF2g

maturation. Of note, the cell-permeable CMK and SSM3 inhibitors showed a strong

inhibition of ORF2g and αV-pro-integrin maturation, whereas the cell membrane

impermeable D6R inhibitor showed a moderate effect on ORF2g maturation. Together,

these results indicate that a furin/PC present in the secretory pathway is likely involved

in the ORF2g/c maturation process.

The ORF2 ARM is the regulator of ORF2 addressing

To further analyze the molecular mechanisms by which ORF2 is differentially

addressed to the cytosolic, nuclear or reticular pathways, we next generated chimeric

and mutant constructs between ORF2 and the CD4 glycoprotein, as a reporter protein.

Therefore, we used a heterologous expression system for the further experiments. The

constructs were expressed in Huh-7 cells stably expressing the T7 RNA-polymerase

(H7-T7-IZ cells) [28]. We selected the 5R/5A and PSG/3R mutations for their marked

phenotype (**Fig 2**) and generated an ARM-deleted mutant (ΔARM, deletion of Gln24

to Arg33, **Fig 5**). The full-length ORF2wt, ORF2$^{5R/5A}$ and ORF2$^{PSG/3R}$ proteins displayed

a similar pattern and phenotype as observed in the infectious system (**Fig 2**), by

immunofluorescence and WB (**Fig 5**). The effect of the 5R/5A mutation or the ARM

deletion on the ORF2 secretion, membrane association and nuclear localization

confirmed that the ARM located downstream of the SP negatively regulates ORF2

reticular translocation but is important for nuclear translocation and membrane

association. Conversely, the PSG/3R mutation showed an increased nuclear

localization and membrane association, whereas ORF2 secretion was fully blocked,

confirming that positively charged residues negatively regulate the functionality of

ORF2 SP but mediate nuclear translocation and membrane association.

Next, to define the impact of ARM on another SP, we exchanged the ORF2 SP by the

one of CD4 that is a functional model SP [29] (**Fig 5,** Chimeras C1). Interestingly, the

chimera C1 displayed a subcellular distribution similar to that of ORF2wt but was no

longer secreted (**Fig 5C**, SN), indicating that the ARM inhibits the functionality of CD4

SP. Indeed, the mutation (C1$^{5R/5A}$) or deletion of ARM (C1$^{\Delta ARM}$) restored secretion of

the chimera C1. The observation that ORF2wt is secreted in the presence of the ARM

whereas it is not with the CD4 SP, suggests the existence of an interplay between

ORF2 SP and ARM. The chimera C1$^{PSG/3R}$ showed an increased nuclear localization

and membrane association, whereas its secretion was abolished (**Fig 5**), confirming

that positively charged residues downregulate the functionality of CD4 SP and mediate

nuclear translocation and membrane association. It should be noted that the chimera

C1$^{PSG/3R}$ showed a marked reticular staining (**Fig 5A**). As described below, we

hypothesized that the CD4 SP might disturb the maturation of ORF2i and anchors the

protein into the membrane on the cytosolic side.

We also generated an additional group of ORF2 constructs in which the SP of ORF2

was partially deleted (C2) (**S5 Fig**). The characterization of these constructs confirmed

that ARM mediates ORF2 nuclear translocation and membrane association

independently of the reticular translocation.

In order to specifically study the impact of ARM on the functionality of the ORF2 SP

independently of the ORF2 ectodomain, CD4 chimeras containing the SP (CD4^{SPORF2}),

the N-terminus (Chimeras C4) or the ARM of ORF2 (Chimeras C5) were also

generated and characterized as previously (**Fig 6**). Thanks to the CD4^{SPORF2} construct,

we confirmed that the SP of ORF2 is a functional SP, as illustrated by its subcellular

pattern (**Fig 6A**) and its efficient secretion (**Fig 6D**). Interestingly, the chimera C4

showed an intracellular distribution different from that of CD4wt, with a significant

nuclear localization (**Figs 6A** and **6B**). In addition, WB analysis revealed a major

decrease in C4 secretion as well as the appearance of a 40kDa band in the soluble

fraction (**Fig 6D**). This band corresponds to the non-N-glycosylated CD4 ectodomain,

which contains 2 N-glycosylation sites, indicating that the CD4 ectodomain is poorly

translocated into the ER lumen when fused to ORF2 N-terminus. The same

observations were made for the chimera C4$^{PSG/3R}$. In contrast, the 5R/5A mutations

restored the secretion but abolished the nuclear translocation of C4 (C4$^{5R/5A}$, **Figs 6A**,

B and **D**). Characterization of the chimeras C5, which contain only the ORF2 ARM,

showed results similar to the chimeras C4 (**Figs 6B-D)**. However, unlike C4, the

chimera C5 was no longer secreted (**Fig 6D)**, supporting the hypothesis of an interplay

between ORF2 SP and ARM. Moreover, the chimera C5 displayed a reticular staining

in addition to nuclear staining (**Fig 6C)**. This observation is in line with our hypothesis

that the CD4 SP does not undergo the same maturation as ORF2 SP, and anchors the

protein into the membrane with a cytosolic orientation.

Taken together, these results demonstrate that the ORF2 ARM on its own is capable

of regulating the functionality of ORF2 or CD4 SP, as well as the nuclear translocation

of the protein that carries it.

The ORF2 ARM regulates the topology of ORF2 SP

Positively charged residues, such as arginine residues, are known to function as

determinants of membrane protein topology, which is reflected in a statistical rule of

membrane topology, *i.e.* the positive-inside rule of membrane proteins [30]. Notably,

positive charges determine the orientation of the signal sequences and contribute to

membrane spanning of the SP H-segment translocating through the translocon [30].

Since our results suggest that the ARM regulates the SP functionality and membrane

association of ORF2, we next analyzed the topology of our different ORF2 and CD4

constructs by immunofluorescence (**Figs 7** and **8**). We used a low concentration of

digitonin that selectively permeabilizes the plasma membrane. Triton X-100-

permeabilized cells were analyzed in parallel as a control. The differential detection of

two epitopes on the ER-membrane associated Calnexin (CNX) was used as a control

of permeabilization. We observed that the ORF2wt, the chimera C1 and the PSG/3R

mutants displayed a staining in both Triton X-100 and Digitonin-permeabilized cells

whereas the ΔARM and 5R/5A constructs showed a labelling only in Triton X-100

permeabilized cells (**Fig 7**). These accessibility differences in association with the

secretion efficiencies (**Figs 5C** and **6D**), which reflect the reticular translocation,

allowed us to infer the membrane orientation of each construct as well as the SP

topology (**Figs 7** and **8**). Thus, in the presence of the ARM, the ORF2 SP likely adopts

a double topology (ORF2wt) whereas the CD4 SP is not functional (C1) and sticks the

protein to the cytosolic side of membranes. The ORF2 SP and CD4 SP are fully

functional when the ARM is deleted (ΔARM) or mutated (5R/5A) whereas they are

nonfunctional when the ARM is coupled to the PSG/3R mutations. The same

observations were done for the C4 and C5 chimeras (**Fig 8**).

Thus, our findings demonstrate that the ORF2 SP and the ARM act together to direct

the fate of ORF2 capsid protein. Thanks to the ARM, the ORF2 SP is likely able to

adopt a dual topology leading to either reticular translocation or membrane integration

to the cytosolic side.

Discussion

In the present study, we analyzed a series of mutants of the HEV ORF2 capsid protein to gain insight into how a same primary sequence can generate several ORF2 isoforms with distinctive sequences, post-translational modifications, subcellular localizations and functions in the HEV lifecycle. Several important conclusions can be drawn from our analyses. The first is that the ORF2 protein early transits through the nucleus during infection to control specific cellular functions *i.e.* antiviral responses of the infected cell. We identified the determinants of the ORF2 nuclear import and export. Notably, an ARM in the N-terminal region of ORF2 mediates nuclear import. Importantly, we showed that the mutation of this motif abolishes ORF2 nuclear translocation but also affects ORF2 addressing into membrane, cytosolic and reticular compartments, which was deleterious for the HEV lifecycle. This brings us to the second important finding, the ARM is pivotal in the fine-tuning of the partitioning and stoichiometry of the ORF2 protein between the nuclear, cytosolic and reticular pathways that are essential to the HEV lifecycle. The last significant finding in this study is the manner by which the SP and ARM cooperate to control the fate of ORF2 protein. Indeed, in addition to mediate the targeting of ORF2 to the ER membrane, the SP is likely able to adopt a reverse signal-anchor topology. This topology inversion would be driven by flanking charged residues of ARM according to the positive-inside rule [31,32] and leads to the anchoring of the ORF2i protein to the cytosolic side of membranes.

Previously, we [12] and others [15] demonstrated that HEV produces several isoforms of the ORF2 capsid protein. The ORF2i protein is the structural component of infectious particles. It is likely derived from the assembly of the ORF2intra form present in the

cytosolic compartment. The ORF2i and ORF2intra proteins are not glycosylated and

display the same sequence starting at Leu14 corresponding to the middle of the SP,

indicating that an intra-membrane protease might be involved in their maturation.

Further investigation is required to identify this intramembrane protease. In contrast,

ORF2g/c proteins are highly glycosylated and secreted, but are not associated with

infectious material. We identified the first residues of ORF2g/c as Ser34 and Ser102,

respectively [12,13]. The nature of the sequences upstream of the ORF2g/c N-termini

and our experiments using PC inhibitors, suggest that a PC, such as Furin, might be

involved in their maturation[33]. Intriguingly, the ARM is right upstream of the N-

terminus of ORF2g, indicating that this motif also serves as recognition site for ORF2g

maturation. Further experiments are needed to thoroughly investigate the processing

of ORF2 forms.

We demonstrated that early during infection, the ORF2 protein transits through the

nucleus likely to control antiviral responses of the infected cell. Although we identified

the determinants of nuclear import and export in the ORF2 sequence, further studies

are now required to identify the cellular partners of nuclear ORF2 and define precisely

the impact of ORF2 host regulation on immune cell responses. A previous study

demonstrated the significance of ORF2 N-terminal Arginine residues in the inhibition

of TBK1-mediated IRF3 phosphorylation [34] Although they did not take into account

the effect of Arginine residues on ORF2 partitioning, their results are in line with our

observations on ORF2 interference with the host innate immunity.

Based on our findings, we propose a model of ORF2 production. Firstly, when engaged

with the translocon, the ORF2 SP initially inserts head-on in an N_{exo}/C_{cyt} orientation

(**Fig 9A**). Two mechanisms can then take place. In one side, ORF2 SP inverts

orientation to N_{cyt}/C_{exo} to integrate the ER membrane as cleavable signal. The C-

terminal end of signal is exposed to the ER lumen and cleaved by signal peptidase,

liberating the ORF2 ectodomain in the ER lumen where it undergoes glycosylation and

protease maturation. This pathway leads to the production of ORF2g/c proteins that

are abundantly secreted and likely serve as immunological baits (**Fig 9B**). On the other

side, ORF2 SP does not invert, keeps a N_{exo}/C_{cyt} orientation and serves as a reverse

signal anchor. Next, the ORF2 protein anchored to the cytosolic side of membranes is

likely processed by an intramembrane protease (**Fig 9C**). This pathway leads to the

production of the ORF2intra protein that is early translocated into the nucleus to play

immunomodulatory functions and/or is then assembled into viral particles in the

cytosolic compartment. Of note, the dual topology was exclusively observed for the

ORF2 and CD4 constructs containing both ORF2 SP and ARM (ORF2wt and chimera

C4), reflecting the specific interplay between ORF2 SP and ARM and no involvement

of other sequence determinant in this process.

Due to the size constraint of their extracellular phase, viruses are under strong

pressure to minimize the size of their genome. Overlapping genes represent an

adaptive strategy developed by many viruses to condense a maximum amount of

information into short nucleotide sequences. This gene overlap strategy is used by

HEV for the ORF2 and ORF3 expression[35] and ORF4 in gt1 [36]. Here, our study

enabled the pioneering demonstration that HEV also developed a master strategy to

condense multiple information into only five amino acids of its capsid protein. This

strategy is likely also exploited by other pathogens. Thus, the ORF2 ARM is a central

regulator of the HEV lifecycle that controls (i) the ORF2 nuclear localization and hereby

controls cellular functions promoting regulation of host antiviral responses, (ii) the

functionality of ORF2 SP leading to the production of either viral particles or humoral

immune decoys, (iii) maturation of glycosylated ORF2 protein by cellular proteases,

and (iv) membrane association that is likely essential to particle assembly. Hence,

ARM controls the fate and function of ORF2 capsid protein isoforms and is likely a key

player in the hijacking of cellular processes.

Materials and Methods

Cell cultures. PLC3 [12]and Huh-7.5 [37] cells were cultured as previously described [13]. The Huh-7-derived H7-T7-IZ cells stably expressing the T7 RNA polymerase ([28]; kindly provided by Ralf Bartenschlager, University of Heidelberg, Germany) were maintained in a medium supplemented with 50 µg/ml of Zeocin. They were used for the transfection of the T7 promoter-driven pTM expression vectors.

Plasmids and transfection. The plasmid pBlueScript SK(+) carrying the DNA of the full length genome of adapted gt3 Kernow C-1 p6 strain, (GenBank accession number JQ679013, kindly provided by S.U Emerson) was used as a template[11]. Mutants of the ORF2 ARM or NES sites were generated by site directed mutagenesis. Individual mutations were introduced by sequential PCR steps, as described previously [13], using the Q5 High-Fidelity 2X Master Mix (New England Biolabs, NEB), then digestions with restriction enzymes and ligation were performed. All the mutations were verified by DNA sequencing. The primers used for the generation of ORF2 mutants are listed in **S1 Table.** The ORF3-null mutant of HEV-p6 (HEV-p6-ΔORF3) was generated as described in [38]

Capped genomic HEV RNAs were prepared with the mMESSAGE mMACHINE kit (Ambion) and delivered to PLC3 cells by electroporation using a Gene Pulser Xcell[TM] apparatus (Bio-Rad) [12].

The plasmids pTM-ORF2 (kindly provided by J. Gouttenoire, University of Lausanne, Switzerland) [14] and pTM/CD4 have been previously described [14,39]. The pTM/CD4 contains the DNA sequence coding for the secreted ectodomain of CD4 (aa 1-371). The primers used for the generation of ORF2/CD4 chimeras/mutants are listed in **S1 Table.** For some constructs, PCR amplifications were performed by multiple heat

pulses [40]. The pTM plasmids were transfected into H7-T7-IZ cells using ViaFect™

Transfection Reagent (Promega) following the manufacturer's recommendations.

Antibodies. Primary antibodies used in this study are listed in **Table 1**. Secondary

antibodies were from Jackson ImmunoResearch.

Table 1: Primary antibodies used for Western-blot and indirect

immunofluorescence experiments.

Target	Clone / Species / Isotype	Reference	Supplier	Antibody registry reference
HEV ORF2	1E6 / Mouse / IgG2b	MAB8002	Millipore	AB-827236
HEV ORF2	P1H1 / Mouse / IgG3	Patent # PCT/EP2019/068338	Home-made	x
HEV ORF2	P3H2 / Mouse / IgG3	Unpublished	Home-made	x
HEV ORF3	Rabbit	Graff *et al.* [38]	Gift of S. Emerson (NIH)	x
CD4	Rabbit	ab133616	Abcam	AB-270883
Importin α-1	Goat	ab6036	Abcam	AB-30524
CRM1	Rabbit	#46249	Cell Signaling	AB-2799298
γ-Tubulin	Mouse / IgG1	#T-5326	Sigma-Aldrich	AB-532292
CNX-N	Rabbit mAb	ab92573	Abcam	AB-10563673
CNX-C	Rabbit	ab22595	Abcam	AB-2069006
SP1	Rabbit	#PA5-29165	Thermofisher	AB-2546641
GM130	Mouse / IgG1	#610822	BD Biosciences	AB-398141
Lamin B1	Rabbit	ab16048	Abcam	AB-443298
αV-integrin	Rabbit	ab124968	Abcam	AB-11129746

485
486
Indirect immunofluorescence. Immunostainings were performed as previously

described [13]For selective permeabilization experiments, cells were fixed with 2% of

PFA, washed twice with PBS and permeabilized for 30 min at 4°C with either 0.01% of

Digitonin (Sigma) in buffer containing 20 mM HEPES pH6.9, 0.3 M sucrose, 0.1 M KCl,

2.5 mM MgCl2 and 1 mM EDTA or 0.5% Triton X-100 in PBS. Cells were next stained

with buffers containing Digitonin.

The nuclei were stained with DAPI and cell outlines with CellMask™ Green

(Invitrogen). Cells were analyzed with a LSM 880 confocal laser-scanning microscope

(Zeiss) using Plan Apochromat 63xOil/1.4N.A. and EC Plan Neofluar 40xOil/1.4N.A.

objectives. The images were processed using ImageJ and Fiji softwares.

Quantification of the ORF2 protein nuclear fluorescence. The method was adapted

from McCloy *et al*. [41]. Briefly, the ORF2 protein nuclear fluorescence was determined

using the ImageJ software. The regions of interest (ROI) were drawn around cell

nuclei. Area, integrated density and mean gray values were measured. Then,

corrected total cell fluorescence (CTCF) was calculated by the following formula: CTCF

= integrated density – (area of selected electroporated cells x mean of background

fluorescence around the cells). The exact nuclear fluorescence was = CTCF-the mean

of the integrated density of non-infected cells.

For nuclear/cytosolic fluorescence intensity ratio, cells were co-stained with

CellMask™ Green (Invitrogen) and analyzed using ImageJ software. ROI were drawn

around the whole cells and the nuclei. Area, integrated density, mean values and the

exact cell and nuclear fluorescence were measured and calculated as described

above. For each cell, the nuclear/cytosolic fluorescence intensity ratio was calculated

by the following formula: exact fluorescence intensity of nucleus / (exact fluorescence

intensity of whole cell - exact fluorescence intensity of nucleus).

Pearson's correlation coefficient (PCC) determination. Colocalization studies were

performed by calculating the PCC using the JACoP plugin of ImageJ software. For

each calculation, at least 30 cells were analyzed.

Virus production and intracellular viral particles preparation. PLC3 cells were electroporated with HEV-p6 RNAs as previously described [12]. Intracellular viral particles were prepared by osmotic shock, as previously described [13]

Infectious titer determination. Infectious titers in focus forming unit (FFU/mL) were determined as described previously [13] by using Huh7.5 cells. ORF2-positive cells were quantified using an InCell 6000 confocal analyzer (GE Healthcare) and the Columbus image analysis software (Perkin Elmer).

Chemicals and Viability assay. Leptomycin B (Cell Signaling), Verdinexor (AdooQ® Biosciences), Mycolactone A/B toxin [42]Decanoyl-RVKR-chloromethylketone [CMK] (Sigma), hexa-D-arginine amide [D6R] (Sigma) and SSM3 trifluoroacetate (Tocris) were used in this study. Dose-response curves of PLC3 cells treated with the different drugs are shown in **S6 Fig.** Cell viability was assessed using a CellTiter 96® AQueous Non-Radioactive Cell Proliferation Assay (Promega).

RNA extraction and quantification. HEV RNA and cellular gene RNA levels were quantified by RT-qPCR as described in **Supporting Information**.

Western blotting analysis. Western blotting analyses were performed as described previously [12]. Target proteins were detected with specific antibodies (**Table 1**) and corresponding peroxidase-conjugated secondary antibodies.

Immunoprecipitations. P1H1 and P3H2 antibodies were bound to magnetic Dynabeads® M-270 Epoxy beads (Thermofisher) overnight at 37°C following the manufacturer's recommendations. Beads were washed and then incubated for 1h at

RT with heat-inactivated supernatants. Beads were washed and then heated at 80°C

for 20 min in Laemmli buffer. Proteins were separated by SDS-PAGE and ORF2

proteins were detected by WB using the 1E6 MAb.

Subcellular extraction. Extractions were performed with the Subcellular Protein

Fractionation Kit for Cultured Cells (Thermo scientific) following the manufacturer's

recommendations. The cytoplasmic extracts were ultra-centrifuged at 100 000 g

during 1h at 4°C. Anti-β-tubulin (cytoplasmic), anti-Calnexin (membranous) and anti-

SP1 or anti-Lamin B1 (nuclear soluble) antibodies were used to control the quality of

extractions.

Transcriptomic analysis. Gene expression analysis was performed at 18 h.p.e. on

PLC3 cells electroporated with HEV-p6-wt, HEV-p6-5R/5A, HEV-p6-ΔORF3 RNAs or

no RNA (mock). Oligonucleotide microarrays for human whole genome (G4858A

design 072363, 8x60k chips SurePrint G3 unrestricted GE, Agilent Technologies) were

used for global gene expression analysis, as described in **Supporting Information**.

Statistical analyses. Statistical analyses were performed with the software RStudio

version 1.2.5001 combined with R version 3.6.1. For all statistical tests, reported p

values were two-sided. A test was declared statistically significant for any p value

below 0.05. For comparing more than three groups of unpaired data, ANOVA or its

non-parametric equivalent test, the Kruskal-Wallis test, was used. ANOVA was

preferred when the distributions in each group followed a normal distribution, and the

assumption of equality of the variances between each group was verified. When tests

showed a significant difference between the groups, post hoc tests were performed.

The Dunnett test (dunnettTest function with default option) followed an ANOVA, and

the Conover's test (kwManyOneConoverTest function with pvalues adjusted by the

Benjamini-Hochberg procedure) followed the Kruskal-Wallis test. Each group was

compared to a reference. For the kinetic experiment, the Friedman test was preferred

because of a lack of normality and homogeneity of variance. The post hoc Nemenyi

test (frdManyOneNemenyiTest function from the package PMCMRplus) was used and

each time point was compared to data observed at 18 h.p.e.

Acknowledgments

This work was supported by a grant from the French agency ANRS-Maladies infectieuses émergentes. This work was also supported by the Pasteur Institute of Lille, Région Hauts-de-France and Inserm-transfert. K.H., M.A. and T.V. were supported by a fellowship from the ANRS. M.F. was supported by a fellowship from the Pasteur Institute of Lille and Région Hauts-de-France. The PLBS platform used in this work was supported by the ANR (ANR-10-EQPX-04-01) and Feder (12001407 [D-AL] EquipEx ImagInEx BioMed). We thank Olivia Beseme for her technical contribution. We thank Suzanne U. Emerson (NIH, USA), Jérôme Gouttenoire (University of Lausanne) and Ralph Bartenschlager (University of Heidelberg) for providing us with reagents. We thank François-Loïc Cosset (University of Lyon) for critical reading of the manuscript.

Author Contributions

K.H., M.F., M.A., C.M., C.C., V.A., A.D., C.L., A.T.F., P.B., D.H., T.V., J-M.S., S.S-D., A.V., P.B., M.D., Y.R., J.D., C-M.A., and L.C. performed research and/or analyzed data.

L.M., M.D., Y.R. contributed to reagents or analytic tool.

L.C. wrote the paper.

Declaration of Interests

The authors declare no competing interests.

References

1. Horvatits T, Wiesch JS zur, Lütgehetmann M, Lohse AW, Pischke S. The Clinical Perspective on Hepatitis E. Viruses. 2019;11: 617–19. doi:10.3390/v11070617

2. Wang S, Cheng X, Dai X, Dong C, Xu M, Liang J, et al. Rabbit and human hepatitis E virus strains belong to a single serotype. Virus research. 2013;176: 101–106. doi:10.1016/j.virusres.2013.05.013

3. Doceul V, Bagdassarian E, Demange A, Pavio N. Zoonotic Hepatitis E Virus: Classification, Animal Reservoirs and Transmission Routes. Viruses. 2016;8: 270. doi:10.3390/v8100270

4. Feng Z, Hensley L, McKnight KL, Hu F, Madden V, Ping L, et al. A pathogenic picornavirus acquires an envelope by hijacking cellular membranes. Nature. 2013;496: 367–371. doi:10.1038/nature12029

5. Yin X, Li X, Feng Z. Role of Envelopment in the HEV Life Cycle. Viruses. 2016;8. doi:10.3390/v8080229

6. Tam AW, Smith MM, Guerra ME, Huang CC, Bradley DW, Fry KE, et al. Hepatitis E virus (HEV): molecular cloning and sequencing of the full-length viral genome. Virology. 1991;185: 120–131.

7. LeDesma R, Nimgaonkar I, Ploss A. Hepatitis E Virus Replication. Viruses. 2019;11: 719–17. doi:10.3390/v11080719

8. Nimgaonkar I, Ding Q, Schwartz RE, Ploss A. Hepatitis E virus: advances and challenges. Nat. Rev. Gastroenterol. Hepatol. 2018;15: 96–110. doi:10.1038/nrgastro.2017.150

9. Yang Y-L, Nan Y-C. Open reading frame 3 protein of hepatitis E virus: Multi-function protein with endless potential. World J Gastroentero. 2021;27: 2458–2473. doi:10.3748/wjg.v27.i20.2458

10. Wißing MH, Brüggemann Y, Steinmann E, Todt D. Virus–Host Cell Interplay during Hepatitis E Virus Infection. Trends Microbiol. 2020;29: 309–319. doi:10.1016/j.tim.2020.07.002

11. Shukla P, Nguyen HT, Faulk K, Mather K, Torian U, Engle RE, et al. Adaptation of a genotype 3 hepatitis E virus to efficient growth in cell culture depends on an inserted human gene segment acquired by recombination. J Virol. 2012;86: 5697–5707. doi:10.1128/jvi.00146-12

12. Montpellier C, Wychowski C, Sayed IM, Meunier J-C, Saliou J-M, Ankavay M, et al. Hepatitis E Virus Lifecycle and Identification of 3 Forms of the ORF2 Capsid Protein. Gastroenterology. 2018;154: 211-223.e8. doi:10.1053/j.gastro.2017.09.020

13. Ankavay M, Montpellier C, Sayed IM, Saliou J-M, Wychowski C, Saas L, et al.
New insights into the ORF2 capsid protein, a key player of the hepatitis E virus
lifecycle. Sci Rep. 2019;9: 6243. doi:10.1038/s41598-019-42737-2

14. Lenggenhager D, Gouttenoire J, Malehmir M, Bawohl M, Honcharova-Biletska H,
Kreutzer S, et al. Visualization of hepatitis E virus RNA and proteins in the human
liver. J Hepatol 2017;67: 471–479. doi:10.1016/j.jhep.2017.04.002

15. Yin X, Ying D, Lhomme S, Tang Z, Walker CM, Xia N, et al. Origin, antigenicity,
and function of a secreted form of ORF2 in hepatitis E virus infection. Proc Natl Acad
Sci USA 2018;3: 201721345–6. doi:10.1073/pnas.1721345115

16. Ba ANN, Pogoutse A, Provart N, Moses AM. NLStradamus: a simple Hidden
Markov Model for nuclear localization signal prediction. BMC Bioinformatics.
2009;10: 202. doi:10.1186/1471-2105-10-202

17. Nakada R, Matsuura Y. Crystal structure of importin-α bound to the nuclear
localization signal of Epstein-Barr virus EBNA-LP protein. Protein Science. 2017;26:
1231–1235. doi:10.1002/pro.3173

18. Dong C, Zafrullah M, Mixson-Hayden T, Dai X, Liang J, Meng J, et al.
Suppression of interferon-α signaling by hepatitis E virus. Hepatology. 2012;55:
1324–1332. doi:10.1002/hep.25530

19. Lei Q, Li L, Zhang S, Li T, Zhang X, Ding X, et al. HEV ORF3 downregulates
TLR7 to inhibit the generation of type I interferon via impairment of multiple signaling
pathways. Sci Rep. 2018;8: 1–10. doi:10.1038/s41598-018-26975-4

20. Wang M, Huang Y, He M, Peng W-J, Tian D-Y. Effects of hepatitis E virus
infection on interferon production via ISG15. World J Gastroentero. 2018;24: 2173–
2180. doi:10.3748/wjg.v24.i20.2173

21. He M, Wang M, Huang Y, Peng W, Zheng Z, Xia N, et al. The ORF3 Protein of
Genotype 1 Hepatitis E Virus Suppresses TLR3-induced NF-κB Signaling via TRADD
and RIP1. Sci Rep. 2016; 1–13. doi:10.1038/srep27597

22. Nan Y, Ma Z, Wang R, Yu Y, Kannan H, Fredericksen B, et al. Enhancement of
Interferon Induction by ORF3 Product of Hepatitis E Virus. J Virol. 2014;88: 8696–
8705. doi:10.1128/jvi.01228-14

23. Xu J, Wu F, Tian D, Wang J, Zheng Z, Xia N. Open reading frame 3 of genotype
1 hepatitis E virus inhibits nuclear factor-карра B signaling induced by tumor
necrosis factor-α in human A549 lung epithelial cells. PLoS ONE. 2014;9: e100787.
doi:10.1371/journal.pone.0100787

24. Mathew C, Ghildyal R. CRM1 Inhibitors for Antiviral Therapy. Front Microbiol
2017;8: 10995–20. doi:10.3389/fmicb.2017.01171

25. Demangel C, High S. Sec61 blockade by mycolactone: A central mechanism in
Buruli ulcer disease. Biology of the cell / under the auspices of the European Cell
Biology Organization. 2018;110: 237–248. doi:10.1111/boc.201800030

26. Seidah NG, Prat A. The biology and therapeutic targeting of the proprotein
convertases. Nat Rev Drug Discov. 2012;11: 367–383. doi:10.1038/nrd3699

27. Cheng Y-W, Chao T-L, Li C-L, Chiu M-F, Kao H-C, Wang S-H, et al. Furin
Inhibitors Block SARS-CoV-2 Spike Protein Cleavage to Suppress Virus Production
and Cytopathic Effects. Cell Reports. 2020;33: 108254.
doi:10.1016/j.celrep.2020.108254

28. Romero-Brey I, Merz A, Chiramel A, Lee J-Y, Chlanda P, Haselman U, et al.
Three-dimensional architecture and biogenesis of membrane structures associated
with hepatitis C virus replication. PLoS Pathog. 2012;8: e1003056.
doi:10.1371/journal.ppat.1003056

29. Lumangtad LA, Bell TW. The signal peptide as a new target for drug design.
Bioorganic & Medicinal Chemistry Letters. 2020;30: 127115–8.
doi:10.1016/j.bmcl.2020.127115

30. Heijne G von. Membrane-protein topology. Nat. Rev. Mol. Cell Biol. 2006;7: 909–
918. doi:10.1038/nrm2063

31. Nilsson J, Persson B, Heijne G von. Comparative analysis of amino acid
distributions in integral membrane proteins from 107 genomes. Proteins. 2005;60:
606–616. doi:10.1002/prot.20583

32. Heijne G von. Control of topology and mode of assembly of a polytopic
membrane protein by positively charged residues. Nature. 1989;341.
doi:10.1038/341456a0

33. Izaguirre G. The Proteolytic Regulation of Virus Cell Entry by Furin and Other
Proprotein Convertases. Viruses. 2019;11: 837–19. doi:10.3390/v11090837

34. Lin S, Yang Y, Nan Y, Ma Z, Yang L, Zhang Y-J. The Capsid Protein of Hepatitis
E Virus Inhibits Interferon Induction via Its N-terminal Arginine-Rich Motif. Viruses.
2019;11: 1050–17. doi:10.3390/v11111050

35. Graff J, Torian U, Nguyen H, Emerson SU. A Bicistronic Subgenomic mRNA
Encodes both the ORF2 and ORF3 Proteins of Hepatitis E Virus. J Virol. 2006;80:
5919–5926. doi:10.1128/jvi.00046-06

36. Nair VP, Anang S, Subramani C, Madhvi A, Bakshi K, Srivastava A, et al.
Endoplasmic Reticulum Stress Induced Synthesis of a Novel Viral Factor Mediates
Efficient Replication of Genotype-1 Hepatitis E Virus. Meng X-J, editor. PLoS Pathog.
2016;12: e1005521. doi:10.1371/journal.ppat.1005521.s009

37. Blight KJ, Mckeating JA, Rice CM. Highly Permissive Cell Lines for Subgenomic
and Genomic Hepatitis C Virus RNA Replication. J Virol. 2002;76: 13001–13014.
doi:10.1128/jvi.76.24.13001-13014.2002

38. Graff J, Nguyen H, Yu C, Elkins WR, Claire MS, Purcell RH, et al. The Open
Reading Frame 3 Gene of Hepatitis E Virus Contains a cis-Reactive Element and
Encodes a Protein Required for Infection of Macaques. J Virol. 2005;79: 6680–6689.
doi:10.1128/jvi.79.11.6680-6689.2005

39. Cocquerel L, Wychowski C, Minner F, Penin F, Dubuisson J. Charged Residues
in the Transmembrane Domains of Hepatitis C Virus Glycoproteins Play a Major Role
in the Processing, Subcellular Localization, and Assembly of These Envelope
Proteins. J Virol. 2000;74: 3623–3633. doi:10.1128/jvi.74.8.3623-3633.2000

40. Orpana AK, Ho TH, Stenman J. Multiple Heat Pulses during PCR Extension
Enabling Amplification of GC-Rich Sequences and Reducing Amplification Bias. Anal
Chem. 2012;84: 2081–2087. doi:10.1021/ac300040j

41. McCloy RA, Rogers S, Caldon CE, Lorca T, Castro A, Burgess A. Partial
inhibition of Cdk1 in G 2phase overrides the SAC and decouples mitotic events. Cell
Cycle. 2014;13: 1400–1412. doi:10.4161/cc.28401

42. Marsollier L, Brodin P, Jackson M, Korduláková J, Tafelmeyer P, Carbonnelle E,
et al. Impact of Mycobacterium ulcerans Biofilm on Transmissibility to Ecological
Niches and Buruli Ulcer Pathogenesis. PLoS Pathog. 2007;3: e62-13.
doi:10.1371/journal.ppat.0030062

Supporting information

Supporting figures

S1 Fig. Colocalization analysis of ORF2 with the Golgi marker 130 (GM130) in PLC3/HEV-p6 cells expressing ORF2wt or ARM/SP mutants.

S2 Fig. Colocalization analysis of ORF2 with the Importin-α1 in PLC3/HEV-p6 cells expressing ORF2wt and ARM/SP mutants.

S3 Fig. Transcriptomic analysis.

S4 Fig. Colocalization analysis of ORF2 with the exportin CRM1 in PLC3/HEV-p6 cells expressing ORF2wt or NES mutants.

S5 Fig. Addressing of C2 constructs.

S6 Fig. Dose-response curves of PLC3 cells treated with the different drugs used in this study.

Supporting Materials and Methods

RNA extraction and quantification.
Transcriptomic analysis.

Supporting tables

756 **S1 Table: Primers of use to generate ORF2/CD4 chimeras/mutants.**

757 **S2 Table: Primers used for RT-qPCR of cellular genes.**

Figure legends

Fig 1. Kinetics of the subcellular localization of ORF2 protein in PLC3/HEV-p6 cells. PLC3 cells were electroporated with HEV-p6 RNA and fixed at the indicated timepoints p.e. Indirect immunofluorescence analysis was performed using the 1E6 anti-ORF2 antibody. Cells were analyzed by confocal microscopy (magnification x40). Scale bar, 20 μm. Nuclear fluorescence intensity quantification was done using ImageJ software (mean $\pm$ S.D., $n \geq 30$ cells, Friedman with Nemenyi test). ****$p < 0.0001$. Mock electroporated PLC3 cells were used as controls.

Fig 2. ORF2 contains an Arginine-Rich Motif (ARM) that is important for its nuclear localization and host gene expression regulation. (A) Schematic sequence alignment of ORF2wt and ARM/SP mutants. (B) Subcellular localization of ORF2wt and ARM/SP mutants. PLC3 cells were electroporated with wt and mutant HEV-p6 RNAs. At 18 h.p.e, cells were processed for indirect immunofluorescence using the 1E6 anti-ORF2 antibody (Ab) and analyzed by confocal microscopy (magnification x63). Red = ORF2; Blue = DAPI. Scale bar, 20μm. Nuclear fluorescence intensity quantification was done using ImageJ software (mean $\pm$ S.D., $n \geq 30$ cells, Kruskal-Wallis with Conover's test). *$p < 0.05$, ****$p < 0.0001$. (C) Subcellular fractionation of PLC3/HEV-p6 expressing ORF2wt and ARM/SP mutants at 10 d.p.e. Fractionation was done using a subcellular protein fractionation kit for cultured cells. ORF2 proteins were detected by WB with 1E6 Ab. Glycosylated ORF2 (ORF2g), cleaved ORF2 (ORF2c), intracellular ORF2 (ORF2intra), nuclear ORF2intra (ORF2ni), nuclear and cleaved ORF2intra (ORF2nc) are indicated. ORF3 protein in cell lysates was detected with a rabbit anti-ORF3 Ab. Tubulin, ER marker Calnexin (CNX) and the

transcription factor SP1 used as a nuclear marker, were also detected to check the

quality of fractionation. Molecular mass markers are indicated on the right (kDa). (D)

Infectious titer determination and HEV RNA quantification in PLC3/HEV-p6 expressing

ORF2wt or mutant proteins. Extra- and intracellular viral particles were extracted at 10

d.p.e and used to infect naïve Huh7.5 cells for 3 days. Cells were next processed for

indirect immunofluorescence. ORF2-positive cells were counted and each positive cell

focus was considered as one FFU. Results were expressed in FFU/ml (n=4). Extra-

and intracellular viral RNAs were quantified at 10 d.p.e by RT-qPCR ($n \geq 5$) (mean $\pm$

S.D., Kruskal-Wallis with Conover's test). *p < 0.05, **p < 0.01, ****p < 0.0001. (E)

Transcriptomic analysis of PLC3 cells expressing HEV-p6 wt, mutants or mock cells

performed with microarrays (Agilent SurePrint). Left - Heatmap of gene expression in

PLC3 cells expressing HEV-p6 wt, mutants or mock cells at 18 h.p.e. Color-code

represents the log fold-change (logFC) of gene expression in the indicated

comparisons. Right – STRING representation of the gene network specifically

modulated by ORF2. Bottom – Signaling pathways preferentially induced by the

nuclear translocation of ORF2 at 18 h.p.e. Transcriptomic results stem from 4

independent electroporation experiments.

Fig 3. Nuclear export of the ORF2 protein. (A) Analysis of ORF2 nuclear export in

inhibitors treated-PLC3/HEV-p6 cells. Cells were treated at 32 h.p.e with 20nM of

Leptomycin B (LepB), 100nM of Verdinexor (Verd) or diluent (EtOH or DMSO,

respectively) for 16h. Cells were processed for indirect immunofluorescence with the

1E6 anti-ORF2 Ab and analyzed by confocal microscopy (magnification x63). Red =

ORF2; Blue = DAPI. (B) Schematic representation of HEV-p6 ORF2 protein sequence

highlighting the three studied NES motifs (*i.e.,* NES9, NES10 and NES12). (C)

Subcellular localization of ORF2 NES mutants at 48 h.p.e. Red = ORF2; Blue = DAPI. In A and C, the scale bars correspond to 20μm, and nuclear/cytosolic fluorescence intensity quantification was done using ImageJ software (mean ± S.D., $n \geq 30$ cells, Kruskal-Wallis with Conover's test). *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$. (D) Subcellular fractionation of PLC3/HEV-p6 expressing ORF2wt and NES mutants at 4 d.p.e. Fractionation was done using a subcellular protein fractionation kit for cultured cells. ORF2 proteins were detected by WB with 1E6 Ab. Glycosylated ORF2 (ORF2g), cleaved ORF2 (ORF2c), intracellular ORF2 (ORF2intra), nuclear ORF2intra (ORF2ni), nuclear and cleaved ORF2intra (ORF2nc) are indicated. ORF3 protein in cell lysates was detected with a rabbit anti-ORF3 Ab. Tubulin, ER marker Calnexin (CNX) and the transcription factor SP1 used as a nuclear marker, were also detected to check the quality of fractionation. Molecular mass markers are indicated on the right (kDa). (E) Infectious titer determination in PLC3/HEV-p6 expressing ORF2wt or NES mutants. Extra- and intracellular viral particles were extracted at 10 d.p.e and used to infect naïve Huh7.5 cells for 3 days. Cells were next processed for indirect immunofluorescence. ORF2-positive cells were counted and each positive cell focus was considered as one FFU. Results were expressed in FFU/ml. (F) HEV RNA quantification in PLC3/HEV-p6 expressing ORF2wt or NES mutants. Extra- and intracellular viral RNAs were quantified at 10 d.p.e by RT-qPCR. In E and F, n=6, mean ± S.D., Kruskal-Wallis with Conover's test, ***$p < 0.001$, ****$p < 0.0001$.

Fig 4. Translocation and maturation of the glycosylated ORF2 forms. (A) Dose-response inhibition of ORF2g/c secretion in mycolactone-treated cells. PLC3/HEV-p6 and mock cells were treated for 24h with the indicated concentrations of mycolactone (in nM) or maximal volume of the vehicle, ethanol (indicated as 0 nM). Supernatants

(SN) and lysates (Cells) were collected and ORF2 proteins were detected by WB using

the 1E6 Ab. Tubulin served as control protein loading. (B) Schematic representation of

ORF2i/g/c proteins and recognition sites of P1H1 and P3H2 antibodies used to

discriminate the different ORF2 forms. SP, signal peptide. PC, proprotein convertase.

Glycans are in black. (C) Immunoprecipitation of ORF2 proteins in SN and lysates of

PLC3/HEV-p6 cells by P1H1, P3H2 and isotype control (CTL) antibodies immobilized

on magnetic beads. ORF2 proteins were detected by WB using the 1E6 Ab. (D-F)

PLC3/HEV-p6 cells were treated for 72h with the indicated concentrations of CMK,

D6R or SSM3 (in μM) or DMSO diluent (indicated as 0 μM). Supernatants (SN) and

lysates (Cells) were collected. SN were immunoprecipitated with P1H1 and P3H2

antibodies and ORF2 proteins were detected by WB using the 1E6 Ab. ORF2intra, αV-

Integrin (IntαV) and Tubulin (Tub) were detected in cell lysates. αV-pro-integrin

(ProintαV) corresponds to the non-maturated αV-integrin. ORF2g* corresponds to the

ORF2g immunoprecipitated by the P1H1 Ab. Molecular mass markers are indicated

on the right (kDa).

Fig 5. The ORF2 ARM is the regulator of ORF2 addressing. Schematic

representation of ORF2wt and CD4wt proteins. ORF2 sequences are in blue. ARM

residues are highlighted in red. CD4 sequences are in green. (A) H7-T7-IZ cells were

transfected with pTM plasmids expressing wt, mutant or chimeric ORF2 proteins.

Twenty-four hours post-transfection, cells were fixed and processed for ORF2 staining

(in red). Nuclei are in blue. Representative confocal images are shown together with

ORF2/DAPI merge images (magnification x63). Blue dots observed in some pictures

are DAPI-stained transfected plasmids. A schematic representation of each construct

is shown on the left. Scale bar, 20μm. (B) Nuclear-to-cytoplasmic ORF2 staining ratio

in H7-T7-IZ cells expressing mutant and chimeric ORF2 proteins. Quantification was

done using ImageJ software (mean $\pm$ S.D., $n \geq 30$ cells, Kruskal-Wallis with Conover's

test). **p < 0.01, ****p < 0.0001. (C) Subcellular fractionation of H7-T7-IZ cells

expressing mutant and chimeric ORF2 proteins at 24h post-transfection. Fractionation

was done using a subcellular protein fractionation kit for cultured cells. ORF2 proteins

were detected by WB with 1E6 Ab. Tubulin, Calnexin (CNX) and Lamin B1 were also

detected to control the quality of fractionation. Molecular mass markers are indicated

on the right (kDa).

Fig 6. CD4 addressing by the ORF2 ARM. Schematic representation of ORF2wt and

CD4wt proteins. SP ORF2 residues are shown in blue. ARM residues are highlighted

in red. CD4 sequences are in green. (A) and (C), H7-T7-IZ cells were transfected with

pTM plasmids expressing CD4wt or chimeric C4 and C5 proteins. Twenty-four hours

post-transfection, cells were fixed and processed for CD4 staining (in green). Nuclei

are in blue. Representative confocal images are shown together with CD4/DAPI merge

images. Blue dots observed in some pictures are DAPI-stained transfected plasmids.

A schematic representation of each construct is shown on the left. Scale bar, 20μm.

(B) Nuclear-to-cytoplasmic CD4 staining ratio in H7-T7-IZ cells expressing CD4wt or

chimeric proteins. Quantification was done using ImageJ software (mean $\pm$ S.D., $n \geq$

30 cells, Kruskal-Wallis with Conover's test). ****p < 0.0001. (D) Subcellular

fractionation of H7-T7-IZ cells expressing CD4wt or chimeric proteins at 24h post-

transfection. Fractionation was done using the subcellular protein fractionation kit for

cultured cells. CD4 proteins were detected by WB with a rabbit polyclonal anti-CD4

Ab. Tubulin, Calnexin (CNX) and Lamin B1 were also detected to control the quality of

fractionation. Molecular mass markers are indicated on the right (kDa).

Fig 7. The ORF2 ARM regulates the topology of ORF2 SP. A schematic representation of differential permeabilization process with Triton X-100 and Digitonin is shown. Representative images of the differential detection of two epitopes on the ER-membrane associated Calnexin (CNX) used to assess the permeabilization conditions are shown. H7-T7-IZ cells were transfected with pTM plasmids expressing wt, mutant or chimeric ORF2 proteins. Twenty-four hours post-transfection, cells were fixed, permeabilized with either Triton X-100 or Digitonin, and processed for ORF2 staining (in red). Nuclei are in blue. Representative confocal merge ORF2/DAPI images are shown (magnification x63). Blue dots observed in some pictures are DAPI-stained transfected plasmids. A schematic representation of each construct is shown on the left and its predicted topology on the right. Blue and red asterisks correspond to 5R/5A and PSG/3R mutations, respectively. Scale bar, 20μm.

Fig 8. The ORF2 ARM is also able to control the topology of CD4 SP. A schematic representation of differential permeabilization process with Triton X-100 and Digitonin is shown. Representative images of the differential detection of two epitopes on the ER-membrane associated Calnexin (CNX) in the used permeabilization conditions are shown. H7-T7-IZ cells were transfected with pTM plasmids expressing CD4wt or chimeric C4 and C5 proteins. Twenty-four hours post-transfection, cells were fixed, permeabilized with either Triton X-100 or Digitonin, and processed for CD4 staining (in green). Nuclei are in blue. Representative confocal CD4/DAPI merge images are shown. Blue dots observed in some pictures are DAPI-stained transfected plasmids. A schematic representation of each construct is shown on the left and its predicted

topology on the right. Blue and red asterisks correspond to 5R/5A and PSG/3R

mutations, respectively. Scale bar, 20μm.

Fig 9. Model of ORF2 addressing regulation by ARM. (A) The signal recognition

particle (SRP) recognizes the hydrophobic signal peptide (SP) of the ORF2 nascent

chain as it emerges from a translating ribosome. The ribosome-nascent chain-SRP

complex is targeted to the membrane and interacts with the SRP receptor, resulting in

the release of the SP and docking of the ribosome–nascent chain complex to the

Sec61 translocon. The ORF2 SP initially inserts head-on in an N$_{exo}$/C$_{cyt}$ orientation,

then inverts its orientation to N$_{cyt}$/C$_{exo}$. (B) The C-terminal end of SP is exposed to ER

lumen and is cleaved by signal peptidase, generating a new N-terminus. Translation

then resumes, and the nascent ORF2 protein is translocated into the ER lumen where

it is glycosylated and likely undergoes maturation by a proprotein convertase. This

pathway generates the ORF2g/c forms. (C) For a fraction of ORF2 nascent polypeptide

chains, the ARM leads the ORF2 SP to retain its N$_{exo}$/C$_{cyt}$ orientation and integrates as

reverse signal-anchor, according to the positive-inside rule. The ORF2 protein

anchored to the cytosolic side of membrane is likely processed by an intramembrane

protease to generate the ORF2i/ORF2 intra protein that is translocated into the nucleus

and assembles into viral particles.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9