Title: Crossing design shapes patterns of genetic variation in synthetic recombinant populations

of *Saccharomyces cerevisiae*

Authors: Mark A. Phillips[1*], Ian C. Kutch[1], and Molly K. Burke[1*]

Affiliations:

[1] Department of Integrative Biology,

Oregon State University

Corvallis, OR 97331

*Corresponding authors: Mark A. Phillips (philmark@oregonstate.edu), Molly K. Burke

(molly.burke@oregonstate.edu)

Abstract

Multiparent or "synthetic recombinant" populations – those created by combining distinct isogenic founders to establish a single recombinant background – have emerged as a useful tool for dissecting the genetics of complex traits. Synthetic recombinant populations can be used to derive inbred lines in which quantitative traits can be mapped, or the recombinant populations themselves can be sampled for experimental evolution. Especially for the latter application, investigators generally value maximizing genetic variation in a recombinant population; in other words, a population harboring relatively equal contributions of the genetic backgrounds of each isogenic founder strain is a desirable resource. It is well-documented that in evolution experiments initiated from recombinant or outbred ancestral populations, the subsequent adaptation that occurs in evolved populations is driven by standing genetic variation, rather than *de novo* mutations. Despite the demonstrated importance of initial genetic variation to the adaptive process, little has been done to systematically evaluate methods of constructing a synthetic recombinant population, for creating resources for evolution experiments. Here we seek to address this issue by comparing patterns of genetic variation in different synthetic recombinant populations of *Saccharomyces cerevisiae* created using one of two combination strategies: pairwise crossing of isogenic strains or simple mixing of strains in equal proportion. We also explore the impact of the varying the number of parental strains used in each strategy. We find that more genetic variation is initially present and subsequently maintained over generations when population construction includes a round of pairwise crossing. We also observe that when using a given crossing strategy, increasing the number of parental strains typically increases genetic diversity. In summary, we suggest that when creating recombinant populations for use in experimental evolution studies, simply mixing founder strains in equal proportion may limit the adaptive potential of that population.

Introduction

Despite major advances in DNA sequencing technologies, deciphering the genetic basis

of complex traits remains a major challenge in modern biology. Efforts to fully characterize the

genetic variants underlying traits like human height have had limited success (Manolio et al.

2009), and there is no clear consensus regarding what models best describe the genetic

architecture of complex traits as a whole (Boyle et al. 2017). Alongside traditional QTL mapping

and genome-wide association studies, the "evolve-and-resequence" (E&R) approach has

emerged as powerful technique that can generate insight into the genetic basis of complex

phenotypes, at least in model organisms (Schlötterer et al. 2014; Long et al. 2015; Bailey and

Bataillon 2016). This approach involves subjecting experimental populations to conditions that

target a specific trait, and simply monitoring the genotypic and phenotypic changes that occur

over generations. As these experiments typically feature replicate populations evolving in

parallel, researchers can use the data they generate to make powerful statistical associations

between genomic variants and traits of interest.

In order to most appropriately model the evolution of complex traits in humans, E&R

studies should employ sexually-reproducing eukaryotic organisms (Burke 2012, Long et al.

2015). However, despite the promise of the E&R approach for dissecting complex traits, there

are still unanswered questions regarding what constitutes optimal experimental design. To that

end, simulations have assessed the effects of particular experimental parameters on the outcomes

of E&R experiments designed to model the evolution of complex traits (Baldwin-Brown et al.

2014; Kofler and Schlötterer 2014; Vlachos and Kofler 2019). These simulations support

intuitive practices such as maximizing replication and experimental duration, and they

recommend maximizing levels of starting genetic variation in experimental populations (i.e. by

including more founder haplotypes). Our work here aims to determine how to best achieve this

recommendation in experiments with outcrossing *Saccharomyces cerevisiae*.

Most E&R experiments with sexually-reproducing model organisms use *Drosophila* (e.g.

Burke et al. 2010; Turner et al. 2011; Orozco-ter Wengel et al. 2012; Tobler et al. 2014; Huang

et al. 2014; Jha et al. 2015; Franssen et al. 2015; Graves et al. 2017; Phillips et al. 2018; Barghi

et al. 2019). However, outcrossing *S. cerevisiae* populations have emerged as an attractive

alternative due to shorter generation times, ease of replication, the ability to freeze and

"resurrect" samples taken from different timepoints, and comparable genetic resources (e.g.

Burke et al. 2014; Phillips et al. 2020). As such, we have chosen to focus our efforts on

determining how to best construct populations of outcrossing *S. cerevisiae* for use in E&R

studies with respect to maximizing standing genetic variation and haplotype diversity.

As *S. cerevisiae* populations isolated from natural or industrial settings typically lack

genetic heterogeneity, generating laboratory populations with standing genetic variation requires

crossing distinct isogenic strains. These "synthetic recombinant" or multiparent populations have

traditionally been used for QTL mapping (e.g. Cubillos et al. 2013; Linder et al. 2020), but they

can also serve as the ancestral population in evolution experiments (e.g. Burke et al. 2014;

Phillips et al. 2020). Notably, synthetic recombinant populations have also been used for E&R in

other model organisms such as *D. melanogaster* (e.g the *Drosophila* Synthetic Population

Resource *cf.* King et al. 2012) and *Caenorhabditis elegans* (e.g. the *C. elegans* Multiparent

Experimental Evolution panel, *cf.* Teotónio et al. 2012). In this context, having clearly defined

founder genotypes enables the estimation of haplotype frequencies in evolved populations from

"Pool-SEQ" data (Michalak et al. 2018, Linder et al. 2020, Phillips et al. 2020). And this ability

to detect haplotypes responding to selection, as opposed to individual markers, allows for better

characterization of linkage and genetic hitchhiking in evolved populations (Barghi and

Schlötterer 2019) than is possible in studies where founder genotypes are unknown (e.g. Graves

et al. 2017). All of this said, to our knowledge little has been done to explicitly assess how to

best generate a recombinant population for downstream use in an evolution experiment. While

simulations predict that maximizing the number of founding haplotypes and levels of genetic

variation will improve experimental outcomes, there are practical and biological limits to the

number of haplotypes that can be combined into a single recombinant background and

effectively maintained. One of our goals here is to evaluate the costs and benefits of increasing

the number of founding haplotypes in a synthetic recombinant population for E&R.

Another of our goals is to evaluate how different methods of crossing founder strains

impact levels of genetic variation in synthetic recombinant populations. There are two general

approaches to this task across model organisms. In the more common approach, individuals

sampled from various lines are simply mixed in equal proportions and allowed to mate freely

prior to the start of the experiment (e.g. Orozco-ter Wengel et al. 2012). With this strategy, the

genetic makeup of the resulting population will depend on mating efficiency between individuals

from different founding strains, reproductive output, and chance. However, work with

Drosophila has shown there is no significant allele frequency differentiation between independently constructed synthetic populations using the same sets of isofemale lines (Nouhaud et al. 2016). Such a high level of reproducibility could indicate that mating efficiency and differences in reproductive output among founding lines are neglible, or it could indicate that these differences exist and shape the genetic makeup of recombinant populations in a parallel manner. So, while this approach is relatively simple and practical to implement, it may be difficult to maximize the amount of genetic variation and founding haplotype representation in synthetic populations when there are drastic differences in reproductive output and/or mating efficiencies among founding strains. In other words, if there is substantial variation in mating efficiencies among founding strains, this could lead to the over-representation of certain haplotypes belonging to strains that mate most efficiently. By contrast, the second approach for constructing synthetic populations is more complex and involves some level of pairwise crossing between founders (Teotonio et al. 2012; King et al. 2012; Cubillos et al. 2013; Harbison et al 2017; Linder et al 2020). While this approach is significantly more labor-intensive and time-consuming, it perhaps has advantages in terms of producing populations that have more equal founder haplotype representation and, consequently, higher levels of genetic variation.

Here we assess which of these methods produces populations most well-suited for E&R studies in outcrossing *S. cerevisiae*. We have constructed sets of synthetic populations using both approaches and the same founder populations. In addition to crossing approach, we also evaluate how the number of founder populations used impacts the genetic makeup of the resulting population. Specifically, we created sets of populations using 4, 8 and 12 isogenic strains and both construction methods. Our objective is to characterize patterns of genetic variation and haplotype diversity across these populations to provide recommendations on how to best construct synthetic recombinant populations in outcrossing yeast for the specific application of using them in an E&R study.

Materials and Methods

Population creation

All yeast strains used in this study originated from heterothallic, haploid, barcoded derivatives of the SGRP yeast strain collection (Cubillos et al. 2009). A subset of 12 of these haploid strains, originally isolated from distinct geographic locations worldwide, were used to create the

synthetic populations we describe here (See Supplementary Figure 1 for phylogeny). These 12

strains were all genetically modified as described in detail by Linder et al. (2020) to enable easy

crossing and diploid recovery; these modified strains were kindly provided by Anthony D. Long

(UC Irvine) in 2017. Briefly, these strains were modified so that *MAT*a and *MAT*α strains both

contain *ho* deletions to prevent mating-type switching, but each contain a different drug-

resistance marker in a pseudogene (YCR043C) tightly linked to the mating type locus (*MAT*a,

*ho*Δ, ura3::KANMX-barcode, *ycr043C*::NatMX and *MAT*α, *ho*Δ, ura3::KANMX-barcode,

ycr043C::HygMX). These genotypes enable haploids of each mating type to be recovered using

media supplemented with either hygromycin B or nourseothricin sulfate, and they enable newly

mated a/α diploids to be recovered in media supplemented with both drugs.

Two different crossing strategies were used to create genetically diverse populations

using 4, 8, and 12 strains as founders (Table 1). "K-type" populations (named for what we call

the "kitchen sink method", or the practice of pooling isogenic strains together without careful

focus on representation) were created by simply pooling equal volumes of saturated overnight

cultures of the respective haploid founders and allowing those cells to mate. To accomplish this,

single colonies of each haploid founder strain were sampled and grown overnight (at 30ºC / 200

rpm) in 1 mL of rich media consisting of 1% yeast extract, 2% peptone, and 2% dextrose (YPD).

After ~24 hours, cultures were washed in fresh YPD media, pooled with the relevant other

overnight cultures in a 50 mL conical tube, vortexed, and now-mixed cultures were allowed to

settle and mate for 90 minutes at room temperature. These cultures were then transferred in

200uL aliquots to agar plates containing 100mg/mL nourseothricin sulfate ("NTC"), 300mg/mL

hygromycin B ("hyg") as well as 200mg/mL G418; this strategy ensured that that only newly

mated diploids would grow. The resulting lawns of mated diploid cells were collected by

scraping with a sterile glass slide into a fresh YPD media. This "cell bank" was archived as a

frozen stock at -80ºC for each of the K-type populations made with 4, 8, and 12 haploid founders

(4K, 8K, and 12K respectively).

"S-type" populations (named S due to the manipulation of spores to achieve better

representation of founder genotypes) were built to mimic more careful crossing designs in which

founding lines are crossed in pairs and/or a round-robin. To accomplish this, each haploid strain

was paired with a different strain of the opposite mating type and mated as described above.

Successful diploid colonies were isolated, grown overnight in 1 mL of YPD, washed and

resuspended in sporulation media (1mL 1% potassium acetate), then cultured for 72 hours at

30ºC / 200 rpm. Tetrads from these diploid cells were dissected using the Spore Play dissecting

microscope (Singer). The four meiotic products (spores) were then collected, allowed to grow for

2 days, and replica plated to plates containing either NTC or hyg to verify the proper segregation

of drug resistance markers and thus mating types. Once validated, the meiotic products were

grown overnight in 1 mL of YPD. Overnight cultures were standardized to the same optical

density (OD_{600}) before being pooled in equal volumes in a 50 mL conical tube. Populations were

given 90 minutes to mate at room temperature, then were plated on agar plates supplemented

with both NTC/hyg/G418 so that only newly mated diploid cells could grow. The resulting lawns

of mated diploid cells were collected by scraping with a sterile glass slide into fresh YPD media.

This "cell bank" was archived as a frozen stock at -80ºC for each of the S-type populations made

with 4, 8 and 12 founders (S4, S8, and S12 respectively; see Supplementary Figure 2 for crossing

schematics).

Population maintenance and 12 cycles of outcrossing

After the creation of the 3 "K-type" and 3 "S-type" synthetic recombinant populations described

above, all populations were taken through 12 consecutive cycles of intentional outcrossing; in

other words, the populations were subjected in parallel to a series of steps that induced regular

sporulation, spore isolation, and mating. Detailed methods are described by Burke et al. (2020).

Briefly, newly mated diploid cells from the last step of the population creation protocol (i.e. the

"cell banks") were grown overnight in 10 mL YPD media. These cultures were washed and

resuspended in sporulation media, and incubated with shaking for 72 hours (30°C/200 rpm).

Cells then underwent a number of methods to disrupt asci and isolate/randomize spores,

including incubation with Y-PER yeast protein extraction reagent (Thermo) to kill vegetative

diploids, digestion with 1% zymolyase (Zymo Research) to weaken ascus walls, and well as

high-speed shaking with 0.5 mm silica beads (BioSpec) to mechanically agitate the asci. After

these steps spores were resuspended in 10 mL YPD and allowed to settle and mate for 90

minutes at room temperature. Diploids were recovered as described above; cultures were

transferred to 10 individual YPD agar plates supplemented with NTC/hyg/G418 in 200 μL

aliquots and incubated at 30°C for 48 hours. The resulting lawns of mated diploid cells were

collected by scraping with a sterile glass slide into fresh YPD media. This "cell bank" was again

sampled for archiving at −80°C, and used to initiate an overnight culture for the next outcrossing

cycle. We estimate that 15–20 asexual generations occurred between every outcrossing cycle of

the experiment. Based on counting colonies from dilutions of cultures plated at various

benchmarks during the protocol, we expect that 7.5–11 generations elapse during the overnight

culture in YPD media, and another 7.5-11 generations elapse during the period of diploid

recovery on agar plates. Thus, a minimum of 15*12 = 180 cell doublings likely took place over

the 12 cycles of outcrossing in each synthetic recombinant population.

Genome sequencing and SNP identification

Each of the founding SGRP strains were plated as haploids on plates containing either

NTC or hygromycin to verify the presence of the appropriate drug resistance markers. Individual

colonies were isolated from each strain for verification of identifying barcodes at the URA3

locus using Sanger sequencing (Cubillos et al. 2009 provide barcode and primer sequences).

Once validated, single colonies were again isolated for whole-genome sequencing. One milliliter

of YPD media was inoculated with single colonies, grown overnight, and the resulting culture

was harvested for gDNA extracted using the Qiagen Puregene Yeast/Bact. Kit. Purified gDNA

from each haploid founder was then prepared for sequencing using the Nextera DNA Sample

Preparation Kit (Illumina). Some minor modifications to the manufacturer's protocol were

implemented to optimize throughput (*cf*. Baym et al. 2015). Genomic DNA libraries were

prepared for experimental recombinant populations in the same way and all samples were pooled

to generate a single multiplexed library. Because the recombinant (i.e. genetically variable)

populations require significantly higher coverage to accurately estimate allele frequencies at

variable sites, these populations were added to the library at 10X the molarity of each haploid

founder sample.   The multiplexed library was run on two SE150 lanes on the HiSeq3000 at the

OSU Center for Genomic Research and Biocomputing (CGRB). Data for the 4S populations

were previously published in Burke et al. (2020) and raw fastq files are available through NCBI

SRA (BioProject ID: PRJNA678990). Raw fastqs for all other populations are also available

through NCBI SRA (BioPorject ID: PRJNA732717).

We have developed a processing pipeline for estimating allele frequencies in each

population directly from our pooled sequence data. We used GATK v4.0 (McKenna et al. 2010;

Poplin et al. 2018) to align raw data to the *S. cerevisiae* S288C reference genome (R64-2-1) and

create a single VCF file for all variants identified across all replicate populations. We also

downloaded and indexed a reference VCF file with SNP information for a number of distinct

natural isolates of S. cerevisiae (Bergström et al. 2014); this is a recommended best practice for

calibrating base quality with GATK v4.0. This VCF file was converted into a SNP frequency

table with a custom python script (available upon request) suitable for downstream analysis with

R (www.R-project.org). In terms of our final SNP calling, we essentially portioned the data to

create three separate SNP tables with each table corresponding to a set of founders and

populations derived from them (e.g. a table containing with the S4 and K4 populations and their

founders). In each table, we chose to only include sites with a minimum coverage >20X in the in

synthetic populations as a quality control measure. Next, sites were filtered based on data from

the founder populations. We excluded all sites that appeared to be polymorphic within a given

founder, and sites where a single nucleotide was fixed across all founders. This was done as such

occurrences could indicate sequencing error given that our founder strains are haploid and

isogenic, and a site is unlikely to be polymorphic in our synthetic populations if it is fixed across

all of the founders.

SNP Variation

Our main objective was to evaluate how crossing strategy and the number of founder

strains impacts patterns of SNP variation in synthetic recombinant populations. To that end, we

assessed SNP-level variation in our recombinant populations using several metrics. First, we

simply determined the number of polymorphic sites segregating in each population immediately

following their creation (cycle 0), and monitored how that number changed over time (i.e. at

subsequent cycles 6 and 12). This approach of tracking the total number of polygenic SNPs

should reveal whether particular crossing strategies – i.e. using a certain number of founders,

and/or one of the two crossing strategies – consistently produced populations with more SNPs,

and whether these SNPs were maintained or lost over 12 outcrossing cycles. We also generated

UpSet plots using the UpsetR package in R (Conway et al. 2017) to visualize patterns of overlap

between the total number of SNPs possible for a given combination of founder strains, and the

SNPs we observed in our actual populations. Here we define the total number of possible SNPs

as all loci for which at least one of the founding strains used has an allele different from the

others; this number will therefore differ among the 4-way, 8-way, and 12-way crosses.

In addition to SNP number, we also characterized the distribution of SNP frequencies in

each population, which allows more direct comparisons between populations with different

numbers of founders but the same crossing strategy, or the same number of founders but

different crossing strategy. To do this, we focused on two metrics: the site frequency spectrum

(SFS), and genome-wide heterozygosity. Here heterozygosity refers to *2pq*, the product of the

reference (i.e. the S288C allele) and alternate allele frequency at a given site multiplied by 2.

It is worth noting that our ability to assess levels of genetic variation across our synthetic

populations is limited by the fact we have only collected Pool-SEQ data. Given the complex life-

history of the yeast populations in this experiment, which involves periods of 7-15 generations of

asexual growth punctuated by discrete outcrossing events, it is not possible for the genotypes of

all individuals in the population to be shuffled by recombination every generation. Therefore,

asexual lineages will evolve by clonal interference for relatively short periods of time, until the

next outcrossing event decouples individual adaptive alleles from a particular genetic

background. It is possible that during these periods of clonal interference, particular diploid

lineages will dominate, and if these lineages are heterozygous at a given locus, that will lead to

an artificially elevated heterozygosity value at that SNP. But we do not believe this is a major

complication in the understanding of nucleotide diversity in our experiment for several reasons;

namely, that our outcrossing protocol includes several measures that maximize outcrossing

efficiency (i.e. any asexual diploids that fail to sporulate are killed), the generally high rate of

recombination in yeast, and that the periods of asexual growth are short and unlikely to exceed

~20 cell doublings.

SNP frequency changes over 12 cycles of outcrossing

Although statistical power in this power is limited due to a lack of replication, we

attempted to identify regions of the genome showing obvious responses to selection in each

synthetic population. Specifically, we used Pearson's χ^2 test as implemented in the poolSeq

package in R (Taus et al. 2017) to compare SNP frequencies between cycle 0 and cycle 12 in

each synthetic population. We chose this particular test based on a benchmarking effort that

suggests it is well-suited to detecting selection in E&R experiments lacking replication (Vlachos

et al. 2019). After results were generated for each synthetic population, log transformed p-values

were plotted across using the GenWin package in R (Beissinger et al. 2015) using the following

parameters: "smoothness = 2000, method = 3." GenWin itself uses a smoothing spline technique

to define windows based on breakpoints in the data. Plots were then examined to see if there

were any genomic regions showing signs of selection based on significance levels relative to the

background.

Haplotype representation

In addition to describing SNP diversity, we also describe the diversity of founder

haplotypes represented in our synthetic populations. We were particularly interested in

evaluating whether the S-type strategy might produce populations in which founder haplotypes

are more evenly represented (at intermediate frequency) compared to the K-strategy. Given the

stochasticity inherent in the K-type strategy, we thought it probable that founder genotypes with

especially high sporulation and/or mating efficiencies (i.e. those with the highest reproductive

outputs in the outcrossing context) might come to dominate. To this end, we estimated haplotype

frequencies in all experimental populations after 12 cycles of outcrossing to determine how

evenly haplotypes were represented. We used the sliding-window haplotype caller described in

Linder et al. (2020) and software the authors have made available as a community resource:

https://github.com/tdlong/yeast_SNP-HAP. Our results were generated by using the

haplotyper.limSolve.code.R script and estimates were made across 30KB windows with a 1KB

stepsize. This particular haplotype caller was developed specifically to estimate haplotype

frequencies in multiparent populations when founder haplotypes are known. A full description of

the algorithm being used, and results of empirical validation can be found in Linder et al. (2020).

Results

SNP Variation

To assess how crossing strategy and number of founder strains impacts SNP variation, we

began by simply counting the number of SNPs present in each of our synthetic populations upon

their creation and how that changes over several cycles of recombination (Table 2). As expected,

the total number of possible SNPs that can possibly contribute to segregating genetic variation

increases with the number of founders used. Looking at our actual populations at cycle 0 and

focusing on those created using the same crossing strategy, we also generally find the observed

number of SNPs in each population to increase with the number of founders used. The only

exception to this pattern is the K12 population where we see dramatic losses in polymorphic sites

relative to all other populations. We also typically observe reductions in the number of SNPs in

all experimental populations over time. However, we do note higher "stability" (i.e. smaller

losses) in the 8-founder populations, and in population S8, we actually observe higher SNP

counts in cycle 12 than in cycle 6. This discrepancy is most likely due to a relatively small

number of sites at very low frequency in cycle 6 (i.e. too low for our SNP calling to pick up),

increasing to detectable levels by cycle 12. Nevertheless, the overall trend still appears to be

reductions in the number of polymorphic sites over time. Our data also suggest these reductions

are typically more pronounced in populations created using the K-type strategy, and that

populations created using the S-type strategy have more polymorphic sites than those created

with the K-type strategy.

We also examined patterns of overlap between our synthetic populations and what is

possible given their respective founders using UpSet plots (Figure 1). UpSet plots are useful

because they allow visualization of observations unique to groups; in other words, they highlight

observations that are included in a specific group and excluded from all others. As an illustrative

example, the fourth vertical bar in Figures 1A-C represents all of the SNPs that could possibly

segregate in a population but do not, and this reveals that a greater proportion of possible SNPs

are lost in synthetic populations with 12 founders compared to those with 4 or 8 (this result is

also noted in Table 2). We were interested in using this visualization approach to evaluate

whether there might be particular SNP groups that are hallmarks of the S-type or K-type strategy.

While many more of the possible SNPs appear in S-type populations relative to the K-type

populations for a given set of founders, the UpSet plots indicate that very few SNPS are unique

to a particular crossing strategy. We interpret this result as evidence that neither crossing strategy

favors specific alleles, and this is true regardless of the number of founder haplotypes used.

Next, we looked at how the SFS varied across different populations and how they

changed over time. As shown in Figure 2, the S-type populations tend to exhibit more even SNP

frequency distributions than the K-type populations, and they are also more stable over time.

This is particularly evident in the 12-founder populations. In these we see that by the final cycle

of recombination in the K12 populations, there is an extreme skew in the SFS with most sites

exhibiting very high or low SNP frequencies. This is also consistent with the results shown in

Table 2 where we see much higher levels of fixation over time in K12 than in any other

population. This contrast is present but far less extreme in populations created with 4 or 8

founder strains. In general, at cycle 0 all populations deviate substantially from the SFS that we

would expect if the respective founders combined in perfectly equal proportions (Supplementary

Figure 3). These deviations suggest that drift and/or selection are impacting the genetic makeup

of synthetic populations from the moment they are established.

Finally, we assessed the effects of crossing strategy and number of founders on genome-

wide heterozygosity (Table 3). The clearest pattern we observed is that after 12 cycles of

recombination, S-type populations exhibited greater overall levels of heterozygosity across the

genome compared to their K-type counterparts. Other patterns in the data are less clear, however.

Looking at the S-type populations alone, we find that by cycle 12 the S4 populations have

experienced a greater loss of heterozygosity than the S8 and S12 populations. However,

differences between S8 and S12 populations are far less severe with the former having slightly

higher mean heterozygosity. As such, there is no clear positive relationship between

heterozygosity and number of founders. This pattern largely breaks down in the K populations.

The K4 population experiences a greater loss of heterozygosity than K8 by cycle 12. However,

the K12 population experiences a near total loss of heterozygosity by cycle 12 as expected given

the other measures of SNP variation we have looked at thus far.

SNP frequency changes over 12 cycles of outcrossing

We used Pearson's χ^2 test to compare SNP frequencies between cycle 0 and cycle 12 of

each individual population to see if there were any regions of the genome showing obvious

responses to selection imposed by our outcrossing maintenance protocol. Looking across results

for each population, we do not find any genomic regions that show consistent responses to

selection (Figure 3). However, we do find instances in individual populations where there are

clear peaks in significance relative to the rest of the genome (e.g. Figure 3A and C-E).

Widespread fixation in K4 and K12 make it difficult to identify such peaks (Figure 3B and F).

We also compared our results to prior work which identified regions of the genome and

genes associated with selection for frequent outcrossing (Cubillos et al. 2013; Burke et al. 2014).

Our rationale for doing this is that as our current study features no within-treatment replication, it

is weak to implicate signatures of adaptation by itself; but, obvious overlap with other studies

would provide indirect evidence for adaptation. Similarly, if a particular population (or

population type) implicated more candidate regions from the literature than the others, this could

provide evidence that a particular crossing strategy might lead to the best outcomes in an E&R

study, with respect to identifying regions of interest. We do find instances where peaks in

significance align with previously identified candidate regions (e.g. Figure 3A, C-D), but there is

no clear pattern where peaks consistently line up with a given candidate region across multiple

populations. There are also many more instances where we do not find peaks that align with the

previously identified candidate regions. So generally speaking, while this comparative approach

provides opportunities for describing particular regions, perhaps those of *a priori* interest, we

cannot conclude that observed changes in SNP frequencies in any population are likely

signatures of selection for forced outcrossing. On the other hand, we also observe that the 8-

founder populations generally implicate more candidate genes from the literature, compared to

any other population (Figure 3C,D). This suggests that perhaps using 8 founding haplotypes in a

population for an E&R project leads to better outcomes, in terms of identifying candidate

regions, than using 4 or 12. Notably, the 8K population and 8S population implicated similar

numbers of peaks, so it is not clear that one crossing strategy is better than the other in this

respect.

Haplotype Representation

Using sequence data from cycle 12 and our founder strains, we estimated haplotype

frequencies across the genome in all populations to assess how crossing strategy and number of

founders impact haplotype representation (Note: estimates for K12 are made using far fewer

SNPs due to the extreme levels of fixation seen in this population). Due to recombination, we

find that individual haplotype representation fluctuates across the genome as haplotype blocks

are broken up (Figures 4-5 for 4-founder populations, and Supplementary Figures 4-7 for 8- and

12-founder populations). However, when looking at mean genome-wide haplotype frequency

estimates, we find that the S-type populations have haplotype frequencies that are more evenly

distributed compared to the K-type populations (Table 4). Population 4S represents the only

scenario where we find close to equal representation of all founding haplotypes across the

genome (Figure 4). By contrast, in population K4 we observe that two of the founder haplotypes

dominate (Figure 5). The observation of more evenly distributed haplotypes in S-type

populations largely holds when more founders are used (Table 4 and Supplementary Figures 4-

7). However, increasing the number of founders in and of itself appears to be associated with less

balanced haplotype representation.

Next, we evaluated whether any of our observed patterns of haplotype representation

could be explained by selection. To do this we compared our findings to the results of Burke et

al. (2014). In this study, twelve replicate populations were created using the same four founding

strains as the S4 and K4 populations, and were subjected to a similar outcrossing regime for 18

cycles (Note: the synthetic population the replicates were derived from was created using a S-

type strategy and underwent 12 cycles of recombination before replicates were generated).

Based on their analysis of genomic data taken from each population across three timepoints,

Burke et al. (2014) identified five major candidate regions associated with adaptation to forced

outcrossing. Four of these regions (regions A, B, C, and E in the paper) were also found to be

clearly defined by a given haplotype at the end of the experiment (See Supplementary Figure S4

in Burke et al. 2014). As the founders used to create the synthetic populations of Burke et al.

(2014) are present in all of the populations featured in this study, we can reasonably interpret

changes in haplotype frequency in these four candidate regions in our populations as evidence of

selection. In other words, as our populations here are unreplicated, and we cannot use strong-

inference approaches with them directly, we sought parallels between haplotype frequency

differences observed here and those described as adaptive in a very similar experiment.

As shown in Table 5, we find some evidence that the haplotypes driving adaptation in the

Burke et al. (2014) study also show substantial increases in frequency in our populations. In

candidate regions A (C9: 950,000-975,000) and E (C13:445,000-460,000), we find clear

evidence of the Y12 haplotype frequency increasing across all of our populations as it did in the

Burke et al. (2014) study. Given how dramatic some of these changes are from cycle 0 to cycle

12, we think it reasonable to speculate that these haplotypes might also be driving adaptation

here. However, this pattern in not corroborated for candidate regions B (C9: 65,000-80,000) and

C (C11:615,000-620,000). Here the haplotypes driving adaptation in the Burke et al. (2014)

study actually decrease in frequency in most instances.

Discussion

Careful crossing of founder strains increases SNP-level variation

Here we primarily sought to assess how crossing strategy - pairwise crossing of founder strains versus mixing populations in equal proportion - impacted levels of SNP and haplotype variation. We considered both the total number of polymorphic sites and levels of heterozygosity at those sites in synthetic populations created using these two approaches. We consistently found that populations generated by with a round of careful crossing that account for differences in mating efficiencies have more polymorphic sites (Table 2) and less variation is lost at those sites over time when compared to populations created by simple mixing founders in equal proportion (Table 3). We also found that increasing the number of founder strains used to create a given synthetic population also generally resulted in higher levels of SNP variation. However, the total percentage of potential polymorphic sites lost was typically higher when more populations were used (Table 2 and Figures 1). And combining the K-type crossing strategy with a large number of founders resulted in dramatic skews in the site frequency spectrum and losses in SNP variation (Figure 2). Lastly, we find no evidence that specific sets of variants are favored in one crossing strategy versus another when the same founders are used (Figure 1). Instead, we suggest that differences in outcomes between crossing strategies are most likely due to strain-specific differences in mating efficiencies and reproductive output (i.e. when populations are simply mixed together, the most compatible strains dominate).

Overall, these findings have led us to two general recommendations. First, if one endeavors to produce populations with the highest possible levels of total SNP variation, many founders and a crossing strategy that involves at least one round of pairwise crossing should be considered. However, if one's goal is to preserve as much of the variation in founding genotypes as possible, a crossing strategy with fewer founders might be more desirable.

Careful crossing of founder strains results in more balanced haplotype representation

Haplotype frequency estimates for our synthetic populations suggest that a strategy involving pairwise crosses results in more balanced representation of founder strains (Figures 4-5, Supplementary Figures 4-7, and Table 4). It is clear than when haploid strains are simply pooled, there is potential for subsets of the founder strains to dominate and skew haplotype representation in the resulting populations. As such, we recommend a careful (i.e. pairwise)

crossing strategy when seeking to maximize founder haplotype representation. In addition, using

fewer founder haplotypes also generally results in a more even representation among them,

though we would not necessarily recommend against using more founders unless it is crucial to

achieve an even haplotype representation. In fact, we would argue that there are empirical

benefits to a more varied distribution of haplotype frequencies segregating within a synthetic

population. Specifically, in an E&R context, such a population creates opportunities to observe

selection on both rare and common haplotypes, and the relative contributions each to the

evolution of quantitative traits remains an unresolved question (e.g. Bloom et al. 2019).

Evidence for adaptation to outcrossing

While we interpret the differences between the different crossing strategies (i.e. between

all S-type and K-type populations) as being primarily driven by initial differences in sporulating

and mating efficiencies between strains and subsequent genetic drift, selection may also play a

role. For instance, we do find evidence that two specific regions of the genome exhibit similar

dynamics in haplotype frequencies, compared to similar prior work; Burke et al. (2014)

previously implicated these regions as potentially driving adaptation for frequent outcrossing.

We also compared our results to those of Linder et al. (2020) which features synthetic

populations constructed using the same founder strains present in this study. However, here we

did not find it was the case that haplotypes most common in their populations on average are also

the most common in our populations. For instance, Y12 is a common haplotype across all of our

populations (Table 4) with mean genome-wide frequencies ranging from 0.2 to ~0.5, but it does

not exceed a frequency of 0.06 in either of the synthetic populations described by Linder et al.

(*cf.* Table 3 of Linder et al. 2020). Similarly, YPS128 which has a mean frequency of 0.41 in one

of the Linder et al. (2020) populations appears at frequencies below 0.11 in all of our populations

except S4. That being said, it is worth noting that while there is overlap in the strains used

between these studies, maintenance protocols are different, and their populations include several

strains absent in our study. So, we can only say that no common founders appear to universally

favored when selection for frequent outcrossing is imposed.

Our analysis comparing initial SNP frequencies to those observed after 12 rounds of

outcrossing in each population also yields possible evidence for adaptation, though our

interpretation here is cautious. We do observe regions that produce peaks in significance in some

populations that overlap regions described by other studies to underlie traits related to

outcrossing (e.g. Figure 3D; peak in C16 in K8). However, peaks are rarely recapitulated across

two or more populations (Figure 3) and in some populations we observe no clearly defined peaks

at all (e.g. Figure 3B; population K4). As such, our results are at best suggestive. But, we find it

notable that a crossing strategy using 8 founding haplotypes leads to the most overlap with other

candidate regions identified in the literature. While this experiment has limited ability to

associate particular genomic regions and phenotypes related to outcrossing, this observation

leads us to speculate that using an intermediate number of founding haplotypes (e.g. 8) may lead

to an increased ability to localize candidate regions in an E&R experiment.

Given that any adaptation we did observe can only be ascribed to the outcrossing process

and associated handling protocols, we conclude that there is likely a great deal of value in

allowing newly-established synthetic populations to experience several cycles of outcrossing –

this can also be thought of as laboratory domestication – before any sort of new selective

pressure is imposed. To be explicit, if some other selection pressure was intentionally imposed

on our populations immediately following cycle 0, it would be very difficult to dissect the

specific genetic changes that might occur due to that pressure, other laboratory handling steps, or

general selection for reproduction via outcrossing.

Conclusions

The results of simulated E&R studies in which populations are sexually-reproducing and

adaptation is driven by standing genetic variation have led to general experimental design

recommendations that maximize genetic variation in the ancestral population, specifically by

increasing the number of starting haplotypes (Baldwin-Brown et al. 2014; Kofler and Schlötterer

2014). Here, we provide empirical results that also these recommendations. Across the metrics

we examined, we consistently find that a crossing strategy involving careful pairwise crosses

leads to populations with more standing genetic variation than those produced by simply mixing

founder genotypes in equal proportion. As such, using this sort of strategy would be our primary

recommendation for researchers aiming to establish recombinant populations from clonal strains

or isogenic lines for use in E&R studies. Using more founders also results in greater total levels

of genetic variation but comes at the cost of maximizing representation of all possible alleles

from a given set of founders. So, the number of founders to used should be chosen based upon

the specific goals of an E&R study, and the questions it hopes to test. Finally, our results should

be placed in the context of yeast biology. Meaning, given the high degree of genetic

differentiation between our yeast strains and the variation among them for mating and

sporulation efficiencies, the patterns we observe are likely more extreme than what might be

expected when creating a *Drosophila* synthetic population using either the K-type or S-type

approach. However, it is certainly conceivable that mating preferences and genetic

incompatibilities among isogenic *Drosophila* lines could impact levels of genetic variation and

haplotype representation when they are crossed. Our findings therefore perhaps warrant

consideration even when creating synthetic populations in non-yeast systems.

Data Availability

The raw sequence files generated over the course of this project are available through NCBI

SRA (BioProject ID: PRJNA732717). Core data files (tables with SNP and haplotype

frequencies, results of statistical analysis, etc.) are available through Dryad (XXXXXX). Core

scripts necessary to reproduce our results are available through Github

(https://github.com/mphillips67/Build-Paper).

Acknowledgements

We thank Anthony Long for his feedback the haplotype estimating analysis. We also thank

OSU's Center for Genome Research and Biocomputing for use of their computational and

sequencing resources. This work was supported by startup funds to M.K.B. (OSU's College of

Science), and M.A.P. is supported by an NSF Postdoctoral Fellowship (NSF 1906246).

References

Bailey, S.F. & Bataillon, T. (2016). Can the experimental evolution programme help us elucidate
the genetic basis of adaptation in nature? *Molecular Ecology,* 25, 203-216.

Baldwin-Brown, J.G., Long, A.D., & Thornton, K.R. (2014). The power to detect quantitative
trait loci using resequenced, experimentally evolved populations of diploid, sexual organisms.
Molecular Biology and Evolution, 31, 1040-1055.

Barghi. N., & Schlötterer, C. (2019). Shifting the paradigm in evolve and resequence studies,
from analysis of single nucleotide polymorphism to selected haplotype blocks. *Molecular*
Ecology, 28, 521-524.

Barghi, N., Tobler, R., Nolte, V., Jaksic, A.M., Mallard, F., Otte, K.A., … Schlötterer, C. (2019).
Genetic redundancy fuels polygenic adaptation in *Drosophila*. *PLOS Biology,* 17, e3000128.

Baym, M., Kryazhimskiy, S., Liberman, T.D., Chung, H., Desai, M.M., & Kishony, R. (2015).
Inexpensive multiplexed library preparation for megabase-sizes genomes. *PLoS One*, 10,
e0131262.

Beissinger, T.M., Rosa, G.J.M, Kaeppler, S.M., Gianola, D., & de Leon, N. Defining window-
boundaries for genomic analyses using smoothing spline techniques. (2015). *Genetics Selection*
Evolution, 47, 30.

Bergström, A., Simpson, J.T., Salinas, F., Barré, B., Parts, L., Zia, A., … Liti, G. (2014). A high-
definition view of functional genetic variation from natural yeast genomes. *Molecular Biology*
and Evolution, 31, 872-888.

Bloom, J.S., Boocock, J., Treusch, S., Sadhu, M.J., Day, L., Oates-Barker, H., Kruglyak, L.
(2019). Rare variants contribute disproportionately to quantitiative trait variation in yeast. eLife,
8,e49212

Boyle, E.A., Li, Y.I., & Pritchard, J.K. (2017). An expanded view of complex traits: from
polygenic to omnigenic. *Cell*, 169:1177-1186.

Burke, M.K., Dunham, J.P., Sharestani, P., Thornton, K.R., Rose, M.R., & Long, A.D.
(2010). Genome-wide analysis of a long-term evolution experiment
with *Drosophila*. *Nature,* 467, 587-590.

Burke, M.K. (2012). How does adaptation sweep through the genome? Insights from long-term
selection experiments. *Proceedings of the Royal Society B,* 279, 5029-5038.

Burke, M.K., Liti, G., & Long, A.D. (2014). Long standing genetic variation drives repeatable
experimental evolution in outcrossing populations of *Saccharomyces cerevisiae*. *Molecular*
Biology and Evolution, 32, 3228-3239.

Burke, M.K., McHugh, K.M., Kutch, I.C. 2020. Heat shock improves random spore isolation in
diverse strains of *Saccharomyces cerevisiae. Frontiers in Genetics*, 11: 597482

Conway, J.R., Lex, A., & Gehlenborg, N. (2017). UpSetR: an R package for the visualization of
intersecting sets and their properties. *Bioinformatics*, 33:,2938-2940.

Cubillos, F.A., Louis, E.J., & Liti, G. (2009). Generation of a large set of genetically tractable
haploid and diploid *Saccharomyces* strains. *FEMS Yeast Research*, 9, 1217-1225.

Cubillos, F.A., Parts, L., Salinas, F., Bergstrom, A., Scovacricchi, E.,… Liti, G. (2013). High-
resolution mapping of complex traits with a four-parent advanced intercross yeast population.
Genetics, 195, 1141-1155.

Franssen, S.U., Nolte, V., Tobler, R., & Schlötterer C. (2015). Patterns of linkage disequilibrium
and long range hitchhiking in evolving experimental *Drosophila* melanogaster
populations. *Molecular Biology and Evolution,* 32, 495-509

Graves, J.L., Hertweck, K.L., Phillips, M.A., Han, M.V., Cabral, L.G., Barter, T.T., … Rose
MR. 2017. Genomics of parallel experimental evolution in Drosophila. *Molecular Biology and*
Evolution, 34, 831–842.

Harbison, S.T., Negron Serrano, Y.L., Hansen, N.F., & Lobell, A.S. (2017). Selection for long
and short sleep duration in *Drosophila melanogaster* reveals the complex genetic network
underlying natural variation in sleep. *PLoS Genetics*, 13, e1007098.

Huang, Y., Wright, S.I., & Agrawal, A.F. (2014). Genome-wide patterns of genetic variation
within and among alternative selective regimes. *PLoS Genetics,* 10, e1004527.

Jha, A.R., Miles, C.M., Lippert, N.R., Brown, C.D., White, KP, & Kretiman, M. (2015). Whole-
genome resequencing of experimental populations reveals polygenic basis of egg-size variation
in *Drosophila melanogaster*. *Molecular Biology and Evolution,* 32, 2616-2632.

King, E.G., Merkes, C.M., McNeil, C.L., Hoofer, S.R., Sen, S., Broman, K.W., Long, A.D.,
Macdonald, S.J. (2012). *Genome Research*, 22, 1558-1566.

Kofler, R., & Schlötterer, C. (2014). A guide for the design of evolve and resquencing studies.
Molecular Biology and Evolution, 31,473-482.

Linder, R.A., Majumbder, A., Chakraborty, M., & Long, A.D. (2020). Two synthetic 18-
wayoutcrossed populations of diploid budding yeast with utility for complex trait dissection.
Genetics, 215, 323-342.

Long AD, Liti G, Luptak A, Tenaillon O. (2015). Elucidating the molecular architecture of
adaptation via evolve and resequence experiments. *Nature Reveiews Genetics,* 16, 567-582.

Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., … Visscher, P.M.
(2009). Finding the missing heritability of complex diseases. *Nature*, 461, 747-753.

Michalak, P., Kang, L., Schou, M.F., Garner, H., & Loeschke, V. (2018). Genomic signatures of
experimental adaptive radiation in Drosophila. *Molecular Ecology,* 28, 600– 614.

Nouhaud, P., Tobler, R. Nolte, V., Schlötterer, C. (2016). Ancestral population reconstitution
from isofemale lines as a tool for experimental evolution. *Ecology and Evolution*, 6, 7169-7175.

Orozco-ter Wengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T., & Schlötterer C. (2012).
Adaptation of *Drosophila* to a novel laboratory environment reveals temporally heterogeneous
trajectories of selected traits. *Molecular Ecology, 21*, 4931–4941.

Phillips, M.A., Rutledge, G.A., Kezos, J.N., Greenspan, Z.S., Talbott, A., Matty, S., …
Shahrestani, P. (2018). Effects of evolutionary history on genome wide and phenotypic
convergence in Drosophila populations. *BMC Genomics*, 19, 743.

Phillips, M.A., Kutch, I.C., Long, A.D., & Burke, M.K. (2020). Increased time sampling in an
eolve-and-resequence experiment with outcrossing *Saccharomyces cerevisiae* reveals multiple
paths of adaptive change. *Molecular Ecology*, 29, 4898-4912.
661
662 Schlötterer, C., Tolber, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of inidviduals –
663 mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics,* 15,
664 749-763.

665 Taus, T., Futschik, A., & Schlötterer, C. (2017). Quantifying selection with pool-seq time series
666 data. *Molecular Biology and Evolution,* 34, 3023-3034.

Teotónio, H., Carvalho, S., Manoel, D., Roque, M., & Chelo, I.M. (2012) Evolution of outcrossing in experimental populations of *Caenorhabditis elegans*. *PLoS One*, 7, e35811.

Tobler, R., Franssen, S.U., Kofler, R., Orozco-terWengel, P., Nolte, V., Hermisson, J., & Schlötterer, C. (2014). Massive habitat-specific genomic response in D. *melanogaster* populations during experimental evolution in hot and cold environments. *Molecular Biology and Evolution,* 31, 364-375.

Turner, T.L., Stewart, A., Fields, A.T., Rice, W.R., & Tarone, A.M. (2011). Population-based resequencing of experimentally evolved populations reveals the genetic basis of body size variation in *Drosophila melanogaster*. *PLoS Genetics,* 7, e10001336.

Vlachos, C., & Kofler, R. (2019) Optimizing the power to identify the genetic basis of complex traits with evolve and resequence studies. *Molecular Biology and Evolution*, 35, 2890-2905.

Zhou, D., Gersten, M., Visk, D.W., Bashir, A., Xue, J., Frazer, K.A., … Haddad, G.G. (2011). Experimental selection of hypoxia-tolerant *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences USA,* 108, 2349–2354.

Figures

Figure 1. UpSet plot for the two synthetic populations made using 4 (A), 8 (B), and 12 (C)

founders. The horizontal bars indicate the size of the SNP sets for each population at cycles 0, 6

and 12, and the total possible number of SNPs given the founder strains used to make them. The

vertical bars represent distinct intersections between the different highlighted sets (as indicated

by the black circles). For a given plot, the first vertical bar represents SNPs found in all

experimental populations, the second bar represents SNPs found only in the S-type populations,

the third bar represents SNPs found only in the K-type populations, and the fourth bar represents

possible SNPs, given relevant founder genotypes, that are not observed in any experimental

population.

717   Figure 2. Site frequency spectra (SFS) for all populations at each timepoint samples were taken

718   for DNA sequencing; immediately after construction, or "cycle 0" (left panels), after 6 cycles of

719   outcrossing (middle panels), and after 12 cycles of outcrossing (right panels).

Figure 3. Results from Pearson's χ^2 test comparing SNP frequencies initially (cycle 0), and after 12 cycles of outcrossing for all populations featured in this study. In each panel, red and green boxes indicate regions of genome and genes associated with selection for frequent outcrossing from Cubillos et al. (2013) and Burke et al. (2014), respectively.

Figure 4. Haplotype frequency estimates for population S4 based on SNP frequencies after 12

cycles of outcrossing. Frequencies were estimated using sliding windows with a length of 30KB

and a 1KB step size. Panels are ordered to match pairings in the first round of crossing (YPS128

was crossed with DBVPG6765, Y12 was crossed with DBVPG6044)

Figure 5. Haplotype frequency estimates for population K4 based on SNP frequencies after 12

cycles of outcrossing. Frequencies were estimated using sliding windows with a length of 30KB

and a 1KB step size. Panels are ordered to match pairings in the first round of crossing (YPS128

was crossed with DBVPG6765, Y12 was crossed with DBVPG6044).

746 Tables

747 Table 1. Strains used to create the synthetic populations featured in this study. Strains are

748 arranged in the rows in the table to indicate which specific pairs were crossed in the S-type

749 populations.

Populations	Founder Strains	
	Mat a	Mat *alpha*
S4 & K4	DBVPG6765	YPS128
	DBVPG6044	Y12
S8 & K8	DBVPG6765	YJM981
	DBVPG6044	L_1528
	BC187	Y12
	L_1374	YPS128
S12 & K12	DBVPG6765	YJM981
	DBVPG6044	Y12
	BC187	L_1528
	SK1	273614N
	L_1374	YPS128
	YJM975	UWOPS05_217_3

762 Table 2. Number of possible SNPs and those actually observed in all synthetic populations at

763 each cycle where samples were taken for sequencing. Percentages are relative to total possible

764 number of SNPs for a given set of founders (n=4, 8, or 12). We estimate that approximately 15-

765 20 asexual generations occur between each cycle of outcrossing.

	S4	K4	S8	K8	S12	K12
Possible SNPs	112,738		122,663		155,516	
Cycle 0	91,658 **(81%)**	71,771 **(64%)**	98,864 **(81%)**	94,965 **(77%)**	101,134 **(65%)**	50,352 **(32%)**
Cycle 6	75,712 **(67%)**	69,954 **(62%)**	78,078 **(64%)**	62,537 **(51%)**	79,070 **(51%)**	38,853 **(25%)**
Cycle 12	57,405 **(51%)**	39,085 **(35%)**	80,941 **(66%)**	61,636 **(50%)**	73,894 **(48%)**	5,171 **(3%)**

785 Table 3. Mean genome-wide heterozygosity for all synthetic populations at each cycle where

786 samples were taken for sequencing. Mean heterozygosity is often, but not always, higher in the

787 S-type populations (shaded gray) relative to their K-type counterparts at a given timepoint.

	S4	K4	S8	K8	S12	K12
Cycle 0	0.38	0.47	0.36	0.28	0.36	0.31
Cycle 6	0.32	0.41	0.33	0.25	0.32	0.25
Cycle 12	0.20	0.17	0.31	0.25	0.28	0.04

Table 4. Mean genome-wide haplotype frequencies for all founder populations in each synthetic
population based on SNP frequencies observed after 12 cycles of outfrossing (*note*: shaded cells
indicate that a founder not used for that particular population).

Founder strain	K4	S4	K8	S8	K12	S12
DBVPG6765	0.486	0.237	0.019	0.027	0.001	0.127
DBVPG6044	0.004	0.291	0.004	0.144	0.169	0.186
BC187			0.178	0.016	0.001	0.01
SK1					0.353	0.12
L_1374			0.23	0.23	0.001	0.011
YJM975					0.001	0.105
YPS128	0.012	0.249	0.009	0.111	0.002	0.005
Y12	0.498	0.222	0.331	0.325	0.468	0.21
273614N					0.001	0.065
L_1528			0.203	0.023	0.001	0.012
UWOPS05_217_3					0.002	0.018
YJM981			0.027	0.197	0.001	0.131

Table 5. Comparison of putatively adaptive haplotypes from Burke et al. (2014) with observed haplotype frequency changes in synthetic populations used in this study. While both studies employed similar protocols Burke et al. (2014) implemented 18 outcrossing cycles while we only implemented 12. The 2014 study is used as a point of reference here because it features 12-fold replication, while in the present study features no within-treatment replication. We specifically targeted the window containing the most significant marker in each region based on the Burke et al. (2014) results. Notable (> 10%) increases in focal haplotype frequency are shown in green.

Candidate region	Haplotype change in Burke et al. (2014)		Haplotype change in populations of the present study					
	Driving haplotype	Mean increase	S4	K4	S8	K8	S12	K12
A- C07:950000-975000	Y12	+0.23	+0.49	+0.29	+0.36	+0.09	+0.16	+0.66
B- C09:65000-80000	DBVPG6044	+0.24	+0.70	-0.01	-0.20	-0.16	+0.15	+0.49
C- C11:615000-620000	DBVPG6044	+0.29	-0.16	+0.01	-0.06	-0.15	-0.11	+0.49
E- C16:445000-460000	Y12	+0.59	+0.29	+0.51	+0.11	+0.93	+0.50	+0.98

Supplementary Figure 1. Unrooted phylogeny showing the general evolutionary relationships

among the strains used in this work, as well as their geographical (colored boxes) and contextual

(colored circles) origins. This tree was constructed using 180,276 bi-allelic SNPs that we

observed to vary across the 12 haploid founder strains sequenced in this study. These SNPs were

used to construct a Newick file which was plotted with the R package *ggtree*.

Supplementary Figure 2. Crossing schematics used to create S4 (A), S8 (B), and S12 (C).

Supplementary Figure 3. Expected cycle 0 SFS for populations created using four (A), eight (B)

and twelve founder strains (C). For specific founders used in each scenario, see Table 1.

874 Supplementary Figure 4. Haplotype frequency estimates for population S8 based on SNP

875 frequencies in cycle 12. Haplotype frequencies were estimated using sliding windows with a

876 length of 30KB and a 1KB step size. Panels are ordered to match pairings in the first round of

877 crossing (e.g. YJM981 was crossed with DBVPG6765, etc.).

Supplementary Figure 5. Haplotype frequency estimates for population K8 based on SNP

frequencies in cycle 12. Haplotype frequencies were estimated using sliding windows with a

length of 30KB and a 1KB step size. Panels are ordered to match pairings in the first round of

crossing (e.g. YJM981 was crossed with DBVPG6765, etc.).

Supplementary Figure 6. Haplotype frequency estimates for population S12 based on SNP

frequencies in cycle 12. Haplotype frequencies were estimated using sliding windows with a

length of 30KB and a 1KB step size. Panels are ordered to match pairings in the first round of

crossing (e.g. YPS128 was crossed with DBVPG6765, etc.).

Supplementary Figure 7. Haplotype frequency estimates for population K12 based on SNP

frequencies in cycle 12. Haplotype frequencies were estimated using sliding windows with a

length of 30KB and a 1KB step size. Panels are ordered to match pairings in the first round of

crossing (e.g. YPS128 was crossed with DBVPG6765, etc.).