Title: Predicting Bird Distributions Under Global Change

Authors & Affiliations:

MacPherson[1], M.P., Burgio[2,3], K.R., DeSaix[4], M.G., Freeman[5], B.G., Herbert[6], J., Herman[7], R., Jirinec[8], V., Shonfield[9], J., Slager[10], D.L., van Rees[11], C.B., and J.E. Jankowski[5]

1. Louisiana State University Museum of Natural Sciences, Louisiana State University, Baton Rouge, LA, USA
2. New York City Audubon Society, New York, NY, USA
3. Department of Ecology and Evolutionary Ecology, University of Connecticut, Storrs, CT, USA
4. Department of Biology, Colorado State University, Fort Collins, CO, USA
5. Biodiversity Research Centre and Department of Zoology, University of British Columbia, Vancouver, BC, Canada
6. Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA
7. Department of Ecology and Evolution, Stony Brook University, NY
8. School of Renewable Natural Resources, Louisiana State University and LSU AgCenter, Baton Rouge, LA, USA
9. LGL Limited, Cambridge, ON, Canada
10. Department of Biology, University of Washington, Seattle, WA, USA
11. River Basin Center & Odum School of Ecology, University of Georgia

Author Contributions:

Maggie MacPherson - concept, symposium organizer, symposium participant, writing & revising, editing full manuscript, topic editor (abstract, theory) and co-editor (temporal resolution of data types), figures, tables.

Kevin R. Burgio - concept, overall project support, symposium participant, writing & revising, conceptualizing and constructing figures, editing full manuscript, topic co-editor (methods & data types for spatial data)

Matthew G. DeSaix – symposium participant, writing & revising, topic co-editor (temporal resolution of data types)

Ben Freeman - symposium participant, writing, publication approach/ideas, topic editor (discussion)

Vitek Jirinec - symposium participant, writing & revising, topic co-editor (logistical methods and data types for occurrence data), figures.

John Herbert - symposium participant, writing, topic co-editor (methods & data types for spatial data)

Julia Shonfield - symposium participant, writing & revising, topic editor (introduction)

David Slager – symposium participant, writing & revising, topic editor (literature cited), topic co-editor (logistical methods and data types for occurrence data)

Rachael Herman - writing, topic co-editor (logistical methods and data types for occurrence ddata)

Charles van Rees – writing & revising, topic editor (analytical methods), figures.

Jill Jankowski – concept, symposium participant, writing & revising, editing full manuscript

ABSTRACT

Global change creates an urgent need to predict spatial responses of biota to support the

conservation of sufficient habitat to maintain biodiversity. We present species distribution model

theory and a synthesis of avian literature on approaches to collecting occurrence data, selecting

explanatory variables and analytical processes currently in use to predict future distributions. We

find that interpreting the validity of current predictive distributions is hindered by variation in

spatio-temporal resolution of data sets that force hypothesis testing under the Grinnellian niche

concept. Broadly, the capacity of species to shift their geographic ranges under land use and

climate change is expected to be limited by both large scale (i.e., the physiological or

fundamental niche) and small scale (i.e., the realized or tolerance niche) factors. We highlight the

strengths and weaknesses of widely used explanatory variables and analytical approaches

tailored to macrohabitat characteristics and the Grinnellian niche concept. This synthesis

addresses if and how current approaches align with theory and makes recommendations for

future directions to improve the accuracy of predictive distribution modelling.

KEY WORDS: species distribution model, environmental niche model, predictive distribution

modeling, global change, climate change.

INTRODUCTION

Anthropogenic pressures on climate and land cover lead to altered ecosystems across, and have

consequences for the geography of species distributions (Parmesan & Yohe, 2003). This creates

an urgent need for understanding species spatial responses to global change and ensuring

conservation of suitable habitat that supports population persistence and conserves biodiversity.

The accuracy of current predictive species distribution models, however, is limited due to

variation in temporal and spatial resolution of datasets. The capacity of species to change their

geographic ranges is limited by both large-scale factors (i.e., things that distinguish range-wide

patterns like temperature and rainfall patterns - macrohabitat) and small-scale factors (i.e., things

that drive responses by individuals like biotic interactions and resource availability -

microhabitat). In this review, we explore the utility of predictive distribution modelling in

guiding conservation action in birds. We focus on avian biodiversity because birds are excellent

indicators of environmental change, may be more vulnerable to climate change than some other

taxa (e.g., mammals), and they have the most widespread and in-depth databases on distributions

(Morrison, 1986; Riddell et al., 2021). Insights from avian literature may also be generalizable to

other taxa because birds display a wide variety of life history strategies, and citizen science

initiatives and long-term research programs have yielded an abundance of predictive distribution

literature (Şekercioğlu et al., 2019).

A multitude of factors can ultimately influence how species, populations and individuals

respond to changing climate regimes. Individuals may lack flexibility in tracking relevant

environmental cues to adjust the timing of life history events and to accommodate novel

ecological circumstances. Alternatively, even when species are flexible in timing life history

events, they may be unable to modulate behavior in response to environmental anomalies that

affect recruitment. For example, advancing egg laying dates to match earlier spring warming

temperatures may have negative consequences for nestling survival when the chances of cold

snaps are decoupled from warming trends caused by climate change (Shipley et al., 2020).

Predicting how species distributions could be altered (shift, contract or expand) with global

change may be species-specific (Hallman & Robinson, 2020; Radosavljevic & Anderson, 2014),

but research into how particular axes of climate change affect species with different life history

strategies, foraging guilds or habitat affinities provides promising insight that may apply to a

wider number of unstudied species (e.g., changing rainfall regimes - Brawn et al., 2017;

advances in spring phenology and breeding success - Sander et al., 2020).

A major gap in our understanding of how organisms may respond to changing

environments lies in movement behavior. In birds, research on migration routes and resource

tracking is well established in the Northern Hemisphere (Faaborg et al., 2010; Thorup et al.,

2017), although much less information is available for species that undergo seasonal movements

in the Southern Hemisphere. In the speciose Neotropics, >50% of bird species are known to be

migratory, and an increase in knowledge of seasonal movements in remaining species could

augment this estimate this by up to 15% (Jahn et al., 2020; Rappole, 1995). Complexity is added

by varied seasonal movements that avian species undertake, including austral migrants (Jahn et

al., 2004, 2013), elevational migrants (Barçante et al., 2017; Hsiung et al., 2018), partial

migrants (Sekercioglu, 2010) and others with intratropical migration routes (e.g., Resplendent

Quetzal - Papeş et al., 2012; Three-wattled Bellbird - Powell & Bjork, 1995) or extended daily

foraging routines (e.g., Oilbirds - Holland et al., 2009). Migratory bird species have been shown

to decline when the timing of annual life history events are mismatched with peaks in food

availability (Møller et al., 2008; Sander et al., 2020). On the other hand, multiple species of

Nearctic-Neotropical migrants are advancing dates of spring migration and breeding behavior in

response to warming spring temperatures (Dunn & Møller, 2019; Pecl et al., 2017; Shipley et al.,

2020), offering support to the idea of "evolutionary rescue" such that existing variation within

annual programs may yield previously unknown adaptive potential (Helm et al., 2019). In this

review, we use literature on predictive bird distribution modelling to inform future work

predicting the importance of dispersal limitation, biotic interactions and estimates of the

fundamental niche to understand climate-driven distributional shifts in birds and other taxa.

With specific attention on global temperature increases caused by climate change,

responses to shifting thermal environments will be varied and complex. Janzen's "seasonality

hypothesis" suggests that tropical organisms should be particularly sensitive to anomalies in

ambient conditions because their physiology is adapted to a relatively constant environment

(Janzen, 1967). This idea has since received much empirical support (Sheldon et al., 2018), and

highlighted the tropics for having a high concentration of thermal specialists that are vulnerable

to climate change (Mora et al., 2013). Tropical ectotherms (i.e., arthropod prey) tend to live near

the upper limits of their thermal optima and are therefore sensitive to even small temperature

increases (Deutsch et al., 2008). Endotherms (like birds), on the other hand, have the capacity to

tolerate ambient temperatures past their thermoneutral zone by raising metabolic activity to

produce evaporative cooling, but this increases energy and water demands at temperature

extremes (Gardner et al., 2011; Huey et al., 2012). Still, few studies have considered the

potential effects on presumably more thermally flexible endotherms due to changes in a more

thermally sensitive ectothermic prey base (Grady et al., 2019). An important consequence

stemming from Janzen's hypothesis is that lowland tropical species could be particularly affected

by novel conditions supplied by global change, resulting in broad-scale biotic attrition, except for

populations immediately adjacent to montane regions, which would have some access to upslope

thermal refugia (*sensu* Brodie et al., 2017; Colwell et al., 2008). At least some studies have

shown that tropical birds are unlikely to experience acute thermal limits to their distributions

(Londoño et al., 2015, 2017). While species may have a sufficient thermal margin to defend

against climate warming (Pollock et al., 2021), they will likely experience added

thermoregulatory costs in response to shifting thermal regimes or movements (Londoño et al.,

2017). As the effects of increased temperature on populations can interact with critical ecological

and life-history parameters, temperature is expected to impact both current distributions and

species' responses to changing thermal environments.

Our review aims to synthesize theory, methods, and data sources widely used to predict

bird distributions under global change, and to highlight best-practices and opportunities for

innovation and refinement. Understanding where species occur and why some areas are occupied

but not others is essential for developing conservation and management plans. For most species,

knowledge of their distribution is limited to areas that have been surveyed. However, predictive

distribution modelling is important to identify areas for protection or management. Modelling

wide-ranging, unevenly distributed, cryptic, or difficult to survey species all present problems for

understanding factors driving occupancy. Understanding how behaviors and life history traits are

linked to patterns of occupancy requires focused investigation but are nevertheless critical layers

for informative distribution models. Here, we bring together the issues complicating predictive

distribution modelling to offer a solutions-oriented approach to habitat conservation under global

change.

PREDICTIVE DISTRIBUTION MODEL THEORY

The current dogma of species distribution modelling falls into two categories: 1) species

distribution models (hereafter SDMs) that comprise presence records and abiotic data, and 2)

ecological niche models (hereafter ENMs) that explicitly estimate the accessible environment

(Soberón & Peterson, 2005). These approaches are meant to have high spatial accuracy but are

not meant to inform on cause-and-effect species-habitat relationships (Merow et al., 2013).

Models can be based on different niche perspectives, which present unique frameworks for

estimating drivers of occupancy across a species' range. The Grinnellian class of niche reflects

abiotic drivers of site occupancy (Elton, 1927; Soberón, 2007; Wisz et al., 2013) and SDMs are

mainly built under the Grinnellian niche concept (Figure 1; Saupe et al., 2012). Coarse-scale

variables describing ecosystem characteristics are often the most relevant for predicting shifting

distributions across large spatial extents. In comparison, the Eltonian class of niche includes

estimates of biotic interactions and resource-consumer dynamics that are only quantifiable at

local scales (Figure 1; Elton, 1927; Soberón, 2007). Finally, the Hutchinsonian class of niche

reflects the functional role of a species, which is often estimated using functional traits and

species' habitat requirements based on functional traits (Rosado et al., 2016), and projects the

probability of site occupancy beyond study areas, including estimates of dispersal probabilities in

ENMs (Figure 1). A species' ability to move beyond their current realized niche depends on the

relative importance of abiotic factors, biotic interactions, and dispersal probability that operate at

different spatial scales (Jankowski et al., 2013).

Figure 1. The "BAM" diagram. The relationship between "B" - biotic, "A" - abiotic, and "M" -

movement (also thought of as "access" and defined by dispersal ability) with respect to the

authors of various niche theories (left). The location of realized, tolerance, and fundamental

niche along the intersection of two hypothetical climate variables (right). Sax et al. (2013)

present an update to niche theory that considers three components of a species niche: 1) the

"realized niche", i.e., conditions within the native range, 2) the "fundamental niche" - conditions,

in which a species could thrive if it were introduced there, and 3) the "tolerance niche" -

conditions, in which individuals could survive, but likely unable to maintain populations over the

long-term. The fundamental niche is the set of appropriate abiotic conditions, and the realized

niche is a smaller area where both abiotic and biotic environments are suitable. Theoretically not

all of the realized niche could be accessible and so the potential range is yet smaller because it

reflects only the area that is suitable and also accessible to the species. Occurrence is not

expected outside of the colored areas. The occupied area is represented by G_0, and G_1 reflects

appropriate conditions that have not or cannot be accessed by the species (as described by Saupe

et al., 2012).

Assessing occurrence or abundance along abiotic gradients is important for predicting

future range shifts under global change, as this relates directly to species' and populations'

responses to the environment. The gradual shifts in temperature and rainfall regimes can be

expected to align with changes in species responses, reflecting the shifts in optimal ambient

conditions and resulting in altered patterns of probability of occurrence or abundance. It is

important to note that ENM products are only able to predict the probability of site occupancy

within the range of abiotic conditions measured within current range boundaries, so will be

limited where climate change introduces novel climates (e.g., Williams et al., 2007).

Spatial ecology theory dictates that species respond to continuous environmental

gradients through gradual changes in abundance, as individuals experience shifting conditions

moving toward or away from environmental optima over space (Austin, 2005, 2007). However,

populations may also show thresholds in abiotic tolerances or in response to biotic factors that

change their response shape (e.g., creating asymmetric or skewed response functions; e.g.,

Oksanen & Minchin, 2002). For example, the asymmetric abiotic stress (AASL) hypothesis

predicts a steeper decrease in a species' probability of occurrence toward the more stressful end

of a species' distribution, which has been supported in a number of vascular plant species

(Dvorský et al., 2017). Other threshold-type effects can result in asymmetric responses shown

across multiple species in communities (e.g., sharp ecotone boundaries, appearance of dominant

predators or competitors; Jankowski et al., 2013). Identifying response shapes is of practical

interest to understand the impact of abiotic and biotic factors on occupancy. Comparisons across

species distinguish commonalities and differences in responses to gradients and can point to key

environmental variables driving patterns in community organization.

Heterogeneity within landscapes is important to consider in determining whether

populations would be able to respond (in time and space) to changing conditions. Landscapes

can vary in ways that affect both the phenology of resources (time) or availability of habitat

(space); such variability can be tracked if conditions change at a pace that populations could

respond to. Examples of this include species adapted to annually ephemeral or temporally patchy

habitats such as wetlands and grasslands; species accustomed to such environments may have

dispersal strategies that predispose them to better track climate (e.g., *Elaenia cristata* - Ritter et

al., 2021). In other cases, habitat and resources are spatially patchily distributed, and the ability

of species to respond to changing conditions may be more taxon-specific for those that rely on

spatiotemporal heterogeneity that is naturally patchy. Examples of this include habitat specialists

in naturally patchy habitats such as Amazonian white sand habitats (e.g., *Elaenia ruficeps* -

Ritter et al., 2021).

Microclimate refugia allow animals to buffer themselves from acutely unfavorable

conditions. One example of this is through behavioral thermoregulation. Ambient temperature is

almost never constant in terrestrial environments—it varies by time, and by habitat, often at very

small scales (Scheffers et al., 2014, 2017). Biologists have long understood that mobile animals

exploit thermal heterogeneity to maintain optimal body temperature (Angilletta Jr & Angilletta,

2009; Angilletta et al., 2009; Cowles & Bogert, 1944; Porter et al., 1973; Stevenson, 1985). For

cold-adapted species in a heating world, this can be achieved by shifting activity times to cooler

periods of the year and day, and by moving to microhabitats that supply cooler microclimates.

This capacity to compensate for unfavorable ambient temperature by behavioral

thermoregulation, known as the "Bogert Effect" (Huey et al., 2003), can at least partly mitigate

the harmful effects of climate warming (Huey et al., 2012). However, behavioral

thermoregulation is contingent on access to cooler areas and periods; animals already occurring

in the most buffered environments have limited options for escape when conditions change.

Furthermore, individuals could experience additional pressured from biotic interactions with

species moving into their habitat to thermoregulate (Huey et al., 2012).

Dispersal limitation is one of three fundamental aspects of distribution theory (Figure 1).

Understanding barriers to distributions caused by dispersal limitation has been of great interest,

for example, in bird species unable to cross areas of unsuitable habitat, and has been studied both

experimentally (e.g., Moore et al., 2008) and theoretically (e.g., Ribas et al., 2012, 2018).

Selection pressures driving morphology, such as migratory compared to sedentary life histories,

may also play a role in physically limiting dispersal distance (Capurucho et al., 2020; Claramunt

et al., 2012; MacPherson et al., 2021; Sheard et al., 2020). Life history strategies also dictate the

demographic trends in dispersal; for example, in many species it is the inexperienced young that

disperse into new areas away from the territories of their parents (e.g., Florida Scrub Jay - Suh et

al., 2020). Dispersal limitation is thus a complex and difficult-to-quantify aspect of predictive

distribution modelling that is nevertheless fundamental for predicting the probability of range

shifts with global change. Coupling studies of distribution modelling with quantifiable measures

of dispersal capacity is an important next step toward making more informed predictions of

species responses to changing environments (Travis et al., 2013; Urban et al., 2013).

Generally, when confronted with climate change, species change in one of three ways: 1)

by going extinct, 2) surviving by shifting their geographic range, or 3) evolving in place.

Evidence from the fossil record suggests all three responses are common (Graham et al., 1996;

Smith et al., 1995). The capacity to adapt or evolve *in situ* in response to climate change has

been relatively understudied (Parmesan & Matthews, 2005). A recent meta-analysis of existing

studies of phenological and morphological adaptations to climate change concluded that

adaptation, while variable among taxa, will not likely work on a sufficient scale to mitigate the

worst effects of ongoing climate change, whereas phenological adaptation (e.g., timing of

reproduction or migration) may mitigate such effects, albeit imperfectly (Radchuk et al., 2019).

The expectation that species with short generation times should be able to survive by evolving in

place remains understudied. Advancing such research is hampered by the difficulty of

conducting experiments on large numbers of species at sufficient time scales needed to develop a

comprehensive framework (Holt et al., 2005; Pierson et al., 2015). Variability among taxa and

short time scales signal that selection pressure has yet to be strong enough to produce measurable

changes, though it appears that species shift their phenology faster where climate change velocity

is faster (Poloczanska et al., 2013). Among abiotic drivers, mainly precipitation changes (but

also temperature) seem to underlie adaptive responses to climate change (Caruso et al., 2017;

Siepielski et al., 2017). Dramatic changes in climate over the remainder of this century are

expected (Figure 2) with many species likely unable to survive in the areas they presently occupy

(Ceballos et al., 2015; Sax et al., 2013). Ramping up efforts that prioritize long-term monitoring

of populations will be important for informing predictions of the resiliency of species to climate

change.

Figure 2. **Current climate and predicted change in climate of global terrestrial biomes**.

Climographs of (A) temperate and (B) tropical and subtropical global terrestrial biomes.

Projected changes in temperature and precipitation regimes for (C) temperate and (D) tropical

and subtropical global terrestrial biomes. While some areas are predicted to have increased or

decreased precipitation in the future, no areas are expected to have lower or stable temperatures.

Temperature increases are predicted to be higher in temperate biomes, while changes in

precipitation are greater for tropical biomes. Climographs built from The World Wildlife Fund

terrestrial ecoregions (https://www.worldwildlife.org/publications/terrestrial-ecoregions-of-the-

world) and current climate (2021-2040) from CanESM5 middle of the road scenario (SSP3-7.0)

accessed from Worldclim (https://www.worldclim.org/data/cmip6/cmip6climate.html). The

projected change is modeled as the difference between current climate (2021-2040, described

above) and future climate using the worst-case scenario (SSP4-6.0; 2081-2100) from CanESM5

global climate model data (Meinshausen et al., 2020). Species' presence, and thus biodiversity, is

highly influenced by biome type, which shifts from one form to another according to temperature

and rainfall regimes (Whittaker 1975). Projected changes in temperature and precipitation may

cause global terrestrial biomes to shift in space, but spatial gains and losses by different biomes

over time may not occur in synchrony with habitat needs to maintain biodiversity (Dorazio et al.,

2015). R code for producing this figure can be found at

https://github.com/mmacphe/Global_Change_Biomes.

Extinction risk (and conversely, population viability) centers on declining population and

small population paradigms. Population declines are driven by deterministic (demographic,

abiotic, and biotic) factors that cause reduction to small numbers (e.g., the Allee effect), and also

by stochastic factors, which dominate the extinction dynamics of small populations (Caughley,

1994; Morris & Doak, 2002; Smith et al., 2021). The abiotic shifts in conditions due to climate

change and the resulting shifts in species distributions play a role in both types of population

dynamics. Shifting distributions can cause temporary or permanent reductions in habitat area

(Thomas et al., 2006; van Rees & Reed, 2018), lowering carrying capacity and driving

population declines at local or regional scales, or reducing connectivity and creating isolated

subpopulations that are more susceptible to small-population dynamics (Anderson et al., 2009).

The degree to which changing distributions result in commonly observed conservation impacts

depends on a diversity of species traits including habitat preference and associations, phenotypic

plasticity, and current range and population size (Jiguet et al., 2007; Visser, 2008). Notably,

research based solely on changes in distributions tends to predict widespread extinctions due to

the aforementioned mechanisms, but plasticity, dispersal, and adaptation appear to have been key

for many species' resilience to past and current climate change (Moritz & Agudo, 2013). For

example, the Galapagos finches are a well-known example of phenotypic plasticity in bill

morphology in response to environmental change (Grant & Grant, 2020; Grant & Weiner, 2017;

Weiner, 1994).

OCCURRENCE DATA FOR PREDICTIVE DISTRIBUTION MODELLING

Occurrence data through time is the foundation for understanding species responses to global

change. To assess whether a change in distribution has occurred or will occur, researchers

require historical records to establish a baseline for comparison, followed by surveys at one or

more subsequent time points. Although birds are among the best studied animals, early baseline

inventories are still rare, and often consist of field notes (Reznick et al., 1994), which present

challenges particularly in reconstructing past abundances (Shaffer et al., 1998). Alternatively,

museum records are among the best sources of historical occurrence data, though they primarily

supply presence/absence data only (Shaffer et al., 1998).

In contrast to sparse information from the past, modern ornithologists benefit from

technological advances that offer unprecedented information on bird occurrence. As birds are

such mobile organisms, these include devices that can track individuals beyond specific field

research sites (satellite imagery - Borowicz et al., 2018; sophisticated automated biodiversity

data collection - Kitzes & Schricker, 2019; for birds within appropriate size ranges - McKinnon

& Love, 2018; and online citizen science efforts - Sullivan et al., 2009, 2014). They improve our

understanding of avian ecology by identifying previously unknown species-habitat relationships

(Jirinec et al., 2016), migratory routes (Jahn et al., 2016, 2021; Stanley et al., 2012, 2015),

foraging areas and behaviors (including accessing microhabitat refuges - Wolfson et al., 2020),

and wintering distributions (Renfrew et al., 2013). These advances facilitate discovery of species

responses to global change by revealing locations outside of direct observation. Aside from

emerging technologies, simpler bird monitoring approaches implemented in the past at large

scales have amassed rich longitudinal data (e.g., breeding bird surveys and regional atlases), and

band recoveries across the globe continue to supply information on bird population trends and

distributions (MacPherson et al., in prep).

A major challenge for predictive models is rarity or sparseness of point occurrence data.

Even with many potential sources of occurrence data, knowledge of bird distributions still varies

greatly by species, habitat, and geography. For example, rare, nocturnal, and/or secretive species

and those in remote areas are difficult to detect and model accurately (MacKenzie et al., 2005;

Stralberg et al., 2015). Rarity is a common characteristic of species in diverse tropical

communities and remains a major challenge for constructing species distribution and

environmental niche models for tropical species. Still, there are approaches to modelling rare

species, for example, using Bayesian network analysis (Hamilton et al., 2015) or "ensemble

small models" (ESMs; Breiner et al., 2015; see "Analytical Methods" section, below).

Decisions about which type of occurrence data is best for predicting distributions under

global change varies based on the trade-off between addressing processes affecting the core

distribution versus a more holistic understanding that includes unique responses at range

boundaries, during different life history activities, or between different sex and age classes.

Understanding of distributions could be improved by increasing the number of studies examining

range edges, and at finer temporal resolutions; "snapshots" in time do not provide a complete

picture of spatial resource requirements for individuals or populations. Passive acoustic

monitoring and long-term studies remain promising approaches for capturing relative abundance

data to estimate population dynamics with respect to global change (Pérez-Granados et al., 2019;

Pérez-Granados & Schuchmann, 2020; Sugai et al., 2020). Trait databases are gaining

momentum as sources of functional diversity to test hypotheses under the Hutchinsonian niche

concept (Gallagher et al., 2020; Leclerc et al., 2020; Matuoka et al., 2020). In particular,

exploring how functional roles of species change across abiotic gradients can dramatically

improve our understanding of abiotic tolerances to more accurately predict range dynamics. For

example, bill size has been shown to be important for thermoregulation in birds (e.g., Danner &

Greenberg, 2015) and a study of functional trait structure along a tropical elevational gradient in

Malaysian Borneo linked larger bills in low elevation communities with thermal tolerance

(Boyce et al., 2019). Current shortfalls in avian occurrence data (summarized by Lees et al.,

2020) require an integrated approach to predicting bird distributions with global change.

Although most efforts to collect occurrence data are from small spatial areas (e.g., from

individual field research sites) and coarse resolutions (Table 1), even incomplete general natural

history knowledge gained across spatial and temporal resolutions remains key to identifying

important explanatory variables responsible for supporting site occupancy now and into the

future.

Table 1. Geospatial context of common occurrence data sources for birds and explanatory

variables. Most occurrence data and explanatory variable data layers are of coarse spatial and

temporal resolution, making them useful for testing hypotheses under the Grinellian class of

niche. Occurrence data sources with fine spatial and temporal resolution are not as common and

require higher economic investment for acquiring location data (i.e., GPS tags) or physically

tracking individual birds (i.e., radio-tracking) but could be important for testing hypotheses

under the Eltonian class of niche. For a more detailed assessment of temporal categories for

avian occurrence data see MacPherson et al. (in prep).

Geospatial data qualities	Examples of occurrence data sources	Examples of explanatory variables
Large spatial area, coarse spatial and/or temporal resolution	•Remote sensing assets (satellite imagery, aerial imagery etc.) •Automated radiotelemetry •Geolocators •eBird •Stable hydrogen isotopes •Genetic data	•Long-term averages in stable climates (e.g., WorldClim) •Land cover (e.g., MODIS, GlobCover, SPOT) •Phenology and vegetation indices (e.g., MODIS, Landsat) •Climate record (e.g., NCDC, ClimateNA, ClimateSA) •Atmospheric weather reanalyses (e.g., ERA, GPM, TRMM, NCEP)
Small spatial area, coarse spatial and/or temporal resolution	•Citizen science (e.g., Breeding Bird Surveys, Christmas Bird Counts) •Natural history collections •Field notes •Point count surveys •Passive acoustic monitoring •Long-term datasets •Bird banding data	
Large spatial area, fine spatial and/or temporal resolution	•GPS tags	•Digital elevation models (e.g., ASTER) •Vegetation indices (e.g., SPOT)
Small spatial area, fine spatial and/or temporal resolution	•VHF radio-tags	•Personal weather data loggers •Weather station data

SELECTING APPROPRIATE EXPLANATORY VARIABLES

Predicting distributions into the future relies on accurate assessments of current life history

requirements. In this section, we synthesize the spatial and temporal resolution of explanatory

variables commonly used to match distributions with habitat and resource needs. Producing

sound predictive models requires suitable proxies for resource or habitat needs that are

ecologically relevant (e.g., Burns et al., 2020; Randin et al., 2020). To scale up to entire

distributions, there must be an underlying knowledge of a species' life history that effectively

links the lives of individuals to the explanatory variable (or proxy variable) associated with site

occupancy. Predicting distributions into a future where the distributions of food webs will shift

requires a breadth in knowledge of life history information to predict the probability of species

ranges shifting individually or with their entire biotic community. This task is complicated by the

potentially differing needs for each life history stage or demographic, which is often the case for

birds (e.g., habitat requirements may differ between fledglings and adults). While the main

considerations are the spatial and temporal resolutions of explanatory variables, biotic

interactions can also play a critical role, and may need to be considered on a species-specific

basis.

SPATIAL RESOLUTION OF EXPLANATORY VARIABLES

Selecting the correct spatial resolution of explanatory variables is key to producing the most

accurate assessments of range shifts under global change. Passively remotely sensed spatial

products (e.g., NDVI and BIOCLIM variables) can be proxies for features of the environment

being used by individuals. Active remote sensing systems, for example light detection and

ranging (LiDAR) has also been used to predict occupancy by light-demanding versus shade-

tolerant plant species (Wüest et al., 2020). LiDAR can also offer direct measurements of

structural features of habitat (e.g., vertical distribution of forest canopy elements and foliage

density) useful in predicting species occupancy (e.g., Goetz et al., 2010). Remotely sensed

variables are thought to reflect a contingency of resources assuming the data product conveys

mutual information on a shared spatiotemporal scale with the amount of food resources (Riotte-

Lambert & Matthiopoulos, 2020). There are some satisfying interpretations of the ecological

relevance of these remotely sensed products (see Bridge et al., 2016 (EVI); Renfrew et al., 2013

(NDVI); Tøttrup et al., 2012 (drought)), although the typical purpose of using these large scale

variables is for spatial accuracy and not causal relationships, in that remotely sensed variables

generally serve as proxies for features of the environment used by individuals.

Nevertheless, many remotely sensed abiotic products do intend to reflect a direct habitat

resource or type. Habitat characteristics (i.e., land cover) are most often identified with remotely

sensed data, which depending on the sensor, can have a variety of resolutions. Medium-high

spatial resolutions ($\leq$ 30m, e.g., Landsat, Sentinel) are useful to pair with more precise

occurrence data (e.g., GPS tags, point counts; Shirley et al., 2013), whereas low spatial

resolution (e.g., MODIS) can be useful for occurrence data such as geolocators or weather radar

(Heim et al., 2020). The temporal resolution of remotely sensed data is a limitation, especially

when modelling the dynamic nature of migratory birds and habitat changes (MacPherson et al.,

2018; Roslin et al., 2021). The limitations in remotely sensed spatio-temporal resolution should

not be a deterrent in SDMs, however, because the information gained is not possible with other

data sources (He et al., 2015; Randin et al., 2020). Still, applied interpretations when spatial

and/or temporal resolutions do not match should be made with caution.

TEMPORAL RESOLUTION OF EXPLANATORY VARIABLES

The temporal predictability of resources is important for predicting future distributions because

annual life history strategies are dependent upon correctly timing life cycle events with required

resources. For example, it is widely held that birds breeding at high latitudes must correctly time

the hatching of young with annual insect emergence to maximize fledging success, and early

arrival to breeding grounds may enhance reproductive success (Alerstam, 2011; Kokko et al.,

2006; Nilsson et al., 2013; Smith & Moore, 2005). Predictive distribution models are expected to

be the most accurate when species-habitat correlations are assessed using occurrence and

environmental data gathered from the same time-period. However, the temporal resolution of

occurrence data is much finer than that at which environmental data is typically collected; this

can create a mismatch that limits the questions that can be answered using current approaches.

The habitat requirements for fulfilling life history needs may vary depending upon the particular

life history event (i.e., mate acquisition, rearing young, molting, migrating, etc.) and the species

life history strategy. Life history strategies vary in birds from those that have evolved to rely on

consistent resource availability (e.g., in the case of dietary or habitat specialists that cannot live

outside of narrow environmental conditions), to periodic resource abundance (e.g., seasonal,

annual, inter-annual weather patterns that drive migratory or irruptive population movements), or

unreliable resource availability (e.g., in nomadic taxa), and it is important to match locality data

in time to environmental data for accurate species-habitat assessment.

The temporal resolutions for widely used climate data varies, with trade-offs between

temporal and spatial resolution in addition to the state of products limiting the types of

distribution questions that can be informed by them (Table 1). Here, we demonstrate this point

by explaining temperature and rainfall data sources because these are the two most widely used

and generally important environmental factors in species distribution modelling (Bradie &

Leung, 2017). The finest spatial and temporal resolution of environmental data is typically from

weather station data from the nearest station to the research site (e.g., from

https://www.wunderground.com/history) or by collecting weather data alongside occurrence data

using data loggers available from various manufacturers (e.g., Meter, Truebner). The ERA5-

Land dataset provides fine scale (hourly) temperature and precipitation data, but at a 9 km spatial

resolution from January 1981 to present

(https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview). National

Centers for Environmental Prediction (NCEP) provides global temperature and precipitation data

(four-times daily) data at a 0.25 x 0.25 degree grid resolution from 1948-present

(https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.surface.html). WorldClim is a commonly

used source of environmental data and the current version (2.1) provides global monthly values

of min and max temperature, and precipitation averaged from 1970-2000 (Fick & Hijmans,

2017). Historical monthly min and max temperature, and total precipitation data used for

hindcasting are averaged by decade from 1960-2018 (Fick & Hijmans, 2017; Harris et al., 2014).

BioClim data are 19 quarterly estimates derived from WorldClim monthly temperature and

rainfall values that are thought to be more biologically meaningful than raw temperature or

rainfall data (Busby, 1991). Bioclimatic variables have been averaged for the years 1970-2000 to

generate monthly estimates (Fick & Hijmans, 2017). Despite being powerful tools for assessing

macroclimate associations, caution should be used when using the most accessible climate

products to assess the probability of future range locations.

To build robust predictive models, it is necessary to match the spatial and temporal

resolution of the occurrence data with that of the habitat characteristics. The temporal resolution

of remotely sensed data is a limitation, especially when modelling the dynamic nature of

migratory birds where dynamic seasonal habitat changes drive short-term habitat quality (e.g.,

seasonally flooded mudflat habitat for migrating shorebirds; Twedt, 2013). Google Earth Engine

(a java script based platform where one can write their own code to integrate many satellite

products and independently calculate indices such as NDVI) is a flexible tool that can facilitate

identifying the correct temporal resolution of land cover data to robustly test explanatory power

of land cover variables. Examples of data sets in Google Earth Engine include vegetation indices

(Landsat, ~2 weeks temporal resolution), products for creating land cover classifications (SPOT,

~1 month), and land cover datasets (e.g., GlobCover, Cropland Data Layers, National Land

Cover Database; Table 1).

BIOTIC FACTORS

Modeling future range shifts in response to global change rarely considers biotic factors. This

omission is the main reason why an assumption of SDMs - that the species is at equilibrium with

their environment - is rarely, if ever, met (Pearson & Dawson, 2003). If this assumption were

true, a species' realized and fundamental niche would completely overlap as long as the species

could access all available suitable niche space (Figure 1). Consideration of biotic factors has the

potential to significantly enrich the field of predictive distribution modelling with more accurate

forecasts, and here we synthesize leading research contributing to this end at both the

intraspecific and interspecific level.

There remains a gap in statistical approaches that account for biotic interactions in

forecasting species distributions. It is thought that biotic interactions vary along abiotic gradients

such that biotic interactions can either enhance or reduce predicted ranges (Louthan et al., 2015).

However, because capturing this information requires both high spatial and high temporal

accuracy across all abiotic scenarios within the realized niche, there are few examples to draw

from. One promising approach to including biotic interactions in predictive spatial modelling is

the development of causal models that estimate the influence of interspecific competition using

co-occurrence data (e.g., Staniczenko et al., 2017). The scope and strength of biotic factors may

be correlated with abiotic pressures (Louthan et al., 2015), and may differ depending upon which

range edge is being considered. For example, much work has been done in the bird literature to

improve our understanding of species-habitat relationships through the study of hybrid zones,

which typically occupy only one part of a species range (e.g., Taylor et al., 2014). Quantification

of biotic factors, like the ones described here, requires targeted species-specific research to test

hypotheses within the Eltonian and Hutchinsonian classes of niche. This research is mainly done

within small spatial extents and without consideration of temporal or spatial resolution (e.g., co-

occurrence records from different time periods or datasets with different spatial resolutions -

Atauchi et al., 2018; Palacio & Girini, 2018; see also König et al., 2021), but some research has

been done at large spatial extents and coarse resolutions (e.g., Acorn Woodpecker occurrence

with Colombian Oaks in the Northern Andes - Freeman & Mason, 2015).

Modelling future range shifts in response to global change must consider the role of

intraspecific biotic factors that shape current distributions. Intraspecific competition can drive

density-dependent range shifts in migratory species such that interannual selection of wintering

locations are dependent on group size and food availability (Corre et al., 2020). Dispersal

limitation can contribute to inadequacies in recruitment to new or population sink sites, limiting

population recovery or range shifts despite available habitat (Palma et al., 2020). The ability to

update behaviors when circumstances change (behavioral flexibility) may be key for driving

range shifts or expansions under global change (Blaisdell et al., 2020). Further, bird species

capable of behavioral innovation (a.k.a. plasticity) often have lower risk of extinction, as they are

better able to adapt to changing ecosystems and habitat destruction (Ducatez et al., 2020; Reed et

al., 1999). Additionally, metapopulation dynamics can affect the introgression of adaptive traits

such that maladaptive traits restrict range shifts under global change (e.g., Garcia-R & Matzke,

2021; Lavretsky et al., 2020). Each of these examples are implicitly expected to vary across

abiotic gradients, highlighting the importance of assessing the probability of site occupancy

beyond the core of a species distribution.

Perhaps more difficult is estimating the strength of interspecific biotic factors in shaping

distributions because this requires a breadth of natural history knowledge beyond that of the

species in question. However, interspecific biotic factors are thought to have strong influences on

the probability of site occupancy. Species may be excluded through predator/prey dynamics

where patches of habitat occupied by a species' predator may be less likely to be occupied by the

species being studied (Léandri-Breton & Bêty, 2020). Interspecific competition can also exclude

species and is often assessed using co-occurrence data (Elsen et al., 2017; Freeman et al., 2016,

2019; Jankowski et al., 2010). Commensalisms, where one organism benefits from the presence

of another organism and the other unaffected, can also facilitate site occupancy (Aitken &

Martin, 2007). Including data on the presence of predators, competitors, and commensalisms in

SDMs can be a useful addition to improving models (Jankowski et al., 2013). Alternatively, the

spatial overlap with other taxa may have little influence if the focal species exhibits behavioral

avoidance (e.g., via interference competition - Jaeger, 1970; or predator-prey dynamics - Lukas

et al., 2021) or the spatio-temporal scope of data are not aligned.

The utility of SDMs to guide predictions of distributions under global change will

certainly be limited by the spatio-temporal resolution of environmental variables. Even when

researchers are mindful when selecting environmental data suited to answer questions about their

occurrence data (i.e., using ecologically relevant environmental and climate data across

biologically relevant space and timescales), a challenge remains in combining and contrasting

the Grinnellian class of niche (e.g., climate variables) against other niche classes that require

finer temporal and spatial data (e.g., density of interacting species). This problem can further

complicate predictive models if site occupancy is dependent on time-lags, as is the case in some

migratory species (Bridge et al., 2016), and is explored further in the following "Analytical

Methods" section.

ANALYTICAL METHODS

Predictive models of species distributions can be divided into correlative versus mechanistic

approaches (Figure 3). Correlative models elucidate relationships between species occurrence

and spatially explicit explanatory variables and can be viewed as hypothesis generating tools.

Data to parameterize correlative models are more readily available, making them the dominant

source of predictive distribution models. Mechanistic models are hypothesis-testing tools that

incorporate physiological tolerance to predict where species will be capable of persisting within

their physiological limits (Guisan & Zimmermann, 2000).

Figure 3. Analytical processes for species distribution models. Left: important terms used in

describing distribution model data. Right: Parallel workflows for example correlative versus

mechanistic predictive modelling.

Descriptions of range characteristics that are the outcome of correlative SDMs can help

us better understand species and their sensitivities to global change. MAXENT is one of the most

popular SDM approaches, widely used for its ability to use presence-only datasets where true

absence or non-detection data are typically hard to find (Elith & Leathwick, 2007; Merow et al.,

2013). MAXENT has been useful for modelling distributions of rare or cryptic species due to its

apparent high performance with small or incomplete datasets (Aubry et al., 2017; Pearson et al.,

2007; Phillips et al., 2006; Wisz et al., 2008; but see also Marini et al., 2009). Boosted regression

trees (BRTs) are one of many machine learning techniques well-suited to modelling complex

ecological data because they can handle different types of predictor variables, fit complex

nonlinear relationships, accommodate missing data and automatically handle interaction effects

between predictors (Elith et al., 2008; Graham et al., 2008). BRTs have been used to predict bird

distributions using occurrence data for waterfowl (Barker et al., 2014), shorebirds (Dalgarno et

al., 2017), seabirds (Oppel et al., 2012), and owls (Domahidi et al., 2019). Hierarchical Bayesian

models can integrate various data types (e.g., presence-absence, presence-only, count data) to

create reliable spatio-temporal distribution models (Hefley & Hooten, 2015). Using continuous

spatial predictor variables such as land cover, the abundance and distribution of species can be

predicted over a continuous latent surface (Chakraborty et al., 2010), which has been useful with

waterfowl aerial survey data (Herbert et al., 2018, 2021). Generalized linear models (GLMs) and

generalized additive models (GAMs) can be employed to understand distributions across

multiple environmental conditions and also conceptualize patterns in distributions of abundances

across environmental gradients (Augustin et al., 1996; Smith & Edwards, 2021). GAMs can be

used to estimate species' response curves with smoothing functions, which offer more flexibility

to capture complex shapes not possible with GLMs (e.g., Maggini et al., 2011). One drawback to

GAMs, however, is that evaluation of response shapes (e.g., optima, skewness) is done through

visual inspection rather than statistically. Huisman-Olff-Fresco (HOF) models are used to predict

occupancy along abiotic gradients and are particularly useful in evaluating alternative shapes of

species responses to gradients, for example, if changes in abundance do not exhibit the widely

assumed symmetric Gaussian functions, but instead show asymmetric, threshold-like changes in

occupancy over space. HOF response curves can be used to highlight distinct distribution

patterns such as species replacements, allowing inference of potential biotic interactions that are

otherwise difficult to measure (Jansen & Oksanen, 2013). The choice of predictive SDM

technique and quality of SDM product are based predominantly on the type of occurrence data

available and secondarily on spatiotemporal matching of explanatory variables.

Mechanistic approaches to SDMs model physico-chemical processes that drive species'

presence. More specifically, biophysical models predict areas where physico-chemical processes

meet life history needs (Kearney, 2006; Michael Kearney & Porter, 2009). For example, a

biophysical model for the endangered Night Parrot (*Pezoporus occidentalis*) shows how air

temperature increases of 3 °C would lead to lethal levels of dehydration (Kearney et al., 2016).

Bayesian networks (BNs) are another type of mechanistic model that go beyond species-habitat

correlations by also considering processes that influence occupancy across space and time (i.e.,

site access and selection - Jones, 2001). Originally employed to model human judgement (Pearl,

1985), BNs have only recently been adapted for predictive SDMs (MacPherson et al., 2018;

Staniczenko et al., 2017). Mechanistic models are informed by causal relationships based on

empirical research or expert knowledge, the latter of which are referred to as belief networks

(Drew & Collazo, 2014; MacPherson et al., 2018). Mechanistic models enable the identification

of the most important variables driving distribution patterns through mapping the fundamental

niche, which is helpful to inform conservation and management decisions when circumstances

change.

Conservation plans rely on SDMs for future species distributions, despite the small, but

growing, body of literature on ways to incorporate climate change into conservation planning

(Jones & Cheung, 2015; Loyola et al., 2013; Nakao et al., 2013; Terribile et al., 2012). Using

correlative predictive models is likely to overestimate range shifts and extinction risk due to the

violations of common assumptions in correlative distribution modeling. For example, SDMs tend

to violate the assumption that the individual species is currently at equilibrium within its

environment (Early & Sax, 2014; Sax et al., 2013) and do not take into account species

interactions (Pearson & Dawson, 2003). Despite these criticisms, SDMs remain the "best"

widely accessible approach currently in use for identifying potential future habitat-area (Tingley

et al., 2014).

Ensemble modeling strategies, in which the predictions of multiple approaches are

combined or used simultaneously, are often suggested to better encompass the range of

uncertainty in prediction (Araújo & New, 2007). This method accounts for the fact that model

choice is often the greatest source of quantifiable uncertainty in species distribution modeling,

and reduces sources of bias from the use of a single algorithm (Dormann et al., 2008; Jarnevich

et al., 2017). Implementing ensembles of five or more of the algorithms and approaches

described above has been greatly facilitated by software packages like BIOMOD (Thuiller et al.,

2009) and VisTrails SAHM (Morisette et al., 2013), which bring methods together into a single

analytical environment. Ensembles can be used to assess a range of potential projected outcomes,

forming a "bounding box" or "consensus" across algorithm predictions (Araújo & New, 2007).

Analysis with model ensembles typically involves scaling and averaging model outputs and,

often weighting these by some measurement of model performance such as "area under the

curve" score.

The ongoing "big data revolution" in many fields including ornithology (La Sorte et al.,

2018) has increased use of artificial intelligence for data handling and analysis (Xia et al., 2020),

especially in correlative distribution modeling. Machine learning algorithms like MAXENT

(Merow et al., 2013), random forests (Mi et al., 2017), neural networks (e.g., Manel et al., 1999),

deep learning (Benkendorf & Hawkins, 2020) and boosted regression trees (Elith et al., 2006) are

now commonly applied SDMs, and are valued for prediction and interpolation. They are

excellent tools for insight into potential future shifts under global change (Elith, 2017).

DISCUSSION

Predicting where species will live in a warmer future is a central goal of modern ecology. This

effort has recast the basic research question of what limits distribution of species – why a species

lives *here* but not *there* – as an applied question. Species are already on the move in response to

recent decades of warming (and, depending on the region, predicted trends of reduced or

increased rainfall; Figure 2), supporting long-held theory that climate limits species' geographic

ranges (Tingley et al., 2009). However, the match between species' distributions and climatic

conditions is often weak; species vary considerably in their observed responses to recent climate

change. These observations indicate that we are still in the early stages of ecological forecasting.

In this research review, we have highlighted the theory underpinning how climate is related to

species' distributions and focused especially on the types of data sources and analytic approaches

that are useful for predictive modeling of species' geographic responses to ongoing climate

change.

Here we briefly highlight three motivating questions to guide future research; the answers

to these questions are particularly likely to generate useful insights.

1. Is dispersal a rate-limiting step in range expansions?

The proximate driver of range shifts is dispersal; ranges expand when individuals move beyond

the existing range limit. Dispersal constraints are thus one obvious explanation for cases where

species' observed range shifts are failing to track changing climate. Birds have incredible

capacities for dispersal (e.g., Slager, 2020), making it tempting to gloss over a possible role for

dispersal constraints in avian range shifts, including unexplored topics such as site fidelity.

However, tropical species often show extreme limitations on dispersal, suggesting that longer-

distance range shifts (e.g., across latitudes) may be slow, particularly when landscapes are

patchy, or when species show high specialization to their associated habitats. In the temperate

zone, even some long-distance migrants may show incredible site fidelity to breeding (and

wintering) territories (Winger et al., 2019), which suggests the possibility that rapid range shifts

may be difficult to achieve. Lastly, dispersal may be possible but may introduce a new trade-off

with other components of the annual life cycle, such as migration. For example, the ability of

boreal breeding birds to take advantage of newly available forest habitat in the Arctic may be

constrained by a trade-off with ever-increasing migration distances.

2. Are species particularly rare at their range limits?

Ecological theory assumes that species are most abundant at the center or core of their range and

become progressively rarer towards their range limits (the "abundant-center hypothesis; Brown,

1984). However, empirical patterns of abundance often fail to fit these expectations. Whether the

abundant-center hypothesis generally holds or not is now a matter of debate (e.g., Dallas et al.,

2020), and the answer may have consequences for rates of species' range shifts. For example,

low abundances at range edges could lead to slower rates of range shifts than if abundance

distributions are more uniform across a core-to-limit transect through a species' geographic

range. This could be due to simple numeric effects – range shifts are easier when there are more

individuals at the range edge to start with – or to genetic effects. That is, low abundances (and

hence lower genetic diversity) could reduce local rates of adaptation at range edges, with gene

flow from the abundant range center swamping any effect of local adaptation at the range edge

that would facilitate range expansion when conditions change (Haldane, 1956; but see Kottler et

al., 2021). Studies on invasive species have the potential to improve our understanding of the

mechanisms that promote expansion from original range boundaries (e.g., the tolerance niche of

the Asian Openbill Bird - Lei & Liu, 2021; and climate matching in the European Bee-eater -

Stiels et al., 2021).

3. How does land use change interact with climate change?

Like most of the range shift literature, we have focused in this review on predicting species'

responses to changes in climate per se. But species are shifting (or failing to shift) their ranges in

landscapes that, more often than not, are dominated by human activities. Studies that

simultaneously incorporate both land use and climate change as drivers of distributional change

are few in number but hold great promise for several reasons. First, land use change is an

obvious driver of species' distributions; many species simply do not persist in landscapes with

extreme levels of human control. Second, appropriate habitat for most species in human-

dominated landscapes consists of habitat patches of varying sizes with varying levels of

connectivity (e.g., Neilan et al., 2019). This fact elevates the role of dispersal in determining

whether patches (and, scaled-up to a larger geographic scale, regions) are occupied by a given

species. Third, habitat change directly affects local-scale climate. Fragmented forests, for

example, average substantially warmer and drier than primary forests (Kapos, 1989), and "urban

heat islands" alter local temperatures that are known to affect bird distributions (in migrants -

Bonnet-Lebrun et al., 2020; and residents - Latimer & Zuckerberg, 2021). Hence, land use

change can act as a multiplier of temperature effects on species.

The consequences of climate change for species are potentially severe, with widespread

predictions of extinctions. Some evidence suggests these dire predictions may be coming true.

For example, warming temperatures have led to local extinctions in mountaintop communities in

southeastern Peru as species shift their geographic ranges to track climate, a potential harbinger

of the possible extinction of high elevation tropical species (Feeley et al., 2012; Freeman et al.,

2018; Rehm & Feeley, 2016). Yet species may be more resilient than models assume; for

example, genetic data indicate that many species were able to persist through dramatic climate

fluctuations in the Pleistocene (Bocalini et al., 2021; Song et al., 2020; Wogan et al., 2020).

CONCLUSION

Humans have already made Earth hotter than it has been since before the Pleistocene (~ 2 million

years ago). This rapid change in Earth's climate has set species on the move and led to a plethora

of scientific research aimed at predicting where species will live in the coming decades as

warming continues. These predictions, though frequently made, are seldom tested. Without this

crucial step of model validation, it is impossible to assess the usefulness of predictive models.

Here, we have presented theory and highlighted a range of data sources and analytical

approaches that can be used to predict species' geographic responses to climate change. Ever-

greater computational power, combined with increasingly large datasets of species occurrence

and landscape covariates, permit the use of greater complexity in models. However, more

complex does not necessarily equate to better. The litmus test for any predictive model is how

well it predicts reality. To this end, we advocate for an increasing focus on collecting empirical

data that matches the spatio-temporal resolution of occurrence data with environmental variables,

tests hypotheses beyond the Grinnellian class of niche, and directly evaluates model predictions

(e.g., using mechanistic SDMs or resurveys). For example, the long-term predictions from

models whose predictions are not met over the short-term are unlikely to be helpful. Determining

species' resiliency will depend on accurate estimates of the fundamental niche including which

attributes make some species more vulnerable than others, which abiotic gradients are the most

important to consider for the promotion of population persistence, and including biotic

interactions in predictive, validated models. The ongoing pursuit of meaningful predictive

distribution models under land use and climate change will be of great utility.

ACKNOWLEDGEMENTS

The authors declare that we have no sources of conflict of interest affecting the objectivity of the
presented topic. This manuscript is the result of a synthesis on how ornithologists predict bird
distributions under global change that was presented in a series of talks at the North American
Ornithological Congress, 2020. We sincerely thank the organizers of this conference for their
support throughout the COVID-19 pandemic. Thank you also to the participants of the Predicting
Bird Distributions Under Global Change symposium, including William Lewis. This research
was funded in part by a U.S. Geological Survey Northwest Climate Adaptation Science Center
award G17AC000218 to C.B. van Rees. The manuscript was approved by the Director of the

Louisiana State University Agricultural Center as manuscript number 2021-XXX-XXXX. The
authors would also like to thank (in alphabetical order by last name) Nicholas Mason, Subir
Shakya, Philip Stouffer, David Vander Pluym, and Brenna Wells who provided important
discussion of the topics presented in this manuscript during its development.
716

717 LITERATURE CITED

Aitken, K. E. H., & Martin, K. (2007). The importance of excavators in hole-nesting communities:

Availability and use of natural tree holes in old mixed forests of western Canada. *Journal of*

Ornithology, *148*(2), 425–434. https://doi.org/10.1007/s10336-007-0166-9

Alerstam, T. (2011). Optimal bird migration revisited. *Journal of Ornithology*, *152*(1), 5–23.

https://doi.org/10.1007/s10336-011-0694-1

Anderson, B. j, Akçakaya, H. r, Araújo, M. b, Fordham, D. a, Martinez-Meyer, E., Thuiller, W., & Brook,

B. w. (2009). Dynamics of range margins for metapopulations under climate change. *Proceedings*

of the Royal Society B: Biological Sciences, *276*(1661), 1415–1420.

https://doi.org/10.1098/rspb.2008.1681

Angilletta Jr, M. J., & Angilletta, M. J. (2009). *Thermal adaptation: A theoretical and empirical*

synthesis.

Angilletta, M. J., Sears, M. W., & Pringle, R. M. (2009). Spatial dynamics of nesting behavior: Lizards

shift microhabitats to construct nests with beneficial thermal properties. *Ecology*, *90*(10), 2933–

2939. https://doi.org/10.1890/08-2224.1

Araújo, M. B., & New, M. (2007). Ensemble forecasting of species distributions. *Trends in Ecology &*

Evolution, *22*(1), 42–47. https://doi.org/10.1016/j.tree.2006.09.010

Atauchi, P. J., Peterson, A. T., & Flanagan, J. (2018). Species distribution models for Peruvian plantcutter

improve with consideration of biotic interactions. *Journal of Avian Biology*, *49*(3), jav-01617.

https://doi.org/10.1111/jav.01617

Aubry, K. B., Raley, C. M., & McKelvey, K. S. (2017). The importance of data quality for generating

reliable distribution models for rare, elusive, and cryptic species. *PLOS ONE*, *12*(6), e0179152.

https://doi.org/10.1371/journal.pone.0179152

Augustin, N. H., Mugglestone, M. A., & Buckland, S. T. (1996). An autologistic model for the spatial

distribution of wildlife. *Journal of Applied Ecology*, *33*(2), 339–347.

https://doi.org/10.2307/2404755

Austin, M. (2005). Vegetation and environment: Discontinuous and continuous. *Vegetation Ecology*, 52–

84.

Austin, M. (2007). Species distribution models and ecological theory: A critical assessment and some

possible new approaches. *Ecological Modelling*, *200*(1), 1–19.

https://doi.org/10.1016/j.ecolmodel.2006.07.005

Barçante, L., Vale, M. M., & Alves, M. A. S. (2017). Altitudinal migration by birds: A review of the

literature and a comprehensive list of species. *Journal of Field Ornithology*, *88*(4), 321–335.

https://doi.org/10.1111/jofo.12234

Barker, N. K. S., Cumming, S. G., & Darveau, M. (2014). Models to predict the distribution and

abundance of breeding ducks in Canada. *Avian Conservation and Ecology*, *9*(2), art7.

https://doi.org/10.5751/ACE-00699-090207

Benkendorf, D. J., & Hawkins, C. P. (2020). Effects of sample size and network depth on a deep learning

approach to species distribution modeling. *Ecological Informatics*, *60*, 101137.

https://doi.org/10.1016/j.ecoinf.2020.101137

Blaisdell, A., Seitz, B., Rowney, C., Folsom, M., MacPherson, M., Deffner, D., & Logan, C. (2020). *Do*

the more flexible individuals rely more on causal cognition? Observation versus intervention in

causal inference in great-tailed grackles. PsyArXiv. https://doi.org/10.31234/osf.io/z4p6s

Bocalini, F., Bolívar-Leguizamón, S. D., Silveira, L. F., & Bravo, G. A. (2021). Comparative

phylogeographic and demographic analyses reveal a congruent pattern of sister relationships

between bird populations of the northern and south-central Atlantic Forest. *Molecular*

Phylogenetics and Evolution, *154*, 106973. https://doi.org/10.1016/j.ympev.2020.106973

Bonnet-Lebrun, A.-S., Manica, A., & Rodrigues, A. S. L. (2020). Effects of urbanization on bird

migration. *Biological Conservation*, *244*, 108423. https://doi.org/10.1016/j.biocon.2020.108423

Borowicz, A., McDowall, P., Youngflesh, C., Sayre-McCord, T., Clucas, G., Herman, R., Forrest, S.,
767    Rider, M., Schwaller, M., Hart, T., Jenouvrier, S., Polito, M. J., Singh, H., & Lynch, H. J. (2018).
768    Multi-modal survey of Adélie penguin mega-colonies reveals the Danger Islands as a seabird
769    hotspot. *Scientific Reports*, *8*(1), 3926. https://doi.org/10.1038/s41598-018-22313-w

770 Boyce, A. J., Shakya, S., Sheldon, F. H., Moyle, R. G., & Martin, T. E. (2019). Biotic interactions are the
771    dominant drivers of phylogenetic and functional structure in bird communities along a tropical
772    elevational gradient. *The Auk*, *136*(ukz054). https://doi.org/10.1093/auk/ukz054

773 Bradie, J., & Leung, B. (2017). A quantitative synthesis of the importance of variables used in MaxEnt
774    species distribution models. *Journal of Biogeography*, *44*(6), 1344–1361.
775    https://doi.org/10.1111/jbi.12894

776 Brawn, J. D., Benson, T. J., Stager, M., Sly, N. D., & Tarwater, C. E. (2017). Impacts of changing rainfall
777    regime on the demography of tropical birds. *Nature Climate Change*, *7*(2), 133–136.
778    https://doi.org/10.1038/nclimate3183

779 Breiner, F. T., Guisan, A., Bergamini, A., & Nobis, M. P. (2015). Overcoming limitations of modelling
780    rare species by using ensembles of small models. *Methods in Ecology and Evolution*, *6*(10),
781    1210–1218. https://doi.org/10.1111/2041-210X.12403

782 Bridge, E. S., Ross, J. D., Contina, A. J., & Kelly, J. F. (2016). Do molt-migrant songbirds optimize
783    migration routes based on primary productivity? *Behavioral Ecology*, *27*(3), 784–792.
784    https://doi.org/10.1093/beheco/arv199

785 Brodie, J. F., Strimas-Mackey, M., Mohd-Azlan, J., Granados, A., Bernard, H., Giordano, A. J., &
786    Helmy, O. E. (2017). Lowland biotic attrition revisited: Body size and variation among climate
787    change 'winners' and 'losers.' *Proceedings of the Royal Society B: Biological Sciences*,
788    *284*(1847), 20162335. https://doi.org/10.1098/rspb.2016.2335

789 Brown, J. H. (1984). On the Relationship between Abundance and Distribution of Species. *The American
790    Naturalist*, *124*(2), 255–279. https://doi.org/10.1086/284267

Burns, P., Clark, M., Salas, L., Hancock, S., Leland, D., Jantz, P., Dubayah, R., & Goetz, S. J. (2020).

Incorporating canopy structure from simulated GEDI lidar into bird species distribution models.

Environmental Research Letters, *15*(9), 095002. https://doi.org/10.1088/1748-9326/ab80ee

Busby, J. R. (1991). BIOCLIM – A bioclimate analysis and prediction system. In C. R. Margules & M. P.

Austin (Eds.), *Nature Conservation: Cost Effective Biological Surveys and Data Analysis* (pp.

64–68). CSIRO.

Capurucho, J. M. G., Ashley, M. V., Tsuru, B. R., Cooper, J. C., & Bates, J. M. (2020). Dispersal ability

correlates with range size in Amazonian habitat-restricted birds. *Proceedings of the Royal Society*

B: Biological Sciences, *287*(1939), 20201450. https://doi.org/10.1098/rspb.2020.1450

Caruso, C. M., Martin, R. A., Sletvold, N., Morrissey, M. B., Wade, M. J., Augustine, K. E., Carlson, S.

M., MacColl, A. D. C., Siepielski, A. M., & Kingsolver, J. G. (2017). What Are the

Environmental Determinants of Phenotypic Selection? A Meta-analysis of Experimental Studies.

The American Naturalist, *190*(3), 363–376. https://doi.org/10.1086/692760

Caughley, G. (1994). Directions in conservation biology. *Journal of Animal Ecology*, *63*(2), 215–244.

https://doi.org/10.2307/5542

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M. (2015).

Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Science*

Advances, *1*(5), e1400253. https://doi.org/10.1126/sciadv.1400253

Chakraborty, A., Gelfand, A. E., Wilson, A. M., Latimer, A. M., & John A. Silander, J. (2010). Modeling

large scale species abundance with latent spatial processes. *The Annals of Applied Statistics*, *4*(3),

1403–1429. https://doi.org/10.1214/10-AOAS335

Claramunt, S., Derryberry, E. P., Remsen, J. V., & Brumfield, R. T. (2012). High dispersal ability inhibits

speciation in a continental radiation of passerine birds. *Proceedings of the Royal Society B:*

Biological Sciences, *279*(1733), 1567–1574. https://doi.org/10.1098/rspb.2011.1922

Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C., & Longino, J. T. (2008). Global warming,

816   elevational range shifts, and lowland biotic attrition in the wet tropics. *Science*, *322*(5899), 258–

817   261. https://doi.org/10.1126/science.1162547

818   Corre, M. L., Dussault, C., & Côté, S. D. (2020). Where to spend the winter? The role of intraspecific

819   competition and climate in determining the selection of wintering areas by migratory caribou.

820   *Oikos*, *129*(4), 512–525. https://doi.org/10.1111/oik.06668

821   Cowles, R. B., & Bogert, C. M. (1944). A preliminary study of the thermal requirements of desert

822   reptiles. *Bulletin of the American Museum of Natural History*, *83*(5), 261–296.

823   Dalgarno, S., Mersey, J. E., Gedalof, Z., & Lemon, M. (2017). Species-environment associations and

824   predicted distribution of Black Oystercatcher breeding pairs in Haida Gwaii, British Columbia,

825   Canada. *Avian Conservation and Ecology*, *12*(2), art9. https://doi.org/10.5751/ACE-01094-

826   120209

827   Dallas, T. A., Santini, L., Decker, R., & Hastings, A. (2020). Weighing the Evidence for the Abundant-

828   Center Hypothesis. *Biodiversity Informatics*, *15*(3), 81–91.

829   https://doi.org/10.17161/bi.v15i3.11989

830   Danner, R. M., & Greenberg, R. (2015). A critical season approach to Allen's rule: Bill size declines with

831   winter temperature in a cold temperate environment. *Journal of Biogeography*, *42*(1), 114–120.

832   https://doi.org/10.1111/jbi.12389

833   Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin,

834   P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of

835   *the National Academy of Sciences*, *105*(18), 6668–6672.

836   Domahidi, Z., Shonfield, J., Nielsen, S. E., Spence, J. R., & Bayne, E. M. (2019). Spatial distribution of

837   the Boreal Owl and Northern Saw-whet Owl in the boreal region of Alberta, Canada. *Avian

838   Conservation and Ecology*, *14*(2), art14. https://doi.org/10.5751/ACE-01445-140214

Dormann, C. F., Purschke, O., Márquez, J. R. G., Lautenbach, S., & Schröder, B. (2008). Components of

Uncertainty in Species Distribution Analysis: A Case Study of the Great Grey Shrike. *Ecology*,

89(12), 3371–3386. https://doi.org/10.1890/07-1772.1

Drew, A., & Collazo, J. (2014). *Bayesian networks as a framework to step-down and support Strategic*

Habitat Conservation of data-poor species: A case study with King Rail (Rallus elegans) in

Eastern North Carolina and Southeastern Virginia. US Fish and Wildlife Service.

https://doi.org/10.13140/2.1.4430.5920

Ducatez, S., Sol, D., Sayol, F., & Lefebvre, L. (2020). Behavioural plasticity is associated with reduced

extinction risk in birds. *Nature Ecology & Evolution*, *4*(6), 788–793.

https://doi.org/10.1038/s41559-020-1168-8

Dunn, P. O., & Møller, A. P. (2019). *Effects of climate change on birds*. Oxford University Press.

Dvorský, M., Macek, M., Kopecký, M., Wild, J., & Doležal, J. (2017). Niche asymmetry of vascular

plants increases with elevation. *Journal of Biogeography*, *44*(6), 1418–1425.

https://doi.org/10.1111/jbi.13001

Early, R., & Sax, D. F. (2014). Climatic niche shifts between species' native and naturalized ranges raise

concern for ecological forecasts during invasions and climate change. *Global Ecology and*

Biogeography, *23*(12), 1356–1365. https://doi.org/10.1111/geb.12208

Elith, J. (2017). Predicting Distributions of Invasive Species. In *Invasive Species: Risk Assessment and*

Management (pp. 93–129). Cambridge University Press.

Elith, J., H. Graham, C., P. Anderson, R., Dudík, M., Ferrier, S., Guisan, A., J. Hijmans, R., Huettmann,

F., R. Leathwick, J., Lehmann, A., Li, J., G. Lohmann, L., A. Loiselle, B., Manion, G., Moritz,

C., Nakamura, M., Nakazawa, Y., McC. M. Overton, J., Townsend Peterson, A., … E.

Zimmermann, N. (2006). Novel methods improve prediction of species' distributions from

occurrence data. *Ecography*, *29*(2), 129–151. https://doi.org/10.1111/j.2006.0906-7590.04596.x

Elith, J., & Leathwick, J. (2007). Predicting species distributions from museum and herbarium records

using multiresponse models fitted with multivariate adaptive regression splines. *Diversity and*

Distributions, *13*(3), 265–275. https://doi.org/10.1111/j.1472-4642.2007.00340.x

Elith, J., Leathwick, J. R., & Hastie, T. (2008). A working guide to boosted regression trees. *Journal of*

Animal Ecology, *77*(4), 802–813. https://doi.org/10.1111/j.1365-2656.2008.01390.x

Elsen, P. R., Tingley, M. W., Kalyanaraman, R., Ramesh, K., & Wilcove, D. S. (2017). The role of

competition, ecotones, and temperature in the elevational distribution of Himalayan birds.

Ecology, *98*(2), 337–348. https://doi.org/10.1002/ecy.1669

Elton, C. (1927). *Animal ecology*. Sidgwick & Jackson.

Faaborg, J., Holmes, R. T., Anders, A. D., Bildstein, K. L., Dugger, K. M., Gauthreaux, S. A., Heglund,

P., Hobson, K. A., Jahn, A. E., Johnson, D. H., Latta, S. C., Levey, D. J., Marra, P. P., Merkord,

C. L., Nol, E., Rothstein, S. I., Sherry, T. W., Sillett, T. S., Thompson, F. R., & Warnock, N.

(2010). Recent advances in understanding migration systems of New World land birds.

Ecological Monographs, *80*(1), 3–48. https://doi.org/10.1890/09-0395.1

Feeley, K. J., Rehm, E. M., & Machovina, B. (2012). perspective: The responses of tropical forest species

to global climate change: acclimate, adapt, migrate, or go extinct? *Frontiers of Biogeography*,

4(2). https://doi.org/10.21425/F5FBG12621

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Freeman, B. G., Freeman, A. M. C., & Hochachka, W. M. (2016). Asymmetric interspecific aggression in

New Guinean songbirds that replace one another along an elevational gradient. *Ibis*, *158*(4), 726–

737. https://doi.org/10.1111/ibi.12384

Freeman, B. G., & Mason, N. A. (2015). The geographic distribution of a tropical montane bird is limited

by a tree: Acorn Woodpeckers (Melanerpes formicivorus) and Colombian oaks (Quercus

humboldtii) in the Northern Andes. *PLOS ONE*, *10*(6), e0128675.

https://doi.org/10.1371/journal.pone.0128675

Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change causes

upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the*

National Academy of Sciences, *115*(47), 11982–11987. https://doi.org/10.1073/pnas.1804224115

Freeman, B. G., Tobias, J. A., & Schluter, D. (2019). Behavior influences range limits and patterns of

coexistence across an elevational gradient in tropical birds. *Ecography*, *42*(11), 1832–1840.

https://doi.org/10.1111/ecog.04606

Gallagher, R. V., Falster, D. S., Maitner, B. S., Salguero-Gómez, R., Vandvik, V., Pearse, W. D.,

Schneider, F. D., Kattge, J., Poelen, J. H., Madin, J. S., Ankenbrand, M. J., Penone, C., Feng, X.,

Adams, V. M., Alroy, J., Andrew, S. C., Balk, M. A., Bland, L. M., Boyle, B. L., … Enquist, B.

J. (2020). Open Science principles for accelerating trait-based science across the Tree of Life.

Nature Ecology & Evolution, *4*(3), 294–303. https://doi.org/10.1038/s41559-020-1109-6

Garcia-R, J. C., & Matzke, N. J. (2021). Trait-dependent dispersal in rails (Aves: Rallidae): Historical

biogeography of a cosmopolitan bird clade. *Molecular Phylogenetics and Evolution*, *159*,

107106. https://doi.org/10.1016/j.ympev.2021.107106

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body size: A

third universal response to warming? *Trends in Ecology & Evolution*, *26*(6), 285–291.

https://doi.org/10.1016/j.tree.2011.03.005

Goetz, S. J., Steinberg, D., Betts, M. G., Holmes, R. T., Doran, P. J., Dubayah, R., & Hofton, M. (2010).

Lidar remote sensing variables predict breeding habitat of a Neotropical migrant bird. *Ecology*,

91(6), 1569–1576. https://doi.org/10.1890/09-1670.1

Grady, J. M., Maitner, B. S., Winter, A. S., Kaschner, K., Tittensor, D. P., Record, S., Smith, F. A.,

Wilson, A. M., Dell, A. I., Zarnetske, P. L., Wearing, H. J., Alfaro, B., & Brown, J. H. (2019).

Metabolic asymmetry and the global diversity of marine predators. *Science*, *363*(6425).

https://doi.org/10.1126/science.aat4220

Graham, C. H., Elith, J., Hijmans, R. J., Guisan, A., Peterson, A. T., & Loiselle, B. A. (2008). The

influence of spatial errors in species occurrence data used in distribution models. *Journal of*

Applied Ecology, *45*(1), 239–247. https://doi.org/10.1111/j.1365-2664.2007.01408.x

Graham, R. W., Lundelius, E. L., Graham, M. A., Schroeder, E. K., Toomey, R. S., Anderson, E.,

Barnosky, A. D., Burns, J. A., Churcher, C. S., Grayson, D. K., Guthrie, R. D., Harington, C. R.,

Jefferson, G. T., Martin, L. D., McDonald, H. G., Morlan, R. E., Semken, H. A., Webb, S. D.,

Werdelin, L., & Wilson, M. C. (1996). Spatial Response of Mammals to Late Quaternary

Environmental Fluctuations. *Science*, *272*(5268), 1601–1606.

https://doi.org/10.1126/science.272.5268.1601

Grant, P. R., & Grant, B. R. (2020). How and Why Species Multiply. In *How and Why Species Multiply*.

Princeton University Press.

https://www.degruyter.com/document/doi/10.1515/9781400837946/html

Grant, P. R., & Weiner, J. (2017). *Ecology and Evolution of Darwin's Finches (Princeton Science*

Library Edition): Princeton Science Library Edition.

Guisan, A., & Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology. *Ecological*

Modelling, *135*(2), 147–186. https://doi.org/10.1016/S0304-3800(00)00354-9

Haldane, J. B. S. (1956). The relation between density regulation and natural selection. *Proceedings of the*

Royal Society of London. Series B - Biological Sciences, *145*(920), 306–308.

https://doi.org/10.1098/rspb.1956.0039

Hallman, T. A., & Robinson, W. D. (2020). Comparing multi- and single-scale species distribution and

abundance models built with the boosted regression tree algorithm. *Landscape Ecology*, *35*(5),

1161–1174. https://doi.org/10.1007/s10980-020-01007-7

Hamilton, S. H., Pollino, C. A., & Jakeman, A. J. (2015). Habitat suitability modelling of rare species

using Bayesian networks: Model evaluation under limited data. *Ecological Modelling*, *299*, 64–

78. https://doi.org/10.1016/j.ecolmodel.2014.12.004

Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of monthly

climatic observations – the CRU TS3.10 Dataset. *International Journal of Climatology*, *34*(3),

623–642. https://doi.org/10.1002/joc.3711

He, K. S., Bradley, B. A., Cord, A. F., Rocchini, D., Tuanmu, M.-N., Schmidtlein, S., Turner, W.,

Wegmann, M., & Pettorelli, N. (2015). Will remote sensing shape the next generation of species

distribution models? *Remote Sensing in Ecology and Conservation*, *1*(1), 4–18.

https://doi.org/10.1002/rse2.7

Hefley, T. J., & Hooten, M. B. (2015). On the existence of maximum likelihood estimates for presence-

only data. *Methods in Ecology and Evolution*, *6*(6), 648–655. https://doi.org/10.1111/2041-

210X.12340

Heim, W., Heim, R. J., Beermann, I., Burkovskiy, O. A., Gerasimov, Y., Ktitorov, P., Ozaki, K., Panov,

I., Sander, M. M., Sjöberg, S., Smirenski, S. M., Thomas, A., Tøttrup, A. P., Tiunov, I. M.,

Willemoes, M., Hölzel, N., Thorup, K., & Kamp, J. (2020). Using geolocator tracking data and

ringing archives to validate citizen-science based seasonal predictions of bird distribution in a

data-poor region. *Global Ecology and Conservation*, *24*, e01215.

https://doi.org/10.1016/j.gecco.2020.e01215

Helm, B., Van Doren, B. M., Hoffmann, D., & Hoffmann, U. (2019). Evolutionary response to climate

change in migratory Pied Flycatchers. *Current Biology*, *29*(21), 3714-3719.e4.

https://doi.org/10.1016/j.cub.2019.08.072

Herbert, J. A., Chakraborty, A., Naylor, L. W., Beatty, W. S., & Krementz, D. G. (2018). Effects of

landscape structure and temporal habitat dynamics on wintering mallard abundance. *Landscape*

Ecology, *33*(8), 1319–1334. https://doi.org/10.1007/s10980-018-0671-7

Herbert, J. A., Chakraborty, A., Naylor, L. W., & Krementz, D. G. (2021). Habitat associations of

wintering dabbling ducks in the Arkansas Mississippi Alluvial Valley: Implications for waterfowl

management beyond the mallard. *Wildlife Biology*, *2021*(1). https://doi.org/10.2981/wlb.00792

Holland, R. A., Wikelski, M., Kümmeth, F., & Bosque, C. (2009). The secret life of oilbirds: New

965   insights into the movement ecology of a unique avian frugivore. *PLOS ONE*, *4*(12), e8264.

966   https://doi.org/10.1371/journal.pone.0008264

967   Holt, R. D., Keitt, T. H., Lewis, M. A., Maurer, B. A., & Taper, M. L. (2005). Theoretical models of

968   species' borders: Single species approaches. *Oikos*, *108*(1), 18–27. https://doi.org/10.1111/j.0030-

969   1299.2005.13147.x

970   Hsiung, A. C., Boyle, W. A., Cooper, R. J., & Chandler, R. B. (2018). Altitudinal migration: Ecological

971   drivers, knowledge gaps, and conservation implications. *Biological Reviews*, *93*(4), 2049–2070.

972   https://doi.org/10.1111/brv.12435

973   Huey, R. B., Hertz, P. E., & Sinervo, B. (2003). Behavioral drive versus behavioral inertia in evolution: A

974   null model approach. *The American Naturalist*, *161*(3), 357–366. https://doi.org/10.1086/346135

975   Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., & Williams, S. E. (2012).

976   Predicting organismal vulnerability to climate warming: Roles of behaviour, physiology and

977   adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1596),

978   1665–1679. https://doi.org/10.1098/rstb.2012.0005

979   Jaeger, R. G. (1970). Potential Extinction Through Competition Between Two Species of Terrestrial

980   Salamanders. *Evolution*, *24*(3), 632–642. https://doi.org/10.2307/2406842

981   Jahn, A. E., Buechley, E., Bugoni, L., Cereghetti, J., Repenning, M., Marra, P. P., & Ryder, T. B. (2021).

982   Variable Seasonal Movement Dynamics Among Individual Snail Kites (Rostrhamus sociabilis) In

983   South America. *Journal of Raptor Research*. https://doi.org/10.3356/JRR-20-77

984   Jahn, A. E., Cueto, V. R., Fontana, C. S., Guaraldo, A. C., Levey, D. J., Marra, P. P., & Ryder, T. B.

985   (2020). Bird migration within the Neotropics. *The Auk*, *137*(ukaa033).

986   https://doi.org/10.1093/auk/ukaa033

987   Jahn, A. E., Cueto, V. R., Fox, J. W., Husak, M. S., Kim, D. H., Landoll, D. V., Ledezma, J. P., Lepage,

988   H. K., Levey, D. J., Murphy, M. T., & Renfrew, R. B. (2013). Migration timing and wintering

areas of three species of flycatchers (Tyrannus) breeding in the Great Plains of North America.

The Auk, *130*(2), 247–257. https://doi.org/10.1525/auk.2013.13010

Jahn, A. E., Levey, D. J., & Smith, K. G. (2004). Reflections across hemispheres: A system-wide

approach to New World bird migration. *The Auk*, *121*(4), 1005–1013.

https://doi.org/10.1093/auk/121.4.1005

Jahn, A. E., Seavy, N., Bejarano, V., Guzmán, M. B., Provinciato, I. C. C., Pizo, M., & MacPherson, M.

(2016). *Intra-tropical migration and wintering areas of Fork-tailed Flycatchers (Tyrannus*

savana) breeding in São Paulo, Brazil. https://doi.org/10.1007/BF03544339

Jankowski, J. E., Londoño, G. A., Robinson, S. K., & Chappell, M. A. (2013). Exploring the role of

physiology and biotic interactions in determining elevational ranges of tropical animals.

Ecography, *36*, 001–012.

Jankowski, J. E., Robinson, S. K., & Levey, D. J. (2010). Squeezed at the top: Interspecific aggression

may constrain elevational ranges in tropical birds. *Ecology*, *91*(7), 1877–1884.

https://doi.org/10.1890/09-2063.1

Jansen, F., & Oksanen, J. (2013). How to model species responses along ecological gradients – Huisman–

Olff–Fresco models revisited. *Journal of Vegetation Science*, *24*(6), 1108–1117.

https://doi.org/10.1111/jvs.12050

Janzen, D. H. (1967). Why mountain passes are higher in the tropics. *The American Naturalist*, *101*(919),

233–249.

Jarnevich, C. S., Talbert, M., Morisette, J., Aldridge, C., Brown, C. S., Kumar, S., Manier, D., Talbert, C.,

& Holcombe, T. (2017). Minimizing effects of methodological decisions on interpretation and

prediction in species distribution studies: An example with background selection. *Ecological*

Modelling, *363*, 48–56. https://doi.org/10.1016/j.ecolmodel.2017.08.017

Jiguet, F., Gadot, A.-S., Julliard, R., Newson, S. E., & Couvet, D. (2007). Climate envelope, life history

traits and the resilience of birds facing global change. *Global Change Biology*, *13*(8), 1672–1684.

https://doi.org/10.1111/j.1365-2486.2007.01386.x

Jirinec, V., Isdell, R. E., & Leu, M. (2016). Prey availability and habitat structure explain breeding space use of a migratory songbird. *The Condor*, *118*(2), 309–328. https://doi.org/10.1650/CONDOR-15-140.1

Jones, J. (2001). Habitat selection studies in avian ecology: A critical review. *The Auk*, *118*(2), 557–562. https://doi.org/10.1093/auk/118.2.557

Jones, M. C., & Cheung, W. W. L. (2015). Multi-model ensemble projections of climate change effects on global marine biodiversity. *ICES Journal of Marine Science*, *72*(3), 741–752. https://doi.org/10.1093/icesjms/fsu172

Kapos, V. (1989). Effects of isolation on the water status of forest patches in the Brazilian Amazon. *Journal of Tropical Ecology*, *5*(2), 173–185. https://doi.org/10.1017/S0266467400003448

Kearney, M. (2006). Habitat, environment and niche: What are we modelling? *Oikos*, *115*(1), 186–191. https://doi.org/10.1111/j.2006.0030-1299.14908.x

Kearney, M., & Porter, W. (2009). Mechanistic niche modelling: Combining physiological and spatial data to predict species' ranges. *Ecology Letters*, *12*(4), 334–350. https://doi.org/10.1111/j.1461-0248.2008.01277.x

Kearney, M. R., Porter, W. P., & Murphy, S. A. (2016). An estimate of the water budget for the endangered night parrot of Australia under recent and future climates. *Climate Change Responses*, *3*(1), 14. https://doi.org/10.1186/s40665-016-0027-y

Kitzes, J., & Schricker, L. (2019). The necessity, promise and challenge of automated biodiversity surveys. *Environmental Conservation*, *46*(4), 247–250. https://doi.org/10.1017/S0376892919000146

Kokko, H., Gunnarsson, T. G., Morrell, L. J., & Gill, J. A. (2006). Why do female migratory birds arrive later than males? *Journal of Animal Ecology*, *75*(6), 1293–1303. https://doi.org/10.1111/j.1365-2656.2006.01151.x

König, C., Wüest, R. O., Graham, C. H., Karger, D. N., Sattler, T., Zimmermann, N. E., & Zurell, D. (2021). Scale dependency of joint species distribution models challenges interpretation of biotic interactions. *Journal of Biogeography*, *0*, 1–11. https://doi.org/10.1111/jbi.14106

Kottler, E. J., Dickman, E. E., Sexton, J. P., Emery, N. C., & Franks, S. J. (2021). Draining the Swamping Hypothesis: Little Evidence that Gene Flow Reduces Fitness at Range Edges. *Trends in Ecology & Evolution*, *36*(6), 533–544. https://doi.org/10.1016/j.tree.2021.02.004

La Sorte, F. A., Lepczyk, C. A., Burnett, J. L., Hurlbert, A. H., Tingley, M. W., & Zuckerberg, B. (2018). Opportunities and challenges for big data ornithology. *The Condor*, *120*(2), 414–426. https://doi.org/10.1650/CONDOR-17-206.1

Latimer, C. E., & Zuckerberg, B. (2021). Habitat loss and thermal tolerances influence the sensitivity of resident bird populations to winter weather at regional scales. *Journal of Animal Ecology*, *90*(2), 317–329. https://doi.org/10.1111/1365-2656.13332

Lavretsky, P., McInerney, N. R., Mohl, J. E., Brown, J. I., James, H. F., McCracken, K. G., & Fleischer, R. C. (2020). Assessing changes in genomic divergence following a century of human-mediated secondary contact among wild and captive-bred ducks. *Molecular Ecology*, *29*(3), 578–595. https://doi.org/10.1111/mec.15343

Léandri-Breton, D.-J., & Bêty, J. (2020). Vulnerability to predation may affect species distribution: Plovers with broader arctic breeding range nest in safer habitat. *Scientific Reports*, *10*(1), 5032. https://doi.org/10.1038/s41598-020-61956-6

Leclerc, C., Villéger, S., Marino, C., & Bellard, C. (2020). Global changes threaten functional and taxonomic diversity of insular species worldwide. *Diversity and Distributions*, *26*(4), 402–414. https://doi.org/10.1111/ddi.13024

Lees, A. C., Rosenberg, K. V., Ruiz-Gutierrez, V., Marsden, S., Schulenberg, T. S., & Rodewald, A. D. (2020). A roadmap to identifying and filling shortfalls in Neotropical ornithology. *The Auk*, *137*(4), NA. https://doi.org/10.1093/auk/ukaa048

Lei, Y., & Liu, Q. (2021). Tolerance niche expansion and potential distribution prediction during Asian

openbill bird range expansion. *Ecology and Evolution*, *00*, 1–13.

https://doi.org/10.1002/ece3.7456

Londoño, G. A., Chappell, M. A., Castañeda, M. del R., Jankowski, J. E., & Robinson, S. K. (2015).

Basal metabolism in tropical birds: Latitude, altitude, and the 'pace of life.' *Functional Ecology*,

29(3), 338–346. https://doi.org/10.1111/1365-2435.12348

Londoño, G. A., Chappell, M. A., Jankowski, J. E., & Robinson, S. K. (2017). Do thermoregulatory costs

limit altitude distributions of Andean forest birds? *Functional Ecology*, *31*(1), 204–215.

https://doi.org/10.1111/1365-2435.12697

Louthan, A. M., Doak, D. F., & Angert, A. L. (2015). Where and when do species interactions set range

limits? *Trends in Ecology & Evolution*, *30*(12), 780–792.

https://doi.org/10.1016/j.tree.2015.09.011

Loyola, R. D., Lemes, P., Nabout, J. C., Trindade-Filho, J., Sagnori, M. D., Dobrovolski, R., & Diniz-

Filho, J. A. F. (2013). A straightforward conceptual approach for evaluating spatial conservation

priorities under climate change. *Biodiversity and Conservation*, *22*(2), 483–495.

https://doi.org/10.1007/s10531-012-0424-x

Lukas, J., Auer, F., Goldhammer, T., Krause, J., Romanczuk, P., Klamser, P., Arias-Rodriguez, L., &

Bierbach, D. (2021). *Diurnal Changes in Hypoxia Shape Predator-Prey Interaction in a Bird-

Fish System*. https://doi.org/10.14279/depositonce-11735

MacKenzie, D. I., Nichols, J. D., Sutton, N., Kawanishi, K., & Bailey, L. L. (2005). Improving inferences

in population studies of rare species that are detected imperfectly. *Ecology*, *86*(5), 1101–1113.

https://doi.org/10.1890/04-1060

MacPherson, M., Burgio, K., van Rees, C., DeSaix, M., Slager, D., Jirinec, V., Herman, R., Herbert, J.,

Shonfield, J., Freeman, B., & Jankowski, J. (In Prep). *Assessing the use of occurrence data in

predicting avian responses to climate change.*

MacPherson, M. P., Jahn, A. E., & Mason, N. A. (2021). Morphology of migration: Associations between

1090   wing shape, bill morphology, and migration in kingbirds (Tyrannus). *BioRxiv*,

1091   2021.04.06.438708. https://doi.org/10.1101/2021.04.06.438708

1092   MacPherson, M. P., Jahn, A. E., Murphy, M. T., Kim, D. H., Cueto, V. R., Tuero, D. T., & Hill, E. D.

1093   (2018). Follow the rain? Environmental drivers of Tyrannus migration across the New World.

1094   *The Auk*, *135*(4), 881–894. https://doi.org/10.1642/AUK-17-209.1

1095   MacPherson, M. P., Webb, E. B., Raedeke, A., Mengel, D., & Nelson, F. (2018). A review of Bayesian

1096   belief network models as decision-support tools for wetland conservation: Are water birds

1097   potential umbrella taxa? *Biological Conservation*, *226*, 215–223.

1098   https://doi.org/10.1016/j.biocon.2018.08.001

1099   Maggini, R., Lehmann, A., Kéry, M., Schmid, H., Beniston, M., Jenni, L., & Zbinden, N. (2011). Are

1100   Swiss birds tracking climate change?: Detecting elevational shifts using response curve shapes.

1101   *Ecological Modelling*, *222*(1), 21–32. https://doi.org/10.1016/j.ecolmodel.2010.09.010

1102   Manel, S., Dias, J. M., Buckton, S. T., & Ormerod, S. J. (1999). Alternative methods for predicting

1103   species distribution: An illustration with Himalayan river birds. *Journal of Applied Ecology*,

1104   *36*(5), 734–747. https://doi.org/10.1046/j.1365-2664.1999.00440.x

1105   Marini, M. Â., Barbet-Massin, M., Lopes, L. E., & Jiguet, F. (2009). Predicted climate-driven bird

1106   distribution changes and forecasted conservation conflicts in a Neotropical savanna. *Conservation*

1107   *Biology*, *23*(6), 1558–1567. https://doi.org/10.1111/j.1523-1739.2009.01258.x

1108   Matuoka, M. A., Benchimol, M., Almeida-Rocha, J. M. de, & Morante-Filho, J. C. (2020). Effects of

1109   anthropogenic disturbances on bird functional diversity: A global meta-analysis. *Ecological*

1110   *Indicators*, *116*, 106471. https://doi.org/10.1016/j.ecolind.2020.106471

1111   McKinnon, E. A., & Love, O. P. (2018). Ten years tracking the migrations of small landbirds: Lessons

1112   learned in the golden age of bio-logging. *The Auk*, *135*(4), 834–856.

1113   https://doi.org/10.1642/AUK-17-202.1

Merow, C., Smith, M. J., & Silander, J. A., Jr. (2013). A practical guide to MaxEnt for modeling species'

distributions: What it does, and why inputs and settings matter. *Ecography*, *36*(10), 1058–1069.

https://doi.org/10.1111/j.1600-0587.2013.07872.x

Mi, C., Huettmann, F., Guo, Y., Han, X., & Wen, L. (2017). Why choose Random Forest to predict rare

species distribution with few samples in large undersampled areas? Three Asian crane species

models provide supporting evidence. *PeerJ*, *5*, e2849. https://doi.org/10.7717/peerj.2849

Møller, A. P., Rubolini, D., & Lehikoinen, E. (2008). Populations of migratory bird species that did not

show a phenological response to climate change are declining. *Proceedings of the National*

Academy of Sciences, *105*(42), 16195–16200. https://doi.org/10.1073/pnas.0803825105

Moore, R. P., Robinson, W. D., Lovette, I. J., & Robinson, T. R. (2008). Experimental evidence for

extreme dispersal limitation in tropical forest birds. *Ecology Letters*, *11*(9), 960–968.

https://doi.org/10.1111/j.1461-0248.2008.01196.x

Mora, C., Frazier, A. G., Longman, R. J., Dacks, R. S., Walton, M. M., Tong, E. J., Sanchez, J. J., Kaiser,

L. R., Stender, Y. O., Anderson, J. M., Ambrosino, C. M., Fernandez-Silva, I., Giuseffi, L. M., &

Giambelluca, T. W. (2013). The projected timing of climate departure from recent variability.

Nature, *502*(7470), 183–187. https://doi.org/10.1038/nature12540

Morisette, J. T., Jarnevich, C. S., Holcombe, T. R., Talbert, C. B., Ignizio, D., Talbert, M. K., Silva, C.,

Koop, D., Swanson, A., & Young, N. E. (2013). VisTrails SAHM: Visualization and workflow

management for species habitat modeling. *Ecography*, *36*(2), 129–135.

https://doi.org/10.1111/j.1600-0587.2012.07815.x

Moritz, C., & Agudo, R. (2013). The future of species under climate change: Resilience or decline?

Science, *341*(6145), 504–508. https://doi.org/10.1126/science.1237190

Morris, W. F., & Doak, D. F. (2002). *Quantitative conservation biology: Theory and practice of*

population viability analysis. Sinauer Associates.

Morrison, M. L. (1986). Bird populations as indicators of environmental change. In R. F. Johnston (Ed.),

Current Ornithology: Volume 3 (pp. 429–451). Springer US. https://doi.org/10.1007/978-1-4615-

6784-4_10

Nakao, K., Higa, M., Tsuyama, I., Matsui, T., Horikawa, M., & Tanaka, N. (2013). Spatial conservation

planning under climate change: Using species distribution modeling to assess priority for adaptive

management of Fagus crenata in Japan. *Journal for Nature Conservation*, *21*(6), 406–413.

https://doi.org/10.1016/j.jnc.2013.06.003

Neilan, W. L., Barton, P. S., McAlpine, C. A., Wood, J. T., & Lindenmayer, D. B. (2019). Contrasting

effects of mosaic structure on alpha and beta diversity of bird assemblages in a human-modified

landscape. *Ecography*, *42*(1), 173–186. https://doi.org/10.1111/ecog.02981

Nilsson, C., Klaassen, R. H. G., & Alerstam, T. (2013). Differences in speed and duration of bird

migration between spring and autumn. *The American Naturalist*, *181*(6), 837–845.

https://doi.org/10.1086/670335

Oksanen, J., & Minchin, P. R. (2002). Continuum theory revisited: What shape are species responses

along ecological gradients? *Ecological Modelling*, *157*(2), 119–129.

https://doi.org/10.1016/S0304-3800(02)00190-4

Oppel, S., Meirinho, A., Ramírez, I., Gardner, B., O'Connell, A. F., Miller, P. I., & Louzao, M. (2012).

Comparison of five modelling techniques to predict the spatial distribution and abundance of

seabirds. *Biological Conservation*, *156*, 94–104. https://doi.org/10.1016/j.biocon.2011.11.013

Palacio, F. X., & Girini, J. M. (2018). Biotic interactions in species distribution models enhance model

performance and shed light on natural history of rare birds: A case study using the straight-billed

reedhaunter Limnoctites rectirostris. *Journal of Avian Biology*, *49*(11), e01743.

https://doi.org/10.1111/jav.01743

Palma, A. C., Goosem, M., Fensham, R. J., Goosem, S., Preece, N. D., Stevenson, P. R., & Laurance, S.

G. W. (2020). Dispersal and recruitment limitations in secondary forests. *Journal of Vegetation*

Science, *32*, e12975. https://doi.org/10.1111/jvs.12975

Papeş, M., Peterson, A. T., & Powell, G. V. N. (2012). Vegetation dynamics and avian seasonal

migration: Clues from remotely sensed vegetation indices and ecological niche modelling.

Journal of Biogeography, *39*(4), 652–664. https://doi.org/10.1111/j.1365-2699.2011.02632.x

Parmesan, C., & Matthews, J. (2005). *Biological Impacts of Climate Change*.

https://doi.org/10.17226/12491

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across

natural systems. *Nature*, *421*(6918), 37–42. https://doi.org/10.1038/nature01286

Pearl, J. (1985). Bayesian networks: A model of self-activated memory for evidential reasoning.

Proceedings of the 7th Conference of the Cognitive Science Society.

Pearson, R. G., & Dawson, T. P. (2003). Predicting the impacts of climate change on the distribution of

species: Are bioclimate envelope models useful? *Global Ecology and Biogeography*, *12*(5), 361–

371. https://doi.org/10.1046/j.1466-822X.2003.00042.x

Pearson, R. G., Raxworthy, C. J., Nakamura, M., & Peterson, A. T. (2007). Predicting species

distributions from small numbers of occurrence records: A test case using cryptic geckos in

Madagascar. *Journal of Biogeography*, *34*(1), 102–117. https://doi.org/10.1111/j.1365-

2699.2006.01594.x

Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I.-C., Clark, T. D., Colwell,

R. K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., Garcia, R. A., Griffis, R.

B., Hobday, A. J., Janion-Scheepers, C., Jarzyna, M. A., Jennings, S., … Williams, S. E. (2017).

Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being.

Science, *355*(6332). https://doi.org/10.1126/science.aai9214

Pérez-Granados, C., Bota, G., Giralt, D., Barrero, A., Gómez-Catasús, J., Rosa, D. B.-D. L., & Traba, J.

(2019). Vocal activity rate index: A useful method to infer terrestrial bird abundance with

acoustic monitoring. *Ibis*, *161*(4), 901–907. https://doi.org/10.1111/ibi.12728

Pérez-Granados, C., & Schuchmann, K.-L. (2020). Monitoring the annual vocal activity of two enigmatic

nocturnal Neotropical birds: The Common Potoo (Nyctibius griseus) and the Great Potoo

1190    (Nyctibius grandis). *Journal of Ornithology*, *161*(4), 1129–1141. https://doi.org/10.1007/s10336-

1191    020-01795-4

1192  Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species

1193    geographic distributions. *Ecological Modelling*, *190*(3), 231–259.

1194    https://doi.org/10.1016/j.ecolmodel.2005.03.026

1195  Pierson, J. C., Beissinger, S. R., Bragg, J. G., Coates, D. J., Oostermeijer, J. G. B., Sunnucks, P.,

1196    Schumaker, N. H., Trotter, M. V., & Young, A. G. (2015). Incorporating evolutionary processes

1197    into population viability models. *Conservation Biology*, *29*(3), 755–764.

1198    https://doi.org/10.1111/cobi.12431

1199  Pollock, H. S., Brawn, J. D., & Cheviron, Z. A. (2021). Heat tolerances of temperate and tropical birds

1200    and their implications for susceptibility to climate warming. *Functional Ecology*, *35*(1), 93–104.

1201    https://doi.org/10.1111/1365-2435.13693

1202  Poloczanska, E. S., Brown, C. J., Sydeman, W. J., Kiessling, W., Schoeman, D. S., Moore, P. J., Brander,

1203    K., Bruno, J. F., Buckley, L. B., Burrows, M. T., Duarte, C. M., Halpern, B. S., Holding, J.,

1204    Kappel, C. V., O'Connor, M. I., Pandolfi, J. M., Parmesan, C., Schwing, F., Thompson, S. A., &

1205    Richardson, A. J. (2013). Global imprint of climate change on marine life. *Nature Climate*

1206    *Change*, *3*(10), 919–925. https://doi.org/10.1038/nclimate1958

1207  Porter, W. P., Mitchell, J. W., Beckman, W. A., & DeWitt, C. B. (1973). Behavioral implications of

1208    mechanistic ecology. *Oecologia*, *13*(1), 1–54. https://doi.org/10.1007/BF00379617

1209  Powell, G. V. N., & Bjork, R. (1995). Implications of intratropical migration on reserve design: A case

1210    study using Pharomachrus mocinno. *Conservation Biology*, *9*(2), 354–362.

1211    https://doi.org/10.1046/j.1523-1739.1995.9020354.x

1212  Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P.,

1213    Adriaensen, F., Ahola, M. P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K. S., Borras, A.,

1214    Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A. A., … Kramer-Schadt, S. (2019).

Adaptive responses of animals to climate change are most likely insufficient. *Nature*

Communications, *10*(1), 3109. https://doi.org/10.1038/s41467-019-10924-4

Radosavljevic, A., & Anderson, R. P. (2014). Making better Maxent models of species distributions:

Complexity, overfitting and evaluation. *Journal of Biogeography*, *41*(4), 629–643.

https://doi.org/10.1111/jbi.12227

Randin, C. F., Ashcroft, M. B., Bolliger, J., Cavender-Bares, J., Coops, N. C., Dullinger, S., Dirnböck, T.,

Eckert, S., Ellis, E., Fernández, N., Giuliani, G., Guisan, A., Jetz, W., Joost, S., Karger, D.,

Lembrechts, J., Lenoir, J., Luoto, M., Morin, X., … Payne, D. (2020). Monitoring biodiversity in

the Anthropocene using remote sensing in species distribution models. *Remote Sensing of*

Environment, *239*, 111626. https://doi.org/10.1016/j.rse.2019.111626

Rappole, J. H. (1995). *The ecology of migrant birds: A neotropical perspective*. Smithsonian institution

Press.

Reed, J. M., Boulinier, T., Danchin, E., & Oring, L. W. (1999). Informed Dispersal. In V. Nolan, E. D.

Ketterson, & C. F. Thompson (Eds.), *Current Ornithology* (pp. 189–259). Springer US.

https://doi.org/10.1007/978-1-4757-4901-4_5

Rehm, E., & Feeley, K. J. (2016). Many species risk mountain top extinction long before they reach the

top. *Frontiers of Biogeography*, *8*(1). https://doi.org/10.21425/F5FBG27788

Renfrew, R. B., Kim, D., Perlut, N., Smith, J., Fox, J., & Marra, P. P. (2013). Phenological matching

across hemispheres in a long-distance migratory bird. *Diversity and Distributions*, *19*(8), 1008–

1019. https://doi.org/10.1111/ddi.12080

Reznick, D., Baxter, R. J., & Endler, J. (1994). Long-term studies of tropical stream fish communities:

The use of field notes and museum collections to reconstruct communities of the past. *Integrative*

and Comparative Biology, *34*(3), 452–462. https://doi.org/10.1093/icb/34.3.452

Ribas, C. C., Aleixo, A., Gubili, C., d'Horta, F. M., Brumfield, R. T., & Cracraft, J. (2018). Biogeography

and diversification of Rhegmatorhina (Aves: Thamnophilidae): Implications for the evolution of

Amazonian landscapes during the Quaternary. *Journal of Biogeography*, *45*(4), 917–928.

https://doi.org/10.1111/jbi.13169

Ribas, C. C., Aleixo, A., Nogueira, A. C. R., Miyaki, C. Y., & Cracraft, J. (2012). A palaeobiogeographic

model for biotic diversification within Amazonia over the past three million years. *Proceedings of*

the Royal Society B: Biological Sciences, *279*(1729), 681–689.

https://doi.org/10.1098/rspb.2011.1120

Riddell, E. A., Iknayan, K. J., Hargrove, L., Tremor, S., Patton, J. L., Ramirez, R., Wolf, B. O., &

Beissinger, S. R. (2021). Exposure to climate change drives stability or collapse of desert

mammal and bird communities. *Science*, *371*(6529), 633–636.

https://doi.org/10.1126/science.abd4605

Riotte-Lambert, L., & Matthiopoulos, J. (2020). Environmental Predictability as a Cause and

Consequence of Animal Movement. *Trends in Ecology and Evolution*, *35*(2), 163–174.

https://doi.org/10.1016/j.tree.2019.09.009

Ritter, C. D., Coelho, L. A., Capurucho, J. M., Borges, S. H., Cornelius, C., & Ribas, C. C. (2021). Sister

species, different histories: Comparative phylogeography of two bird species associated with

Amazonian open vegetation. *Biological Journal of the Linnean Society*, *132*(1), 161–173.

https://doi.org/10.1093/biolinnean/blaa167

Rosado, B. H. P., Figueiredo, M. S. L., Mattos, E. A. de, & Grelle, C. E. V. (2016). Eltonian shortfall due

to the Grinnellian view: Functional ecology between the mismatch of niche concepts. *Ecography*,

39(11), 1034–1041. https://doi.org/10.1111/ecog.01678

Roslin, T., Antão, L., Hällfors, M., Meyke, E., Lo, C., Tikhonov, G., Delgado, M. del M., Gurarie, E.,

Abadonova, M., Abduraimov, O., Adrianova, O., Akimova, T., Akkiev, M., Ananin, A.,

Andreeva, E., Andriychuk, N., Antipin, M., Arzamascev, K., Babina, S., … Ovaskainen, O.

(2021). Phenological shifts of abiotic events, producers and consumers across a continent. *Nature*

Climate Change, *11*(3), 241–248. https://doi.org/10.1038/s41558-020-00967-7

Sander, M., Chamberlain, D., Mermillon, C., Alba, R., Jähnig, S., Rosselli, D., & Lisovski, S. (2020).

Reduced breeding success suggests trophic mismatch despite timely arrival in an alpine songbird

[Preprint]. https://assets.researchsquare.com/files/rs-137126/v1/650443e6-45ba-470a-814c-

47dde4d049d7.pdf

Saupe, E. E., Barve, V., Myers, C. E., Soberón, J., Barve, N., Hensz, C. M., Peterson, A. T., Owens, H.

L., & Lira-Noriega, A. (2012). Variation in niche and distribution model performance: The need

for a priori assessment of key causal factors. *Ecological Modelling*, *237–238*, 11–22.

https://doi.org/10.1016/j.ecolmodel.2012.04.001

Sax, D. F., Early, R., & Bellemare, J. (2013). Niche syndromes, species extinction risks, and management

under climate change. *Trends in Ecology & Evolution*, *28*(9), 517–523.

https://doi.org/10.1016/j.tree.2013.05.010

Scheffers, B. R., Edwards, D. P., Diesmos, A., Williams, S. E., & Evans, T. A. (2014). Microhabitats

reduce animal's exposure to climate extremes. *Global Change Biology*, *20*(2), 495–503.

https://doi.org/10.1111/gcb.12439

Scheffers, B. R., Edwards, D. P., Macdonald, S. L., Senior, R. A., Andriamahohatra, L. R., Roslan, N.,

Rogers, A. M., Haugaasen, T., Wright, P., & Williams, S. E. (2017). Extreme thermal

heterogeneity in structurally complex tropical rain forests. *Biotropica*, *49*(1), 35–44.

https://doi.org/10.1111/btp.12355

Sekercioglu, C. H. (2010). Partial migration in tropical birds: The frontier of movement ecology. *Journal*

of Animal Ecology, *79*(5), 933–936. https://doi.org/10.1111/j.1365-2656.2010.01739.x

Şekercioğlu, C., Wenny, D. G., & Whelan, C. J. (2019). *Why birds matter: Avian ecological function and*

ecosystem services.

http://chicago.universitypressscholarship.com/10.7208/chicago/9780226382777.001.0001/upso-

9780226382463

Shaffer, H. B., Fisher, R. N., & Davidson, C. (1998). The role of natural history collections in documenting species declines. *Trends in Ecology & Evolution*, *13*(1), 27–30. https://doi.org/10.1016/S0169-5347(97)01177-4

Sheard, C., Neate-Clegg, M. H. C., Alioravainen, N., Jones, S. E. I., Vincent, C., MacGregor, H. E. A., Bregman, T. P., Claramunt, S., & Tobias, J. A. (2020). Ecological drivers of global gradients in avian dispersal inferred from wing morphology. *Nature Communications*, *11*(1), 2463. https://doi.org/10.1038/s41467-020-16313-6

Sheldon, K. S., Huey, R. B., Kaspari, M., & Sanders, N. J. (2018). Fifty years of mountain passes: A perspective on Dan Janzen's classic article. *The American Naturalist*, *191*(5), 553–565. https://doi.org/10.1086/697046

Shipley, J. R., Twining, C. W., Taff, C. C., Vitousek, M. N., Flack, A., & Winkler, D. W. (2020). Birds advancing lay dates with warming springs face greater risk of chick mortality. *Proceedings of the National Academy of Sciences*, *117*(41), 25590–25594. https://doi.org/10.1073/pnas.2009864117

Shirley, S. M., Yang, Z., Hutchinson, R. A., Alexander, J. D., McGarigal, K., & Betts, M. G. (2013). Species distribution modelling for the people: Unclassified landsat TM imagery predicts bird occurrence at fine resolutions. *Diversity and Distributions*, *19*(7), 855–866. https://doi.org/10.1111/ddi.12093

Siepielski, A. M., Morrissey, M. B., Buoro, M., Carlson, S. M., Caruso, C. M., Clegg, S. M., Coulson, T., DiBattista, J., Gotanda, K. M., Francis, C. D., Hereford, J., Kingsolver, J. G., Augustine, K. E., Kruuk, L. E. B., Martin, R. A., Sheldon, B. C., Sletvold, N., Svensson, E. I., Wade, M. J., & MacColl, A. D. C. (2017). Precipitation drives global variation in natural selection. *Science*, *355*(6328), 959–962. https://doi.org/10.1126/science.aag2773

Slager, D. L. (2020). Seasonal and directional dispersal behavior in an ongoing dove invasion. *Journal of Avian Biology*, *51*(2). https://doi.org/10.1111/jav.02332

Smith, A. C., & Edwards, B. P. M. (2021). North American Breeding Bird Survey status and trend estimates to inform a wide range of conservation needs, using a flexible Bayesian hierarchical

generalized additive model. *Ornithological Applications*, *123*(duaa065).

https://doi.org/10.1093/ornithapp/duaa065

Smith, B. T., Gehara, M., & Harvey, M. G. (2021). The demography of extinction in eastern North

American birds. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1944), 20201945.

https://doi.org/10.1098/rspb.2020.1945

Smith, F. A., Betancourt, J. L., & Brown, J. H. (1995). Evolution of Body Size in the Woodrat over the

Past 25,000 Years of Climate Change. *Science*, *270*(5244), 2012–2014.

https://doi.org/10.1126/science.270.5244.2012

Smith, R. J., & Moore, F. R. (2005). Arrival timing and seasonal reproductive performance in a long-

distance migratory landbird. *Behavioral Ecology and Sociobiology*, *57*(3), 231–239.

https://doi.org/10.1007/s00265-004-0855-9

Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species. *Ecology*

Letters, *10*(12), 1115–1123. https://doi.org/10.1111/j.1461-0248.2007.01107.x

Soberón, J., & Peterson, A. T. (2005). INTERPRETATION OF MODELS OF FUNDAMENTAL

ECOLOGICAL NICHES AND SPECIES' DISTRIBUTIONAL AREAS. In *Biodiversity*

Informatics (Vol. 2, pp. 1–10).

https://kuscholarworks.ku.edu/bitstream/handle/1808/20560/soberon_interpretation.pdf?sequence

=1&isAllowed=y

Song, G., Zhang, R., Machado-Stredel, F., Alström, P., Johansson, U. S., Irestedt, M., Mays, H. L.,

McKay, B. D., Nishiumi, I., Cheng, Y., Qu, Y., Ericson, P. G. P., Fjeldså, J., Peterson, A. T., &

Lei, F. (2020). Great journey of Great Tits (Parus major group): Origin, diversification and

historical demographics of a broadly distributed bird lineage. *Journal of Biogeography*, *47*(7),

1585–1598. https://doi.org/10.1111/jbi.13863

Staniczenko, P. P. A., Sivasubramaniam, P., Suttle, K. B., & Pearson, R. G. (2017). Linking

macroecology and community ecology: Refining predictions of species distributions using biotic

interaction networks. *Ecology Letters*, *20*(6), 693–707. https://doi.org/10.1111/ele.12770

Stanley, C. Q., MacPherson, M., Fraser, K. C., McKinnon, E. A., & Stutchbury, B. J. M. (2012). Repeat

tracking of individual songbirds reveals consistent migration timing but flexibility in route. *PLOS*

ONE, *7*(7), e40688. https://doi.org/10.1371/journal.pone.0040688

Stanley, C. Q., Mckinnon, E. A., Fraser, K. C., Macpherson, M. P., Casbourn, G., Friesen, L., Marra, P.

P., Studds, C., Ryder, T. B., Diggs, N. E., & Stutchbury, B. J. M. (2015). Connectivity of wood

thrush breeding, wintering, and migration sites based on range-wide tracking. *Conservation*

Biology, *29*(1). https://doi.org/10.1111/cobi.12352

Stevenson, R. D. (1985). The relative importance of behavioral and physiological adjustments controlling

body temperature in terrestrial ectotherms. *The American Naturalist*, *126*(3), 362–386.

https://doi.org/10.1086/284423

Stiels, D., Bastian, H.-V., Bastian, A., Schidelko, K., & Engler, J. O. (2021). An iconic messenger of

climate change? Predicting the range dynamics of the European Bee-eater (Merops apiaster).

Journal of Ornithology. https://doi.org/10.1007/s10336-021-01867-z

Stralberg, D., Matsuoka, S. M., Hamann, A., Bayne, E. M., Sólymos, P., Schmiegelow, F. K. A., Wang,

X., Cumming, S. G., & Song, S. J. (2015). Projecting boreal bird responses to climate change:

The signal exceeds the noise. *Ecological Applications*, *25*(1), 52–69. https://doi.org/10.1890/13-

2289.1

Sugai, L. S. M., Desjonquères, C., Silva, T. S. F., & Llusia, D. (2020). A roadmap for survey designs in

terrestrial acoustic monitoring. *Remote Sensing in Ecology and Conservation*, *6*(3), 220–235.

https://doi.org/10.1002/rse2.131

Suh, Y. H., Pesendorfer, M. B., Tringali, A., Bowman, R., & Fitzpatrick, J. W. (2020). Investigating

social and environmental predictors of natal dispersal in a cooperative breeding bird. *Behavioral*

Ecology, *31*(3), 692–701. https://doi.org/10.1093/beheco/araa007

Sullivan, B. L., Aycrigg, J. L., Barry, J. H., Bonney, R. E., Bruns, N., Cooper, C. B., Damoulas, T.,

Dhondt, A. A., Dietterich, T., Farnsworth, A., Fink, D., Fitzpatrick, J. W., Fredericks, T.,

Gerbracht, J., Gomes, C., Hochachka, W. M., Iliff, M. J., Lagoze, C., La Sorte, F. A., … Kelling,

S. (2014). The eBird enterprise: An integrated approach to development and application of citizen

science. *Biological Conservation*, *169*, 31–40. https://doi.org/10.1016/j.biocon.2013.11.003

Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird: A citizen-

based bird observation network in the biological sciences. *Biological Conservation*, *142*(10),

2282–2292. https://doi.org/10.1016/j.biocon.2009.05.006

Taylor, S. A., White, T. A., Hochachka, W. M., Ferretti, V., Curry, R. L., & Lovette, I. (2014). Climate-

Mediated Movement of an Avian Hybrid Zone. *Current Biology*, *24*(6), 671–676.

https://doi.org/10.1016/j.cub.2014.01.069

Terribile, L. C. (2012). Areas of climate stability of species ranges in the Brazilian Cerrado:

Disentangling uncertainties through time. *Natureza & Conservação*, *10*(2), 152–159.

https://doi.org/10.4322/natcon.2012.025

Thomas, C. D., Franco, A. M. A., & Hill, J. K. (2006). Range retractions and extinction in the face of

climate warming. *Trends in Ecology & Evolution*, *21*(8), 415–416.

https://doi.org/10.1016/j.tree.2006.05.012

Thorup, K., Tøttrup, A. P., Willemoes, M., Klaassen, R. H. G., Strandberg, R., Vega, M. L., Dasari, H. P.,

Araújo, M. B., Wikelski, M., & Rahbek, C. (2017). Resource tracking within and across

continents in long-distance bird migrants. *Science Advances*, *3*(1), e1601360.

https://doi.org/10.1126/sciadv.1601360

Thuiller, W., Lafourcade, B., Engler, R., & Araújo, M. B. (2009). BIOMOD – a platform for ensemble

forecasting of species distributions. *Ecography*, *32*(3), 369–373. https://doi.org/10.1111/j.1600-

0587.2008.05742.x

Tingley, M. W., Darling, E., & Wilcove, D. S. (2014). Fine- and coarse-filter conservation strategies in a

time of climate change. *Annals of the New York Academy of Sciences*, *1322*(The Year in Ecology

and Conservation Biology), 92–109. https://doi.org/10.1111/nyas.12484

Tingley, M. W., Monahan, W. B., Beissinger, S. R., & Moritz, C. (2009). Birds track their Grinnellian

niche through a century of climate change. *Proceedings of the National Academy of Sciences*,

106(Supplement 2), 19637–19643. https://doi.org/10.1073/pnas.0901562106

Tøttrup, A. P., Klaassen, R. H. G., Kristensen, M. W., Strandberg, R., Vardanis, Y., Lindström, Å.,

Rahbek, C., Alerstam, T., & Thorup, K. (2012). Drought in Africa caused delayed arrival of

European songbirds. *Science*, *338*(6112), 1307–1307. https://doi.org/10.1126/science.1227548

Travis, J. M. J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., Boulangeat, I., Hodgson, J.

A., Kubisch, A., Penteriani, V., Saastamoinen, M., Stevens, V. M., & Bullock, J. M. (2013).

Dispersal and species' responses to climate change. *Oikos*, *122*(11), 1532–1540.

https://doi.org/10.1111/j.1600-0706.2013.00399.x

Twedt, D. J. (2013). Foraging Habitat for Shorebirds in Southeastern Missouri and its Predicted Future

Availability. *Wetlands*, *33*(4), 667–678. https://doi.org/10.1007/s13157-013-0422-0

Urban, M. C., Zarnetske, P. L., & Skelly, D. K. (2013). Moving forward: Dispersal and species

interactions determine biotic responses to climate change. *Annals of the New York Academy of*

Sciences, *1297*, 44–60. https://doi.org/10.1111/nyas.12184

van Rees, C. B., & Reed, J. M. (2018). Predicted effects of landscape change, sea level rise, and habitat

management on the extirpation risk of the Hawaiian common gallinule (Gallinula galeata

sandvicensis) on the island of Oʻahu. *PeerJ*, *6*, e4990. https://doi.org/10.7717/peerj.4990

Visser, M. E. (2008). Keeping up with a warming world; assessing the rate of adaptation to climate

change. *Proceedings of the Royal Society B: Biological Sciences*, *275*(1635), 649–659.

https://doi.org/10.1098/rspb.2007.0997

Weiner, J. (1994). *The Beak of the Finch: A Story of Evolution in Our Time*.

https://doi.org/10.5860/choice.32-1538

Williams, J. W., Jackson, S. T., & Kutzbach, J. E. (2007). Projected distributions of novel and

disappearing climates by 2100 AD. *Proceedings of the National Academy of Sciences*, *104*(14),

5738–5742. https://doi.org/10.1073/pnas.0606292104

Winger, B. M., Auteri, G. G., Pegan, T. M., & Weeks, B. C. (2019). A long winter for the Red Queen:

Rethinking the evolution of seasonal migration. *Biological Reviews*, *94*(3), 737–752.

https://doi.org/10.1111/brv.12476

Wisz, M. S., Hijmans, R. J., Li, J., Peterson, A. T., Graham, C. H., & Guisan, A. (2008). Effects of

sample size on the performance of species distribution models. *Diversity and Distributions*, *14*(5),

763–773. https://doi.org/10.1111/j.1472-4642.2008.00482.x

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F.,

Forchhammer, M. C., Grytnes, J.-A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto,

M., Maiorano, L., Nilsson, M.-C., Normand, S., Öckinger, E., Schmidt, N. M., … Svenning, J.-C.

(2013). The role of biotic interactions in shaping distributions and realised assemblages of

species: Implications for species distribution modelling. *Biological Reviews*, *88*(1), 15–30.

https://doi.org/10.1111/j.1469-185X.2012.00235.x

Wogan, G. O. U., Voelker, G., Oatley, G., & Bowie, R. C. K. (2020). Biome stability predicts population

structure of a southern African aridland bird species. *Ecology and Evolution*, *10*(9), 4066–4081.

https://doi.org/10.1002/ece3.6175

Wolfson, D. W., Fieberg, J. R., & Andersen, D. E. (2020). Juvenile Sandhill Cranes exhibit wider ranging

and more exploratory movements than adults during the breeding season. *Ibis*, *162*(2), 556–562.

https://doi.org/10.1111/ibi.12786

Wüest, R. O., Bergamini, A., Bollmann, K., & Baltensweiler, A. (2020). LiDAR data as a proxy for light

availability improve distribution modelling of woody species. *Forest Ecology and Management*,

456, 117644. https://doi.org/10.1016/j.foreco.2019.117644

Xia, J., Wang, J., & Niu, S. (2020). Research challenges and opportunities for using big data in global

change biology. *Global Change Biology*, *26*(11), 6040–6061. https://doi.org/10.1111/gcb.15317

1440
1441