TITLE: An introduction to predictive distribution modelling for conservation

to encourage novel perspectives

Authors: MacPherson[1,2], M.P.*, Burgio[3,4], K.R., DeSaix[5], M.G., Freeman[6], B.G., Herbert[7,8], J.,

Herman[9], R., Jirinec[10,11], V., Shonfield[12], J., Slager[13], D.L., van Rees[14], C.B., and J.E. Jankowski[6]

ABSTRACT

An introduction to predictive distribution modelling for conservation to encourage novel

perspectives. The rapid pace and potentially irreversible consequences of global change create an

urgent need to predict the spatial responses of biota for conservation to better inform the

prioritization and management of terrestrial habitats and prevent future extinctions. Here, we

provide an accessible entry point to the field to guide near-future work building predictive

species distribution models (SDMs) by synthesizing a technical framework for the proactive

conservation of avian biodiversity. Our framework offers a useful approach to navigate the

challenges surrounding the large spatio-temporal resolution of datasets and datasets that favor

hypothesis testing at broad spatio-temporal scales and coarse resolutions, which can affect our

ability to assess the validity of current predicted distributions. We explain how to improve the

accuracy of predictive models by determining the extent to which: 1) dispersal limitation impacts

the rate of range shifts, 2) taxa are rare at their range limits, and 3) land use and climate change

interact. Finally, we offer approaches to filling knowledge gaps by creatively leveraging existing

methods and data sources.

KEY WORDS: species distribution model, environmental niche model, predictive distribution

modeling, climate change, Grinnellian niche, Eltonian niche.

RESUMEN

Una introducción a la modelización predictiva de la distribución para la conservación con el fin de fomentar nuevas perspectivas. El rápido ritmo y las consecuencias potencialmente irreversibles del cambio global crean una necesidad urgente de predecir las respuestas espaciales de la biota para la conservación, con el fin de informar mejor la priorización y gestión de los hábitats terrestres y prevenir futuras extinciones. Aquí proporcionamos un punto de entrada accesible al campo para guiar el trabajo del futuro próximo en la construcción de modelos predictivos de distribución de especies (SDM), sintetizando un marco técnico para la conservación proactiva de la biodiversidad aviar. Nuestro marco ofrece un enfoque útil para navegar por los retos que rodean a la gran resolución espacio-temporal de los conjuntos de datos y a los conjuntos de datos que favorecen la comprobación de hipótesis a escalas espacio-temporales amplias y resoluciones gruesas, lo que puede afectar a nuestra capacidad para evaluar la validez de las distribuciones predichas actuales. Explicamos cómo mejorar la precisión de los modelos predictivos determinando hasta qué punto 1) la limitación de la dispersión influye en el ritmo de los cambios de área de distribución, 2) los taxones son raros en los límites de su área de distribución, y 3) el uso del suelo y el cambio climático interactúan. Por último, proponemos enfoques para colmar las lagunas de conocimiento aprovechando de forma creativa los métodos y fuentes de datos existentes.

PALABRAS CLAVES: modelo de distribución de especies, modelo de nicho ambiental, modelización predictiva de la distribución, cambio climático, nicho grinnelliano, nicho eltoniano.

Author Affiliations:

M.P. MacPherson (https://orcid.org/0000-0002-9522-5688)

1. Louisiana State University Museum of Natural Sciences, Louisiana State University, Baton Rouge, LA, USA

2. Département des sciences du bois et de la forêt, Université Laval, Québec City, Québec, Canada

K.R. Burgio (https://orcid.org/0000-0002-8375-2501)

3. New York City Audubon Society, New York, NY, USA

4. Department of Ecology and Evolutionary Ecology, University of Connecticut, Storrs, CT, USA

M.G. DeSaix (https://orcid.org/0000-0002-5721-0311)

5. Department of Biology, Colorado State University, Fort Collins, CO, USA

B.G. Freeman (https://orcid.org/0000-0001-6131-6832)

J.E. Jankowski (https://orcid.org/0000-0003-3273-1388)

6. Biodiversity Research Centre and Department of Zoology, University of British Columbia, Vancouver, BC, Canada

J. Herbert (https://orcid.org/0000-0002-2912-2226)

7. Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA

8. Rhode Island Department of Environmental Management, Division of Fish & Wildlife, West Kingston, RI, USA

R. Herman (https://orcid.org/0000-0002-5914-6907)

9. Department of Ecology and Evolution, Stony Brook University, NY

V. Jirinec (https://orcid.org/0000-0001-9856-9681)

10. School of Renewable Natural Resources, Louisiana State University and LSU AgCenter, Baton Rouge, LA, USA

11. Integral Ecology Research Center, 239 Railroad Ave., Blue Lake, CA, USA

J. Shonfield (https://orcid.org/0000-0002-7159-4016)

12. LGL Limited, Cambridge, ON, Canada

D.L. Slager (https://orcid.org/0000-0003-2525-2039)

13. Department of Biology, University of Washington, Seattle, WA, USA

C.B. van Rees (https://orcid.org/0000-0003-0558-3674)

14. River Basin Center & Odum School of Ecology, University of Georgia

*Corresponding author: maggie.macpherson@gmail.com

Author Contributions:

Maggie MacPherson - concept, symposium organizer, symposium participant, writing & revising, editing full manuscript, topic editor (abstract, theory) and co-editor (temporal resolution of data types), figures, tables.

Kevin R. Burgio - concept, overall project support, symposium participant, writing & revising, conceptualizing and constructing figures, editing full manuscript, topic co-editor (methods & data types for spatial data)

Matthew G. DeSaix – symposium participant, writing & revising, topic co-editor (temporal resolution of data types)

Ben Freeman - symposium participant, writing, publication approach/ideas, topic editor

(discussion)

Vitek Jirinec - symposium participant, writing & revising, topic co-editor (logistical methods and

data types for occurrence data), figures.

John Herbert - symposium participant, writing, topic co-editor (methods & data types for spatial

data)

Julia Shonfield - symposium participant, writing & revising, topic editor (introduction)

David Slager – symposium participant, writing & revising, topic editor (literature cited), topic

co-editor (logistical methods and data types for occurrence data)

Rachael Herman - writing, topic co-editor (logistical methods and data types for occurrence data)

Charles van Rees – writing & revising, topic editor (analytical methods), figures.

Jill Jankowski – concept, symposium participant, writing & revising, editing full manuscript

INTRODUCTION

Anthropogenic pressures on climate and land cover lead to altered ecosystems and species

distributions (Parmesan and Yohe, 2003). This creates an urgent need for understanding species

spatial responses to global change and ensuring conservation of suitable habitat that supports

population persistence and conserves biodiversity (De Frenne et al., 2021; Roberts et al., 2019).

Species distribution models link species' occurrence with ecological explanatory variables and

can be used to predict range dynamics for proactive conservation measures (van Rees et al.,

2022; Zurell et al., 2016); however, the accuracy of current predictive species distribution

models where there is a lot of occurrence data is still limited by the variation in temporal and

spatial resolution of datasets used to build them (König et al., 2021). This is because the capacity

of species to change their geographic ranges is limited by both large-scale factors (i.e., range-

wide drivers like temperature and rainfall that define macrohabitat) and small-scale factors (i.e.,

individual-specific biotic interactions and resource availability that affect microhabitat use). This

issue is exacerbated for data-poor species, such as rare, cryptic and consequently, many tropical

species. The limited availability of high-resolution datasets capable of capturing fine-scale

dynamics thus decreases our predictive capacity for many such taxa.

A multitude of factors can ultimately influence how species, populations, and individuals

respond to changing climate regimes driven by global change (reviewed in Peterson et al., 2019).

Individuals may lack flexibility in tracking relevant environmental cues to adjust the timing of

life history events or to otherwise accommodate novel ecological circumstances. Alternatively,

even when species are flexible in timing life history events, they may be unable to modulate

behavior in response to environmental anomalies that affect recruitment. For example, advancing

egg-laying dates to match earlier spring warming temperatures may have negative consequences

for nestling survival when the chances of cold snaps are decoupled from warming trends (Sauser

et al., 2021; Shipley et al., 2020). Predictions for how animal distributions could be altered

(shifted, contracted or expanded) with climate change may be species-specific (Hallman and

Robinson, 2020; Radosavljevic and Anderson, 2014), but research into how particular axes of

climate change affect species with different life history strategies, foraging guilds or habitat

affinities provides promising insights that may apply to a wider number of unstudied species

(changing rainfall regimes -Brawn et al., 2017; e.g., rearrangements of ecosystems - Huntley et

al., 2008; advances in spring phenology and breeding success - Sander et al., 2020). Finally, the

potential for range shifts is ultimately limited by whether potential habitat is available,

accessible, and whether population density is high enough at the appropriate range edge to

facilitate a shift (Block and Levine, 2021; Stiels et al., 2021).

A major gap in the current understanding of how organisms are likely to respond to

global change remains in the effect of land use change and its interaction with climate change.

Dispersal limitation is a critical component of distribution model theory, but is difficult to

estimate in practice (Sousa et al., 2021). Species with high site fidelity (Merkle et al., 2022) or

inflexible migratory routes (Stanley et al., 2012) may not be able to adopt prospecting behaviors

(searching for new sites; Cooper and Marra, 2020) to accommodate dispersal to appropriate

habitat following change. In the Northern Hemisphere, avian migration routes and resource

tracking are well established (Faaborg et al., 2010; Thorup et al., 2017), but individuals still may

not be able to correctly time migratory movements to keep up with the pace of climate change, as

evidenced in declining populations that now experience mismatches in the timing of annual life

history events and peaks in food availability (Møller et al., 2008; Sander et al., 2020). On the

other hand, multiple Nearctic-Neotropical migrants have advanced their spring migration and

breeding behavior in response to warming spring temperatures (Dunn and Møller, 2019; Pecl et

al., 2017; Shipley et al., 2020), offering support to the idea of "evolutionary rescue", where

existing variation within annual cycles may yield previously unknown adaptive potential (Helm

et al., 2019). Recent research on the interaction between land use and climate change points to

the need to improve our understanding of interacting mechanisms underpinning risks to

population persistence under climate and habitat stressors (Schulte to Bühne et al., 2021).

Here, we provide an overview of predictive bird distribution modeling with the aim to

inform future work focused on how dispersal limitation, biotic factors, and abiotic factors affect

distributional shifts in birds and other taxa driven by global change. Birds are excellent

indicators of environmental change, and they have the most widespread and in-depth databases

on distributions (Morrison, 1986; Riddell et al., 2021), making them excellent focal taxa for

predictive conservation modeling. Citizen science initiatives and long-term research programs

have yielded an abundance of predictive distribution literature, and insights from avian literature

should be generalizable to other taxa because birds display a wide variety of life history

strategies (Şekercioğlu et al., 2019).

We synthesize theory, methods, and data sources widely used to predict bird distributions

under global change to highlight best practices and opportunities for innovation and refinement.

Understanding where species occur and why some areas are occupied but not others is essential

for developing effective conservation plans. Predictive distribution modeling is important to

build upon our knowledge of distributions based on surveyed areas and to identify potential areas

for protection or management. Modeling wide ranging, unevenly distributed, cryptic, or difficult

to survey species all present problems for understanding factors driving occupancy. Here, we

bring together the issues that often complicate predictive distribution modeling to offer a

solutions-oriented approach to habitat conservation under global change.

ECOLOGICAL & EVOLUTIONARY THEORY AS A BACKBONE TO

APPROACHING PREDICTIVE SDMs

The first step in our synthesis is to illuminate the importance of a taxon's ecology and evolution

in determining their spatial response to global change. Dramatic changes in climate over the

remainder of this century are expected (Figure 1), with many species likely unable to survive in

all of the areas they presently occupy (Ceballos et al., 2015; Sax et al., 2013). Species typically

respond to climate change in one of three ways: 1) by going extinct, 2) persisting via shifts in

geographic range, or 3) adapting in place via evolutionary change. Evidence from the fossil

record suggests all three responses are common (Graham et al., 1996; Smith et al., 1995). The

capacity to adapt *in situ* in response to climate change has been relatively understudied (Jirinec et

al., 2021; Parmesan and Matthews, 2005). A meta-analysis of observed phenological and

morphological adaptations to climate change concluded that morphological trait adaptation,

while variable among taxa, will not likely work on a sufficient scale to mitigate the worst effects

of ongoing climate change, whereas phenological adaptation (e.g., timing of reproduction or

migration) could mitigate such effects, albeit imperfectly (Radchuk et al., 2019). Whether

species with relatively shorter generation times should be better able to survive by evolving in

place remains understudied. Advancing such research is hampered by the difficulty of

conducting experiments on large numbers of species at sufficient time scales needed to develop a

comprehensive framework (Holt et al., 2005; Pierson et al., 2015). Variability among taxa and

the short time scales in which we have been making observations together indicate that selection

pressure has yet to be strong enough to produce measurable changes, though it appears that

species shift their phenology faster where the rate of climate change is higher (Poloczanska et al.,

2013). Precipitation, and to a lesser extent temperature, are the primary drivers of adaptive

responses to climate change (Caruso et al., 2017; Jirinec et al., 2021; Siepielski et al., 2017).

Relevant ecological covariates inform one of two main modeling categories to predict

where a taxon could occur: 1) species distribution models (hereafter SDMs) that comprise

presence records and abiotic data, and 2) ecological niche models (hereafter ENMs) that

explicitly estimate the accessible environment (Soberón and Peterson, 2005). These approaches

are meant to have high spatial accuracy but are not intended to inform on cause-and-effect

species-habitat relationships (Merow et al., 2013). Whether using SDMs or ENMs, these models

can be based on different niche perspectives, which present unique frameworks for estimating

drivers of occupancy across a species' range. Here, we list simplified descriptions of three niche

concepts to provide a basic introductory framework for understanding the domain of predictive

distribution modeling as all are used in modern predictive modeling (see *An Overview of*

Analytical Methods section for more information). Models based on the Grinnellian niche

concept focus on abiotic drivers of site occupancy (Elton, 1927; Soberón, 2007; Wisz et al.,

2013) and SDMs are mainly built under this framework (Figure 1; Saupe et al., 2012). Coarse-

scale variables describing ecosystem characteristics are often the most relevant for predicting

shifting distributions across large spatial extents. In comparison, the Eltonian niche concept

includes estimates of biotic interactions and resource-consumer dynamics that are only

quantifiable at local scales (Figure 1; Elton, 1927; Soberón, 2007). Finally, the Hutchinsonian

niche concept reflects the functional role of a species, which is often estimated using functional

traits and habitat requirements based on functional traits (Rosado et al., 2016), and projects the

probability of site occupancy beyond study areas, including estimates of dispersal probabilities in

ENMs (Figure 2). Because a species' ability to move beyond their current realized niche depends

on the relative importance of abiotic factors, biotic interactions, and dispersal probability, which

can operate at different spatial scales (Jankowski et al., 2013), knowledge of these categories is

important for making sound inferences from whatever data and analytical approach are used.

While ENM tools can be used to predict spatially beyond the current range of a taxon,

they are only able to predict the probability of site occupancy within the range of abiotic

conditions measured within current range boundaries. Thus, assessing occurrence or abundance

along abiotic gradients is important for predicting future range shifts under global change, as this

relates directly to the response of species or populations to the environment. The gradual shifts in

temperature and rainfall regimes can be expected to align with changes in species responses,

reflecting the shifts in optimal ambient conditions and resulting in altered patterns of probability

of occurrence or abundance. However, ENM tools are limited where climate change introduces

novel conditions (e.g., Williams et al., 2007), and more mechanistic (rather than correlative)

analytical approaches may provide more accurate predictions.

Demographic information is also useful to incorporate for more accurate predictions

because spatial ecology theory dictates that species respond to continuous environmental

gradients through gradual changes in abundance, as individuals experience shifting conditions

toward or away from environmental optima over space (Austin, 2007, 2005). Populations may

also show thresholds in abiotic tolerances or in response to biotic factors that change their

response shape (e.g., creating asymmetric or skewed response functions; e.g., Oksanen and

Minchin, 2002). For example, the asymmetric abiotic stress limitation (AASL) hypothesis

predicts a steeper decrease in a species' probability of occurrence toward the more stressful end

of a species' distribution, which has been supported in several vascular plant species (Dvorský et

al., 2017). Other threshold-type effects can result in asymmetric responses shown across multiple

species in communities (e.g., sharp ecotone boundaries, appearance of dominant predators or

competitors; Jankowski et al., 2013). Identifying response shapes is of practical interest to

understand the impact of abiotic and biotic factors on occupancy. The evaluation of responses

along abiotic gradients are becoming more common but often rely on high spatial resolution

survey data (Bani et al., 2019; Burner et al., 2019; e.g., Maggini et al., 2011; Urli et al., 2014)

that is difficult to amass. However, with more studies, comparisons across species would enable

distinguishing commonalities and differences in responses to gradients and could point to key

environmental variables driving patterns in community organization.

Whether populations would be able to respond (in time and space) to changing conditions

also depends on heterogeneity within landscapes; this is the realm of landscape ecology land use

change. Landscapes can vary naturally or due to anthropogenic alterations in ways that affect

both the phenology of resources (time) or availability of habitat (space); such variability can be

tracked if conditions change at a pace that populations could respond to. Examples of this

include species adapted to annually ephemeral or temporally patchy habitats such as wetlands

and grasslands; species accustomed to such environments may have dispersal strategies that

predispose them to better track climate (e.g., *Elaenia cristata* - Ritter et al., 2021). In other cases,

habitat and resources are spatially patchily distributed, and the ability of species to respond to

changing conditions may be more taxon-specific for those that rely on spatiotemporal

heterogeneity that is naturally patchy. Examples of this include habitat specialists in naturally

patchy habitats such as Amazonian white sand habitats (e.g., Rufous-crowned Elaenia (*Elaenia*

ruficeps) - Ritter et al., 2021).

Alternatively, animals may avoid the problem of range shifting altogether by buffering

themselves from acutely unfavorable conditions using microclimate refugia. One example of this

is through behavioral thermoregulation. Ambient temperature is almost never constant in

terrestrial environments—it varies by time, and by habitat, often at very small scales (Scheffers

et al., 2017, 2014). Biologists have long understood that mobile animals exploit thermal

heterogeneity to maintain optimal body temperature (Angilletta Jr and Angilletta, 2009;

Angilletta et al., 2009; Cowles and Bogert, 1944; Porter et al., 1973; Stevenson, 1985). For cold-

adapted species in a warming world, this can be achieved by shifting activity times to cooler

periods of the year and day, and by moving to microhabitats with cooler microclimates. This

capacity to compensate for unfavorable ambient temperature by behavioral thermoregulation,

known as the "Bogert Effect" (Huey et al., 2003), can at least partly mitigate the harmful effects

of climate warming (Huey et al., 2012). However, behavioral thermoregulation is contingent on

access to cooler areas and periods; animals already occurring in the most buffered environments

have limited options for escape when conditions change. Furthermore, individuals could

experience additional pressure from biotic interactions with species moving into their habitat to

thermoregulate (Huey et al., 2012).

Whether taxa will be able to respond to global change is ultimately a taxon-specific

question driven by the constraints of dispersal limitation; one of three fundamental aspects of

distribution theory (Figure 2). Understanding barriers to distributions caused by dispersal

limitation has been of great interest, for example, in bird species unable to cross areas of

unsuitable habitat, and has been studied both experimentally (e.g., Moore et al., 2008; Naka et

al., 2022) and theoretically (Ribas et al., 2018, e.g., 2012). Selection pressures driving

morphology, such as migratory compared to sedentary life histories, may play a role in

physically limiting dispersal distance (Capurucho et al., 2020; Claramunt et al., 2012;

MacPherson et al., 2022; Sheard et al., 2020), even in birds. Life history strategies also dictate

the demographic trends in dispersal; for example, in many species it is the inexperienced young

that disperse into new areas away from the territories of their parents (e.g., Florida Scrub Jay

(*Aphelocoma coerulescens*) - Suh et al., 2020). Dispersal limitation is thus a complex and

difficult-to-quantify aspect of predictive distribution modeling that is nevertheless fundamental

for predicting the probability of range shifts with global change (reviewed in Zurell, 2017).

Coupling studies of distribution modeling with quantifiable measures of dispersal capacity is an

important next step toward making more informed predictions of species responses to changing

environments (Sousa et al., 2021; Travis et al., 2013; Urban et al., 2013).

Ultimately, extinction risk (and conversely, population viability) centers on declining

population and small population paradigms; population ecology. Population declines are driven

by deterministic (demographic, abiotic, and biotic) factors that cause reduction to small numbers

(e.g., the Allee effect), and also by stochastic factors, which dominate the extinction dynamics of

small populations (Caughley, 1994; Morris and Doak, 2002; Smith et al., 2021). The abiotic

shifts in conditions due to climate change and the resulting shifts in species distributions play a

role in both types of population dynamics. Shifting distributions can cause temporary or

permanent reductions in habitat area (Thomas et al., 2006; van Rees and Reed, 2018), lowering

carrying capacity and driving population declines at local or regional scales, or reducing

connectivity and creating isolated subpopulations that are more susceptible to small-population

dynamics (Anderson et al., 2009). The degree to which changing distributions result in

commonly observed conservation impacts depends on a diversity of species traits including

habitat preferences and associations, phenotypic plasticity, and current range and population size

(Jiguet et al., 2007; Visser, 2008). Notably, research based solely on changes in distributions

tends to predict widespread extinctions due to the aforementioned mechanisms, but plasticity,

dispersal, and adaptation appear to have been key for many species' resilience to past and current

climate change (Moritz and Agudo, 2013). For example, the Galapagos finches are a well-known

example of phenotypic plasticity in bill morphology in response to environmental change (Grant

and Grant, 2020; Grant and Weiner, 2017; Weiner, 1994).

Conservation action that occurs at small spatial scales requires deep natural history

knowledge for decision-making. In this section we described the domain of how ecological

theory informs spatial responses to land use and climate change. When designing research

projects that aim to predict future distributions under the umbrella of conservation ecology, we

urge modelers to deeply consider the ecology of focal taxa by referring to this section again in

the future, and the references cited within it. **Box 1** summarizes essential broad considerations

for predictive distribution modeling by the novice modeler.

335   A TECHNICAL OVERVIEW OF OCCURRENCE DATA FOR

336   PREDICTIVE DISTRIBUTION MODELING

Occurrence data through time is the foundation for applied predictive distribution modeling.

Here, we present technical aspects of different sources of avian occurrence data to inform which

questions can be answered with different data sets. To assess whether a change in distribution

has occurred or will occur, researchers require historical records to establish a baseline for

comparison, followed by surveys at one or more subsequent time points. Although birds are

among the best studied class of animals, early baseline inventories are still rare, and often consist

of field notes (Reznick et al., 1994), which present challenges particularly in reconstructing past

abundances (Shaffer et al., 1998). The aim of general collecting for birds by museums is to

document with vouchered specimens the complete taxonomic representation of a location, thus

supplying presence and absence information. A caveat to this is that complete information is

limited to small areas of intense investigation, thus limiting the widespread application of data

gathered by museums in presence/absence modelling (e.g., Loiselle et al., 2003). Although

general collecting for birds is no longer the norm at some museums (Ferguson, 2020; for reasons,

please see Remsen, 1995), museum records remain the best sources of historical occurrence data,

though they primarily supply presence or presence/absence data only and not abundance data

(Shaffer et al., 1998).

A major challenge for predictive models is rarity or sparseness of point occurrence data.

Even with many potential sources of occurrence data, knowledge of bird distributions still varies

greatly by species, habitat, and geography. For example, rare, nocturnal, and/or secretive species

and those in remote areas are difficult to detect and model accurately (MacKenzie et al., 2005;

Stralberg et al., 2015). Rarity is a common characteristic of species in diverse tropical

communities and remains a major challenge for constructing species distribution and

environmental niche models for tropical species (e.g., Marini et al., 2010).

In contrast to sparse information from the past, modern ornithologists benefit from

technological advances that offer unprecedented information on bird occurrence including from

devices that track individuals beyond specific field sites (satellite imagery - Borowicz et al.,

2018; and sophisticated automated biodiversity data collection - Kitzes and Schricker, 2019; for

birds within appropriate size ranges - McKinnon and Love, 2018; Sullivan et al., 2014, online

citizen science efforts - 2009). They improve our understanding of avian ecology by identifying

previously unknown species-habitat relationships (Jirinec et al., 2016), migratory routes (Jahn et

al., 2021, 2016; Stanley et al., 2015, 2012), foraging areas and behaviors (including accessing

microhabitat refuges - Wolfson et al., 2020), and wintering distributions (Renfrew et al., 2013).

These advances facilitate discovery of species responses to global change by revealing locations

outside of direct observation. Aside from emerging technologies, simpler monitoring approaches

implemented in the past at large scales have amassed rich long-term legacy datasets (e.g.,

breeding bird surveys, regional atlases, and the 5 million bird eggs housed at natural history

museums - Marini et al., 2020), and band recoveries across the globe continue to supply

information on bird population trends and distributions.

Decisions about which type of occurrence data is best for predictive modeling hinges on

the trade-off between addressing processes affecting the core distribution versus a more holistic

understanding including unique responses at range boundaries, during different life history

activities, or between different sex and age classes. Passive acoustic monitoring and long-term

studies remain promising approaches for capturing relative abundance data to estimate

population dynamics with respect to global change (Pérez-Granados et al., 2019; Pérez-Granados

and Schuchmann, 2020; Sugai et al., 2020). Trait databases are gaining momentum as sources of

functional diversity to test hypotheses under the Hutchinsonian niche concept (Gallagher et al.,

2020; Leclerc et al., 2020; Matuoka et al., 2020; Tobias et al., 2022). Exploring how functional

roles of species change across abiotic gradients can dramatically improve our understanding of

abiotic tolerances to more accurately predict range dynamics. For example, bill size has been

shown to be important for thermoregulation in birds (e.g., Danner and Greenberg, 2015) and a

study of functional trait structure along a tropical elevational gradient in Malaysian Borneo

linked larger bills in low elevation communities with thermal tolerance (Boyce et al., 2019).

Current shortfalls in avian occurrence data (summarized by Lees et al., 2020) require an

integrated approach to predicting bird distributions with global change. Although most efforts to

collect occurrence data are from small spatial areas (e.g., from individual field research sites) and

coarse resolutions (Table 1), this data contributes meaningfully to our understanding of species-

habitat relationships that can be applied to predictive modelling.

SELECTING APPROPRIATE EXPLANATORY VARIABLES

Predicting distributions relies on accurate assessments of life history requirements. In this

section, we synthesize the spatial and temporal resolution of explanatory variables commonly

used to match distributions with habitat and resource needs. Producing sound predictive models

requires ecologically relevant, suitable proxies for resource or habitat needs (e.g., Burns et al.,

2020; Randin et al., 2020). To scale up to entire distributions, there must be an underlying

knowledge of a species' life history that effectively links the lives of individuals to the

explanatory variable (or proxy variable) associated with site occupancy.

Predicting distributions influenced by changing food webs requires a breadth in

knowledge of life history information to predict the probability of species' ranges shifting

individually or with their entire biotic community. This task is complicated by the potentially

differing needs for each life history stage or demographic, which is often the case for birds (e.g.,

habitat requirements may differ between fledglings and adults). While main considerations are

the spatial and temporal resolutions of explanatory variables, biotic interactions can also play a

critical role (see Fig. 1) and may need to be considered on a taxon-specific basis.

CONSIDERING THE SPATIAL RESOLUTION OF EXPLANATORY VARIABLES

Selecting the correct spatial resolution of explanatory variables to use with occurrence data is

key to producing accurate assessments of species-habitat relationships. First, matching the spatial

resolution of explanatory variables with our understanding of the scale at which individuals

make occupancy decisions (e.g., whether they are micro- or macrohabitat based) is important for

making proper predictions of where organisms could move in the future. Second, matching

spatial resolution between explanatory variables and occurrence data prevents pseudoreplication.

Specifically, occurrence data may be pseudoreplicated if many samples occur in the same pixel

of explanatory data. Data spatial resolution also affects the possibility of model overfitting (i.e.,

when the importance of a narrow set of conditions is overestimated because occurrence data

come from a limited portion of conditions actually experienced by the taxon).

Remotely sensed spatial products (e.g., NDVI and BIOCLIM variables) can be proxies

for features of the environment being used by individuals at macrohabitat scales that require

occurrence data from a wider distribution. Remote sensing systems with finer spatial resolution,

for example light detection and ranging (LiDAR), have been used to predict occupancy by light-

demanding versus shade-tolerant plant species (Wüest et al., 2020), and direct measurements of

structural features of habitat (e.g., vertical distribution of forest canopy elements and foliage

density) used to predict occupancy (e.g., Goetz et al., 2010). Remotely sensed variables are

thought to reflect a contingency of resources assuming the data product conveys mutual

information on a shared spatiotemporal scale with the amount of food resources (Riotte-Lambert

& Matthiopoulos, 2020). There are some satisfying interpretations of the ecological relevance of

these remotely sensed products (Bridge et al., 2016 (EVI); Renfrew et al., 2013 (NDVI); see

Tøttrup et al., 2012 (drought)), but the typical purpose of using these large scale variables is for

spatial accuracy and not to infer causal relationships.

However, explanatory variables that reflect some ecological relevancy is important for

building accurate predictions of how taxa may respond to global change. Some remotely sensed

abiotic products do reflect a direct habitat resource or type. Habitat characteristics (i.e., land

cover) are most often identified with remotely sensed data, which depending on the sensor, can

have a variety of resolutions. Medium-high spatial resolutions (≤30m, e.g., Landsat, Sentinel) are

useful to pair with more precise occurrence data (e.g., GPS tags, point counts; Shirley et al.,

2013), whereas low spatial resolution (e.g., MODIS) can be useful for occurrence data from

geolocators or weather radar (Heim et al., 2020). The temporal resolution of remotely sensed

data is a limitation, especially when modeling the dynamic nature of migratory animals in

seasonal environments (MacPherson et al., 2018a; Roslin et al., 2021).

CONSIDERING THE TEMPORAL RESOLUTION OF EXPLANATORY VARIABLES

The temporal predictability of resources is important for predicting future distributions because

annual life history strategies are dependent upon correctly timing life cycle events with required

resources. For example, it is widely held that birds breeding at high latitudes must correctly time

the hatching of young with annual insect emergence to maximize fledging success, and early

arrival to breeding grounds may enhance reproductive success (Alerstam, 2011; Kokko et al.,

2006; Nilsson et al., 2013; Smith & Moore, 2005). Predictive distribution models are expected to

be the most accurate when species-habitat correlations are assessed using occurrence and

environmental data gathered from the same time-period. However, the temporal resolution of

occurrence data is much finer than that of environmental data; this can create a mismatch that

limits the questions that can be answered using current approaches. The habitat requirements for

fulfilling life history needs may vary depending upon the life history event (i.e., mate acquisition,

rearing young, molting, migrating) and the species life history strategy. Life history strategies

vary in birds from those that have evolved to rely on consistent resource availability (e.g., in the

case of dietary or habitat specialists that cannot live outside of narrow environmental conditions),

to periodic resource abundance (e.g., seasonal, annual, inter-annual weather patterns that drive

migratory or irruptive population movements), or irregular resource availability (e.g., in nomadic

taxa), and it is important to match locality data in time to environmental data for accurate

species-habitat assessment.

The temporal resolutions for widely used climate data varies, with trade-offs between

temporal and spatial resolution in addition to the state of products limiting the types of

distribution questions that they can inform (Table 1). We expand on this using the examples of

temperature and rainfall explanatory variables in Appendix 1 because these are the two most

widely used and generally important factors in species distribution modeling (Bradie and Leung,

2017).

To build robust predictive models, it is necessary to match the spatial and temporal

resolution of the occurrence data with that of the habitat characteristics. The temporal resolution

of remotely sensed data is a limitation, especially when modeling the dynamic nature of

migratory birds where dynamic seasonal habitat changes drive short-term habitat quality (e.g.,

seasonally flooded mudflat habitat for migrating shorebirds; Twedt, 2013). Google Earth Engine

(a javascript-based platform where one can write their own code to integrate many satellite

products and independently calculate indices such as NDVI) is a flexible tool that can facilitate

identifying the correct temporal resolution of land cover data to robustly test explanatory power

of land cover variables. Examples of data sets in Google Earth Engine include vegetation indices

(Landsat, ~2 weeks temporal resolution), products for creating land cover classifications (SPOT,

~1 month), and static land cover datasets (e.g., GlobCover, Cropland Data Layers, National Land

Cover Database; Table 1).

CONSIDERING BIOTIC FACTORS

Modeling future range shifts in response to global change rarely considers biotic factors. This

omission is the main reason why an assumption of SDMs - that the species is at equilibrium with

their environment - is rarely, if ever, met (Pearson and Dawson, 2003). If this assumption were

supported, the realized and fundamental niche would completely overlap as long as the species

could access all available suitable niche space (Figure 1). Consideration of biotic factors has the

potential to significantly enrich the field of predictive distribution modeling with more accurate

forecasts, and here we synthesize leading research contributing to this end at both the

intraspecific and interspecific level.

It is thought that biotic interactions vary along abiotic gradients such that they can either

enhance or reduce predicted ranges (Louthan et al., 2015). However, because capturing this

information requires both high spatial and temporal accuracy across all abiotic scenarios within

the realized niche, there are few examples to draw from. One promising approach to including

biotic interactions in predictive spatial modeling is the development of causal models that

estimate the influence of interspecific competition using co-occurrence data (e.g., Staniczenko et

al., 2017). The scope and strength of biotic factors may be correlated with abiotic pressures

(Louthan et al., 2015), and may differ depending upon which part of the range is being

considered. For example, much work has been done in the bird literature to improve our

understanding of species-habitat relationships through the study of hybrid zones, which typically

occupy only one part of a species range (eg., Taylor et al., 2014). Quantification of biotic factors,

like the ones described here, requires targeted species-specific research to test hypotheses within

the Eltonian and Hutchinsonian niche concepts. This research is mainly done within small spatial

extents and without consideration of temporal or spatial resolution (e.g., co-occurrence records

from different time periods or datasets with different spatial resolutions - Atauchi et al., 2018;

Palacio and Girini, 2018; see also König et al., 2021), but some research has been done at large

spatial extents and coarse resolutions (e.g., Acorn Woodpecker (*Melanerpes formicivorus*)

occurrence with Colombian Oaks in the Northern Andes - Freeman and Mason, 2015).

Intraspecific competition can drive density-dependent range shifts in migratory species

such that interannual selection of locations are dependent on group size and food availability

(Corre et al., 2020). Dispersal limitation can contribute to inadequacies in recruitment to new or

population sink sites, limiting population recovery or range shifts despite available habitat

(Palma et al., 2020; Zurell et al., 2016). The ability to update behaviors when circumstances

change (behavioral flexibility) may be key for driving range shifts or expansions under global

change (Blaisdell et al., 2021). Further, bird species capable of behavioral innovation (a.k.a.

plasticity) often have lower risk of extinction, as they are better able to adapt to changing

ecosystems and habitat destruction (Ducatez et al., 2020; Reed et al., 1999). However,

metapopulation dynamics can affect the introgression of adaptive traits such that maladaptive

traits restrict range shifts under global change (e.g., Garcia-R and Matzke, 2021; Lavretsky et al.,

2020). Each of these examples are implicitly expected to vary across abiotic gradients,

highlighting the importance of assessing the probability of site occupancy beyond the core of a

species distribution.

Perhaps more difficult is estimating the strength of interspecific biotic factors in shaping

distributions because this requires a breadth of natural history knowledge beyond that of the

species in question. However, interspecific biotic factors are thought to have strong influences on

the probability of site occupancy. Species may be excluded through predator/prey dynamics

where patches of habitat occupied by a species' predator may be less likely to be occupied by the

species being studied (Léandri-Breton and Bêty, 2020). Interspecific competition can also

exclude species and is often assessed using co-occurrence data (Elsen et al., 2017; Freeman et

al., 2016; Jankowski et al., 2010). Commensalisms, where one organism benefits from the

presence of another organism and the other unaffected, can also facilitate site occupancy (Aitken

and Martin, 2007). Including data on the presence of predators, competitors, and commensalisms

in SDMs can be a useful addition to improving models (Jankowski et al., 2013). Alternatively,

the spatial overlap with other taxa may have little influence if the focal species exhibits

behavioral avoidance (e.g., via interference competition - Jaeger, 1970; or predator-prey

dynamics - Lukas et al., 2021) or the spatio-temporal scope of data are not aligned.

The utility of SDMs to guide predictions of distributions under global change will

certainly be limited by the spatio-temporal resolution of environmental variables. Even when

researchers are mindful when selecting environmental data suited to answer questions about their

occurrence data (i.e., using ecologically relevant environmental and climate data across

biologically relevant space and timescales), a challenge remains in combining and contrasting

the Grinnellian niche concept (e.g., climate variables) against other niche concepts that require

finer temporal and spatial data (e.g., density of interacting species). This problem can further

complicate predictive models if site occupancy is dependent on time-lags, as is the case in some

migratory species (Bridge et al., 2016), and is explored further in the following section.

AN OVERVIEW OF ANALYTICAL METHODS

Predictive models of species distributions can be divided into correlative versus mechanistic

approaches (Figure 3). Correlative models elucidate relationships between species occurrence

and spatially explicit explanatory variables and can be viewed as hypothesis-generating tools.

Data to parameterize correlative models are readily available (e.g., correlating occurrence data

with land-cover or weather data as described above), making them the dominant approach to

predictive distribution modeling. Mechanistic models (a.k.a. causal models) are hypothesis-

testing tools that incorporate physiological tolerance to predict where species will be capable of

persisting within their physiological limits (Guisan and Zimmermann, 2000); data to support

these approaches is more limited. Here, we summarize popular SDM tools with important

information to help novice modelers identify which tool might be appropriate given their data

and question. We provide examples that students can refer to for accessing more detailed

methodologies.

Range estimates from correlative SDMs are valuable, helping us better understand

species and their sensitivities to global change. MAXENT is one of the most popular SDM

approaches, widely used for its ability to use presence-only datasets where true absence or non-

detection data are typically hard to find (Elith and Leathwick, 2007; Merow et al., 2013).

MAXENT has been useful for modeling distributions of rare or cryptic species due to its

apparent high performance with small or incomplete datasets (Aubry et al., 2017; Pearson et al.,

2007; Phillips et al., 2006; Wisz et al., 2008; but see also Marini et al., 2009). Boosted regression

trees (BRTs) are one of many machine learning techniques well-suited to modeling complex

ecological data because they can handle different types of predictor variables, fit complex

nonlinear relationships, accommodate missing data and automatically handle interaction effects

between predictors (Elith et al., 2008; Graham et al., 2008). BRTs have been used to predict bird

distributions using occurrence data for land birds (Veloz et al., 2015), waterfowl (Barker et al.,

2014), shorebirds (Dalgarno et al., 2017), seabirds (Oppel et al., 2012), and owls (Domahidi et

al., 2019). Hierarchical Bayesian models can also integrate various occurrence data types (e.g.,

presence-absence, presence-only, count data) to create reliable spatio-temporal distribution

models (Hefley and Hooten, 2015). Using continuous spatial predictor variables such as land

cover, the abundance and distribution of species can be predicted over a continuous latent

surface (Chakraborty et al., 2010), which has been useful with waterfowl aerial survey data

(Herbert et al., 2021, 2018). Generalized linear models (GLMs) and generalized additive models

(GAMs) can be employed to understand distributions across multiple environmental conditions

and also conceptualize patterns in distributions of abundances across environmental gradients

(Augustin et al., 1996; Smith & Edwards, 2021). GAMs are used to model nonlinear

relationships (e.g., Maggini et al., 2011). One drawback to GAMs, however, is that evaluation of

response shapes (e.g., optima, skewness) is done through visual inspection rather than

statistically. Huisman-Olff-Fresco (HOF) models are used to predict occupancy along abiotic

gradients and are particularly useful in evaluating alternative shapes of species responses to

gradients, for example, if changes in abundance do not exhibit the widely assumed symmetric

Gaussian functions, but instead show asymmetric, threshold-like changes in occupancy over

space (Huisman et al., 1993). HOF response curves can be used to highlight distinct distribution

patterns such as species replacements, allowing inference of potential biotic interactions that are

otherwise difficult to measure (Jansen and Oksanen, 2013). Species climate response surfaces

(CRSs) model the probability of occurrence using bioclimatic variables (Huntley et al., 1995).

Given that explanatory variables in correlative SDMs cannot be used to infer causation, the CRS

approach can accommodate interactions between abiotic variables to predict distributions under

future climate scenarios (Huntley et al., 2006). The choice of predictive SDM technique and

quality of SDM product are based predominantly on the type of occurrence data available and

secondarily on spatiotemporal matching of explanatory variables.

Mechanistic approaches to SDMs predict areas where physico-chemical processes meet

life history needs (Kearney, 2006; Kearney & Porter, 2009). For example, a biophysical model

for the endangered Night Parrot (*Pezoporus occidentalis*) in Australia shows how air temperature

increases of 3 °C would lead to lethal levels of dehydration (Kearney et al., 2016). Bayesian

networks (BNs) are another type of mechanistic model that go beyond species-habitat

correlations by also considering processes that influence occupancy across space and time (i.e.,

site access and selection - Jones, 2001). Originally employed to model human judgement (Pearl,

1985), BNs have only recently been adapted for predictive SDMs (MacPherson et al., 2018b;

Staniczenko et al., 2017). Mechanistic models are informed by causal relationships based on

empirical research or expert knowledge. The latter are referred to as belief networks (Drew and

Collazo, 2014; MacPherson et al., 2018b). Mechanistic models enable the identification of the

most important variables driving distribution patterns through mapping the fundamental niche,

which is helpful to inform conservation and management decisions when circumstances change.

Ensemble modeling strategies, in which the predictions of multiple approaches are

combined or used simultaneously, are often suggested to better encompass the range of

uncertainty in prediction (Araújo and New, 2007). This method accounts for the fact that model

choice is often the greatest source of quantifiable uncertainty in species distribution modeling

and reduces sources of bias from the use of a single algorithm (Dormann et al., 2008; Jarnevich

et al., 2017). Implementing ensembles of five or more of the algorithms and approaches

described above has been greatly facilitated by software packages like BIOMOD (Thuiller et al.,

2009) and VisTrails SAHM (Morisette et al., 2013), which bring methods together into a single

analytical environment. Ensembles can be used to assess a range of potential projected outcomes,

forming a "bounding box" or "consensus" across algorithm predictions (Araújo and New, 2007).

Analysis with model ensembles typically involves scaling and averaging model outputs and,

often weighting these by some measurement of model performance such as "area under the

curve" (AUC) score.

The ongoing "big data revolution" in many fields including ornithology (La Sorte et al.,

2018) has increased use of artificial intelligence for data handling and analysis (Xia et al., 2020),

especially in correlative distribution modeling. Machine learning algorithms like MAXENT

(Merow et al., 2013), random forests (Mi et al., 2017), neural networks (e.g., Manel et al., 1999),

deep learning (Benkendorf and Hawkins, 2020) and boosted regression trees (Elith et al., 2006)

are now commonly applied SDMs, and are valued for prediction and interpolation. They are

excellent tools for insight into potential future shifts under global change (Elith, 2017).

Conservation plans rely on SDMs for future species distributions, despite the small, but

growing, body of literature on ways to incorporate climate change into conservation planning

(Alexander et al., 2017; Hole et al., 2011; Jones & Cheung, 2015; Loyola et al., 2013; Nakao et

al., 2013; Terribile et al., 2012; Willis et al., 2009). The predictive power of SDMs for informing

conservation decisions remains limited by a lack of tool capability to calculate confidence

intervals around predictions (Marini et al., 2010), a lack of validation to reduce uncertainties and

the need for agreed-upon standards for guiding model building (Araújo et al., 2019, 2005). Using

correlative predictive models is likely to overestimate range shifts and extinction risk due to the

violations of common assumptions in correlative distribution modeling. For example, SDMs tend

to violate the assumption that the individual species is currently at equilibrium within its

environment (Early and Sax, 2014; Sax et al., 2013) and do not take into account species

interactions (Pearson and Dawson, 2003). Despite these criticisms, SDMs remain the "best"

644 widely accessible approach currently in use for identifying potential future habitat-area (Tingley

645 et al., 2014).

647 **DISCUSSION**

648 To invite novel perspectives into the field of predictive distribution modelling, we reviewed the

649 theory underpinning how global change shapes range dynamics and outlined the technical details

650 of data sources and analytic approaches needed for creating sound tests of ecological theory to

651 improve predictive models. Predicting where species will live in an uncertain future is a central

652 goal of modern ecology. This effort has recast the basic research question of what limits

653 distribution of species – why a species lives *here* but not *there* – as an applied question. Species

654 are already on the move in response to recent decades of global change (and, depending on the

655 region, predicted trends of reduced or increased rainfall; Figure 2), supporting long-held theory

656 that climate limits species' geographic ranges (Tingley et al., 2009). However, the match

657 between species' distributions and climatic conditions is often weak (Chapman, 2010; Suggitt et

658 al., 2012); species vary considerably in their observed responses to recent climate change

659 (Lehmann et al., 2020; Mamantov et al., 2021). These observations indicate that we are still in

660 the early stages of ecological forecasting. To guide future research, we identified three core

661 motivating questions that build from our technical introduction as the answers to these questions

662 are particularly likely to generate important advancements in the field of predictive distribution

663 modeling.

664 *1. Is dispersal a rate-limiting step in range expansions?* The proximate driver of range shifts is

665 dispersal; ranges expand when individuals move beyond the existing range limit. Dispersal

666 constraints are thus one obvious explanation for cases where the rate of species' observed range

shifts are failing to track with the rate of changing climate. Birds have incredible capacities for

dispersal (e.g., Slager, 2020), making it tempting to underestimate the possible role for dispersal

constraints in avian range shifts, including unexplored factors such as site fidelity. However,

even bird species often show extreme limitations in dispersal, suggesting that longer-distance

range shifts (e.g., across latitudes) may be slow, particularly in heterogeneous landscapes, or

when species show high specialization to their associated habitats. In the temperate zone, even

some long-distance migrants may show strong site fidelity to breeding (and wintering) territories

(Winger et al., 2019), which suggests the possibility that rapid range shifts may be difficult to

achieve. Lastly, dispersal may be possible but may introduce a new trade-off with other

components of the annual life cycle, such as migration. For example, the ability of boreal

breeding birds to take advantage of newly available forest habitat in the Arctic may be

constrained by a trade-off with ever-increasing migration distances. We can improve our

understanding of dispersal limitation by creatively pairing historical with contemporary datasets

to identify patterns of dispersal limitation in places where global change has already altered

species distributions.

2. Are species particularly rare at their range limits? Ecological theory assumes that species are

most abundant at the center or core of their range and become progressively rarer towards their

range limits (the "abundant-center hypothesis"; Brown, 1984). However, empirical patterns of

abundance often fail to fit these expectations. Whether the abundant-center hypothesis generally

holds or not is now a matter of debate (e.g., Dallas et al., 2020), and the answer may have

consequences for rates of species' range shifts. For example, low abundances at range edges

could lead to slower rates of range shifts than if abundance distributions are more uniform across

a core-to-limit transect through a species' geographic range. This could be due to simple numeric

effects – range shifts are easier when there are more individuals at the range edge to start with –

or to genetic effects. That is, low abundances (and hence lower genetic diversity) could reduce

local rates of adaptation at range edges, with gene flow from the abundant range center

swamping any effect of local adaptation at the range edge that would facilitate range expansion

when conditions change (Haldane, 1956; but see Kottler et al., 2021). Pairing occurrence data

with natural history knowledge of species expanding their ranges has the potential to improve

our understanding of the mechanisms that promote expansion from original range boundaries

(e.g., the tolerance niche of the Asian Openbill (*Anastomus oscitans*) - Lei and Liu, 2021; and

climate matching in the European Bee-eater (*Merops apiaster*) - Stiels et al., 2021). Additionally,

assessing occupancy from the perspective of abiotic gradients rather than focal taxa would

improve our ability to identify significant geographic barriers to range shifts with climate

change.

3. How does land use change interact with climate change? Like most of the range shift

literature, we have focused in this review on predicting species' responses to changes in climate

per se. But species are shifting (or failing to shift) their ranges in landscapes that are increasingly

dominated by human activities (e.g., Fumy and Fartmann, 2021). Studies that simultaneously

incorporate both land use and climate change as drivers of distributional change are few in

number but hold great promise for several reasons. First, land use change is an obvious driver of

species' distributions; many species simply do not persist in landscapes with extreme levels of

human control. Second, appropriate habitat for most species in human-dominated landscapes

consists of habitat patches of varying sizes with varying levels of connectivity (e.g., Neilan et al.,

2019). This fact elevates the role of dispersal in determining whether patches (and scaled-up to a

larger geographic scale, regions) are occupied by a given species. Third, habitat change directly

affects local-scale climate. Fragmented forests, for example, average substantially warmer and

drier than primary forests (Kapos, 1989; Nunes et al., 2022), and "urban heat islands" alter local

temperatures that are known to affect bird distributions (in migrants - Bonnet-Lebrun et al.,

2020; and residents - Latimer and Zuckerberg, 2021). Hence, land use change can act as a

multiplier of temperature effects on species. Here we emphasize the importance of high spatial

and temporal resolution datasets for identifying mechanisms of global change that affect

individual organisms or populations. In light of this, mechanistic (rather than correlative) models

are likely to be better suited to identifying the interaction of land use and climate change when

environmental conditions can be linked to the biological processes of individuals (or the loss of

migration with increased urbanization - Bonnet-Lebrun et al., 2020; e.g., thermoregulatory

behaviors of individuals in unshaded desert areas - van de Ven et al., 2019).

The consequences of climate change for species are potentially severe, with widespread

predictions of extinctions (Thomas et al., 2006), yet the application of predictive models to

conservation and management are still limited (but see Casazza et al., 2021). Some evidence

suggests these dire predictions may be coming true. For example, warming temperatures have

led to local extinctions in mountaintop communities in southeastern Peru as species shift their

geographic ranges to track climate, a potential harbinger of the possible extinction of high

elevation tropical species (Feeley et al., 2012; Freeman et al., 2018; Rehm and Feeley, 2016).

Yet species may be more resilient than models assume; for example, genetic data indicate that

many species were able to persist through dramatic climate fluctuations in the Pleistocene

(Bocalini et al., 2021; Song et al., 2020; Wogan et al., 2020). Application of predictive

distribution modes in conservation and management should become more widespread by

including the development of tools for calculating confidence intervals (Marini et al., 2010) and

validating models (i.e., comparing past to present distributions with respect to global change) for

correlative models, and increasing the use of mechanistic models.

Humans have already made Earth hotter than it has been since before the Pleistocene (~ 2

mya). This rapid change in Earth's climate has set species on the move and led to a plethora of

scientific research aimed at predicting where species will live in the coming decades as warming

continues. These predictions, though frequently made, are seldom tested (Tayleur et al., 2015;

but see Tingley et al., 2009; Wilson et al., 2018). Without this crucial step of model validation, it

is impossible to assess the usefulness of predictive models. Here, we have presented theory and

highlighted a range of data sources and analytical approaches that can be used to predict species'

geographic responses to climate change. Ever-greater computational power, combined with

increasingly large datasets of species occurrence and landscape covariates, permit the use of

greater complexity in models (e.g., Lurgi et al., 2015). However, more complex does not

necessarily equate to better. The litmus test for any predictive model is how well it predicts

reality. To this end, we advocate for an increasing focus on collecting empirical data that

matches the spatio-temporal resolution of occurrence data with environmental variables, tests

hypotheses beyond the Grinnellian niche concept, and directly evaluates model predictions (e.g.,

using mechanistic SDMs or resurveys). For example, the long-term predictions from models

whose predictions are not met over the short-term are unlikely to be helpful (see Willis et al.,

2009). Determining species' resiliency will depend on accurate estimates of the fundamental

niche including which attributes make some species more vulnerable than others, which abiotic

gradients are the most important to consider for the promotion of population persistence, and

including biotic interactions in predictive, validated models. Our review aims to support the

ongoing pursuit of more meaningful predictive distribution models under land use and climate

change that will be of great near-term utility.

ACKNOWLEDGEMENTS

The authors declare that we have no sources of conflict of interest affecting the objectivity of the

presented topic. This manuscript is the result of a synthesis on how ornithologists predict bird

distributions under global change that was presented in a series of talks at the North American

Ornithological Congress, 2020. We sincerely thank the organizers of this conference for their

support throughout the COVID-19 pandemic. Thank you also to the participants of the Predicting

Bird Distributions Under Global Change symposium, including William Lewis. This research

was funded in part by a U.S. Geological Survey Northwest Climate Adaptation Science Center

award G17AC000218 to C.B. van Rees. The manuscript was approved by the Director of the

Louisiana State University Agricultural Center as manuscript number 20213-XXX-XXXX. The

authors would also like to thank (in alphabetical order by last name) Anna Borne, Nicholas

Mason, Quinn McCallum, Diego Ocampo Vargas, Samantha Rutledge, Subir Shakya, Philip

Stouffer, David Vander Pluym, and Brenna Wells who provided important discussion of the

topics presented in this manuscript during its development.

LITERATURE CITED

Aitken, K.E.H., Martin, K., 2007. The importance of excavators in hole-nesting communities:
availability and use of natural tree holes in old mixed forests of western Canada. J
Ornithol 148, 425–434. https://doi.org/10.1007/s10336-007-0166-9
Alerstam, T., 2011. Optimal bird migration revisited. J Ornithol 152, 5–23.
https://doi.org/10.1007/s10336-011-0694-1

Alexander, J.D., Stephens, J.L., Veloz, S., Salas, L., Rousseau, J.S., Ralph, C.J., Sarr, D.A.,
2017. Using regional bird density distribution models to evaluate protected area
networks and inform conservation planning. Ecosphere 8, e01799.
https://doi.org/10.1002/ecs2.1799
Anderson, B. j, Akçakaya, H. r, Araújo, M. b, Fordham, D. a, Martinez-Meyer, E., Thuiller, W.,
Brook, B. w, 2009. Dynamics of range margins for metapopulations under climate
change. Proceedings of the Royal Society B: Biological Sciences 276, 1415–1420.
https://doi.org/10.1098/rspb.2008.1681
Angilletta Jr, M.J., Angilletta, M.J., 2009. Thermal adaptation: a theoretical and empirical
synthesis.
Angilletta, M.J., Sears, M.W., Pringle, R.M., 2009. Spatial dynamics of nesting behavior: Lizards
shift microhabitats to construct nests with beneficial thermal properties. Ecology 90,
2933–2939. https://doi.org/10.1890/08-2224.1
Araújo, M.B., Anderson, R.P., Márcia Barbosa, A., Beale, C.M., Dormann, C.F., Early, R.,
Garcia, R.A., Guisan, A., Maiorano, L., Naimi, B., O'Hara, R.B., Zimmermann, N.E.,
Rahbek, C., 2019. Standards for distribution models in biodiversity assessments.
Science Advances 5, eaat4858. https://doi.org/10.1126/sciadv.aat4858
Araújo, M.B., New, M., 2007. Ensemble forecasting of species distributions. Trends in Ecology
& Evolution 22, 42–47. https://doi.org/10.1016/j.tree.2006.09.010
Araújo, M.B., Pearson, R.G., Thuiller, W., Erhard, M., 2005. Validation of species–climate
impact models under climate change. Global Change Biology 11, 1504–1513.
https://doi.org/10.1111/j.1365-2486.2005.01000.x
Atauchi, P.J., Peterson, A.T., Flanagan, J., 2018. Species distribution models for Peruvian
plantcutter improve with consideration of biotic interactions. Journal of Avian Biology 49,
jav-01617. https://doi.org/10.1111/jav.01617
Aubry, K.B., Raley, C.M., McKelvey, K.S., 2017. The importance of data quality for generating
reliable distribution models for rare, elusive, and cryptic species. PLOS ONE 12,
e0179152. https://doi.org/10.1371/journal.pone.0179152
Augustin, N.H., Mugglestone, M.A., Buckland, S.T., 1996. An autologistic model for the spatial
distribution of wildlife. Journal of Applied Ecology 33, 339–347.
https://doi.org/10.2307/2404755
Austin, M.P., 2007. Species distribution models and ecological theory: A critical assessment and
some possible new approaches. Ecological Modelling 200, 1–19.
https://doi.org/10.1016/j.ecolmodel.2006.07.005
Austin, M.P., 2005. Vegetation and environment: discontinuous and continuous. Vegetation
ecology 52–84.
Bani, L., Luppi, M., Rocchia, E., Dondina, O., Orioli, V., 2019. Winners and losers: How the
elevational range of breeding birds on Alps has varied over the past four decades due to
climate and habitat changes. Ecology and Evolution 9, 1289–1305.
https://doi.org/10.1002/ece3.4838
Barker, N.K.S., Cumming, S.G., Darveau, M., 2014. Models to predict the distribution and
abundance of breeding ducks in Canada. ACE 9, art7. https://doi.org/10.5751/ACE-
00699-090207
Benkendorf, D.J., Hawkins, C.P., 2020. Effects of sample size and network depth on a deep
learning approach to species distribution modeling. Ecological Informatics 60, 101137.
https://doi.org/10.1016/j.ecoinf.2020.101137
Blaisdell, A., Seitz, B., Rowney, C., Folsom, M., MacPherson, M., Deffner, D., Logan, C., 2021.
Do the more flexible individuals rely more on causal cognition? Observation versus
intervention in causal inference in great-tailed grackles. Peer Community Journal 1, e50.
https://doi.org/10.24072/pcjournal.44

Block, S., Levine, J.M., 2021. How Dispersal Evolution and Local Adaptation Affect the Range
Dynamics of Species Lagging Behind Climate Change. The American Naturalist 197,
E173–E187. https://doi.org/10.1086/714130
Bocalini, F., Bolívar-Leguizamón, S.D., Silveira, L.F., Bravo, G.A., 2021. Comparative
phylogeographic and demographic analyses reveal a congruent pattern of sister
relationships between bird populations of the northern and south-central Atlantic Forest.
Molecular Phylogenetics and Evolution 154, 106973.
https://doi.org/10.1016/j.ympev.2020.106973
Bonnet-Lebrun, A.-S., Manica, A., Rodrigues, A.S.L., 2020. Effects of urbanization on bird
migration. Biological Conservation 244, 108423.
https://doi.org/10.1016/j.biocon.2020.108423
Borowicz, A., McDowall, P., Youngflesh, C., Sayre-McCord, T., Clucas, G., Herman, R., Forrest,
S., Rider, M., Schwaller, M., Hart, T., Jenouvrier, S., Polito, M.J., Singh, H., Lynch, H.J.,
2018. Multi-modal survey of Adélie penguin mega-colonies reveals the Danger Islands
as a seabird hotspot. Scientific Reports 8, 3926. https://doi.org/10.1038/s41598-018-
22313-w
Boyce, A.J., Shakya, S., Sheldon, F.H., Moyle, R.G., Martin, T.E., 2019. Biotic interactions are
the dominant drivers of phylogenetic and functional structure in bird communities along a
tropical elevational gradient. The Auk 136. https://doi.org/10.1093/auk/ukz054
Bradie, J., Leung, B., 2017. A quantitative synthesis of the importance of variables used in
MaxEnt species distribution models. Journal of Biogeography 44, 1344–1361.
https://doi.org/10.1111/jbi.12894
Brawn, J.D., Benson, T.J., Stager, M., Sly, N.D., Tarwater, C.E., 2017. Impacts of changing
rainfall regime on the demography of tropical birds. Nature Climate Change 7, 133–136.
https://doi.org/10.1038/nclimate3183
Bridge, E.S., Ross, J.D., Contina, A.J., Kelly, J.F., 2016. Do molt-migrant songbirds optimize
migration routes based on primary productivity? Behav Ecol 27, 784–792.
https://doi.org/10.1093/beheco/arv199
Brown, J.H., 1984. On the Relationship between Abundance and Distribution of Species. The
American Naturalist 124, 255–279. https://doi.org/10.1086/284267
Burner, R.C., Styring, A.R., Rahman, M.A., Sheldon, F.H., 2019. Occupancy patterns and upper
range limits of lowland Bornean birds along an elevational gradient. Journal of
Biogeography 46, 2583–2596. https://doi.org/10.1111/jbi.13691
Burns, P., Clark, M., Salas, L., Hancock, S., Leland, D., Jantz, P., Dubayah, R., Goetz, S.J.,
2020. Incorporating canopy structure from simulated GEDI lidar into bird species
distribution models. Environ. Res. Lett. 15, 095002. https://doi.org/10.1088/1748-
9326/ab80ee
Busby, J.R., 1991. BIOCLIM – A bioclimate analysis and prediction system, in: Margules, C.R.,
Austin, M.P. (Eds.), Nature Conservation: Cost Effective Biological Surveys and Data
Analysis. CSIRO, Melbourne, Australia, pp. 64–68.
Capurucho, J.M.G., Ashley, M.V., Tsuru, B.R., Cooper, J.C., Bates, J.M., 2020. Dispersal ability
correlates with range size in Amazonian habitat-restricted birds. Proceedings of the
Royal Society B: Biological Sciences 287, 20201450.
https://doi.org/10.1098/rspb.2020.1450
Caruso, C.M., Martin, R.A., Sletvold, N., Morrissey, M.B., Wade, M.J., Augustine, K.E., Carlson,
S.M., MacColl, A.D.C., Siepielski, A.M., Kingsolver, J.G., 2017. What Are the
Environmental Determinants of Phenotypic Selection? A Meta-analysis of Experimental
Studies. The American Naturalist 190, 363–376. https://doi.org/10.1086/692760
Casazza, G., Abeli, T., Bacchetta, G., Dagnino, D., Fenu, G., Gargano, D., Minuto, L.,
Montagnani, C., Orsenigo, S., Peruzzi, L., Varaldo, L., Rossi, G., 2021. Combining
conservation status and species distribution models for planning assisted colonisation

under climate change. Journal of Ecology 109, 2284–2295. https://doi.org/10.1111/1365-
2745.13606

Caughley, G., 1994. Directions in conservation biology. Journal of Animal Ecology 63, 215–244.
https://doi.org/10.2307/5542

Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., Palmer, T.M., 2015.
Accelerated modern human–induced species losses: Entering the sixth mass extinction.
Science Advances 1, e1400253. https://doi.org/10.1126/sciadv.1400253

Chakraborty, A., Gelfand, A.E., Wilson, A.M., Latimer, A.M., John A. Silander, J., 2010.
Modeling large scale species abundance with latent spatial processes. The Annals of
Applied Statistics 4, 1403–1429. https://doi.org/10.1214/10-AOAS335

Chapman, D.S., 2010. Weak climatic associations among British plant distributions. Global
Ecology and Biogeography 19, 831–841. https://doi.org/10.1111/j.1466-
8238.2010.00561.x

Claramunt, S., Derryberry, E.P., Remsen, J.V., Brumfield, R.T., 2012. High dispersal ability
inhibits speciation in a continental radiation of passerine birds. Proceedings of the Royal
Society B: Biological Sciences 279, 1567–1574. https://doi.org/10.1098/rspb.2011.1922

Cooper, N.W., Marra, P.P., 2020. Hidden Long-Distance Movements by a Migratory Bird.
Current Biology 30, 4056-4062.e3. https://doi.org/10.1016/j.cub.2020.07.056

Corre, M.L., Dussault, C., Côté, S.D., 2020. Where to spend the winter? The role of intraspecific
competition and climate in determining the selection of wintering areas by migratory
caribou. Oikos 129, 512–525. https://doi.org/10.1111/oik.06668

Cowles, R.B., Bogert, C.M., 1944. A preliminary study of the thermal requirements of desert
reptiles. Bulletin of the American Museum of Natural History 83, 261–296.

Dalgarno, S., Mersey, J.E., Gedalof, Z., Lemon, M., 2017. Species-environment associations
and predicted distribution of Black Oystercatcher breeding pairs in Haida Gwaii, British
Columbia, Canada. ACE 12, art9. https://doi.org/10.5751/ACE-01094-120209

Dallas, T.A., Santini, L., Decker, R., Hastings, A., 2020. Weighing the Evidence for the
Abundant-Center Hypothesis. Biodiv. Inf. 15, 81–91.
https://doi.org/10.17161/bi.v15i3.11989

Danner, R.M., Greenberg, R., 2015. A critical season approach to Allen's rule: Bill size declines
with winter temperature in a cold temperate environment. Journal of Biogeography 42,
114–120. https://doi.org/10.1111/jbi.12389

De Frenne, P., Lenoir, J., Luoto, M., Scheffers, B.R., Zellweger, F., Aalto, J., Ashcroft, M.B.,
Christiansen, D.M., Decocq, G., De Pauw, K., Govaert, S., Greiser, C., Gril, E., Hampe,
A., Jucker, T., Klinges, D.H., Koelemeijer, I.A., Lembrechts, J.J., Marrec, R., Meeussen,
C., Ogée, J., Tyystjärvi, V., Vangansbeke, P., Hylander, K., 2021. Forest microclimates
and climate change: Importance, drivers and future research agenda. Global Change
Biology 27, 2279–2297. https://doi.org/10.1111/gcb.15569

Domahidi, Z., Shonfield, J., Nielsen, S.E., Spence, J.R., Bayne, E.M., 2019. Spatial distribution
of the Boreal Owl and Northern Saw-whet Owl in the boreal region of Alberta, Canada.
ACE 14, art14. https://doi.org/10.5751/ACE-01445-140214

Dormann, C.F., Purschke, O., Márquez, J.R.G., Lautenbach, S., Schröder, B., 2008.
Components of Uncertainty in Species Distribution Analysis: A Case Study of the Great
Grey Shrike. Ecology 89, 3371–3386. https://doi.org/10.1890/07-1772.1

Drew, A., Collazo, J., 2014. Bayesian networks as a framework to step-down and support
Strategic Habitat Conservation of data-poor species: A case study with King Rail (Rallus
elegans) in Eastern North Carolina and Southeastern Virginia. US Fish and Wildlife
Service, Raleigh, NC. https://doi.org/10.13140/2.1.4430.5920

Ducatez, S., Sol, D., Sayol, F., Lefebvre, L., 2020. Behavioural plasticity is associated with
reduced extinction risk in birds. Nature Ecology & Evolution 4, 788–793.
https://doi.org/10.1038/s41559-020-1168-8

Dunn, P.O., Møller, A.P., 2019. Effects of climate change on birds. Oxford University Press, New York, NY.

Dvorský, M., Macek, M., Kopecký, M., Wild, J., Doležal, J., 2017. Niche asymmetry of vascular plants increases with elevation. Journal of Biogeography 44, 1418–1425. https://doi.org/10.1111/jbi.13001

Early, R., Sax, D.F., 2014. Climatic niche shifts between species' native and naturalized ranges raise concern for ecological forecasts during invasions and climate change. Global Ecology and Biogeography 23, 1356–1365. https://doi.org/10.1111/geb.12208

Elith, J., 2017. Predicting Distributions of Invasive Species, in: Invasive Species: Risk Assessment and Management. Cambridge University Press, Cambridge, UK, pp. 93–129.

Elith, J., H. Graham, C., P. Anderson, R., Dudík, M., Ferrier, S., Guisan, A., J. Hijmans, R., Huettmann, F., R. Leathwick, J., Lehmann, A., Li, J., G. Lohmann, L., A. Loiselle, B., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., McC. M. Overton, J., Townsend Peterson, A., J. Phillips, S., Richardson, K., Scachetti-Pereira, R., E. Schapire, R., Soberón, J., Williams, S., S. Wisz, M., E. Zimmermann, N., 2006. Novel methods improve prediction of species' distributions from occurrence data. Ecography 29, 129–151. https://doi.org/10.1111/j.2006.0906-7590.04596.x

Elith, J., Leathwick, J., 2007. Predicting species distributions from museum and herbarium records using multiresponse models fitted with multivariate adaptive regression splines. Diversity and Distributions 13, 265–275. https://doi.org/10.1111/j.1472-4642.2007.00340.x

Elith, J., Leathwick, J.R., Hastie, T., 2008. A working guide to boosted regression trees. Journal of Animal Ecology 77, 802–813. https://doi.org/10.1111/j.1365-2656.2008.01390.x

Elsen, P.R., Tingley, M.W., Kalyanaraman, R., Ramesh, K., Wilcove, D.S., 2017. The role of competition, ecotones, and temperature in the elevational distribution of Himalayan birds. Ecology 98, 337–348. https://doi.org/10.1002/ecy.1669

Elton, C., 1927. Animal ecology. Sidgwick & Jackson, London.

Faaborg, J., Holmes, R.T., Anders, A.D., Bildstein, K.L., Dugger, K.M., Gauthreaux, S.A., Heglund, P., Hobson, K.A., Jahn, A.E., Johnson, D.H., Latta, S.C., Levey, D.J., Marra, P.P., Merkord, C.L., Nol, E., Rothstein, S.I., Sherry, T.W., Sillett, T.S., Thompson, F.R., Warnock, N., 2010. Recent advances in understanding migration systems of New World land birds. Ecological Monographs 80, 3–48. https://doi.org/10.1890/09-0395.1

Feeley, K.J., Rehm, E.M., Machovina, B., 2012. perspective: The responses of tropical forest species to global climate change: acclimate, adapt, migrate, or go extinct? Frontiers of Biogeography 4. https://doi.org/10.21425/F5FBG12621

Ferguson, A.W., 2020. On the role of (and threat to) natural history museums in mammal conservation: an African small mammal perspective. fozo.1 69, 20028.1. https://doi.org/10.25225/jvb.20028

Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37, 4302–4315. https://doi.org/10.1002/joc.5086

Freeman, B.G., Freeman, A.M.C., Hochachka, W.M., 2016. Asymmetric interspecific aggression in New Guinean songbirds that replace one another along an elevational gradient. Ibis 158, 726–737. https://doi.org/10.1111/ibi.12384

Freeman, B.G., Mason, N.A., 2015. The geographic distribution of a tropical montane bird is limited by a tree: Acorn Woodpeckers (Melanerpes formicivorus) and Colombian oaks (Quercus humboldtii) in the Northern Andes. PLOS ONE 10, e0128675. https://doi.org/10.1371/journal.pone.0128675

Freeman, B.G., Scholer, M.N., Ruiz-Gutierrez, V., Fitzpatrick, J.W., 2018. Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. PNAS 115, 11982–11987. https://doi.org/10.1073/pnas.1804224115

Fumy, F., Fartmann, T., 2021. Climate and land-use change drive habitat loss in a mountain bird species. Ibis 163, 1189–1206. https://doi.org/10.1111/ibi.12954

Gallagher, R.V., Falster, D.S., Maitner, B.S., Salguero-Gómez, R., Vandvik, V., Pearse, W.D., Schneider, F.D., Kattge, J., Poelen, J.H., Madin, J.S., Ankenbrand, M.J., Penone, C., Feng, X., Adams, V.M., Alroy, J., Andrew, S.C., Balk, M.A., Bland, L.M., Boyle, B.L., Bravo-Avila, C.H., Brennan, I., Carthey, A.J.R., Catullo, R., Cavazos, B.R., Conde, D.A., Chown, S.L., Fadrique, B., Gibb, H., Halbritter, A.H., Hammock, J., Hogan, J.A., Holewa, H., Hope, M., Iversen, C.M., Jochum, M., Kearney, M., Keller, A., Mabee, P., Manning, P., McCormack, L., Michaletz, S.T., Park, D.S., Perez, T.M., Pineda-Munoz, S., Ray, C.A., Rossetto, M., Sauquet, H., Sparrow, B., Spasojevic, M.J., Telford, R.J., Tobias, J.A., Violle, C., Walls, R., Weiss, K.C.B., Westoby, M., Wright, I.J., Enquist, B.J., 2020. Open Science principles for accelerating trait-based science across the Tree of Life. Nature Ecology & Evolution 4, 294–303. https://doi.org/10.1038/s41559-020-1109-6

Garcia-R, J.C., Matzke, N.J., 2021. Trait-dependent dispersal in rails (Aves: Rallidae): Historical biogeography of a cosmopolitan bird clade. Molecular Phylogenetics and Evolution 159, 107106. https://doi.org/10.1016/j.ympev.2021.107106

Goetz, S.J., Steinberg, D., Betts, M.G., Holmes, R.T., Doran, P.J., Dubayah, R., Hofton, M., 2010. Lidar remote sensing variables predict breeding habitat of a Neotropical migrant bird. Ecology 91, 1569–1576. https://doi.org/10.1890/09-1670.1

Graham, C.H., Elith, J., Hijmans, R.J., Guisan, A., Peterson, A.T., Loiselle, B.A., 2008. The influence of spatial errors in species occurrence data used in distribution models. Journal of Applied Ecology 45, 239–247. https://doi.org/10.1111/j.1365-2664.2007.01408.x

Graham, R.W., Lundelius, E.L., Graham, M.A., Schroeder, E.K., Toomey, R.S., Anderson, E., Barnosky, A.D., Burns, J.A., Churcher, C.S., Grayson, D.K., Guthrie, R.D., Harington, C.R., Jefferson, G.T., Martin, L.D., McDonald, H.G., Morlan, R.E., Semken, H.A., Webb, S.D., Werdelin, L., Wilson, M.C., 1996. Spatial Response of Mammals to Late Quaternary Environmental Fluctuations. Science 272, 1601–1606. https://doi.org/10.1126/science.272.5268.1601

Grant, P.R., Grant, B.R., 2020. How and Why Species Multiply, How and Why Species Multiply. Princeton University Press.

Grant, P.R., Weiner, J., 2017. Ecology and Evolution of Darwin's Finches (Princeton Science Library Edition): Princeton Science Library Edition.

Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in ecology. Ecological Modelling 135, 147–186. https://doi.org/10.1016/S0304-3800(00)00354-9

Haldane, J.B.S., 1956. The relation between density regulation and natural selection. Proceedings of the Royal Society of London. Series B - Biological Sciences 145, 306–308. https://doi.org/10.1098/rspb.1956.0039

Hallman, T.A., Robinson, W.D., 2020. Comparing multi- and single-scale species distribution and abundance models built with the boosted regression tree algorithm. Landscape Ecol 35, 1161–1174. https://doi.org/10.1007/s10980-020-01007-7

Harris, I., Jones, P.D., Osborn, T.J., Lister, D.H., 2014. Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset. International Journal of Climatology 34, 623–642. https://doi.org/10.1002/joc.3711

Hefley, T.J., Hooten, M.B., 2015. On the existence of maximum likelihood estimates for presence-only data. Methods in Ecology and Evolution 6, 648–655. https://doi.org/10.1111/2041-210X.12340

Heim, W., Heim, R.J., Beermann, I., Burkovskiy, O.A., Gerasimov, Y., Ktitorov, P., Ozaki, K.,
Panov, I., Sander, M.M., Sjöberg, S., Smirenski, S.M., Thomas, A., Tøttrup, A.P.,
Tiunov, I.M., Willemoes, M., Hölzel, N., Thorup, K., Kamp, J., 2020. Using geolocator
tracking data and ringing archives to validate citizen-science based seasonal predictions
of bird distribution in a data-poor region. Global Ecology and Conservation 24, e01215.
https://doi.org/10.1016/j.gecco.2020.e01215
Helm, B., Van Doren, B.M., Hoffmann, D., Hoffmann, U., 2019. Evolutionary response to climate
change in migratory Pied Flycatchers. Current Biology 29, 3714-3719.e4.
https://doi.org/10.1016/j.cub.2019.08.072
Herbert, J.A., Chakraborty, A., Naylor, L.W., Beatty, W.S., Krementz, D.G., 2018. Effects of
landscape structure and temporal habitat dynamics on wintering mallard abundance.
Landscape Ecol 33, 1319–1334. https://doi.org/10.1007/s10980-018-0671-7
Herbert, J.A., Chakraborty, A., Naylor, L.W., Krementz, D.G., 2021. Habitat associations of
wintering dabbling ducks in the Arkansas Mississippi Alluvial Valley: implications for
waterfowl management beyond the mallard. wbio 2021.
https://doi.org/10.2981/wlb.00792
Hole, D.G., Huntley, B., Arinaitwe, J., Butchart, S.H.M., Collingham, Y.C., Fishpool, L.D.C.,
Pain, D.J., Willis, S.G., 2011. Toward a Management Framework for Networks of
Protected Areas in the Face of Climate Change. Conservation Biology 25, 305–315.
https://doi.org/10.1111/j.1523-1739.2010.01633.x
Holt, R.D., Keitt, T.H., Lewis, M.A., Maurer, B.A., Taper, M.L., 2005. Theoretical models of
species' borders: single species approaches. Oikos 108, 18–27.
https://doi.org/10.1111/j.0030-1299.2005.13147.x
Huey, R.B., Hertz, P.E., Sinervo, B., 2003. Behavioral drive versus behavioral inertia in
evolution: A null model approach. The American Naturalist 161, 357–366.
https://doi.org/10.1086/346135
Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A.M., Jess, M., Williams, S.E., 2012.
Predicting organismal vulnerability to climate warming: roles of behaviour, physiology
and adaptation. Philosophical Transactions of the Royal Society B: Biological Sciences
367, 1665–1679. https://doi.org/10.1098/rstb.2012.0005
Huisman, J., Olff, H., Fresco, L. f. m., 1993. A hierarchical set of models for species response
analysis. Journal of Vegetation Science 4, 37–46. https://doi.org/10.2307/3235732
Huntley, B., Berry, P.M., Cramer, W., McDonald, A.P., 1995. Special Paper: Modelling Present
and Potential Future Ranges of Some European Higher Plants Using Climate Response
Surfaces. Journal of Biogeography 22, 967–1001. https://doi.org/10.2307/2845830
Huntley, B., Collingham, Y.C., Green, R.E., Hilton, G.M., Rahbek, C., Willis, S.G., 2006.
Potential impacts of climatic change upon geographical distributions of birds. Ibis 148,
8–28. https://doi.org/10.1111/j.1474-919X.2006.00523.x
Huntley, B., Collingham, Y.C., Willis, S.G., Green, R.E., 2008. Potential Impacts of Climatic
Change on European Breeding Birds. PLOS ONE 3, e1439.
https://doi.org/10.1371/journal.pone.0001439
Jaeger, R.G., 1970. Potential Extinction Through Competition Between Two Species of
Terrestrial Salamanders. Evolution 24, 632–642. https://doi.org/10.2307/2406842
Jahn, A.E., Buechley, E., Bugoni, L., Cereghetti, J., Repenning, M., Marra, P.P., Ryder, T.B.,
2021. Variable Seasonal Movement Dynamics Among Individual Snail Kites
(Rostrhamus sociabilis) In South America. Journal of Raptor Research.
https://doi.org/10.3356/JRR-20-77
Jahn, A.E., Seavy, N., Bejarano, V., Guzmán, M.B., Provinciato, I.C.C., Pizo, M., MacPherson,
M., 2016. Intra-tropical migration and wintering areas of Fork-tailed Flycatchers
(Tyrannus savana) breeding in São Paulo, Brazil. https://doi.org/10.1007/BF03544339

Jankowski, J.E., Londoño, G.A., Robinson, S.K., Chappell, M.A., 2013. Exploring the role of
physiology and biotic interactions in determining elevational ranges of tropical animals.
Ecography 001–012.
Jankowski, J.E., Robinson, S.K., Levey, D.J., 2010. Squeezed at the top: Interspecific
aggression may constrain elevational ranges in tropical birds. Ecology 91, 1877–1884.
https://doi.org/10.1890/09-2063.1
Jansen, F., Oksanen, J., 2013. How to model species responses along ecological gradients –
Huisman–Olff–Fresco models revisited. Journal of Vegetation Science 24, 1108–1117.
https://doi.org/10.1111/jvs.12050
Jarnevich, C.S., Talbert, M., Morisette, J., Aldridge, C., Brown, C.S., Kumar, S., Manier, D.,
Talbert, C., Holcombe, T., 2017. Minimizing effects of methodological decisions on
interpretation and prediction in species distribution studies: An example with background
selection. Ecological Modelling 363, 48–56.
https://doi.org/10.1016/j.ecolmodel.2017.08.017
Jiguet, F., Gadot, A.-S., Julliard, R., Newson, S.E., Couvet, D., 2007. Climate envelope, life
history traits and the resilience of birds facing global change. Global Change Biology 13,
1672–1684. https://doi.org/10.1111/j.1365-2486.2007.01386.x
Jirinec, V., Burner, R.C., Amaral, B.R., Bierregaard Jr., R.O., Fernandez-Arellano, G.,
Hernandez-Palma, A., Johnson, E.I., Lovejoy, T.E., Powell, L.L., Rutt, C.L., Wolfe, J.D.,
Stouffer, P.C., 2021. Morphological consequences of climate change for resident birds in
intact Amazonian rainforest. Science Advances 7, eabk1743.
https://doi.org/10.1126/sciadv.abk1743
Jirinec, V., Isdell, R.E., Leu, M., 2016. Prey availability and habitat structure explain breeding
space use of a migratory songbird. Condor 118, 309–328.
https://doi.org/10.1650/CONDOR-15-140.1
Jones, J., 2001. Habitat selection studies in avian ecology: a critical review. Auk 118, 557–562.
https://doi.org/10.1093/auk/118.2.557
Jones, M.C., Cheung, W.W.L., 2015. Multi-model ensemble projections of climate change
effects on global marine biodiversity. ICES Journal of Marine Science 72, 741–752.
https://doi.org/10.1093/icesjms/fsu172
Kapos, V., 1989. Effects of isolation on the water status of forest patches in the Brazilian
Amazon. Journal of Tropical Ecology 5, 173–185.
https://doi.org/10.1017/S0266467400003448
Kearney, M., 2006. Habitat, environment and niche: what are we modelling? Oikos 115, 186–
191. https://doi.org/10.1111/j.2006.0030-1299.14908.x
Kearney, M., Porter, W., 2009. Mechanistic niche modelling: combining physiological and spatial
data to predict species' ranges. Ecology Letters 12, 334–350.
https://doi.org/10.1111/j.1461-0248.2008.01277.x
Kearney, M.R., Porter, W.P., Murphy, S.A., 2016. An estimate of the water budget for the
endangered night parrot of Australia under recent and future climates. Clim Chang
Responses 3, 14. https://doi.org/10.1186/s40665-016-0027-y
Kitzes, J., Schricker, L., 2019. The necessity, promise and challenge of automated biodiversity
surveys. Environmental Conservation 46, 247–250.
https://doi.org/10.1017/S0376892919000146
Kokko, H., Gunnarsson, T.G., Morrell, L.J., Gill, J.A., 2006. Why do female migratory birds
arrive later than males? Journal of Animal Ecology 75, 1293–1303.
https://doi.org/10.1111/j.1365-2656.2006.01151.x
König, C., Wüest, R.O., Graham, C.H., Karger, D.N., Sattler, T., Zimmermann, N.E., Zurell, D.,
2021. Scale dependency of joint species distribution models challenges interpretation of
biotic interactions. Journal of Biogeography 1–11. https://doi.org/10.1111/jbi.14106

Kottler, E.J., Dickman, E.E., Sexton, J.P., Emery, N.C., Franks, S.J., 2021. Draining the Swamping Hypothesis: Little Evidence that Gene Flow Reduces Fitness at Range Edges. Trends in Ecology & Evolution 36, 533–544. https://doi.org/10.1016/j.tree.2021.02.004

La Sorte, F.A., Lepczyk, C.A., Burnett, J.L., Hurlbert, A.H., Tingley, M.W., Zuckerberg, B., 2018. Opportunities and challenges for big data ornithology. The Condor 120, 414–426. https://doi.org/10.1650/CONDOR-17-206.1

Latimer, C.E., Zuckerberg, B., 2021. Habitat loss and thermal tolerances influence the sensitivity of resident bird populations to winter weather at regional scales. Journal of Animal Ecology 90, 317–329. https://doi.org/10.1111/1365-2656.13332

Lavretsky, P., McInerney, N.R., Mohl, J.E., Brown, J.I., James, H.F., McCracken, K.G., Fleischer, R.C., 2020. Assessing changes in genomic divergence following a century of human-mediated secondary contact among wild and captive-bred ducks. Molecular Ecology 29, 578–595. https://doi.org/10.1111/mec.15343

Léandri-Breton, D.-J., Bêty, J., 2020. Vulnerability to predation may affect species distribution: plovers with broader arctic breeding range nest in safer habitat. Scientific Reports 10, 5032. https://doi.org/10.1038/s41598-020-61956-6

Leclerc, C., Villéger, S., Marino, C., Bellard, C., 2020. Global changes threaten functional and taxonomic diversity of insular species worldwide. Diversity and Distributions 26, 402–414. https://doi.org/10.1111/ddi.13024

Lees, A.C., Rosenberg, K.V., Ruiz-gutierrez, V., Marsden, S., Schulenberg, T.S., Rodewald, A.D., 2020. A roadmap to identifying and filling shortfalls in Neotropical ornithology. The Auk 137, 1–17. https://doi.org/10.1093/auk/ukaa048

Lehmann, P., Ammunét, T., Barton, M., Battisti, A., Eigenbrode, S.D., Jepsen, J.U., Kalinkat, G., Neuvonen, S., Niemelä, P., Terblanche, J.S., Økland, B., Björkman, C., 2020. Complex responses of global insect pests to climate warming. Frontiers in Ecology and the Environment 18, 141–150. https://doi.org/10.1002/fee.2160

Lei, Y., Liu, Q., 2021. Tolerance niche expansion and potential distribution prediction during Asian openbill bird range expansion. Ecology and Evolution 1–13. https://doi.org/10.1002/ece3.7456

Loiselle, B.A., Howell, C.A., Graham, C.H., Goerck, J.M., Brooks, T., Smith, K.G., Williams, P.H., 2003. Avoiding Pitfalls of Using Species Distribution Models in Conservation Planning. Conservation Biology 17, 1591–1600. https://doi.org/10.1111/j.1523-1739.2003.00233.x

Louthan, A.M., Doak, D.F., Angert, A.L., 2015. Where and when do species interactions set range limits? Trends in Ecology & Evolution 30, 780–792. https://doi.org/10.1016/j.tree.2015.09.011

Loyola, R.D., Lemes, P., Nabout, J.C., Trindade-Filho, J., Sagnori, M.D., Dobrovolski, R., Diniz-Filho, J.A.F., 2013. A straightforward conceptual approach for evaluating spatial conservation priorities under climate change. Biodivers Conserv 22, 483–495. https://doi.org/10.1007/s10531-012-0424-x

Lukas, J., Auer, F., Goldhammer, T., Krause, J., Romanczuk, P., Klamser, P., Arias-Rodriguez, L., Bierbach, D., 2021. Diurnal Changes in Hypoxia Shape Predator-Prey Interaction in a Bird-Fish System. https://doi.org/10.14279/depositonce-11735

Lurgi, M., Brook, B.W., Saltré, F., Fordham, D.A., 2015. Modelling range dynamics under global change: which framework and why? Methods in Ecology and Evolution 6, 247–256. https://doi.org/10.1111/2041-210X.12315

MacKenzie, D.I., Nichols, J.D., Sutton, N., Kawanishi, K., Bailey, L.L., 2005. Improving inferences in population studies of rare species that are detected imperfectly. Ecology 86, 1101–1113. https://doi.org/10.1890/04-1060

MacPherson, M.P., Jahn, A.E., Mason, N.A., 2022. Morphology of migration: associations
between wing shape, bill morphology and migration in kingbirds (*Tyrannus*). Biological
Journal of the Linnean Society 135, 71–83. https://doi.org/10.1093/biolinnean/blab123
MacPherson, M.P., Jahn, A.E., Murphy, M.T., Kim, D.H., Cueto, V.R., Tuero, D.T., Hill, E.D.,
2018a. Follow the rain? Environmental drivers of Tyrannus migration across the New
World. Auk 135, 881–894. https://doi.org/10.1642/AUK-17-209.1
MacPherson, M.P., Webb, E.B., Raedeke, A., Mengel, D., Nelson, F., 2018b. A review of
Bayesian belief network models as decision-support tools for wetland conservation: Are
water birds potential umbrella taxa? Biological Conservation 226, 215–223.
https://doi.org/10.1016/j.biocon.2018.08.001
Maggini, R., Lehmann, A., Kéry, M., Schmid, H., Beniston, M., Jenni, L., Zbinden, N., 2011. Are
Swiss birds tracking climate change?: Detecting elevational shifts using response curve
shapes. Ecological Modelling 222, 21–32.
https://doi.org/10.1016/j.ecolmodel.2010.09.010
Mamantov, M.A., Gibson-Reinemer, D.K., Linck, E.B., Sheldon, K.S., 2021. Climate-driven
range shifts of montane species vary with elevation. Global Ecology and Biogeography
30, 784–794. https://doi.org/10.1111/geb.13246
Manel, S., Dias, J.M., Buckton, S.T., Ormerod, S.J., 1999. Alternative methods for predicting
species distribution: an illustration with Himalayan river birds. Journal of Applied Ecology
36, 734–747. https://doi.org/10.1046/j.1365-2664.1999.00440.x
Marini, Miguel Â., Barbet-Massin, M., Lopes, L.E., Jiguet, F., 2010. Predicting the occurrence of
rare Brazilian birds with species distribution models. J Ornithol 151, 857–866.
https://doi.org/10.1007/s10336-010-0523-y
Marini, M.Â., Barbet-Massin, M., Lopes, L.E., Jiguet, F., 2009. Predicted climate-driven bird
distribution changes and forecasted conservation conflicts in a Neotropical savanna.
Conservation Biology 23, 1558–1567. https://doi.org/10.1111/j.1523-1739.2009.01258.x
Marini, Miguel Ângelo, Barbet-Massin, M., Martinez, J., Prestes, N.P., Jiguet, F., 2010. Applying
ecological niche modelling to plan conservation actions for the Red-spectacled Amazon
(Amazona pretrei). Biological Conservation 143, 102–112.
https://doi.org/10.1016/j.biocon.2009.09.009
Marini, M.Â., Hall, L., Bates, J., Steinheimer, F.D., McGowan, R., Silveira, L.F., Lijtmaer, D.A.,
Tubaro, P.L., Córdoba-Córdoba, S., Gamauf, A., Greeney, H.F., Schweizer, M.,
Kamminga, P., Cibois, A., Vallotton, L., Russell, D., Robinson, S.K., Sweet, P.R.,
Frahnert, S., Corado, R., Heming, N.M., 2020. The five million bird eggs in the world's
museum collections are an invaluable and underused resource. The Auk 137, ukaa036.
https://doi.org/10.1093/auk/ukaa036
Matuoka, M.A., Benchimol, M., Almeida-Rocha, J.M. de, Morante-Filho, J.C., 2020. Effects of
anthropogenic disturbances on bird functional diversity: A global meta-analysis.
Ecological Indicators 116, 106471. https://doi.org/10.1016/j.ecolind.2020.106471
McKinnon, E.A., Love, O.P., 2018. Ten years tracking the migrations of small landbirds:
Lessons learned in the golden age of bio-logging. tauk 135, 834–856.
https://doi.org/10.1642/AUK-17-202.1
Merkle, J.A., Abrahms, B., Armstrong, J.B., Sawyer, H., Costa, D.P., Chalfoun, A.D., 2022. Site
fidelity as a maladaptive behavior in the Anthropocene. Frontiers in Ecology and the
Environment 20, 187–194. https://doi.org/10.1002/fee.2456
Merow, C., Smith, M.J., Silander, J.A., Jr., 2013. A practical guide to MaxEnt for modeling
species' distributions: what it does, and why inputs and settings matter. Ecography 36,
1058–1069. https://doi.org/10.1111/j.1600-0587.2013.07872.x
Mi, C., Huettmann, F., Guo, Y., Han, X., Wen, L., 2017. Why choose Random Forest to predict
rare species distribution with few samples in large undersampled areas? Three Asian

crane species models provide supporting evidence. PeerJ 5, e2849.
https://doi.org/10.7717/peerj.2849

Møller, A.P., Rubolini, D., Lehikoinen, E., 2008. Populations of migratory bird species that did not show a phenological response to climate change are declining. PNAS 105, 16195–16200. https://doi.org/10.1073/pnas.0803825105

Moore, R.P., Robinson, W.D., Lovette, I.J., Robinson, T.R., 2008. Experimental evidence for extreme dispersal limitation in tropical forest birds. Ecology Letters 11, 960–968. https://doi.org/10.1111/j.1461-0248.2008.01196.x

Morisette, J.T., Jarnevich, C.S., Holcombe, T.R., Talbert, C.B., Ignizio, D., Talbert, M.K., Silva, C., Koop, D., Swanson, A., Young, N.E., 2013. VisTrails SAHM: visualization and workflow management for species habitat modeling. Ecography 36, 129–135. https://doi.org/10.1111/j.1600-0587.2012.07815.x

Moritz, C., Agudo, R., 2013. The future of species under climate change: resilience or decline? Science 341, 504–508. https://doi.org/10.1126/science.1237190

Morris, W.F., Doak, D.F., 2002. Quantitative conservation biology: theory and practice of population viability analysis. Sinauer Associates, Sunderland, Mass.

Morrison, M.L., 1986. Bird populations as indicators of environmental change, in: Johnston, R.F. (Ed.), Current Ornithology: Volume 3, Current Ornithology. Springer US, Boston, MA, pp. 429–451. https://doi.org/10.1007/978-1-4615-6784-4_10

Naka, L.N., Costa, B.M. da S., Lima, G.R., Claramunt, S., 2022. Riverine Barriers as Obstacles to Dispersal in Amazonian Birds. Frontiers in Ecology and Evolution 10.

Nakao, K., Higa, M., Tsuyama, I., Matsui, T., Horikawa, M., Tanaka, N., 2013. Spatial conservation planning under climate change: Using species distribution modeling to assess priority for adaptive management of Fagus crenata in Japan. Journal for Nature Conservation 21, 406–413. https://doi.org/10.1016/j.jnc.2013.06.003

Neilan, W.L., Barton, P.S., McAlpine, C.A., Wood, J.T., Lindenmayer, D.B., 2019. Contrasting effects of mosaic structure on alpha and beta diversity of bird assemblages in a human-modified landscape. Ecography 42, 173–186. https://doi.org/10.1111/ecog.02981

Nilsson, C., Klaassen, R.H.G., Alerstam, T., 2013. Differences in speed and duration of bird migration between spring and autumn. The American Naturalist 181, 837–845. https://doi.org/10.1086/670335

Nunes, M.H., Campana Camargo, J.L., Vincent, G., Calders, K., Oliveira, R.S., Huete, A., Mendes de Moura, Y., Nelson, B., Smith, M.N., Stark, S.C., Maeda, E.E., 2022. Forest fragmentation impacts the seasonality of Amazonian evergreen canopies | Nature Communications. Nature Communications 13. https://doi.org/10.1038/s41467-022-28490-7

Oksanen, J., Minchin, P.R., 2002. Continuum theory revisited: what shape are species responses along ecological gradients? Ecological Modelling 157, 119–129. https://doi.org/10.1016/S0304-3800(02)00190-4

Oppel, S., Meirinho, A., Ramírez, I., Gardner, B., O'Connell, A.F., Miller, P.I., Louzao, M., 2012. Comparison of five modelling techniques to predict the spatial distribution and abundance of seabirds. Biological Conservation 156, 94–104. https://doi.org/10.1016/j.biocon.2011.11.013

Palacio, F.X., Girini, J.M., 2018. Biotic interactions in species distribution models enhance model performance and shed light on natural history of rare birds: a case study using the straight-billed reedhaunter Limnoctites rectirostris. Journal of Avian Biology 49, e01743. https://doi.org/10.1111/jav.01743

Palma, A.C., Goosem, M., Fensham, R.J., Goosem, S., Preece, N.D., Stevenson, P.R., Laurance, S.G.W., 2020. Dispersal and recruitment limitations in secondary forests. Journal of Vegetation Science e12975. https://doi.org/10.1111/jvs.12975

1284   Parmesan, C., Matthews, J., 2005. Biological Impacts of Climate Change.
1285   https://doi.org/10.17226/12491
1286   Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change impacts across
1287   natural systems. Nature 421, 37–42. https://doi.org/10.1038/nature01286
1288   Pearl, J., 1985. Bayesian networks: A model of self-activated memory for evidential reasoning,
1289   in: Proceedings of the 7th Conference of the Cognitive Science Society. University of
1290   California, Irvine, CA.
1291   Pearson, R.G., Dawson, T.P., 2003. Predicting the impacts of climate change on the distribution
1292   of species: are bioclimate envelope models useful? Global Ecology and Biogeography
1293   12, 361–371. https://doi.org/10.1046/j.1466-822X.2003.00042.x
1294   Pearson, R.G., Raxworthy, C.J., Nakamura, M., Peterson, A.T., 2007. Predicting species
1295   distributions from small numbers of occurrence records: a test case using cryptic geckos
1296   in Madagascar. Journal of Biogeography 34, 102–117. https://doi.org/10.1111/j.1365-
1297   2699.2006.01594.x
1298   Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., Clark, T.D.,
1299   Colwell, R.K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., Garcia,
1300   R.A., Griffis, R.B., Hobday, A.J., Janion-Scheepers, C., Jarzyna, M.A., Jennings, S.,
1301   Lenoir, J., Linnetved, H.I., Martin, V.Y., McCormack, P.C., McDonald, J., Mitchell, N.J.,
1302   Mustonen, T., Pandolfi, J.M., Pettorelli, N., Popova, E., Robinson, S.A., Scheffers, B.R.,
1303   Shaw, J.D., Sorte, C.J.B., Strugnell, J.M., Sunday, J.M., Tuanmu, M.-N., Vergés, A.,
1304   Villanueva, C., Wernberg, T., Wapstra, E., Williams, S.E., 2017. Biodiversity
1305   redistribution under climate change: Impacts on ecosystems and human well-being.
1306   Science 355. https://doi.org/10.1126/science.aai9214
1307   Pérez-Granados, C., Bota, G., Giralt, D., Barrero, A., Gómez-Catasús, J., Rosa, D.B.-D.L.,
1308   Traba, J., 2019. Vocal activity rate index: a useful method to infer terrestrial bird
1309   abundance with acoustic monitoring. Ibis 161, 901–907. https://doi.org/10.1111/ibi.12728
1310   Pérez-Granados, C., Schuchmann, K.-L., 2020. Monitoring the annual vocal activity of two
1311   enigmatic nocturnal Neotropical birds: the Common Potoo (Nyctibius griseus) and the
1312   Great Potoo (Nyctibius grandis). J Ornithol 161, 1129–1141.
1313   https://doi.org/10.1007/s10336-020-01795-4
1314   Peterson, M.L., Doak, D.F., Morris, W.F., 2019. Incorporating local adaptation into forecasts of
1315   species' distribution and abundance under climate change. Global Change Biology 25,
1316   775–793. https://doi.org/10.1111/gcb.14562
1317   Phillips, S.J., Anderson, R.P., Schapire, R.E., 2006. Maximum entropy modeling of species
1318   geographic distributions. Ecological Modelling 190, 231–259.
1319   https://doi.org/10.1016/j.ecolmodel.2005.03.026
1320   Pierson, J.C., Beissinger, S.R., Bragg, J.G., Coates, D.J., Oostermeijer, J.G.B., Sunnucks, P.,
1321   Schumaker, N.H., Trotter, M.V., Young, A.G., 2015. Incorporating evolutionary
1322   processes into population viability models. Conservation Biology 29, 755–764.
1323   https://doi.org/10.1111/cobi.12431
1324   Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J.,
1325   Brander, K., Bruno, J.F., Buckley, L.B., Burrows, M.T., Duarte, C.M., Halpern, B.S.,
1326   Holding, J., Kappel, C.V., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F.,
1327   Thompson, S.A., Richardson, A.J., 2013. Global imprint of climate change on marine life.
1328   Nature Climate Change 3, 919–925. https://doi.org/10.1038/nclimate1958
1329   Porter, W.P., Mitchell, J.W., Beckman, W.A., DeWitt, C.B., 1973. Behavioral implications of
1330   mechanistic ecology. Oecologia 13, 1–54. https://doi.org/10.1007/BF00379617
1331   Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P.,
1332   Adriaensen, F., Ahola, M.P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K.S.,
1333   Borras, A., Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A.A.,
1334   Dingemanse, N.J., Doi, H., Eeva, T., Fickel, J., Filella, I., Fossøy, F., Goodenough, A.E.,

Hall, S.J.G., Hansson, B., Harris, M., Hasselquist, D., Hickler, T., Joshi, J., Kharouba, H., Martínez, J.G., Mihoub, J.-B., Mills, J.A., Molina-Morales, M., Moksnes, A., Ozgul, A., Parejo, D., Pilard, P., Poisbleau, M., Rousset, F., Rödel, M.-O., Scott, D., Senar, J.C., Stefanescu, C., Stokke, B.G., Kusano, T., Tarka, M., Tarwater, C.E., Thonicke, K., Thorley, J., Wilting, A., Tryjanowski, P., Merilä, J., Sheldon, B.C., Pape Møller, A., Matthysen, E., Janzen, F., Dobson, F.S., Visser, M.E., Beissinger, S.R., Courtiol, A., Kramer-Schadt, S., 2019. Adaptive responses of animals to climate change are most likely insufficient. Nature Communications 10, 3109. https://doi.org/10.1038/s41467-019-10924-4

Radosavljevic, A., Anderson, R.P., 2014. Making better Maxent models of species distributions: complexity, overfitting and evaluation. Journal of Biogeography 41, 629–643. https://doi.org/10.1111/jbi.12227

Randin, C.F., Ashcroft, M.B., Bolliger, J., Cavender-Bares, J., Coops, N.C., Dullinger, S., Dirnböck, T., Eckert, S., Ellis, E., Fernández, N., Giuliani, G., Guisan, A., Jetz, W., Joost, S., Karger, D., Lembrechts, J., Lenoir, J., Luoto, M., Morin, X., Price, B., Rocchini, D., Schaepman, M., Schmid, B., Verburg, P., Wilson, A., Woodcock, P., Yoccoz, N., Payne, D., 2020. Monitoring biodiversity in the Anthropocene using remote sensing in species distribution models. Remote Sensing of Environment 239, 111626. https://doi.org/10.1016/j.rse.2019.111626

Reed, J.M., Boulinier, T., Danchin, E., Oring, L.W., 1999. Informed Dispersal, in: Nolan, V., Ketterson, E.D., Thompson, C.F. (Eds.), Current Ornithology, Current Ornithology. Springer US, Boston, MA, pp. 189–259. https://doi.org/10.1007/978-1-4757-4901-4_5

Rehm, E., Feeley, K.J., 2016. Many species risk mountain top extinction long before they reach the top. Frontiers of Biogeography 8. https://doi.org/10.21425/F5FBG27788

Remsen, J.V., 1995. The importance of continued collecting of bird specimens to ornithology and bird conservation. Bird Conservation International 5, 146–180. https://doi.org/10.1017/S095927090000099X

Renfrew, R.B., Kim, D., Perlut, N., Smith, J., Fox, J., Marra, P.P., 2013. Phenological matching across hemispheres in a long-distance migratory bird. Diversity and Distributions 19, 1008–1019. https://doi.org/10.1111/ddi.12080

Reznick, D., Baxter, R.J., Endler, J., 1994. Long-term studies of tropical stream fish communities: The use of field notes and museum collections to reconstruct communities of the past. Integr Comp Biol 34, 452–462. https://doi.org/10.1093/icb/34.3.452

Ribas, C.C., Aleixo, A., Gubili, C., d'Horta, F.M., Brumfield, R.T., Cracraft, J., 2018. Biogeography and diversification of Rhegmatorhina (Aves: Thamnophilidae): Implications for the evolution of Amazonian landscapes during the Quaternary. Journal of Biogeography 45, 917–928. https://doi.org/10.1111/jbi.13169

Ribas, C.C., Aleixo, A., Nogueira, A.C.R., Miyaki, C.Y., Cracraft, J., 2012. A palaeobiogeographic model for biotic diversification within Amazonia over the past three million years. Proceedings of the Royal Society B: Biological Sciences 279, 681–689. https://doi.org/10.1098/rspb.2011.1120

Riddell, E.A., Iknayan, K.J., Hargrove, L., Tremor, S., Patton, J.L., Ramirez, R., Wolf, B.O., Beissinger, S.R., 2021. Exposure to climate change drives stability or collapse of desert mammal and bird communities. Science 371, 633–636. https://doi.org/10.1126/science.abd4605

Riotte-Lambert, L., Matthiopoulos, J., 2020. Environmental Predictability as a Cause and Consequence of Animal Movement. Trends in Ecology and Evolution 35, 163–174. https://doi.org/10.1016/j.tree.2019.09.009

Ritter, C.D., Coelho, L.A., Capurucho, J.M., Borges, S.H., Cornelius, C., Ribas, C.C., 2021. Sister species, different histories: comparative phylogeography of two bird species

associated with Amazonian open vegetation. Biological Journal of the Linnean Society 132, 161–173. https://doi.org/10.1093/biolinnean/blaa167

Roberts, C.P., Allen, C.R., Angeler, D.G., Twidwell, D., 2019. Shifting avian spatial regimes in a changing climate. Nat. Clim. Chang. 9, 562–566. https://doi.org/10.1038/s41558-019-0517-6

Rosado, B.H.P., Figueiredo, M.S.L., Mattos, E.A. de, Grelle, C.E.V., 2016. Eltonian shortfall due to the Grinnellian view: functional ecology between the mismatch of niche concepts. Ecography 39, 1034–1041. https://doi.org/10.1111/ecog.01678

Roslin, T., Antão, L., Hällfors, M., Meyke, E., Lo, C., Tikhonov, G., Delgado, M. del M., Gurarie, E., Abadonova, M., Abduraimov, O., Adrianova, O., Akimova, T., Akkiev, M., Ananin, A., Andreeva, E., Andriychuk, N., Antipin, M., Arzamascev, K., Babina, S., Babushkin, M., Bakin, O., Barabancova, A., Basilskaja, I., Belova, N., Belyaeva, N., Bespalova, T., Bisikalova, E., Bobretsov, A., Bobrov, V., Bobrovskyi, V., Bochkareva, E., Bogdanov, G., Bolshakov, V., Bondarchuk, S., Bukharova, E., Butunina, A., Buyvolov, Y., Buyvolova, A., Bykov, Y., Chakhireva, E., Chashchina, O., Cherenkova, N., Chistjakov, S., Chuhontseva, S., Davydov, E.A., Demchenko, V., Diadicheva, E., Dobrolyubov, A., Dostoyevskaya, L., Drovnina, S., Drozdova, Z., Dubanaev, A., Dubrovsky, Y., Elsukov, S., Epova, L., Ermakova, O., Ermakova, O.S., Ershkova, E., Esengeldenova, A., Evstigneev, O., Fedchenko, I., Fedotova, V., Filatova, T., Gashev, S., Gavrilov, A., Gaydysh, I., Golovcov, D., Goncharova, N., Gorbunova, E., Gordeeva, T., Grishchenko, V., Gromyko, L., Hohryakov, V., Hritankov, A., Ignatenko, E., Igosheva, S., Ivanova, U., Ivanova, N., Kalinkin, Y., Kaygorodova, E., Kazansky, F., Kiseleva, D., Knorre, A., Kolpashikov, L., Korobov, E., Korolyova, H., Korotkikh, N., Kosenkov, G., Kossenko, S., Kotlugalyamova, E., Kozlovsky, E., Kozsheechkin, V., Kozurak, A., Kozyr, I., Krasnopevtseva, A., Kruglikov, S., Kuberskaya, O., Kudryavtsev, A., Kulebyakina, E., Kulsha, Y., Kupriyanova, M., Kurbanbagamaev, M., Kutenkov, A., Kutenkova, N., Kuyantseva, N., Kuznetsov, A., Larin, E., Lebedev, P., Litvinov, K., Luzhkova, N., Mahmudov, A., Makovkina, L., Mamontov, V., Mayorova, S., Megalinskaja, I., Meydus, A., Minin, A., Mitrofanov, O., Motruk, M., Myslenkov, A., Nasonova, N., Nemtseva, N., Nesterova, I., Nezdoliy, T., Niroda, T., Novikova, T., Panicheva, D., Pavlov, A., Pavlova, K., Podolski, S., Polikarpova, N., Polyanskaya, T., Pospelov, I., Pospelova, E., Prokhorov, I., Prokosheva, I., Puchnina, L., Putrashyk, I., Raiskaya, J., Rozhkov, Y., Rozhkova, O., Rudenko, M., Rybnikova, I., Rykova, S., Sahnevich, M., Samoylov, A., Sanko, V., Sapelnikova, I., Sazonov, S., Selyunina, Z., Shalaeva, K., Shashkov, M., Shcherbakov, A., Shevchyk, V., Shubin, S., Shujskaja, E., Sibgatullin, R., Sikkila, N., Sitnikova, E., Sivkov, A., Skok, N., Skorokhodova, S., Smirnova, E., Sokolova, G., Sopin, V., Spasovski, Y., Stepanov, S., Stratiy, V., Strekalovskaya, V., Sukhov, A., Suleymanova, G., Sultangareeva, L., Teleganova, V., Teplov, V., Teplova, V., Tertitsa, T., Timoshkin, V., Tirski, D., Tolmachev, A., Tomilin, A., Tselishcheva, L., Turgunov, M., Tyukh, Y., Van, P., Van, V., Vasin, A., Vasina, A., Vekliuk, A., Vetchinnikova, L., Vinogradov, V., Volodchenkov, N., Voloshina, I., Xoliqov, T., Yablonovska-Grishchenko, E., Yakovlev, V., Yakovleva, M., Yantser, O., Yarema, Y., Zahvatov, A., Zakharov, V., Zelenetskiy, N., Zheltukhin, A., Zubina, T., Kurhinen, J., Ovaskainen, O., 2021. Phenological shifts of abiotic events, producers and consumers across a continent. Nature Climate Change 11, 241–248. https://doi.org/10.1038/s41558-020-00967-7

Sander, M., Chamberlain, D., Mermillon, C., Alba, R., Jähnig, S., Rosselli, D., Lisovski, S., 2020. Reduced breeding success suggests trophic mismatch despite timely arrival in an alpine songbird.

Saupe, E.E., Barve, V., Myers, C.E., Soberón, J., Barve, N., Hensz, C.M., Peterson, A.T., Owens, H.L., Lira-Noriega, A., 2012. Variation in niche and distribution model

performance: The need for a priori assessment of key causal factors. Ecological Modelling 237–238, 11–22. https://doi.org/10.1016/j.ecolmodel.2012.04.001

Sauser, C., Delord, K., Barbraud, C., 2021. Sea ice and local weather affect reproductive phenology of a polar seabird with breeding consequences. Ornithological Applications 123, duab032. https://doi.org/10.1093/ornithapp/duab032

Sax, D.F., Early, R., Bellemare, J., 2013. Niche syndromes, species extinction risks, and management under climate change. Trends in Ecology & Evolution 28, 517–523. https://doi.org/10.1016/j.tree.2013.05.010

Scheffers, B.R., Edwards, D.P., Diesmos, A., Williams, S.E., Evans, T.A., 2014. Microhabitats reduce animal's exposure to climate extremes. Global Change Biology 20, 495–503. https://doi.org/10.1111/gcb.12439

Scheffers, B.R., Edwards, D.P., Macdonald, S.L., Senior, R.A., Andriamahohatra, L.R., Roslan, N., Rogers, A.M., Haugaasen, T., Wright, P., Williams, S.E., 2017. Extreme thermal heterogeneity in structurally complex tropical rain forests. Biotropica 49, 35–44. https://doi.org/10.1111/btp.12355

Schulte to Bühne, H., Tobias, J.A., Durant, S.M., Pettorelli, N., 2021. Improving Predictions of Climate Change–Land Use Change Interactions. Trends in Ecology & Evolution 36, 29–38. https://doi.org/10.1016/j.tree.2020.08.019

Şekercioğlu, C., Wenny, D.G., Whelan, C.J., 2019. Why birds matter: avian ecological function and ecosystem services.

Shaffer, H.B., Fisher, R.N., Davidson, C., 1998. The role of natural history collections in documenting species declines. Trends in Ecology & Evolution 13, 27–30. https://doi.org/10.1016/S0169-5347(97)01177-4

Sheard, C., Neate-Clegg, M.H.C., Alioravainen, N., Jones, S.E.I., Vincent, C., MacGregor, H.E.A., Bregman, T.P., Claramunt, S., Tobias, J.A., 2020. Ecological drivers of global gradients in avian dispersal inferred from wing morphology. Nature Communications 11, 2463. https://doi.org/10.1038/s41467-020-16313-6

Shipley, J.R., Twining, C.W., Taff, C.C., Vitousek, M.N., Flack, A., Winkler, D.W., 2020. Birds advancing lay dates with warming springs face greater risk of chick mortality. PNAS 117, 25590–25594. https://doi.org/10.1073/pnas.2009864117

Shirley, S.M., Yang, Z., Hutchinson, R.A., Alexander, J.D., McGarigal, K., Betts, M.G., 2013. Species distribution modelling for the people: unclassified landsat TM imagery predicts bird occurrence at fine resolutions. Diversity and Distributions 19, 855–866. https://doi.org/10.1111/ddi.12093

Siepielski, A.M., Morrissey, M.B., Buoro, M., Carlson, S.M., Caruso, C.M., Clegg, S.M., Coulson, T., DiBattista, J., Gotanda, K.M., Francis, C.D., Hereford, J., Kingsolver, J.G., Augustine, K.E., Kruuk, L.E.B., Martin, R.A., Sheldon, B.C., Sletvold, N., Svensson, E.I., Wade, M.J., MacColl, A.D.C., 2017. Precipitation drives global variation in natural selection. Science 355, 959–962. https://doi.org/10.1126/science.aag2773

Slager, D.L., 2020. Seasonal and directional dispersal behavior in an ongoing dove invasion. Journal of Avian Biology 51. https://doi.org/10.1111/jav.02332

Smith, A.C., Edwards, B.P.M., 2021. North American Breeding Bird Survey status and trend estimates to inform a wide range of conservation needs, using a flexible Bayesian hierarchical generalized additive model. Ornithological Applications 123. https://doi.org/10.1093/ornithapp/duaa065

Smith, B.T., Gehara, M., Harvey, M.G., 2021. The demography of extinction in eastern North American birds. Proceedings of the Royal Society B: Biological Sciences 288, 20201945. https://doi.org/10.1098/rspb.2020.1945

Smith, F.A., Betancourt, J.L., Brown, J.H., 1995. Evolution of Body Size in the Woodrat over the Past 25,000 Years of Climate Change. Science 270, 2012–2014. https://doi.org/10.1126/science.270.5244.2012

Smith, R.J., Moore, F.R., 2005. Arrival timing and seasonal reproductive performance in a long-
distance migratory landbird. Behav Ecol Sociobiol 57, 231–239.
https://doi.org/10.1007/s00265-004-0855-9
Soberón, J., 2007. Grinnellian and Eltonian niches and geographic distributions of species.
Ecology Letters 10, 1115–1123. https://doi.org/10.1111/j.1461-0248.2007.01107.x
Soberón, J., Peterson, A.T., 2005. INTERPRETATION OF MODELS OF FUNDAMENTAL
ECOLOGICAL NICHES AND SPECIES' DISTRIBUTIONAL AREAS, Biodiversity
Informatics.
Song, G., Zhang, R., Machado-Stredel, F., Alström, P., Johansson, U.S., Irestedt, M., Mays,
H.L., McKay, B.D., Nishiumi, I., Cheng, Y., Qu, Y., Ericson, P.G.P., Fjeldså, J., Peterson,
A.T., Lei, F., 2020. Great journey of Great Tits (Parus major group): Origin,
diversification and historical demographics of a broadly distributed bird lineage. Journal
of Biogeography 47, 1585–1598. https://doi.org/10.1111/jbi.13863
Sousa, N.O. de M., Lopes, L.E., Costa, L.M., Motta-Junior, J.C., Freitas, G.H.S. de, Dornas, T.,
Vasconcelos, M.F. de, Nogueira, W., Tolentino, V.C. de M., De-Carvalho, C.B., Barbosa,
M.O., Ubaid, F.K., Nunes, A.P., Malacco, G.B., Marini, M.Â., 2021. Adopting habitat-use
to infer movement potential and sensitivity to human disturbance of birds in a
Neotropical Savannah. Biological Conservation 254, 108921.
https://doi.org/10.1016/j.biocon.2020.108921
Staniczenko, P.P.A., Sivasubramaniam, P., Suttle, K.B., Pearson, R.G., 2017. Linking
macroecology and community ecology: refining predictions of species distributions using
biotic interaction networks. Ecology Letters 20, 693–707.
https://doi.org/10.1111/ele.12770
Stanley, C.Q., MacPherson, M., Fraser, K.C., McKinnon, E.A., Stutchbury, B.J.M., 2012. Repeat
tracking of individual songbirds reveals consistent migration timing but flexibility in route.
PLOS ONE 7, e40688. https://doi.org/10.1371/journal.pone.0040688
Stanley, C.Q., McKinnon, E.A., Fraser, K.C., Macpherson, M.P., Casbourn, G., Friesen, L.,
Marra, P.P., Studds, C., Ryder, T.B., Diggs, N.E., Stutchbury, B.J.M., 2015. Connectivity
of wood thrush breeding, wintering, and migration sites based on range-wide tracking.
Conservation Biology 29, 164–174. https://doi.org/10.1111/cobi.12352
Stevenson, R.D., 1985. The relative importance of behavioral and physiological adjustments
controlling body temperature in terrestrial ectotherms. The American Naturalist 126,
362–386. https://doi.org/10.1086/284423
Stiels, D., Bastian, H.-V., Bastian, A., Schidelko, K., Engler, J.O., 2021. An iconic messenger of
climate change? Predicting the range dynamics of the European Bee-eater (Merops
apiaster). J Ornithol. https://doi.org/10.1007/s10336-021-01867-z
Stralberg, D., Matsuoka, S.M., Hamann, A., Bayne, E.M., Sólymos, P., Schmiegelow, F.K.A.,
Wang, X., Cumming, S.G., Song, S.J., 2015. Projecting boreal bird responses to climate
change: the signal exceeds the noise. Ecological Applications 25, 52–69.
https://doi.org/10.1890/13-2289.1
Sugai, L.S.M., Desjonquères, C., Silva, T.S.F., Llusia, D., 2020. A roadmap for survey designs
in terrestrial acoustic monitoring. Remote Sensing in Ecology and Conservation 6, 220–
235. https://doi.org/10.1002/rse2.131
Suggitt, A.J., Stefanescu, C., Páramo, F., Oliver, T., Anderson, B.J., Hill, J.K., Roy, D.B.,
Brereton, T., Thomas, C.D., 2012. Habitat associations of species show consistent but
weak responses to climate. Biology Letters 8, 590–593.
https://doi.org/10.1098/rsbl.2012.0112
Suh, Y.H., Pesendorfer, M.B., Tringali, A., Bowman, R., Fitzpatrick, J.W., 2020. Investigating
social and environmental predictors of natal dispersal in a cooperative breeding bird.
Behavioral Ecology 31, 692–701. https://doi.org/10.1093/beheco/araa007

Sullivan, B.L., Aycrigg, J.L., Barry, J.H., Bonney, R.E., Bruns, N., Cooper, C.B., Damoulas, T., Dhondt, A.A., Dietterich, T., Farnsworth, A., Fink, D., Fitzpatrick, J.W., Fredericks, T., Gerbracht, J., Gomes, C., Hochachka, W.M., Iliff, M.J., Lagoze, C., La Sorte, F.A., Merrifield, M., Morris, W., Phillips, T.B., Reynolds, M., Rodewald, A.D., Rosenberg, K.V., Trautmann, N.M., Wiggins, A., Winkler, D.W., Wong, W.-K., Wood, C.L., Yu, J., Kelling, S., 2014. The eBird enterprise: An integrated approach to development and application of citizen science. Biological Conservation 169, 31–40. https://doi.org/10.1016/j.biocon.2013.11.003

Sullivan, B.L., Wood, C.L., Iliff, M.J., Bonney, R.E., Fink, D., Kelling, S., 2009. eBird: A citizen-based bird observation network in the biological sciences. Biological Conservation 142, 2282–2292. https://doi.org/10.1016/j.biocon.2009.05.006

Tayleur, C., Caplat, P., Massimino, D., Johnston, A., Jonzén, N., Smith, H.G., Lindström, Å., 2015. Swedish birds are tracking temperature but not rainfall: evidence from a decade of abundance changes. Global Ecology and Biogeography 24, 859–872. https://doi.org/10.1111/geb.12308

Taylor, S.A., White, T.A., Hochachka, W.M., Ferretti, V., Curry, R.L., Lovette, I., 2014. Climate-Mediated Movement of an Avian Hybrid Zone. Current Biology 24, 671–676. https://doi.org/10.1016/j.cub.2014.01.069

Terribile, L.C. et al., 2012. Areas of climate stability of species ranges in the Brazilian Cerrado: disentangling uncertainties through time. NatCon 10, 152–159. https://doi.org/10.4322/natcon.2012.025

Thomas, C.D., Franco, A.M.A., Hill, J.K., 2006. Range retractions and extinction in the face of climate warming. Trends in Ecology & Evolution 21, 415–416. https://doi.org/10.1016/j.tree.2006.05.012

Thorup, K., Tøttrup, A.P., Willemoes, M., Klaassen, R.H.G., Strandberg, R., Vega, M.L., Dasari, H.P., Araújo, M.B., Wikelski, M., Rahbek, C., 2017. Resource tracking within and across continents in long-distance bird migrants. Science Advances 3, e1601360. https://doi.org/10.1126/sciadv.1601360

Thuiller, W., Lafourcade, B., Engler, R., Araújo, M.B., 2009. BIOMOD – a platform for ensemble forecasting of species distributions. Ecography 32, 369–373. https://doi.org/10.1111/j.1600-0587.2008.05742.x

Tingley, M.W., Darling, E., Wilcove, D.S., 2014. Fine- and coarse-filter conservation strategies in a time of climate change. Annals of the New York Academy of Sciences 1322, 92–109. https://doi.org/10.1111/nyas.12484

Tingley, M.W., Monahan, W.B., Beissinger, S.R., Moritz, C., 2009. Birds track their Grinnellian niche through a century of climate change. PNAS 106, 19637–19643. https://doi.org/10.1073/pnas.0901562106

Tobias, J.A., Sheard, C., Pigot, A.L., Devenish, A.J.M., Yang, J., Sayol, F., Neate-Clegg, M.H.C., Alioravainen, N., Weeks, T.L., Barber, R.A., Walkden, P.A., MacGregor, H.E.A., Jones, S.E.I., Vincent, C., Phillips, A.G., Marples, N.M., Montaño-Centellas, F.A., Leandro-Silva, V., Claramunt, S., Darski, B., Freeman, B.G., Bregman, T.P., Cooney, C.R., Hughes, E.C., Capp, E.J.R., Varley, Z.K., Friedman, N.R., Korntheuer, H., Corrales-Vargas, A., Trisos, C.H., Weeks, B.C., Hanz, D.M., Töpfer, T., Bravo, G.A., Remeš, V., Nowak, L., Carneiro, L.S., Moncada R., A.J., Matysioková, B., Baldassarre, D.T., Martínez-Salinas, A., Wolfe, J.D., Chapman, P.M., Daly, B.G., Sorensen, M.C., Neu, A., Ford, M.A., Mayhew, R.J., Fabio Silveira, L., Kelly, D.J., Annorbah, N.N.D., Pollock, H.S., Grabowska-Zhang, A.M., McEntee, J.P., Carlos T. Gonzalez, J., Meneses, C.G., Muñoz, M.C., Powell, L.L., Jamie, G.A., Matthews, T.J., Johnson, O., Brito, G.R.R., Zyskowski, K., Crates, R., Harvey, M.G., Jurado Zevallos, M., Hosner, P.A., Bradfer-Lawrence, T., Maley, J.M., Stiles, F.G., Lima, H.S., Provost, K.L., Chibesa, M., Mashao, M., Howard, J.T., Mlamba, E., Chua, M.A.H., Li, B., Gómez, M.I., García, N.C.,

Päckert, M., Fuchs, J., Ali, J.R., Derryberry, E.P., Carlson, M.L., Urriza, R.C., Brzeski, K.E., Prawiradilaga, D.M., Rayner, M.J., Miller, E.T., Bowie, R.C.K., Lafontaine, R.-M., Scofield, R.P., Lou, Y., Somarathna, L., Lepage, D., Illif, M., Neuschulz, E.L., Templin, M., Dehling, D.M., Cooper, J.C., Pauwels, O.S.G., Analuddin, K., Fjeldså, J., Seddon, N., Sweet, P.R., DeClerck, F.A.J., Naka, L.N., Brawn, J.D., Aleixo, A., Böhning-Gaese, K., Rahbek, C., Fritz, S.A., Thomas, G.H., Schleuning, M., 2022. AVONET: morphological, ecological and geographical data for all birds. Ecology Letters 25, 581–597. https://doi.org/10.1111/ele.13898

Tøttrup, A.P., Klaassen, R.H.G., Kristensen, M.W., Strandberg, R., Vardanis, Y., Lindström, Å., Rahbek, C., Alerstam, T., Thorup, K., 2012. Drought in Africa caused delayed arrival of European songbirds. Science 338, 1307–1307. https://doi.org/10.1126/science.1227548

Travis, J.M.J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., Boulangeat, I., Hodgson, J.A., Kubisch, A., Penteriani, V., Saastamoinen, M., Stevens, V.M., Bullock, J.M., 2013. Dispersal and species' responses to climate change. Oikos 122, 1532–1540. https://doi.org/10.1111/j.1600-0706.2013.00399.x

Twedt, D.J., 2013. Foraging Habitat for Shorebirds in Southeastern Missouri and its Predicted Future Availability. Wetlands 33, 667–678. https://doi.org/10.1007/s13157-013-0422-0

Urban, M.C., Zarnetske, P.L., Skelly, D.K., 2013. Moving forward: dispersal and species interactions determine biotic responses to climate change. Ann N Y Acad Sci 1297, 44–60. https://doi.org/10.1111/nyas.12184

Urli, M., Delzon, S., Eyermann, A., Couallier, V., García-Valdés, R., Zavala, M.A., Porté, A.J., 2014. Inferring shifts in tree species distribution using asymmetric distribution curves: a case study in the Iberian mountains. Journal of Vegetation Science 25, 147–159. https://doi.org/10.1111/jvs.12079

van de Ven, T.M.F.N., McKechnie, A.E., Cunningham, S.J., 2019. The costs of keeping cool: behavioural trade-offs between foraging and thermoregulation are associated with significant mass losses in an arid-zone bird. Oecologia 191, 205–215. https://doi.org/10.1007/s00442-019-04486-x

van Rees, C.B., Hand, B.K., Carter, S., Bargeron, C., Cline, T., Daniel, W., Ferrante, J., Gaddis, K., Hunter, M., Jarnevich, C., McGeoch, M., Morisette, J., Neilson, M., Rozance, M., Sepulveda, A., Wallace, R., Whited, D., Wilcox, T., Kimball, J., Luikart, G., 2022. A Framework to Integrate Innovations in Invasion Biology for Proactive Management. Biological Reviews in press.

van Rees, C.B., Reed, J.M., 2018. Predicted effects of landscape change, sea level rise, and habitat management on the extirpation risk of the Hawaiian common gallinule (Gallinula galeata sandvicensis) on the island of Oʻahu. PeerJ 6, e4990. https://doi.org/10.7717/peerj.4990

Veloz, S., Salas, L., Altman, B., Alexander, J., Jongsomjit, D., Elliott, N., Ballard, G., 2015. Improving effectiveness of systematic conservation planning with density data. Conservation Biology 29, 1217–1227. https://doi.org/10.1111/cobi.12499

Visser, M.E., 2008. Keeping up with a warming world; assessing the rate of adaptation to climate change. Proceedings of the Royal Society B: Biological Sciences 275, 649–659. https://doi.org/10.1098/rspb.2007.0997

Weiner, J., 1994. The Beak of the Finch: A Story of Evolution in Our Time. https://doi.org/10.5860/choice.32-1538

Williams, J.W., Jackson, S.T., Kutzbach, J.E., 2007. Projected distributions of novel and disappearing climates by 2100 AD. PNAS 104, 5738–5742. https://doi.org/10.1073/pnas.0606292104

Willis, S.G., Thomas, C.D., Hill, J.K., Collingham, Y.C., Telfer, M.G., Fox, R., Huntley, B., 2009. Dynamic distribution modelling: predicting the present from the past. Ecography 32, 5–12. https://doi.org/10.1111/j.1600-0587.2008.05711.x

Wilson, S., Smith, A.C., Naujokaitis-Lewis, I., 2018. Opposing responses to drought shape
spatial population dynamics of declining grassland birds. Diversity and Distributions 24,
1687–1698. https://doi.org/10.1111/ddi.12811
Winger, B.M., Auteri, G.G., Pegan, T.M., Weeks, B.C., 2019. A long winter for the Red Queen:
rethinking the evolution of seasonal migration. Biological Reviews 94, 737–752.
https://doi.org/10.1111/brv.12476
Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H., Guisan, A., 2008. Effects of
sample size on the performance of species distribution models. Diversity and
Distributions 14, 763–773. https://doi.org/10.1111/j.1472-4642.2008.00482.x
Wisz, M.S., Pottier, J., Kissling, W.D., Pellissier, L., Lenoir, J., Damgaard, C.F., Dormann, C.F.,
Forchhammer, M.C., Grytnes, J.-A., Guisan, A., Heikkinen, R.K., Høye, T.T., Kühn, I.,
Luoto, M., Maiorano, L., Nilsson, M.-C., Normand, S., Öckinger, E., Schmidt, N.M.,
Termansen, M., Timmermann, A., Wardle, D.A., Aastrup, P., Svenning, J.-C., 2013. The
role of biotic interactions in shaping distributions and realised assemblages of species:
implications for species distribution modelling. Biological Reviews 88, 15–30.
https://doi.org/10.1111/j.1469-185X.2012.00235.x
Wogan, G.O.U., Voelker, G., Oatley, G., Bowie, R.C.K., 2020. Biome stability predicts
population structure of a southern African aridland bird species. Ecology and Evolution
10, 4066–4081. https://doi.org/10.1002/ece3.6175
Wolfson, D.W., Fieberg, J.R., Andersen, D.E., 2020. Juvenile Sandhill Cranes exhibit wider
ranging and more exploratory movements than adults during the breeding season. Ibis
162, 556–562. https://doi.org/10.1111/ibi.12786
Wüest, R.O., Bergamini, A., Bollmann, K., Baltensweiler, A., 2020. LiDAR data as a proxy for
light availability improve distribution modelling of woody species. Forest Ecology and
Management 456, 117644. https://doi.org/10.1016/j.foreco.2019.117644
Xia, J., Wang, J., Niu, S., 2020. Research challenges and opportunities for using big data in
global change biology. Global Change Biology 26, 6040–6061.
https://doi.org/10.1111/gcb.15317
Zurell, D., 2017. Integrating demography, dispersal and interspecific interactions into bird
distribution models. Journal of Avian Biology 48, 1505–1516.
https://doi.org/10.1111/jav.01225
Zurell, D., Thuiller, W., Pagel, J., Cabral, J.S., Münkemüller, T., Gravel, D., Dullinger, S.,
Normand, S., Schiffers, K.H., Moore, K.A., Zimmermann, N.E., 2016. Benchmarking
novel approaches for modelling species range dynamics. Global Change Biology 22,
2651–2664. https://doi.org/10.1111/gcb.13251
1673

1674 **DATA ACCESSIBILITY STATEMENT**

1675 The R code for building Figure 1 is available at the authors github page

1676 [https://github.com/mmacphe/Global_Change_Biomes]. Data layer files were obtained from

1677 public online repositories.

1678

1679 **FIGURES & BOXES**

1680

Figure 1. **Current climate and predicted change in climate of global terrestrial biomes**.

Climographs of (A) temperate and (B) tropical and subtropical global terrestrial biomes.

Projected changes in temperature and precipitation regimes for (C) temperate and (D) tropical

and subtropical global terrestrial biomes. While some areas are predicted to have increased or

decreased precipitation in the future, no areas are expected to have lower or stable temperatures.

Temperature increases are predicted to be higher in temperate biomes, while changes in

precipitation are greater for tropical biomes. Climographs built from The World Wildlife Fund

terrestrial ecoregions (https://www.worldwildlife.org/publications/terrestrial-ecoregions-of-the-

world) and current climate (2021-2040) from CanESM5 middle of the road scenario (SSP3-7.0)

accessed from Worldclim (https://www.worldclim.org/data/cmip6/cmip6climate.html). The

projected change is modeled as the difference between current climate (2021-2040, described

above) and future climate using the worst-case scenario (SSP4-6.0; 2081-2100) from CanESM5

global climate model data (Meinshausen et al., 2020). Species' presence, and thus biodiversity, is

highly influenced by biome type, which shifts from one form to another according to temperature

and rainfall regimes (Whittaker 1975). Projected changes in temperature and precipitation may

cause global terrestrial biomes to shift in space, but spatial gains and losses by different biomes

over time may not occur in synchrony with habitat needs to maintain biodiversity (Dorazio et al.,

2015). R code for producing this figure can be found at

[https://github.com/mmacphe/Global_Change_Biomes].

1700

1701

Figure 2. The "BAM" diagram. The relationship between "B" - biotic, "A" - abiotic, and "M" -

movement (also thought of as "access" and defined by dispersal ability) with respect to the

authors of various niche theories (left). The location of realized, tolerance, and fundamental

niche along the intersection of two hypothetical climate variables (right). Sax et al. (2013)

present an update to niche theory that considers three components of a species niche: 1) the

"realized niche", i.e., conditions within the native range, 2) the "fundamental niche" - conditions,

in which a species could thrive if it were introduced there, and 3) the "tolerance niche" -

conditions, in which individuals could survive, but likely unable to maintain populations over the

long-term. The fundamental niche is the set of appropriate abiotic conditions, and the realized

niche is a smaller area where both abiotic and biotic environments are suitable. Theoretically not

all of the realized niche could be accessible and so the potential range is yet smaller because it

reflects only the area that is suitable and also accessible to the species. Occurrence is not

expected outside of the colored areas. The occupied area is represented by G_0, and G_1 reflects

appropriate conditions that have not or cannot be accessed by the species (as described by Saupe

et al., 2012).

1717

1718

1719   **Figure 3.** Analytical processes for species distribution models. Left: important terms used in

1720   describing distribution model data. Right: Parallel workflows for example correlative versus

1721   mechanistic predictive modelling.

1722

1723

Essential Broad Considerations for Model-building

Common pitfalls of novice modelers:
- The timing of occurrence points does not match when explanatory variable data was collected.
- An artificial species limit is used (e.g., a political boundary) rather than the complete distribution which would reflect more accurately the range of each explanatory variable.
- Projecting beyond the area reachable by the taxon (e.g., predicting into a distant location unlikely to be reached, or across an ecological barrier).

1724

1725 **Box 1**. **Essential Broad Considerations for Predictive Distribution Modeling by Novice**

1726 **Modelers.** Here we summarize this section by highlighting common pitfalls observed by the

1727 authors while acting as reviewers in the peer-review process.

1728

1729 **TABLES**

1730

Table 1. Geospatial context of common occurrence data sources for birds and explanatory

variables. Most occurrence data and explanatory variable data layers are of coarse spatial and

temporal resolution, making them useful for testing hypotheses under the Grinnellian class of

niche. Occurrence data sources with fine spatial and temporal resolution are not as common and

require higher investment for acquiring location data (i.e., GPS tags) or physically tracking

individual birds (i.e., radio-tracking) but could be important for testing hypotheses under the

Eltonian class of niche.

Geospatial data qualities	Examples of occurrence data sources	Examples of explanatory variables

Large spatial area, coarse spatial and/or temporal resolution	•Remote sensing assets (satellite imagery, aerial imagery etc.) •Automated radiotelemetry •Geolocators •Citizen Science: eBird, European Bird Census Council, European Breeding Bird Atlas, South African Bird Atlas Project •Stable hydrogen isotopes •Genetic data •Natural History collections	•Long-term averages in stable climates (e.g., WorldClim) •Land cover (e.g., MODIS, GlobCover, SPOT) •Phenology and vegetation indices (e.g., MODIS, Landsat) •Climate record (e.g., NCDC, ClimateNA, ClimateSA) •Atmospheric weather reanalyses (e.g., ERA – European Centre for Medium-Range Weather Forecasts Re-Analysis, GPM – Global Precipitation Measurement, TRMM – Tropical Rainfall Measuring Mission, NCEP – National Centers for Environmental Prediction)
Small spatial area, coarse spatial and/or temporal resolution	•Citizen science (e.g., Breeding Bird Surveys, Christmas Bird Counts) •Natural history collections •Field notes	

	•Point count surveys	
	•Passive acoustic monitoring	
	•Long-term datasets	
	•Bird banding data	
Large spatial area, fine spatial and/or temporal resolution	•GPS tags	•Digital elevation models (e.g., ASTER) •Vegetation indices (e.g., SPOT)
Small spatial area, fine spatial and/or temporal resolution	•VHF radio-tags	•Personal weather data loggers •Weather station data

1738

APPENDIX 1

Predicting future distributions from current relationships with explanatory variables is

often done using freely available future climate projections (e.g., various "business as usual",

"middle of the road", and "worst case scenario" models downloaded from

https://www.worldclim.org), whereas hindcasting distributions is often done using paleoclimate

models downloaded from the Intercomparison Project Phase II at https://www.pmip2.lsce.ipsl.fr.

Here, we elaborate on the challenge of matching occurrence data with temperature and rainfall

explanatory variable data sources because these are the two most widely used and generally

important environmental factors in species distribution modeling (Bradie and Leung, 2017). The

finest spatial and temporal resolution of environmental data is typically from weather station data

from the nearest station to the research site (e.g., from https://www.wunderground.com/history)

or by collecting weather data alongside occurrence data using data loggers available from

various manufacturers (e.g., Meter, Truebner). The ERA5-Land dataset provides fine scale

(hourly) temperature and precipitation data, but at a 9 km spatial resolution from January 1950 to

present (https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview).

National Centers for Environmental Prediction (NCEP) provides global temperature and

precipitation data (four-times daily) data at a 0.25 x 0.25 degree grid resolution from 1948-

present (https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.surface.html). WorldClim is a

commonly used source of environmental data, and the current version (2.1) provides global

monthly values of min and max temperature, and precipitation averaged from 1970-2000 (Fick

and Hijmans, 2017). Historical monthly min and max temperature, and total precipitation data

used for hindcasting are averaged by decade from 1960-2018 (Fick and Hijmans, 2017; Harris et

al., 2014). BioClim data are 19 quarterly estimates derived from WorldClim monthly

temperature and rainfall values that are thought to be more biologically meaningful than raw

temperature or rainfall data (Busby, 1991). Despite being powerful tools for assessing

macroclimate associations, caution should be used when using the most accessible climate

products to inform conservation decisions by assessing the probability of future range locations

with empirical validations that climate products (e.g., Bioclim variable 1 - annual mean

temperature, Bioclim variable 5 - max temperature of the warmest month) are ecologically

relevant to site occupancy.