Histone acetyltransferase activity of CREB-binding protein is essential for synaptic plasticity in *Lymnaea*

Dai Hatakeyama[1,2,*], Hiroshi Sunada[3], Yuki Totani[4], Takayuki Watanabe[5,6], Ildikó Felletár[1], Adam Fitchett[1], Murat Eravci[1], Aikaterini Anagnostopoulou[1], Ryosuke Miki[2], Takashi Kuzuhara[2], Ildikó Kemenes[1], Etsuro Ito[3,4], György Kemenes[1,*]

[1]Sussex Neuroscience, School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK. [2]Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Tokushima 770-8514, Japan. [3]Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University, Sanuki 769-2193, Japan. [4]Department of Biology, Waseda University, Tokyo 162-8480, Japan. [5]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan. [6]Laboratory of Neuroethology, Sokendai-Hayama, Hayama 240-0193, Japan.

Keywords: Long-term memory; *Lymnaea*; CREB-binding protein (CBP); Histone Acetyl Transferase (HAT); Cerebral Giant Cell (CGC); synaptic plasticity

*Correspondence should be addressed to Dr. Dai Hatakeyama, Tokushima Bunri University, 180 Nishihama-Houji, Yamashiro-cho, Tokushima City, Tokushima 770-8514, Japan, daihatake926@ph.bunri-u.ac.jp; and Prof. György Kemenes, University of Sussex, Brighton, BN1 9QG, United Kingdom, G.Kemenes@sussex.ac.uk.

Abstract

In eukaryotes, CREB-binding protein (CBP), a coactivator of CREB, functions both as a platform for recruiting other components of the transcriptional machinery and as a histone acetyltransferase (HAT) that alters chromatin structure. We previously showed that the transcriptional activity of cAMP-responsive element binding protein (CREB) plays a crucial role in neuronal plasticity in the pond snail *Lymnaea stagnalis*. However, there is no information on the role CBP plays in CREB-initiated plastic changes in *Lymnaea*. In this study, we characterized the *Lymnaea* CBP (LymCBP) gene and investigated the roles it plays in synaptic plasticity involved in regulating feeding behaviors. Similar to CBPs of other species, LymCBP possesses functional domains, such as KIX domain, which is essential for interaction with CREB and was shown to regulate long-term memory (LTM). *In situ* hybridization showed that the staining patterns of LymCBP mRNA in the central nervous system were very similar to those of *Lymnaea* CREB1 (LymCREB1). A particularly strong LymCBP mRNA signal was observed in the Cerebral Giant Cell (CGC), an identified extrinsic modulatory interneuron of the feeding circuit, key to both appetitive and aversive LTM for taste. Biochemical experiments using the recombinant protein of LymCBP HAT domain showed that its enzymatic activity was blocked by classical HAT inhibitors such as curcumin, anacardic acid and garcinol. Preincubation of *Lymnaea* CNSs with these HAT inhibitors blocked cAMP-induced long-term potentiation between the CGC and the follower B1 motoneuron. We therefore suggest that HAT activity of LymCBP in the CGCs is a key factor in synaptic plasticity contributing to LTM after classical conditioning.

Introduction

　　De novo gene transcription is required for consolidation of long-term memory (LTM) (Kandel, 2001) and cAMP-responsive element binding protein (CREB)-dependent gene expression is one of the crucial steps of this process in mammals (Matos et al., 2019; Laviv et al., 2020) as well as invertebrates (Lakhina et al., 2015; Zhou et al., 2015; Hirano et al., 2016). Transcriptional activation requires the recruitment of multifunctional coactivators, resulting in a variety of different types of epigenetic modifications of histones and genomic DNA (Cedar and Bergman, 2009). In CREB-dependent control of gene expression, CREB-binding protein (CBP) functions as a coactivator of CREB. To date, CBP has been reported to be critical for LTM consolidation in mammals (Alarcón et al., 2004; Korzus et al., 2004; Chatterjee et al., 2013) and the gastropod *Aplysia* (Guan et al., 2002; Zhou et al., 2015). One of the most important functions of CBP is due to its histone acetyltransferase (HAT) activity, which stimulates gene transcription (Martinez-Balbás et al., 1998). Several studies have shown that histone acetylation by CBP is necessary for hippocampal long-term potentiation (LTP) (Korzus et al., 2004; Vecsey et al., 2007). Pharmacological inhibition of CBP HAT activity using the inhibitors curcumin and garcinol was reported to block memory consolidation (Zhao et al., 2012; Monsey et al., 2015; Merschbaecher et al., 2016) and memory-associated neuronal plasticity (Maddox et al., 2013a). These studies demonstrated that HAT activity of CBP plays an essential role in regulating synaptic plasticity associated with memory consolidation.

　　The pond snail *Lymnaea stagnalis* is a widely used organism to understand evolutionarily conserved molecular mechanisms of the consolidation of LTM for taste-related associations (Kemenes et al., 2006; Hatakeyama et al., 2013a; Murakami et al., 2013; Totani et al., 2020; Nakai et al., 2020a; 2020b). Sadamoto et al. (2004) first succeeded in the cloning of an isoform of CREB from the central nervous system (CNS) of *Lymnaea* and defined it as LymCREB1. They identified 7 different isoforms of LymCREB1 by alternative splicing and

found that aversive taste conditioning significantly increased LymCREB1 gene expression (Sadamoto et al., 2010). Ribeiro et al. (2003) showed that appetitive taste conditioning selectively increased phosphorylated LymCREB1 in the 'learning ganglia' (the cerebral and buccal ganglia) of the *Lymnaea* CNS.

An identified extrinsic modulatory neuron type of the feeding system, the cerebral giant cell (CGC) was reported to play crucial roles in LTM after both aversive and appetitive taste conditioning (Kemenes et al., 2006; Ito et al., 2012; Nikitin et al., 2013). Injection of LymCREB1 siRNA into CGCs reduced the amplitude of excitatory postsynaptic potential (EPSP) in monosynaptic follower neurons of the CGCs, the B1 motoneurons (Wagatsuma et al., 2006), suggesting that regulation of gene expression by LymCREB1 was required for synaptic enhancement in memory consolidation. Based on these previous findings, we hypothesized that HAT activity of the *Lymnaea* CBP (LymCBP) plays an important role in the LymCREB1 initiated molecular processes of synaptic consolidation.

In the present study, we first cloned the LymCBP cDNA from the *Lymnaea* CNSs and identified neurons expressing the LymCBP mRNA. Focusing on the CGC, we pharmacologically and electrophysiologically analyzed the relationship between the HAT activity of LymCBP and the synaptic plasticity involved in the aversively conditioned feeding behavior of *Lymnaea*. Our findings provide new insights into the functions of LymCBP in synaptic plasticity underlying the consolidation of associative LTM.

Materials and Methods

Molecular cloning of LymCBP. To clone LymCBP, a series of degenerate PCR was performed with TaKaRa Ex Taq® (Takara Clontech) and primers, which were designed at the basis of highly conserved domains, such as Taz1 (transcriptional adapter zinc-binding 1), KIX domain and Taz2 domain, of *Aplysia* CBP (ApCBP; GenBank accession number: AY064470). After the sequential analyses of the Taz1, KIX and Taz2 domains of LymCBP, we performed 5' and

3'RACE, and the PCR for the internal region between Taz1 and KIX domains and between

KIX and Taz2 domains. The FirstChoice® RLM-RACE Kit (ThermoFisher) was used for

amplification of 5' and 3' ends of LymCBP. All PCR products were subcloned with TA

Cloning® Kit (ThermoFisher) or pGEM®-T Easy Vector System (Promega). Nucleotide

sequences of primers were summarized in Table 1 (No. 1-14).

Phylogenetic tree of LymCBP. To construct a phylogenetic tree of the CBP/p300 proteins, we

aligned the full-length deduced amino acid sequence of the *Lymnaea* CBP with those of the

known homologs of other species (listed with each Accession Number in Table 2) by using

the MUSCLE algorisms (Edgar, 2004) on the Geneious (v9.1) program (available from

http://www.geneious.com/). Maximum likelihood tree was constructed from the aligned

sequences using the MEGA 6 program (Tamura et al., 2013) with default settings of the

program. 1000 bootstrap replications were conducted to evaluate the reliabilities of the

reconstructed trees. The obtained tree was visualized with the FigTree (v1.4.2) program

(available   from   http://tree.bio.ed.ac.uk/software/figtree/).   A   CBP   homolog   of   the

Choanoflagellate *Salpingoeca rosetta* was used as an outgroup.

TaqMan-based quantitative PCR (qPCR). To quantify the absolute copy number of LymCBP

mRNA in several *Lymnaea* tissues, we performed TaqMan-based qPCR by establishing RNA

standard curves derived from a dilution series of known template concentrations. The

procedures for this experiment were modified from previous reports (Hatakeyama et al.,

2004b, 2006; Wagatsuma et al., 2005). Partial LymCBP gene (from 303rd to 722nd

nucleotides, which contains the open reading flame of the *N*-terminal region) was amplified

by RT-PCR with primers shown in Table 1 (No. 15, 16), and the PCR product was subcloned

into pGEM®-T Easy Vector (Promega). The plasmid was purified with Plasmid Midi Kit

(QIAGEN) and digested with restriction enzyme *Spe* I. The LymCBP RNA was synthesized

with MAXIscript^{TM} (ThermoFisher), purified with RNeasy$^®$ Mini Kit (QIAGEN), quantified

the concentration, and used as the standard RNA. Serially diluted standard RNA ($5 \times 10^1 - 5$

$\times 10^6$ copies/µl) were reverse-transcribed in 10 µl of reaction mixture to prepare the first

strand cDNA with 2.5 µM of the specific primer for LymCBP diluted to 1:5 and used as the

standard cDNAs.

Total RNA from different types of *Lymnaea* tissue (CNS, buccal mass, penis,

ovotestis, gut and mantle) was purified with TRIzol^{TM} Reagent (ThermoFisher). Reverse

transcription (RT) of standard series ($5 \times 10^1 - 5 \times 10^6$ copies) and total RNA of *Lymnaea*

tissues (each 300 ng) were performed in the buffer containing 0.2 µg yeast tRNA, 1× PCR

Buffer II (ThermoFisher), 5.5 mM MgCl$_2$, 0.5 mM dNTP, 0.2 µM gene specific primer for

LymCBP, 0.3 unit/ml Prime RNase inhibitor (Eppendorf), and 0.9 units/ml MultiScribe^{TM}

Reverse Transcriptase (ThermoFisher) for 10 min at 25^{o}C, for 30 min at 48^{o}C, and for 5 min

at 95^{o}C. All RT samples were diluted to 5 times by adding RNase-free distilled water. The

diluted RT solution of samples and standards (1 µl) were added to PCR-reaction solution

[final concentration: 1× TaqMan$^®$ Gene Expression Master Mix (ThermoFisher), 50 nM each

primer, and 50 nM probe]. Nucleotide sequences of primers and probes were summarized in

Table 1 (No. 17-20). Reaction was carried out at 95^{o}C for 10 min, 45 cycles of 95^{o}C for 15

sec, and 60^{o}C for 1 min with real-time PCR system (StepOnePlus Real-Time PCR System,

ThermoFisher). In the assay, several doses of standard cDNA ($1 \times 10^1 - 1 \times 10^6$ copies) were

applied in triplicate to estimate inter-assay coefficients of variation between runs.

In situ *hybridization of LymCBP*. The procedures for this experiment were modified from

previous reports (Hatakeyama et al., 2004a; 2006; 2010). Non-functional region (from 278th

to 1146th of nucleotide sequence) of LymCBP was isolated by RT-PCR (see Table 1 (No. 21,

22) for nucleotide sequences of primers). After insertion of this PCR product into pGEM$^®$-T

Easy Vector, the plasmid was linearized with *Eco*R V and *Bam*H I to generate

digoxigenin-labeled antisense and sense RNA probes, respectively. The procedures for

generating the probes, hybridization and coloring were the same as those that were previously

reported (Hatakeyama et al., 2004a; 2006).

Western blotting, co-precipitation and mass spectrometry.

Experimental animals. Pond snails, *Lymnaea stagnalis,* were bred at the University of Sussex,

Brighton, United Kingdom and were maintained in large holding tanks containing copper-free

water at 18-20^{0}C on a 12-hour light - dark cycle. The animals were fed on a vegetable-based

fish food (Tetra-Phyll; TETRA Werke, Melle, Germany) twice a week and lettuce three times

a week.

Preparation of snail CNS samples for Western blotting, co-precipitation and mass

spectrometry. The whole CNS of four-month-old snails were dissected. Briefly, the shell of

the snail was removed, and the preparation was then pinned down in a Sylgard-coated dish

containing HEPES buffered saline and dissected under a stereomicroscope (E-Zoom6,

Edmund Optics, Barrington NJ, USA). The CNS was accessed by a dorsal body incision and

isolated from the buccal mass by severing all the peripheral nerves and immediately placed in

Eppendorf tubes on dry ice and stored at -80^{o}C.

Western blotting. *Lymnaea* CNS lysates were prepared in RIPA buffer. Protein (10-20 µg) was

electrophoresed on homemade 6% Bis-Tris protein gels and transferred to polyvinylidene

difluoride (PVDF) membrane. The polyclonal antibody to anti-human-CBP (C20, sc-583,

Santa Cruz) was used at 1:500 dilution in 4% skimmed milk in TBS-Tween overnight at 4°C

with rotation. Following washes with TBS-Tween (10 min ×3), the membranes were

incubated with the anti-rabbit-IgG conjugated with horseradish peroxidase (Sigma Aldrich) at

room temperature at 1:10,000 dilution for 2 hr. Finally, the membranes were cleaned from the

excess of secondary antibody by washing with TBS-Tween (10 min ×3) and TBS (10 min ×1).

Detection of blot was carried out with a LumiSensor^{TM} HRP Substrate Kit (GenScript

Technology).

<u>Immunoprecipitation assay and immunoblotting.</u> Sixty pooled *Lymnae*a CNSs were

homogenised in RIPA lysis buffer supplemented with 2 mM PMSF, 1 mmol/L Na_3VO_4 and

protease inhibitors, (Santa Cruz, sc-24948) by using a micropestelle.   50 µl of the

pre-immunoprecipitated lysate was saved as "input" to be used for immunoblotting. The

homogenate was pre-cleared in a Pierce Spin column (Thermofisher Scientific, 69705)

containing 40 µl of Protein A/G PLUS-Agarose (Santa Cruz, sc-2003) for one hour at 4^0C on

a rotor. Four Pierce Spin columns containing 40 µl of Protein A/G PLUS-Agarose suspension

were prepared. Three of the Pierce Spin columns were used to prepare Antibody/Agarose

complex and one of the columns was used as a negative control without containing the

antibody. 500 µl of HEPES buffer was added to all of the Protein A/G PLUS-Agarose spin

columns. For the Antibody/Agarose complex preparation, 10 µg of anti-mouse CREB2

antibody (Santa Cruz, sc-390063) was added to each of the Protein A/G PLUS-Agarose Spin

column.   The columns were incubated on a rotor at RT for 3 hrs. The columns were

centrifuged at 2500 RPM for 1 min and the flow-through was removed. The columns were

then washed/centrifuged four times at 2500 RPM with HEPES buffer at 4^0C. After the final

wash, 500 µg of CNS homogenised lysate was transferred into each column containing the

Antibody/Agarose complex and the one column without containing the antibody. The

columns were placed on a rotor and incubated overnight at 4^0C. The following day, the

columns were centrifuged at 2500 RPM for 1 min and the lysate was removed. The columns

were washed/centrifuged four times at 2500 RPM with HEPES buffer. The eluates for each

column were eluted in 50 µl of Glycine, pH 2.8. The pH of the eluates was neutralized with 5

µl of 1M Tris, pH 9.5.   The IP products were pooled together and 100 µl of the IP product

was used for LC/MS analysis and 45 µl of the IP product was prepared for Western Blotting.

45 µl of the IP product was resuspended in 15 µl of 4 x Loading sample buffer and boiled at

90^0C for 10 mins. Immunoprecipitates, negative control and 25 µg of the

pre-immunoprecipitated lysate (5% input) were resolved in 10% SDS-polyacrylamide gel and

transferred to a nitrocellulose membrane.   Non-specific binding sites were blocked with 5%

milk in TBS solution with 0.1% Tween-20 (1xTBST) for 1 hour at RT.   Membranes were

incubated with anti-mouse CREB2 antibody (1:1000, Santa Cruz, sc-390063) overnight at 4^0C.

The following day, the membrane was washed three times for 10 mins in TBST and incubated

with highly cross-adsorbed Infra-Red Dye 680RD goat anti-mouse secondary antibody (1:5000,

LI-COR, 926-68070) for 1 h at RT, followed by two washes with 1xTBST and a final wash of

1xPBS. Membranes were developed using the LI-COR Odyssey Fc Fluorescence Imaging

system.

<u>Protein precipitation for detergent removal.</u> In order to remove the detergents used   in our

immunoprecipitation protocol, which are incompatible with the downstream MS analysis we

have used the methanol/chloroform precipitation method as previously described (Wessel and

Flugge, 1984). Protein precipitates were then resuspended in 50 μl of 5 X Invitrosol^{TM}

LC/MS Protein Solubilizer. The samples were incubated at 60^oC for 5 mins, vortexed and

incubated at 60^oC for another 10 mins. 200 μl of 25 mM ammonium bicarbonate (ABC Buffer)

was added per sample in order to obtain an Invitrosol^{TM} concentration of 1X. The samples were

separated into 5 tubes containing 50 μl of sample per tube.

<u>In-solution Trypsin digestion with the MS detergents.</u> Samples were reduced in 10 mM

Dithiothreitol (DTT) for 30 mins at RT and cysteines were alkylated in the dark in 60 mM

Iodoacetamide (IAA) for 20 min at RT. 2 μl of Trypsin/Lys-C mix (0.5μg/μl, Promega, V5073)

was added to each sample and incubated for 3 hrs at RT in the dark.   After 3 hrs, 150 μl of ABC

buffer per tube was added and incubated overnight at RT.   The following day, the digestion was

stopped by adding 200 μl of 5% Acetonitrile, 3% Trifluoroacetic acid. Tryptic peptides were

desalted with stage tips as previously described (Rappsilber et al., 2003) The desalted peptides

were concentrated in a Speed-Vac for 10 mins at low power.   The peptides were reconstituted

with 15 μl of 5% Acetonitrile, 0.1% Formic Acid and transferred to a microplate for LC/MS

analysis.

<u>LC/MS analysis.</u> Peptides were separated by reverse-phase chromatography using a Dionex

Ultimate 3000 nanoLC (Dionex/Thermo Fisher Scientific) on a C18 capillary column (Acclaim

PepMap100 C18, 2 μm, 100 Å, 75 μm i.d.×20 cm / Thermo Fisher Scientific) at an eluent flow

rate of 300 nl/min. Mobile phase A contained 5% acetonitrile, 0.1% formic acid and mobile

phase B contained 80% acetonitrile 0.1% formic acid. The column was preequilibrated with 5%

mobile phase B followed by an increase to 65% mobile phase B in 75 min. The eluted peptides

were ionized online by electrospray ionization (ESI) and transferred into an LTQ Orbitrap XL

mass spectrometer (Thermo Fisher Scientific) which was operated in the positive mode to

measure full scan MS spectra (300–1700 m/z in the Orbitrap analyzer at resolution R = 30,000)

followed by isolation and fragmentation (MS/MS) of the ten most intense ions (in the LTQ

iontrap) by collision-induced dissociation (CID).

<u>MS/MS data analysis.</u> The analysis of the raw MS and MS/MS Spectra was performed using

the MaxQuant software (version. 1.6.3.4). Initial maximum precursor and fragment mass

deviations were set to 7 ppm and 0.5 Da, respectively. Variable modification (oxidation of

methionine and N-terminal acetylation) and fixed modification (cysteine

carbamidomethylation) were set for the search against a UniProt FASTA formatted database of

the taxonomic clade *euthyneura* (including *Lymnaea stagnalis*). Trypsin with a maximum of

two missed cleavages was chosen for digestion. The minimum peptide length was set to 7

amino acids and the false discovery rate (FDR) for peptide and protein identification was set to

0.01.

Recombinant protein of LymCBP. The partial recombinant protein of the LymCBP HAT

domain (from 1240th arginine to 1641st lysine, GenBank accession number BAE02656.1)

with His$_6$-tag was synthesized in *Escherichia coli* and purified as previously described with

slight modifications (Hatakeyama et al., 2014, 2018a). LymCBP catalytic domain was

amplified by RT-PCR with cDNA prepared from *Lymnaea* CNSs, PrimeSTAR® HS DNA

Polymerase (Takara Clontech) and primers, whose nucleotide sequences were summarized in

Table 1 (No. 23, 24). The PCR product was ligated into pET28a(+) plasmid (Novagen), which

was linearized with *Nhe* I and *Bam*H I, using In-Fusion® HD Cloning Plus kit (Takara

Clontech). Then, the Rosetta-gami™ 2(DE3) competent cells (Novagen) was transformed

with this ligated plasmid. Expression of the partial recombinant protein of the LymCBP

catalytic domain was induced by treatment with 0.32 mM of isopropyl

β-D-thiogalactopyranoside (IPTG) in TBG-M9 medium [0.8% Bacto Tryptone (BD) and 0.4%

NaCl], followed by purification using nickel-nitrilotriacetic acid (Ni-NTA) agarose resin

(QIAGEN). The recombinant protein was desalted with PD-10 column (Sephadex™ G-25M,

GE Healthcare), purified using a cation exchange column [HiTrap carboxymethyl (CM) FF

column (GE Healthcare)] installed in the ÄKTAprime plus system (GE Healthcare), and

condensed with Amicon® Ultra-15 column (Merck Millipore).

Acetylation assays using radioisotope-labeled acetyl-CoA. The procedures followed to

perform this experiment were modified from previous reports (Hatakeyama et al., 2018a;

2018b). The recombinant LymCBP HAT domain (200 ng) were incubated with the histone H1

purified from calf thymus (Sigma-Aldrich, 1 μg) and 7.4 kBq of [^{14}C]-acetyl-CoA (Moravek,

Inc.) for 4 hr in buffer containing 50 mM Tris-HCl (pH 8.0), 10% glycerol, 1 mM

dithiothreitol, and 10 mM sodium butyrate. Curcumin (MedChemExpress), anacardic acid

(MedChemExpress) and garcinol (Enzo) prepared in dimethyl sulfoxide (DMSO) were

premixed at each 50 μM. Reactions were separated in 14% SDS-PAGE gels. To detect β

radiation released from radiocarbon, the gels were dried on pieces of filter papers and exposed

on imaging plates for several days. Signals were observed using a fluoroimage analyzer

(FLA-2000; Fuji Film).

Intracellular recording. The whole CNS, including the cerebral ganglia and the buccal

ganglia was dissected out from the snail in *Lymnaea* saline [50 mM NaCl, 1.6 mM KCl, 2.0

mM $MgCl_2$, 3.5 mM $CaCl_2$, and 10 mM HEPES (pH 7.9)]. The isolated CNS was

immobilized on a Sylgard-coated dish using stainless-steel pins. In order to physically

separate the cerebral giant cells and B1 motoneurons, a Vaseline dam was constructed around

the cerebral ganglia. This enabled us to apply a solution that contains the inhibitors to the

CGCs. The CGC and the B1 motoneuron were impaled with glass microelectrodes filled with

3 M KCl giving tip resistances of 20–50 MΩ. A train of 5 spikes in the CGC, which was

produced by a current injection (0.8–1.8 nA) for 1 sec, evoked a single large compound EPSP

in the B1 motoneuron (electric stimulator: SEN-7203, Nihon Kohden, Tokyo, Japan;

intracellular recording amplifier: MEZ-8300, Nihon Kohden; AD converter: DIGIDATA

1322A, Axon Instruments, Foster City, CA, USA). We used the neurons located in the

ipsilateral side.

Injection of cAMP or 5'AMP to the CGC. Based on a previous study (Nakamura et al., 1999),

cyclic AMP (cAMP) or 5'AMP (Sigma-Aldrich), which was filled into a glass microelectrode

(50–100 MΩ) as a 200 mM solution dissolved in 20 mM Tris buffer (pH7.5), was injected

into the CGC by passing hyperpolarizing current pulses (50 msec on, 50 msec off) of 4 nA for

20 min. Before and after the injection of hyperpolarizing current into the CGC, the EPSP

recorded in the B1 motoneuron is not changed by activation of the CGC. The data were

recorded in the same preparation at 30 min and 3 hr after cAMP or 5'AMP injection. The data

at 0 hr (i.e., pre) were recorded before cAMP injection. For the analysis of EPSP changes

recorded in the B1 motoneuron, the peak amplitude of the EPSP was measured.

Application of HAT inhibitors to the CGC. We used the following three different inhibitors of

HATs: curcumin (Sigma), anacardic acid (Sigma) and garcinol (Enzo). Each of these drugs

was dissolved in DMSO as a stock solution at 67.9 mM (curcumin), 50 mM (anacardic acid)

and 20 mM (garcinol). Stock solutions were diluted 1000-fold in *Lymnaea* saline then applied

to the CGC area where the cerebral ganglia were surrounded by Vaseline wall and incubation

for 1 hr. After incubation with each drug, intracellular recordings for the measurement of

EPSPs were made. The data was expressed as the mean ± SEM. Significant differences at $p <$

0.05 between pre-injection, 30 min and 3 hr after injection cAMP or 5'AMP were examined

by one-way repeated measures ANOVA and post hoc Tukey's test. For the normalized EPSP

amplitude, the values after 30 min and 3 hr for each group compared to the pre-injection were

examined with one-way repeated measured ANOVA and post hoc Tukey's test.

Results

Properties* of the *LymCBP gene and protein

The arrangement of the functional domains in LymCBP was the same as that of other

p300/CBPs (Fig. 1A). The full sequence of LymCBP (GenBank accession number:

AB217914) was about 75% identical to *Aplysia californica* CBP (ApCBP; GenBank

accession number: AAL54859) in amino acid sequence level. In functional domains, such as

Taz1 domain, KIX domain, bromodomain, HAT domain, and Taz2 domain of LymCBP, the

amino acid sequences were individually over 90% identical to those of ApCBP and human

CBP (GenBank accession number: U47741) (Fig. 1B-D). Comparing amino acid sequences

among LymCBP, human CBP and human p300, the functional amino acid residues in

bromodomain for interacting with acetylated histones (Dhalluin et al., 1999; Mujtaba et al.,

2004; Plotnikov et al., 2014; Xu et al., 2017), those in HAT domain for intramolecular

interactions (Delvecchio et al., 2013; Ortega et al., 2018; Zhang et al., 2018), and interaction

with acetyl-CoA (Liu et al., 2008; Maksimoska et al., 2014) were highly conserved (Table S1).

Molecular phylogenetic analysis of the p300/CBP proteins revealed that *Lymnaea* CBP is

closely related to CBP proteins found in other molluscan species such as *Aplysia* and the

oyster *Crassostrea giga* (Fig. 2).

Localization of LymCBP mRNA in the **Lymnaea** *CNS*

Absolute copy numbers of LymCBP mRNA in whole CNS and other tissues were

analyzed by quantitative PCR (qPCR) (Fig. 3). More than 15,000 copies of LymCBP mRNA

were contained in 6 ng of total RNA of CNSs (Fig. 3). This value was 1.4 to 2.9 times higher

compared to other tissues (Fig. 3).

In situ hybridization showed the localization of LymCBP mRNA in the *Lymnaea*

CNSs ($n = 8$). Weak intensity of the positive signal was detected in some neurons in the

buccal ganglia (Fig. 4A). The neurons localized at the ventral edge of the ganglia seemed to

be parts of the B4 cluster (B4CL) cells. Strong signal was observed in the CGCs in the

cerebral ganglia (Fig. 4B,C). The same signal was also seen in identifiable large neurons, the

RPeD1 (Fig. 4D) and the LPeD1 (Fig. 4E), in the pedal ganglia. Notably, the staining patterns

of LymCBP mRNA in the buccal, cerebral and pedal ganglia were very similar to those of

LymCREB1 (Sadamoto et al., 2004). The same positive signal of LymCBP was also detected

in unidentified neurons in parietal and visceral ganglia (Fig. 4F). No positive signals were

detected in negative control experiments with sense LymCBP cRNA probe (data not shown).

Existence of the protein form of LymCBP

Western blotting demonstrated the existence of the protein form of LymCBP in the

CNS. The epitope of the anti-human-CBP antibody is the *C*-terminal region of human CBP,

which is highly conserved in LymCBP (Fig. 1D). The western blotting method detected two

positive bands (Fig. 5A). The larger band, indicated with a filled arrowhead around the

marker of 245 kDa, was considered to be LymCBP, whose molecular weight was estimated as

252.3 kDa from the amino acid sequence (Fig. 1). The smaller band shown with an open

arrowhead may be an isoform of LymCBP (Fig. 5A). In the EMBL-EBI database, twenty

transcript variants of CBP are found in human, and, of these isoforms, seven transcripts can be

translated (Gene: "CREBBP", ENSG00000005339). Therefore, it is suggested that *Lymnaea*

also expresses the smaller isoform of CBP.

Phosphorylation of CREB1 at Ser 133 is triggering its kinase inducible domain

(KID) to bind to the KID interacting domain (KIX) of CREB binding protein (CBP) (Mayr

and Montminy, 2001; Radhakrishnan et al., 1997; Shaywitz et al., 2000). To identify the

interaction of CBP with CREB proteins in our model organism *Lymnaea stagnalis* we have

utilized immunoprecipitation (IP) and liquid chromatography mass spectrometry (LC/MS).

Although in pilot experiments we found that none of our CREB1 antibodies worked

well for IP assays with total CNS lysates from *Lymnaea stagnalis*, previous studies have

shown that human CREB2 (human activating transcription 4, hATF4), a transcriptional

repressor of CREB1, can directly interact with multiple domains of the CBP including i. the

KIX domain, ii. a region that contains the third zinc finger and the E1A-interacting domain, iii.

the C-terminal region containing the p160/SRC-1-interacting domain and iii. the histone

acetyltransferase domain which can directly interact with histone acetyltransferase p300

transcriptional coactivator of CBP (Liang and Hai, 1997). In addition, CBP and p300 can

acetylate CREB2 in its bZIP domain and enhance its transcriptional activity (Gachon et al.,

2002; Lassot et al., 2005; Liang and Hai, 1997). In order to confirm these reported

interactions of CBP and CREB2, we have performed IP assays with a specific antibody for

CREB2 and CNS homogenates from *Lymnaea stagnalis* (Fig. 5B), followed by identification

of the CREB2 interaction partners by LC/MS (Fig. 5C). As shown in Fig. 5C, after LC/MS,

the data analysis of the CID fragmentation spectrum from a peptide with the mass of 559.81

m/z revealed the amino-acid sequence of a unique peptide (KVFNPDELR) from the

UniprotKB entry E2QC33 of *Lymnaea stagnalis*, validating that *Lymnaea* CBP is an

interacting partner of CREB2 in the CNS. Importantly, these results further confirmed that

LymCBP is expressed in the *Lymnaea* CNS.

Effects of HAT inhibitors on LymCBP enzymatic activity

To investigate the enzymatic properties of LymCBP as a functional HAT, we

produced a partial recombinant protein of its HAT domain (from 1240th to 1641st, Fig. 1D)

and investigated its acetyltransferase activity. The recombinant protein, to which an His_6-tag

was attached at its *N*-terminus, could be purified by precipitation using Ni-NTA agarose resin.

To remove contaminated proteins derived from bacterial cells, the recombinant protein

solution was desalted and then purified by cation-exchange chromatography. This purification

procedure was successful, and only the bands of the recombinant protein were observed in the

12th to 19th fractions (arrowhead in Fig. 6A). However, the concentrations of these

preparations were too low to be measured by Bradford protein assay. Then, the eight fractions

were mixed and treated with a centrifugal filter to condense the recombinant protein, resulting

its concentration increased to around 80 ng/μL (lane 2 in Fig. 6B), which was sufficient for

the biochemical experiments.

We first determined the incubation temperature for the *in vitro* acetylation assay. A

reaction mixture containing recombinant LymCBP HAT domain, bovine histone H1, and

[^{14}C]-acetyl-CoA were incubated for 4 hr at different temperatures; 15ºC, 20ºC, 25ºC, 30ºC

and 37ºC. Separation by SDS-PAGE and Coomassie Brilliant Blue (CBB) staining of the gel

showed that same amount of histone H1 was loaded (lanes 3-7 in Fig. 6C). However, the band

of recombinant LymCBP HAT domain was notably decreased after incubation at 37ºC,

suggesting that LymCBP was degraded (lane 7 in Fig. 6C). By autoradiography, the positive

signal indicating acetylation of histone H1 was observed when incubated at 15ºC, 20ºC, 25ºC

and 30ºC (lanes 8-11 in Fig. 6C), and this signal was almost eliminated after incubation at

37ºC (lane 12 in Fig. 6C). Therefore, we decided to incubate samples at 30ºC in following

acetylation assays.

Next, the effects of HAT inhibitors, namely curcumin, anacardic acid and garcinol,

on the HAT activity of LymCBP was investigated. After adding each HAT inhibitor (each at

50 μM final concentration) or DMSO as negative control, samples were incubated for 4 h at

30°C and separated by SDS-PAGE. Although the amount of each protein in each tube should

be same, CBB staining showed that the band intensities of histone H1 were decreased by

incubating with curcumin and garcinol (lanes 14 and 16 in Fig. 6D), suggesting that these

inhibitors degraded the protein. Autoradiography showed that LymCBP acetylated histone H1

without any HAT inhibitors (lane 17 in Fig. 6D). In the presence of HAT inhibitors, LymCBP

HAT activity was significantly blocked and acetylation signals were eliminated (lanes 18-20

in Fig. 6D). These results showed that similar to human CBP, LymCBP had HAT activity, and

this enzymatic activity was significantly blocked by HAT inhibitors.

Synaptic plasticity between a buccal motoneuron and its regulatory interneuron induced by

cAMP

The isolated *Lymnaea* CNS was incubated with DMSO saline for 1 h before the

injection of cAMP into the CGC cell body. We tested the changes in the EPSP in the B1

motoneuron by activation of the CGC after the cAMP injection. We measured the amplitudes

of the compound EPSPs 30 min and 3 h after the onset of cAMP injection. The CGC was

depolarized by current injection to fire action potentials, evoking the compound EPSP in the

B1 motoneuron. We used a a train of 5 spikes in the CGC to evoke large compound EPSPs in

the B1 motoneuron because the single EPSPs in the B1 motoneuron were very small

(approximately 1 mV). These compound EPSPs at 30 min and 3 h after in preparations with

cAMP-injected CGCs were about twice as large as pre injection EPSPs [Fig. 7A, $F_{(2,18)}$ =

10.35, pre injection vs 30 min: $p < 0.05$; pre injection vs 3 hr: $p < 0.05$, $n = 10$]. In contrast,

there were no significant changes in the 5'AMP-injected preparations [Fig. 7B, $F_{(2,12)}$ =

1.192, pre injection vs 30 min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$, $n = 7$]. Thus, the effect

of HAT inhibitors on the compound EPSPs 30 min and 3 hr after cAMP injection were

examined in the following experiments.

Effects of HAT inhibitors on the EPSP in the B1 motoneuron

HAT inhibitors were shown to possess significant inhibitory effects on HAT activity

of LymCBP (Fig. 6C). Therefore, we examined the effects of the application of these HAT

inhibitors to CGCs on the compound B1 EPSPs that were found to be increased after cAMP

injection into the CGC. As reported previously (Nakamura et al., 1999; Sadamoto et al., 2004),

the EPSP in the B1 motoneuron is facilitated by injection of cAMP into the CGC (Fig. 7A).

This increase in EPSP amplitude was not observed when 5'AMP was injected into CGCs

instead of cAMP (Fig. 7B). Next, we investigated changes to the cAMP-induced EPSP in the

presence of the HAT inhibitors curcumin, anacardic acid and garcinol. Incubation with 67.9

μM curcumin of the cerebral ganglion for 1 h did not change the amplitude of the compound

EPSPs in the B1 motoneuron (Fig. 7C). After 1-hr incubation with curcumin prior to the

cAMP injection there was no significant change in the amplitude of the compound EPSPs 30

min and 3 hr after the cAMP injection compared with pre injection [Fig. 7C, $F(2,22) = 6.001$,

pre injection vs 30 min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$, $n = 12$]. However, a

significant difference was observed between 30 min and 3 hr. The incubation with 50 μM

anacardic acid on the cerebral ganglion for 1 hr that did not enlarge of the compound EPSPs

in the B1 motoneuron (Fig. 7D). The 1-hr incubation with anacardic acid prior to the cAMP

injection, in this group, there was no significant differences in the amplitude of the compound

EPSPs 30 min and 3 hr after the cAMP injection compared with pre injection [Fig. 7D,

$F(2,10) = 1.563$, pre injection vs 30 min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$, $n = 6$]. The

incubation with 20 μM garcinol on the cerebral ganglion for 1 hr that did not enlarge of the

compound EPSPs in the B1 motoneuron (Fig. 7E). The incubation for 1 hr with garcinol prior

to the cAMP injection, in this group, there was no significant differences in the amplitude of

the compound EPSPs 30 min and 3 hr after the cAMP injection compared with pre injection

[Fig. 5E, $F(2,14) = 0.01214$, pre injection vs 30 min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$,

$n = 8$].

463   The results from all these groups are summarized in Fig. 7F. The effect of HAT

inhibitors on increasing compound EPSPs amplitude was tested by taking a 0 min baseline

recording followed by a 30 min and a 3 hr recording after cAMP was injected. The compound

EPSPs were normalized to the initial baseline value. Incubation with the HAT inhibitors

before the injection of cAMP, did not result in increased compound EPSPs amplitudes, as

seen in the DMSO saline treatment group. In the host hoc Tukey's test, there were significant

differences in the following comparisons: DMSO saline treatment cAMP injection group; pre

injection vs 30 min; pre injection vs 3 hr, curcumin treatment group; 30 min vs 3 hr).

Discussion

473   In the present study we, for the first time, identified a conserved CREB-binding

protein in the CNS of *Lymnaea* (LymCBP) and showed the importance of its HAT activity for

synaptic plasticity between an identified interneuron (CGC) and a postsynaptic motoneuron

(B1) of the feeding system. The CGC is a key interneuron for consolidating both aversive and

appetitive LTM for taste, and both LymCBP and LymCREB1 were transcribed in this neuron

(Fig. 4). A previous report showed that inhibiting functions of LymCREB1 by pre-injection of

excess amount of a CRE oligonucleotide and siRNA of LymCREB1 into the CGC blocked

cAMP-induced synaptic facilitation between the CGCs and the B1 motoneurons (Sadamoto et

al., 2004; Wagatsuma et al., 2006), and suggested a crucial role of LymCREB1 as a

transcriptional factor in synaptic facilitation. Here, inhibition of LymCBP HAT activity using

HAT inhibitors similarly blocked the synaptic potentiation (Fig. 7), suggesting that this

synaptic facilitation requires the co-operation of LymCREB1 and LymCBP HAT activity.

Gene structure of LymCBP

We succeeded in the cloning of the full-length cDNA of LymCBP from the *Lymnaea*

CNS. Comparing its deduced amino acid sequence with those of *Aplysia* and human CBPs,

several functional domains were found to be highly conserved (Fig. 1). This result suggests

that LymCBP has similar functions to mammalian CBP and p300 (Fig. 2). The KIX domain of

CBP associates with the kinase-inducible domain (KID) of CREB, and they construct a

complex together (Chrivia et al., 1993). KID, which is also called as P-box, is highly

conserved in LymCREB1 (Sadamoto et al., 2004). High conservation of the KIX domain in

LymCBP and KID in LymCREB1 suggested that LymCBP and LymCREB could interact via

these domains. Additionally, a previous report showed that the KIX domain of CBP is

involved in the formation of spatial memory following hippocampus-dependent learning

(Chatterjee et al., 2020). Here, we demonstrated that these two mRNAs were co-localized in

the CGCs, key neurons for LTM consolidation (Fig. 4). Taken together, the consolidation of

LTM in *Lymnaea* is suggested to be regulated by LymCBP and LymCREB1. The Taz1 and

Taz2 domains of LymCBP were also highly conserved (Fig. 1). These domains function as

interfaces to bind with a variety of transcription factors and HATs, such as Fos and

p300/CBP-associated factor (pCAF) (Glass et al., 1997). It was previously reported that the

bromodomain of mammalian CBP is important for acetylation of histones (Raisner et al.,

2018) and interaction with acetylated histones (Zeng et al., 2008; Plotnikov et al., 2014). At

least 15 amino acid residues in the bromodomain regulate interaction with acetylated histones

(Dhalluin et al., 1999; Plotnikov et al., 2014; Xu et al., 2017), and, compared with amino acid

sequences among human p300, human CBP and LymCBP, 14 out of 15 amino acid residues

were conserved in LymCBP (Table S1). Other functional amino acid residues are highly

conserved in the RING domain (Table S2), the acetyl-CoA-binding domain (Table S3), and

the ZZ domain (Table S4). The RING domain in human p300/CBP has an autoinhibitory

function for the HAT domain (Delvecchio et al., 2013; Ortega et al., 2019). The ZZ domain of

human p300/CBP recognizes histone tails and modulates its enzymatic activity (Legge et al.,

2004; Zhang et al., 2018). The most important domain in CBP is the KIX domain, which is

well known as an interface for interacting with CREB (Kwok et al., 1994), and this domain is

highly conserved in LymCBP (Fig. 1). These results suggested that not only the HAT activity,

which was shown in Fig. 6, but also other functions of CBP are highly conserved in LymCBP.

Our immunoprecipitation assays indicated an interaction between LymCBP and

LymCREB2 (Fig. 5B). A similar interaction also has been shown in the human homologues of

these proteins (Liang and Hai, 1997). The interfaces in CBP for binding with CREB2 include

the KIX domain, Taz2 domain and HAT domain (Liang and Hai, 1997). These domains were

highly conserved in LymCBP (Fig. 1). In addition, acetylation of CREB2 by CBP was

previously reported (Gachon et al., 2002). Therefore, the interaction between LymCBP and

LymCREB2 shown in Fig. 5B may be needed for LymCREB2 acetylation by LymCBP.

Expression of LymCBP in the Lymnaea CNSs

In situ hybridization showed that LymCBP mRNA was transcribed in a limited

number of identifiable neurons (Fig. 4F). In particular, the CGC in the cerebral ganglia

exhibited a strong LymCBP signal (Fig. 4B,C). Both mRNA and protein forms of LymCREB1

exist in the CGC (Ribeiro et al., 2003; Sadamoto et al., 2004), showing the co-localization of

LymCBP and LymCREB1 in this neuron. The CGC was identified to be a key neuron for the

long-lasting conditioned responses after taste aversion learning (Kojima et al., 2001;

Yamanaka et al., 2000; Ito et al., 2012; Sunada et al., 2017) and appetitive classical

conditioning (Kemenes et al., 2006; Nikitin et al., 2008; Vavoulis et al., 2010; Marra et al.,

2013; Nikitin et al. 2013). The same signal was also seen in identified large neurons, the

RPeD1 and the LPeD1, in the pedal ganglia (Fig. 4D,E). The RPeD1 was especially reported

to be necessary for LTM formation in operant conditioning of aerial respiratory behavior

(Scheibenstock et al., 2002). The staining patterns of LymCBP mRNA in the buccal, cerebral

and pedal ganglia were very similar to those of LymCREB1 (Sadamoto et al., 2004), which

possesses KID for interacting with LymCBP and plays a role in the enhancement of

transcriptional activity (Sadamoto et al., 2004).

Our proteomics experiments confirmed the existence of translated LymCBP in the

Lymnaea CNS, corroborating the findings from the DNA and RNA level assays of LymCBP

expression.

In addition, we previously indicated the existence of both mRNA and protein forms

of another transcriptional factor, CCAAT/enhancer binding protein of *Lymnaea* (LymC/EBP),

in the CGC and the RPeD1 (Hatakeyama et al., 2004a). In mammalian cells, C/EBP is

acetylated by p300/CBP, its acetylation boosts C/EBP-mediated transcriptional activity

(Ceseña et al., 2007). Taken together, the interaction between LymCBP and transcriptional

factors, such as LymCREB1 and LymC/EBP, may be involved in the consolidation of LTM of

Lymnaea.

Involvement of LymCBP HAT activity in synaptic plasticity

In this study, we used HAT inhibitors (curcumin, anacardic acid and garcinol;

reviewed by Hatakeyama et al., 2013b) for biochemical and electrophysiological experiments

(Figs. 6 and 7). Biochemical analyses using the recombinant LymCBP HAT domain showed

that the HAT activity of LymCBP was significantly blocked by HAT inhibitors. Molecular

docking simulation indicated the amino acid residues of the human p300, whose amino acid

sequence of the HAT domain was highly identical with human CBP, for interaction with

garcinol (Coste et al., 2020). For interaction between human p300 and garcinol, five amino

acid residues (S1400, Y1414, D1444, Y1446 and Q1455) in its HAT domain were required,

and these were completely conserved in LymCBP (S1356, Y1370, D1400, Y1402 and Q1411)

(Table S5). Therefore, HAT activity of LymCBP might be inhibited by similar interaction

mechanisms with garcinol.

Electrophysiological experiments showed that these chemicals inhibited
cAMP-dependent synaptic plasticity between the CGCs and their follower neurons (Fig. 7).
Although the block of synaptic plasticity by curcumin has not been reported in other species,
its inhibitory effects on newly acquired or reactivated fear memories were indicated (Monsey
et al., 2015). Similarly, garcinol was also reported to block newly acquired memories in
mammals (Zhao et al., 2012; Maddox et al., 2013a; Monsey et al., 2017) and honeybees
(Merschbaecher et al., 2016). Another HAT inhibitor, C646, which was not used in this study,
significantly impaired memory consolidation (Maddox et al., 2013b; Merschbaecher et al.,
2016). These inhibitory effects might be induced by blocking HAT activities of CBP and other
HATs. On the other hand, there are many reports showing that curcumin has ameliorative
effects from the impairment of memory and synaptic plasticity caused by amyloid peptides
(Ahmed et al., 2011; Zhang et al., 2015), chronic stress (Liu et al., 2014), drugs (Soukhaklari
et al., 2018), heavy metals (Namgyal et al., 2020), aging (Belviranli et al., 2013; Cheng et al.,
2013) and viral proteins (Tang et al., 2009; Shen et al., 2015), suggesting that other molecular
mechanisms, except for inhibiting HAT activity, rescue memory and synaptic plasticity.

Acknowledgements
This work was partly supported by The Japan Society for the Promotion of Science (JSPS), a
Grant-in-Aid for Scientific Research (C) to D.H. (No. 26460562 and 17K08867), Takeda
Science Foundation to D.H, Waseda University Grants for Special Research Projects
(2018K-141 and 2020C-135) to E.I. Work at the University of Sussex, UK, was funded by
grants from the Medical Research Council to G.K. (MRC/G0400551), and the Biotechnology
and Biological Sciences Research Council (BBSRC) to G.K. and I.K. (BB/H009906/1) and to
I.K. and G.K. (BB/P00766X/1), respectively.

Author Contributions

Experiments performed in the UK: D.H., I.F., A.A., M.E. I.K. and G.K. conceived and designed the experiments. I.F. and A.F. performed the Western blot assays. A.A. and M.E. performed the immunoprecipitation and LC-MS experiments, respectively. I.F., A.F., A.A., M.E. and G.K. analyzed and visualized the data. A.A., M.E. I.K. and G.K. wrote the relevant results sections of the paper. G.K. and I.K. obtained the funding for the work performed at the University of Sussex.

Experiments performed in Japan: D.H. conceived and designed the experiments. D.H. and Y.T. purified total RNA from *Lymnaea* tissues. D.H., R.M., and T.K. prepared the LymCBP recombinant protein and performed acetylation assay. H.S. and E.I. performed electrophysiological experiments. T.W. created the phylogenetic tree.

D.H. wrote the first version of the paper and all the authors contributed to the final version.

Competing Interests

The authors declare no competing financial interests.

References

Ahmed T, Gilani AH, Hosseinmardi N, Semnanian S, Enam SA, Fathollahi Y (2011) Curcuminoids rescue long-term potentiation impaired by amyloid peptide in rat hippocampal slices. SYNAPSE 65:572-582.

Alarcón JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER, Barco A (2004) Chromatin acetylation, memory, and LTP are impaired in CBP$^{+/-}$ mice: a model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. Neuron 42:947–959.

Bedford DC, Kasper LH, Fukuyama T, Brindle PK (2010) Target gene context influences the transcriptional requirement for the KAT3 family of CBP and p300 histone acetyltransferases. Epigenetics 5:9–15.

Belviranli M, Okudan N, Esra K, Atalik N, Öz M (2013) Curcumin improves spatial memory and decreases oxidative damage in aged female rats. Biogerontology 14:187–196.

Cedar H, Bergman Y (2009) Linking DNA methylation and histone modification: patterns and paradigms. Nat Rev Genet 10:295–304.

Ceseña TI, Cardinaux JR, Kwok R, Schwartz J (2007) CCAAT/enhancer-binding protein (C/EBP) β is acetylated at multiple lysines: Acetylation of C/EBPβ at lysine 39 modulates its ability to activate transcription. J Biol Chem 282:956–967.

Chatterjee S, Mizar P, Cassel R, Neidl R, Selvi BR, Mohankrishna DV, Vedamurthy BM, Schneider A, Bousiges O, Mathis C, Cassel JC, Eswaramoorthy M, Kundu TK, Boutillier AL (2013) A novel activator of CBP/p300 acetyltransferases promotes neurogenesis and extends memory duration in adult mice. J Neurosci 33:10698–10712.

Chatterjee S, Angelakos CC, Bahl E, Hawk JD, Gaine ME, Poplawski SG, Schneider-Anthony A, Yadav M, Porcari GS, Cassel JC, Giese KP, Michaelson JJ, Lyons LC, Boutillier AL, Abel T. (2020) The CBP KIX domain regulates long-term memory and circadian activity. BMC Biol 18:155.

Cheng YF, Guo L, Xie YS, Liu YS, Zhang J, Wu QW, Li JM (2013) Curcumin rescues

aging-related loss of hippocampal synapse input specificity of long term potentiation in

mice. Neurochem Res 38:98–107.

Chrivia JC, Kwok RP, Lamb N, Hagiwara M, Montminy MR, Goodman RH (1993)

Phosphorylated CREB binds specifically to the nuclear protein CBP. Nature

365:855–859.

Coste C, Gérard N, Dinh CP, Bruguière A, Rouger C, Leong ST, Awang K, Richomme P,

Derbré S, Charreau B (2020) Targeting MHC regulation using polycyclic polyprenylated

acylphloroglucinols isolated from *Garcinia bancana*. Biomolecules 10:1266.

Delvecchio M, Gaucher J, Aguilar-Gurrieri C, Ortega E, Panne D (2013) Structure of the p300

catalytic core and implications for chromatin targeting and HAT regulation. Nat Struct

Mol Biol 20:1040–1046.

Dhalluin C, Carlson JE, Zeng L, He C, Aggarwal AK, Zhou MM (1999) Structure and ligand

of a histone acetyltransferase bromodomain. Nature 399:491–496.

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

Gachon F, Devaux C, Mesnard JM (2002) Activation of HTLV-I transcription in the presence

of Tax is independent of the acetylation of CREB-2 (ATF-4). Virology 299:271–278.

Glass CK, Rose DW, Rosenfeld MG (1997) Nuclear receptor coactivators. Curr Opin Cell

Biol 9:222–232.

Guan Z, Giustetto M, Lomvardas S, Kim JH, Miniaci MC, Schwartz JH, Thanos D, Kandel

ER (2002) Integration of long-term-memory-related synaptic plasticity involves

bidirectional regulation of gene expression and chromatin structure. Cell 111:483–493.

Hatakeyama D, Fujito Y, Sakakibara M, Ito E (2004a) Expression and distribution of

transcription factor CCAAT/enhancer-binding protein in the central nervous system of

Lymnaea stagnalis. Cell Tissue Res 318:631–641.

Hatakeyama D, Sadamoto H, Ito E (2004b) Real-time quantitative RT-PCR method for

estimation of mRNA level of CCAAT/enhancer binding protein in the central nervous system of *Lymnaea stagnalis*. Acta Biol Hung 55:157–161.

Hatakeyama D. Sadamoto H, Watanabe T, Wagatsuma A, Kobayashi S, Fujito Y, Yamashita M, Sakakibara M, Kemenes G, Ito E (2006) Requirement of new protein synthesis of a transcription factor for memory consolidation: Paradoxical changes in mRNA and protein levels of C/EBP. J Mol Biol 356:569–577.

Hatakeyama D, Mita K, Kobayashi S, Sadamoto H, Fujito Y, Hiripi L, Elekes K, Ito E (2010) Glutamate transporters in the central nervous system of a pond snail. J Neurosci Res 88:1374–1386.

Hatakeyama D, Okuta A, Otsuka E, Lukowiak K, Ito E (2013a) Consolidation of long-term memory by insulin in *Lymnaea* is not brought about by changing the number of insulin receptors. Commun Integr Biol 6:e23955.

Hatakeyama D, Tierling S, Kuzuhara T, Müller U (2013b) Epigenetic regulation of gene expression in the nervous system. pp. 151–171. In "Methods in Neuroethological Research", Oka K and Ogawa H, Eds. Springer.

Hatakeyama D, Shoji M, Yamayoshi S, Hirota T, Nagae M, Yanagisawa S, Nakano M, Ohmi N, Noda T, Kawaoka Y, Kuzuhara T (2014) A novel functional site in the PB2 subunit of influenza A virus essential for acetyl-CoA interaction, RNA polymerase activity, and viral replication. J Biol Chem 289:24980–24994.

Hatakeyama D, Shoji M, Yamayoshi S, Yoh R, Ohmi N, Takenaka S, Saitoh A, Arakaki Y, Masuda A, Komatsu T, Nagano R, Nakano N, Noda T, Kawaoka Y, Kuzuhara T (2018a) Influenza A virus nucleoprotein is acetylated by histone acetyltransferases, PCAF and GCN5. J Biol Chem 293:7126–7138.

Hatakeyama D, Ohmi N, Saitoh A, Makiyama K, Morioka M, Okazaki H, Kuzuhara T (2018b) Acetylation of lysine residues in the recombinant nucleoprotein and VP40 matrix

protein of Zaire Ebolavirus by eukaryotic histone acetyltransferases. Biochem Biophys

Res Comm 504:635–640.

Hirano Y, Ihara K, Masuda T, Yamamoto T, Iwata I, Takahashi A, Awata H, Nakamura N,

Takakura M, Suzuki Y, Horiuchi J, Okuno H, Saitoe M (2016) Shifting transcriptional

machinery is required for long-term memory maintenance and modification in

Drosophila mushroom bodies. Nat Commun 7:13471.

Ito E, Otsuka E, Hama N, Aonuma H, Okada R, Hatakeyama D, Fujito Y, Kobayashi S (2012)

Memory trace in feeding neural circuitry underlying conditioned taste aversion in

Lymnaea. PLoS One, 7:e43151.

Kandel ER (2001) The molecular biology of memory storage: A dialogue between genes and

synapses. Science 294:1030–1038.

Kemenes I, Straub VA, Nikitin ES, Staras K, O'Shea M, Kemenes G, Benjamin PR. (2006)

Role of delayed nonsynaptic neuronal plasticity in long-term associative memory. Curr

Biol 16:1269–1279.

Kojima S, Hosono T, Fujito Y, Ito E (2001) Optical detection of neuromodulatory effects of

conditioned taste aversion in the pond snail *Lymnaea stagnalis*. J Neurobiol 49:118–128.

Korzus E, Rosenfeld MG, Mayford M (2004) CBP histone acetyltransferase activity is a

critical component of memory consolidation. Neuron 42:961–972.

Kwok RP, Lundblad JR, Chrivia JC, Richards JP, Bächinger HP, Brennan RG, Roberts SG,

Green MR, Goodman RH (1994) Nuclear protein CBP is a coactivator for the

transcription factor CREB. Nature 370:223–226.

Lakhina V, Arey RN, Kaletsky R, Kauffman A, Stein G, Keyes W, Xu D, Murphy CT (2015)

Genome-wide functional analysis of CREB/long-term memory-dependent transcription

reveals distinct basal and memory gene expression programs. Neuron 85:330–345.

Lassot I, Estrabaud E, Emiliani S, Benkirane M, Benarous R, Margottin-Goguet F. (2005)

p300 modulates ATF4 stability and transcriptional activity independently of its

acetyltransferase domain. J Biol Chem 280:41537–41545.

Laviv T, Scholl B, Parra-Bueno P, Foote B, Zhang C, Yan L, Hayano Y, Chu J, Yasuda R.

(2020) *In vivo* imaging of the coupling between neuronal and CREB activity in the mouse

brain. Neuron 105:799–812.

Legge GB, Martinez-Yamout MA, Hambly DM, Trinh T, Lee BM, Dyson HJ, Wright PE

(2004) ZZ domain of CBP: an unusual zinc finger fold in a protein interaction module. J

Mol Biol 343:1081–1093.

Liang G, Hai T (1997) Characterization of human activating transcription factor 4, a

transcriptional activator that interacts with multiple domains of cAMP-responsive

element-binding protein (CREB)-binding protein. J Biol Chem 272:24088–24095.

Liu X, Wang L, Zhao K, Thompson PR, Hwang Y, Marmorstein R, Cole PA (2008) The

structural basis of protein acetylation by the p300/CBP transcriptional coactivator. Nature

451:846–850.

Liu D, Wang Z, Gao Z, Xie K, Zhang Q, Jiang H, Pang Q (2014) Effects of curcumin on

learning and memory deficits, BDNF, and ERK protein expression in rats exposed to

chronic unpredictable stress. Behav Brain Res 271:116–121.

Maddox SA, Watts CS, Doyère V, Schafe GE (2013a) A naturally-occurring histone

acetyltransferase inhibitor derived from *Garcinia indica* impairs newly acquired and

reactivated fear memories. PLoS One 8:e54463.

Maddox SA, Watts CS, Schafe GE (2013b) p300/CBP histone acetyltransferase activity is

required for newly acquired and reactivated fear memories in the lateral amygdala. Learn

Mem 20:109–119.

Maksimoska J, Segura-Peña D, Cole PA, Marmorstein R (2014) Structure of the p300 histone

acetyltransferase bound to acetyl-coenzyme A and its analogues. Biochemistry

53:3415–3422.

Marra V, O'Shea M, Benjamin PR, Kemenes I (2013) Susceptibility of memory consolidation

during lapses in recall. Nat Commun 4:1578.

Martinez-Balbás MA, Bannister AJ, Martin K, Haus-Seuffert P, Meisterernst M, Kouzarides T (1998) The acetyltransferase activity of CBP stimulates transcription. EMBO J 17:2886-2893.

Matos MR, Visser E, Kramvis I, van der Loo RJ, Gebuis T, Zalm R, Rao-Ruiz P, Mansvelder HD, Smit AB, van den Oever MC. (2019) Memory strength gates the involvement of a CREB-dependent cortical fear engram in remote memory. Nat Commun 10:2315.

Mayr B, Montminy M (2001) Transcriptional regulation by the phosphorylation-dependent factor CREB. Nat Rev Mol Cell Biol 2:599–609.

Merschbaecher K, Hatko L, Folz J, Mueller U (2016) Inhibition of different histone acetyltransferases (HATs) uncovers transcription-dependent and -independent acetylation-mediated mechanisms in memory formation. Learn Mem 23:83–89.

Michel M, Kemenes I, Müller U, Kemenes G (2008) Different phases of long-term memory require distinct temporal patterns of PKA activity after single-trial classical conditioning. Learn Mem 15:694–702.

Monsey MS, Gerhard DM, Boyle LM, Briones MA, Seligsohn M, Schafe GE (2015) A diet enriched with curcumin impairs newly acquired and reactivated fear memories. Neuropsychopharmacology 40:1278–1288.

Monsey MS, Sanchez H, Taylor JR (2017) The naturally occurring compound *Garcinia indica* selectively impairs the reconsolidation of a cocaine-associated memory. Neuropsychopharmacology 42:587–597.

Mujtaba S, He Y, Zeng L, Yan S, Plotnikova O, Sachchidanand, Sanchez R, Zeleznik-Le NJ, Ronai Z, Zhou MM (2004) Structural mechanism of the bromodomain of the coactivator CBP in p53 transcriptional activation. Mol Cell 13:251–263.

Murakami J, Okada R, Sadamoto H, Kobayashi S, Mita K, Sakamoto Y, Yamagishi M, Hatakeyama D, Otsuka E, Okuta A, Sunada H, Takigami S, Sakakibara M, Fujito Y,

Awaji M, Moriyama S, Lukowiak K, Ito E (2013) Involvement of insulin-like peptide in

long-term synaptic plasticity and long-term memory of the pond snail *Lymnaea stagnalis*.

J Neurosci 33:371–383.

Nakai J, Totani Y, Hatakeyama D, Dyakonova VE, Ito E (2020a) Another example of

conditioned taste aversion: Case of snails. Biology (Basel) 9:422.

Nakai J, Totani Y, Kojima S, Sakakibara M, Ito E (2020b) Features of behavioral changes

underlying conditioned taste aversion in the pond snail *Lymnaea stagnalis*. Invert

Neurosci 20:8.

Nakamura H, Kobayashi S, Kojima S, Urano A, Ito E (1999) PKA-dependent regulation of

synaptic enhancement between a buccal motor neuron and its regulatory interneuron in

Lymnaea stagnalis. Zool Sci 16:387–394.

Namgyal D, Ali S, Mehta R, Sarwat M (2020) The neuroprotective effect of curcumin against

Cd-induced neurotoxicity and hippocampal neurogenesis promotion through

CREB-BDNF signaling pathway. Toxicology 442:152542.

Nikitin ES, Vavoulis DV, Kemenes I, Marra V, Piger Z, Michel M, Feng J, O'Shea M,

Benjamin PR, Kemenes G. (2008) Persistent sodium current is a nonsynaptic substrate for

long-term associative memory. Curr Biol 18:1221–1226.

Nikitin ES, Balaban PM, Kemenes G (2013) Nonsynaptic plasticity underlies a

compartmentalized increase in synaptic efficacy after classical conditioning. Curr Biol

23:614–619.

Ortega E, Rengachari S, Ibrahim Z, Hoghoughi N, Gaucher J, Holehouse AS, Khochbin S,

Panne D (2018) Transcription factor dimerization activates the p300 acetyltransferase.

Nature 562:538–544.

Plotnikov AN, Yang S, Zhou TJ, Rusinova E, Frasca A, Zhou MM (2014) Structural insights

into acetylated-histone H4 recognition by the bromodomain-PHD finger module of

human transcriptional coactivator CBP. Structure 22:353–360.

Radhakrishnan I, Perez-Alvarado GC, Parker D, Dyson HJ, Montminy MR, Wright PE (1997)

Solution structure of the KIX domain of CBP bound to the transactivation domain of

CREB: a model for activator:coactivator interactions. Cell 91:741–752.

Raisner R, Kharbanda S, Jin L, Jeng E, Chan E, Merchant M, Haverty PM, Bainer R, Cheung

T, Arnott D, Flynn EM, Romero FA, Magnuson S, Gascoigne KE (2018) Enhancer

activity requires CBP/P300 bromodomain-dependent histone H3K27 acetylation. Cell

Rep 24:1722–1729.

Rappsilber J, Ishihama Y, Mann M (2003) Stop and go extraction tips for matrix-assisted laser

desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics.

Anal Chem 75:663–670.

Ribeiro MJ, Serfőző Z, Papp A, Kemenes I, O'Shea M, Yin JC, Benjamin PR, Kemenes G

(2003) Cyclic AMP response element-binding (CREB)-like proteins in a molluscan brain:

cellular localization and learning-induced phosphorylation. Eur J Neurosci

18:1223–1234.

Sadamoto H, Sato H, Kobayashi S, Murakami J, Aonuma H, Ando H, Fujito Y, Hamano K,

Awaji M, Lukowiak K, Urano A, Ito E (2004) CREB in the pond snail *Lymnaea*

stagnalis: cloning, gene expression, and function in identifiable neurons of the central

nervous system. J Neurobiol 58:455–466.

Sadamoto H, Kitahashi T, Fujito Y, Ito E (2010) Learning-dependent gene expression of

CREB1 isoforms in the molluscan brain. Front Behav Neurosci 4:25.

Scheibenstock A, Krygier D, Haque Z, Syed N, Lukowiak K (2002) The soma of RPeD1 must

be present for long-term memory formation of associative learning in *Lymnaea*. J

Neurophysiol 88:1584–1591.

Senthil R, Angel KJ, Malathi R, Venkatesan D (2011) Isolation, identification and

computational studies on *Pseudomonas aeruginosa* sp. strain MPC1 in tannery effluent.

Bioinformation 6:187–190.

Shaywitz AJ, Dove SL, Kornhauser JM, Hochschild A, Greenberg ME (2000) Magnitude of

the CREB-dependent transcriptional response is determined by the strength of the

interaction between the kinase-inducible domain of CREB and the KIX domain of

CREB-binding protein. Mol Cell Biol 20:9409–9422.

Shen LL, Jiang ML, Liu SS, Cai MC, Hong ZQ, Lin LQ, Xing YY, Chen GL, Pan R, Yang LJ,

Xu Y, Dong J (2015) Curcumin improves synaptic plasticity impairment induced by

HIV-1gp120 V3 loop. Neural Regen Res 10:925–931.

Soukhaklari R, Moezi L, Pirsalami F, Ashjazadeh N, Moosavi M (2018) Curcumin

ameliorates scopolamine-induced mice memory retrieval deficit and restores

hippocampal p-Akt and p-GSX-3β. Eur J Pharmacol 841:28-32.

Sunada H, Lukowiak K, Ito E (2017) Cerebral giant cells are necessary for the formation and

recall of memory of conditioned taste aversion in *Lymnaea*. Zoolog Sci 34:72–80.

Tang H, Lu D, Pan R, Qin X, Xiong H, Dong J (2009) Curcumin improves spatial memory

impairment induced by human immunodeficirncy virus type i glycoprotein V3 loop

peptide in rats. Life Sci 85:1–10.

Totani Y, Nakai J, Dyakonova VE, Lukowiak K, Sakakibara M, Ito E (2020) Induction of

LTM following an insulin injection. eNeuro 7:ENEURO.0088-20.2020.

Vavoulis DV, Nikitin ES, Kemenes I, Marra V, Feng J, Benjamin PR, Kemenes G. (2010)

Balanced plasticity and stability of the electrical properties of a molluscan modulatory

interneuron after classical conditioning: a computational study. Front Behav Neurosci

4:19.

Vecsey CG, Hawk JD, Lattal KM, Stein JM, Fabian SA, Attner MA, Cabrera SM,

McDonough CB, Brindle PK, Abel T, Wood MA (2007) Histone deacetylase inhibitors

enhance memory and synaptic plasticity via CREB:CBP-dependent transcriptional

activation. J Neurosci 27:6128–6140.

Xu L, Cheng A, Huang M, Zhang J, Jiang Y, Wang C, Li F, Bao H, Gao J, Wang N, Liu J, Wu

J, Wong CCL, Ruan K (2017) Structural insight into the recognition of acetylated histone

H3K56ac mediated by the bromodomain of CREB-binding protein. FEBS J

284:3422–3436.

Wagatsuma A, Sadamoto H, Kitahashi T, Lukowiak K, Urano A, Ito E (2005) Determination

of the exact copy numbers of particular mRNAs in a single cell by quantitative real-time

RT-PCR. J Exp Biol 208:2389–2398.

Wagatsuma A, Azami S, Sakura M, Hatakeyama D, Aonuma H, Ito E (2006) De Novo

synthesis of CREB in a presynaptic neuron is required for synaptic enhancement involved

in memory consolidation. J Neurosci Res 84:954–960.

Wessel D, Flugge UI (1984) A method for the quantitative recovery of protein in dilute

solution in the presence of detergents and lipids. Anal Biochem 138:1417–143.

Yamanaka M, Hatakeyama D, Sadamoto H, Kimura T, Ito E. (2000) Development of key

neurons for learning stimulates learning ability in *Lymnaea stagnalis*. Neurosci Lett

278:113–116.

Zeng L, Zhang Q, Gerona-Navarro G, Moshkina N, Zhou MM (2008) Structural basis of

site-specific histone recognition by the bromodomains of human coactivators PCAF and

CBP/p300. Structure 16:643–652.

Zhang L, Fang Y, Xu Y, Lian Y, Xie N, Wu T, Zhang H, Sun L, Zhang R, Wang Z (2015)

Curcumin improves amyloid β-peptide (1-42) induced spatial memory deficits through

BDNF-ERK signaling pathway. PLoS One 10:e0131525.

Zhang Y, Xue Y, Shi J, Ahn JW, Mi W, Ali M, Wang X, Klein BJ, Wen H, Li W, Shi X,

Kutateladze TG (2018) The ZZ domain of p300 mediates specificity of the adjacent HAT

domain for histone H3. Nat Struct Mol Biol 25:841–849.

Zhao Z, Fan L, Fortress AM, Boulware MI, Frick KM (2012) Hippocampal histone

acetylation regulates object recognition and the estradiol-induced enhancement of object

recognition. J Neurosci 32:2344–2351.

Zhou L, Zhang Y, Liu RY, Smolen P, Cleary LJ, Byrne JH (2015) Rescue of impaired

long-term facilitation at sensorimotor synapses of *Aplysia* following siRNA knockdown

of CREB1. J Neurosci 35:1617–1626.

 Figure Legends

 Fig. 1. Predicted amino acid sequence of *Lymnaea* CBP. (A) Diagram of the structure of

 Lymnaea CBP (GenBank accession number: AB217914). Main functional domains (Taz1

 domain, KIX domain, bromodomain, acetylation domain and Taz2 domain) are highlighted

 with patterned bars. Alignments of amino acid sequences of (B) Taz1 domain, (C) KIX

 domain and (D) compound of the bromodomain, HAT domain and Taz2 domain between

 Lymnaea CBP and *Aplysia* CBP (GenBank accession number: AY064470) and human CBP

 (GenBank accession number: U47741). The patterns of bars in (D) correspond to those shown

 in (A).

 Fig. 2. Molecular phylogenetic tree of *Lymnaea* CBP and other p300/CBP proteins. The scale

 bar indicates 0.1 substitutions per site. Bootstrap values are shown at the nodes. A CBP

 homologue of the Choanoflagellate *Salpingoeca rosetta* was used as an outgroup. The

 GenBank accession numbers of p300/CBP-like proteins are listed in Table 1.

 Fig. 3. Quantification of LymCBP mRNA in the whole CNS of *Lymnaea*. LymCBP is more

 abundantly transcribed in the CNS than other tissues of *Lymnaea*. The data are expressed as

 the mean ± SEM; $p < 0.01$, **$p < 0.01$, Tukey's post-hoc test.

 Fig. 4. Localization of LymCBP mRNA in identifiable neurons shown by *in situ* hybridization.

 (A) Weak signal of LymCBP could be observed in several neurons (arrowheads), especially in

 the B4CL feeding motoneurons (arrows), in the buccal ganglia. Bu. G.: buccal ganglion, L:

 left; R: right. (B-E) Strong signal of LymCBP mRNA was observed in both CGCs

 (arrowheads in B and C), RPeD1 (arrowhead in D) and LPeD1 (arrowhead in E). Ce. G.:

 cerebral ganglion, Pe. G.: pedal ganglion. (F) Schematic drawing of the distribution of

neurons containing LymCBP mRNA. Pa. G.: parietal ganglion, Vi. G.: visceral ganglion. Scale

bars = 100 μm (A, D and E) and 20 μm (B and C).

Fig. 5. Identification of the protein form of LymCBP. (A) Existence of LymCBP protein in

CNSs was confirmed by western blotting. Molecular weight of the positive band (245 kDa;

filled arrowhead) was consistent with theoretical value calculated from amino acid sequence

of LymCBP (252.3 kDa). A putative smaller isoform of LymCBP was also detected (open

arrowhead). (B, C) CREB-Binding Protein identified as an interacting partner of CREB2 by

MS/MS analysis in the CNS of *Lymnaea stagnalis*. (B) Sixty extracted CNS of *Lymnaea*

stagnalis were pooled, homogenised and subjected to immunoprecipitation using a specific

CREB2 antibody. Immune complexes were immunoblotted for CREB2. (C) MS/MS

Spectrum of the CBP identified as one of the interacting partners of CREB2 in the CNS of

Lymnaea stagnalis. CREB2 IP product was subjected for MS/MS analysis and a CBP specific

peptide for *Lymnaea* CBP (KVFNPDELR) was identified.

Fig. 6. HAT activity of LymCBP was blocked by inhibitors. (A) The partial recombinant

protein of LymCBP HAT domain purified by cation-exchange chromatography was detected

in fractions No. 12–19 (bands shown with arrowhead). The concentrations of these

preparations were too low to be measured by Bradford protein assay. (B) All fractions

obtained by cation-exchange chromatography were mixed (lane 1), and the recombinant

protein was condensed using a centrifugal filter. A high enough concentration of the

recombinant protein for biochemical experiments was obtained (lane 2). (C) Condensed

recombinant LymCBP HAT domain was incubated with histone H1 at different temperatures;

15^{o}C, 20^{o}C, 25^{o}C, 30^{o}C and 37^{o}C. CBB staining of the gel showed that same amount of

histone H1 was loaded (open arrowhead, lanes 3-7). However, the band of recombinant

LymCBP HAT domain (filled arrowhead) was drastically decreased after incubation at 37^{o}C,

suggesting that LymCBP was denatured (lane 7). A positive signal of acetylation was observed after incubation at 15°C, 20°C, 25°C and 30°C (lanes 8-11). This signal was almost eliminated after incubation at 37°C (lane 12). (D) Effects of the HAT inhibitors curcumin, anacardic acid and garcinol. Reaction mixtures were incubated at 30°C. Although the same amount of recombinant LymCBP HAT domain (filled arrowhead) and histone H1 (open arrowhead) was incubated, CBB staining showed that the band signals of histone H1 were attenuated after incubating with curcumin (lane 14) and garcinol (lane 16), suggesting that these inhibitors degraded the protein. Autoradiography indicated that, without any HAT inhibitor, positive signal of acetylation was detected on the band of histone H1 (lane 17). This signal was eliminated by addition of each HAT inhibitor (lanes 18-20), showing that the acetylation activity of LymCBP recombinant protein was significantly blocked by HAT inhibitors.

Fig. 7. Long-term synaptic changes at synapses between the CGC and the B1 motoneuron in the isolated CNS are blocked by HAT inhibitors. EPSPs were evoked in the B1 motoneuron by the injection of depolarizing current into the CGC. The isolated CNS was incubated with DMSO saline or HAT inhibitors for 1 h prior to injection of cAMP. (A) cAMP injection into the CGC cell body increased EPSP amplitude in B1 motoneuron at both 30 min and 3 h after the injection. (B) In contrast, EPSP amplitude increase was not observed in the group which was injected with 5'AMP. (C, D, E) The CNS was incubated with saline containing one of the HAT inhibitors curcumin (67.9 μM), or anacardic acid (50 μM), or garcinol (20 μM) for 1 h prior to injection of the cAMP. In these groups, the EPSP was not increased either at 30 min or 3h after the injection of cAMP. (F) The enhancement of B1 EPSPs is shown as the summarized data. Injection of cAMP into the CGC cell body increased EPSP amplitude in the B1 motoneuron, but pre-treatment with either HAT inhibitor did not elicit enlargement of EPSP. The data are expressed as the mean ± SEM, $*p < 0.05$, Tukey's post-hoc test.

947 **Table 1.** Nucleotide sequences of primers used for cloning (No. 1-14), synthesis for *in situ*

948 hybridization probe (No. 15-16), synthesis of recombinant protein (No. 17-18), and qRT-PCR

949 (No. 19-24) of LymCBP. Amino acid sequences of ApCBP for designing degenerate PCR

950 (DGPCR) was shown with nucleotide sequences.

| 952 **Names of primers** | **Nucleotide sequences** |
953 **and probes**	
955 1. DGPCR_Taz1_fw	5'-CTS ATC CAG CAG CAG CTB GTB CTS C-3' (LIQQQLVL)
956 2. DGPCR_Taz1_rv	5'-ATT TGW CGA GAW GAR GCA CAR TGR G-3' (AHCASSRQII)
957 3. DGPCR_KIX_fw	5'-CTG GTS CAR GCY ATY TTY CC-3' (KLVQAIFP)
958 4. DGPCR_KIX_rv	5'-GTA SAT YTT CTC RGC SAG CAA RTG-3' (HLLAEKIY)
959 5. DGPCR_Taz2_fw	5'-TCC ATT CAR CGC TGY ATC TCN TCH C-3' (SIQRCISSL)
960 6. DGPCR_Taz2_rv	5'-ACA GAA TGG GAC CKG RCA YTT RTT YTC-3' (ENKCQVPFC)
961 7. 5'RACE_outer	5'-TTA CTG GCT GGT ATA GAT TTA ACA ATA TCT GGA GC-3'
962 8. 5'RACE_inner	5'-TAT GTT ACC CAA AAG ATT ACT CAT ATG ACT GCT AGG-3'
963 9. 3'RACE_outer	5'-CTA ACT GTC GTA TCA ACT CGT GCA TCA AG-3'
964 10. 3'RACE_inner	5'-GCA TCA AGA TGA AAC GTG TTG TTT CAC-3'
965 11. Taz1-KIX_fw	5'-ACA TGC AAC AAA GGA AAA TCT TG-3'
966 12. Taz1-KIX_rv	5'-TGA TAA TAT TCT CCT CTA TTG TTA GCA G-3'
967 13. KIX-Taz2_fw	5'-ACA TGC AAC AAA GGA AAA TCT TGT GAA GTG GCC C-3'
968 14. KIX-Taz2_rv	5'-CGT ACA ATG TTT GGC ATG GTA ACA ACA GAG TGC-3'
969 15. qPCR_cloning_fw	5'-GAT GTA AAC ATG GCC GAC CAC CAA GTT G-3'
970 16. qPCR_cloning_rv	5'-AGA CGA TGT AGC AAC TTT GTT TGT ATT GGC-3'
971 17. qPCR_RT	5'-TGT TAA GAG ACA TCG GCA TTG ATG-3'
972 18. qPCR_fw	5'-GCC CTC CGG CCA ACA AGA A-3'
973 19. qPCR_rv	5'-TAT TAT CGC TGG GTG TAT TGA GAG AT-3'

974	20.	qPCR_probe	5'-Fam-ACC CAG AAT CGG ATC AC-Tamra-3'

975   21.   in-situ-probe_fw   5'-GAT ATC TCT TGA AAT CTT AGT GCT GTG TTG ATG-3' (*Eco*R V)

976   22.   in-situ-probe_rv   5'-GGA TCC ATC AGC AGA AGT CTG TCG AGT ACC-3' (*Bam*H I)

977   23.   recombinant_fw   5'-CGC GGC AGC CAT ATG **GCT AGC** AGA CGA AAA GAA AAC

978   AAA TTT ACA GCC-3' [nucleotide sequence of pET28a(+) plasmid,

979   **the restriction site of *Nhe* I**, nucleotide sequence of LymCBP]

980   24.   recombinant_rv   5'-ACG GAG CTC GAA TTC **GGA TCC** TTA ATC AGG TTC TGT GAT

981   TGG TGG CAG-3' [nucleotide sequence of pET28a(+) plasmid,

982   **the restriction site of *Bam*H I**, nucleotide sequence of LymCBP]

984 **Table 2.** GenBank accession numbers of various CBP and p300 used for molecular

985 phylogenetic tree shown in Fig. 2.

	Species	Protein	GenBank Accession No.
987	**Species**	**Protein**	**GenBank Accession No.**
988	*Alligator mississippiensis*	CBP	XP_006264860
989	*Alligator mississippiensis*	p300	XP_014463094
990	*Apis mellifera*	CBP	XP_006568897
991	*Aplysia californica*	CBP	NP_001191640.1
992	*Blattella germanica*	CBP	CUT08824
993	*Brugia malayi*	CBP	CTP80907
994	*Caenorhabditis elegans*	CBP	NP_499161
995	*Capitella teleta*	CBP	ELU13629
996	*Ciona intestinalis*	CBP	XP_009860762.1
997	*Crassostrea gigas*	CBP	XP_011427368.1
998	*Danio rerio*	CBP	XP_009297440.1
999	*Drosophila melanogaster*	nejire	NP_001188575.1
1000	*Gallus gallus*	CBP	XP_015150108.1
1001	*Gallus gallus*	p300	XP_004937767
1002	*Homo sapiens*	CBP	NP_004371.2
1003	*Homo sapiens*	p300	NP_001420
1004	*Hydra vulgaris*	CBP	XP_002156492.2
1005	*Hymenolepis microstoma*	CBP	CDS30104
1006	*Limulus polyphemus*	CBP	XP_013776092
1007	*Loa loa*	CBP	EJD74639
1008	*Lymnaea stagnalis*	CBP	AB217914
1009	*Mus musculus*	CBP	NP_001020603.1

1010	*Mus musculus*	p300	NP_808489
1011	*Opisthorchis viverrini*	CBP	XP_009171514
1012	*Parasteatoda tepidariorum*	CBP	XP_015922458
1013	*Priapulus caudatus*	CBP	XP_014664166
1014	*Rattus norvegicus*	CBP	NP_596872.3
1015	*Rattus norvegicus*	p300	XP_017450856
1016	*Salpingoeca rosetta*	CBP	XP_004998720
1017	*Schistosoma mansoni*	CBP	CCD78413
1018	*Strongylocentrotus purpuratus*	CBP	XP_011677827.1
1019	*Takifugu rubripes*	CBP	XP_011602271.1
1020	*Xenopus laevis*	CBP1	NP_001088637.1
1021	*Xenopus laevis*	CBP2	NP_001192159.1

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7