Histone acetyltransferase activity of CREB-binding protein is essential for synaptic plasticity in *Lymnaea*

Dai Hatakeyama[1,2,*], Hiroshi Sunada[3], Yuki Totani[4], Takayuki Watanabe[5,6], Ildikó Felletár[1], Adam Fitchett[1], Murat Eravci[1], Aikaterini Anagnostopoulou[1], Ryosuke Miki[2], Takashi Kuzuhara[2], Ildikó Kemenes[1], Etsuro Ito[3,4], György Kemenes[1,*]

[1]Sussex Neuroscience, School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK. [2]Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Tokushima 770-8514, Japan. [3]Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University, Sanuki 769-2193, Japan. [4]Department of Biology, Waseda University, Tokyo 162-8480, Japan. [5]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan. [6]Laboratory of Neuroethology, Sokendai-Hayama, Hayama 240-0193, Japan.

Keywords: Long-term memory; *Lymnaea*; CREB-binding protein (CBP); histone acetyl transferase (HAT); Cerebral Giant Cell (CGC); synaptic plasticity

*Correspondence should be addressed to Dr. Dai Hatakeyama, Tokushima Bunri

University, 180 Nishihama-Houji, Yamashiro-cho, Tokushima City, Tokushima

770-8514, Japan, daihatake926@ph.bunri-u.ac.jp; and Prof. György Kemenes,

University of Sussex, Brighton, BN1 9QG, United Kingdom,

G.Kemenes@sussex.ac.uk.

Abstract

In eukaryotes, CREB-binding protein (CBP), a coactivator of CREB, functions both as a

platform for recruiting other components of the transcriptional machinery and as a

histone acetyltransferase (HAT) that alters chromatin structure. We previously showed

that the transcriptional activity of cAMP-responsive element binding protein (CREB)

plays a crucial role in neuronal plasticity in the pond snail *Lymnaea stagnalis*. However,

there is no information on the role how CBP plays in CREB-initiated plastic changes in

Lymnaea. In this study, we characterized the *Lymnaea* CBP (LymCBP) gene and

investigated the roles it plays in synaptic plasticity involved in regulating feeding

behaviors. Similar to CBPs of other species, LymCBP possesses functional domains,

such as KIX domain, which is essential for interaction with CREB and was shown to

regulate long-term memory (LTM). *In situ* hybridization showed that the staining

patterns of LymCBP mRNA in the central nervous system were very similar to those of

Lymnaea CREB1 (LymCREB1). A particularly strong LymCBP mRNA signal was

observed in the Cerebral Giant Cell (CGC), an identified extrinsic modulatory

interneuron of the feeding circuit, key to both appetitive and aversive LTM for taste.

Biochemical experiments using the recombinant protein of LymCBP HAT domain

showed that its enzymatic activity was blocked by classical HAT inhibitors; i.e.

curcumin, anacardic acid and garcinol. Preincubation of *Lymnaea* CNSs with these HAT

inhibitors blocked cAMP-induced long-term potentiation between the CGC and the

follower B1 motoneuron. We therefore suggest that HAT activity of LymCBP in the

CGCs is a key factor in synaptic plasticity contributing to LTM after classical

conditioning.

Introduction

De novo gene transcription is required for consolidation of long-term memory

(LTM) (Kandel, 2001) and cAMP-responsive element binding protein

(CREB)-dependent gene expression is one of the crucial steps of this process in

mammals (Matos et al., 2019; Laviv et al., 2020) as well as invertebrates (Lakhina et al.,

2015; Zhou et al., 2015; Hirano et al., 2016). Transcriptional activation requires the

recruitment of multifunctional coactivators, resulting in a variety of different types of

epigenetic modifications of histones and genomic DNA (Cedar and Bergman, 2009). In

CREB-dependent control of gene expression, CREB-binding protein (CBP) functions as

a coactivator of CREB. To date, CBP has been reported to be critical for LTM

consolidation in mammals (Alarcón et al., 2004; Korzus et al., 2004; Chatterjee et al.,

2013) and the gastropod *Aplysia* (Guan et al., 2002; Zhou et al., 2015). One of the most

important functions of CBP is due to its histone acetyltransferase (HAT) activity, which

stimulates gene transcription (Martinez-Balbás et al., 1998). Several studies have shown

that histone acetylation by CBP is necessary for hippocampal long-term potentiation

(LTP) (Korzus et al., 2004; Vecsey et al., 2007). Pharmacological inhibition of CBP

HAT activity using the inhibitors curcumin and garcinol was reported to block memory

consolidation (Zhao et al., 2012; Monsey et al., 2015; Merschbaecher et al., 2016) and

memory-associated neuronal plasticity (Maddox et al., 2013a). These studies

demonstrated that HAT activity of CBP plays an essential role in regulating synaptic

plasticity associated with memory consolidation.

The pond snail *Lymnaea stagnalis* is a widely used organism to understand

evolutionarily conserved molecular mechanisms of the consolidation of LTM for

taste-related associations (Kemenes et al., 2006; Hatakeyama et al., 2013a; Murakami et

al., 2013; Totani et al., 2020; Nakai et al., 2020a; 2020b). Sadamoto et al. (2004) first

succeeded in the cloning of an isoform of CREB from central nervous systems (CNSs)

of *Lymnaea* and defined it as LymCREB1. They identified 7 different isoforms of

LymCREB1 by alternative splicing and found that aversive taste conditioning

significantly increased LymCREB1 gene expression (Sadamoto et al., 2010). Ribeiro et

al. (2003) showed that appetitive taste conditioning selectively increased

phosphorylated LymCREB1 in the 'learning ganglia' (the cerebral and buccal ganglia)

of the *Lymnaea* CNS.

An identified extrinsic modulatory neuron type of the feeding system, the

cerebral giant cell (CGC) was reported to play crucial roles in LTM after both aversive

and appetitive taste conditioning (Kemenes et al., 2006; Ito et al., 2012; Nikitin et al.,

2013). Injection of LymCREB1 siRNA into CGCs reduced the amplitude of excitatory

postsynaptic potential (EPSP) in monosynaptic follower neurons of the CGCs, the B1

motoneurons (Wagatsuma et al., 2006), suggesting that regulation of gene expression by

LymCREB1 was required for synaptic enhancement in memory consolidation. Based on

these previous findings, we hypothesized that HAT activity of the *Lymnaea* CBP

(LymCBP) plays an important role in the LymCREB1 initiated molecular processes of

synaptic consolidation.

In the present study, we first cloned the LymCBP cDNA from the *Lymnaea*

CNSs and identified neurons expressing the LymCBP mRNA. Focusing on the CGC,

we pharmacologically and electrophysiologically analyzed the relationship between the

HAT activity of LymCBP and the synaptic plasticity involved in the aversively

classically conditioned feeding behavior of *Lymnaea*. Our findings provide new insights

into the functions of LymCBP in synaptic plasticity underlying the consolidation of

associative LTM.

Materials and Methods

Experimental animals. Pond snails, *Lymnaea stagnalis,* were bred at the University of

Sussex (Brighton, UK) and Waseda University (Tokyo, Japan) were maintained in large

holding tanks containing copper-free water at 18-22°C on a 12-hr light - dark cycle. The

animals were fed on a vegetable-based fish food (Tetra-Phyll; TETRA Werke, Melle,

Germany), lettuce and turnip leaves.

Molecular cloning of LymCBP. To clone LymCBP, a series of degenerate PCR were

performed with TaKaRa Ex Taq® (Takara Clontech) and primers, which were designed

at the basis of highly conserved domains, such as Taz1 (transcriptional adapter

zinc-binding 1), KIX domain and Taz2 domain, of *Aplysia* CBP (ApCBP; GenBank

accession number: AY064470). After the sequential analyses of the Taz1, KIX and Taz2

domains of LymCBP, we performed 5' and 3'RACE, and the PCR for the internal region

between Taz1 and KIX domains and between KIX and Taz2 domains. The FirstChoice®

RLM-RACE Kit (ThermoFisher) was used for amplification of 5' and 3' ends of

LymCBP. All PCR products were subcloned with TA Cloning® Kit (ThermoFisher) or

pGEM®-T Easy Vector System (Promega). Nucleotide sequences of primers were

summarized in Table 1 (No. 1-14).

Phylogenetic tree of LymCBP. To construct a phylogenetic tree of the CBP/p300

proteins, we aligned the full-length deduced amino acid sequence of the *Lymnaea* CBP

with those of the known homologs of other species (listed with each Accession Number

in Table 2) by using the MUSCLE algorisms (Edgar, 2004) on the Geneious (v9.1)

program (available from http://www.geneious.com/). Maximum likelihood tree was

constructed from the aligned sequences using the MEGA 6 program (Tamura et al.,

2013) with default settings of the program. 1000 bootstrap replications were conducted

to evaluate the reliabilities of the reconstructed trees. The obtained tree was visualized

with the FigTree (v1.4.2) program (available from

http://tree.bio.ed.ac.uk/software/figtree/). A CBP homolog of the Choanoflagellate

Salpingoeca rosetta was used as an outgroup.

TaqMan-based quantitative PCR (qPCR). To quantify the absolute copy number of

LymCBP mRNA in several *Lymnaea* tissues, we performed TaqMan-based qPCR by

establishing RNA standard curves derived from a dilution series of known template

concentrations. The procedures for this experiment were modified from previous reports

(Hatakeyama et al., 2006; Wagatsuma et al., 2005). Partial LymCBP gene (from 303rd

to 722nd nucleotides, which contains the open reading flame of the *N*-terminal region)

was amplified by RT-PCR with primers shown in Table 1 (No. 15, 16), and the PCR

product was subcloned into pGEM®-T Easy Vector (Promega). The plasmid was

purified with Plasmid Midi Kit (QIAGEN) and digested with restriction enzyme *Spe* I.

The LymCBP RNA was synthesized with MAXIscript[TM] (ThermoFisher), purified with

RNeasy® Mini Kit (QIAGEN), quantified the concentration, and used as the standard

RNA. Serially diluted standard RNA ($5 \times 10^1 - 5 \times 10^6$ copies/µl) were

reverse-transcribed in 10 µl of reaction mixture to prepare the first strand cDNA with

2.5 µM of the specific primer for LymCBP diluted to 1:5 and used as the standard

cDNAs.

Total RNA from different types of *Lymnaea* tissue (CNS, buccal mass, penis,

ovotestis, gut and mantle) was purified with TRIzol[TM] Reagent (ThermoFisher).

Reverse transcription (RT) of standard series ($5 \times 10^1 - 5 \times 10^6$ copies) and total RNA of

Lymnaea tissues (each 300 ng) were performed in the buffer containing 0.2 µg yeast

tRNA, $1\times$ PCR Buffer II (ThermoFisher), 5.5 mM MgCl$_2$, 0.5 mM dNTP, 0.2 µM gene

specific primer for LymCBP, 0.3 unit/ml Prime RNase inhibitor (Eppendorf), and 0.9

units/ml MultiScribe[TM] Reverse Transcriptase (ThermoFisher) for 10 min at 25°C, for

30 min at 48°C, and for 5 min at 95°C. All RT samples were diluted to 5 times by

adding RNase-free distilled water. The diluted RT solution of samples and standards (1

µl) were added to PCR-reaction solution [final concentration: $1\times$ TaqMan® Gene

Expression Master Mix (ThermoFisher), 50 nM each primer, and 50 nM probe].

Nucleotide sequences of primers and probes were summarized in Table 1 (No. 17-20).

Reaction was carried out at 95°C for 10 min, 45 cycles of 95°C for 15 sec, and 60°C for

1 min with real-time PCR system (StepOnePlus Real-Time PCR System, ThermoFisher).

In the assay, several doses of standard cDNA (1×10^1 – 1×10^6 copies) were applied in

triplicate to estimate inter-assay coefficients of variation between runs.

In situ *hybridization of LymCBP.* The procedures for this experiment were modified

from previous reports (Hatakeyama et al., 2004a; 2006; 2010). Non-functional region

(from 278th to 1146th of nucleotide sequence) of LymCBP was isolated by RT-PCR

(see Table 1 (No. 21, 22) for nucleotide sequences of primers). After insertion of this

PCR product into pGEM®-T Easy Vector, the plasmid was linearized with *Eco*R V and

*Bam*H I to generate digoxigenin-labeled antisense and sense RNA probes, respectively.

The procedures for generating the probes, hybridization and coloring were same as

those that were previously reported (Hatakeyama et al., 2004; 2006).

Western blotting. Lysates of *Lymnaea* CNS dissected from four-month-old snails were

prepared in RIPA buffer. Protein (10-20 µg) was electrophoresed on 6% Bis-Tris protein

gels and transferred to polyvinylidene difluoride (PVDF) membrane. The polyclonal

antibody to anti-human-CBP (C20, sc-583, Santa Cruz) was used at 1:500 dilution in

4% skimmed milk in TBS-Tween overnight at 4°C with rotation. Following washes

with TBS-Tween (10 min ×3), the membranes were incubated with the anti-rabbit-IgG

conjugated with horseradish peroxidase (Sigma Aldrich) at room temperature at

1:10,000 dilution for 2 hr. Finally, the membranes were cleaned from the excess of

secondary antibody by washing with TBS-Tween (10 min ×3) and TBS (10 min ×1).

Detection of blot was carried out with a LumiSensor™ HRP Substrate Kit (GenScript

Technology).

Immunoprecipitation assay and immunoblotting. Sixty pooled *Lymnae*a CNSs were

homogenised in RIPA lysis buffer supplemented with 2 mM PMSF, 1 mmol/L Na_3VO_4

and protease inhibitors, (Santa Cruz, sc-24948) by using a micropestelle. Fifty μl of the

pre-immunoprecipitated lysate was saved as "input" to be used for immunoblotting. The

homogenate was pre-cleared in a Pierce Spin column (Thermofisher Scientific, 69705)

containing 40 μl of Protein A/G PLUS-Agarose (Santa Cruz, sc-2003) for 1 hr at 4°C on

a rotor. Four Pierce Spin columns containing 40 μl of Protein A/G PLUS-Agarose

suspension were prepared. Three of the Pierce Spin columns were used to prepare

antibody/agarose complex and one of the columns was used as a negative control

without containing the antibody. HEPES buffer (500 μl) was added to all of the Protein

A/G PLUS-Agarose spin columns. For the Antibody/Agarose complex preparation, 10

μg of anti-mouse CREB2 antibody (Santa Cruz, sc-390063) was added to each of the

Protein A/G PLUS-Agarose Spin column. The columns were incubated on a rotor at RT

for 3 hrs. The columns were centrifuged at 2500 RPM for 1 min and the flow-through

was removed. The columns were then washed/centrifuged four times at 2500 RPM with

HEPES buffer at 4^{o}C . After the final wash, 500 μg of CNS homogenised lysate was

transferred into each column containing the antibody/agarose complex and the one

column without containing the antibody. The columns were placed on a rotor and

incubated overnight at 4^{o}C. The following day, the columns were centrifuged at 2500

RPM for 1 min and the lysate was removed. The columns were washed/centrifuged four

times at 2500 RPM with HEPES buffer. The eluates for each column was eluted in 50 μl

of glycine, pH 2.8. The pH of the eluates was neutralized with 5 μl of 1 M Tris, pH 9.5.

The immuneprecipitated (IP) products were pooled together and 100 μl of the IP

product was used for LC/MS analysis and 45 μl of the IP product was prepared for

western blotting. The IP product (45 μl) was resuspended in 15 μl of 4× loading sample

buffer and boiled at 90^{o}C for 10 mins. Immunoprecipitates, negative control and 25 μg

of the pre-immunoprecipitated lysate (5% input) were resolved in 10%

SDS-polyacrylamide gel and transferred to a nitrocellulose membrane. Non-specific

binding sites were blocked with 5% skimmed milk in TBS solution with 0.1%

Tween-20 (TBST) for 1 hr at RT. Membranes were incubated with anti-mouse CREB2

antibody (1:1000, Santa Cruz, sc-390063) overnight at 4°C. The following day, the

membrane was washed three times for 10 mins in TBST and incubated with highly

cross-adsorbed Infra-Red Dye 680RD goat anti-mouse secondary antibody (1:5000,

LI-COR Inc., 926-68070) for 1 hr at RT, followed by two washes with TBST and a final

wash of 1× PBS. Membranes were developed using the LI-COR Odyssey Fc

Fluorescence Imaging system (LI-COR Inc.).

Protein precipitation for detergent removal. In order to remove the detergents used in

our immunoprecipitation protocol, which are incompatible with the downstream MS

analysis, we have used the methanol/chloroform precipitation method as previously

described (Wessel and Flugge, 1984). Protein precipitates were then resuspended in 50

μl of 5× Invitrosol^{TM} LC/MS Protein Solubilizer (ThermoFisher). The samples were

incubated at 60°C for 5 mins, vortexed and incubated at 60°C for another 10 mins.

Ammonium bicarbonate (ABC buffer, 200 μl of 25 mM) was added per sample in order to

obtain 1× Invitrosol^{TM} concentration. The samples were separated into 5 tubes containing

50 μl of sample per tube.

LC/MS analysis. Samples were reduced in 10 mM dithiothreitol (DTT) for 30 mins at RT

and cysteines were alkylated in the dark in 60 mM iodoacetamide (IAA) for 20 min at RT.

Two μl of Trypsin/Lys-C mix (0.5 μg/μl, Promega, V5073) was added to each sample and

incubated for 3 hr at RT in the dark. After 3 hr, 150 μl of ABC buffer per tube was added

and incubated overnight at RT. The following day, the digestion was stopped by adding

200 μl of 5% acetonitrile, 3% trifluoroacetic acid. Tryptic peptides were desalted with

stage tips as previously described (Rappsilber et al., 2003). The desalted peptides were

concentrated in a Speed-Vac for 10 mins at low power. The peptides were reconstituted

with 15 μl of 5% acetonitrile, 0.1% formic acid and transferred to a microplate for LC/MS

analysis. Peptides were separated by reverse-phase chromatography using a Dionex

Ultimate 3000 nanoLC (Dionex/ThermoFisher) on a C18 capillary column (Acclaim

PepMap100 C18, 2 μm, 100 Å, 75 μm i.d. ×20 cm, ThermoFisher) at an eluent flow rate

of 300 nl/min. Mobile phase A contained 5% acetonitrile, 0.1% formic acid and mobile

phase B contained 80% acetonitrile 0.1% formic acid. The column was preequilibrated

with 5% mobile phase B followed by an increase to 65% mobile phase B in 75 min. The

eluted peptides were ionized online by electrospray ionization (ESI) and transferred into

an LTQ Orbitrap XL mass spectrometer (ThermoFisher) which was operated in the

positive mode to measure full scan MS spectra (300-1700 m/z in the Orbitrap analyzer at

resolution R = 30,000) followed by isolation and fragmentation (MS/MS) of the ten most

intense ions (in the LTQ iontrap) by collision-induced dissociation (CID).

MS/MS data analysis. The analysis of the raw MS and MS/MS Spectra was performed

using the MaxQuant software (version. 1.6.3.4). Initial maximum precursor and fragment

mass deviations were set to 7 ppm and 0.5 Da, respectively. Variable modification

(oxidation of methionine and N-terminal acetylation) and fixed modification (cysteine

carbamidomethylation) were set for the search against a UniProt FASTA formatted

database of the taxonomic clade *euthyneura* (including *Lymnaea stagnalis*). Trypsin with

a maximum of two missed cleavages was chosen for digestion. The minimum peptide

length was set to 7 amino acids and the false discovery rate (FDR) for peptide and protein

identification was set to 0.01.

Recombinant protein of LymCBP. The partial recombinant protein of the LymCBP HAT

domain (from 1240th arginine to 1641st lysine, GenBank accession number

BAE02656.1) with His_6-tag was synthesized in *Escherichia coli* and purified as

previously described with slight modifications (Hatakeyama et al., 2014, 2018a).

LymCBP catalytic domain was amplified by RT-PCR with cDNA prepared from

Lymnaea CNSs, PrimeSTAR® HS DNA Polymerase (Takara Clontech) and primers,

whose nucleotide sequences were summarized in Table 1 (No. 23, 24). The PCR

product was ligated into pET28a(+) plasmid (Novagen), which was linearized with *Nhe*

I and *Bam*H I, using In-Fusion® HD Cloning Plus kit (Takara Clontech). Then, the

Rosetta-gami™ 2(DE3) competent cells (Novagen) was transformed with this ligated

plasmid. Expression of the partial recombinant protein of the LymCBP catalytic domain

was induced by treatment with 0.32 mM of isopropyl β-D-thiogalactopyranoside (IPTG)

in TBG-M9 medium [0.8% Bacto Tryptone (BD) and 0.4% NaCl], followed by

purification using nickel-nitrilotriacetic acid (Ni-NTA) agarose resin (QIAGEN). The

recombinant protein was desalted with PD-10 column (Sephadex™ G-25M, GE

Healthcare), purified using a cation exchange column [HiTrap carboxymethyl (CM) FF

column (GE Healthcare)] installed in the ÄKTAprime plus system (GE Healthcare), and

condensed with Amicon® Ultra-15 column (Merck Millipore).

Acetylation assays using radioisotope-labeled acetyl-CoA. The procedures followed to

perform this experiment were modified from previous reports (Hatakeyama et al.,

2018a; 2018b). The recombinant LymCBP HAT domain (200 ng) were incubated with

the histone H1 purified from calf thymus (Sigma-Aldrich, 1 μg) and 7.4 kBq of

[^{14}C]-acetyl-CoA (Moravek, Inc.) for 4 hr in buffer containing 50 mM Tris-HCl (pH

8.0), 10% glycerol, 1 mM dithiothreitol, and 10 mM sodium butyrate. Curcumin

(MedChemExpress), anacardic acid (MedChemExpress) and garcinol (Enzo) prepared

in dimethyl sulfoxide (DMSO) were premixed at each 50 μM. Reactions were separated

in 14% SDS-PAGE gels. To detect β radiation released from radiocarbon, the gels were

dried on pieces of filter papers and exposed on imaging plates for several days. Signals

were observed using a fluoroimage analyzer (FLA-2000; Fuji Film).

Intracellular recording. The whole CNS, including the cerebral ganglia and the buccal

ganglia was dissected out from the snail in *Lymnaea* saline [50 mM NaCl, 1.6 mM KCl,

2.0 mM MgCl$_2$, 3.5 mM CaCl$_2$, and 10 mM HEPES (pH 7.9)]. The isolated CNS was

immobilized on a Sylgard-coated dish using stainless-steel pins. In order to physically

separate the cerebral giant cells and B1 motoneurons, a Vaseline dam was constructed

around the cerebral ganglia. This enabled us to apply a solution that contains the

inhibitors to the CGCs. The CGC and the B1 motoneuron were impaled with glass

microelectrodes filled with 3 M KCl giving tip resistances of 20–50 MΩ. A train of 5

spikes in the CGC, which was produced by a current injection (0.8–1.8 nA) for 1 sec,

evoked a single large compound EPSP in the B1 motoneuron (electric stimulator:

SEN-7203, Nihon Kohden, Tokyo, Japan; intracellular recording amplifier: MEZ-8300,

Nihon Kohden; AD converter: DIGIDATA 1322A, Axon Instruments, Foster City, CA,

USA). We used the neurons located in the ipsilateral side.

Injection of cAMP or 5'AMP to the CGC. Based on a previous study (Nakamura et al.,

1999), cyclic AMP (cAMP) or 5'AMP (Sigma-Aldrich), which was filled into a glass

microelectrode (50–100 MΩ) as a 200 mM solution dissolved in 20 mM Tris buffer

(pH7.5), was injected into the CGC by passing hyperpolarizing current pulses (50 msec

on, 50 msec off) of 4 nA for 20 min. Before and after the injection of hyperpolarizing

current into the CGC, the EPSP recorded in the B1 motoneuron is not changed by

activation of the CGC. The data were recorded in the same preparation at 30 min and 3

hr after cAMP or 5'AMP injection. The data at 0 hr (i.e. pre) were recorded before

cAMP injection. For the analysis of EPSP changes recorded in the B1 motoneuron, the

peak amplitude of the EPSP was measured.

Application of HAT inhibitors to the CGC. We used the following three different

inhibitors of HATs: curcumin (Sigma), anacardic acid (Sigma) and garcinol (Enzo).

Each of these drugs was dissolved in DMSO as a stock solution at 67.9 mM (curcumin),

50 mM (anacardic acid) and 20 mM (garcinol). Stock solutions were diluted 1000-fold

in *Lymnaea* saline then applied to the CGC area where the cerebral ganglia were

surrounded by Vaseline wall and incubation for 1 hr. After incubation with each drug,

intracellular recordings for the measurement of EPSPs were made. The data were

expressed as the mean ± SEM. Significant differences at $p < 0.05$ between pre-injection,

30 min and 3 hr after injection cAMP or 5'AMP were examined by one-way repeated

measures ANOVA and post hoc Tukey's test. For the normalized EPSP amplitude, the

values after 30 min and 3 hr for each group compared to the pre-injection were

examined with one-way repeated measured ANOVA and post hoc Tukey's test.

Results

Properties **of the** *LymCBP gene and protein*

The arrangement of the functional domains in LymCBP was the same as that of

other p300/CBPs (Fig. 1A). The full sequence of LymCBP (GenBank accession

number: AB217914) was about 75% identical to *Aplysia californica* CBP (ApCBP;

GenBank accession number: AAL54859) in amino acid sequence level. In functional

domains, such as Taz1 domain, KIX domain, bromodomain, HAT domain, and Taz2

domain of LymCBP, the amino acid sequences were individually over 90% identical to

those of ApCBP and human CBP (GenBank accession number: U47741) (Fig. 1B-D).

Comparing amino acid sequences among LymCBP, human CBP and human p300, the

functional amino acid residues in bromodomain for interacting with acetylated histones

(Dhalluin et al., 1999; Mujtaba et al., 2004; Plotnikov et al., 2014; Xu et al., 2017),

those in HAT domain for intramolecular interactions (Delvecchio et al., 2013; Ortega et

al., 2018; Zhang et al., 2018), and interaction with acetyl-CoA (Liu et al., 2008;

Maksimoska et al., 2014) were highly conserved (Table S1). Molecular phylogenetic

analysis of the p300/CBP proteins revealed that *Lymnaea* CBP is closely related to CBP

proteins found in other molluscan species such as *Aplysia* and the oyster *Crassostrea*

giga (Fig. 2).

Localization of LymCBP mRNA in the* Lymnaea *CNS

Absolute copy number of LymCBP mRNA in whole CNS and other tissues

were analyzed by quantitative PCR (qPCR) (Fig. 3). More than 15,000 copies of

LymCBP mRNA was contained in 6 ng of total RNA of CNSs (Fig. 3). This value was

1.4 to 2.9 times as those in other tissues (Fig. 3).

In situ hybridization showed the localization of LymCBP mRNA in *Lymnaea*

CNSs ($n = 8$). Weak intensity of the positive signal was detected in some neurons in the

buccal ganglia (Fig. 4A). The neurons localized at the ventral edge of the ganglia

seemed to be parts of the B4 cluster (B4CL) cells. Strong signal was observed in the

CGCs in the cerebral ganglia (Fig. 4B,C). Same signal was also seen in identifiable

large neurons, the RPeD1 (Fig. 4D) and the LPeD1 (Fig. 4E), in pedal ganglia. The

staining patterns of LymCBP mRNA in buccal, cerebral and pedal ganglia were very

similar with those of LymCREB1 (Sadamoto et al., 2004). The same positive signal of

LymCBP was also detected in unidentified neurons in parietal and visceral ganglia (Fig.

4F). No positive signals were detected in negative control experiments with sense

LymCBP cRNA probe (data not shown).

Existence of the protein form of LymCBP

Western blotting demonstrated the existence of the protein form of LymCBP in

the CNS. The epitope of the anti-human-CBP antibody is the *C*-terminal region of

human CBP, which is highly conserved in LymCBP (Fig. 1D). The western blotting

method detected two positive bands (Fig. 5A). The larger band, indicated with a filled

arrowhead around the marker of 245 kDa, was considered to be LymCBP, whose

molecular weight was estimated as 252.3 kDa from the amino acid sequence (Fig. 1).

The smaller band shown with an open arrowhead may be an isoform of LymCBP (Fig.

5A). In the EMBL-EBI database, twenty transcript variants of CBP are found in human,

and, of these isoforms, seven transcripts can be translated (Gene: "CREBBP",

ENSG00000005339). Therefore, it is suggested that *Lymnaea* also expresses the smaller

isoform of CBP.

Phosphorylation of CREB1 at Ser 133 is triggering its kinase inducible domain

(KID) to bind to the KID interacting domain (KIX) of CREB binding protein (CBP)

(Mayr and Montminy, 2001; Radhakrishnan et al., 1997; Shaywitz et al., 2000). To

identify the interaction of CBP with CREB proteins in our model organism *Lymnaea*

stagnalis we have utilized immunoprecipitation (IP) and liquid chromatography mass

spectrometry (LC/MS).

Although in pilot experiments we found that none of our CREB1 antibodies

worked well for IP assays with total CNS lysates from *Lymnaea stagnalis*, previous

studies have shown that human CREB2 (human activating transcription 4, hATF4), a

transcriptional repressor of CREB1, can directly interact with multiple domains of the

CBP including i. the KIX domain, ii. a region that contains the third zinc finger and the

E1A-interacting domain, iii. the C-terminal region containing the

p160/SRC-1-interacting domain and iii. the histone acetyltransferase domain which can

directly interact with histone acetyltransferase p300 transcriptional coactivator of CBP

(Liang and Hai, 1997). In addition, CBP and p300 can acetylate CREB2 in its bZIP

domain and enhance its transcriptional activity (Gachon et al., 2002; Lassot et al., 2005;

Liang and Hai, 1997). In order to confirm these reported interactions of CBP and

CREB2, we have performed IP assays with a specific antibody for CREB2 and CNS

homogenates from *Lymnaea stagnalis* (Fig. 5B), followed by identification of the

CREB2 interaction partners by LC/MS (Fig. 5C). As shown in Fig. 5C, after LC/MS,

the data analysis of the CID fragmentation spectrum from a peptide with the mass of

559.81 m/z revealed the amino-acid sequence of a unique peptide (KVFNPDELR) from

the UniprotKB entry E2QC33 of *Lymnaea stagnalis*, validating that *Lymnaea* CBP is an

interacting partner of CREB2 in the CNS. Importantly, these results further confirmed

that LymCBP is expressed in the *Lymnaea* CNS.

Effects of HAT inhibitors on LymCBP enzymatic activity

405   To investigate the enzymatic properties of LymCBP as a functional HAT, we

produced a partial recombinant protein of its HAT domain (from 1240th to 1641st, Fig.

1D) and investigated its acetyltransferase activity. The recombinant protein, to which an

His$_6$-tag was attached at its *N*-terminus, could be purified by precipitation using Ni-NTA

agarose resin. To remove contaminated proteins derived from bacterial cells, the

recombinant protein solution was desalted and then purified by cation-exchange

chromatography. This purification procedure was successful, and only the bands of the

recombinant protein were observed in the 12th to 19th fractions (arrowhead in Fig. 6A).

However, the concentrations of these preparations were too low to be measured by

Bradford protein assay. Then, the eight fractions were mixed and treated with a

centrifugal filter to condense the recombinant protein, resulting its concentration

increased to around 80 ng/μL (lane 2 in Fig. 6B), which was sufficient for the

biochemical experiments.

We first determined the incubation temperature for the *in vitro* acetylation assay.

A reaction mixture containing recombinant LymCBP HAT domain, bovine histone H1,

and [^{14}C]-acetyl-CoA were incubated for 4 hr at different temperatures; 15^{o}C, 20^{o}C,

25^{o}C, 30^{o}C and 37^{o}C. Separation by SDS-PAGE and Coomassie Brilliant Blue (CBB)

staining of the gel showed that same amount of histone H1 was loaded (lanes 3-7 in Fig.

6C). However, the band of recombinant LymCBP HAT domain was notably decreased

after incubation at 37^{o}C, suggesting that LymCBP was degraded (lane 7 in Fig. 6C). By

autoradiography, the positive signal indicating acetylation of histone H1 was observed

when incubated at 15^{o}C, 20^{o}C, 25^{o}C and 30^{o}C (lanes 8-11 in Fig. 6C), and this signal

was almost eliminated after incubation at 37°C (lane 12 in Fig. 6C). Therefore, we

decided to incubate samples at 30°C in following acetylation assays.

Next, the effects of HAT inhibitors, namely curcumin, anacardic acid and

garcinol, on the HAT activity of LymCBP was investigated. After adding each HAT

inhibitor (each at 50 μM final concentration) or DMSO as negative control, samples

were incubated for 4 h at 30°C and separated by SDS-PAGE. Although the amount of

each protein in each tube should be same, CBB staining showed that the band intensities

of histone H1 were decreased by incubating with curcumin and garcinol (lanes 14 and

16 in Fig. 6D), suggesting that these inhibitors degraded the protein. Autoradiography

showed that LymCBP acetylated histone H1 without any HAT inhibitors (lane 17 in Fig.

6D). In the presence of HAT inhibitors, LymCBP HAT activity was significantly

blocked and acetylation signals were eliminated (lanes 18-20 in Fig. 6D). These results

showed that similar to human CBP, LymCBP had HAT activity, and this enzymatic

activity was significantly blocked by HAT inhibitors.

Synaptic plasticity between a buccal motoneuron and its regulatory interneuron

induced by cAMP

The isolated *Lymnaea* CNS was incubated with DMSO saline for 1 h before the

injection of cAMP into the CGC cell body. We tested the changes in the EPSP in the B1

motoneuron by activation of the CGC after the cAMP injection. We measured the

amplitudes of the compound EPSPs 30 min and 3 h after the onset of cAMP injection.

The CGC was depolarized by current injection to fire action potentials, evoking the

compound EPSP in the B1 motoneuron. We used a train of 5 spikes in the CGC to evoke

large compound EPSPs in the B1 motoneuron because the single EPSPs in the B1

motoneuron were very small (approximately 1 mV). These compound EPSPs at 30 min

and 3 h after in preparations with cAMP-injected CGCs were about twice as large as pre

injection EPSPs [Fig. 7A, $F(2,18) = 10.35$, pre injection vs 30 min: $p < 0.05$; pre

injection vs 3 hr: $p < 0.05$, $n = 10$]. In contrast, there were no significant changes in the

5'AMP-injected preparations [Fig. 7B, $F(2,12) = 1.192$, pre injection vs 30 min: $p >$

0.05; pre injection vs 3 hr: $p > 0.05$, $n = 7$]. Thus, the effect of HAT inhibitors on the

compound EPSPs 30 min and 3 hr after cAMP injection were examined in the following

experiments.

Effects of HAT inhibitors on the EPSP in the B1 motoneuron

461   HAT inhibitors were shown to possess significant inhibitory effects on HAT

activity of LymCBP (Fig. 6C). Therefore, we examined the effects of the application of

these HAT inhibitors to CGCs on the compound B1 EPSPs that were found to be

increased after cAMP injection into the CGC. As reported previously (Nakamura et al.,

1999; Sadamoto et al., 2004), the EPSP in the B1 motoneuron is facilitated by injection

of cAMP into the CGC (Fig. 7A). This increase in EPSP amplitude was not observed

when 5'AMP was injected into CGCs instead of cAMP (Fig. 7B). Next, we investigated

changes to the cAMP-induced EPSP in the presence of the HAT inhibitors curcumin,

anacardic acid and garcinol. Incubation with 67.9 μM curcumin of the cerebral ganglion

for 1 h did not change the amplitude of the compound EPSPs in the B1 motoneuron (Fig.

7C). After 1-hr incubation with curcumin prior to the cAMP injection there was no

significant change in the amplitude of the compound EPSPs 30 min and 3 hr after the

cAMP injection compared with pre injection [Fig. 7C, $F(2,22) = 6.001$, pre injection vs

30 min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$, $n = 12$]. However, a significant

difference was observed between 30 min and 3 hr. The incubation with 50 μM anacardic

acid on the cerebral ganglion for 1 hr that did not enlarge of the compound EPSPs in the

B1 motoneuron (Fig. 7D). The 1-hr incubation with anacardic acid prior to the cAMP

injection, in this group, there was no significant differences in the amplitude of the

compound EPSPs 30 min and 3 hr after the cAMP injection compared with pre injection

[Fig. 7D, $F(2,10) = 1.563$, pre injection vs 30 min: $p > 0.05$; pre injection vs 3 hr: $p >$

0.05, $n = 6$]. The incubation with 20 μM garcinol on the cerebral ganglion for 1 hr that

did not enlarge of the compound EPSPs in the B1 motoneuron (Fig. 7E). The incubation

for 1 hr with garcinol prior to the cAMP injection, in this group, there was no significant

differences in the amplitude of the compound EPSPs 30 min and 3 hr after the cAMP

injection compared with pre injection [Fig. 5E, $F_{(2,14)} = 0.01214$, pre injection vs 30

min: $p > 0.05$; pre injection vs 3 hr: $p > 0.05$, $n = 8$].

487   The results from all these groups are summarized in Fig. 7F. The effect of HAT

inhibitors on increasing compound EPSPs amplitude was tested by taking a 0 min

baseline recording followed by a 30 min and a 3 hr recording after cAMP was injected.

The compound EPSPs were normalized to the initial baseline value. Incubation with the

HAT inhibitors before the injection of cAMP, did not result in increased compound

EPSPs amplitudes, as seen in the DMSO saline treatment group. In the host hoc Tukey's

test, there were significant differences in the following comparisons: DMSO saline

treatment cAMP injection group; pre injection vs 30 min; pre injection vs 3 hr,

curcumin treatment group; 30 min vs 3 hr).

Discussion

498   In the present study we, for the first time, identified a conserved CREB-binding

protein in the CNS of *Lymnaea* (LymCBP) and showed the importance of its HAT

activity for synaptic plasticity between an identified interneuron (CGC) and a

postsynaptic motoneuron (B1) of the feeding system. The CGC is a key interneuron for

consolidating both aversive and appetitive LTM for taste, and both LymCBP and

LymCREB1 were transcribed in this neuron (Fig. 4). A previous report showed that

inhibiting functions of LymCREB1 by pre-injection of excess amount of a CRE

oligonucleotide and siRNA of LymCREB1 into the CGC blocked cAMP-induced

synaptic facilitation between the CGCs and the B1 motoneurons (Sadamoto et al., 2004;

Wagatsuma et al., 2006), and suggested a crucial role of LymCREB1 as a transcriptional

factor in synaptic facilitation. Here, inhibition of LymCBP HAT activity using HAT

inhibitors similarly blocked the synaptic potentiation (Fig. 7), suggesting that this

synaptic facilitation requires the co-operation of LymCREB1 and LymCBP HAT

activity.

Gene structure of LymCBP

We succeeded in the cloning of the full-length cDNA of LymCBP from the

Lymnaea CNS. Comparing its deduced amino acid sequence with those of *Aplysia* and

human CBPs, several functional domains were found to be highly conserved (Fig. 1).

517 This result suggests that LymCBP has similar functions to mammalian CBP and p300

518 (Fig. 2). The KIX domain of CBP associates with the kinase-inducible domain (KID) of

519 CREB, and they construct a complex together (Chrivia et al., 1993). KID, which is also

520 called as P-box, is highly conserved in LymCREB1 (Sadamoto et al., 2004). High

521 conservation of the KIX domain in LymCBP and KID in LymCREB1 suggested that

522 LymCBP and LymCREB could interact via these domains. Additionally, a previous

523 report showed that the KIX domain of CBP is involved in the formation of spatial

524 memory following hippocampus-dependent learning (Chatterjee et al., 2020). Here, we

525 demonstrated that these two mRNAs were co-localized in the CGCs, key neurons for

526 LTM consolidation (Fig. 4). Taken together, the consolidation of LTM in *Lymnaea* is

527 suggested to be regulated by LymCBP and LymCREB1. The Taz1 and Taz2 domains of

528 LymCBP were also highly conserved (Fig. 1). These domains function as interfaces to

529 bind with a variety of transcription factors and HATs, such as Fos and

530 p300/CBP-associated factor (pCAF) (Glass et al., 1997). It was previously reported that

531 the bromodomain of mammalian CBP is important for acetylation of histones (Raisner

532 et al., 2018) and interaction with acetylated histones (Zeng et al., 2008; Plotnikov et al.,

533 2014). At least 15 amino acid residues in the bromodomain regulate interaction with

534 acetylated histones (Dhalluin et al., 1999; Plotnikov et al., 2014; Xu et al., 2017), and,

compared with amino acid sequences among human p300, human CBP and LymCBP,

14 out of 15 amino acid residues were conserved in LymCBP (Table S1). Other

functional amino acid residues are highly conserved in the RING domain (Table S2), the

acetyl-CoA-binding domain (Table S3), and the ZZ domain (Table S4). The RING

domain in human p300/CBP has an autoinhibitory function for the HAT domain

(Delvecchio et al., 2013; Ortega et al., 2019). The ZZ domain of human p300/CBP

recognizes histone tails and modulates its enzymatic activity (Legge et al., 2004; Zhang

et al., 2018). The most important domain in CBP is the KIX domain, which is well

known as an interface for interacting with CREB (Kwok et al., 1994), and this domain

is highly conserved in LymCBP (Fig. 1). These results suggested that not only the HAT

activity, which was shown in Fig. 6, but also other functions of CBP are highly

conserved in LymCBP.

Our immunoprecipitation assays indicated an interaction between LymCBP and

LymCREB2 (Fig. 5B). A similar interaction also has been shown in the human

homologues of these proteins (Liang and Hai, 1997). The interfaces in CBP for binding

with CREB2 include the KIX domain, Taz2 domain and HAT domain (Liang and Hai,

1997). These domains were highly conserved in LymCBP (Fig. 1). In addition,

acetylation of CREB2 by CBP was previously reported (Gachon et al., 2002). Therefore,

the interaction between LymCBP and LymCREB2 shown in Fig. 5B may be needed for

LymCREB2 acetylation by LymCBP.

Expression of LymCBP in the Lymnaea CNSs

In situ hybridization showed that LymCBP mRNA was transcribed in a limited

number of identifiable neurons (Fig. 4F). In particular, the CGC in the cerebral ganglia

exhibited a strong LymCBP signal (Fig. 4B,C). Both mRNA and protein forms of

LymCREB1 exist in the CGC (Ribeiro et al., 2003; Sadamoto et al., 2004), showing the

co-localization of LymCBP and LymCREB1 in this neuron. The CGC was identified to

be a key neuron for the long-lasting conditioned responses after taste aversion learning

(Kojima et al., 2001; Yamanaka et al., 2000; Ito et al., 2012; Sunada et al., 2017) and

appetitive classical conditioning (Kemenes et al., 2006; Nikitin et al., 2008; Vavoulis et

al., 2010; Marra et al., 2013; Nikitin et al. 2013). The same signal was also seen in

identified large neurons, the RPeD1 and the LPeD1, in the pedal ganglia (Fig. 4D,E).

The RPeD1 was especially reported to be necessary for LTM formation in operant

conditioning of aerial respiratory behavior (Scheibenstock et al., 2002). The staining

patterns of LymCBP mRNA in the buccal, cerebral and pedal ganglia were very similar

to those of LymCREB1 (Sadamoto et al., 2004), which possesses KID for interacting

with LymCBP and plays a role in the enhancement of transcriptional activity (Sadamoto

et al., 2004).

Our proteomics experiments confirmed the existence of translated LymCBP in

the *Lymnaea* CNS, corroborating the findings from the DNA and RNA level assays of

LymCBP expression.

In addition, we previously indicated the existence of both mRNA and protein

forms of another transcriptional factor, CCAAT/enhancer binding protein of *Lymnaea*

(LymC/EBP), in the CGC and the RPeD1 (Hatakeyama et al., 2004a). In mammalian

cells, C/EBP is acetylated by p300/CBP, its acetylation boosts C/EBP-mediated

transcriptional activity (Ceseña et al., 2007). Taken together, the interaction between

LymCBP and transcriptional factors, such as LymCREB1 and LymC/EBP, may be

involved in the consolidation of LTM of *Lymnaea*.

Involvement of LymCBP HAT activity in synaptic plasticity

In this study, we used HAT inhibitors (curcumin, anacardic acid and garcinol;

reviewed by Hatakeyama et al., 2013b) for biochemical and electrophysiological

experiments (Figs. 6 and 7). Biochemical analyses using the recombinant LymCBP HAT

domain showed that the HAT activity of LymCBP was significantly blocked by HAT

inhibitors. Molecular docking simulation indicated the amino acid residues of the

human p300, whose amino acid sequence of the HAT domain was highly identical with

human CBP, for interaction with garcinol (Coste et al., 2020). For interaction between

human p300 and garcinol, five amino acid residues (S1400, Y1414, D1444, Y1446 and

Q1455) in its HAT domain were required, and these were completely conserved in

LymCBP (S1356, Y1370, D1400, Y1402 and Q1411) (Table S5). Therefore, HAT

activity of LymCBP might be inhibited by similar interaction mechanisms with garcinol.

Electrophysiological experiments showed that these chemicals inhibited

cAMP-dependent synaptic plasticity between the CGCs and their follower neurons (Fig.

7). Although the block of synaptic plasticity by curcumin has not been reported in other

species, its inhibitory effects on newly acquired or reactivated fear memories were

indicated (Monsey et al., 2015). Similarly, garcinol was also reported to block newly

acquired memories in mammals (Zhao et al., 2012; Maddox et al., 2013a; Monsey et al.,

2017) and honeybees (Merschbaecher et al., 2016). Another HAT inhibitor, C646, which

was not used in this study, significantly impaired memory consolidation (Maddox et al.,

2013b; Merschbaecher et al., 2016). These inhibitory effects might be induced by

blocking HAT activities of CBP and other HATs. On the other hand, there are many

reports showing that curcumin has ameliorative effects from the impairment of memory

and synaptic plasticity caused by amyloid peptides (Ahmed et al., 2011; Zhang et al.,

2015), chronic stress (Liu et al., 2014), drugs (Soukhaklari et al., 2018), heavy metals

(Namgyal et al., 2020), aging (Belviranli et al., 2013; Cheng et al., 2013) and viral

proteins (Tang et al., 2009; Shen et al., 2015), suggesting that other molecular

mechanisms, except for inhibiting HAT activity, rescue memory and synaptic plasticity.

Acknowledgements

This work was partly supported by The Japan Society for the Promotion of Science

(JSPS), a Grant-in-Aid for Scientific Research (C) to D.H. (No. 26460562 and

17K08867), Takeda Science Foundation to D.H., Waseda University Grants for Special

Research Projects (2018K-141 and 2020C-135) to E.I. Work at the University of Sussex,

UK, was funded by grants from the Medical Research Council to G.K.

(MRC/G0400551), and the Biotechnology and Biological Sciences Research Council

(BBSRC) to G.K. and I.K. (BB/H009906/1) and to I.K. and G.K. (BB/P00766X/1),

respectively.

Author Contributions

Experiments performed in the UK: D.H., I.F., A.A., M.E. I.K. and G.K. conceived and

designed the experiments. I.F. and A.F. performed the Western blot assays. A.A. and

M.E. performed the immunoprecipitation and LC-MS experiments, respectively. I.F.,

A.F., A.A., M.E. and G.K. analyzed and visualized the data. A.A., M.E. I.K. and G.K.

wrote the relevant results sections of the paper. G.K. and I.K. obtained the funding for

the work performed at the University of Sussex.

Experiments performed in Japan: D.H. conceived and designed the experiments. D.H.,

Y.T. and E.I. purified total RNA from *Lymnaea* tissues. D.H., R.M., and T.K. prepared

the LymCBP recombinant protein and performed acetylation assay. H.S. and E.I.

performed electrophysiological experiments. T.W. created the phylogenetic tree.

D.H. wrote the first version of the paper and all the authors contributed to the final

version.

Competing Interests

The authors declare no competing financial interests.

References

Ahmed T, Gilani AH, Hosseinmardi N, Semnanian S, Enam SA, Fathollahi Y (2011)

Curcuminoids rescue long-term potentiation impaired by amyloid peptide in rat

hippocampal slices. SYNAPSE 65:572-582.

Alarcón JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER, Barco A (2004)

Chromatin acetylation, memory, and LTP are impaired in CBP[+/-] mice: a model for

the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. Neuron

42:947–959.

Belviranli M, Okudan N, Esra K, Atalik N, Öz M (2013) Curcumin improves spatial

memory and decreases oxidative damage in aged female rats. Biogerontology

14:187–196.

Cedar H, Bergman Y (2009) Linking DNA methylation and histone modification:

patterns and paradigms. Nat Rev Genet 10:295–304.

Ceseña TI, Cardinaux JR, Kwok R, Schwartz J (2007) CCAAT/enhancer-binding

protein (C/EBP) β is acetylated at multiple lysines: Acetylation of C/EBPβ at lysine

39 modulates its ability to activate transcription. J Biol Chem 282:956–967.

Chatterjee S, Mizar P, Cassel R, Neidl R, Selvi BR, Mohankrishna DV, Vedamurthy BM,

Schneider A, Bousiges O, Mathis C, Cassel JC, Eswaramoorthy M, Kundu TK,

Boutillier AL (2013) A novel activator of CBP/p300 acetyltransferases promotes

neurogenesis and extends memory duration in adult mice. J Neurosci

33:10698–10712.

Chatterjee S, Angelakos CC, Bahl E, Hawk JD, Gaine ME, Poplawski SG,

Schneider-Anthony A, Yadav M, Porcari GS, Cassel JC, Giese KP, Michaelson JJ,

Lyons LC, Boutillier AL, Abel T. (2020) The CBP KIX domain regulates long-term

memory and circadian activity. BMC Biol 18:155.

Cheng YF, Guo L, Xie YS, Liu YS, Zhang J, Wu QW, Li JM (2013) Curcumin rescues

aging-related loss of hippocampal synapse input specificity of long term

potentiation in mice. Neurochem Res 38:98–107.

Chrivia JC, Kwok RP, Lamb N, Hagiwara M, Montminy MR, Goodman RH (1993)

Phosphorylated CREB binds specifically to the nuclear protein CBP. Nature

365:855–859.

Coste C, Gérard N, Dinh CP, Bruguière A, Rouger C, Leong ST, Awang K, Richomme P,

Derbré S, Charreau B (2020) Targeting MHC regulation using polycyclic

polyprenylated acylphloroglucinols isolated from *Garcinia bancana*. Biomolecules

10:1266.

Delvecchio M, Gaucher J, Aguilar-Gurrieri C, Ortega E, Panne D (2013) Structure of

the p300 catalytic core and implications for chromatin targeting and HAT regulation.

Nat Struct Mol Biol 20:1040–1046.

Dhalluin C, Carlson JE, Zeng L, He C, Aggarwal AK, Zhou MM (1999) Structure and

ligand of a histone acetyltransferase bromodomain. Nature 399:491–496.

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

Gachon F, Devaux C, Mesnard JM (2002) Activation of HTLV-I transcription in the

presence of Tax is independent of the acetylation of CREB-2 (ATF-4). Virology

299:271–278.

Glass CK, Rose DW, Rosenfeld MG (1997) Nuclear receptor coactivators. Curr Opin

Cell Biol 9:222–232.

Guan Z, Giustetto M, Lomvardas S, Kim JH, Miniaci MC, Schwartz JH, Thanos D,

Kandel ER (2002) Integration of long-term-memory-related synaptic plasticity

involves bidirectional regulation of gene expression and chromatin structure. Cell

111:483–493.

Hatakeyama D, Fujito Y, Sakakibara M, Ito E (2004) Expression and distribution of

transcription factor CCAAT/enhancer-binding protein in the central nervous system

of *Lymnaea stagnalis*. Cell Tissue Res 318:631–641.

Hatakeyama D. Sadamoto H, Watanabe T, Wagatsuma A, Kobayashi S, Fujito Y,

Yamashita M, Sakakibara M, Kemenes G, Ito E (2006) Requirement of new protein

synthesis of a transcription factor for memory consolidation: Paradoxical changes

in mRNA and protein levels of C/EBP. J Mol Biol 356:569–577.

Hatakeyama D, Mita K, Kobayashi S, Sadamoto H, Fujito Y, Hiripi L, Elekes K, Ito E

(2010) Glutamate transporters in the central nervous system of a pond snail. J

Neurosci Res 88:1374–1386.

Hatakeyama D, Okuta A, Otsuka E, Lukowiak K, Ito E (2013a) Consolidation of

long-term memory by insulin in *Lymnaea* is not brought about by changing the

number of insulin receptors. Commun Integr Biol 6:e23955.

Hatakeyama D, Tierling S, Kuzuhara T, Müller U (2013b) Epigenetic regulation of gene

expression in the nervous system. pp. 151–171. In "Methods in Neuroethological

Research", Oka K and Ogawa H, Eds. Springer.

Hatakeyama D, Shoji M, Yamayoshi S, Hirota T, Nagae M, Yanagisawa S, Nakano M,

Ohmi N, Noda T, Kawaoka Y, Kuzuhara T (2014) A novel functional site in the PB2

subunit of influenza A virus essential for acetyl-CoA interaction, RNA polymerase

activity, and viral replication. J Biol Chem 289:24980–24994.

Hatakeyama D, Shoji M, Yamayoshi S, Yoh R, Ohmi N, Takenaka S, Saitoh A, Arakaki

Y, Masuda A, Komatsu T, Nagano R, Nakano N, Noda T, Kawaoka Y, Kuzuhara T

(2018a) Influenza A virus nucleoprotein is acetylated by histone acetyltransferases,

PCAF and GCN5. J Biol Chem 293:7126–7138.

Hatakeyama D, Ohmi N, Saitoh A, Makiyama K, Morioka M, Okazaki H, Kuzuhara T

(2018b) Acetylation of lysine residues in the recombinant nucleoprotein and VP40

matrix protein of Zaire Ebolavirus by eukaryotic histone acetyltransferases.

Biochem Biophys Res Comm 504:635–640.

Hirano Y, Ihara K, Masuda T, Yamamoto T, Iwata I, Takahashi A, Awata H, Nakamura N,

Takakura M, Suzuki Y, Horiuchi J, Okuno H, Saitoe M (2016) Shifting

transcriptional machinery is required for long-term memory maintenance and

modification in *Drosophila* mushroom bodies. Nat Commun 7:13471.

Ito E, Otsuka E, Hama N, Aonuma H, Okada R, Hatakeyama D, Fujito Y, Kobayashi S

(2012) Memory trace in feeding neural circuitry underlying conditioned taste

aversion in *Lymnaea*. PLoS One, 7:e43151.

Kandel ER (2001) The molecular biology of memory storage: A dialogue between genes

and synapses. Science 294:1030–1038.

Kemenes I, Straub VA, Nikitin ES, Staras K, O'Shea M, Kemenes G, Benjamin PR.

(2006) Role of delayed nonsynaptic neuronal plasticity in long-term associative

memory. Curr Biol 16:1269–1279.

Kojima S, Hosono T, Fujito Y, Ito E (2001) Optical detection of neuromodulatory

effects of conditioned taste aversion in the pond snail *Lymnaea stagnalis*. J

Neurobiol 49:118–128.

Korzus E, Rosenfeld MG, Mayford M (2004) CBP histone acetyltransferase activity is a

critical component of memory consolidation. Neuron 42:961–972.

Kwok RP, Lundblad JR, Chrivia JC, Richards JP, Bächinger HP, Brennan RG, Roberts

SG, Green MR, Goodman RH (1994) Nuclear protein CBP is a coactivator for the

transcription factor CREB. Nature 370:223–226.

Lakhina V, Arey RN, Kaletsky R, Kauffman A, Stein G, Keyes W, Xu D, Murphy CT

(2015) Genome-wide functional analysis of CREB/long-term memory-dependent

transcription reveals distinct basal and memory gene expression programs. Neuron

85:330–345.

Lassot I, Estrabaud E, Emiliani S, Benkirane M, Benarous R, Margottin-Goguet F.

(2005) p300 modulates ATF4 stability and transcriptional activity independently of

its acetyltransferase domain. J Biol Chem 280:41537–41545.

Laviv T, Scholl B, Parra-Bueno P, Foote B, Zhang C, Yan L, Hayano Y, Chu J, Yasuda R.

(2020) *In vivo* imaging of the coupling between neuronal and CREB activity in the

mouse brain. Neuron 105:799–812.

Legge GB, Martinez-Yamout MA, Hambly DM, Trinh T, Lee BM, Dyson HJ, Wright

PE (2004) ZZ domain of CBP: an unusual zinc finger fold in a protein interaction

module. J Mol Biol 343:1081–1093.

Liang G, Hai T (1997) Characterization of human activating transcription factor 4, a

transcriptional activator that interacts with multiple domains of cAMP-responsive

element-binding protein (CREB)-binding protein. J Biol Chem 272:24088–24095.

Liu X, Wang L, Zhao K, Thompson PR, Hwang Y, Marmorstein R, Cole PA (2008) The

structural basis of protein acetylation by the p300/CBP transcriptional coactivator.

Nature 451:846–850.

Liu D, Wang Z, Gao Z, Xie K, Zhang Q, Jiang H, Pang Q (2014) Effects of curcumin on

learning and memory deficits, BDNF, and ERK protein expression in rats exposed

to chronic unpredictable stress. Behav Brain Res 271:116–121.

Maddox SA, Watts CS, Doyère V, Schafe GE (2013a) A naturally-occurring histone

acetyltransferase inhibitor derived from *Garcinia indica* impairs newly acquired

and reactivated fear memories. PLoS One 8:e54463.

Maddox SA, Watts CS, Schafe GE (2013b) p300/CBP histone acetyltransferase activity

is required for newly acquired and reactivated fear memories in the lateral

amygdala. Learn Mem 20:109–119.

Maksimoska J, Segura-Peña D, Cole PA, Marmorstein R (2014) Structure of the p300

histone acetyltransferase bound to acetyl-coenzyme A and its analogues.

Biochemistry 53:3415–3422.

Marra V, O'Shea M, Benjamin PR, Kemenes I (2013) Susceptibility of memory

consolidation during lapses in recall. Nat Commun 4:1578.

Martinez-Balbás MA, Bannister AJ, Martin K, Haus-Seuffert P, Meisterernst M,

Kouzarides T (1998) The acetyltransferase activity of CBP stimulates transcription.

EMBO J 17:2886-2893.

Matos MR, Visser E, Kramvis I, van der Loo RJ, Gebuis T, Zalm R, Rao-Ruiz P,

Mansvelder HD, Smit AB, van den Oever MC. (2019) Memory strength gates the

involvement of a CREB-dependent cortical fear engram in remote memory. Nat

Commun 10:2315.

Mayr B, Montminy M (2001) Transcriptional regulation by the

phosphorylation-dependent factor CREB. Nat Rev Mol Cell Biol 2:599–609.

Merschbaecher K, Hatko L, Folz J, Mueller U (2016) Inhibition of different histone

acetyltransferases (HATs) uncovers transcription-dependent and -independent

acetylation-mediated mechanisms in memory formation. Learn Mem 23:83–89.

Monsey MS, Gerhard DM, Boyle LM, Briones MA, Seligsohn M, Schafe GE (2015) A

diet enriched with curcumin impairs newly acquired and reactivated fear memories.

Neuropsychopharmacology 40:1278–1288.

Monsey MS, Sanchez H, Taylor JR (2017) The naturally occurring compound *Garcinia*

indica selectively impairs the reconsolidation of a cocaine-associated memory.

Neuropsychopharmacology 42:587–597.

Mujtaba S, He Y, Zeng L, Yan S, Plotnikova O, Sachchidanand, Sanchez R, Zeleznik-Le

NJ, Ronai Z, Zhou MM (2004) Structural mechanism of the bromodomain of the

coactivator CBP in p53 transcriptional activation. Mol Cell 13:251–263.

Murakami J, Okada R, Sadamoto H, Kobayashi S, Mita K, Sakamoto Y, Yamagishi M,

Hatakeyama D, Otsuka E, Okuta A, Sunada H, Takigami S, Sakakibara M, Fujito Y,

Awaji M, Moriyama S, Lukowiak K, Ito E (2013) Involvement of insulin-like

peptide in long-term synaptic plasticity and long-term memory of the pond snail

Lymnaea stagnalis. J Neurosci 33:371–383.

Nakai J, Totani Y, Hatakeyama D, Dyakonova VE, Ito E (2020a) Another example of

conditioned taste aversion: Case of snails. Biology (Basel) 9:422.

Nakai J, Totani Y, Kojima S, Sakakibara M, Ito E (2020b) Features of behavioral

changes underlying conditioned taste aversion in the pond snail *Lymnaea stagnalis*.

Invert Neurosci 20:8.

Nakamura H, Kobayashi S, Kojima S, Urano A, Ito E (1999) PKA-dependent regulation

of synaptic enhancement between a buccal motor neuron and its regulatory

interneuron in *Lymnaea stagnalis*. Zool Sci 16:387–394.

Namgyal D, Ali S, Mehta R, Sarwat M (2020) The neuroprotective effect of curcumin

against Cd-induced neurotoxicity and hippocampal neurogenesis promotion

through CREB-BDNF signaling pathway. Toxicology 442:152542.

Nikitin ES, Vavoulis DV, Kemenes I, Marra V, Piger Z, Michel M, Feng J, O'Shea M,

Benjamin PR, Kemenes G. (2008) Persistent sodium current is a nonsynaptic

substrate for long-term associative memory. Curr Biol 18:1221–1226.

Nikitin ES, Balaban PM, Kemenes G (2013) Nonsynaptic plasticity underlies a

compartmentalized increase in synaptic efficacy after classical conditioning. Curr

Biol 23:614–619.

Ortega E, Rengachari S, Ibrahim Z, Hoghoughi N, Gaucher J, Holehouse AS, Khochbin

S, Panne D (2018) Transcription factor dimerization activates the p300

acetyltransferase. Nature 562:538–544.

Plotnikov AN, Yang S, Zhou TJ, Rusinova E, Frasca A, Zhou MM (2014) Structural

insights into acetylated-histone H4 recognition by the bromodomain-PHD finger

module of human transcriptional coactivator CBP. Structure 22:353–360.

Radhakrishnan I, Perez-Alvarado GC, Parker D, Dyson HJ, Montminy MR, Wright PE

(1997) Solution structure of the KIX domain of CBP bound to the transactivation

domain of CREB: a model for activator:coactivator interactions. Cell 91:741–752.

Raisner R, Kharbanda S, Jin L, Jeng E, Chan E, Merchant M, Haverty PM, Bainer R,

Cheung T, Arnott D, Flynn EM, Romero FA, Magnuson S, Gascoigne KE (2018)

Enhancer activity requires CBP/P300 bromodomain-dependent histone H3K27

acetylation. Cell Rep 24:1722–1729.

Rappsilber J, Ishihama Y, Mann M (2003) Stop and go extraction tips for

matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample

pretreatment in proteomics. Anal Chem 75:663–670.

Ribeiro MJ, Serfőző Z, Papp A, Kemenes I, O'Shea M, Yin JC, Benjamin PR, Kemenes

G (2003) Cyclic AMP response element-binding (CREB)-like proteins in a

molluscan brain: cellular localization and learning-induced phosphorylation. Eur J

Neurosci 18:1223–1234.

Sadamoto H, Sato H, Kobayashi S, Murakami J, Aonuma H, Ando H, Fujito Y, Hamano

K, Awaji M, Lukowiak K, Urano A, Ito E (2004) CREB in the pond snail *Lymnaea*

837 *stagnalis*: cloning, gene expression, and function in identifiable neurons of the

838 central nervous system. J Neurobiol 58:455–466.

839 Sadamoto H, Kitahashi T, Fujito Y, Ito E (2010) Learning-dependent gene expression of

840 CREB1 isoforms in the molluscan brain. Front Behav Neurosci 4:25.

841 Scheibenstock A, Krygier D, Haque Z, Syed N, Lukowiak K (2002) The soma of

842 RPeD1 must be present for long-term memory formation of associative learning in

843 *Lymnaea*. J Neurophysiol 88:1584–1591.

844 Shaywitz AJ, Dove SL, Kornhauser JM, Hochschild A, Greenberg ME (2000)

845 Magnitude of the CREB-dependent transcriptional response is determined by the

846 strength of the interaction between the kinase-inducible domain of CREB and the

847 KIX domain of CREB-binding protein. Mol Cell Biol 20:9409–9422.

848 Shen LL, Jiang ML, Liu SS, Cai MC, Hong ZQ, Lin LQ, Xing YY, Chen GL, Pan R,

849 Yang LJ, Xu Y, Dong J (2015) Curcumin improves synaptic plasticity impairment

850 induced by HIV-1gp120 V3 loop. Neural Regen Res 10:925–931.

851 Soukhaklari R, Moezi L, Pirsalami F, Ashjazadeh N, Moosavi M (2018) Curcumin

852 ameliorates scopolamine-induced mice memory retrieval deficit and restores

853 hippocampal p-Akt and p-GSX-3β. Eur J Pharmacol 841:28-32.

854 Sunada H, Lukowiak K, Ito E (2017) Cerebral giant cells are necessary for the

formation and recall of memory of conditioned taste aversion in *Lymnaea*. Zoolog

Sci 34:72–80.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular

Evolutionary Genetics Analysis version 6.0. Mol Biol Evol 30:2725–2729.

Tang H, Lu D, Pan R, Qin X, Xiong H, Dong J (2009) Curcumin improves spatial

memory impairment induced by human immunodeficirncy virus type i glycoprotein

V3 loop peptide in rats. Life Sci 85:1–10.

Totani Y, Nakai J, Dyakonova VE, Lukowiak K, Sakakibara M, Ito E (2020) Induction

of LTM following an insulin injection. eNeuro 7:ENEURO.0088-20.2020.

Vavoulis DV, Nikitin ES, Kemenes I, Marra V, Feng J, Benjamin PR, Kemenes G.

(2010) Balanced plasticity and stability of the electrical properties of a molluscan

modulatory interneuron after classical conditioning: a computational study. Front

Behav Neurosci 4:19.

Vecsey CG, Hawk JD, Lattal KM, Stein JM, Fabian SA, Attner MA, Cabrera SM,

McDonough CB, Brindle PK, Abel T, Wood MA (2007) Histone deacetylase

inhibitors enhance memory and synaptic plasticity via CREB:CBP-dependent

transcriptional activation. J Neurosci 27:6128–6140.

Xu L, Cheng A, Huang M, Zhang J, Jiang Y, Wang C, Li F, Bao H, Gao J, Wang N, Liu

J, Wu J, Wong CCL, Ruan K (2017) Structural insight into the recognition of

acetylated histone H3K56ac mediated by the bromodomain of CREB-binding

protein. FEBS J 284:3422–3436.

Wagatsuma A, Sadamoto H, Kitahashi T, Lukowiak K, Urano A, Ito E (2005)

Determination of the exact copy numbers of particular mRNAs in a single cell by

quantitative real-time RT-PCR. J Exp Biol 208:2389–2398.

Wagatsuma A, Azami S, Sakura M, Hatakeyama D, Aonuma H, Ito E (2006) De Novo

synthesis of CREB in a presynaptic neuron is required for synaptic enhancement

involved in memory consolidation. J Neurosci Res 84:954–960.

Wessel D, Flugge UI (1984) A method for the quantitative recovery of protein in dilute

solution in the presence of detergents and lipids. Anal Biochem 138:1417–143.

Yamanaka M, Hatakeyama D, Sadamoto H, Kimura T, Ito E. (2000) Development of

key neurons for learning stimulates learning ability in *Lymnaea stagnalis*. Neurosci

Lett 278:113–116.

Zeng L, Zhang Q, Gerona-Navarro G, Moshkina N, Zhou MM (2008) Structural basis of

site-specific histone recognition by the bromodomains of human coactivators PCAF

and CBP/p300. Structure 16:643–652.

Zhang L, Fang Y, Xu Y, Lian Y, Xie N, Wu T, Zhang H, Sun L, Zhang R, Wang Z

(2015) Curcumin improves amyloid β-peptide (1-42) induced spatial memory

deficits through BDNF-ERK signaling pathway. PLoS One 10:e0131525.

Zhang Y, Xue Y, Shi J, Ahn JW, Mi W, Ali M, Wang X, Klein BJ, Wen H, Li W, Shi X,

Kutateladze TG (2018) The ZZ domain of p300 mediates specificity of the adjacent

HAT domain for histone H3. Nat Struct Mol Biol 25:841–849.

Zhao Z, Fan L, Fortress AM, Boulware MI, Frick KM (2012) Hippocampal histone

acetylation regulates object recognition and the estradiol-induced enhancement of

object recognition. J Neurosci 32:2344–2351.

Zhou L, Zhang Y, Liu RY, Smolen P, Cleary LJ, Byrne JH (2015) Rescue of impaired

long-term facilitation at sensorimotor synapses of *Aplysia* following siRNA

knockdown of CREB1. J Neurosci 35:1617–1626.

Figure Legends

Fig. 1. Predicted amino acid sequence of *Lymnaea* CBP. (A) Diagram of the structure of

Lymnaea CBP (GenBank accession number: AB217914). Main functional domains

(Taz1 domain, KIX domain, bromodomain, acetylation domain and Taz2 domain) are

highlighted with patterned bars. Alignments of amino acid sequences of (B) Taz1

domain, (C) KIX domain and (D) compound of the bromodomain, HAT domain and

Taz2 domain between *Lymnaea* CBP and *Aplysia* CBP (GenBank accession number:

AY064470) and human CBP (GenBank accession number: U47741). The patterns of

bars in (D) correspond to those shown in (A).

Fig. 2. Molecular phylogenetic tree of *Lymnaea* CBP and other p300/CBP proteins. The

scale bar indicates 0.1 substitutions per site. Bootstrap values are shown at the nodes. A

CBP homologue of the Choanoflagellate *Salpingoeca rosetta* was used as an outgroup.

The GenBank accession numbers of p300/CBP-like proteins are listed in Table 1.

Fig. 3. Quantification of LymCBP mRNA in the whole CNS of *Lymnaea*. LymCBP is

more abundantly transcribed in the CNS than other tissues of *Lymnaea*. The data are

expressed as the mean ± SEM; $p < 0.01$, $**p < 0.01$, Tukey's post-hoc test.

Fig. 4. Localization of LymCBP mRNA in identifiable neurons shown by *in situ*

hybridization. (A) Weak signal of LymCBP could be observed in several neurons

(arrowheads), especially in the B4CL feeding motoneurons (arrows), in the buccal

ganglia. Bu. G.: buccal ganglion, L: left; R: right. (B-E) Strong signal of LymCBP

mRNA was observed in both CGCs (arrowheads in B and C), RPeD1 (arrowhead in D)

and LPeD1 (arrowhead in E). Ce. G.: cerebral ganglion, Pe. G.: pedal ganglion. (F)

Schematic drawing of the distribution of neurons containing LymCBP mRNA. Pa. G.:

parietal ganglion, Vi. G.: visceral ganglion. Scale bars = 100 μm (A, D and E) and 20

μm (B and C).

Fig. 5. Identification of the protein form of LymCBP. (A) Existence of LymCBP protein

in CNSs was confirmed by western blotting. Molecular weight of the positive band (245

kDa; filled arrowhead) was consistent with theoretical value calculated from amino acid

sequence of LymCBP (252.3 kDa). A putative smaller isoform of LymCBP was also

detected (open arrowhead). (B, C) CREB-Binding Protein identified as an interacting

partner of CREB2 by MS/MS analysis in the CNS of *Lymnaea stagnalis*. (B) Sixty

extracted CNS of *Lymnaea stagnalis* were pooled, homogenised and subjected to

immunoprecipitation using a specific CREB2 antibody. Immune complexes were

immunoblotted for CREB2. (C) MS/MS Spectrum of the CBP identified as one of the

interacting partners of CREB2 in the CNS of *Lymnaea stagnalis*. CREB2 IP product

was subjected for MS/MS analysis and a CBP specific peptide for *Lymnaea* CBP

(KVFNPDELR) was identified.

Fig. 6. HAT activity of LymCBP was blocked by inhibitors. (A) The partial recombinant

protein of LymCBP HAT domain purified by cation-exchange chromatography was

detected in fractions No. 12–19 (bands shown with arrowhead). The concentrations of

these preparations were too low to be measured by Bradford protein assay. (B) All

fractions obtained by cation-exchange chromatography were mixed (lane 1), and the

recombinant protein was condensed using a centrifugal filter. A high enough

concentration of the recombinant protein for biochemical experiments was obtained

(lane 2). (C) Condensed recombinant LymCBP HAT domain was incubated with histone

H1 at different temperatures; 15ºC, 20ºC, 25ºC, 30ºC and 37ºC. CBB staining of the gel

showed that same amount of histone H1 was loaded (open arrowhead, lanes 3-7).

However, the band of recombinant LymCBP HAT domain (filled arrowhead) was

drastically decreased after incubation at 37°C, suggesting that LymCBP was denatured

(lane 7). A positive signal of acetylation was observed after incubation at 15°C, 20°C,

25°C and 30°C (lanes 8-11). This signal was almost eliminated after incubation at 37°C

(lane 12). (D) Effects of the HAT inhibitors curcumin, anacardic acid and garcinol.

Reaction mixtures were incubated at 30°C. Although the same amount of recombinant

LymCBP HAT domain (filled arrowhead) and histone H1 (open arrowhead) was

incubated, CBB staining showed that the band signals of histone H1 were attenuated

after incubating with curcumin (lane 14) and garcinol (lane 16), suggesting that these

inhibitors degraded the protein. Autoradiography indicated that, without any HAT

inhibitor, positive signal of acetylation was detected on the band of histone H1 (lane 17).

This signal was eliminated by addition of each HAT inhibitor (lanes 18-20), showing

that the acetylation activity of LymCBP recombinant protein was significantly blocked

by HAT inhibitors.

Fig. 7. Long-term synaptic changes at synapses between the CGC and the B1

motoneuron in the isolated CNS are blocked by HAT inhibitors. EPSPs were evoked in

the B1 motoneuron by injection of the depolarizing current into the CGC. The isolated

CNS was incubated with DMSO saline or HAT inhibitors for 1 hr prior to injection of

cAMP. (A) cAMP injection into the CGC cell body increased EPSP amplitude in B1

motoneuron at both 30 min and 3 h after the injection of the cAMP. (B) In contrast,

EPSP amplitude increase was not observed in the group which was injected with 5'AMP.

(C, D, E) The CNS was incubated with saline containing one of the HAT inhibitors

curcumin (67.9 μM) or anacardic acid (50 μM) or garcinol (20 μM)for 1 h prior to

injection of the cAMP. In these groups, the EPSP was not increased either at 30 min or

3h after the injection of cAMP. (F) The enhancement of B1 EPSPs is shown as the

summarized data. Injection of cAMP into the CGC cell body increased EPSP amplitude

in the B1 motoneuron, but pre-treatment with either HAT inhibitor did not elicit

enlargement of EPSP. The data are expressed as the mean ± SEM, *$p < 0.05$, Tukey's

post-hoc test.

Table 1. Nucleotide sequences of primers used for cloning (No. 1-14), synthesis for *in*

situ hybridization probe (No. 15-16), synthesis of recombinant protein (No. 17-18), and

qRT-PCR (No. 19-24) of LymCBP. Amino acid sequences of ApCBP for designing

degenerate PCR (DGPCR) was shown with nucleotide sequences.

Names of primers and probes		Nucleotide sequences
1.	DGPCR_Taz1_fw	5'-CTS ATC CAG CAG CAG CTB GTB CTS C-3' (LIQQQLVL)
2.	DGPCR_Taz1_rv	5'-ATT TGW CGA GAW GAR GCA CAR TGR G-3' (AHCASSRQII)
3.	DGPCR_KIX_fw	5'-CTG GTS CAR GCY ATY TTY CC-3' (KLVQAIFP)
4.	DGPCR_KIX_rv	5'-GTA SAT YTT CTC RGC SAG CAA RTG-3' (HLLAEKIY)
5.	DGPCR_Taz2_fw	5'-TCC ATT CAR CGC TGY ATC TCN TCH C-3' (SIQRCISSL)
6.	DGPCR_Taz2_rv	5'-ACA GAA TGG GAC CKG RCA YTT RTT YTC-3' (ENKCQVPFC)
7.	5'RACE_outer	5'-TTA CTG GCT GGT ATA GAT TTA ACA ATA TCT GGA GC-3'
8.	5'RACE_inner	5'-TAT GTT ACC CAA AAG ATT ACT CAT ATG ACT GCT AGG-3'
9.	3'RACE_outer	5'-CTA ACT GTC GTA TCA ACT CGT GCA TCA AG-3'
10.	3'RACE_inner	5'-GCA TCA AGA TGA AAC GTG TTG TTT CAC-3'
11.	Taz1-KIX_fw	5'-ACA TGC AAC AAA GGA AAA TCT TG-3'
12.	Taz1-KIX_rv	5'-TGA TAA TAT TCT CCT CTA TTG TTA GCA G-3'
13.	KIX-Taz2_fw	5'-ACA TGC AAC AAA GGA AAA TCT TGT GAA GTG GCC C-3'
14.	KIX-Taz2_rv	5'-CGT ACA ATG TTT GGC ATG GTA ACA ACA GAG TGC-3'
15.	qPCR_cloning_fw	5'-GAT GTA AAC ATG GCC GAC CAC CAA GTT G-3'
16.	qPCR_cloning_rv	5'-AGA CGA TGT AGC AAC TTT GTT GTT ATT GGC-3'
17.	qPCR_RT	5'-TGT TAA GAG ACA TCG GCA TTG ATG-3'
18.	qPCR_fw	5'-GCC CTC CGG CCA ACA AGA A-3'
19.	qPCR_rv	5'-TAT TAT CGC TGG GTG TAT TGA GAG AT-3'
20.	qPCR_probe	5'-[Fam]-ACC CAG AAT CGG ATC AC-[Tamra]-3'
21.	in-situ-probe_fw	5'-<u>GAT ATC</u> TCT TGA AAT CTT AGT GCT GTG TTG ATG-3' (*Eco*R V)
22.	in-situ-probe_rv	5'-<u>GGA TCC</u> ATC AGC AGA AGT CTG TCG AGT ACC-3' (*Bam*H I)
23.	recombinant_fw	5'-<u>CGC GGC AGC CAT ATG</u> **GCT AGC** <u>AGA CGA AAA GAA AAC</u> <u>AAA TTT ACA GCC</u>-3' [nucleotide sequence of pET28a(+) plasmid, **the restriction site of *Nhe* I**, nucleotide sequence of LymCBP]
24.	recombinant_rv	5'-<u>ACG GAG CTC GAA TTC</u> **GGA TCC** TTA <u>ATC AGG TTC TGT GAT</u> <u>TGG TGG CAG</u>-3' [nucleotide sequence of pET28a(+) plasmid, **the restriction site of *Bam*H I**, nucleotide sequence of LymCBP]

Table 2. GenBank accession numbers of various CBP and p300 used for molecular phylogenetic tree shown in Fig. 2.

Species	Protein	GenBank Accession No.
Alligator mississippiensis	CBP	XP_006264860
Alligator mississippiensis	p300	XP_014463094
Apis mellifera	CBP	XP_006568897
Aplysia californica	CBP	NP_001191640.1
Blattella germanica	CBP	CUT08824
Brugia malayi	CBP	CTP80907
Caenorhabditis elegans	CBP	NP_499161
Capitella teleta	CBP	ELU13629
Ciona intestinalis	CBP	XP_009860762.1
Crassostrea gigas	CBP	XP_011427368.1
Danio rerio	CBP	XP_009297440.1
Drosophila melanogaster	nejire	NP_001188575.1
Gallus gallus	CBP	XP_015150108.1
Gallus gallus	p300	XP_004937767
Homo sapiens	CBP	NP_004371.2
Homo sapiens	p300	NP_001420
Hydra vulgaris	CBP	XP_002156492.2
Hymenolepis microstoma	CBP	CDS30104
Limulus polyphemus	CBP	XP_013776092
Loa loa	CBP	EJD74639
Lymnaea stagnalis	CBP	AB217914
Mus musculus	CBP	NP_001020603.1
Mus musculus	p300	NP_808489
Opisthorchis viverrini	CBP	XP_009171514
Parasteatoda tepidariorum	CBP	XP_015922458
Priapulus caudatus	CBP	XP_014664166
Rattus norvegicus	CBP	NP_596872.3
Rattus norvegicus	p300	XP_017450856
Salpingoeca rosetta	CBP	XP_004998720
Schistosoma mansoni	CBP	CCD78413
Strongylocentrotus purpuratus	CBP	XP_011677827.1
Takifugu rubripes	CBP	XP_011602271.1

| 1060 | *Xenopus laevis* | CBP1 | NP_001088637.1 |
| 1061 | *Xenopus laevis* | CBP2 | NP_001192159.1 |

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7