Title

A global dataset for the projected impacts of climate

change on four major crops

Authors

Toshihiro Hasegawa[1], Hitomi Wakatsuki[1], Hui Ju[2], Shalika Vyas[3], Gerald C. Nelson[4], Aidan

Farrell[5], Delphine Deryng[6], Francisco Meza[7], David Makowski[8]

Affiliations

1. Institute for Agro-Environmental Sciences, National Agricultural and Food Research

Organization, Tsukuba, Ibaraki, 305-8604, Japan

2. Institute of Environment and sustainable Development in Agriculture, Chinese Academy of

Agricultural Sciences (IEDA,CAAS), Beijing, 100081, China

3. Borlaug Institute for South Asia, CIMMYT, New Delhi, 110012, India

4. University of Illinois, Urbana, IL.,USA

5. The University of the West Indies, St. Augustine, Trinidad

6. IRI THESys, Humboldt-Universität zu Berlin, Berlin, 10099, Germany

7. Pontificia Universidad Católica de Chile, Santiago, Chile

8. Applied mathematics and computer science (MIA 518), INRAE AgroParisTech, Université

Paris-Saclay, 75231, Paris, France.

corresponding author(s): Toshihiro Hasegawa (thase@affrc.go.jp)

Abstract

Reliable estimates of the impacts of climate change on crop production are critical for

assessing the sustainability of food systems. Global, regional, and site-specific crop

simulation studies have been conducted for nearly four decades, representing valuable

sources of information for climate change impact assessments. However, the wealth of data

produced by these studies has not been made publicly available. Here, we develop a global

dataset by consolidating previously published meta-analyses and data collected through a

new literature search covering recent crop simulations. The new global dataset builds on

8314 simulations from 203 studies published between 1984 and 2020. It contains projected

yields of four major crops (maize, rice, soybean, and wheat) in 91 countries under major

emission scenarios for the 21st century, with and without adaptation measures, along with

geographical coordinates, current temperatures, local and global warming levels. This

dataset provides a basis for a comprehensive understanding of the impacts of climate

change on crop production and will facilitate the rapidly developing data-driven machine

learning applications.

Background & Summary

Climate change affects many processes of food systems directly and indirectly[1], but the

primary effects often appear in crop production. Projections of crop production under future

climate change have been studied since the early 1980s. From the 1990s onward,

researchers have used future climate data and crop simulation models to project the impacts

of climate change on crop yields under various scenarios[2]. Since then, crop simulation

models have been used in hundreds of studies to simulate yields for different crops under a

range of climate scenarios and growing conditions[3]. The results have been periodically

reviewed and assessed by various national and international organisations, in particular by

the Intergovernmental Panel on Climate Change (IPCC) working group II, which provides

policy-relevant scientific evidence for the impacts of and adaptation to climate change[3].

Review studies covering the last five IPCC assessment cycles confirm that the overall effects
are negative but vary significantly among regions[4,5].
53
Before 2010, simulation studies were conducted mainly by individual research groups using
different climate models, target years, spatial resolution with local management and cultivar
conditions. Since 2010, however, significant efforts have been made to coordinate modelling
studies through Agricultural Model Intercomparison and Improvement Project (AgMIP)[6],
which intercompares multiple crop models using the standardised inputs. Early AgMIP
activities have successfully disentangled sources of uncertainties in crop yield projections
and revealed that yield projections are variable among crop models and that model
ensemble mean or median often works better than a single model[7-10], underpinning the
importance of datasets based on multiple crop models.
63
Data sets including crop model simulations produced by AgMIP were subjected to statistical
analysis and the results were used to quantify the impacts of climate change on major
crops[11,12]. A versatile tool to aggregate simulated results is already available for global
gridded studies[13] to facilitate access to the data. Besides these coordinated efforts, however,
many simulation results are scattered and not readily available for meta-analysis. To deliver
policy-relevant quantitative information, we need to develop a shared and well-documented
database that can be used to assess the impacts of different climate and adaptation
scenarios on crop yields.
72
Here, we compile and develop a global database for potential use for the IPCC Working
Group II assessment, drawing on two data sources. The first source of information is the
dataset used in the meta-analysis of Aggarwal, et al. [5], which includes studies considered in
the previous five cycles of IPCC assessments[4,14]. The second source is based on a new
literature search of studies published during the sixth IPCC assessment cycle (covering the
period 2014-2020) reporting crop simulations produced for several contrasting climate
change scenarios. Based on these two sources of information, the combined dataset covers
all six cycles of the IPCC assessment and can serve as a strong basis for the analysis from the
sixth IPCC assessment onward.
82
83 The dataset contains the most relevant variables for evaluating climate change impacts on
84 yields of maize, rice, soybean, and rice for the 21st century. They include geographical
85 coordinates, crop species, CO_2 emission scenarios, CO_2 concentrations, current temperature
86 levels, local and global warming degrees, the relative changes in yield as a percentage of the
87 baseline period, and the projected effect size of adaptation options.
88
89
Methods
Data collection. As shown in the PRISMA diagram (Fig. 1), we used two data sources to
develop this dataset. The first source is based on the previous meta-analysis by Aggarwal et
al.[5], which includes studies published before 2016 (Aggarwal-DS, hereafter). This meta-
analysis builds on the dataset used for the 5[th] IPCC assessment report[4,14] and an additional
search through three types of databases; Scientific database (Scopus, Web of Science, CAB
Direct, JSTOR, Agricola etc), journals and open access repositories, and institutional Websites
(FAO database, AgMIP Database, World Bank, etc.) and Google Scholar. See Aggarwal et al.[5]
for details. Briefly, the search terms used by Aggarwal et al.[5] include "agriculture" or "crop
"or "farm" or "crop yield" or "crop yields" or "farm yields" or "crop productivity" or
"agricultural productivity" or "maize" or "rice" or "wheat" and "climate change assessment"
or "climate impacts" or "impact assessments" or "climate change impact" or "climate impact"

or "effect of climate" or "impact of climate change". The number of selected papers covering
the four major crops is 166. We further screened them according to the availability of local
temperature rise and geographical information, and traceability, resulting in 99 studies
published between 1984 and 2016.

The second source is based on a new recent literature search conducted using Scopus in
March 2020 for four major crops (maize, rice, soybean, and wheat) for peer-review papers
published from 2014 onward in line with the sixth assessment cycle of IPCC. For maize, the
following search equation was used:

PUBYEAR > 2013 AND TITLE-ABS-KEY((maize OR corn) AND (("greenhouse gas" OR "global
warming" OR "climate change" OR "climate variability" OR "climate warming")) AND NOT
(emissions OR mitigation OR REDD OR MRV))
113
114 Similar search equations were used for the other crops. Collectively, this search returned a
115 total of 4703 references between 2014 and 2020: 1899 for maize, 1790 for wheat, 757 for
116 rice, and 257 for soybean with some duplications because some papers studied multiple
117 crops. Removing the duplicates, the number is down to 3816 studies.
118
To collect climate-scenario-based simulations, we then selected a subset of studies including
the following terms related to climate scenarios in titles, abstracts, or authors' keywords;
"RCP", "RCP2.6", "RCP6.0", "RCP4.5", "RCP8.5", "CMIP5", and "CMIP6". RCP stands for the
Representative Concentration Pathways[15], and each RCP corresponds to a greenhouse gas
concentration trajectory describing different future greenhouse gas emission levels. The
number followed by RCP is the level of radiative forcing (Wm^{-2}) reached at the end of the 21st
century, which increases with the volume of greenhouse gas emitted to the atmosphere[16].
CMIP5[17] and CMIP6[18] are the Coupled Model Intercomparison Project Phase 5 and Phase 6,
respectively, where groups of different earth system models (ESMs) provide global-scale
climate projections based on different RCPs. Additionally, "process-based model" was used
to search in the authors' keywords to select for studies that use crop simulation models
under CMIP5 or CMIP6 climate scenarios. As of March 2020, no results were found for CMIP6
in any search results.
132
This screening process resulted in a total of 207 references all together for four major crops.
These studies were further evaluated for their variables and data availability; studies not
reporting yield data were excluded. Projected yields with and without adaptations and yields
of the baseline period were extracted, along with geographical coordinates, crop species,
greenhouse gas emission scenarios, and adaptation options. We also tried to obtain local and
global temperature changes and CO_2 concentrations as much as possible. The authors of the
two grid simulation studies provided the aggregated results for countries or regions[19,20]. The
results from different ESMs were averaged.
141
We consolidated these two sources and removed duplicates, resulting in a total of 203
studies. Both datasets include studies with different spatial scales: site-based, regional, and
global. Among these, the results from the global gridded crop models were aggregated to
country levels, and we focused on top-producing countries, which account for 95% of the
world's   production   of   each   commodity   as   of   2010   (FAOSTAT,
http://www.fao.org/faostat/en/, accessed on September 4, 2020). As a result, the dataset
contains 8,314 sets of yield projections during the 21st century from studies published
between 1984 and 2020 (Online-only Table 1).

150
151
152 **Fig. 1.** A diagram depicting paper collection and selection from the two data sources. N is the
153 number of studies.
154
155 **Relative yield impacts.** Simulated grain mass per unit land area is used to derive the impact
156 of climate change on yield (YI), which is defined as:
157
158 $$YI(\%) = (Y_f/Y_b - 1) \times 100$$
159
160 Where Y_f is the future yield, and Y_b is the baseline yield. However, one study[20] accounts for
161 technological factors change over time and simulations under both climate change and
162 counterfactual non-climate change scenarios and, for this study, YI is expressed as:
163
$$YI(\%) = ((Y_{f\_cc} - Y_{bcc} - Y_{f\_ncc} - Y_{b\_ncc})/Y_{b\_cc} - 1) \times 100$$

Where Y_{f_cc} and Y_{b_cc} are the future and baseline yield with climate change, Y_{f_ncc} and Y_{b_ncc} is
the future yield under counterfactual no climate change scenario.

Adaptation to climate changes. Various management or cultivar options are tested in the
simulations. If the authors of the article consider these options as ways to adapt crops to
climate change, we treat them as adaptation options, which are categorised into fertiliser,
irrigation, cultivar, soil organic matter management, planting time, tillage, and others.
Specifically, in the fertiliser option, if the amount and timing of fertiliser application are
changed from the current conventional method, we treat them as adaptation. In the
irrigation option, if the simulation program determines the irrigation scheduling based on
the crop growth, climatic and soil moisture conditions, we treat this as adaptation because
the management is adjusted to the future climatic conditions. If rainfed and irrigated
conditions are simulated separately, we do not consider irrigation as an adaptation. We
define cultivar option as the use of cultivars of different maturity groups and/or higher heat
tolerance compared to conventional cultivars. The planting time option corresponds to a
shift of planting time from conventional timing. If multiple planting times are tested, we
select the one that gives the best yield. The soil organic matter management option
corresponds to application of compost and/or crop residue. The tillage option corresponds
to reduced- or no-till cultivation compared to no conventional tillage. When studies consider
adaptation options, we compute YI from the ratio of yield with adaptation under climate
change to baseline yield without adaptation.

Warming level. Both local temperature rise (ΔT_l) and global mean temperature rise (ΔT_g)
from the baseline period have important implications. The former directly affects crop
growth and yield, and the latter represents a global target associated with the mitigation
activities. We extracted both ΔT_l and ΔT_g from the literature as much as possible, but ΔT_g is
not available in many studies. In such cases, we estimated ΔT_g using the Warming Attribution
Calculator (http://wlcalc.climateanalytics.org/choices). In the dataset, we provide two
estimates for ΔT_g: one from the current baseline period (2001-2010) and the other from the
preindustrial era (1850-1900).

Current temperature level. Current annual mean temperatures were obtained from the
W5E5 dataset[21], which was compiled to support the bias adjustment of climate input data
for the impact assessments performed in Phase 3b of the Inter-Sectoral Impact Model
Intercomparison Project (ISIMIP3b, https://www.isimip.org/protocol/3/). The W5E5 dataset
includes half-degree grid resolution annual mean temperature data, which we extracted for
each simulation site or region using the geographic information.

CO_2 concentrations. Several studies report two series of yield simulations obtained using
two CO_2 levels to infer the CO_2 fertilization effects: one obtained with CO_2 concentrations
fixed at the current levels and the other obtained with increased future CO_2 concentrations
provided by the emission scenario considered. In the dataset, we make this explicit in the
following two variables:
1. CO_2: Binary variable equal to "Yes" if future CO_2 concentrations from the emission
scenarios were used and "No" if the current CO_2 concentration was used for the
yield simulations.
2. CO_2 ppm: if available, CO_2 concentration was extracted from the original paper. If
not, we calculated it from projected changes in CO_2 concentrations based on the
scenarios and periods studied. CO_2 concentration data were obtained from
https://www.ipcc-data.org/observ/ddc_co2.html for CMIP3 and Meinshausen, et al.
[16] (http://www.pik-potsdam.de/~mmalte/rcps/) for CMIP5.

Baseline correction. Because baseline periods differed between studies, we corrected YI, ΔT_l,
and ΔT_g to the 2001-2010 baseline period by a linear interpolation method following
Aggarwal et al[22]. Namely, the impacts YI were first divided by the year gap between the
future period midpoint year and the baseline period midpoint year of the original study. The
impact per year was then multiplied by the year gap from our reference baseline period
midpoint year (2005). The same method was applied to express ΔT_l relatively to 2001-2010.
We also retain original YI and

226 Data Records

Data availability. All the data are stored in the figshare repository, which are freely available
at https://doi.org/10.6084/m9.figshare.14691579.v1, where the following files are uploaded:

1. "Projected_impacts_datasheet.xlsx" includes the final dataset after screening.
2. "Meta-data.xlsx" contains the summary of the dataset, such as the definition and
unit of the variables used in "Projected_impacts_datatasheet.xlsx".
3. "Online_only_summary_table.xlsx" contains data distribution, median, and mean
impacts of climate change, presented in the online-only tables.
4. "Supplementary_materials.pdf" contains methods for estimating local temperature
rise and summary distribution of climate change impacts on four crop yields.
5. "References.docx" provides a list of references that provided data.

Coverage of the data. A total of 8252 yield simulations are registered in the consolidated
dataset. The number of simulations grows exponentially with publication year: 18 in the
1980s, 179 in 1990s, 724 in 2000s and 7331 in 2010s (Online-only Table 1).
About 83% of the simulations use CMIP5 climate scenarios, and 10% use CMIP3. From
CMIP5, RCP2.6, RCP4.5 and RCP8.5 are the most used concentration pathways (Online-only
Table 2, Fig. 2a). ΔT_g from the baseline period (2001-2010) ranges from 0 to 4.8 °C (0.8 to 5.6
°C from the preindustrial period). Almost all simulations with ΔT_g > 3°C use RCP8.5, resulting
in a greater ΔT_g range under CMIP5 (RCPs) than under previous scenarios (SRES and others).

Projected time periods span widely in the 21[st] century, but the midpoint years peak at 2020
for the near future, 2050 for mid-century, and 2080 for end-century (Fig.2b). Major emission
scenarios such as RCP2.6, 4.5 and 8.5 are almost equally distributed across time periods.
About 8 % of the simulations assume no CO_2 fertilisation effects.

Relative frequency of the regions studied generally reflects harvested areas of the four crops
in each region (Fig. 2c). About 41% of the simulations were performed in Asia, which
accounts for about 47% of the harvested area of the four major crops (mean of 2017-2019,
FAOSTAT, http://www.fao.org/faostat/en/, accessed on April 28, 2021). Europe is slightly
overstudied for its world share of the harvested areas (12%). Central and South Americas is
slightly under-researched (9%) for the regional share of harvested areas (15%), whereas
Africa's share is comparable to the area harvested (10%). Altogether global harvested areas
for these four major crops amount 7×10^8 ha: wheat represents 31% of this area, followed by
maize (28%), rice (23%) and soybean (18%). Nevertheless, maize studies are over
represented, accounting for about half of the simulations (52%), followed by wheat (26%)
and rice (18%); soybean accounts only for 4% of the simulations (Fig.2c). Regionally, maize
and wheat are harvested across almost all regions, and simulations follow the actual
distribution of these crops. Rice is predominantly studied in Asia, reflecting actual
distribution (85 % of the harvested area is in Asia). Soybean remains understudied compared
to the other three crops despite its large cultivated area (about 75% of the rice harvested

area). Regionally, simulation sites or regions for soybean are mostly in the Americas, which
account for 76% of the total soybean harvested area.
270
About 36 % of the simulations (2992) are performed under current management practices,
and the rest (5260) considered different management or cultivars as adaptation options (Fig.
1d). More than half of the simulations are run with multiple options. Among these options,
fertiliser accounts for 32 % followed by irrigation (29%), cultivar and planting date (17%
each). There are 1853 pairs of yield simulations available for comparing results obtained with
and without adaptation. These pairs of yield data can be used to compute the adaptation
potentials of the different options considered.
278

279
280
Fig. 2. Data availability of crop yield simulations and its breakdown.
(a) By global temperature rise from the preindustrial era and climate scenarios, **(b)** By
projected time periods (midpoint years) and climate scenarios, **(c)** IPCC regions[23] and crop
species, and **(d)** adaptation options and crop species. n is the total number. Note that
n=9687 in adaptation options (d) exceeds the total number of simulations (8314) because we
collectively add all the options used in the simulations, including those that use multiple
options. n is the number of simulation results.

288
Technical Validation
Data quality check. We repeatedly checked the data with multiple authors for the new
dataset. For the Aggarwal-DS, we reviewed the sources of references. In case of missing
information such as climate scenarios, CO_2 concentration, or temperature increase, we
came back to the original reference. Inconsistencies between the dataset and original papers
were corrected when possible. Overall, corrections were made on 163 simulations from 5
studies, which we flag with "*" in the remark column of the dataset. We removed all data of
the Aggarwal-DS that were untraceable in the original paper. This quality control excluded 47
simulations from 9 articles listed in the "Excluded" sheet.
298
We first examined the distribution of the climate change impacts on crop yields, which span
from −100 to 136% (Fig. 3). This distribution is skewed to the left, as indicated by the
negative skewness. The kurtosis shows that the distribution is salient around the mean
compared to the normal distribution. We tested the effects of potential outliers outside the
1.5-fold interquartile range (IQR) on the summary statistics of the climate change impacts on
crop yields[24]. Removing values outside the 1.5-fold IQR decreases the number of simulations
by 706 (8.5%) and the negative effects of climate change on crop yields by 3.5% for the mean
and 0.6% for the median, suggesting that the deletion affected the original distribution. We,
therefore, keep all the simulation results in the dataset.
308

variables	With_Adapt	Without_Adapt
Min	-100.0	-100.0
Max	135.9	89.7
mean	-4.6	-12.0
q1	-9.3	-20.0
median	-1.6	-6.4
q3	4.0	1.6
sd	19.7	24.3
skewness	-1.5	-1.1
kurtosis	8.8	5.3

309
310 **Fig. 3.** Distribution of climate change impacts on crop yields with and without adaptation
311
Methods to estimate local temperature changes. Local temperature change (ΔT_l) was not
available for about half of the simulations. To estimate ΔT_l for these studies, we examined
the relationship between ΔT_l and the following six input variables in 4143 simulations where
ΔT_l was available: ΔT_g, CO_2 concentrations, latitudes, longitudes, time periods, and emission
scenarios. Values of ΔT_l were estimated using random forest algorithms trained to establish a
function relating local temperature rise to the six inputs considered. We tested and
compared six models based on different combinations of the input variables. The full model
explains about 97.5% of the variation in ΔT_l, but the reduced model with three variables (ΔT_g,
latitude, and longitude) was close in term of percentage of explained variance (97.0%), and
led to a cross-validated RMSE as low as 0.18 °C (Supplementary Table S1 and Fig. S2). We,
therefore, used the reduced model to impute ΔT_l for the 4110 missing data.

323
Comparison with previous studies. The overall effects of climate change on crop yields are
negative, with the mean and median of −12% and −6.7% without adaptation and −4.6% and
−1.6% with adaptation, respectively (Online-only Table 3 and 4). The median per-decade
yield impact without adaptation is −2.3% for maize, −3.3% for soybean, −0.7% for rice, and
−1.3% for wheat (Table 1), which are consistent with previous IPCC assessments[14]. The
median per-warming-degree impact is −7.6% for maize, −11.7% for soybean, −2.2% for rice,
and −3.8% for wheat (Table 2). Per-degree yield impacts for each crop are generally within
the range reported in the previous meta-analysis[11]. Among the four crops, soybean has the
least number of simulations, resulting in a greater variation in both per-decade and per-
degree impacts. Maize consistently shows the largest negative impacts, while rice shows the
least.
335
336 Table 1. Summary statistics of climate change impacts on four major crops expresses as per
337 decade impact and per degree impact without adaptation.
338

	Per decade impact (% decade^{-1})				Per degree impact (% °C^{-1})			
	Maize	Rice	Soybean	Wheat	Maize	Rice	Soybean	Wheat
Minimum	-40.0	-40.8	-30.0	-35.4	-158.7	-71.7	-112.7	-122.7
Maximum	14.2	26.2	13.8	21.2	47.4	118.9	40.4	64.5
Mean	-4.4	-1.3	-3.9	-2.1	-15.5	-2.5	-13.2	-6.7
1st quartile	-6.5	-2.4	-7.7	-3.8	-20.1	-6.8	-21.9	-11.2
Median	-2.2	-0.7	-3.3	-1.2	-7.3	-2.2	-11.7	-3.6
3rd quartile	-0.3	0.8	-0.2	0.5	-1.2	2.9	-0.5	1.8
Standard deviation	7.4	4.7	6.9	5.2	26.3	11.9	24.0	17.2
Skewness	-1.7	-2.5	-0.6	-1.7	-2.0	0.9	-1.1	-2.2
Kurtosis	7.7	21.7	4.9	10.6	7.9	23.7	5.6	13.1

339
340
341 The climate change impacts by IPCC regional groups[23] reveals that Europe and North America
342 are expected to be less affected by climate change in the mid-century (MC) and the end-
343 century (EC) than Africa, Central and South America, particularly for maize and soybean.
344 Both positive and negative effects are mixed in all regions (Fig. 4, Supplementary Fig. S2, Fig.
345 S3).
346

349 **Fig. 4** Climate change impacts (% of yield change) on four crops without adaptation under

350 RCP8.5. Upper two panels, Mid-century; Lower two panels, End-Century. Maps with bluish

351 symbols, positive effects (yield gain); Maps with reddish symbols, negative effects (yield loss).

352 Projections under RCP2.6 and RCP4.5 are given in Supplementary Fig. S3.

(a)

(c)

Fig.5 Projected yield changes relative to the baseline period (2001-2005).

(a) Mid-century (MC) projections without adaptation under RCP8.5 scenario, upper panels

showing positive impacts and lower panels negative impacts, **(b)** End-century (EC)

projections under three RCP scenarios by current annual temperature (T_{ave}), and **(c)** Yield

change as a function of global temperature rise from the pre-industrial period by three T_{ave}

levels. The box is the interquartile range (IQR) and the middle line in the box represents the

median. The upper- and lower-end of whiskers are median 1.5 × IQR ± median. Open circles

are values outside the 1.5 × IQR.

Regional differences in the impacts in MC and EC are associated with the current
temperature level. In MC, positive or neutral effects are projected when current annual
average temperatures (T_{ave}) are below 10-15 °C, but the effects become negative as T_{ave}
increases beyond these levels regardless of RCPs (Fig. 5a). This accounts for the regional
differences as a function of latitude reported in previous meta-analyses[4,5]. In EC, the
threshold T_{ave} shifts lower, and the negative effects become more severe, particularly under
a high emission scenario (RCP8.5) (Fig. 5b). The effect of ΔT_g from the baseline period on YI
differs depending on the T_{ave} (Fig. 5c); At $T_{ave}<10°C$, YI is generally neutral even where ΔT_g
$>2°C$ in most crops, but at $T_{ave}>20°C$, YI is negative even with small ΔT_g, notably in maize. The
difference in the YI dependence on ΔT_g between regions is also consistent with the previous
study[4].
376
377 **(a)** Adaptation options

395 **(b)** Regions

415 **Fig. 6** Adaptation potential, defined as the difference between yield impacts with and
416 without adaptation in projected yield impacts, for three RCPs by mid- and end-century (MC,

417 EC). The box is the interquartile range (IQR) and the middle line in the box represents the
418 median. The upper- and lower-end of whiskers are median 1.5 × IQR ± median. Open circles
419 are values outside the 1.5 × IQR. **(a)** By adaptation options and **(b)** by IPCC regions.
420
Adaptation potential, defined as the difference between yield impacts with and without
adaptation, is an important metric to measure our capacity and limit to adapt to climate
change. The options tested with simulation models show an average effect of 8% in MC and
11% in EC (Fig. 6, Supplementary Fig.S4), which is not sufficient to offset the negative
impacts, particularly in currently warmer regions. A residual damages will thus likely remain
even with adaptation, underpinning several lines of evidence[25,26].
427
428
429 Usage Notes

Crop yield simulation studies can provide a narrative of when, where, and what will happen
to crop production under different GHG emissions and climate scenarios. They are also
expected to provide quantitative information on the potential and limits to adaptation.
However, robust estimates covering different temporal and spatial scales need to draw on
multiple results obtained from various simulation studies. Nearly four decades have passed
since the model projections based on future climate scenarios started. This dataset
covers the entire period of simulation studies using climate scenarios, which can help update
the quantitative review of climate change impacts on crops. The full list of references is
provided in the reference file (https://doi.org/10.6084/m9.figshare.14691579.v1).

Currently, studies are heavily biased towards major cereals such as maize, rice, and wheat,
but this can be expanded to include other crops. As of 2020, our literature search failed to
find published reports using CMIP6 climate scenarios, but this dataset can be easily updated
when new simulations using new climate scenarios or other crop species become available.
The next IPCC assessment cycle can fully utilise this dataset by adding the latest simulation
results.

One of the caveats to the current dataset is that it only includes crop yield data,
notwithstanding crop simulation studies are expected to produce other results than yield.
Because of the recent progress in crop modelling, grain quality projections are emerging[27].
We have extensively included the temperature levels to account for the impacts concerning
the warming and current temperature, but there is a need to include other key climatic
variables such as precipitation and soil moisture. It will be useful to expand our dataset in
the future to include this type of data.

452 Code Availability

453 A script file was created using the R statistical programming to estimate ΔT_i and is available
454 from the corresponding author upon request.
455
456 Competing interests

457 The authors declare no competing interests.
458
459
460 References

1   Davis, K. F., Downs, S. & Gephart, J. A. Towards food supply chain resilience to
environmental shocks. *Nature Food* **2**, 54-65, doi:10.1038/s43016-020-00196-3
(2021).

2   Parry, M. L. *et al.* in *The IPCC Impacts Assessment* (eds W.J.M Tegart, G.W. Sheldon,
& D.C. Griffiths) 1-45 (AGPS Press, Canberra, Australia, 1990).
3   Porter, J. R. *et al.* Invited review: Intergovernmental Panel on Climate Change,
agriculture, and food—A case of shifting cultivation and history. *Global Change*
Biology **25**, 2518-2529, doi:10.1111/gcb.14700 (2019).
4   Challinor, A. J. *et al.* A meta-analysis of crop yield under climate change and
adaptation. *Nature Climate Change* **4**, 287-291, doi:10.1038/nclimate2153 (2014).
5   Aggarwal, P., Vyas, S., Thornton, P. & Campbell, B. M. How much does climate
change add to the challenge of feeding the planet this century? *Environmental*
Research Letters **14**, doi:10.1088/1748-9326/aafa3e (2019).
6   Rosenzweig, C. *et al.* The Agricultural Model Intercomparison and Improvement
Project (AgMIP): Protocols and pilot studies. *Agricultural and Forest Meteorology*
170, 166-182, doi:10.1016/j.agrformet.2012.09.011 (2013).
7   Asseng, S. *et al.* Uncertainty in simulating wheat yields under climate change. *Nature*
climate change **3**, 827-832 (2013).
8   Li, T. *et al.* Uncertainties in predicting rice yield by current crop models under a wide
range of climatic conditions. *Global Change Biology* **21**, 1328-1341,
doi:https://doi.org/10.1111/gcb.12758 (2015).
9   Bassu, S. *et al.* How do various maize crop models vary in their responses to climate
change factors? *Global Change Biology* **20**, 2301-2320,
doi:https://doi.org/10.1111/gcb.12520 (2014).
10   Wallach, D. *et al.* Multimodel ensembles improve predictions of crop–environment–
management interactions. *Global Change Biology*, 5072-5083,
doi:10.1111/gcb.14411 (2018).
11   Zhao, C. *et al.* Temperature increase reduces global yields of major crops in four
independent estimates. *Proceedings of the National Academy of Sciences* **114**, 9326-
9331, doi:10.1073/pnas.1701762114 (2017).
12   Makowski, D., Marajo-Petitzon, E., Durand, J.-l. & Ben-Ari, T. Quantitative synthesis
of temperature, CO2, rainfall, and adaptation effects on global crop yields. *European*
Journal of Agronomy **115**, 126041-126041, doi:10.1016/j.eja.2020.126041 (2020).
13   Villoria, N. B. *et al.* Rapid aggregation of global gridded crop model outputs to
facilitate cross-disciplinary analysis of climate change impacts in agriculture.
Environmental Modelling & Software **75**, 193-201,
doi:10.1016/j.envsoft.2015.10.016 (2016).
14   Porter, J. R. *et al.* in *Climate Change 2014: Impacts, Adaptation, and Vulnerability.*
Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth
Assessment Repor+ (eds C B Field *et al.*) 485-533 (Cambridge University Press,
Cambridge, United Kingdom and New York, NY, USA, 2014).
15   van Vuuren, D. P. *et al.* The representative concentration pathways: an overview.
Climatic Change **109**, 5, doi:10.1007/s10584-011-0148-z (2011).
16   Meinshausen, M. *et al.* The RCP greenhouse gas concentrations and their extensions
from 1765 to 2300. *Climatic Change* **109**, 213-241, doi:10.1007/s10584-011-0156-z
(2011).
17   Taylor, K. E., Stouffer, R. J. & Meehl, G. A. An Overview of CMIP5 and the Experiment
Design. *Bulletin of the American Meteorological Society* **93**, 485-498,
doi:10.1175/bams-d-11-00094.1 (2012).
18   Eyring, V. *et al.* Overview of the Coupled Model Intercomparison Project Phase 6
(CMIP6) experimental design and organization. *Geosci. Model Dev.* **9**, 1937-1958,
doi:10.5194/gmd-9-1937-2016 (2016).
19   Ishigooka, Y., Hasegawa, T., Kuwagata, T. & Nishimori, M. Evaluation of the most
appropriate spatial resolution of input data for assessing the impact of climate

change on rice productivity in Japan. *Journal of Agricultural Meteorology* **76**, 61--68,
doi:10.2480/agrmet.D-19-00021 (2020).
20   Iizumi, T. *et al.* Responses of crop yield growth to global temperature and
socioeconomic changes. *Scientific Reports* **7**, 7800-7800, doi:10.1038/s41598-017-
08214-4 (2017).
21   Lange, S. NWFDE5 over land merged with ERA5 over the ocean (W5E5). V. 1.0. *GFZ*
Data Services, doi:10.5880/pik.2019.023 (2019).
22   Paudyal, B. R., Chanana, N., Khatri-Chhetri, A., Sherpa, L., Kadariya, I., & Aggarwal, P.
Gender integration in climate change and agricultural policies: The case of Nepal.
Frontiers in Sustainable Food Systems **3**, doi:10.3389/fsufs.2019.00066 (2019).
23   Hewitson, B. *et al.* in *Climate Change 2014: Impacts, Adaptation, and Vulnerability.*
Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment
Report of the Intergovernmental Panel on Climate Change   1133-1197 (Cambridge
University Press, Cambrige, UK, 2014).
24   Upton, G. & Cook, I. *Understanding Statistics.* (Oxford University Press, 1997).
25   Minoli, S. *et al.* Global Response Patterns of Major Rainfed Crops to Adaptation by
Maintaining Current Growing Periods and Irrigation. *Earth's Future* **7**, 1464-1480,
doi:10.1029/2018EF001130 (2019).
26   Iizumi, T. *et al.* Climate change adaptation cost and residual damage to global crop
production. *Climate Research* **80**, 203-218, doi:10.3354/cr01605 (2020).
27   Asseng, S. *et al.* Climate change impact and adaptation for wheat protein. *Global*
Change Biology **25**, 155-173, doi:10.1111/GCB.14481 (2018).
537

Acknowledgements

539 This study was performed by the Environment Research and Technology Development Fund
540 (JPMEERF20S11820) of the Environmental Restoration and Conservation Agency of Japan. TH
541 and DM would like to thank Joint-Linkage-Call between INRAE and NARO for supporting this
542 collaborative study. We also thank Dr. T. Iizumi and Y. Ishigooka for providing the aggregated
543 simulation results.

Author contributions

Toshihiro Hasegawa and Hitomi Wakatsuki designed the dataset.
Hitomi Wakatsuki and Hui Ju collected simulation results from the SCOPUS search.
Shalika Vyas designed and collected the Aggarwal dataset.
Gerald C. Nelson conducted literature search and provided global temperature dataset.
David Makowski and Hitomi Wakatsuki developed a statistical imputation for missing data on
the local temperature rise.
All authors worked on data analysis and drafting the final version of the manuscript.

Additional Formatting Information

555 Correspondence and requests for materials should be addressed to TH.