1 # Expression of putative effectors of different *Xylella fastidiosa*
2 # subspecies/strains reveals recognition and defense activation
3 # in various model plants

4 *Matthaios Sertedakis[1§], Konstantinos Kotsaridis[1§], Dimitra Tsakiri[1§], Ana Dominguez-*
5 *Ferreras[3], Vardis Ntoukakis[3], Panagiotis F. Sarris[1,2,4]**
6
7 [1] Department of Biology, University of Crete, 714 09 Heraklion, Crete, Greece,
8 [2] Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology-Hellas,
9 Heraklion, Crete, Greece,
10 [3] School of Life Sciences, University of Warwick, Coventry CV4 7AL, United Kingdom,
11 [4] Biosciences, University of Exeter, Exeter, United Kingdom.
12
13
14
15 [§] Equal contribution
16
Corresponding Author:
Professor Panagiotis F. Sarris, p.sarris@imbb.forth.gr; p.sarris@uoc.gr

SUMMARY

The re-emergence of Gram-negative bacterium *Xylella fastidiosa* in Europe in 2013
impelled the scientific community to discover novel strategies for crop protection. The
wide host range of *Xylella* indicates the existence of yet not characterized pathogenic
mechanisms to overcome plant defenses. The recent uprising accuracy of a variety of
bioinformatics tools, with the ability to predict the function of putative microbial protein
represent a useful approach for understanding which of these proteins are associated
with pathogens virulence. In this study we collected a number of putative effectors from
two *X. fastidiosa* strains: Temecula1 and CoDiRo and the subspecies (ssp.) Sandyi
Ann-1. We designed an *in-planta Agrobacterium* based expression system that drives
the expressed proteins to the cell apoplast, in order to investigate their ability to
activate defense in various model plants. Furthermore, we organized the resulted
proteins according to their sequential and structural similarities via the I-TASSER
online tool. We identified that various *X. fastidiosa* proteins were able to differentially
elicit cell death-like phenotypes in *Nicotiana tabacum, N. sylvestris* and *N.*
benthamiana. These proteins are members of different enzymatic groups: a)
hydrolases/hydrolases inhibitors, b) serine proteases and c) metal transferases.
Collectively, we identified structurally similar proteins that were able to differentially
elicit cell death-like phenotypes in different cultivars of the same species. Our findings
provide the bases for further studies on the mechanisms that underlie host-defense
activation by *X. fastidiosa* putative effectors, as well as, pathogens adaptation in
susceptible hosts.

INTRODUCTION

Plants respond to invading pathogens by exploiting their innate immunity system. microbe-associated molecular patterns (MAMP)-triggered immunity (MTI) and effectors triggered immunity (ETI) have been described as the main two layers of defense during the infection of a host (Jones, J.D.G., 2006; Duxbury et al., 2016; Mermigka et al., 2019; Cui et al., 2015). Recent studies have proposed a revised version of the zig-zag model of plant innate immunity introduced by Jones, J.D.G., 2006 (Jones & Dangl, 2006); strongly indicating the existence of a crosstalk between MTI and ETI. ETI potentiates MTI immune responses and vice versa (Katagiri & Tsuda, 2010; Ngou et al., 2021). MTI is responsible for the detection of pathogen/microbe associated molecular patterns (P/MAMPs) and/or danger associated molecular patterns (DAMPs), via specific cell surface-localized pattern-recognition receptors (PRRs) (Couto & Zipfel, 2016). MTI shields the plant cell from various pathogen-derived molecules or by recognizing self-made elicitors (Malukani et al., 2020) and triggering downstream signaling events to activate defense responses (Schwessinger & Zipfel, 2008; Apama et al., 2009).

MTI defensive outcomes include production of reactive oxygen species (ROS), calcium influx, activation of mitogen-activated protein kinases (MAPKs), chromatin remodeling, differential regulation of gene expression and callose deposition. Collectively, these responses restrict pathogens on the site of the infection and prevent disease development (Lu et al., 2015; Mur et al., 2008; Pardal et al., 2021; Stotz et al., 2014).

In order to reach their nutritional needs and proliferate effectively inside the host, several pathogens have evolved to secrete virulence factors - known as effectors – directly into the host-cell cytoplasm or into the extremely hostile apoplastic space. In several cases, interaction of apoplastic pathogen effectors with plant PRRs has been associated with induction of immune responses and development of Programmed Cell Death (PCD) (van der Burgh & Joosten, 2019). For instance, the apoplastic effectors Avr2 and Avr4 from the fungal pathogen *Cladosporium fulvum* are recognized by the receptor like proteins (RLPs) *Cf-2* and *Cf-4* respectively in *Solanum lycopersicum* and trigger a strong defense response, including PCD (Song et al., 2009; Postma et al., 2016; Ilyas et al., 2015; Kourelis & Van Der Hoorn, 2018). Similarly, the apoplastic effectors Chp-7 and ChpG of *Clavibacter michiganensis*, elicit PCD when they are secreted to the apoplast, but not when they are expressed in the host-cell cytoplasm (Lu et al., 2015). There are additional virulent extracellular effectors related to PCD phenotypes in plants, however the exact mechanisms of innate immunity underlying their perception remain deeply uncharacterized (Nissinen et al., 2009; Lu et al., 2015).

X. fastidiosa (*Xf*) was first described as the causal agent of "Pierce's disease" (PD) in

grapes and it is an extremely dangerous plant pathogenic bacterium worldwide

(Mollenhauer & Hopkins, 1974; Hopkins & Purcell, 2002; Chatterjee et al., 2008). *Xf* is

a gram negative, slow growing and strictly aerobic bacterium that has been a subject

of interest due to its economic impact. *Xf* has an extremely extended host range which

consists of more than 300 plant species (Baldi & La Porta, 2017), including *Nerium*

oleander, *Olea europaea* and *Vitis vinifera* species (Schneider et al., 2020; Food &

Authority, 2018; Huang et al., 2020). While there are emerging studies assessing the

life style of this pathogen, host specificity and colonization strategies, less progress

has been accomplished in the field of the molecular host-pathogen interactions.

Similarly, the individual role(s) of putative virulence proteins secreted by *Xf*, in order to

subvert host's immune machinery and how this leads to disease development and

finally plant death, is still poorly understood (Roper et al., 2019; Rapicavoli et al., 2018;

Chatterjee et al., 2008; Zhang et al., 2015; Nascimento et al., 2016; Gouran et al.,

2016).

Xf lacks a Type III translocation system (T3SS); the common bacterial transporter of

virulence factors from the pathogen's cytosol directly into the host's intracellular

environment. However, *Xf* possesses Type I, II, IV and V secretion systems (Simpson

et al., 2000; Van Sluys et al., 2003). The *Xf* 12- protein Type II secretion system (T2SS)

with origins to its close relatives of the *Xanthomonas* group, considerably acts as the

main source of its pathogenicity (Rapicavoli et al., 2018). Proteases and cell wall

degrading enzymes (CWDEs) are often secreted by T2SS, while mutations on

essential components of the secretion mechanism, usually lead to avirulent

phenotypes (Rapicavoli et al., 2018).

In this study, using the KEGG database, we searched for homologues of various

known type II effector genes of pathogenic microorganisms that could be present in

several *Xf* sequenced genomes (**Suppl. Table S1**). This process resulted in selection

of nineteen putative *Xf* type II effectors originating from two strains and one spp. for

further study (**Suppl. Table S1**).

Gene evolution is a process that involves mechanisms such as gene duplications and

horizontal gene transfers, which resulted in the hypothesis that sequence unrelated

genes may have high similarity in their tertiary folding and furthermore have the same

function in pathogen virulence (de Guillen et al., 2015; Andrie et al., 2008). Based on

this hypothesis and using I-TASSER online server, we compared the protein with the

highest sequence similarity with the one, which was used as a template for the

predicted structures of the selected *Xf* proteins (Roy et al., 2010; Yang et al., 2015;
Yang & Zhang, 2015) (**Fig. 1** & **Suppl. Fig. S1**).

114
115
116

MATERIALS & METHODS

Plant material

120 Three wild type (WT) *Nicotiana* species were used for our study; *N. tabacum* cultivars
121 N34/4, Xanthi, Petit Gerard, *N. sylvestris* ecotypes ITB626, NIC6, A34750352,
122 A04750326, NS25 and TW136 and *Nicotiana benthamiana*. All plants were grown
123 under greenhouse conditions, at 23°C and under a 16-hour photoperiod.

Bacterial strains

Escherichia coli (strains Stellar and DH10b) were routinely grown on LB medium with
the appropriate antibiotics and incubation at 37°C for 16 hours (h). Liquid cultures were
grown at 37°C for 16 h applying shaking at 200 rpm.

Agrobacterium tumefaciens (strains AGL1, C58C1 and GV3101) were grown on LB
medium plates with selective antibiotics and incubation at 28°C for two days. Liquid
cultures were grown at 28°C for 24h while shaking at 200 rpm.

Available genomic data banks for three strains of *Xylella fastidiosa* (strains Temecula1,
CoDiRo and Sandyi ann-1) allowed DNA synthesis of effector genes.

Cloning & Constructs

All 19 effectors of *Xylella fastidiosa* (originating from Temecula1, CoDiRo, Sandyi ann-
1 strains) were synthesized and introduced in plasmid vector pICH41308, kindly
provided by Dr. Vardis Ntoukakis (Department of Life Sciences, University of Warwick).
Effector genes along with a 35S promoter, fused to the coding sequence of secretion
peptide of tobacco PR1a (Pathogenesis-Related protein 1a) and NOS terminator were
transferred in binary vector pICH86966 and cloned using Golden Gate cloning (Engler
et al., 2008). pMDC:spC7HPB construct, kindly provided by Professor Jane
Glazebrook (Department of Plant Biology, University of Minnesota), was introduced in
Agrobacterium tumefaciens AGL1 and was used in this study as a positive marker of
apoplastic HR-like cell for *in planta* assays. pBluescript::PR1a_sp,
pICH86988::XopQ:YFP and pICH86988::GUS:YFP plasmids that were used in this

work had already been constructed for a previous study in our lab (Michalopoulou et
al., 2020).

Agrobacterium-mediated transient expression *in planta*

Transformed *Agrobacterium* cells were grown on LB agar plates with selective
antibiotics for 2 days. Single colonies were used to inoculate LB liquid medium
containing selective antibiotics and were cultured for 24h. Cells were then harvested
by centrifugation, washed with 5 ml of 10mM $MgCl_2$ and re-suspended in 1 ml MM
solution (10mM $MgCl_2$, 10mM MES [pH 5.6]). OD_{600} was adjusted to 0.5 with MM and
the bacteria were then used for infiltration of plant leaves with a blunt end syringe. Six
week old plants were challenged with Agrobacterium, leading to *in planta* transient
expression of *Xf* effectors, as well as positive and negative markers for identification
of HR-like Programmed Cell Death (PCD). Infiltrated plants were incubated under
greenhouse conditions (23°C, 16-hour photoperiod) and HR-like cell death was
assessed 4dpi.

In silico structural predictions of proteins

Structural predictions were made using the I-TASSER online server. Comparisons
were performed between the protein with the highest sequence similarity with the one
used as a template for the predicted structures of the selected *Xf* proteins (Roy et al.,
2010; Yang et al., 2015; Yang & Zhang, 2015). The predicted structure with TM-score
>5 was considered reliable. Visualization of the structures was carried out through
Pymol v2.3.1 (Schrodinger & DeLano., 2020).

RESULTS & DISCUSSION

To test whether these proteins can elicit PCD after their delivery into the plant apoplast,
we first synthesized the corresponding genes including silent mutations where needed
to domesticate the sequences making them compatible with the Golden Gate system.
Then, we cloned the synthesized genes of interest in an *Agrobacterium*-mediated
transient expression system (**Fig. 2a**). The gene expression in this system was under
the transcriptional regulation of the constitutive CaMV 35S promoter. To ensure
secretion of the protein into the apoplast we fused the selected *Xf* proteins to the
secretion peptide of tobacco PR1a (Pathogenesis-Related protein 1a) (Lu et al., 2015).
The PR1a secretion peptide is cleaved upon secretion to the apoplast (Lu et al., 2015).
We generated 19 such constructs to screen for PCD symptoms by the selected *Xf*
proteins. For our screening we used three distinct *Nicotiana tabacum* cultivars (N34'4,

Petit Gerard, Xanthi); the *N. benthamiana* and six *N. sylvestris* ecotypes. As a positive

control we used the *Clavibacter michiganensis* apoplastic effector Chp7 that has been

shown to elicit an PCD response upon its secretion to the apoplast by the PR1a

secretion peptide (Lu et al., 2015) (**Fig. 2b** & **Suppl. Fig. S2**). Furthermore, the type

III effector XopQ of *Xanthomonas campestris* pv. *vesicatoria*, served as a secondary

positive control, due to its known ability to elicit Hypersensitive Response (HR) in both

N. tabacum and *N. benthamiana* (Adlung et al., 2016). *Agrobacterium*-mediated

transient expression of the GUS reporter gene in these species did not elicit cell death

and it was used as a negative control for our transient expression assays.

Our screening revealed nine proteins that are known or predicted to be type II-secreted

by *X. fastidiosa* and were able to elicit PCD phenotypes in different *Nicotiana* species

(**Fig. 2b**). Most of these proteins successfully elicited PCD four days' post inoculation

(4 dpi) in all three *N. tabacum* cultivars used and four out of six *N. sylvestris* ecotypes.

Interestingly, all nine effectors tested for induction of PCD displayed divergence in

distinct *N. sylvestris* ecotypes, while the protein encoded by D934_09300 was able to

elicit cell death in *N. tabacum* cv. "N34'4" and cv. "Xanthi" leaves but not in *N. tabacum*

cv. "Petit Gerard", indicating a form of specificity in this response. Moreover, signs of

cell death were entirely missing from *N. benthamiana* leaves suggesting that the

responses observed in *N. sylvestris* and *N. tabacum* are most likely not a result of

cytotoxic effects. Apart from these nine putative effectors, ten more proteins were

studied but did not elicit PCD in any plant species/cultivar tested in this study (**Suppl.**

Fig. S1).

Cell death phenotypes varied in severity and seemed to develop at different rates. In

order to comprehensively evaluate our results, we first assigned a cell death score to

each observed cell death phenotype, based on its intensity (**Suppl. Fig. S2**).

We also reviewed the frequency of a certain score regarding both the studied protein

and the plant cultivar used in each experiment (**Suppl. Fig. S3**). The necrotic

phenotype observed in *N. sylvestris* and *N. tabacum* cv. "N34'4" plants was typically

more pronounced compared to that of tobacco cultivars "Petit Gerard" and "Xanthi".

For this, we have to consider potential differences in the transformation efficiency of

distinct cultivars under the transient expression system we applied. However, these

results could also indicate that the same protein may elicit PCD of varying intensity

when introduced to different *Nicotiana* relatives or cultivars of the same plant species

and hint that the potency of this type of response is possibly host-dependent.

The *in silico* structural prediction presented here, indicates that all three proteins
encoded by PD_0956, RA12_05570 and D934_07885, which successfully elicited
PCD in all studied tobacco cultivars and in *N. sylvestris* ecotype A_34750352, have a
high structural similarity with hydrolases (**Fig.1**; **Table 1**). Hydrolases form a big distinct
enzyme class that includes enzymes which act as biochemical catalysts by using water
molecules to break chemical bonds. This class contains enzymes classified as: a)
lipases; b) phosphatases; c) glycosidases; d) peptidases; and e) nucleosidases.
Specifically, serine proteases/endopeptidases/hydrolases are enzymes where the
nucleophilic serine residue in their active center is used for the hydrolysis of their
substrates (Simon & Cravatt, 2010). Hydrolases group includes proteases that are
secreted by various pathogens having a wide range of functions in virulence. They also
constitute an important group of *X. fastidiosa* including cell wall-degrading enzymes
(CWDEs) (Nascimento et al., 2016). CWDEs presence in the apoplast can trigger
immunity responses, mostly through a modified "self" recognition of degradation
products of these enzymes by plant PRRs (van der Burgh & Joosten, 2019). Similarly,
serine proteases delivered by pathogens into the apoplast have been shown to activate
PCD (Lu et al., 2015). Provided that PD_0956, RA12_05570 and D934_07885
proteins' putative enzymatic activity is valid, their ability to elicit PCD could be
considered a DAMP-recognition event. Serine proteases are also present in large
families of plant extracellular proteins that are often involved in signaling pathways
associated with pathogen resistance (Hou et al., 2019). Therefore, manipulation of
such pathways by bacterial proteases could be a virulence strategy.

Another prominent group of *Xf* proteins is that of PD_1703, RA12_01530 and
D934_08750 that all trigger PCD in three tobacco cultivars and two *N. sylvestris*
ecotypes TW_136, NS_25 but not to ITB_626 (**Fig. 2b**) (Zhang et al., 2015). According
to our structural analysis (**Table 1**), the last two proteins revealed similarities to LipA,
a known *Xanthomonas oryzae* pv. *oryzae* cell wall degrading enzyme (CWDE). While,
for the PD_1703, even if it was previously characterized as a LipA-like protein
(Nascimento et al., 2016), according to our I-TASSER structural prediction, it revealed
similarities to Hydrolase/Serine proteases (**Table 1**). However, hydrolases class is one
of the largest and most diverse enzyme families which among others includes
proteases and lipases, so, this might be a misannotation of the particular protein
database.
LipA homologues are present in all sequenced Xanthomonads and are predicted
lipases, although LipA actually exhibits esterase activity (Apama et al., 2009) (**Fig. 3**).
LipA is known to elicit immune responses in rice and recent findings point to the

possible involvement of a rice Wall-Associated Kinase (WAK) *OsWAKL21.2* in LipA
recognition (Jha et al., 2007; Malukani et al., 2020). Structural similarities of LipA with
PD_1703, RA12_01530 and D934_08750, could mean that these proteins act and are
recognized in a similar manner (this finding is under further investigation by our group).
Notably, PD_1703 has been shown to elicit PCD in grapevine a known *Xf* 'Temecula-
1' host. However, PD_1703 was found to be vital for *Xf* virulence in grapevines,
suggesting that other virulence components of the pathogen could potentially suppress
the PCD induction (Nascimento et al., 2016).

In this study we also focus on three *Xf* proteins encoded by PD_0915, D934_09265
and D934_09300 that were found to elicit apoplastic PCD in tobacco and one *N.
sylvestris* ecotype (**Fig. 2b**; **Table 1**). Structural analysis revealed that this group
consists of proteins with sequence and structural similarity to "*Zonula occludens*" toxins
or "Zot proteins", although PD_0915 was predicted to be more confidently similar to a
metal-transferase. The "Zot" protein was described first in *Vibrio cholera*, where it is
involved in intestinal barrier disturbance, however, "Zot" proteins were identified later
in several other pathogens (Pérez-Reytor et al., 2018, 2020) (**Fig. 3**). Zot proteins have
been associated with high cytotoxicity before (Pérez-Reytor et al., 2018), though this
is not always the case. For instance, in *Vibrio parahaemolyticus,* "Zot" expression did
not positively correlate with cytotoxicity, rather than with an actin disturbance on
infected cells (Pérez-Reytor et al., 2020). *Xf* putative "Zot" proteins studied here appear
not to correlate to cytotoxic effects. Interestingly, the protein encoded by D934_09300
did elicit PCD in the apoplast of *N. tabacum* cv. "N34'4" and cv. "Xanthi", and *N.
sylvestris* ecotype NS_25 but this kind of response was not observed in *N. tabacum*
cv. "Petit Gerard"; or in *N. sylvestris* ecotype ITB_626 and in *N. benthamiana* (**Fig. 2b**).
These data suggest specific recognition of D934_09300 and highlight the complexity
of the plant surveillance system and its possible differentiation among distinct cultivars
of the same species.

Finally, ten putative *Xf* effectors, which were unable to induce necrosis during *in planta*
assays in the selected hosts, were also analyzed for their tertiary structures using I-
TASSER online server and certain predictions could be made for their folding and
function (**Suppl. Fig. S1**; **Table 1**). Notably, LipA-like proteins D934_08755 or
D934_12535, despite their strong correlation with other cell death inducers described
in this study (PD_1703, D934_08750, RA12_01530), were incapable of causing similar
phenotypes when expressed in the apoplast of *Nicotiana* species. This potentially
indicates putative alterations on their active sites that prevent their binding to specific
substrates of plant cell wall.

All the nineteen proteins were used for phylogenetic analysis using homolog proteins
obtained from the KEGG database (**Fig. 3**; **Suppl. Fig. S4**).

Our data, collectively, pinpoint nine proteins belonging to the sparsely studied *X.*
fastidiosa putative "effectorome" that can elicit PCD when transiently expressed and
secreted into the leaf apoplast of different *Nicotiana* species. These proteins are
structurally predicted as putative "CWDEs" or "Zot toxins" that originate from different
X. fastidiosa strains/subspecies. Lack of signs of cytotoxicity, along with the predicted
enzymatic activity of these proteins, hints their possible recognition by the plant innate
immunity system. At least in one case, the protein eliciting the response is a known
required virulence factor of the pathogen, suggesting that it employs other virulence
strategies to suppress immune responses and avoid recognition. The suppression of
immune responses through type III-delivered effector proteins is a common feature
among other members of the *Xanthomonadaceae* family (Jha et al., 2007). However,
since *X. fastidiosa* lacks such a system (Rapicavoli et al., 2018), how this bacterium
avoids recognition by the host's surveillance system, remains to be elucidated. In
summary, our findings, open possibilities and encourage further investigation and
identification of the related PRRs that could be a potential biotechnological tool to
confer broad-spectrum disease resistance against *X. fastidiosa*. Quite recently, the
expression of a PRR receptor in sweet orange has been shown to confer ligand-
dependent activation of defense responses against a citrus infecting strain of *X.*
fastidiosa (Mitre et al., 2021).
306

Acknowledgements

The project was partially supported and D.T. was supported by the "The Vineyard Roads" (project code: 2018ΣΕ01300000) part of the "Emblematic Research Action of National Scope for the exploitation of new technologies in the Agri-food sector, specializing in genomic technologies and pilot application in the value chains of "olive", "grapevine", "honey "and "livestock")» financed by Greek national funds through the Public Investments Program (PIP) of General Secretariat for Research & Technology (GSRT) Duration 2019-2021.
K.K. was supported by the General Secretariat of Research and Innovation (GSRI) through the project "Innovative plant protection technologies for quarantine pathogens of *Xanthomonadaceae* family utilizing tools of Optoacoustic and Molecular Biology - INNOVA-PROTECT", Project code: T1EΔK-01878.

Authors' contributions

P.F.S. designed the research. M.S., K.K, and D.T. performed the research. V.D. and P.F.S. analysed the data. V.D and A.D.F. provided technical support; lab material and tools. M.S., K.K, and D.T. and P.F.S. wrote the paper. All authors have read and approved the manuscript.

FIGURES & TABLES LEGENDS

Table 1. Sequence similar and structural template proteins for the predicted structures

of the nineteen selected *Xylella fastidiosa* putative proteins using I-TASSER.

Figure 1. Predicted model presentation of the selected *Xylella fastidiosa* putative

effectors using I-TASSER online server. The proteins presented here successfully

elicited programed cell death (PCD) in at least one plant cultivar/species tested. The

colors suggest the protein orientation (Blue: N-termini, Red: C-termini). **(a-b)**

Hydrolase/ Esterase (LipA), **(c-f)** Hydrolase and **(g-i)** Zonular Occludens Toxin,

according to their sequence similarities. We used Pymol v2.3.1 to visualize the

structures (Schrodinger & DeLano., 2020).

Figure 2. Putative *Xylella fastidiosa* apoplastic effector proteins elicit programed cell

death (PCD) in *Nicotiana spp.* **(a)** Schematic representation of the cassette that was

cloned in an *Agrobacterium*-mediated plant transient expression system. Genes of

interest were under the control of the constitutive CaMV 35S promoter. Secretion of

the protein into the leaf apoplast was achieved through fusion to the secretion peptide

of tobacco PR1a, which is cleaved during protein secretion (Lu et al., 2015). This figure

was created with BioRender.com. **(b)** Apoplastic effector PR1 sp-Chp7 from

Clavibacter michiganensis along with intracellular acting effector XopQ from

Xanthomonas campestris pv. *vesicatoria* were used as positive Programmed Cell

Death (PCD) - HR markers, while GUS gene served as the negative control for these

assays (Lu et al., 2015; Adlung et al., 2016). *X. fastidiosa* virulence factors which

induced PCD, following overexpression in the plant apoplast, are illustrated above.

PCD occurred 4 days post infiltration under room temperature, in all studied cases.

The assays were repeated at least five times for each putative effector with similar

results.

Figure 3. Phylogenetic trees were constructed for all nineteen putative *Xf* effector

proteins that are presented in this study, which were divided into subgroups based on

their ability to elicit programed cell death (PCD) and on their orthology, according to

KEGG database: **(A-D)** Proteins that induced PCD in this study, with a predicted

orthology of **(A)** lipases, **(B)** peptidases, **(C-D)** Zona Occludens Toxins. These proteins

were correlated with 35 close protein relatives from *Xanthomonas*, *Clavibacter*,

Ralstonia, *Amycolatopsis*, *Pseudarthrobacter*, *Dermatophilus*, *Streptomyces*,

Stenotrophomonas, *Moraxela*, *Azoarcus*, *Collimonas*, *Sulfurimicrobium*,

Chromobacterium genera and viruses *Stenotrophomonas* phage phiSHP2,

Stenotrophomonas phage SMA6. The evolutionary history in each group presented
here was inferred using the Neighbor-Joining method (Saitou & Nei, 1987). The
bootstrap consensus tree inferred from 1500 replicates is taken to represent the
evolutionary history of the different taxa belonging amino-acid sequences as
mentioned before. The evolutionary distances were computed using the Poisson
correction method (Zuckerkandl & Pauling, 1965) and are in the units of the number of
amino acid substitutions per site. Evolutionary analysis was conducted in MEGA X
(Kumar et al., 2018). The abbreviations of microbes and the gene loci used for the
construction of these phylogenetic trees are presented in **Supplementary Table S2**.

SUPPLEMENTARY DATA LEGENDS

Supplementary Table S1. All 19 effector proteins tested for induction of PCD in *Nicotiana* species owe their origins to two pathogenic strains of *Xylella fastidiosa* Temecula1, CoDiRO and ssp. *Sandyi*. Each strain has been associated with infectious diseases in susceptible hosts *Vitis vinifera,* Olive trees and *N. oleander*.

Supplementary Table S2. Bacterial species and gene loci members used for phylogenetic analysis presented in **Figure 3** and **Supplementary Figure S4**.

Supplementary Figure S1. Predicted model presentation of the selected *Xylella fastidiosa* effectors using I-TASSER online server. The proteins presented here did not elicit programed cell death (PCD) in any of the plant cultivar/species tested. The colors suggest the protein orientation (Blue: N-termini, Red: C-termini). **(a-c)** Transport Protein, **(d)** Membrane Protein, **(e-f)** Hydrolase/ Esterase (LipA), **(g)** Cell Adhesion protein, **(h)** Hydrolase, **(i)** Oxidoreductase and **(k)** Gene Regulation. We used Pymol v2.3.1 to visualize the structures. The presented proteins did not induce PCD phenotype in any tested plant cultivars (Schrodinger & DeLano., 2020).

Supplementary Figure S2. *Xylella fastidiosa* virulence factors which did not elicit programed cell death (PCD), following overexpression in the plant apoplast, are illustrated above. Photographs of infiltrated leaves were taken 4 days post infiltration. In all cases, plants were incubated at room temperature. Apoplastic effector PR1 sp-Chp7 from *Clavibacter michiganensis* along with intracellular acting effector XopQ from *Xanthomonas campestris* pv. *vesicatoria* were used as positive Programmed Cell Death (PCD) - HR markers, while GUS gene served as the negative control for these assays (Lu et al., 2015; Adlung et al., 2016). The assays were repeated at least five times for each putative effector with similar results.

Supplementary Figure S3. Cell death intensity score. Cell death score percentage (%) in this diagram is representative of all consistent experimental replicates previously introduced in Figure 2b. Diagram bars are color-coded based on a cell death intensity scale 0-4. Colors in each bar represent the cell death score in percentage out of the total infiltrated panels scored for the nine *Xylella fastidiosa* proteins that elicited programed cell death (PCD) in *Nicotiana spp.*

Supplementary Figure S4. (**A-B**) Proteins that did not induce programed cell death (PCD) in this study, with a predicted orthology of (**A**) lipases and (**B**) haemagglutinins.

These groups of proteins from *Xylella fastidiosa* were correlated with 56 proteins with

origins to bacterial genera of *Pectobacterium*, *Rhodoferax*, *Ralstonia*, *Pseudomonas*,

Stenotrophomonas, *Edwardsiella* and *Lysobacter* (Saitou & Nei, 1987). The

evolutionary history in each group presented here was inferred using the Neighbor-

Joining method (Saitou & Nei, 1987). The bootstrap consensus tree inferred from 1500

replicates is taken to represent the evolutionary history of the different taxa belonging

amino-acid sequences as mentioned before. The evolutionary distances were

computed using the Poisson correction method (Zuckerkandl & Pauling, 1965) and are

in the units of the number of amino acid substitutions per site. Evolutionary analysis

was conducted in MEGA X (Kumar et al., 2018). The abbreviations of microbes and

the gene loci used for the construction of these phylogenetic trees are presented in

Supplementary Table S2.

References

Adlung, N., Prochaska, H., Thieme, S., Banik, A., Blüher, D., John, P., Nagel, O., Schulze, S., Gantner, J., Delker, C., Stuttmann, J., & Bonas, U. (2016). Non-host resistance induced by the Xanthomonas effector XopQ is widespread within the genus Nicotiana and functionally depends on EDS1. *Frontiers in Plant Science*, *7*(NOVEMBER2016). https://doi.org/10.3389/fpls.2016.01796

Andrie, R. M., Schoch, C. L., Hedges, R., Spatafora, J. W., & Ciuffetti, L. M. (2008). Homologs of ToxB, a host-selective toxin gene from Pyrenophora tritici-repentis, are present in the genome of sister-species Pyrenophora bromi and other members of the Ascomycota. *Fungal Genetics and Biology*, *45*(3), 363–377. https://doi.org/https://doi.org/10.1016/j.fgb.2007.10.014

Apama, G., Chatterjee, A., Sonti, R. V., & Sankaranarayanan, R. (2009). A cell wall-degrading esterase of xanthomonas oryzae requires a unique substrate recognition module for pathogenesis on rice. *Plant Cell*, *21*(6), 1860–1873. https://doi.org/10.1105/tpc.109.066886

Baldi, P., & La Porta, N. (2017). Xylella fastidiosa: Host Range and Advance in Molecular Identification Techniques. *Frontiers in Plant Science*, *8*, 944. https://doi.org/10.3389/fpls.2017.00944

Chatterjee, S., Almeida, R. P. P., & Lindow, S. (2008). Living in two worlds: The plant and insect lifestyles of Xylella fastidiosa. *Annual Review of Phytopathology*, *46*, 243–271. https://doi.org/10.1146/annurev.phyto.45.062806.094342

Couto, D., & Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants. *Nature Reviews Immunology*, *16*(9), 537–552. https://doi.org/10.1038/nri.2016.77

Cui, H., Tsuda, K., & Parker, J. E. (2015). Effector-Triggered Immunity: From Pathogen Perception to Robust Defense. *Annual Review of Plant Biology*, *66*(1), 487–511. https://doi.org/10.1146/annurev-arplant-050213-040012

de Guillen, K., Ortiz-Vallejo, D., Gracy, J., Fournier, E., Kroj, T., & Padilla, A. (2015). Structure Analysis Uncovers a Highly Diverse but Structurally Conserved Effector Family in Phytopathogenic Fungi. *PLOS Pathogens*, *11*(10), e1005228.

Duxbury, Z., Ma, Y., Furzer, O. J., Huh, S. U., Cevik, V., Jones, J. D. G., & Sarris, P. F. (2016). Pathogen perception by NLRs in plants and animals: Parallel worlds. *BioEssays*, *38*(8), 769–781. https://doi.org/10.1002/bies.201600046

Engler, C., Kandzia, R., & Marillonnet, S. (2008). A one pot, one step, precision cloning method with high throughput capability. *PLoS ONE*, *3*(11). https://doi.org/10.1371/journal.pone.0003647

Food, E., & Authority, S. (2018). Update of the Xylella spp. host plant database. *EFSA Journal*, *16*(9). https://doi.org/10.2903/j.efsa.2018.5408

Gouran, H., Gillespie, H., Nascimento, R., Chakraborty, S., Zaini, P. A., Jacobson, A., Phinney, B. S., Dolan, D., Durbin-Johnson, B. P., Antonova, E. S., Lindow, S. E., Mellema, M. S., Goulart, L. R., & Dandekar, A. M. (2016). The Secreted Protease PrtA Controls Cell Growth, Biofilm Formation and Pathogenicity in Xylella fastidiosa. *Scientific Reports*, *6*(August), 1–13. https://doi.org/10.1038/srep31098

Hopkins, D. L., & Purcell, a H. (2002). % Cwug Qh 2Kgteg U & Kugcug Qh)
Tcrgxkpg Cpf 1Vjgt ' Ogtigpv & Kugcugu. *Plant Disease*, *86*(10), 1056–1066.

Hou, S., Liu, Z., Shen, H., & Wu, D. (2019). *Damage-Associated Molecular Pattern-*
Triggered Immunity in Plants. *10*(May). https://doi.org/10.3389/fpls.2019.00646

Huang, W., Reyes-Caldas, P., Mann, M., Seifbarghi, S., Kahn, A., Almeida, R. P. P.,
Béven, L., Heck, M., Hogenhout, S. A., & Coaker, G. (2020). Bacterial Vector-
Borne Plant Diseases: Unanswered Questions and Future Directions. *Molecular*
Plant, *13*(10), 1379–1393. https://doi.org/10.1016/j.molp.2020.08.010

Ilyas, M., Hörger, A. C., Bozkurt, T. O., Van Den Burg, H. A., Kaschani, F., Kaiser,
M., Belhaj, K., Smoker, M., Joosten, M. H. A. J., Kamoun, S., & Van Der Hoorn,
R. A. L. (2015). Functional Divergence of Two Secreted Immune Proteases of
Tomato. *Current Biology*, *25*(17), 2300–2306.
https://doi.org/10.1016/j.cub.2015.07.030

Jha, G., Rajeshwari, R., & Sonti, R. V. (2007). Functional interplay between two
Xanthomonas oryzae pv. oryzae secretion systems in modulating virulence on
rice. *Molecular Plant-Microbe Interactions*, *20*(1), 31–40.
https://doi.org/10.1094/MPMI-20-0031

Jones, J. D. G., & Dangl, J. L. (2006). The Plant Immune System. *Nature*, *444*(7117),
323–329.

Katagiri, F., & Tsuda, K. (2010). Understanding the plant immune system. *Molecular*
Plant-Microbe Interactions, *23*(12), 1531–1536. https://doi.org/10.1094/MPMI-
04-10-0099

Kourelis, J., & Hoorn, R. A. L. Van Der. (2018). *Defended to the Nines : 25 Years of*
Resistance Gene Cloning Identi fi es Nine Mechanisms for R Protein Function.
30(February), 285–299. https://doi.org/10.1105/tpc.17.00579

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology*
and Evolution, *35*(6), 1547–1549. https://doi.org/10.1093/molbev/msy096

Lu, Y., Hatsugai, N., Katagiri, F., Ishimaru, C. A., & Glazebrook, J. (2015). Putative
serine protease effectors of clavibacter michiganensis induce a hypersensitive
response in the apoplast of nicotiana species. *Molecular Plant-Microbe*
Interactions, *28*(11), 1216–1226. https://doi.org/10.1094/MPMI-02-15-0036-R

Malukani, K. K., Ranjan, A., Hota, S. J., Patel, H. K., & Sonti, R. V. (2020). Dual
activities of receptor-like kinase OsWAKL21.2 induce immune responses. *Plant*
Physiology, *183*(3), 1345–1363. https://doi.org/10.1104/pp.19.01579

Mermigka, G., Amprazi, M., Mentzelopoulou, A., Amartolou, A., & Sarris, P. F.
(2019). Plant and Animal Innate Immunity Complexes: Fighting Different
Enemies with Similar Weapons. *Trends in Plant Science*, 1–12.
https://doi.org/10.1016/j.tplants.2019.09.008

Michalopoulou, V. A., Kotsaridis, K., & Mermigka, G. (2020). The host exocyst
complex is targeted by a conserved bacterial type III effector protein that
promotes virulence. *BioRxiv*, 1–32.

Mitre, L. K., Sousa Teixeira-Silva, N., Rybak, K., Magalhães, M., Rodrigues de
Souza-Neto, R., Robatzek, S., & Alves de Souza, A. (2021). The Arabidopsis

immune receptor EFR increases resistance to the bacterial pathogens
Xanthomonas and Xylella in transgenic sweet orange. *BioRxiv*,
2021.01.22.427732. https://doi.org/10.1101/2021.01.22.427732

Mollenhauer, H. H., & Hopkins, D. L. (1974). Ultrastructural study of Pierce's disease
bacterium in grape Xylem tissue. *Journal of Bacteriology*, *119*(2), 612–618.
https://doi.org/10.1128/jb.119.2.612-618.1974

Mur, L. A. J., Kenton, P., Lloyd, A. J., Ougham, H., & Prats, E. (2008). The
hypersensitive response; The centenary is upon us but how much do we know?
Journal of Experimental Botany, *59*(3), 501–520.
https://doi.org/10.1093/jxb/erm239

Nascimento, R., Gouran, H., Chakraborty, S., Gillespie, H. W., Almeida-Souza, H. O.,
Tu, A., Rao, B. J., Feldstein, P. A., Bruening, G., Goulart, L. R., & Dandekar, A.
M. (2016). The Type II Secreted Lipase/Esterase LesA is a Key Virulence Factor
Required for Xylella fastidiosa Pathogenesis in Grapevines. *Scientific Reports*,
6(November 2015), 1–17. https://doi.org/10.1038/srep18598

Ngou, B. P. M., Ahn, H. K., Ding, P., & Jones, J. D. G. (2021). Mutual potentiation of
plant immunity by cell-surface and intracellular receptors. *Nature, April 2020*.
https://doi.org/10.1101/2020.04.10.034173

Nissinen, R., Xia, Y., Mattinen, L., Ishimaru, C. A., Knudson, D. L., Knudson, S. E.,
Metzler, M., & Pirhonen, M. (2009). The putative secreted serine protease chp-7
Is required for full virulence and induction of a nonhost hypersensitive response
by clavibacter michiganensis subsp. sepedonicus. *Molecular Plant-Microbe*
Interactions, *22*(7), 809–819. https://doi.org/10.1094/MPMI-22-7-0809

Pardal, A. J., Piquerez, S. J. M., Id, A. D., Id, L. F., Mastorakis, E., Id, E. R., Latrasse,
D., Id, L. C., Gimenez-ibanez, S., Id, S. H. S., Benhamed, M., & Id, V. N. (2021).
Immunity onset alters plant chromatin and utilizes EDA16 to regulate oxidative
homeostasis. 1–26. https://doi.org/10.1371/journal.ppat.1009572

Pérez-Reytor, D., Jaña, V., Pavez, L., Navarrete, P., & García, K. (2018). Accessory
toxins of vibriopathogens and their role in epithelial disruption during infection.
Frontiers in Microbiology, *9*(SEP), 1–11.
https://doi.org/10.3389/fmicb.2018.02248

Pérez-Reytor, D., Pavón, A., Lopez-Joven, C., Ramírez-Araya, S., Peña-Varas, C.,
Plaza, N., Alegría-Arcos, M., Corsini, G., Jaña, V., Pavez, L., del Pozo, T.,
Bastías, R., Blondel, C. J., Ramírez, D., & García, K. (2020). Analysis of the
Zonula occludens Toxin Found in the Genome of the Chilean Non-toxigenic
Vibrio parahaemolyticus Strain PMC53.7. *Frontiers in Cellular and Infection*
Microbiology, *10*(September), 1–13. https://doi.org/10.3389/fcimb.2020.00482

Postma, J., Liebrand, T. W. H., Bi, G., Evrard, A., Bye, R. R., Mbengue, M., Kuhn, H.,
Joosten, M. H. A. J., & Robatzek, S. (2016). Avr4 promotes Cf-4 receptor-like
protein association with the BAK1/SERK3 receptor-like kinase to initiate
receptor endocytosis and plant immunity. *The New Phytologist*, *210*(2), 627–
642. https://doi.org/10.1111/nph.13802

Rapicavoli, J., Ingel, B., Blanco-Ulate, B., Cantu, D., & Roper, C. (2018a). Xylella
fastidiosa: an examination of a re-emerging plant pathogen. *Molecular Plant*
Pathology, *19*(4), 786–800. https://doi.org/10.1111/mpp.12585

Rapicavoli, J., Ingel, B., Blanco-Ulate, B., Cantu, D., & Roper, C. (2018b). Xylella

fastidiosa: an examination of a re-emerging plant pathogen. *Molecular Plant*
Pathology, *19*(4), 786–800. https://doi.org/10.1111/mpp.12585

Roper, C., Castro, C., & Ingel, B. (2019). Xylella fastidiosa: bacterial parasitism with
hallmarks of commensalism. *Current Opinion in Plant Biology*, *50*, 140–147.
https://doi.org/10.1016/j.pbi.2019.05.005

Roy, P. P., Paul, S., Mitra, I., & Roy, K. (2010). Roy et al. On Two Novel Parameters
for Validation of Predictive QSAR Models. Molecules, 2009, 14, 1660-1701.
Molecules, *15*(1), 604–605. https://doi.org/10.3390/molecules15010604

Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for
reconstructing phylogenetic trees. *Molecular Biology and Evolution*, *4*(4), 406–
425. https://doi.org/10.1093/oxfordjournals.molbev.a040454

Schneider, K., van der Werf, W., Cendoya, M., Mourits, M., Navas-Cortés, J. A.,
Vicent, A., & Lansink, A. O. (2020). Impact of Xylella fastidiosa subspecies
pauca in European olives. *Proceedings of the National Academy of Sciences of*
the United States of America, *117*(17), 9250–9259.
https://doi.org/10.1073/pnas.1912206117

Schrodinger, L. L. C., & DeLano, W. (n.d.). *PyMOL*.

Schwessinger, B., & Zipfel, C. (2008). News from the frontline: recent insights into
PAMP-triggered immunity in plants. *Current Opinion in Plant Biology*, *11*(4),
389–395. https://doi.org/10.1016/j.pbi.2008.06.001

Simon, G. M., & Cravatt, B. F. (2010). Activity-based proteomics of enzyme
superfamilies: Serine hydrolases as a case study. *Journal of Biological*
Chemistry, *285*(15), 11051–11055. https://doi.org/10.1074/jbc.R109.097600

Song, J., Win, J., Tian, M., Schornack, S., Kaschani, F., Ilyas, M., Van Der Hoorn, R.
A. L., & Kamoun, S. (2009). Apoplastic effectors secreted by two unrelated
eukaryotic plant pathogens target the tomato defense protease Rcr3.
Proceedings of the National Academy of Sciences of the United States of
America, *106*(5), 1654–1659. https://doi.org/10.1073/pnas.0809201106

Stotz, H. U., Mitrousia, G. K., de Wit, P. J. G. M., & Fitt, B. D. L. (2014). Effector-
triggered defence against apoplastic fungal pathogens. *Trends in Plant Science*,
19(8), 491–500. https://doi.org/10.1016/j.tplants.2014.04.009

van der Burgh, A. M., & Joosten, M. H. A. J. (2019). Plant Immunity: Thinking
Outside and Inside the Box. *Trends in Plant Science*, *xx*(xx), 1–15.
https://doi.org/10.1016/j.tplants.2019.04.009

Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J., & Zhang, Y. (2015). The I-TASSER
Suite: protein structure and function prediction. *Nature Methods*, *12*(1), 7–8.
https://doi.org/10.1038/nmeth.3213

Yang, J., & Zhang, Y. (2015). I-TASSER server: new development for protein
structure and function predictions. *Nucleic Acids Research*, *43*(W1), W174–
W181. https://doi.org/10.1093/nar/gkv342

Zhang, S., Chakrabarty, P. K., Fleites, L. A., Rayside, P. A., Hopkins, D. L., &
Gabriel, D. W. (2015). Three new pierce's disease pathogenicity effectors
identified using xylella fastidiosa biocontrol strain EB92-1. *PLoS ONE*, *10*(7), 1–
17. https://doi.org/10.1371/journal.pone.0133796

Zuckerkandl, E., & Pauling, L. (1965). Molecules as documents of evolutionary
history. *Journal of Theoretical Biology, 8*(2), 357–366.
https://doi.org/10.1016/0022-5193(65)90083-4

Tables and Figures

Figure 1

(a) D934_08750

(b) RA12_01530

(c) PD_1703

(d) D934_07885

(e) PD_0956

(f) RA12_05570

(g) D934_09300

(h) PD_0915

(i) D934_ 09265

Figure 2

2a.

2b.

Figure 3

A.

B.

C.

D.

Xylella fastidiosa Proteins	Sequence similarity (PDB)	Description	Structural template (PDB)	Description
PD_0956	3WY8	Hydrolase/Protease	3WY8	Hydrolase/Serine protease
PD_0915	2R2A	Zonular Occludens Toxin (Zot)	2DHR	ATP-dependent metalloprotease/Hydrolase
PD_1703	3WY8	Hydrolase/serine protease	1Z8G	Hydrolase/Hydrolase Inhibitor
D934_08750	3H2K	Hydrolase/Esterase (LipA)	3H2K	Hydrolase/ Esterase (LipA)
D934_07885	3WY8	Hydrolase/Serine protease	1Z8G	Hydrolase/Hydrolase Inhibitor
D934_09300	2R2A	Zonular Occludens Toxin (Zot)	2R2A	Zonular Occludens Toxin (Zot)
D934_09265	2R2A	Zonular Occludens Toxin (Zot)	4WWO	Transferase/Transferase Inhibitor
RA12_01530	3H2K	Hydrolase/ Esterase (LipA)	3H2K	Hydrolase/ Esterase (LipA)
RA12_05570	3WY8	HydrolaseSerine protease	1Z8G	Hydrolase/Hydrolase Inhibitor
D934_00810	5N8P	Membrane Protein	3JAV	Transport Protein
D934_05685	5N8P	Membrane Protein	5IJO	Transport Protein
D934_12725	3JAV	Transport Protein	3JAV	Transport Protein
D934_08755	3H2K	Hydrolase/ Esterase (LipA)	3H2K	Hydrolase/ Esterase (LipA)
D934_12535	3H2K	Hydrolase/ Esterase (LipA)	3H2K	Hydrolase/ Esterase (LipA)
D934_12795	7KVE	Blood Clotting	1RWR	Cell Adhesion
RA12_11155	4UIC	Sugar Binding Protein	1G6O	Hydrolase
RA12_11125	3V05	Toxin	5N8P	Membrane Protein
RA12_03930	6VDP	Oxidoreductase	6VDP	Oxidoreductase
RA12_03905	6W1S	Gene Regulation	6WIS	Gene Regulation

Supplementary Tables and Figures

Supplementary Table S1

Xf gene ID	Subspecies/ Strains	Host
PD0915	strain Temecula1	Grapevine
PD0956	strain Temecula1	Grapevine
PD1703	strain Temecula1	Grapevine
RA12_01530	strain CoDiRO	Olive tree
RA12_03905	strain CoDiRO	Olive tree
RA12_03930	strain CoDiRO	Olive tree
RA12_05570	strain CoDiRO	Olive tree
RA12_11125	strain CoDiRO	Olive tree
RA12_11155	strain CoDiRO	Olive tree
D934_00810	subsp. sandyi Ann-1	Oleander
D934_05685	subsp. sandyi Ann-1	Oleander
D934_07885	subsp. sandyi Ann-1	Oleander
D934_08750	subsp. sandyi Ann-1	Oleander
D934_08755	subsp. sandyi Ann-1	Oleander
D934_09265	subsp. sandyi Ann-1	Oleander
D934_09300	subsp. sandyi Ann-1	Oleander
D934_12535	subsp. sandyi Ann-1	Oleander
D934_12725	subsp. sandyi Ann-1	Oleander
D934_12795	subsp. sandyi Ann-1	Oleander

Supplementary Figure S1

(a) D934_00810

(b) D934_05685

(c) D934_12725

(d) RA12_11125

(e) D934_08755

(f) D934_12535

(g) D934_12795

(h) RA12_11155

(i) RA12_03930

(k) RA12_03905

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4