Manuscript title – Fitness Landscapes Reveal Context-dependent Benefits of Oviposition

Choice

Vrinda Ravi Kumar[1]*, Gaurav Agavekar[1,2] and Deepa Agashe[1]*

[1]National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore,

India

[2]Current address: Okinawa Institute of Science and Technology (OIST), Japan

*Correspondence:

V. Ravi Kumar - vrindaravikumar42@gmail.com (Phone: +91 8971 396 960); ORCID: 0000-

0001-6054-2694

D. Agashe - dagashe@ncbs.res.in (Address: National Centre for Biological Sciences (NCBS-

TIFR), GKVK, Bellary Road, Bangalore, Karnataka, India 560065; phone: +91 80 2366 6524;

fax : +91 80 2363 6662); ORCID: 0000-0002-0374-8159

STATEMENT OF AUTHORSHIP

VRK and DA conceived the work; VRK, DA and GA designed experiments; VRK and GA

conducted experiments; VRK and DA analysed data and wrote the manuscript; DA acquired

funding.

Keywords – optimal choice, context-dependence, female oviposition, resource preference,

heterogeneous habitat

Article Type - Research article

Running title – Optimality of oviposition preference

Word count – abstract - 160; main text (excluding title page, acknowledgements, references and

figure legends) – 4363

Reference count – 68

Figure count – 4

ABSTRACT

Resource choice behaviour has enormous fitness consequences and can drive niche expansion. However, individual behavioural choices are often mediated by context, determined by past experience. Are such context-dependent behaviours adaptive? Using *Tribolium castaneum* (the red flour beetle)*,* we demonstrate that context-dependent oviposition choice reflects distinct, context-specific local fitness peaks. Manipulating female egg allocation in a habitat containing optimal and suboptimal resource patches, we measured offspring fitness to generate fitness landscapes as a function of all possible oviposition behaviours (i.e., combinations of fecundity and resource preference). Females from different age and competition contexts exhibit distinct behaviours which optimize different fitness components that are linked in a tradeoff. With increasing age and prior exposure to competition, they produce few but fast-developing offspring that are advantageous under high resource competition. In contrast, young naïve females produce many slow-developing offspring, beneficial under weak competition. Systematically mapping complete context-dependent fitness landscapes is thus critical to infer behavioural optimality and offers predictive power in novel contexts.

INTRODUCTION

Organisms often encounter heterogeneous habitats, and their behavioural choices in such situations ultimately determine their fitness. This is perhaps best exemplified by females' reproductive decisions when faced with alternate resources. For example, females of phytophagous insects can distribute their eggs on different host plants, which affects offspring survival and growth and determines their reproductive success (1-2). Therefore, oviposition preference is predicted to face strong selection on par with other life history traits (3), sometimes leading to genetic correlations between preference and performance (2). However, oviposition choice can also vary dramatically with females' prior experience, reflecting the ecological or internal context determined by their environment or life history (4). For instance, females of the polyphagous moth *Spodoptera littoralis* that develop on a specific host plant as larvae also prefer it subsequently for oviposition (5). More generally, ecological factors such as age and resource competition change across individuals' lifespan, altering key components of oviposition behaviour such as fecundity (6-9) and oviposition choice (10–11). Given the potentially enormous range of contexts experienced by different individuals across generations, how do these context-dependent oviposition behaviours evolve?

Broadly, two kinds of selection pressures may drive context-dependent oviposition choice. First, female choice may evolve under selection for a single globally optimal (context-independent) preference, with context-dependent deviations from this optimum reflecting weak selection, environmental constraints or tradeoffs. For instance, limited perception of resource quality (12) or other search-related constraints (2, 13, 14) may lead to suboptimal preference, as could tradeoffs between maternal and offspring fitness (15) or the presence of competitors or predators (16-17). Alternatively, context-specific female choice may reflect context-specific selection to optimize distinct fitness peaks, with variable female choice perfectly attuned for each context (18). To our knowledge, no study has distinguished between these alternative hypotheses, perhaps because it is challenging to comprehensively connect past context to future fitness. For instance, to identify context-specific fitness optima it is critical to measure multiple fitness components, because many traits contribute to fitness and may trade off with each other (19–22) across contexts. Further, to correctly infer whether an observed behaviour is locally or globally

adaptive (or maladaptive), we must generate a unified fitness landscape that encompasses all

possible behavioural choices, including those outside the range of behaviours observed in nature.

Using females from diverse contexts to create complete fitness landscapes also allows the

prediction of female fitness in novel contexts. Illuminating the entire phenotypic fitness

landscape is thus critical to understand the evolution of context-dependent oviposition behaviour.

We created such fitness landscapes for multiple fitness variables for the red flour beetle

Tribolium castaneum, a widespread generalist pest of cereal flours that has long served as a

model system in ecology and evolutionary biology (23, 24). Prior work on this species

demonstrates significant context-dependent variation in key oviposition behaviours, modulated

by age and population density (indicating strength of resource competition). Here, we explicitly

connect prior female context, behavioural choice, and fitness, by conducting three sets of

experiments focused on a choice between an optimal resource (wheat flour) and a suboptimal

novel resource (finger millet flour) (Figure 1). We first validated the context-dependence of two

aspects of oviposition behaviour (fecundity and preference) of females from six different age and

density contexts (Experiment 1). Second, we generated complete fitness landscapes as a function

of the full range of observed female fecundity and possible oviposition choices. To do this, we

experimentally manipulated egg allocation in a habitat with both resource patches, and measured

multiple components of offspring fitness (Experiment 2). Finally, we validated the fitness

landscape by jointly measuring oviposition behaviours and consequent offspring fitness for

females from three commonly-experienced contexts (Experiment 3). By placing oviposition

behaviour in a comprehensive fitness landscape, we demonstrate that divergent, context-

dependent oviposition behaviours could evolve via selection driving females towards context-

specific local adaptive peaks for distinct fitness components.

METHODS

Collecting females from different age and density contexts

We derived experimental individuals from previously described large, outbred laboratory "stock"

populations of *T. castaneum* (25), provided with either wheat or finger millet flour. Flour beetles

have highest fitness in wheat and show significantly slower development in finger millet (~37%

slower development relative to wheat) (25). Details of population maintenance are given in SI

methods. As required for different experiments (Figure 1), we generated females from six distinct

age and density contexts (Figure S1). We used 'young', 'mid-aged' and 'old' females aged ~2

weeks, ~4 weeks and ~6 weeks after eclosion respectively (SI Methods), that experienced either

low density ("LD") or high density ("HD") (Figure S1, SI Methods). Females from each context

were mated with males from the same context to ensure consistency.

Measuring oviposition behaviour and offspring fitness

To test how prior context changes oviposition behaviour, we measured the oviposition preference

of females from different contexts (Figure 1, Experiment 1) for wheat vs. finger millet flour in

individual 2-choice assays (SI Methods, Figure S2). We used the total number of eggs laid per

female over 2 days to quantify fecundity, and the percentage of eggs deposited in finger millet as

a proportional measure for oviposition preference for finger millet. The small oviposition

window minimized egg cannibalism by larvae and adults. To measure the fitness consequences

of female oviposition behaviours (Figure 1, Experiment 3), we removed adults and allowed

offspring to develop undisturbed for 21 days without restricting larval movement, and quantified

the resultant offspring fitness. We measured the proportion of eggs that survived three weeks of

development ("survival"), the proportion of offspring that had successfully pupated or eclosed

("development rate"), and the total number of live offspring of all stages ("total offspring"). All

measures of fitness were evaluated for the entire habitat, i.e., including both resource patches.

Thus, we assessed the overall optimality of females' reproductive strategies as they would

operate in natural conditions, including all interactions between offspring and with their natal

environment.

Generating and validating fitness landscapes

To generate complete fitness landscapes (Figure 1, Experiment 2), we had to quantify the impact

of the entire range of possible oviposition behaviours on offspring fitness. Hence, we collected

females from three different contexts with distinct ranges of fecundity (young LD, young HD

and old HD; see Results); but we did not allow them to express their oviposition choice. Instead,

we collected eggs from each female and manipulated egg allocation systematically across wheat

and finger millet patches in a 2-patch habitat (as in the oviposition choice assays), mimicking the

full range of possible oviposition preferences (n=16-20 females per allocation). By distributing

females with various fecundity values across egg allocation treatments, we dissociated female

fecundity from resource preference, and then measured female fitness as described above.

Using these data, we generated fitness landscapes for each fitness component as a function of

female fecundity (observed) and "preference" for finger millet (experimentally manipulated egg

allocation). To do this, we extrapolated fitness values using local smoothing ('loess') by

estimating the value of the three fitness measures for all possible combinations of female

fecundity and oviposition preference using the 'panel.2dsmoother' function (26, from the

latticeExtra package) in the levelplot function from the lattice package (27). Fitness was

evaluated in an equally spaced 20 by 20 point grid across the sampled range of fecundity and

preference. We also tested the impact of using a different smoothing function (generalized

additive model, GAM) and altered grid size (ranging from 10 x 10 to 200 x 200 points).

These fitness landscapes allowed us to predict the consequences of any given combination of

fecundity and oviposition preference. To validate the predictions (Figure 1, Experiment 3), we

allowed females from three different contexts (young LD, young HD and old HD) to make

oviposition choices; counted and returned eggs to the patch in which they were laid; and

measured offspring fitness for each female. Using the median values of observed fecundity and

preference for females from each context, we predicted their offspring fitness based on the

fitness landscape and compared the predicted values with the median observed fitness values

using a paired t-test.

Statistical analysis

All analyses were conducted using R (28) in RStudio (29). We assessed the impact of age and

density on female fecundity and of age, density and fecundity on oviposition preference

(Experiment 1) and the impact of context, fecundity, and manipulated egg allocation on offspring

fitness components (Experiment 2). We used ANOVA models for fecundity; generalised linear

models with binomially distributed errors ('binomial GLM') for female oviposition preference,

offspring survival and development rate; and a GLM with Poisson errors ('Poisson GLM') for the

total number of offspring (SI methods).

RESULTS

Prior context alters female oviposition behaviour

Based on earlier work, we expected that female age and prior exposure to population density (an indicator of resource competition) would impact oviposition behaviour, i.e., female fecundity and preference for the novel finger millet resource (Figure 1, Experiment 1). Overall, fecundity ranged from 0 to 48 eggs, and oviposition choice ranged from 0 to 100% preference for finger millet. However, these traits differed significantly across females from different contexts (Figure 2A, 2B). For instance, female fecundity typically decreased with age and prior exposure to high population density ("HD") (Figure 2A; ANOVA, F = 32.34, p < 0.001). Young females at low density ("LD") had the highest fecundity, whereas HD females of the same age laid only half as many eggs (Tukey's HSD, difference ~16 eggs, p < 0.001), similar to mid-aged and older females that experienced HD throughout their adult life (Tukey's HSD, difference (mid-aged HD) = 15.5 eggs, p < 0.001; difference (old HD) = 18.5 eggs, p < 0.001). This broad effect of age was consistent when the same set of young females were allowed to age (Figure S4); and in an independent experimental block with young LD, young HD and old HD females (Figure S5). Concomitant with fecundity, oviposition preference for finger millet also typically decreased with age and density (Figure 2B), except for old HD females which oviposited randomly. Combining data across contexts, females with high fecundity laid more eggs in finger millet (Spearman's rho = 0.303, p < 0.001; Figure 2C). However, fecundity and preference were not correlated within a given context (except for aged LD females; Figure S6 shows more experimental blocks).

Despite these context-specific trends in oviposition behaviour, we noted some exceptions. When young HD females were aged, they continued to show a strong preference for wheat (typical of HD females) despite experiencing LD for 4 weeks (compare young HD vs. aged, Figure 2B), and also failed to exhibit the high fecundity commonly observed at low density (Figure 2A). Thus, the impact of early experience with HD seemed to last for a very long time. Additionally, old HD females did not prefer wheat as predicted by the age- and density- dependent trends in other females. Hence, preference may shift non-monotonically as females experience high density for different time intervals. Nonetheless, it is clear that age and density present appropriate contexts

that influence two key components of oviposition behaviour. Therefore, we proceeded with
measuring the fitness consequences of these divergent context-dependent resource preference
behaviours.

Fitness landscapes predict consequences of divergent oviposition behaviour
To quantify the fitness impacts of potential oviposition behaviours (Figure 1, Experiment 2), we
allowed females to lay eggs in wheat and then manually distributed each female's eggs in wheat
and finger millet patches, covering the entire range of possible oviposition preference. To include
the widest possible range of fecundity (Figure 2A), we sampled females from three different
contexts (young LD, young HD, and old HD females), and measured three metrics of offspring
fitness (Figure S7). This design ensured that we sampled the full range of possible fecundity–
preference combinations to estimate their impact on fitness.

Across female contexts, offspring survival was largely invariable (~75% offspring survived).
Interestingly, young LD females produced twice as many offspring as females from other
contexts (~20 vs. ~10 per female), although they had the slowest development (~25% offspring
completed pupation in 3 weeks, vs. ~50-75% for young HD and old HD females). Thus, distinct
oviposition behaviours associated with different female contexts can dramatically alter offspring
fitness. Notably, offspring development rate was strongly negatively correlated with female
fecundity as well as total offspring (Figure 3A–B, Figure S8), reflecting a well-known density-
regulated tradeoff between offspring number and development time in this species (23).

To visualize the impacts of oviposition behaviour on fitness, we used data for each fitness
component (Figures 3C-E) to construct complete fitness landscapes as a function of the two key
aspects of oviposition behaviours (female fecundity and oviposition choice; Figures 3F-H). Since
the landscapes encompass the entire range of possible oviposition behaviours (all possible
preferences and the entire natural fecundity range), they offer critical insight into the fitness
consequences of oviposition decisions for all females when offered a similar choice. Recall that
females from different contexts exhibit distinct combinations of fecundity and preference (Figure
2), such that they occupy different positions in these landscapes.

As expected, the fitness landscape for survival was relatively flat (focus on the contours,
ignoring triangles; Figure 3F). However, landscapes for the other two fitness components paint a
more interesting dichotomy. Overall, most of the variation in fitness was distributed along the
fecundity axis, suggesting (as expected) that fecundity is a powerful modulator of fitness
outcomes. For development rate, the landscape unsurprisingly predicts a fitness optimum at low
fecundity and low preference for finger millet; with rapid development at low fecundity
regardless of preference (Figure 3G) suggesting that competition among offspring strongly
impacts development rate. In contrast, offspring production is maximized at high fecundity (as
expected) and strong preference for finger millet, with a relatively weak impact of preference
(Figure 3H). These contrasting predicted fitness optima for offspring development rate and total
number of offspring largely reflect the underlying negative tradeoff between the two components
(Figure 3B, independent experimental blocks in Figure S8), primarily mediated by fecundity
(Figure 3A). The estimated fitness maxima predicted by the landscapes were robust to changes in
the fitting method (loess vs. GAM) and the grid density over which fitness was evaluated.

Separate fitness landscapes constructed for each female context generated similar patterns,
indicating that the unified fitness landscape is robust (Figures S9, S10). We also confirmed that
the landscapes accurately predict the fitness consequences of any combination of female
fecundity and oviposition choice. In an independent experiment, we allowed females to express
their oviposition preference (used to predict fitness) and measured the fitness consequences of
their choice (Figure 1, Experiment 3). Observed female fitness matched predictions from the
original landscape (compare the triangle colour with background colour, Figures 3G–I; Table S1,
$p > 0.05$ for all three fitness parameters), even though the oviposition behaviour of old HD
females in this experiment was different from what we had observed in Experiment 1
(comparison shown in Figure S5). Thus, fitness landscapes generated by scrambling the inherent
association between female fecundity and preference and fitness accurately predicted the fitness
consequences of diverse oviposition behaviours.

Females from different contexts use alternate adaptive oviposition strategies
The fitness landscapes generated above suggest two general oviposition strategies that each
increase different fitness components – high preference for wheat with low fecundity, or high

acceptance of finger millet with high fecundity. To explicitly connect prior female context with

these strategies, we asked whether context-dependent behaviours are consistent with either of

these predicted fitness peaks. To do this, we used the observed context-specific oviposition

behaviours described in Figure 2, which represent females from diverse age and density contexts.

We predicted the fitness outcomes for each context by overlaying the median values of female

fecundity and preference onto the landscapes for each fitness component. As expected, females

from different contexts were positioned widely across the landscape (Figures 3I-K). As noted

earlier (Figure 2A), females that were older or experienced high density generally had low

fecundity; whereas younger females or those at low density laid many more eggs. At the same

time, they also had divergent oviposition choices. In light of the fitness landscapes, we can

connect these contrasting patterns, now revealed as alternate adaptive strategies. For instance,

young LD females were close to the peak for total offspring, whereas young HD females produce

fewer, rapidly-developing offspring. Distinct oviposition behaviours thus place females close to

different locally adaptive fitness peaks.

Although the fitness landscapes were generated using females from only three contexts, we could

also use them to predict fitness for other contexts, using only their oviposition behaviours.

Strikingly, females from the two "aged" contexts also present divergent outcomes, though both

(young HD and young LD females) were aged for 4 weeks under identical LD conditions. Their

distinct positions in the fitness landscapes are driven primarily by their distinct fecundity,

indicating a long-lasting impact of early exposure to high density. Aged young HD females

showed a dramatic reduction in offspring production (Figure 3K), but their increased preference

for wheat increased offspring development rate (Figure 3J). In summary, the complete fitness

landscapes reveal that maternal context alters distinct fitness components by differentially

influencing two key aspects of oviposition behaviour.

DISCUSSION

Here, we present the first comprehensive analysis of the linkages between females' prior context,

their oviposition behaviour, and offspring fitness. By governing female behaviour, the contexts

determine females' placement on the fitness landscapes, defining their fitness. Thus, a context-

specific interpretation of fitness is critical to infer optimality of specific behaviours. Importantly, to identify distinct local fitness peaks we need separate landscapes for each fitness component. For instance, if we only measured offspring development rate, we would incorrectly conclude that young LD females behave maladaptively. However, using the ecological selection faced by females in their prior context (here, high vs. low competition) we can better assess the adaptive consequences of their behaviour. We thus infer that under conditions of resource abundance (when their offspring are unlikely to face strong competition), females can effectively maximize the total number of offspring even if they develop slowly. Such context-dependent optimization of different fitness components may explain prior cases of apparent maladaptive resource use. For instance, flour beetles exhibited seemingly maladaptive density-dependent niche contraction in a heterogeneous habitat containing wheat and corn (a poor resource) (31). Based on our current results, we suggest that the observed preference for wheat at higher densities may reflect optimization of offspring development rate or survival; whereas at low density, females maximize total offspring production. Thus, age and density contexts primarily alter female fecundity, and females accordingly modulate their oviposition resource choice to optimize either total offspring or offspring development rate. Thus, our results demonstrate the lack of a single global fitness peak for oviposition behaviour, and show that a change in context can shift female behaviour towards alternative fitness peaks.

How generalisable is the suggested coupling between context-specific behaviour and fitness? Here, we considered two very broadly applicable contexts across species – age and experienced population density (competition) – that differ both between individuals in a population and across an individuals' lifespan. In *Tribolium*, transitions across contexts are expected to occur in association with population growth and dispersal events (32-35). Ageing is universal and change in the strength of competition is a common feature of the life history of many organisms; both factors can impact oviposition. Mechanistically, any change in context that alters female body condition could influence resource allocation towards competing aspects of fitness (reflecting tradeoffs), ultimately determining oviposition behaviour (36–38). Hence, we can broadly frame our results to make generalized predictions about the context-dependent optimization of fitness for any species (Figure 4). While the exact fitness components that face selection or trade off across contexts may vary across species or environments, we suggest that the core axes that

determine the impact of context on behaviour – age and resource competition – should be widely
applicable.

Our results show that females in different contexts can optimize different fitness components that
trade off with each other. How does preference evolve, given that females' behaviour is pulled in
opposing directions across their lifetime? Broadly, frequent changes in selective context during
the evolutionary history of a species could lead to the observed strong, specific associations
between female context and expressed oviposition behaviour. We speculate that in *T. castaneum,*
instead of selection acting independently on both fitness components, only offspring
development rate evolves under strong selection. This is generally expected for species with a
high intrinsic growth rate that often experience high population density, such as pests and
invasives (39,40). At different times during their life (depending on age and density), female
beetles may thus experience either intensified or relaxed selection to produce fast-developing
offspring. For instance, prior work shows that larval development time in *Tribolium* increases
under high larval or adult density (41,42). Thus, one way to increase offspring development rate
under high resource competition is to decrease fecundity, and *T. castaneum* females may employ
this strategy (43). Such selection on development rate may additionally drive high preference for
optimal resources (here, wheat) under high competition. But how do we explain the random
resource preferences of both old HD females and young LD females in this framework? One
possibility is that old females – especially after prolonged exposure to high density – are unable
to make effective choices, and therefore allocate eggs randomly. Meanwhile, under low
competition (e.g., young LD females), selection on offspring development rate should be weak,
allowing females to randomly allocate their eggs across habitats. Thus, distinct mechanisms may
explain the convergent random preference behaviour of young LD and old HD females; this
hypothesis needs to be explicitly tested. Nonetheless, our data suggest that both age and density
have immediate impacts on fecundity, in turn altering female choice. Hence, fecundity and
oviposition choice are inextricable, whereas most prior work on female context has focused on
fecundity alone (38,43,44). We suggest that for dispersing species that often encounter
heterogeneous habitats, joint measurement of fecundity and resource choice is essential to fully
understand oviposition behaviour.

Our results also suggest a long-lasting impact of prior context on behaviour. Aged young HD
females – despite a 4-week exposure to low density – continued to produce fast-developing
offspring (optimal at high density), instead of increasing fecundity to maximize total offspring
(optimal at low density). The failure to increase fecundity in response to a relaxation in adult
competition is costly (~15 fewer offspring than expected under low density) and surprising (45),
because flour beetles routinely experience changes in population density that drive their
population dynamics (46) and dispersal (32,33,35). As a result, we expect flour beetles to
experience strong selection to rapidly respond to changes in competition and alter oviposition
behaviours. We speculate that early physiological changes due to high density (47–51) may be
irreversible, preventing females from optimizing their oviposition behaviours even under
favourable conditions. Such "carry-over effects" are widespread in the animal kingdom (52–54).
Alternatively, the suboptimal suppression of fecundity may persist in the population due to weak
selection, e.g. if old isolated (i.e., recently dispersed) females are rare in the normal life cycle of
T. castaneum (33). Further work is needed to understand this interesting phenomenon and
distinguish between the various hypotheses.

Context-dependent oviposition behaviour has longer-term implications for species' evolution.
For instance, reduced female preference for the ancestral or optimal resource could initiate the
process of niche expansion by exposing offspring to selection in a new habitat, or by influencing
the resource preference and performance of subsequent generations. We showed previously that
in flour beetles, larval experience with novel resources increases later use of the same resources
(25); a pattern that is also observed in many other insects (55–58). Hence, initial female
acceptance of a new resource can increase its use by offspring, exposing offspring to selection
and potentially improving their performance over time. Therefore, preference can lead and
performance may follow when species colonize new habitats (59-61). More generally, random or
no-preference behaviours have been suggested to drive the evolution of generalist species (62-
64). Our work indicates that such effects may occur only when females from specific contexts
colonize new habitats (in our case, old HD and young LD females). Hence, the prior experience
and history of colonizing females may be important determinants of successful niche expansion.

In conclusion, we suggest that it is important to quantify the linkages between female experience, body condition, oviposition behaviour, and offspring fitness to enable context-informed interpretations of optimality. Understanding these links is important to successfully use oviposition behaviours for pest management, such as in the push-pull strategy (65), where experience-dependent behaviour is exploited to reduce oviposition on crops. Although a large body of prior work has partially addressed these links, we have lacked predictive power in new environmental or individual contexts. Here, we show an effective and reliable way to generate broad predictions about the fitness consequences of female oviposition behaviours. By dissociating females' preference from their fecundity, one can systematically cover the entire range of phenotype values that comprise the fitness landscape, including combinations of fecundity and preference that may not be directly or easily observed. One can then predict the fitness consequences of female behaviour in new contexts, without the need to measure fitness in each scenario. Studies documenting evolutionary traps already highlight the power of explicitly considering longer-term historical context to understand behaviour (66,67). We suggest that considering short-term female context is similarly important to understand female behaviours as drivers of niche expansion.

ACKNOWLEDGEMENTS

We thank Laasya Samhita, Pratibha Sanjenbam, Rittik Deb, Shyamsunder Buddh, Sneha Garge and other members of the Agashe lab for discussion and comments on the manuscript. We thank Sumita Nanda for the beetle illustration used in figures. We acknowledge funding and support from the National Centre for Biological Sciences (NCBS-TIFR), the Department of Atomic Energy, Government of India (Project Identification No. RTI 4006) and a SERB Women Research Excellence Award (WEA/2020/000030) to DA.

The authors declare no conflict of interest.

REFERENCES

1. Jaenike J. On Optimal Oviposition Behavior in Phytophagous. Theor Popul Biol. 1978;14(3):350–6.

2. Gripenberg S, Mayhew PJ, Parnell M, Roslin T. A meta-analysis of preference–

performance relationships in phytophagous insects. Ecol Lett. 2010;13(3):383–93.

3. Resetarits WJ. Oviposition site choice and life history evolution. Am Zool.
1996;36(2):205–15.

4. Anderson P, Anton S. Experience-based modulation of behavioural responses to plant
volatiles and other sensory cues in insect herbivores. Plant Cell Environ.
2014;37(8):1826–35.

5. Proffit M, Khallaf MA, Carrasco D, Larsson MC, Anderson P. 'Do you remember the
first time?' Host plant preference in a moth is modulated by experiences during larval
feeding and adult mating. Ecol Lett. 2015;18(4):365–74.

6. Coulson T, Milner-Gulland EJ, Clutton-Brock T. The relative roles of density and
climatic variation on population dynamics and fecundity rates in three contrasting
ungulate species. Proc R Soc B Biol Sci. 2000; 267:1771–1779.

7. Karjalainen J, Urpanen O, Keskinen T, Huuskonen H, Sarvala J, Valkeajärvi
P, Marjomäki TJ. Phenotypic plasticity in growth and fecundity induced by strong
population fluctuations affects reproductive traits of female fish. Ecol Evol. 2016; 6:779-
90.

8. Berggreen IE, Offenberg J, Calis M, Heckmann LH. Impact of density, reproduction
period and age on fecundity of the yellow mealworm *Tenebrio molitor* (Coleoptera:
Tenebrionidae). J Insects Food Feed. 2018;4:43–502018.

9. Khan I, Prakash A, Issar S, Umarani M, Sasidharan R, Masagalli JN, Lama P, Venkatesan
R, Agashe D. Female Density-Dependent Chemical Warfare Underlies Fitness Effects of
Group Sex Ratio in Flour Beetles. Am Nat. 2018; 191:306-17.

10. Yoshioka M, Couret J, Kim F, McMillan J, Burkot TR, Dotson EM, Kitron U, Vozquez-
Prokopec GM. Diet and density dependent competition affect larval performance and
oviposition site selection in the mosquito species *Aedes albopictus* (Diptera: Culicidae).
Parasites and Vectors. 2012; 5(225):1–11.

11. Raitanen J, Forsman JT, Kivelä SM, Mäenpää MI, Välimäki P. Attraction to conspecific
eggs may guide oviposition site selection in a solitary insect. Behav Ecol.
2014;25(1):110–6.

12. Bernays EA. Neural Limitations in Phytophagous Insects : Implications for Diet
Breadth and Evolution of Host Affiliation. Annu Rev Entomol. 2001; 46:703-27.

13. Wood CW, Wice EW, Del Sol J, Paul S, Sanderson BJ, Brodie ED. Constraints imposed
by a natural landscape override offspring fitness effects to shape oviposition decisions in
wild forked fungus beetles. Am Nat. 2018;191(4):524–38.

14. Price PW. Macroevolutionary Theory on Macroecological Patterns. 2003. Cambridge
University Press. Cambridge.

15. García-Robledo C, Horvitz CC. Parent-offspring conflicts, "optimal bad motherhood"
and the "mother knows best" principles in insect herbivores colonizing novel host plants.
Ecol Evol. 2012;2:1446–57.

16. Binckley CA, Resetarits WJ. Oviposition behavior partitions aquatic landscapes along
predation and nutrient gradients. Behav Ecol. 2008;19(3):552–7.

17. Touchon JC, Worley JL. Oviposition site choice under conflicting risks demonstrates that
aquatic predators drive terrestrial egg-laying. Proc R Soc B Biol Sci. 2015;
282:20150376.

18. Dvorak J, Gvozdik L. Adaptive accuracy of temperature oviposition preferences in newts.
Evol Ecol. 2010; 24:1115-27.

19. Schluter D, Price TD, Rowe L. Conflicting selection pressures and life history trade-offs.
Proc R Soc B Biol Sci. 1991;246(1315):11–7.

20. Walsh B, Blows MW. Abundant Genetic Variation + Strong Selection = Multivariate
Genetic Constraints□: A Geometric View of Adaptation. Annu Rev Ecol Evol Syst. 2009;
40:41-59.

21. Lailvaux SP, Kasumovic MM. Defining individual quality over lifetimes and selective
contexts. Proc R Soc B Biol Sci. 2011;278(1704):321–8.

22. Taff CC, Steinberger D, Clark C, Belinsky K, Sacks H, Freeman-Gallant CR, Dunn PO,
Whittingham LA. Multimodal sexual selection in a warbler: Plumage and song are
related to different fitness components. Anim Behav [Internet]. 2012;84(4):813–21.
Available from: http://dx.doi.org/10.1016/j.anbehav.2012.07.002

23. Sokoloff A. The Biology of *Tribolium* (volume 1). Oxford, UK: Oxford University Press.
1972.

24. Pointer MD, Gage MJG, Spurgin LG. *Tribolium* beetles as a model system in evolution
and ecology. Heredity. 2021; 126:869-83.

25. Ravi Kumar V, Issar S, Agashe D. The induction of larval resource preference in

heterogeneous habitats. Ecol Entomol. 2018; 43(6):719–30.

26. Sarkar D, Andrews F. latticeExtra: Extra Graphical Utilities Based on Lattice. 2019. R

package version 0.6-29.

27. Sarkar D. Lattice: Multivariate Data Visualization with R. 2008. Springer, New York

28. R Core Team. R: A language and environment for statistical computing. 2017. R

Foundation for Statistical Computing, Vienna, Austria.

29. RStudio Team. RStudio: Integrated Development for R. 2020. RStudio, Boston,

Massachusetts, USA.

30. Good NE. The Flour Beetles of the genus *Tribolium.* USDA Tech. Bull. 1936; 5, 27-28.

31. Parent CE, Agashe D, Bolnick DI. Intraspecific competition reduces niche width in

experimental populations. Ecol Evol. 2014;3978–90.

32. Ziegler JR. Dispersal and Reproduction in *Tribolium*: The Influence of Food Level. J

Insect Physiol. 1977;23:955–60.

33. Ziegler JR. Evolution of the Migration Response⬚: Emigration by *Tribolium* and the

Influence of Age. Evolution. 1976;30(3):579–92.

34. Gurdasani K, Rafter MA, Daglish GJ, Walter GH. The dispersal flight of *Tribolium*

castaneum - A field test of laboratory generated predictions. J Stored Prod Res.

2019;83:25–33.

35. Gurdasani K, Rafter MA, Daglish GJ, Walter GH. Characterising the variables associated

with *Tribolium castaneum* adults that initiate flight in laboratory tests: Generating

predictions for the field. J Stored Prod Res. 2018;79:123–31.

36. Papaj DR. Ovarian dynamics and host use. Annu Rev Entomol. 2000;45:423–48.

37. Jervis MA, Boggs CL, Ferns PN. Egg maturation strategy and its associated trade-offs⬚:

a synthesis focusing on Lepidoptera. Ecol Entomol. 2005;30:359–75.

38. Berger D, Walters R, Gotthard K. What limits insect fecundity? Body size- and

temperature-dependent egg maturation and oviposition in a butterfly. Funct Ecol.

2008;22(3):523–9.

39. Sakai AK, Allendorf FW, Holt JS, Lodge M, Molofsky J, With KA, Baughman S, Cabin

RJ, Cohen JE, Ellstrand NC, et al. The population biology of invasive species. Annu Rev

Ecol Syst. 2001;35:305–32.

40. Lagos ME, White CR, Marshall DJ. Do invasive species live faster? Mass-specific

metabolic rate depends on growth form and invasion status. Funct Ecol. 2017;31:2080–6.

41. Park T, Miller V, Lutherman CZ. Studies in Population Physiology. IX. The Effect of Imago Population Density on the Duration of the Larval and Pupal Stages of *Tribolium confusum* Duval. Ecology. 1939;20(3):365–73.

42. Janus MC. Phenotypic diversity of *Tribolium confusum* pupae in heterogeneous environments. Entomol Exp Appl. 1989;50:281–6.

43. Halliday WD, Slevan-Tremblay I, Blouin-Demers G. Do Female Red Flour Beetles Assess both Current and Future Competition during Oviposition? J Insect Behav. 2019;32(3):181–7.

44. Awmack CS, Leather SR. Host Plant Quality and Fecundity in Herbivorous Insects. Annu Rev Entomol. 2002;47:817–44.

45. Sonleitner FJ. Factors Affecting Egg Cannibalism and Fecundity in Populations of Adult *Tribolium castaneum* Herbst. Physiol Zool. 1961;34(3):233–55.

46. Costantino RF, Desharnais RA. Population Dynamics and the *Tribolium* Model: Genetics and Demography (volume 13). 2012 Springer Science and Business Media.

47. Reading CJ, Clark RT. The impact of density, rainfall and environmental temperature on body condition and fecundity in the common toad, *Bufo bufo*. Oecologia. 2010;102:453–9.

48. Forchhammer MC, Clutton-Brock TH, Lindström J, Albon SD. Climate and population density induce long-term cohort variation in a northern ungulate. J Anim Ecol. 2001;70(5):721–9.

49. Stewart KM, Bowyer RT, Dick BL, Johnson BK, Kie JG. Density-dependent effects on physical condition and reproduction in North American elk: An experimental test. Oecologia. 2005;143(1):85–93.

50. Ramsay JM, Feist GW, Varga ZM, Westerfield M, Kent ML, Schreck CB. Whole-body cortisol is an indicator of crowding stress in adult zebrafish, *Danio rerio*. Aquaculture. 2006;258:565–74.

51. Assie LK, Brostaux Y, Haubruge E. Density-dependent reproductive success in *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). J Stored Prod Res. 2008;44(3):285–9.

52. Boonstra R, Hik D, Singleton GR, Tinnikov A. The impact of predator-induced stress on the snowshoe hare cycle. Ecol Monogr. 1998;68(3):371–94.

53. Van Allen BG, Rudolf VHW. Ghosts of Habitats Past: Environmental Carry-Over Effects
Drive Population Dynamics in Novel Habitat. Am Nat. 2013 May;181(5):596–608.

54. Burgess SC, Marshall DJ. Adaptive parental effects: the importance of estimating
environmental predictability and offspring fitness appropriately. Oikos. 2014;123:769–
76.

55. del Campo ML, Miles CI, Schroeder FC, Mueller C, Booker R, Renwick JA. Host
recognition by the tobacco hornworm is mediated by a host plant compound. Nature.
2001;411(May):186-9.

56. Henniges-Janssen K, Schofl G, Reineke A, Heckel DG, Groot AT. Oviposition of
diamondback moth in the presence and absence of a novel host plant. Bull Entomol Res.
2011; 101(1):99-105.

57. Soler R, Pineda A, Li Y, Ponzio C, van Loon JJA, Weldegergis BT, Dicke M. Neonates
know better than their mothers when selecting a host plant. Oikos. 2012;121(12):1923–
34.

58. Gretes WC, Stanwyck EA, Hanson FE. Innate and acquired components of oligophagy in
the herbivorous lepidopteran, *Manduca sexta*. Entomol Exp Appl. 2016;160(3):259–71.

59. Dethier V. G. Evolution of Feeding Preferences in Phytophagous Insects. Evolution.
1954;8(1):33–54.

60. Butlin R. A new approach to sympatric speciation. Trends Ecol Evol. 1987;2(10):310–1.

61. Feder JL, Forbes AA. Host Fruit-Odor Discrimination and Sympatric Host-Race
Formation. In : Specialization, Speciation and Radiation : The Evolutionary Biology of
Herbivorous Insects. {ed. Tilmon, K.J.}. 2008. University of California Press, 101-112.

62. Tabashnik BE, Wheelock H, Rainbolt JD, Watt WB. Individual Variation in the
Oviposition Preference in the Butterfly, *Colias eurytheme*. Oecologia. 1981;50(2):225–
30.

63. Singer MC. Determinants of Multiple Host Use by a Phytophagous Insect Population.
Evolution. 1983;37(2):389–403.

64. Agosta SJ. On ecological fitting, plant-insect associations, herbivore host shifts, and host
plant selection. Oikos. 2006;114(3):556–65.

65. Alkema JT, Dicke M, Wertheim B. Context-Dependence and the Development of Push-
Pull Approaches for Integrated Management of *Drosophila suzukii*. Insects. 2019;10:1–

19.

66. Hale R, Swearer SE. Ecological traps: current evidence and future directions. Proc R Soc
B. 2016;283(1824): 20152647

67. Singer MC, Parmesan C. Lethal trap created by adaptive evolutionary response to an
exotic resource. Nature. 2018;557:238–41.

68. Ravi Kumar V, Agavekar G, Agashe D. Fitness Landscapes Reveal Context-Dependent
Benefits of Oviposition Choice. 2021.
bioRxiv 2021.05.27.445916; doi: https://doi.org/10.1101/2021.05.27.445916

Figure 1. Experimental design. (A) Overview of key experiments (B) Details for experiment 2,

used to generate fitness landscapes as a function of female fecundity and egg allocation across

resource patches (W = wheat flour, FM = finger millet flour).

593
594

Figure 2. Oviposition behaviour of females from different prior contexts (Experiment 1). (A) Total fecundity and (B) % Eggs laid in finger millet (FM) in a two-patch habitat (LD: low density, HD: high density). Boxplots are coloured by female context; darker shades of a given colour represent the same set of females after ageing. Sample sizes are noted below boxes in panel A. Letters above boxes indicate results of a binomial generalized linear model, such that boxes with the same letter are not significantly different. (C) Correlation between total fecundity and preference for finger millet for females from each context (for aged LD females, Spearman's rho = 0.315; ns = correlation not significant); lines indicate best-fit linear regressions for each context. Each point represents data for a single female; large filled circles indicate the median fecundity and preference for females from a given context; points and lines are coloured by context, as in panels A and B.

Figure 3. Complete fitness landscapes predict the fitness consequences of female oviposition behaviour in diverse contexts. Linear relationship between offspring development rate and (A) maternal fecundity and (B) total offspring (colour scheme as in Figure 2; n = 96 for young LD; n = 92 for young HD; n = 80 for old HD). Each point indicates data for a single female. (C-E) Measured fitness values (indicated by the z-axis colour bar on the right) for individual females for each egg allocation treatment (Experiment 2; n = 16-20 per egg allocation) spanning their natural fecundity range. Points are slightly jittered along the y axis for visualization. (F–H) Fitness landscapes generated for each fitness component using loess smoothing on data shown in (C-E). For validation, landscapes are overlaid with data from an independent experiment (Experiment 3) where we measured both oviposition preference (which determines the position of each triangle in the landscape) and offspring fitness (which determines the colour fill of each triangle). A close match in the colour of the triangle and its background indicates high predictability of fitness consequences (also see Table S1). (I–K) The same fitness landscapes overlaid with median observed oviposition behaviours for females from different contexts (obtained from Experiment 1, Figure 2), indicated by black filled circles. Lines connect cases where the same set of females were aged (indicated by "A"). The position of each circle shows the predicted fitness values for females from a given context.

625
626

Figure 4. Illustration of the proposed general linkages between female context, oviposition

behaviour and fitness. Young females that experience low competition (as a function of low

population density; bottom left) may reproduce and establish a population as they age, increasing

population density and competition (top right). Further reproduction will generate young females

that also experience high competition (top left). Density-dependent dispersal may then expose

females to low competition while colonizing a new habitat (bottom left). Old females may also

disperse away from high density conditions (bottom right), though we do not have data on this

aspect (indicated by dashed line).