Pathways of savannization in a mesic African savanna-forest mosaic

Heath Beckett[1,2,*], A. Carla Staver[3], Tristan Charles-Dominique[4] and William J. Bond[1]

[1]University of Cape Town, Department of Biological Sciences, Rondebosch 7701, Cape

Town, South Africa.

[2]Global Change Biology Group, Department Botany & Zoology, Stellenbosch University,

Stellenbosch, South Africa.

[3]Ecology and Evolutionary Biology, Yale University, New Haven, CT 06511.

[4]CNRS UMR7618; Sorbonne University; Institute of Ecology and Environmental Sciences

Paris; 4, place Jussieu 75005 PARIS.

E-mail addresses: heath.beckett@uct.ac.za, carla.staver@yale.edu, tristan.charles-

dominique@sorbonne-universite.fr, william.bond@uct.ac.za.

Running Title: Pathways of forest savannization

Keywords: catastrophic regime shifts, Alternative Stable States, savanna, forest, extreme fire

events, savannization

Type of Article: Research Article

*Corresponding author: Heath Beckett, HW Pearson Building, University Ave N,

Rondebosch, Cape Town, 7701, (+27) 72 111 7035, heath.beckett@uct.ac.za

Abstract

Typically, savannas experience frequent fires, which limit tree cover and promote

flammable grass accumulation, whereas forests and thickets form dense canopies that

exclude fires by reducing C4-grass fuel loads and creating a humid microclimate. However,

extreme fires occasionally burn into these closed canopy systems. Although these are

known to cause tree mortality and can make repeat fires more likely, the long-term

consequences of an extreme fire event for closed canopy vegetation structure and potential

savannization remain largely unknown. Here, we analysed whether an extreme fire event

could, alone, alter species composition, vegetation structure, and fire regimes of closed

canopy ecosystems in a natural savanna-forest-thicket mosaic, or whether successive fires

after an initial extreme fire event were necessary to trigger a biome transition between

these alternative stable states. We found that forests that only burned once recovered,

whereas those that burned again following an initial extreme fire event underwent a

transition from closed-canopy forests to open, grassy systems. Thickets had lower levels of

tree mortality following fires than forests, although repeat fires also led to a transition

towards open grassy systems, structurally resembling savannas. Our results suggest that

while extreme fires set the stage for a biome transition, subsequent fires are necessary for

grass composition to resemble neighbouring savanna grass communities, ultimately

resulting in a savanna fire regime. Savannization, following extreme fire events, is possible

in natural mosaic ecosystems where open and closed canopy vegetation types represent

alternative stable states.

Introduction

Fires are becoming more frequent and severe in forested ecosystems globally (Andela *et al.*,

2019, Jolly *et al.*, 2015). In tropical forests, extreme fires associated with droughts and

agriculture are expanding into forests that historically were not flammable (Chen *et al.*,

2014, Morton *et al.*, 2013). Some authors have hypothesized that these tropical forest

understory fires may result in runaway feedbacks, leading to more frequent fires and to

savannization of forests, resulting in eventual forest collapse (Cochrane, 1999, Barlow &

Peres, 2004, Barlow & Peres, 2008, Silverio et al, 2013, Flores *et al*, 2016, van Nes et al,

2018). However, although we know that the occurrence of one forest understory fire can

increase the likelihood of subsequent fires, direct evidence for forest savannization by

runaway feedbacks is sparse. Invasive exotic grasses have been shown to invade the forest

understory only locally (Silverio *et al.*, 2013), limited by dispersal (although in

anthropogenically modified environments there is evidence for seed dispersal by trucks (see

Veldman and Putz, 2010)) and possible biogeochemical feedbacks. As a result, the question

of whether fires can actually convert a tropical forest to a savanna, in a natural setting,

remains open.

Historically and today, fires burn frequently in savannas, at a rate of several per decade, but

rarely penetrate beyond intact forest margins (Kellman & Meave, 1997, Biddulph &

Kellman, 1998, Hennenberg *et al.*, 2008). In some places, this can allow the persistence of a

forest-savanna mosaic, despite the potentially destructive impact of fire on forest trees

(Cochrane & Laurance, 2002, Brando *et al*, 2011, Brando *et al*, 2014), by feedbacks that are

spatially localized (Favier *et al.*, 2004, Beckett & Bond, 2019). Mechanistically, C4-grass-

fuelled fires maintain an open canopy by preventing the establishment of fire-sensitive

forest saplings, but fires stop readily at the savanna-forest boundary, because shade in the

forest understory prevents the accumulation of shade-intolerant C4 grasses within meters

of the boundary (Charles-Dominique *et al.*, 2018, Hennenberg *et al.*, 2008, Hoffmann *et al.*,

2012a). Forest microclimates also reduce wind speeds and increase fuel moisture (Biddulph

& Kellman, 1998, Hoffmann *et al.*, 2012b, Little *et al.*, 2012), further hampering fire spread.

These feedbacks provide support for the idea that savanna and forest are alternative stable

states (Staver *et al.*, 2011a, but see Veenendaal et al, 2015, Hanan et al, 2014). However,

the impact of extreme events (especially extreme fires, see Keeley & Pausas, 2019) on the

dynamics of these mosaics is largely unknown.

One possibility is that mosaics are mostly stable, with distinct distributions and stable edges,

not just on short to medium time scales (ten to several hundred years), but also on longer

time scales (Killeen *et al,* 2006, Breman et al, 2012, Rull *et* al, 2013, Cardoso *et al,* 2020).

This might be the case if savanna-fire feedbacks are very strong (Beckage et al, 2009, Dantas

et al, 2016), or if some other process besides fire – e.g., oligotrophy, seasonal flooding – play

a strong role in stabilizing savanna-forest distributions within a mosaic (Veenendaal et al,

2015, Murphy and Bowman, 2012). In this situation, extreme fires in forests may have fairly

negligible medium-term effects on forest structure, acting as a stand-replacing disturbance

from which forests recover via succession (Fearnside, 1990, Kennard, 2002, Brando et al,

2020) rather than as a trigger of a catastrophic transition.

An alternative possibility is that savanna-forest boundaries may be highly dynamic over

longer time-scales (Schwartz *et al.*, 1986, West *et al.*, 2000, Staver *et* al, 2017, Aleman et al

2019) and in response to changing climate or climate events (Hirota *el al,* 2010, Breman et

al, 2012, Beckett & Bond, 2019, Wright et al, 2021, Sato *et al,* 2021). Spatially explicit

modelling work has argued that fires at the savanna-forest boundary can have major effects

on the biogeographic distribution of biomes (Staver *et al,* 2011b, van de Leemput *et al,*

2015, Goel *et al,* 2020a, Goel *et al,* 2020b, Wuyts *et al*, 2019) and local observations confirm

that boundaries are sometimes dynamic through time (Brook & Bowman, 2006, Silva *et* al,

2008, Ibanez *et al,* 2013, Beckett & Bond, 2019). Modern observations also confirm that

tropical forests can invade savannas and grasslands especially at the boundary between the

two, in Africa (Baccini et al., 2017; Aleman et al., 2018), South America, and Australia

(Marimon et al., 2014; Ondei et al., 2017; Stevens et al., 2017; Rosan et al., 2019).

However, modern evaluations of invasions of forests by savannas has predominantly been

studied in the Neotropics (Cavelier et al, 1998, Veldman and Putz, 2011, Silverio *et al.*, 2013,

Flores et al 2016), typically in the context of invasive exotic grasses, El Niño-fuelled

droughts, and deforestation (Nepstad *et* al. 1996, Barlow and Peres, 2004, Barlow and

Peres, 2008, Veldman and Putz, 2011, Silverio *et al.*, 2013). This work provides invaluable

insights into the potential processes of savannization, such as the appearance of grass-

dominated systems which differ from old growth savannas (Veldman and Putz, 2011), the

susceptibility of forest trees to fire induced mortality (Barlow and Peres, 2008, Staver *et al,*

2020), and the invasion of grassy fuels following repeat fires (Veldman and Putz, 2011,

Silverio *et al*, 2013). However, these processes typically rely upon the introduction of exotic

grass species (which lead to novel fire regimes), and deforestation derived boundaries

(Alencar *et al*. 2015) with unclear historical and palaeoecological precedents (but see Flores

et al 2016).

From its conception in the 1940's (Aubreville, 1949, Budowski, 1956), the term

savannization has been used to refer to fire induced degradation of tropical forest subject to

anthropogenic pressures such as farming and deforestation (Nyerges, 1990, Uhl *et al*, 1982,

Cochrane, 1999, Nepstad *et al*, 1999). The term rose to prominence following predictions of

large-scale Amazonian Forest die off under elevated CO_2 (Cox *et al*, 2004) and is widely used

in the contemporary literature to refer to a reduction of tree biomass and species

associated with forest ecosystems. In doing so, the savanna biome is – unintentionally -

associated with degraded, species-poor forests and identified as an early successional stage

of forest vegetation (Veldman, 2016, Bond, 2016, Schmidt *et al*, 2019). The term

savannization should be used to refer to the conversion of one vegetation state (typically

forests) into a savanna and the establishment of the processes regulating savannas, which is

possible in regions of the world where savannas and forests represent alternative stable

states (Staver & Levin, 2012, Archibald *et al*, 2019, Aleman *et al*, 2020).

Perpetuating the usage of the term to refer to anthropogenic degradation undermines the

value and antiquity of the savanna biome and reinforces a widely held – yet critically

incorrect – belief that savannas are an anthropogenically derived vegetation type (Veldman

et al, 2015a, Veldman, 2016). As such, we believe the question of whether extreme fires can

cause a biome switch between forests and savannas in a natural setting has not yet been

settled but is increasingly relevant in an age of climate extremes. Vegetation mosaics of

alternative stable states offer a rare opportunity for testing the concept in a natural setting.

This question has both theoretical and practical importance. From the perspective of

understanding ecological processes, we have no clear understanding of the possible role of

extreme fires in biogeographic transitions between savanna and forest (*e.g.*, Aleman et al

2019) and in shaping transitions in response to changing climate (Beckett & Bond 2019).

From a practical perspective, recent decades have seen significant increases in deforestation

and fragmentation of intact tropical forests (Watson et al, 2018; Taubert et al, 2018). The

most commonly proposed solution to this is to plant trees in order to restore fragmented

landscapes (e.g., the Bonn Challenge, REDD+; Veldman et al, 2015b), but this can be harmful

to the natural, biodiverse, open systems found in these mosaics (Bond and Parr, 2010,

Veldman et al, 2015b, Ratnam et al, 2011). Understanding the stability and dynamics of

savanna-forest mosaics would help to justify and inform the management of these systems

as mosaics, with a focus on the preservation of savannas as well as forests (Bond & Parr,

2010).

Here, we have examined vegetation trajectories after extreme fire events empirically,

focusing on a mesic savanna-forest-thicket mosaic in Hluhluwe iMfolozi Park, South Africa,

where extreme fires burned into forests and thickets in 2004 and 2008; some of these areas

have burned subsequently, while others have not. These changing fire regimes offer an

opportunity to examine *in situ* the spectrum of possible outcomes for closed canopy

vegetation trajectories. On the one extreme, a forest or thicket experiencing fire might

recover its original state (Forest to Forest (FtoF)). In this case, we might expect to see FtoF

1) a reduction in tree cover and biomass that is only temporary, FtoF 2) the inhibition of a

regular fire regime (e.g. by suppression of grassy fuels), and FtoF 3) a transient shift to early-

successional forest trees species (light-loving but not as fire tolerant as savanna species)

that eventually transitions back to the original forest state. On the other extreme, in a forest

switching to savanna (Forest to Savanna (FtoS)), we instead expect: FtoS 1) a reduction in

tree cover and biomass that persists through time, FtoS 2) the establishment of a regular

fire regime (or at least of sufficient grassy biomass), and FtoS 3) a shift in tree species

composition from shade-tolerant species that compete for light to shade-intolerant species

that require open environments and invest in fire protection. Differentiating between these

possibilities provides us the rare opportunity to test the forest savannization hypothesis

developed for the Amazon (Cochrane et al 1999), allowing us to investigate what happens in

the aftermath of extreme disturbances in forests and thickets.

Materials and Methods

The northern section of Hluhluwe-iMfolozi Park (HiP) receives sufficient annual rainfall

(approx. 1000mm per annum) (Balfour & Howison, 2002) to support a closed canopy forest,

with vegetation dynamics that are driven by fire compared to the more arid (approx.

600mm per annum), herbivore-driven system in the iMfolozi section (Staver *et al.*, 2012).

Closed canopy vegetation states in HiP are differentiated into scarp forests (which we refer

to as forests here onwards) and thicket (*sensu* Charles-Dominique et al 2015a). Forest,

thicket, and savanna definitions are based on the classification scheme used in Charles-

Dominique et al (2015a), using vegetation structure and associated growth forms

(Woodward et al, 2004; Ratnam et al, 2011). In HiP, forests are classified as tall (>10m)

woody vegetation with a shade-tolerant intermediate tree layer, lacking a C4 grass layer.

Occasionally, one finds patches of C3 grasses and herbaceous plants among the litter layer

(Charles-Dominique *et al*, 2015). Thickets are classified as having dense shrub and treelet

vegetation, a canopy that is generally 4 to 6m tall, and a variable understory with dense

patches of herbaceous sub-shrubs, shrubs and occasional patches of shade-tolerant grasses

(Charles-Dominique *et al*, 2015) and savannas are classified as having discontinuous tree

cover with a continuous layer of C4 grasses. Forest and thicket patches naturally abut

savannas with a full complement of savanna herbivores, including elephant and black rhino.

As in most of southern Africa, fires are usually lit by people, pre-empting lightning fires

which generally occur near the end of the dry season (Archibald *et al.*, 2017), and humans

have been burning fires in the region for at least 150 ka with Iron Age farmers arriving in the

area ~ 2 ka (Hall, 1980, Staver *et al.*, 2017). The Hluhluwe section of HiP experiences a

relatively frequent fire regime with a mean and median fire return interval of 2.9 and 1.3

years, respectively (Balfour and Midgley, 2008).

On the 14^th and 15^th of September 2008, an extreme fire event occurred in the Hluhluwe

section of HiP and, unlike typical savanna fires, it burned through large tracts of thicket and

forest. Aerial photography of the region from 1937 to 2013 shows a gradual increase in

closed canopy patch sizes from 1937 to 1992, and major loss events from 1992 to 2013

(Beckett & Bond, 2019), associated with this fire event. These extreme fires resulted from a

combination of air temperatures above 30°C, relative humidity below 30%, and wind speeds

averaging more than 25km/hr with gusts to 30km/h or more (Bradstock et al 2009; Browne

& Bond, 2011). In this region, these conditions are exceedingly rare, over the period of 2001

to 2008, less than three days of these extreme fire weather conditions occurred, with the

longest consecutive record of 11 hours occurring on the day of the extreme fire event in

2008 (Browne & Bond, 2011). Some areas that experienced this extreme fire burned

subsequently, whereas others did not (see below for further detail, Appendix S1).

Vegetation sampling

We used a set of plots in which the structure and species composition of woody plants were

recorded in forest, thicket and savanna vegetation types. A total of 131 plots were sampled

(Fig. 1), 51 of which were first sampled in 2007 by Staver *et al (2012)* and resampled in 2014

by Case and Staver (2017), 31 of which were sampled once in 2014 by Charles-Dominique *et*

al (2015a), and an additional 49 which were established specifically for this study between

2009 and 2015. Staver et al (2012) sampled 253 plots across HiP, 51 of these were chosen as

the savanna baselines as they burnt in the 2008 fire and were near forest and thicket

patches. Intact forest and thicket plots were laid out in mature unburnt vegetation patches

(Fig. 1), as mapped by Whateley and Porter (1983) and by aerial-photograph-derived

vegetation maps (Beckett & Bond, *2019*). These represent our pre-fire baselines for forest

and thicket communities and, along with the 2007 savanna plots, are referred to from here

on as our 'before' dataset, while burnt plots represent the 'after' dataset.

To determine vegetation trajectories after the extreme fires, we made use of a natural

experiment, selecting plot locations based on areas that burned in extreme fire events in

2004 and 2008. Some plots were sampled repeatedly, supplemented with once-sampled

plots allowing a space-for-time chronosequence to evaluate vegetation change. In 2013,

Beckett and Bond (2019) identified regions previously unknown to have burnt in the 2008

fire, we established plots in these areas targeting forest patches that had not burnt since

2008, and others that had burnt repeatedly. Three plots were laid out in a forest that

burned in a separate fire in 2004 and sampled nine and 11 years after the fire. These three

plots (with 20m buffers on all sides) covered approximately 80% of the burnt forest area.

One of these plots was burnt again in 2012 and was excluded from the 11 year resampling

as it had no analogues. The data from nine years after the fire are included in the analyses,

while the repeat sample of 11 years are only included in figures in the Supplementary

Materials for interest. Plot sampling dates, repeat sample dates and vegetation history are

summarized in Appendix S1.

Each plot measured 40m x 10m, within which area we identified every woody plant

(including lianas) above 50cm in height and recorded its height as one of four size classes

(0.5-2m, 2-5m, 5-10m, and >10m). In 103 of the 134 sites, the grass layer was evaluated at

20 points along the plot; grass biomass was measured via disk pasture meter (DPM) and the

dominant grass species recorded beneath the disk. A DPM is a weighted metal disk of a

known diameter, which is dropped on a grass sward from a standard height. The resting

height is a measure of the grass biomass under the disk, which can be calibrated locally

(Bransby and Tainton, 1977, grass biomass = 1.26 + 26.1*[DPM], R2 = 0.73, N = 1745;

Waldram et al, 2007).

Fire occurrence

Fire frequency of each plot from 2001 to 2016 was derived from a combination of the

Ezemvelo KwaZulu Natal Wildlife fire records for HiP and personal observations. The EKZNW

fire records are hand drawn maps from rangers responsible for sections of the park. These

indicate the general location of the fire, but do not include information on fine-scale fire

refugia and unburnt vegetation patches within the burned area (see Fig. 1). Instances where

the EKZNW fire records indicated fires burning into large forest patches were compared

with MODIS Active Fire Detections (MCD14DL, Collection 6). MODIS Active Fires maps

thermal anomalies within 1km pixels using the MODIS MOD14/MYD14 Fire and Thermal

Anomalies algorithm (Giglio *et al.*, 2003). MODIS Active Fires increases the probability that

fires within forests are detected since alternative approaches based on burn scar mapping

are sensitive to obstruction from the forest canopy.

Data analysis and statistics

Aerial photography and vegetation maps were analysed using Quantum GIS 3.0.0 (Quantum

GIS Development Team, 2018). Statistical analyses were performed in R (version 3.4.0, R.

Core Team, 2017), using the packages *vegan* (Oksanen *et al.*, 2013) and *tidyverse* (Wickham,

2017). Fire frequency before and after extreme fire events was compared using a Wilcoxon

signed-rank test for non-parametric paired data. Grass biomass was estimated using the

relationship published by Waldram *et al* (2007) and comparisons between vegetation types

and time periods were investigated using a Wilcoxon signed-rank test for non-parametric

paired data. We used a multivariate analysis of variance (MANOVA) to test for differences in

the total mean cumulative length of the vegetation types at different time periods and

followed this with a series of ANOVA models to test for significant differences between

vegetation types within the size classes. Cumulative length is a composite measure of

species importance, similar to basal area, which combines stem density (values reported in

Appendix S2) and tree height into a single value (Fei *et al.*, 2005). This is done by combining

the height of all individuals of a species and allows for a weighted index where large trees

contribute more to the overall score than small trees. We performed an NMDS Ordination

on a Bray-Curtis species dissimilarity matrix of the cumulative length of each tree species for

the 'before' plots, representing vegetation communities which that had not experienced the

extreme fire event, namely, the intact forest, intact thicket, and 2007 savanna plots. Data

were transformed using a Hellinger transformation to account for large abundances

(Legendre & Legendre, 2012). Species were assigned to a biome (forest, thicket, or savanna)

based on their association with sites in the NMDS ordination; species not found before the

extreme fire event were classified as pioneer species. We then performed a Principal

Coordinates Analysis (PCoA) and a pairwise permutational MANOVA on a Bray-Curtis

dissimilarity matrix of the total cumulative length in each plot, as well as within the size

classes to identify changes in the composition of these species types.

Results

Establishment of a savanna fire regime

Forests and thickets rarely burned, with intact forest and thicket sites experiencing no fires

between 2000 and 2016 (Fig. 2). Savanna sites, in contrast, burned several times in a

decade. In fact, fire frequency (FF) increased from approximately one fire every four years

(FF = 0.24.yr^{-1} (SD = 0.11)) before the 2008 extreme fire event to one fire every three years

(FF = 0.36.yr^{-1} (SD = 0.13); t = -11.8, df = 43, p < 0.05) after 2008. Thicket patches that

burned in the 2008 extreme fire event quickly established a savanna fire regime with sites

burning on average once every three years (FF = 0.34.yr^{-1} (SD = 0.07)) , compared to no fires

from 2000 to 2007.

Some forests that burned in the extreme fire events burned again, whereas others did not.

To investigate how contrasting fire returns affected their post-extreme-fire trajectories, we

split plots into two groups, those that burned only once (hereafter "once-burnt forests")

versus those that burned in an extreme fire event and at least once subsequently (hereafter

"frequently-burnt forests"). Frequently-burnt forests burned on average once every four

years since the extreme fire event (FF = 0.26.yr^{-1} (SD = 0.04) , Fig. 2).

Colonisation by savanna grasses

Intact thicket and forest plots differed from savanna plots in having closed canopies and

little or no grass in the understory (Fig. 3). Savanna plots had on average 233g.m^{-2} (SD =

87)of grass biomass whereas intact thicket and forest patches had 148g.m^{-2} (SD = 84) and

10g.m^{-2} (SD = 2.8) respectively (Fig. 3). Frequently-burnt thicket and frequently-burnt forest

plots showed an increase in grass biomass following successive fires, with grass biomass

four and 22 times higher than intact thicket and forest plots respectively (Fig. 3). Once-burnt

forest plots had relatively high, but variable, grass biomass (130g.m^{-2} (SD = 114)) in the first

year after the extreme fire events, grass biomass by the end of the study, 9 years later, was

less variable (33g.m^{-2} (SD = 11)), approaching the levels of intact forest grass biomass. In

terms of grass species, *Panicum maximum* colonised forests and thickets immediately after

fire, whereas *Themeda triandra* increased in abundance where fires occurred repeatedly

(Fig. 3).

Trajectories in woody structure

Forest, thicket, and savanna plots displayed distinct woody structural profiles (See Appendix

S3), with most savanna trees in the smallest size class, most forest trees in the largest size

class, and an even spread between the two intermediate size classes in thicket (Fig. 4,

Appendix S3). Savanna plots showed little change in structure between 2007 and 2014

despite an increase in fire frequency (Pillai's Trace = 0.039, approx $F_{(3,91)}$ = 1.238, p = 0.301,

Appendix S3).

Thicket plots that burned frequently were different to Intact thicket plots (Pillai's Trace =

0.76, approx $F_{(4,30)}$ = 23.406, p < 0.001, Appendix S3) and had a comparable structure to

savanna plots seven years after the extreme fire (Pillai's Trace = 0.15, approx $F_{(4,51)}$ =

2.2415, p = 0.077, Appendix S3), with most trees in the smallest size class. Similarly, forests

that burned multiple times were different to intact forests (Pillai's Trace = 0.89, approx

$F_{(4,28)}$ = 58.734, p < 0.001) and converged to a savanna-like vegetation structure (Pillai's

Trace = 0.20, approx $F_{(4,53)}$ = 3.3665, p = 0.016) 7 years after the extreme fire event, with

comparable numbers of trees in the smaller two size classes and differences in the larger

two size classes (Appendix S3).

Six years after the fire, once-burnt forests were different to intact forests (Pillai's Trace =

0.902, approx $F_{(4,17)}$ = 38.902, p < 0.001) and savannas (Pillai's Trace = 0.77, approx $F_{(3,43)}$

= 46.72, p < 0.001). The six year once-burnt and the frequently-burnt forests were also

different (Pillai's Trace = 0.61, approx $F_{(4,12)}$ = 4.72, p = 0.016), with the second largest size

class contributing to the majority of the difference between the two ($F_{(1,15)}$ = 21.96, p <

0.001). Nine years after the fire, the relationships between once-burnt forests and intact

forests (Pillai's Trace = 0.761, approx $F_{(4,17)}$ = 13.499, p < 0.0001), savannas (Pillai's Trace =

0.767, approx $F_{(3,43)}$ = 47.296, p < 0.0001) and frequently-burnt forests (Pillai's Trace =

0.716, approx $F_{(4,12)}$ = 7.5689, p = 0.003) were similar to those of the six year once-burnt

forests (Appendix S3, Appendix Fig. S1), however the cumulative length in the second

largest size class of once-burnt forests was approaching that of the intact forests ($F_{(1,20)}$ =

3.641, p = 0.071).

Trajectories in woody species composition

Forest, thicket, and savanna plots were characterized by distinct woody species assemblages

(Fig. 5, Appendix Table S4 & Table S5The overall (Appendix Table S4) and size class specific

(Appendix Table S5) abundance of species types within savannas did not differ between

2007 and 2014 (Fig. 6).

Compositionally, frequently-burnt thickets differed from savannas 7 years after the extreme

fire event (Fig. 6, Appendix Table S4 & S5), this is most evident in the smallest size class

where frequently-burnt thickets were dominated by a mixture of basally resprouting thicket

species, *e.g.*, *Euclea racemosa*, *Euclea divinorum* and *Berchemia zeyheri*, and a proportion of

savanna species, including *Acacia karroo (=Vachellia karoo)*, *Acacia nilotica (=V. nilotica)*,

Dichrostachys cinerea, Cordia caffra, Combretum molle and *Rhus (= Searsia) pentheri* (Fig. 5),

compared to savannas which very rarely had thicket species recruiting.

By contrast, seven years after the extreme fire event, the composition of frequently-burnt

forests had completely switched to dominance by a number of thicket species, including

Euclea natalensis and *Euclea racemosa,* and savanna species (see Fig. 5 & 6). These plots

also differed from savannas (Appendix Table S4 & S5) in the composition of species in the

smallest size class and more closely resembled the frequently-burnt thickets.

Once-burnt forests were dominated by pioneer forest species seldom found in the intact

forests (especially *Trema orientalis, Croton sylvaticus,* and *Dombeya burgessiae*). These

trees grew from seed banks, as opposed to the dominance of basally resprouting tree

species in frequently-burnt thicket plots. After 6 years, these once-burnt forests recovered

some species characteristic of mature forest, including *Englerophytum natalense* and

Chaetacme aristata, in smaller size classes, although the larger size classes remained

dominated by the unique pioneer forest assemblage (Fig. 6). The once-burnt forests (after

six years) were not different to the frequently-burnt forests in overall abundance of species

types (Appendix Table S4), however they were different when considering size class specific

species type abundance (Appendix Table S5). Nine years after fire, the once-burnt forests

were dominated by forest species in the two smaller size classes, and pioneer species in the

second largest size class (Appendix Fig. S2). These plots differed to the frequently-burnt

forests in overall and size class specific species type abundance (Appendix Table S4 & S5, Fig.

S3.).

Discussion

Extreme fire events differ from typical savanna fires in penetrating beyond the biome

boundaries, often opening up the canopy of closed forest or thicket. Here, we found that

forests that burned only once recovered by secondary successional pathways, whereas

forests and thickets that burned again following the initial extreme fire event underwent a

persistent transition from closed-canopy vegetation states to open-canopy grassy systems.

The grass layer was first dominated by *Panicum maximum* but gradually replaced by the

more flammable *Themeda triandra* (Simpson et al, 2016). Woody species composition was

the slowest to respond to woody structural change, and we observed only minimal

colonization of woody savanna species into the new grass layer over a decade of post-burn

observations.

Together, these results suggest that extreme fires, when followed by subsequent fires, can

trigger a biome switch from closed, fire-suppressing vegetation state to open, fire-

dependent vegetation, structurally resembling savannas. This transition occurred on the

time scale of a decade, which suggests that savanna-forest-thicket mosaics are likely

dynamic even in short-to-medium term (Killeen et al, 2006, Rull *et al*, 2013, Beckett and

Bond, 2019). The occurrence of subsequent fires was crucial for maintaining an open

canopy and allowing flammable grasses to colonise (see also Balch *et al.*, 2009, Silverio *et*

al., 2013), ultimately determining whether a biome shift from forest or thicket to savanna

occurred.

In the transition from closed canopy vegetation states to savanna, initial grass colonization

was dominated by *Panicum maximum*, which accumulated biomass rapidly. However, with

seven years of repeated fires, *Themeda triandra* gained in dominance in open-canopy

systems. These replacement patterns are consistent with our understanding of these two

species; *Panicum maximum* (Paniceae) is often observed in relatively shaded savanna

environments and is known to retain moisture longer into the dry season (Simpson et al,

2016), whereas *Themeda triandra* (Andropogoneae) prefers open, full-sun environments

(Downing & Marshall, 1980, Kinyamario *et al.*, 1995) and is highly flammable but slower to

colonise new environments (O'Connor & Everson, 1998).

On the time scale of a decade, however, novel savannas differed from old-growth natural

savannas in their woody-layer composition (see also Veldman, 2016), and colonization of

open habitats by savanna trees was slower than grass colonization. While forest canopies

opened up readily, allowing grasses to colonise, the woody community remained

depauperate of savanna trees (see also Veldman & Putz, 2011 and theoretical predictions

described by Staver and Levin, 2012, Touboul et al, 2018). Thickets differed from forests in

that, even when burned repeatedly, they retained a tree community dominated by thicket

species (probably resprouts). Meanwhile, forests that only burned once were colonised

mostly by a unique pioneer assemblage. However, in novel savannas, a frequent fire regime

eventually depleted thicket and forest remnants, and we hypothesize that savanna species

will eventually colonise (see also Charles-Dominique et al, 2015b); we suggest further

observation of previously closed-canopy areas that continue to experience frequent fires.

Notably, we also found that forests could recover readily from a singular extreme fire event,

as long as a fire-free interval of a decade allowed the canopy to close, shading out grass

fuels and setting the system on a trajectory to its previous state. Forests likely recovered

primarily from a seed bank, with an early-successional forest composed of rapidly growing

pioneers (*i.e., Trema orientalis, Croton sylvaticus, Celtis africana*) and the clonally spreading

liana *Dalbergia armata*. These species excluded light-demanding grasses, likely by shading

and microclimate effects that also make the environment suitable for the establishment of

climax forest species (see also Pammenter et al. 1985).

Thickets also recovered from a single fire readily but differed somewhat from forest in their

post-fire recovery. While forest recovery occurred through pioneer species germination and

was accompanied by a complete species turnover, structural recovery in thickets instead

occurred through basal resprouting. Notably, fire-driven tree mortality in thickets was low,

as in savannas, contrasting with much higher tree mortality in forests (see also Ryan &

Williams, 2011, Staver *et al.*, 2020). We predict that, on longer time scales, thickets (and by

extension other dry forests) may be more resilient or tolerant of fires than wetter forests

(Staal et al, 2020, Staver *et al*, 2020).

Our study is a rare documentation of invasion of grass-fuelled fires into closed forest and

thickets communities in a natural and ancient African savanna-forest-thicket mosaic. Our

results offer a few key insights. Firstly, we provide evidence for extreme fires as a

mechanism by which savannas might have invaded forests before the introduction of

invasive exotic species. Fires are implicated in the rise of the savanna biome from the late

Miocene (Bond *et al.*, 2003, Keeley & Rundel, 2005, Scheiter *et al*, 2012, Behrensmeyer &

Freeman, 2018), by opening up forests and allowing shade-intolerant savanna grasses to

spread, perhaps by gradual invasion at the forest edge (Schertzer & Staver,2018). Fires have

also been implicated in constricting forest distributions in response to millennial scale

changes in climate (Aleman *et al.*, 2019, Sato *et al*, 2021) – probably by increasing forest

canopy openness, then grass invasion, and finally by the arrival of savanna trees. However,

on-the-ground observations of how this savannization process proceeds in a natural

ecosystem are very few.

Secondly, we also provide rare direct support for the longstanding hypothesis that runaway

fire feedbacks, following an initial extreme fire, might plausibly result in the savannization of

tropical forests (Cochrane, 1999) without the need for human interference. As in the

Amazon, these extreme fire events present a rare but real problem. At our site, the

combination of conditions associated with extreme fire development – air temperatures

above 30°C, relative humidity below 30% and wind speeds gusting to 30+km/h (Bradstock *et*

al., 2009) – occurred on only 3 full days over the period between 2001 and 2008, with the

longest consecutive record of 11 hours occurring on the day of the extreme fire event in

2008, on the 15th September. Their rarity makes conditions like these difficult to predict

with any accuracy; however, we can be confident that days like these are likely to increase

in frequency as temperatures rise and aridity increases in the future, making extreme fires

an accelerating concern in savanna-forest mosaics, just as they are elsewhere (Jolly *et al.*,

2015, Flannigan et al, 2013). Transitions from forest to savanna are indeed possible, and

likely to increase in potential extent in future.

The key difference between this study and work published under the savannization

umbrella in the Neotropics revolves around the level of anthropogenic influence in these

systems. Contemporary savannization in the Amazon is recorded in areas where exotic

pasture grasses have invaded degraded forests, increasing fuel loads, and allowing

subsequent fires (Silverio *et al*, 2013). However, paleoecological evidence supports the claim

that savannization has occurred in the Neotropics, predating colonial anthropogenic

activities, and has been attributed to fire and low CO_2 (Rull *et al*, 2013, Sato *et al*, 2021). It is

interesting to note the potential impact the disappearance of Neotropical megafauna may

have had on the probability of savannization occurring under non-anthropogenic conditions

(Owen-Smith, 1987, Johnson, 2009). In HiP, and other African systems, herbivores -

particularly African Elephants – may play a role in weakening forest and thicket boundaries

by opening the canopy and allowing grass invasion. The role of megafauna in savannization

processes has not been sufficiently investigated and offers an interesting avenue for future

research.

Preventing or suppressing extreme fires is challenging in savannas where fires are used as a

management tool under less extreme conditions. In savannas, woody encroachment has

long been a conservation concern, with widespread observations of increases in the woody

component of savannas (Stevens *et al.*, 2017), particularly in savannas where fire is

suppressed instead of actively managed (Silva et al, 2008, Durigan, 2020). This trend is

predicted to intensify as increasing CO_2 favours C3 trees over C4 grasses (Higgins & Scheiter,

2012; Bond & Midgley, 2000, Moncrieff et al, 2014) – suggesting that, while forests are at

increased risk of savannization, savannas are also at increased risk of forest expansion and

woody encroachment. We suggest that widespread fire suppression is clearly not the

solution to preventing extreme fires. We should aim instead for a rationalized fire

management policy, allowing and encouraging fires in fire-dependent savannas but

minimizing fires under extreme conditions, especially in fire-sensitive forests. Most crucially,

we suggest that, barring subsequent fires, which are not inevitable, forests can recover

readily from the effects of extreme fires.

References

Aleman JC, Blarquez O, Elenga H et al. (2019) Palaeo-trajectories of forest savannization in

the southern Congo. Biology letters, 15, 20190284.

Aleman, J. C., & Staver, A. C. (2018). Spatial patterns in the global distributions of savanna

and forest. Global Ecology and Biogeography, 27(7), 792-803.

Aleman, J. C., Fayolle, A., Favier, C., Staver, A. C., Dexter, K. G., Ryan, C. M., ... & Swaine, M.

D. (2020). Floristic evidence for alternative biome states in tropical Africa. Proceedings of

the National Academy of Sciences, 117(45), 28183-28190.

Alencar, A. A., Brando, P. M., Asner, G. P., & Putz, F. E. (2015). Landscape fragmentation,

severe drought, and the new Amazon forest fire regime. Ecological applications, 25(6), 1493-

1505.

Andela N, Morton DC, Giglio L et al. (2019) The Global Fire Atlas of individual fire size,

duration, speed and direction. Earth System Science Data, 11, 529-552.

Archibald S, Beckett H, Bond WJ, Coetsee C, Druce DJ, Staver AC (2017) Interactions

between Fire and Ecosystem Processes. In: Conserving Africa's Mega-Diversity in the

Anthropocene: The Hluhluwe-iMfolozi Park Story. (eds Cromsigt JP, Archibald S, Owen-Smith

N) pp Page.

Archibald, S., Bond, W. J., Hoffmann, W., Lehmann, C., Staver, C., & Stevens, N. (2019).

Distribution and determinants of savannas. Savanna woody plants and large herbivores, 1-

24.

Aubréville, A. (1949). Climats, forêts et désertification de l'Afrique tropicale.

Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., & Houghton, R. A.

(2017). Tropical forests are a net carbon source based on aboveground measurements of

gain and loss. Science, 358(6360), 230-234.

Balch JK, Nepstad DC, Curran LM (2009) Pattern and process: fire-initiated grass invasion at

Amazon transitional forest edges. In: Tropical fire ecology. pp Page., Springer.

Balfour DA, Howison OE (2002) Spatial and temporal variation in a mesic savanna fire

regime: responses to variation in annual rainfall. African Journal of Range & Forage Science,

19, 45-53.

Balfour, D. A., & Midgley, J. J. (2008). A demographic perspective on bush encroachment by

Acacia karroo in Hluhluwe-Imfolozi Park, South Africa. African Journal of Range and Forage

Science, 25(3), 147-151.

Barlow, J., & Peres, C. A. (2004). Ecological responses to El Niño–induced surface fires in

central Brazilian Amazonia: management implications for flammable tropical

forests. Philosophical Transactions of the Royal Society of London. Series B: Biological

Sciences, 359(1443), 367-380.

Barlow, J., & Peres, C. A. (2008). Fire-mediated dieback and compositional cascade in an

Amazonian forest. Philosophical Transactions of the Royal Society B: Biological

Sciences, 363(1498), 1787-1794.

Beckage, B., Platt, W. J., & Gross, L. J. (2009). Vegetation, fire, and feedbacks: a disturbance-

mediated model of savannas. The American Naturalist, 174(6), 805-818.

Beckett H, Bond WJ (2019) Fire refugia facilitate forest and savanna co-existence as

alternative stable states. Journal of Biogeography, 46(12), 2800-2810

Behrensmeyer, A. K., & Freeman, K. H. (2018) Grassland fire ecology has roots in the late

Miocene. Proceedings of the National Academy of Sciences, 115(48), 12130-12135.

Biddulph J, Kellman M (1998) Fuels and fire at savanna-gallery forest boundaries in

southeastern Venezuela. Journal of Tropical Ecology, 14, 445-461.

Bond WJ, Midgley GF, Woodward FI (2003) The importance of low atmospheric CO_2 and fire

in promoting the spread of grasslands and savannas. Global Change Biology, 9, 973-982.

Bond, W. J. (2008). What limits trees in C4 grasslands and savannas?. Annual review of

ecology, evolution, and systematics, 39, 641-659.

Bond, W. J. (2016). Ancient grasslands at risk. Science, 351(6269), 120-122.

Bond, W. J., & Midgley, G. F. (2000). A proposed CO_2-controlled mechanism of woody plant

invasion in grasslands and savannas. Global Change Biology, 6(8), 865-869.

Bond, W. J., & Parr, C. L. (2010). Beyond the forest edge: ecology, diversity and conservation

of the grassy biomes. Biological conservation, 143(10), 2395-2404.

Bradstock RA, Hammill KA, Collins L, Price O (2009) Effects of weather, fuel and terrain on

fire severity in topographically diverse landscapes of south-eastern Australia. Landscape

Ecology, 25, 607-619.

Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., et al. (2014).

Abrupt increases in Amazonian tree mortality due to drought-fire interactions. PNAS,

111(17), 6347–6352.

Brando, P. M., Nepstad, D. C., Balch, J. K., Bolker, B., Christman, M. C., Coe, M., & Putz, F. E.

(2011). Fire-induced tree mortality in a neotropical forest: the roles of bark traits, tree size,

wood density and fire behavior. Global Change Biology, 18(2), 630–641.

Brando, P. M., Soares-Filho, B., Rodrigues, L., Assunção, A., Morton, D., Tuchschneider, D., ...

& Coe, M. T. (2020). The gathering firestorm in southern Amazonia. Science advances, 6(2),

eaay1632.

Bransby, D. I., & Tainton, N. M. (1977). The disc pasture meter: possible applications in

grazing management. Proceedings of the annual congresses of the Grassland Society of

Southern Africa, 12(1), 115-118.

Breman, E., Gillson, L., & Willis, K. (2012). How fire and climate shaped grass-dominated

vegetation and forest mosaics in northern South Africa during past millennia. The Holocene,

22(12), 1427-1439.

Brook, B. W., & Bowman, D. M. (2006). Postcards from the past: charting the landscape-

scale conversion of tropical Australian savanna to closed forest during the 20th century.

Landscape Ecology, 21(8), 1253.

Browne C, Bond W (2011) Firestorms in savanna and forest ecosystems: curse or cure? Veld

& Flora, 97, 62-63.

Budowski, G. 1956. Tropical savannas, a sequence of forest felling and repeated burnings.

Turrialba 6(1-2) :23-33

Cardoso, A. W., Oliveras, I., Abernethy, K. A., Jeffery, K. J., Glover, S., Lehmann, D., ... &

Malhi, Y. (2020). A distinct ecotonal tree community exists at central African forest–savanna

transitions. Journal of Ecology.

Case MF, Staver AC (2017) Fire prevents woody encroachment only at higher-than-historical

frequencies in a South African savanna. Journal of Applied Ecology, 54, 955-962.

Cavelier, J., Aide, T., Santos, C., Eusse, A., & Dupuy, J. (1998). The savannization of moist

forests in the Sierra Nevada de Santa Marta, Colombia. Journal of Biogeography, 25(5), 901-

912.

Charles-Dominique T, Staver AC, Midgley GF, Bond WJ (2015a) Functional differentiation of

biomes in an African savanna/forest mosaic. South African Journal of Botany, 101, 82-90.

Charles-Dominique T, Beckett H, Midgley GF, Bond WJ (2015b) Bud protection: a key trait

for species sorting in a forest-savanna mosaic. New Phytol, 207, 1052-1060.

Charles-Dominique T, Midgley GF, Tomlinson KW, Bond WJ (2018) Steal the light: shade vs

fire adapted vegetation in forest–savanna mosaics. New Phytologist, 218, 1419-1429.

Chen Y, Morton D, Jin Y et al. (2014) Erratum: Long-term trends and interannual variability

of forest, savanna and agricultural fires in South America (Carbon Management (2013) 4: 6

(617-638)).

Cochrane MA (1999) Positive Feedbacks in the Fire Dynamic of Closed Canopy Tropical

Forests. Science, 284, 1832-1835.

Cochrane MA, Laurance WF (2002) Fire as a large-scale edge effect in Amazonian forests.

Journal of Tropical Ecology, 18, 311-325.

Cox, P. M., Betts, R. A., Collins, M., Harris, P. P., Huntingford, C., & Jones, C. D. (2004).

Amazonian forest dieback under climate-carbon cycle projections for the 21st

century. Theoretical and applied climatology, 78(1), 137-156.

Dantas, V. D. L., Hirota, M., Oliveira, R. S., & Pausas, J. G. (2016). Disturbance maintains

alternative biome states. Ecology Letters, 19(1), 12-19.

Downing B, Marshall D (1980) Complementary dominance of Themeda triandra and

Panicum maximum examined through shoot production. Proceedings of the Annual

Congresses of the Grassland Society of Southern Africa, 15, 163-166.

Durigan, G (2020) Zero-fire: not possible nor desirable in the Cerrado of Brazil. Flora,

151612.

Favier C, Chave J, Fabing A, Schwartz D, Dubois MA (2004) Modelling forest–savanna mosaic

dynamics in man-influenced environments: effects of fire, climate and soil heterogeneity.

Ecological Modelling, 171, 85-102.

Fearnside, P. M. (1990). Fire in the tropical rain forest of the Amazon basin. In Fire in the

tropical biota (pp. 106-116). Springer, Berlin, Heidelberg.

Fei S, Gould PJ, Steiner KC, Finley JC, Mcdill ME (2005) Forest regeneration composition and

development in upland, mixed-oak forests. Tree physiology, 25, 1495-1500.

Flannigan M, Cantin AS, De Groot WJ, Wotton M, Newbery A, Gowman LM (2013) Global

wildland fire season severity in the 21st century. Forest Ecology and Management, 294, 54-

61.

Flores, B. M., Fagoaga, R., Nelson, B. W., & Holmgren, M. (2016). Repeated fires trap

Amazonian blackwater floodplains in an open vegetation state. Journal of Applied

Ecology, 53(5), 1597-1603.

Giglio L, Kendall J, Mack R (2003) A multi-year active fire dataset for the tropics derived from

the TRMM VIRS. International Journal of Remote Sensing, 24, 4505-4525.

Goel, N., Guttal, V., Levin, S. A., & Staver, A. C. (2020a). Dispersal increases the resilience of

tropical savanna and forest distributions. The American Naturalist, 195(5), 833–850.

Goel, N., Van Vleck, E. S., Aleman, J. C., & Staver, A. C. (2020b). Dispersal limitation and fire

feedbacks maintain mesic savannas in Madagascar. Ecology, 12(1), 145–12.

Hall M (1980) An iron-smelting site in the Hluhluwe Game Reserve, Zululand. Annals of the

Natal Museum, 24, 165-175.

Hanan, N. P., Tredennick, A. T., Prihodko, L., Bucini, G., & Dohn, J. (2014). Analysis of stable

states in global savannas: is the CART pulling the horse?. Global Ecology and

Biogeography, 23(3), 259-263.

Hennenberg KJ, Goetze D, Szarzynski J, Orthmann B, Reineking B, Steinke I, Porembski S

(2008) Detection of seasonal variability in microclimatic borders and ecotones between

forest and savanna. Basic and Applied Ecology, 9, 275-285.

Higgins, SI, & Scheiter, S (2012). Atmospheric CO 2 forces abrupt vegetation shifts locally,

but not globally. Nature, 488(7410), 209-212.

Hirota, M., Nobre, C., Oyama, M. D., & Bustamante, M. M. (2010). The climatic sensitivity of

the forest, savanna and forest–savanna transition in tropical South America. New

Phytologist, 187(3), 707-719.

Hoffmann WA, Geiger EL, Gotsch SG et al. (2012a) Ecological thresholds at the savanna-

forest boundary: how plant traits, resources and fire govern the distribution of tropical

biomes. Ecol Lett, 15, 759-768.

Hoffmann WA, Jaconis SY, Mckinley KL, Geiger EL, Gotsch SG, Franco AC (2012b) Fuels or

microclimate? Understanding the drivers of fire feedbacks at savanna-forest boundaries.

Austral Ecology, 37, 634-643.

Ibanez, T., Borgniet, L., Mangeas, M., Gaucherel, C., Géraux, H., & Hély, C. (2013). Rainforest

and savanna landscape dynamics in New Caledonia: towards a mosaic of stable rainforest

and savanna states?. Austral Ecology, 38(1), 33-45.

Johnson, C. N. (2009). Ecological consequences of Late Quaternary extinctions of

megafauna. Proceedings of the Royal Society B: Biological Sciences, 276(1667), 2509-2519.

Jolly WM, Cochrane MA, Freeborn PH, Holden ZA, Brown TJ, Williamson GJ, Bowman DM

(2015) Climate-induced variations in global wildfire danger from 1979 to 2013. Nat

Commun, 6, 7537.

Keeley JE, Pausas JG (2019) Distinguishing disturbance from perturbations in fire-prone

ecosystems. International Journal of Wildland Fire, 28, 282-287.

Keeley JE, Rundel PW (2005) Fire and the Miocene expansion of C4 grasslands. Ecology

Letters, 8, 683-690.

Kellman, M., & Meave, J. (1997). Fire in the tropical gallery forests of Belize. Journal of

Biogeography, 24(1), 23-34.

Kennard, D. K. (2002). Secondary forest succession in a tropical dry forest: patterns of

development across a 50-year chronosequence in lowland Bolivia. Journal of tropical

ecology, 18(1), 53-66.

Killeen, T. J., Chavez, E., Peña-Claros, M., Toledo, M., Arroyo, L., Caballero, J., ... & Steininger,

M. (2006). The Chiquitano dry forest, the transition between humid and dry forest in

eastern lowland Bolivia. In Neotropical savannas and seasonally dry forests (pp. 214-233).

CRC Press.

Kinyamario J, Trlica M, Njoka T (1995) Influence of tree shade on plant water status, gas

exchange, and water use efficiency of Panicum maximum Jacq. and Themeda triandra Forsk.

in a Kenya savanna. African Journal of Ecology, 33, 114-123.

Legendre P, Legendre LF (2012) Numerical ecology, Elsevier.

Little JK, Prior LD, Williamson GJ, Williams SE, Bowman DMJS (2012) Fire weather risk differs

across rain forest-savanna boundaries in the humid tropics of north-eastern Australia.

Austral Ecology, 37, 915-925.

Moncrieff GR, Scheiter S, Bond WJ, Higgins SI (2014) Increasing atmospheric CO_2 overrides

the historical legacy of multiple stable biome states in Africa. New Phytol, 201, 908-915.

Morton D, Le Page Y, Defries R, Collatz G, Hurtt G (2013) Understorey fire frequency and the

fate of burned forests in southern Amazonia. Philosophical Transactions of the Royal Society

B: Biological Sciences, 368, 20120163.

Murphy, B. P., & Bowman, D. M. (2012). What controls the distribution of tropical forest and

savanna?. Ecology letters, 15(7), 748-758.

Nepstad, D. C., Uhl, C., Pereira, C. A., & Da Silva, J. M. C. (1996). A comparative study of tree

establishment in abandoned pasture and mature forest of eastern Amazonia. Oikos, 25-39.

Nepstad, D. C., Verssimo, A., Alencar, A., Nobre, C., Lima, E., Lefebvre, P., ... & Brooks, V.

(1999). Large-scale impoverishment of Amazonian forests by logging and

fire. Nature, 398(6727), 505-508.

Nyerges, A. E. (1992). Swidden agriculture and the savannization of forests in Sierra

Leone (Doctoral dissertation, Ann Arbor: UMI).

O'Connor T, Everson T (1998) Population dynamics of perennial grasses in African savanna

and grassland. Population biology of grasses, 399.

Oksanen J, Blanchet FG, Kindt R et al. (2013) Package 'vegan'. Community ecology package,

version, 2.

Ondei, S., Prior, L. D., Vigilante, T., & Bowman, D. M. (2017). Fire and cattle disturbance

affects vegetation structure and rain forest expansion into savanna in the Australian

monsoon tropics. Journal of biogeography, 44(10), 2331-2342.

Owen-Smith, N. (1987). Pleistocene extinctions: the pivotal role of

megaherbivores. Paleobiology, 13(3), 351-362.

Pammenter N, Berjak M, Macdonald I (1985) Regeneration of a Natal coastal dune forest

after fire. South African Journal of Botany, 51, 453-459.

Ratnam J, Bond WJ, Fensham RJ et al. (2011) When is a 'forest' a savanna, and why does it

matter? Global Ecology and Biogeography, 20, 653-660.

Rosan, T. M., Aragão, L. E., Oliveras, I., Phillips, O. L., Malhi, Y., Gloor, E., & Wagner, F. H.

(2019). Extensive 21st-century woody encroachment in South America's

savanna. Geophysical Research Letters, 46(12), 6594-6603.

Rull, V., Montoya, E., Nogue, S., Vegas-Vilarrubia, T., & Safont, E. (2013). Ecological

palaeoecology in the neotropical Gran Sabana region: long-term records of vegetation

dynamics as a basis for ecological hypothesis testing. Perspectives in Plant Ecology,

Evolution and Systematics, 15(6), 338-359.

Ryan CM, Williams M (2011) How does fire intensity and frequency affect miombo

woodland tree populations and biomass? Ecological Applications, 21, 48-60.

Sato, H., Kelley, D. I., Mayor, S. J., Martin Calvo, M., Cowling, S. A., & Prentice, I. C. (2021).

Dry corridors opened by fire and low CO_2 in Amazonian rainforest during the Last Glacial

Maximum. Nature Geoscience, 1-8.

Scheiter S, Higgins SI, Osborne CP, Bradshaw C, Lunt D, Ripley BS, Taylor LL, and Beerling DJ

(2012) Fire and fire-adapted vegetation promoted C4 expansion in the late Miocene. New

Phytologist 195, no. 3: 653-666.Schwartz D, Mariotti A, Lanfranchi R, Guillet B (1986)

13C/12C ratios of soil organic matter as indicators of vegetation changes in the Congo.

Geoderma, 39, 97-103.

Schertzer, E., & Staver, A. C. (2018). Fire spread and the issue of community-level selection

in the evolution of flammability. Journal of The Royal Society Interface, 15(147), 20180444.

Schwartz, D., Mariotti, A., Lanfranchi, R., & Guillet, B. (1986). 13C/12C ratios of soil organic

matter as indicators of vegetation changes in the Congo. Geoderma, 39(2), 97-103.

Silva, LCR, Sternberg L, Haridasan M, Hoffmann WA, Miralles-Wilhelm F, and Franco

AC.(2008) Expansion of gallery forests into central Brazilian savannas. Global Change Biology

14, no. 9: 2108-2118.

Silverio DV, Brando PM, Balch JK, Putz FE, Nepstad DC, Oliveira-Santos C, Bustamante MM

(2013) Testing the Amazon savannization hypothesis: fire effects on invasion of a

neotropical forest by native cerrado and exotic pasture grasses. Philos Trans R Soc Lond B

Biol Sci, 368, 20120427.

Simpson KJ, Ripley BS, Christin PA, Belcher CM, Lehmann CE, Thomas GH, and Osborne CP.

(2016) Determinants of flammability in savanna grass species. Journal of Ecology, 104(1),

138-148.

Schmidt, I. B., Ferreira, M. C., Sampaio, A. B., Walter, B. M., Vieira, D. L., & Holl, K. D. (2019).

Tailoring restoration interventions to the grassland-savanna-forest complex in central

Brazil. Restoration Ecology, 27(5), 942-948.

Staal, A., Fetzer, I., Wang-Erlandsson, L., Bosmans, J. H., Dekker, S. C., van Nes, E. H., ... &

Tuinenburg, O. A. (2020). Hysteresis of tropical forests in the 21st century. Nature

communications, 11(1), 1-8.

Staver AC, Archibald S, Levin SA (2011a) The global extent and determinants of savanna and

forest as alternative biome states. Science, 334, 230-232.

Staver AC, Beckett H, Graf J (2017) 3 r Long-Term Vegetation Dynamics within the Hluhluwe

iMfolozi Park. Conserving Africa's Mega-Diversity in the Anthropocene: The Hluhluwe-

iMfolozi Park Story, 56.

Staver AC, Bond WJ, Cramer MD, Wakeling JL (2012) Top-down determinants of niche

structure and adaptation among African Acacias. Ecol Lett, 15, 673-679.

Staver, A. C., & Levin, S. A. (2012). Integrating theoretical climate and fire effects on savanna

and forest systems. The American Naturalist, 180(2), 211-224.

Staver, A. C., Archibald, S., & Levin, S. (2011b). Tree cover in sub-Saharan Africa: rainfall and

fire constrain forest and savanna as alternative stable states. Ecology, 92(5), 1063-1072.

Staver, A. C., Brando, P. M., Barlow, J., Morton, D. C., Paine, C. T., Malhi, Y., ... & del Aguila

Pasquel, J. (2020). Thinner bark increases sensitivity of wetter Amazonian tropical forests to

fire. Ecology Letters, 23(1), 99-106.

Stevens N, Lehmann CE, Murphy BP, Durigan G (2017) Savanna woody encroachment is

widespread across three continents. Glob Chang Biol, 23, 235-244.

Taubert, F., Fischer, R., Groeneveld, J., Lehmann, S., Müller, M. S., Rödig, E., ... & Huth, A.

(2018). Global patterns of tropical forest fragmentation. Nature, 554(7693), 519-522.

Touboul, J. D., Staver, A. C., & Levin, S. A. (2018). On the complex dynamics of savanna

landscapes. Proceedings of the National Academy of Sciences, 115(7), E1336-E1345.

Uhl, C., Clark, H., Clark, K., & Maquirino, P. (1982). Successional patterns associated with

slash-and-burn agriculture in the upper Rio Negro region of the Amazon basin. Biotropica,

249-254.

van de Leemput, I. A., van Nes, E. H., & Scheffer, M. (2015). Resilience of Alternative States

in Spatially Extended Ecosystems. PLoS ONE, 10(2), e0116859.

http://doi.org/10.1371/journal.pone.0116859.s012

Van Nes, E. H., Staal, A., Hantson, S., Holmgren, M., Pueyo, S., Bernardi, R. E., ... & Scheffer,

M. (2018). Fire forbids fifty-fifty forest. PloS one, 13(1), e0191027.

Veenendaal, E. M., Torello-Raventos, M., Feldpausch, T. R., Domingues, T. F., Gerard, F.,

Schrodt, F., ... & Kemp, J. (2015). Structural, physiognomic and above-ground biomass

variation in savanna-forest transition zones on three continents-how different are co-

occurring savanna and forest formations?. Biogeosciences, 12(10), 2927-2951.

Veldman JW, Buisson E, Durigan G et al. (2015a) Toward an old-growth concept for

grasslands, savannas, and woodlands. Frontiers in Ecology and the Environment, 13, 154-

162.

Veldman, J. W., & Putz, F. E. (2010). Long-distance dispersal of invasive grasses by logging

vehicles in a tropical dry forest. Biotropica, 42(6), 697-703.

Veldman JW, Putz FE (2011) Grass-dominated vegetation, not species-diverse natural

savanna, replaces degraded tropical forests on the southern edge of the Amazon Basin.

Biological Conservation, 144, 1419-1429.

Veldman, J. W. (2016). Clarifying the confusion: old-growth savannahs and tropical

ecosystem degradation. Philosophical Transactions of the Royal Society B: Biological

Sciences, 371(1703), 20150306.

Veldman, J.W., Overbeck, G.E., Negreiros, D., Mahy, G., Le Stradic, S., Fernandes, G.W.,

Durigan, G., Buisson, E., Putz, F.E. and Bond, W.J. (2015b). Where tree planting and forest

expansion are bad for biodiversity and ecosystem services. BioScience, 65(10), 1011-1018.

Waldram MS, Bond WJ, Stock WD (2007) Ecological Engineering by a Mega-Grazer: White

Rhino Impacts on a South African Savanna. Ecosystems, 11, 101-112.

Watson, J. E., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C., ... & McAlpine, C.

(2018). The exceptional value of intact forest ecosystems. Nature ecology & evolution, 2(4),

599-610.

West, A. G., Bond, W. J., & Midgley, J. J. (2000). Soil carbon isotopes reveal ancient grassland

under forest. S. Afr. J. Sci, 96, 253.

Whateley A, Porter R (1983) The woody vegetation communities of the Hluhluwe-Corridor-

Umfolozi game reserve complex. Bothalia, 14, 745-758.

Wickham H (2017) The tidyverse. R package ver. 1.1. 1.

Woodward FI, Lomas MR, Kelly CK (2004) Global climate and the distribution of plant

biomes. Philos Trans R Soc Lond B Biol Sci, 359, 1465-1476.

Wright, J. L., Bomfim, B., Wong, C. I., Marimon-Júnior, B. H., Marimon, B. S., & Silva, L. C.

(2021). Sixteen hundred years of increasing tree cover prior to modern deforestation in

Southern Amazon and Central Brazilian savannas. Global Change Biology, 27(1), 136-150.

Wuyts, B., Champneys, A. R., Verschueren, N., & House, J. I. (2019). Tropical tree cover in a

heterogeneous environment: A reaction-diffusion model. PloS one, 14(6), e0218151.

Acknowledgments

797 The financial assistance of SAEON towards this research is hereby acknowledged. Opinions

798 expressed and conclusions arrived at are those of the author and are not necessarily to be

799 attributed to SAEON.

Author Contributions

801 HB, WJB and ASC conceived of the idea. TCD and ASC contributed data. HB collected the

802 data and designed and performed the analyses with input from TCD and ASC. HB wrote the

803 paper in consultation with WJB, ASC and TCD.

Figure Legends

Figure 1. Fire Frequency map (left) of Hluhluwe iMfolozi Park from 1955 to 2013 based on

Hluhluwe iMfolozi Park records with the Hluhluwe section outlined in red, and (right) the

distribution of plots within the Hluhluwe portion of HiP and biomes based on a

reclassification of Whateley & Porter (1983). Reclassification based on Charles-Dominique *et*

al (2015). Green represents forest, blue represents thickets and grey represents savannas.

Inset map shows the location of Hluhluwe iMfolozi Park within South Africa.

Figure 2. Fire Frequency for plots within each vegetation type (indicated by grey text) before

and after extreme fire events (not including the year of the extreme fire event), calculated

using the MODIS Active Fires Product from 2000 to 2016. Grey lines link plot points before

and after extreme fire events. Asterisks indicate significance level (*** <0.05, **** < 0.001,

NS = no significant difference).

Figure 3. Changes in the (A) total biomass of grass found in all plots for frequently-burnt

forests, once-burnt forest, frequently-burnt thicket and savanna, and two dominant grass

species, (B) Panicum maximum and (C) Themeda triandra. Green bars represent forest plots,

blue represents thicket, and yellow represents savanna. Error bars indicate standard error.

Points (plot grass biomass) are included to display variation. Shared letters indicate lack of a

significant difference within panels based on Wilcoxon signed-rank test, p < 0.05.

Figure 4. Mean cumulative length of all tree species within the four size classes 'before' and

'after' the extreme fire event in A) Savanna, B) Frequently-burnt Thicket, C) Frequently-

burnt Forest, and D) Once-burnt Forest. Error bars indicate standard error. Points represent

individual plots.

Figure 5. NMDS ordination of the cumulative length of all species in all height classes in

plots. Ellipses show the grouping of plots. Species associations used in the composition

analyses are based on species found in plots before the extreme fire events only. Letters

indicate the position of plots, grey dots indicate the species scores.

Figure 6. Initial and final composition and structure of savannas in 2007 (A) and 2014 (B),

intact thicket (C) and frequently-burnt thicket 7 years after extreme fire event (D), intact

forest (E) and frequently-burnt forest 7 years after extreme fire event (F) and intact forest

(G) and once-burnt forest 6 years after fire (H). Panels display the average cumulative length

of tree species within each size class for the different time periods 'before' and 'after' the

extreme fire event. Colours indicate the biome in which these species are predominantly

found. Pioneer species are those which were not found in plots sampled before the extreme

fire event. Data recorded in intact forest and thicket vegetation represent pre-fire space for

time substitutions.

	1-4
	5-8
	9-12
	13-16
	17-20
	21-24
	25-28
	28-32
	>32

Figure 1. Fire Frequency map (left) of Hluhluwe iMfolozi Park from 1955 to 2013

based on Hluhluwe iMfolozi Park records with the Hluhluwe Game Reserve section

outlined in red, and (right) the distribution of plots within the Hluhluwe Game Reserve

portion of HiP and biomes based on a reclassification of Whateley & Porter (1983).

Reclassification based on Charles-Dominique *et al* (2015). Green represents forest,

blue represents thickets and grey represents savannas. Inset map shows the location

of Hluhluwe iMfolozi Park within South Africa.

Figure 2. Fire Frequency for plots within each vegetation type (indicated by grey text) before

and after extreme fire events (not including the year of the extreme fire event), calculated

using the MODIS Active Fires Product. Grey lines link plot points before and after extreme

fire events. Asterisks indicate significance level (*** <0.05, **** < 0.001, NS = no significant

difference).

Figure 3. Changes in the (A) total biomass of grass found in all plots for frequently-burnt

forests, once-burnt forest, frequently-burnt thicket and savanna, and two dominant grass

species, (B) Panicum maximum and (C) Themeda triandra. Green bars represent forest plots,

blue represents thicket, and yellow represents savanna. Error bars indicate standard error.

Points (plot grass biomass) are included to display variation. Shared letters indicate lack of a

significant difference within panels based on Wilcoxon signed-rank test, p < 0.05.

Figure 4. Mean cumulative length of all tree species within the four size classes 'before' and 'after' the extreme fire event in A) Savanna, B) Frequently-burnt Thicket, C) Frequently-burnt Forest, and D) Once-burnt Forest. Error bars indicate standard error. Points represent individual plots.

Figure 5. NMDS ordination of the cumulative length of all species in all height classes in transects. Ellipses show the grouping of transects. Species associations used in the composition analyses are based on species found in transects before the extreme fire events only. Letters indicate the position of transects, grey dots indicate the species scores.

Figure 6. Initial and final composition and structure of savannas in 2007 (A) and 2014 (B), intact thicket (C) and frequently-burnt thicket 7 years after extreme fire event (D), intact forest (E) and frequently-burnt forest 7 years after extreme fire event (F) and intact forest (G) and once-burnt forest 6 years after fire (H). Panels display the average cumulative length of tree species within each size class for the different time periods 'before' and 'after' the extreme fire event. Colours indicate the biome in which these species are predominantly found. Pioneer species are those which were

944   not found in plots sampled before the extreme fire event. Data recorded in intact

945   forest and thicket vegetation represent pre-fire space for time substitutions.